Title

Assessing functional connectivity differences and work-related fatigue in surviving COVID-negative patients.

Running title

Functional Alterations and Fatigue in COVID

Author Names and Affiliations

Rakibul Hafiz [‡1], Tapan Kumar Gandhi[‡2], Sapna Mishra[2], Alok Prasad[3], Vidur Mahajan[4], Benjamin H. Natelson[5], Xin Di[1], Bharat B. Biswal*[1]

[1]: Department of Biomedical Engineering, New Jersey Institute of Technology (NJIT), 323 Dr Martin Luther King Jr Blvd, Newark, NJ 07102, USA

[2]: Department of Electrical Engineering, Indian Institute of Technology (IIT), Block II, IIT Delhi Main Rd, IIT Campus, Hauz Khas, New Delhi, Delhi 110016, India

[3]: Internal Medicine, Irene Hospital & Senior Consultant Medicine, Metro Heart and Super-specialty Hospital, New Delhi, India

[4]: Centre for Advanced Research in Imaging, Neuroscience & Genomics, Mahajan Imaging, New Delhi, India

[5]: Pain and Fatigue Study Center, Department of Neurology, Icahn School of Medicine at Mount Sinai, NY, USA

[‡]: The authors contributed equally for this project as first authors.

* Corresponding Author e-mail

 bharat.biswal@njit.edu

Number of pages: 17

Number of figures: 3

Number of Words

Abstract: 176

Introduction: 1103

Discussion: 868

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgements: This study was supported by an NIH grant (R01AT009829).

Abstract

The recent Coronavirus Disease 2019 (COVID-19) has affected all aspects of life around the world. Neuroimaging evidence suggests the novel coronavirus can attack the central nervous system (CNS), causing cerebro-vascular abnormalities in the brain. This can lead to focal changes in cerebral blood flow and metabolic oxygen consumption rate in the brain. However, the extent and spatial locations of brain alterations in COVID-19 survivors are largely unknown. In this study, we have assessed brain functional connectivity (FC) using resting-state functional MRI (RS-fMRI) in 38 (25 males) COVID patients two weeks after hospital discharge, when PCR negative and 31 (24 males) healthy subjects. FC was estimated using independent component analysis (ICA) and dual regression. The COVID group demonstrated significantly enhanced FC in regions from the Occipital and Parietal Lobes, comparing to the HC group. On the other hand, the COVID group exhibited significantly reduced FC in several vermal layers of the cerebellum. More importantly, we noticed negative correlation of FC with self-reported fatigue within regions from the Parietal lobe, which are known to be associated with fatigue.

Keywords: *COVID, Functional Connectivity, ICA, Fatigue, RS-fMRI*

Significance Statement

Early neuroimaging studies have mostly focused on structural MRI imaging to report brain abnormalities in acutely ill COVID-19 patients. It is not clear whether functional abnormalities co-exist with structural alterations in patients who have survived the infection and have been discharged from the hospital. A few recent studies have emerged which attempted to address the structural/functional alterations. However, further

investigations across different sites are necessary for more conclusive inference. More

importantly, fatigue is a highly prevalent symptom among COVID survivors, therefore, the

relations of brain imaging abnormalities to fatigue should be investigated. In this study,

we try to address these gaps, by collecting imaging data from COVID survivors, now PCR

negative, and healthy subjects from a single site – the Indian Institute of Technology (IIT),

Delhi, India. Furthermore, this is a continuation of an ongoing study. We have already

submitted a manuscript showing structural abnormalities and gray matter volume

correlates of self-reported fatigue among this group of COVID survivors.

Introduction

The novel coronavirus pandemic has taken more than 4.5 million lives across the globe

((WHO), 2020). With efforts of vaccination, mask mandates and social distancing, the

spread of this contagious disease has been mitigated significantly, however, advents of

new strains such as the delta and omicron variants have been setting back progress and

especially affecting densely populated countries (India, being a prime example). While

the initial wave demanded most medical attention towards severe damage to the

respiratory system, recent evidence suggests the novel coronavirus, Severe Acute

Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) can also attack the central nervous

system (CNS). Early pandemic MRI reports from acutely ill patients show evidence of a

wide range of cerebrovascular abnormalities. A review featuring 22 articles (n = 126

patients) from seven countries (Gulko et al., 2020) showed that patients with SARS-CoV-

2 infection showed acute infarcts, posterior reversible encephalopathy syndrome,

hyperintensities from fluid-attenuated inversion recovery (FLAIR) images and

microhemorrhages. Another multicenter study (n = 64) also reported higher rates of

ischemic strokes (27%) and encephalitis (13%). Moreover, inflammatory vascular

pathologies (Keller et al., 2020) and other cerebrovascular abnormalities (Nicholson et

al., 2020) have also been reported. Abnormal FLAIR uptakes were also reported in the

parietal and occipital lobes among others (Kandemirli et al., 2020), as well as in the

hypothalamus and the thalamus in individual patients (Paterson et al., 2020).

These single-case reports played a vital role in informing and shaping recent studies with

primary focus on group level structural differences in moderate (Duan et al., 2021; Qin et

al., 2021) to large sample groups (Douaud et al., 2021). Understandably, the initial target

of most neuroimaging studies was brain abnormalities in severe patients. Therefore, a

large sample of hospitalized survivors were not investigated, especially those with

persistent symptoms. This led to a rise in follow-up studies on a span of 3 to 6 months

after initial infection (Lu et al., 2020; Tu et al., 2021) and longitudinal designs (Douaud et

al., 2021) where structural abnormalities were investigated before and after the pandemic.

We have recently shown gray matter volume differences in survivors after a shorter

interval (2 weeks after hospital discharge), as well as, a relation between gray matter

volume and self-reported fatigue at work (Hafiz et al., 2021) (in press). It is still unclear,

though, if such structural abnormalities are also accompanied by functional brain

alterations in COVID-19 survivors.

To address this gap, functional brain imaging can be incorporated through functional

magnetic resonance imaging (fMRI). Among the earliest literature, a task-based fMRI

study reported loss in task activation in the orbito-frontal cortex (OFC) and strong BOLD

activations in the piriform cortex (Ismail & Gad, 2021) from a single female (25 years)

COVID patient with persistent olfactory dysfunction. They used a simple smell on/off

block design task. A single case resting state fMRI (RS-fMRI) study from an

unresponsive patient reported intact functional connectivity (FC) of the default mode

network (DMN) (Fischer et al., 2020), which was a good prognosis for ultimate recovery.

However, these studies were case reports, which leaves the question of whether there

are generalizable group level functional brain alterations in COVID survivors. To that

end, a few studies have emerged that report various functional abnormalities among

COVID survivors. For example, the initial case report from (Fischer et al., 2020) has

now been followed up with a group level report with specific focus on severe patients

who were initially unresponsive, but recovered completely and were able to return

(Fischer et al., 2021) to pre-COVID level behaviorally. When they compared the

functional connectivity of these unresponsive patients with healthy controls, they found

significantly reduced default mode network (DMN) connectivity and reduction in inter-

network connectivity between DMN and salience (SAL) networks. Currently, a growing

concern among survivors is persistence of a sequela of symptoms (Logue et al., 2021;

Peluso et al., 2021; Tabacof et al., 2020), now commonly called 'Long COVID', which

point to brain as the responsible organ. Fatigue, lack of attention, anxiety, memory loss,

delayed recovery of smell and/or taste, muscle pain and stress are some of the

commonly reported symptoms among many others.

Since several of these symptoms suggest cognitive abnormalities among survivors,

most contemporary neuroimaging studies have turned their attention to behavioral

correlates of functional brain alterations, primarily, post-traumatic stress syndromes

(Benedetti et al., 2021; Fu et al., 2021). On the other hand, several others have

attempted to use functional connectivity (FC) as a neurobiological indicator of higher

stress levels (Liu et al., 2021; Perica et al., 2021), depression (Zhang et al., 2022) and

negative affect (Xiao et al., 2021) among only healthy subjects before and after the

pandemic.

Despite fatigue being one of the most frequently reported symptoms, very little is known

of functional brain correlates of fatigue within survivors. Prior to the current study, we

have shown a positive correlation of gray matter volume within regions from the *ventral*

basal ganglia (BG) and *ventromedial prefrontal cortex (vmPFC)* with self-reported

fatigue at work in survivors two weeks after hospital discharge (Hafiz et al., 2021) (in

press). Here, we continue our investigation to explore the functional correlates of fatigue

among the same set of survivors. Moreover, functional changes can occur across

different networks among survivors, owing to a range of symptoms experienced during

the recovery phase. Therefore, we applied a data driven approach to estimate FC

differences between healthy controls (HCs) and surviving, now COVID-negative,

patients using RS-fMRI.

The earliest 'resting state' study was done by Biswal et al., in 1995 (Biswal et al., 1995),

and subsequent studies have shown that spatially distinct regions that are temporally

synchronized may share information with each other (Cole et al., 2010; De Luca et al.,

2006; Fox et al., 2005; Fox et al., 2006; Greicius et al., 2003; Kalcher et al., 2012; Meier

et al., 2012). Independent Component Analysis (ICA), is a data driven technique which

groups all voxels in the brain into distinct spatial networks based on the similarity of time

courses (McKeown et al., 1998). The large-scale resting-state networks (RSNs) derived

from ICA have been shown to have local and higher level associative hierarchy (Yeo et

al., 2011) and replicate highly reproducible activation maps across subjects (Smith et al., 2009). FC estimates from group ICA and dual regression (C.F. Beckmann, 2009; Filippini et al., 2009) were used to test our hypothesis that surviving COVID-negative patients would demonstrate altered FC in RSNs comprising of cortical regions where hyperintensities have been reported from single cases and group level differences identified from recent neuroimaging studies. We further hypothesized that FC would demonstrate significant correlation with self-reported fatigue scores in brain regions, known to be associated with fatigue.

Materials and Methods

Participants

This is a continuation of our recent publication using the same sample groups where structural brain alterations were reported (Hafiz et al., 2021). 47 COVID patients and 35 healthy controls were recruited from a single site located at Indian Institute of Technology (IIT), Delhi, India. 9 COVID and 4 HC subjects were removed during quality control and motion assessment, leaving with an effective sample of 38 (25 males) COVID and 31 (24 males) HC. The mean age of the COVID group was 34.79 years (SD = $\pm$ 10.41 years); and 32.68 years (SD = $\pm$ 9.78 years) for the HC group. The COVID subjects were scanned two weeks after they were released from the hospital when confirmed to be COVID-negative upon polymerase chain reaction (PCR) retesting. During scanning, all protocols were strictly followed based on the Institutional Review Board (IRB) guidelines at the Indian Institute of Technology (IIT), Delhi, India.

Clinical Assessment

The most frequently observed symptoms from the participants during hospitalization were - fever, cough, body ache, chills, difficulty breathing, bowel irritation, nausea, loss of sense of smell and loss of consciousness. From the day of discharge till the day of scan, we further asked if the participants were experiencing any persistent or new symptoms. Work-related fatigue (65.6%), muscle pain (50%), lack of sleep (50%), lack of attention (43.8%), headache (40.6%), joint pain (40.6%), memory loss (28.1%), delayed recovery of sense of smell (39%) and/or taste (31%), bowel irritation (33%) and interestingly, hair loss (66%) were commonly reported. Please note, most survivors experienced multiple symptoms simultaneously, hence the '%' represents symptoms that overlap within participants. For example, 43.8% of 27 post-COVID participants reporting with lack of attention also reported a work-related fatigue score > 2 on a scale of 0 to 5, with 0 representing no fatigue and 5 representing the highest fatigue possible. The average fatigue score in this sub-set of COVID participants was 2.93/5 ± 1.21 [SD].

Brain Imaging

Anatomical MRI – High-resolution T1-weighted images were acquired on a 3T GE scanner with a 32-channel head coil in 3D imaging mode with a fast BRAVO sequence. The imaging parameters were TI = 450 ms; 244 x 200 matrix; Flip angle = 12 and FOV = 256 mm. The subject was placed in a supine position and the whole brain was scanned in the sagittal configuration where 152 slices were collected, and each slice was 1.00 mm thick. The spatial resolution of all the anatomical scans was 1.0 mm x 1.0 mm x 1.0 mm.

Resting-state fMRI – A gradient echo planar imaging (EPI) was used to obtain 200 whole-brain functional volumes. The parameters were: TR = 2000 ms; TE = 30 ms; Flip angle = 90, 38 slices, matrix = 64x64; FOV = 240 x 240 mm^2; acquisition voxel size = 3.75 x 3.75 x 3 mm^3. The participant was requested to stay as still and motionless as possible with eyes fixed to a cross on an overhead screen.

Data Pre-Processing

The data preprocessing was performed primarily using Statistical Parametric Mapping 12 (SPM12) toolbox (http://www.fil.ion.ucl.ac.uk/spm/) within a MATLAB environment (The MathWorks, Inc., Natick, MA, USA). However, some steps utilized useful tools from FSL (FMRIB Analysis Group, Oxford, UK) and AFNI (http://afni.nimh.nih.gov/afni) (Cox, 1996) for housekeeping, visual inspection and quality control purposes. At the beginning, first five time points were excluded from each subject to account for magnetic stabilization. The functional images were motion corrected for head movement using a least squared approach and 6 parameters (rigid body) spatial transformation with respect to the mean image of the scan. The subjects with excessive head motion were identified using framewise displacement (FWD) (Power et al., 2012). Additionally, time frames with high FWD crossing a threshold of 0.5 mm (Power et al., 2012) were identified along with the previous and the next two frames and added as regressors (Yan et al., 2016) during temporal regression of nuisance signals. If more than 50% of the time series data were affected due to regression of high motion frames the participant was removed from the analysis. Moreover, any participant with the maximum framewise translation or rotation exceeding 2 mm was removed from further analysis. Anatomical image from each subject was coregistered to the mean functional image obtained from the motion correction step.

T1-weighted image from each subject was segmented into gray matter (GM), white matter

(WM), and cerebrospinal fluid (CSF) tissue probability maps and an average template

including all participants was generated using DARTEL (Ashburner, 2007). This template

was used to spatially normalize all functional images to the MNI space and resampled to

isotropic voxel size of 3 mm x 3 mm x 3 mm. Time series, from brain compartments with

high physiological noise signals such as, CSF and WM was extracted by thresholding the

probability maps from the segmentation stage above the 99[th] percentile, and first 5

principial components were obtained using a COMPCOR based (Behzadi et al., 2007)

principal component analysis (PCA) from both tissues. These 10 components along with

Friston's 24- parameter model (6 head motion parameters + 6 previous time point motion

parameters + 12 corresponding quadratic parameters) (Friston et al., 1996) and time

frames with high FWD (> 0.5 mm) were added as regressors in a multiple linear

regression model to remove unwanted signals voxel-wise. The residuals from the

regression step were then bandpass filtered between 0.01 to 0.1 Hz and finally, spatial

smoothing was performed using a Gaussian kernel of 6 mm full width at half maximum

(FWHM).

Head Motion Assessment

We performed in-scanner head movement assessment using mean Framewise

Displacement (FWD) based on the methods depicted in (Power et al., 2012). A two-tailed

two-sample student's t-test revealed no significant differences in mean FWD between the

two groups ($t = -1.57, p = 0.12, α = 0.05$).

ICA and Dual Regression

Group level resting state networks were obtained by applying the 'gica' option of the
'melodic' module from FSL toolbox (FMRIB Analysis Group, Oxford, UK). All subjects' 4D
functional images after pre-processing were temporally concatenated into a 2D matrix of
'space' x 'time' as delineated in (C.F. Beckmann, 2009) and 25 spatial maps were
obtained. Resting State Networks (RSNs) were identified by matching ICs with the 1000
functional connectome project maps (Biswal et al., 2010) using Dice's coefficient and
spatial correlations obtained from AFNI's '3dMatch' program (Taylor & Saad, 2013).
Further visual inspection was performed to make sure all network regions aligned with
the functional network and ROIs depicted in (Altmann et al., 2015; Shirer et al., 2012).
Dual regression (C.F. Beckmann, 2009; Filippini et al., 2009) was performed leveraging
the standardized group ICA output from the 'melodic' step and applying it directly to the
'fsl-glm' module in FSL to obtain subject specific RSN maps. The subject specific network
maps were standardized to Z-scores before consequently applying them in statistical
analysis to infer group level estimates.

Statistical Analysis

To investigate FC differences between COVID and HC groups, we performed an unpaired
two sample t-test between standardized subject-specific RSN maps from the two groups.
Significant clusters were identified and main effect of interest from the corresponding
contrast maps representing the difference in mean beta scores from two groups were
obtained by thresholding the t-score map values that survived the corrected threshold. To
account for confounding effects that may explain some of the variance in the data, age
and sex were also added as covariates of no interest. Cluster-based thresholding was

applied at height threshold of $p_{unc} < 0.01$, with *family wise error* (*FWE*) correction at p_{FWE}

< 0.05 for multiple comparisons. The cluster extent threshold (k_E) obtained from this step

was used to generate corrected statistical maps for the contrasts with significant effects.

We further wanted to evaluate which of the large scale RSNs demonstrates linear

relationship or correlation with self-reported fatigue among the COVID individuals. We

incorporated a multiple linear regression approach where the FC at each voxel was the

response variable (Y), and the self-reported fatigue score was the explanatory variable

(X). We also added age and sex as covariates of no interest. Significant clusters were

obtained in the same manner as described earlier in this section for group level

differences in FC. For visual representation of the significant linear relationship between

the two variables, the average FC within the significant cluster was obtained from each

subject. These average FC values were then linearly regressed against the fatigue scores

and visualized within a scatter plot and a line of best fit with 95% confidence interval. Age

and sex were regressed out during the linear regression step. The correlation analysis

and the graphical plotting was done using 'inhouse' scripts prepared in RStudio (RStudio,

2021).

Results

We identified twenty-two large-scale resting state networks (RSNs) (see Figure1) from

the group ICA analysis. Group level statistical analysis was run for each network using

standardized subject specific RSN maps obtained from the dual regression step.

Significantly enhanced FC was observed in the COVID group compared to the HC group

in particularly regions from the *occipital* and *parietal* lobes. Figures2 and 3 show all

significant clusters from the FC and linear regression analysis, respectively.

Figure2 shows the results from the group level analysis from three RSNs. Figure2 A (top

row) demonstrates regions with significantly enhanced FC in the COVID-19 group

compared to the HC group for the *BGN* network. The FC difference was observed in the

Right – Calcarine Cortex (Calc), Cuneus (Cu) and *Lingual Gyrus (LiG)* regions of the

occipital lobe*.* Similarly, the COVID survivors also demonstrated enhanced FC of the *PRN*

network (Figure2 B) with regions from the *Parietal Lobe*: *Bilateral – Superior Parietal*

Lobule (SPL) and Precuneus (PCu) regions. On the other hand, Figure2 C shows reduced

functional connectivity among COVID participants compared to HCs in several layers of

the *cerebellar vermal lobules (CVL) (I-V, VI-VII)* for the *LANG* network. The cluster peak

information including peak T-scores and *FWE* corrected exact p-values with relevant

spatial regions from each network showing significant differences have been tabulated

for an easy reference in Table1*.*

Figure3 shows brain regions where a significant negative correlation was observed

between FC and self-reported fatigue, from the *PRN* network. Figure3 (A) (left) shows the

cluster where a negative correlation between FC and fatigue scores was observed in the

Left – Superior Parietal Lobule (SPL), Superior Occipital Gyrus (SOG), Angular Gyrus

(AnG) and *Precuneus (PCu).* The graph on the right visually presents this negative

relationship ($r = -0.75$, $p = 0.00001$, $r^2 = 0.56$) between the average FC of this cluster and

fatigue scores.

Discussion

The results from this study support our hypothesis that COVID survivors would

demonstrate altered FC when compared to HCs, even two weeks after discharge from

the hospital and demonstrate significant linear relationship with work-related fatigue. Our

hypothesis was based on both early case-reports and more recent group level

neuroimaging reports of structural and functional brain alterations. Individual case reports

were primarily from acutely ill patients using FLAIR (Kandemirli et al., 2020; Kremer et

al., 2020; Paterson et al., 2020) and Susceptibility Weighted Imaging (SWI) (Conklin et

al., 2021), whereas, group level reports, such as those derived from fMRI, include,

reduced *default mode* and *salience* connectivity (Fischer et al., 2021) and high prevalence

of abnormal time varying and topological organizations between *sensorimotor* and *visual*

networks (Fu et al., 2021). In the current context, we report between group FC alterations

of three large scale RSNs – *BGN*, *PRN* and *LANG* networks and further show negative

correlation of FC from the *PRN* network with self-reported fatigue at work among COVID

survivors.

While initially a single patient showed no differences in FC of *DMN* when compared to

five healthy controls (Fischer et al., 2020), Fischer and colleagues recently reported

reduced FC within *DMN* and between *DMN* and *SAL* networks after group level

assessment (Fischer et al., 2021). In the current study, we did not observe any significant

alterations in posterior or ventral DMN (PDMN and VDMN) networks, but our patient

group was not unresponsive and as acutely ill as the patients reported in (Fischer et al.,

2021). However, we did observe differences in FC for the PRN network which consists of

Precuneus (PCu), Frontal Eye Fields (FEF) and parts of the *Superior Parietal Lobule*

(SPL). Enhanced FC in this network was observed in the *Bilateral SPL* and *PCu* regions.

PCu is a constituent of DMN, and higher functional connectivity with this region may

indicate some compensatory mechanism due to loss in connections in other pathways.

Furthermore, *SPL* is a constituent of the *posterior parietal cortex (PPC)* which has been

shown to have functional association with altered *anterior insula* connectivity in chronic

fatigue syndrome (CFS) (Wortinger et al., 2017). Moreover, these brain regions are also

known to be involved in attention processing, therefore, enhanced FC in these regions

may indicate possible compensatory mechanisms of attention related symptoms that

recovering patients may experience. Therefore, further investigations are necessary to

understand these processes better, especially, from a clinical perspective.

Enhanced FC in the COVID group was also observed for the *BGN* network within the

occipital lobe (*Calc, Cu* and *LiG*). *Calc* and *Cu* are primarily involved in visual processing.

Fu and colleagues reported that COVID survivors had higher connectivity between

Cerebellum, *Sensorimotor* and *Visual* networks, indicating they spent abnormally higher

time in a specific brain state compared to healthy controls (Fu et al., 2021). A recent study

has also suggested *Cu* to be a major hub for mild cognitive impairment in idiopathic REM

sleep behavior disorder (iRBD) (Mattioli et al., 2021). *LiG* and weak *insular* coactivation

with the *occipital* cortex have been shown to be associated with disrupted salience

processing that can lead to loss in motivation in day-to-day tasks (Kim et al., 2018).

Moreover, the *basal ganglia* are known to be associated with fatigue (Miller et al., 2014),

cognitive, emotional and attention processing (Di Martino et al., 2008; van Schouwenburg

et al., 2015). We also observed reduced FC within several layers of the *Cerebellar Vermal*

Lobules among COVID participants when compared to HCs. These lobules have been

suggested to be involved in cognition and emotion processing (Park et al., 2018). The

synergy of these studies to our findings indicates possible functional brain associations

of commonly observed symptoms in survivors with post-acute sequelae SARS-CoV-2

infection (PASC or Long COVID) lasting many months (Carfì et al., 2020; Garrigues et

al., 2020; Logue et al., 2021; Peluso et al., 2021). FC alterations in multiple networks also

suggest that RS-fMRI can be quite useful to investigate multiple brain networks across

the whole brain (Damoiseaux et al., 2006; Raichle & Mintun, 2006; Shulman et al., 2004)

in COVID-19 patients. The results also suggest a possible link between structural and

functional abnormalities in COVID patients since the FC alterations were observed in

regions that align with anatomical regions exhibiting hyperintensities from FLAIR and SWI

studies.

We further evaluated linear relationship between FC of RSNs and self-reported fatigue at

work among COVID participants. We observed a significant negative correlation of FC

with fatigue within the *Left SPL, SOG, AnG* and *PCu,* i.e., brain regions primarily

belonging to the *parietal* lobe (see Table2 for cluster information). Structural atrophy in

the *parietal* lobe has been shown to be associated with fatigue among multiple sclerosis

(MS) patients (Calabrese et al., 2010; Pellicano et al., 2010). An RS-fMRI study of patients

with chronic fatigue syndrome (CFS) used ICA to reveal loss of intrinsic connectivity in

the parietal lobe (Gay et al., 2016). It is interesting that lower FC in the parietal lobe

correlates negatively to higher fatigue scores among COVID survivors. To the best of our

knowledge, this is the first study to show work-related fatigue correlates of FC among

recovering patients 2 weeks after hospital discharge. Therefore, future studies are

necessary to evaluate this avenue further in the surviving cohorts.

337   **Limitations**

Despite our efforts to show group level effects that reflect individual and group level

reports in the recent literature, our study still maintains a cross-sectional design. In cases

like this, a better approach for the future would be to use follow up designs (Fu et al.,

2021; Lu et al., 2020; Tu et al., 2021) or possibly a longitudinal design where patients

could be observed both before and after the pandemic like the one using the UK-biobank

(Douaud et al., 2021). Our effort here, was to show group level effects at an early stage

of recovery (2 weeks after hospital discharge) and determine the relation between work-

related fatigue and FC of RSNs. We believe the results from this study will help

understanding the recovery stage brain alterations and how they might drive fatigue-

related symptoms among COVID survivors.

Author Credit Statement

Rakibul Hafiz: Methodology, Software, Formal Analysis, Data Curation, Writing –

Original Draft, Review and Editing.

Tapan K. Gandhi: Conceptualization, Investigation, Resources, Supervision, Writing –

Review and Editing.

Sapna Mishra: Investigation, Resources, Data Curation, Writing – Review and Editing.

Alok Prasad: Writing – Review and Editing.

Vidur Mahajan: Writing – Review and Editing.

Xin Di: Methodology, Writing – Review and Editing.

Benjamin H. Natelson: Writing – Review and Editing.

Bharat Biswal: Conceptualization, Resources, Project Administration, Supervision,

Writing – Review and Editing.

References

(WHO), W. H. O. (2020, 10/21/2021, 4:46pm CET). *WHO Coronavirus Disease (COVID-19) Dashboard*. Retrieved 10/21/2021 from https://covid19.who.int/table

Altmann, A., Ng, B., Landau, S. M., Jagust, W. J., & Greicius, M. D. (2015). Regional brain hypometabolism is unrelated to regional amyloid plaque burden. *Brain*, *138*(Pt 12), 3734-3746. https://doi.org/10.1093/brain/awv278

Ashburner, J. (2007). A fast diffeomorphic image registration algorithm. *Neuroimage*, *38*(1), 95-113. https://doi.org/10.1016/j.neuroimage.2007.07.007

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage*, *37*(1), 90-101. https://doi.org/10.1016/j.neuroimage.2007.04.042

Benedetti, F., Palladini, M., Paolini, M., Melloni, E., Vai, B., De Lorenzo, R., Furlan, R., Rovere-Querini, P., Falini, A., & Mazza, M. G. (2021). Brain correlates of depression, post-traumatic distress, and inflammatory biomarkers in COVID-19 survivors: A multimodal magnetic resonance imaging study. *Brain, behavior, & immunity - health*, *18*, 100387-100387. https://doi.org/10.1016/j.bbih.2021.100387

Biswal, B., Yetkin, F. Z., Haughton, V. M., & Hyde, J. S. (1995). Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. *Magn Reson Med*, *34*(4), 537-541.

Biswal, B. B., Mennes, M., Zuo, X.-N., Gohel, S., Kelly, C., Smith, S. M., Beckmann, C. F., Adelstein, J. S., Buckner, R. L., Colcombe, S., Dogonowski, A.-M., Ernst, M., Fair, D., Hampson, M., Hoptman, M. J., Hyde, J. S., Kiviniemi, V. J., Kötter, R., Li, S.-J., . . . Milham, M. P. (2010). Toward discovery science of human brain function. *Proceedings of the National Academy of Sciences*, *107*(10), 4734-4739. https://doi.org/10.1073/pnas.0911855107

C.F. Beckmann, C. E. M., N. Filippini, and S.M. Smith. (2009). Group comparison of resting-state FMRI data using multi-subject ICA and dual regression. *OHBM*.

Calabrese, M., Rinaldi, F., Grossi, P., Mattisi, I., Bernardi, V., Favaretto, A., Perini, P., & Gallo, P. (2010). Basal ganglia and frontal/parietal cortical atrophy is associated with fatigue in relapsing-remitting multiple sclerosis. *Mult Scler*, *16*(10), 1220-1228. https://doi.org/10.1177/1352458510376405

Carfì, A., Bernabei, R., Landi, F., & Gemelli Against, C.-P.-A. C. S. G. (2020). Persistent Symptoms in Patients After Acute COVID-19. *JAMA*, *324*(6), 603-605. https://doi.org/10.1001/jama.2020.12603

Cole, D. M., Smith, S. M., & Beckmann, C. F. (2010). Advances and pitfalls in the analysis and interpretation of resting-state FMRI data. *Front Syst Neurosci*, *4*, 8. https://doi.org/10.3389/fnsys.2010.00008

Conklin, J., Frosch, M. P., Mukerji, S. S., Rapalino, O., Maher, M. D., Schaefer, P. W., Lev, M. H., Gonzalez, R. G., Das, S., Champion, S. N., Magdamo, C., Sen, P., Harrold, G. K., Alabsi, H., Normandin, E., Shaw, B., Lemieux, J. E., Sabeti, P. C., Branda, J. A., . . . Edlow, B. L. (2021). Susceptibility-weighted imaging reveals cerebral microvascular injury in severe COVID-19. *Journal of the Neurological Sciences*, *421*, 117308-117308. https://doi.org/10.1016/j.jns.2021.117308

Cox, R. W. (1996). AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. *Comput Biomed Res*, *29*(3), 162-173.

Damoiseaux, J. S., Rombouts, S. A., Barkhof, F., Scheltens, P., Stam, C. J., Smith, S. M., &
Beckmann, C. F. (2006). Consistent resting-state networks across healthy subjects.
Proc Natl Acad Sci U S A, *103*(37), 13848-13853.
https://doi.org/10.1073/pnas.0601417103
De Luca, M., Beckmann, C. F., De Stefano, N., Matthews, P. M., & Smith, S. M. (2006). fMRI
resting state networks define distinct modes of long-distance interactions in the
human brain. *Neuroimage*, *29*(4), 1359-1367.
https://doi.org/10.1016/j.neuroimage.2005.08.035
Di Martino, A., Scheres, A., Margulies, D. S., Kelly, A. M., Uddin, L. Q., Shehzad, Z., Biswal, B.,
Walters, J. R., Castellanos, F. X., & Milham, M. P. (2008). Functional connectivity of
human striatum: a resting state FMRI study. *Cereb Cortex*, *18*(12), 2735-2747.
https://doi.org/10.1093/cercor/bhn041
Douaud, G., Lee, S., Alfaro-Almagro, F., Arthofer, C., Wang, C., Lange, F., Andersson, J. L. R.,
Griffanti, L., Duff, E., Jbabdi, S., Taschler, B., Winkler, A., Nichols, T. E., Collins, R.,
Matthews, P. M., Allen, N., Miller, K. L., & Smith, S. M. (2021). Brain imaging before
and after COVID-19 in UK Biobank. *medRxiv*.
https://doi.org/10.1101/2021.06.11.21258690
Duan, K., Premi, E., Pilotto, A., Cristillo, V., Benussi, A., Libri, I., Giunta, M., Bockholt, H. J., Liu,
J., Campora, R., Pezzini, A., Gasparotti, R., Magoni, M., Padovani, A., & Calhoun, V. D.
(2021). Alterations of frontal-temporal gray matter volume associate with clinical
measures of older adults with COVID-19. *Neurobiol Stress*, *14*, 100326.
https://doi.org/10.1016/j.ynstr.2021.100326
Filippini, N., MacIntosh, B. J., Hough, M. G., Goodwin, G. M., Frisoni, G. B., Smith, S. M.,
Matthews, P. M., Beckmann, C. F., & Mackay, C. E. (2009). Distinct patterns of brain
activity in young carriers of the APOE-epsilon4 allele. *Proc Natl Acad Sci U S A*,
106(17), 7209-7214. https://doi.org/10.1073/pnas.0811879106
Fischer, D., Snider, S. B., Barra, M. E., Sanders, W. R., Rapalino, O., Schaefer, P., Foulkes, A. S.,
Bodien, Y. G., & Edlow, B. L. (2021). Disorders of Consciousness Associated With
COVID-19: A Prospective, Multimodal Study of Recovery and Brain Connectivity.
Neurology. https://doi.org/10.1212/wnl.0000000000013067
Fischer, D., Threlkeld, Z. D., Bodien, Y. G., Kirsch, J. E., Huang, S. Y., Schaefer, P. W., Rapalino,
O., Hochberg, L. R., Rosen, B. R., & Edlow, B. L. (2020). Intact Brain Network Function
in an Unresponsive Patient with COVID-19. *Annals of neurology*, *88*(4), 851-854.
https://doi.org/10.1002/ana.25838
Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E. (2005).
The human brain is intrinsically organized into dynamic, anticorrelated functional
networks. *Proc Natl Acad Sci U S A*, *102*(27), 9673-9678.
https://doi.org/10.1073/pnas.0504136102
Fox, M. D., Snyder, A. Z., Zacks, J. M., & Raichle, M. E. (2006). Coherent spontaneous activity
accounts for trial-to-trial variability in human evoked brain responses. *Nat Neurosci*,
9(1), 23-25. https://doi.org/10.1038/nn1616
Friston, K. J., Williams, S., Howard, R., Frackowiak, R. S., & Turner, R. (1996). Movement-
related effects in fMRI time-series. *Magn Reson Med*, *35*(3), 346-355.
Fu, Z., Tu, Y., Calhoun, V. D., Zhang, Y., Zhao, Q., Chen, J., Meng, Q., Lu, Z., & Hu, L. (2021).
Dynamic functional network connectivity associated with post-traumatic stress

symptoms in COVID-19 survivors. *Neurobiology of Stress*, *15*, 100377.
https://doi.org/https://doi.org/10.1016/j.ynstr.2021.100377
Garrigues, E., Janvier, P., Kherabi, Y., Le Bot, A., Hamon, A., Gouze, H., Doucet, L., Berkani, S.,
Oliosi, E., Mallart, E., Corre, F., Zarrouk, V., Moyer, J. D., Galy, A., Honsel, V., Fantin, B.,
& Nguyen, Y. (2020). Post-discharge persistent symptoms and health-related quality
of life after hospitalization for COVID-19. *J Infect*, *81*(6), e4-e6.
https://doi.org/10.1016/j.jinf.2020.08.029
Gay, C. W., Robinson, M. E., Lai, S., O'Shea, A., Craggs, J. G., Price, D. D., & Staud, R. (2016).
Abnormal Resting-State Functional Connectivity in Patients with Chronic Fatigue
Syndrome: Results of Seed and Data-Driven Analyses. *Brain Connect*, *6*(1), 48-56.
https://doi.org/10.1089/brain.2015.0366
Greicius, M. D., Krasnow, B., Reiss, A. L., & Menon, V. (2003). Functional connectivity in the
resting brain: a network analysis of the default mode hypothesis. *Proc Natl Acad Sci*
U S A, *100*(1), 253-258. https://doi.org/10.1073/pnas.0135058100
Gulko, E., Oleksk, M. L., Gomes, W., Ali, S., Mehta, H., Overby, P., Al-Mufti, F., & Rozenshtein,
A. (2020). MRI Brain Findings in 126 Patients with COVID-19: Initial Observations
from a Descriptive Literature Review. *American Journal of Neuroradiology*.
https://doi.org/10.3174/ajnr.A6805
Hafiz, R., Gandhi, T. K., Mishra, S., Prasad, A., Mahajan, V., Di, X., Natelson, B. H., & Biswal, B.
B. (2021). Higher Limbic and Basal Ganglia volumes in surviving COVID-negative
patients and the relations to fatigue. *medRxiv*, 2021.2011.2023.21266761.
https://doi.org/10.1101/2021.11.23.21266761
Ismail, I. I., & Gad, K. A. (2021). Absent Blood Oxygen Level–Dependent Functional Magnetic
Resonance Imaging Activation of the Orbitofrontal Cortex in a Patient With
Persistent Cacosmia and Cacogeusia After COVID-19 Infection. *JAMA Neurology*.
https://doi.org/10.1001/jamaneurol.2021.0009
Kalcher, K., Huf, W., Boubela, R. N., Filzmoser, P., Pezawas, L., Biswal, B., Kasper, S., Moser,
E., & Windischberger, C. (2012). Fully exploratory network independent component
analysis of the 1000 functional connectomes database. *Front Hum Neurosci*, *6*, 301.
https://doi.org/10.3389/fnhum.2012.00301
Kandemirli, S. G., Dogan, L., Sarikaya, Z. T., Kara, S., Akinci, C., Kaya, D., Kaya, Y., Yildirim, D.,
Tuzuner, F., Yildirim, M. S., Ozluk, E., Gucyetmez, B., Karaarslan, E., Koyluoglu, I.,
Demirel Kaya, H. S., Mammadov, O., Kisa Ozdemir, I., Afsar, N., Citci Yalcinkaya, B., . . .
Kocer, N. (2020). Brain MRI Findings in Patients in the Intensive Care Unit with
COVID-19 Infection. *Radiology*, *297*(1), E232-e235.
https://doi.org/10.1148/radiol.2020201697
Keller, E., Brandi, G., Winklhofer, S., Imbach, L. L., Kirschenbaum, D., Frontzek, K., Steiger, P.,
Dietler, S., Haeberlin, M., Willms, J., Porta, F., Waeckerlin, A., Huber, M., Abela, I. A.,
Lutterotti, A., Stippich, C., Globas, C., Varga, Z., & Jelcic, I. (2020). Large and Small
Cerebral Vessel Involvement in Severe COVID-19: Detailed Clinical Workup of a Case
Series. *Stroke*, *51*(12), 3719-3722. https://doi.org/10.1161/strokeaha.120.031224
Kim, B. H., Shin, Y. B., Kyeong, S., Lee, S. K., & Kim, J. J. (2018). Disrupted salience processing
involved in motivational deficits for real-life activities in patients with
schizophrenia. *Schizophr Res*, *197*, 407-413.
https://doi.org/10.1016/j.schres.2018.01.019

Kremer, S., Lersy, F., Anheim, M., Merdji, H., Schenck, M., Oesterlé, H., Bolognini, F., Messie, J., Khalil, A., Gaudemer, A., Carré, S., Alleg, M., Lecocq, C., Schmitt, E., Anxionnat, R., Zhu, F., Jager, L., Nesser, P., Mba, Y. T., . . . Cotton, F. (2020). Neurologic and neuroimaging findings in patients with COVID-19: A retrospective multicenter study. *Neurology*, *95*(13), e1868-e1882. https://doi.org/10.1212/wnl.0000000000010112

Liu, P., Yang, W., Zhuang, K., Wei, D., Yu, R., Huang, X., & Qiu, J. (2021). The functional connectome predicts feeling of stress on regular days and during the COVID-19 pandemic. *Neurobiol Stress*, *14*, 100285. https://doi.org/10.1016/j.ynstr.2020.100285

Logue, J. K., Franko, N. M., McCulloch, D. J., McDonald, D., Magedson, A., Wolf, C. R., & Chu, H. Y. (2021). Sequelae in Adults at 6 Months After COVID-19 Infection. *JAMA Netw Open*, *4*(2), e210830. https://doi.org/10.1001/jamanetworkopen.2021.0830

Lu, Y., Li, X., Geng, D., Mei, N., Wu, P.-Y., Huang, C.-C., Jia, T., Zhao, Y., Wang, D., Xiao, A., & Yin, B. (2020). Cerebral Micro-Structural Changes in COVID-19 Patients – An MRI-based 3-month Follow-up Study. *EClinicalMedicine*, *25*. https://doi.org/10.1016/j.eclinm.2020.100484

Mattioli, P., Pardini, M., Famà, F., Girtler, N., Brugnolo, A., Orso, B., Meli, R., Filippi, L., Grisanti, S., Massa, F., Bauckneht, M., Miceli, A., Terzaghi, M., Morbelli, S., Nobili, F., & Arnaldi, D. (2021). Cuneus/precuneus as a central hub for brain functional connectivity of mild cognitive impairment in idiopathic REM sleep behavior patients. *Eur J Nucl Med Mol Imaging*, *48*(9), 2834-2845. https://doi.org/10.1007/s00259-021-05205-6

McKeown, M. J., Makeig, S., Brown, G. G., Jung, T. P., Kindermann, S. S., Bell, A. J., & Sejnowski, T. J. (1998). Analysis of fMRI data by blind separation into independent spatial components. *Hum Brain Mapp*, *6*(3), 160-188. https://doi.org/10.1002/(SICI)1097-0193(1998)6:3<160::AID-HBM5>3.0.CO;2-1

Meier, T. B., Wildenberg, J. C., Liu, J., Chen, J., Calhoun, V. D., Biswal, B. B., Meyerand, M. E., Birn, R. M., & Prabhakaran, V. (2012). Parallel ICA identifies sub-components of resting state networks that covary with behavioral indices. *Front Hum Neurosci*, *6*, 281. https://doi.org/10.3389/fnhum.2012.00281

Miller, A. H., Jones, J. F., Drake, D. F., Tian, H., Unger, E. R., & Pagnoni, G. (2014). Decreased basal ganglia activation in subjects with chronic fatigue syndrome: association with symptoms of fatigue. *PLoS One*, *9*(5), e98156. https://doi.org/10.1371/journal.pone.0098156

Nicholson, P., Alshafai, L., & Krings, T. (2020). Neuroimaging Findings in Patients with COVID-19. *American Journal of Neuroradiology*, *41*(8), 1380-1383. https://doi.org/10.3174/ajnr.A6630

Park, I. S., Lee, N. J., & Rhyu, I. J. (2018). Roles of the Declive, Folium, and Tuber Cerebellar Vermian Lobules in Sportspeople. *Journal of clinical neurology (Seoul, Korea)*, *14*(1), 1-7. https://doi.org/10.3988/jcn.2018.14.1.1

Paterson, R. W., Brown, R. L., Benjamin, L., Nortley, R., Wiethoff, S., Bharucha, T., Jayaseelan, D. L., Kumar, G., Raftopoulos, R. E., Zambreanu, L., Vivekanandam, V., Khoo, A., Geraldes, R., Chinthapalli, K., Boyd, E., Tuzlali, H., Price, G., Christofi, G., Morrow, J., . . . Zandi, M. S. (2020). The emerging spectrum of COVID-19 neurology: clinical, radiological and laboratory findings. *Brain*, *143*(10), 3104-3120. https://doi.org/10.1093/brain/awaa240

547 Pellicano, C., Gallo, A., Li, X., Ikonomidou, V. N., Evangelou, I. E., Ohayon, J. M., Stern, S. K.,
548   Ehrmantraut, M., Cantor, F., McFarland, H. F., & Bagnato, F. (2010). Relationship of
549   Cortical Atrophy to Fatigue in Patients With Multiple Sclerosis. *Archives of*
550   *Neurology*, *67*(4), 447-453. https://doi.org/10.1001/archneurol.2010.48
551 Peluso, M. J., Kelly, J. D., Lu, S., Goldberg, S. A., Davidson, M. C., Mathur, S., Durstenfeld, M. S.,
552   Spinelli, M. A., Hoh, R., Tai, V., Fehrman, E. A., Torres, L., Hernandez, Y., Williams, M.
553   C., Arreguin, M. I., Bautista, J. A., Ngo, L. H., Deswal, M., Munter, S. E., . . . Martin, J. N.
554   (2021). Rapid implementation of a cohort for the study of post-acute sequelae of
555   SARS-CoV-2 infection/COVID-19. *medRxiv*, 2021.2003.2011.21252311.
556   https://doi.org/10.1101/2021.03.11.21252311
557 Perica, M. I., Ravindranath, O., Calabro, F. J., Foran, W., & Luna, B. (2021). Hippocampal-
558   Prefrontal Connectivity Prior to the COVID-19 Pandemic Predicts Stress Reactivity.
559   *Biol Psychiatry Glob Open Sci*, *1*(4), 283-290.
560   https://doi.org/10.1016/j.bpsgos.2021.06.010
561 Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012). Spurious
562   but systematic correlations in functional connectivity MRI networks arise from
563   subject motion. *Neuroimage*, *59*(3), 2142-2154.
564   https://doi.org/10.1016/j.neuroimage.2011.10.018
565 Qin, Y., Wu, J., Chen, T., Li, J., Zhang, G., Wu, D., Zhou, Y., Zheng, N., Cai, A., Ning, Q., Manyande,
566   A., Xu, F., Wang, J., & Zhu, W. (2021). Long-term microstructure and cerebral blood
567   flow changes in patients recovered from COVID-19 without neurological
568   manifestations. *J Clin Invest*, *131*(8). https://doi.org/10.1172/jci147329
569 Raichle, M. E., & Mintun, M. A. (2006). Brain work and brain imaging. *Annu Rev Neurosci*, *29*,
570   449-476. https://doi.org/10.1146/annurev.neuro.29.051605.112819
571 RStudio. (2021). *RStudio Team (2021). RStudio: Integrated Development for R.*
572   *http://www.rstudio.com/.* In RStudio.
573 Shirer, W. R., Ryali, S., Rykhlevskaia, E., Menon, V., & Greicius, M. D. (2012). Decoding
574   subject-driven cognitive states with whole-brain connectivity patterns. *Cereb Cortex*,
575   *22*(1), 158-165. https://doi.org/10.1093/cercor/bhr099
576 Shulman, R. G., Rothman, D. L., Behar, K. L., & Hyder, F. (2004). Energetic basis of brain
577   activity: implications for neuroimaging. *Trends Neurosci*, *27*(8), 489-495.
578   https://doi.org/10.1016/j.tins.2004.06.005
579 Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., Filippini, N.,
580   Watkins, K. E., Toro, R., Laird, A. R., & Beckmann, C. F. (2009). Correspondence of the
581   brain's functional architecture during activation and rest. *Proceedings of the*
582   *National Academy of Sciences*, *106*(31), 13040-13045.
583   https://doi.org/10.1073/pnas.0905267106
584 Tabacof, L., Tosto-Mancuso, J., Wood, J., Cortes, M., Kontorovich, A., McCarthy, D., Rizk, D.,
585   Mohammadi, N., Breyman, E., Nasr, L., Kellner, C., & Putrino, D. (2020). Post-acute
586   COVID-19 syndrome negatively impacts health and wellbeing despite less severe
587   acute infection. *medRxiv*, 2020.2011.2004.20226126.
588   https://doi.org/10.1101/2020.11.04.20226126
589 Taylor, P. A., & Saad, Z. S. (2013). FATCAT: (an efficient) Functional and Tractographic
590   Connectivity Analysis Toolbox. *Brain Connect*, *3*(5), 523-535.
591   https://doi.org/10.1089/brain.2013.0154

Tu, Y., Zhang, Y., Li, Y., Zhao, Q., Bi, Y., Lu, X., Kong, Y., Wang, L., Lu, Z., & Hu, L. (2021). Post-
traumatic stress symptoms in COVID-19 survivors: a self-report and brain imaging
follow-up study. *Molecular Psychiatry*. https://doi.org/10.1038/s41380-021-01223-
w
van Schouwenburg, M. R., den Ouden, H. E., & Cools, R. (2015). Selective attentional
enhancement and inhibition of fronto-posterior connectivity by the basal ganglia
during attention switching. *Cereb Cortex*, *25*(6), 1527-1534.
https://doi.org/10.1093/cercor/bht345
Wortinger, L. A., Glenne Øie, M., Endestad, T., & Bruun Wyller, V. (2017). Altered right
anterior insular connectivity and loss of associated functions in adolescent chronic
fatigue syndrome. *PLoS One*, *12*(9), e0184325.
https://doi.org/10.1371/journal.pone.0184325
Xiao, M., Chen, X., Yi, H., Luo, Y., Yan, Q., Feng, T., He, Q., Lei, X., Qiu, J., & Chen, H. (2021).
Stronger functional network connectivity and social support buffer against negative
affect during the COVID-19 outbreak and after the pandemic peak. *Neurobiology of
Stress*, *15*, 100418-100418. https://doi.org/10.1016/j.ynstr.2021.100418
Yan, C.-G., Wang, X.-D., Zuo, X.-N., & Zang, Y.-F. (2016). DPABI: Data Processing & Analysis
for (Resting-State) Brain Imaging. *Neuroinformatics*, *14*(3), 339-351.
https://doi.org/10.1007/s12021-016-9299-4
Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman,
J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fischl, B., Liu, H., & Buckner, R. L. (2011).
The organization of the human cerebral cortex estimated by intrinsic functional
connectivity. *J Neurophysiol*, *106*(3), 1125-1165.
https://doi.org/10.1152/jn.00338.2011
Zhang, S., Cui, J., Zhang, Z., Wang, Y., Liu, R., Chen, X., Feng, Y., Zhou, J., Zhou, Y., & Wang, G.
(2022). Functional connectivity of amygdala subregions predicts vulnerability to
depression following the COVID-19 pandemic. *J Affect Disord*, *297*, 421-429.
https://doi.org/10.1016/j.jad.2021.09.107
620
621

Figure 1. Twenty-two Resting State Networks (RSNs) identified from group ICA using 'melodic'. Abbreviated names of each network are shown at the bottom of each image. Three orthogonal slices are shown for each network along with a volume rendered image to show depth and three-dimensional view of the RSNs. Statistical estimates (Z-scores) are embedded into a colorbar at the bottom-right. PMV = Primary Visual Network, LV = Lateral Visual, OCP = Occipital Pole, MV = Medial Visual, PRN = Precuneus Network, DAN = Dorsal Attention, VDMN = Ventral Default Mode Network (DMN), PDMN = Posterior DMN, RFP = Right Fronto Parietal, LFP = Left Fronto Parietal, AUD = Auditory, TPJN = Temporo-Parietal Junction Network, LANG = Language Network, EXEC = Executive Control Network, INS = Insular Network, MSMN = Medial Sensory-Motor Network (SMN), VSMN = Ventral SMN, SSNR = Somatosensory Network - Right, SMNL = Somatosensory Network - Left, , BGN = Basal Ganglia Network, SCRB = Superior Cerebellar Network, PCRB = Posterior Cerebellar Network.

Figure 2. ΔFC | Functional Connectivity differences between COVID survivors and healthy controls. [top row] (A) COVID > HC: Enhanced FC in COVID compared to HCs observed in the *Basal Ganglia Network (BGN)* network**.** Three orthogonal slices (left) along with a cut-to-depth volume rendered image to show the effects in the *right Calc, Cu* and *LiG*. The colorbar represents t – score values. Cluster information include - cluster peak: *[9 -84 6],* | cluster extent threshold, $k_E = 69$ | cluster size = *69 voxels*. **(B) COVID > HC:** Enhanced FC in COVID compared to HCs observed in the *Precuneus (PRC)* network, demonstrating a significant difference in FC in the *bilateral SPL and PCu* regions. Cluster information include - cluster peak: *[21 -57 54],* | cluster extent threshold, $k_E = 90$ | cluster size = *90 voxels***. (C) HC > COVID:** Enhanced FC in HCs compared to COVID observed in the *Language (LANG)* network demonstrating significant difference in several vermal layers of the *Cerebellum*. Cluster information include - cluster peak: *[9 -63 -24],* | cluster extent threshold, $k_E = 57$ | cluster size = *57 voxels*. [**bottom row**] Corresponding group level ICA networks from which FC differences are shown on the top row. The colorbar represents z-scores.

Figure 3. FC corr. Fatigue | COVID: Negative correlation of FC with self-reported fatigue scores in COVID individuals. (left) For the *PRN* network, three orthogonal slices (left) along with a cut-to-depth volume rendered image showing regions from the *Superior Parietal* and *Occipital* Gyri that demonstrated significantly negative correlation with fatigue. The colorbar represents t-score values. **(right)** The graph shows the linear relationship of the average FC within the significant cluster and self-reported fatigue scores from COVID individuals. The x-axis represents the average FC (z-scores) from the cluster and the y-axis represents the fatigue scores. The shaded gray area represents the 95% confidence interval. The red line represents the least squares regression line of best fit. Cluster information include - cluster peak: *[-21 -75 36], |* cluster extent threshold, $k_E = 58$ and cluster size = *58 voxels.*

ΔFC	RSN Name	Cl. No.	Anatomical Locations	Cl. Ext.	Cl. Size	Peak MNI Coordinates			Peak T, p_FWE Values
						X	Y	Z	
COVID > HC	*BGN*	1	*Right – Calcarine Cortex (Calc)*	69	69	9	-84	6	4.46, 0.004
			Right – Cuneus (Cun)						
			Right – Lingual Gyrus (LiG)						
	PRN	1	*Right – Superior Parietal Lobule (SPL)*	90	90	21	-57	54	4.22, 0.001
			Right – Precuneus (PCu)						
			Left – Superior Parietal Lobule (SPL)						
			Left – Precuneus (PCu)						
HC > COVID	*LANG*	1	*Right – Cerebellar Exterior (CExt.)*	57	57	9	-63	-24	3.56, 0.019
			Right – Cerebellar Vermal Lobules I-V						
			Right – Cerebellar Vermal Lobules VI-VII						

Table 1. List of spatial regions from significant clusters obtained from the contrast – COVID > HC. The regions from three RSNs – *BGN*, *PRN* and *LANG* which demonstrated significant differences are presented with peak MNI coordinates (X Y Z) and corresponding peak t-score values for each cluster. Keys – ΔFC = Direction of change in Functional Connectivity; Cl. = Cluster; Cl. No. = Cluster Number; Cl. Ext. = Cluster Extent Threshold; Cl. Size = Cluster Size; T = peak t-score; p_{FWE} = corrected p-value.

Corr (FC, ftg.)	RSN Name	Cl. No.	Anatomical Locations	Cl. Ext.	Cl. Size	Peak MNI Coordinates			Peak T, p_{FWE} Values
						X	Y	Z	
COVID	*PRN*	1	*Left – Superior Parietal Lobule (SPL)*	58	58	-21	-75	36	5.31, 0.01
			Left – Superior Occipital Gyrus (SOG)						
			Left – Angular Gyrus (AnG)						
			Left – Precuneus (PCu)						

Table 2. List of spatial regions from clusters showing significant correlation with self-reported fatigue among COVID individuals. The regions from *PRN* which demonstrated significant correlation are presented with peak MNI coordinates (X Y Z) and corresponding peak t-score values for each cluster. Keys – FC = Functional Connectivity; ftg. = Fatigue Scores, Cl. = Cluster; Cl. No. = Cluster Number; Cl. Ext. = Cluster Extent Threshold; Cl. Size = Cluster Size; T = peak t-score; p_{FWE} = corrected p-value.

643