Activation of a bacterial flow sensor by hypochlorous acid from stimulated neutrophils

Ilona P. Foik[1,†], Lauren A. Urban[2-4,†], Andy P. Huang[5], Timothy L. Downing[2-4,6,*], Albert Siryaporn[1,5,*]

[1]Department of Physics & Astronomy, University of California Irvine; Irvine, California, USA.

[2]Department of Microbiology & Molecular Genetics, University of California Irvine; Irvine, California, USA.

[3]NSF-Simons Center for Multiscale Cell Fate, University of California Irvine; Irvine, CA, USA.

[4]UCI Edwards Lifesciences Foundation Cardiovascular Innovation and Research Center (CIRC), University of California Irvine; Irvine, CA, USA.

[5]Department of Molecular Biology & Biochemistry, University of California Irvine; Irvine, California, USA.

[6]Department of Biomedical Engineering, University of California Irvine, Irvine; California, USA.

[†]These authors contributed equally.

[*]To which correspondence should be sent:
T. Downing: tim.downing@uci.edu
A. Siryaporn: asirya@uci.edu

ABSTRACT

Bacteria sense environmental stressors and activate responses to improve their survival in harsh growth conditions. Neutrophils respond to the presence of bacteria by producing oxidative antibacterial species including hypochlorous acid (HOCl). However, the extent that bacteria detect activated neutrophils or HOCl has not been known. Here, we report that the opportunistic bacterial pathogen *Pseudomonas aeruginosa* responds to activated neutrophils by activating the *fro* system, which regulates the expression of antioxidative factors. We show that this response is specific to HOCl and that other oxidative factors including H_2O_2, do not trigger a *fro* response. The *fro* system has been previously shown to detect flow that is present in host vasculature, such as in animal circulatory systems. Our data thus suggest a model in which *fro* serves as an early host detection system in *P. aeruginosa* that improves its survival against neutrophil-mediated defenses, which could promote colonization in human tissue and increase pathogenicity.

One Sentence Summary: *P. aeruginosa* primes its defenses against neutrophils by sensing flow and hypochlorous acid via the *fro* system.

INTRODUCTION

P. aeruginosa is a Gram-negative bacterium that is a major cause of acute hospital-acquired infections and a significant threat for antibiotic-resistant infections *(1)*. To survive in a broad range of conditions, this bacterium has evolved signaling pathways that relay information about changes in nutrients, chemical gradients, and mechanical forces such as those generated by flow, and stimulates responses that promote survival *(2, 3)*. Many of these signaling pathways improve the bacterium's survival in hosts in particular. Understanding how such environmental-sensing pathways detect and respond to host cues is critical for improving treatments of bacterial infections.

Mammalian host immune systems respond to the presence of bacterial pathogens by recruiting immune cells to sites of infection. Neutrophils are among the first and most abundant immune cells recruited to inflamed or damaged tissues *(4)*. Neutrophils are stimulated into an activated state by the presence of the bacterial components, including the membrane constituent lipopolysaccharide. Activated neutrophils kill bacteria through a number of mechanisms, including producing respiratory burst products such as reactive oxygen species (ROS) and reactive chlorine species (RCS) *(5, 6)*. While the toxic effects of ROS on bacteria are understood, far less is known about how bacteria are affected by RCS *(7, 8)*. Hypochlorous acid (HOCl) is a major neutrophil RCS that has strong bactericidal activity towards bacteria *(9, 10)*. HOCl is formed through the activity of myeloperoxidase with H_2O_2 and chlorine ions *(5, 6, 11)*. *P. aeruginosa* is a potent activator of neutrophil respiratory bursts *(12)*. However, whether and how *P. aeruginosa* detects and responds to neutrophil respiratory bursts is not understood.

Recently, it was reported that *P. aeruginosa* detects the flow of fluids via the flow-regulated operon *fro (3)*. Flow is ubiquitous in the lung and in the circulatory system, which

neutrophils and other immune cells must travel through to reach infected sites. The *fro* system

could thus enable *P. aeruginosa* to detect the presence of flow-associated host environments and

immune system components. *P. aeruginosa* induces *fro* operon expression during infections in

humans *(3, 14)* (Fig. 1A).The operon consists of the genes *froABCD* (*PA14_21570*,

PA14_21580, PA14_21590, PA14_21600). FroB is predicted to localize in cytoplasm while FroC

and FroD are predicted to localize to the inner membrane *(15)*. The *fro* operon is regulated by

two upstream genes, *froR* (*PA14_21550)* and *froI* (*PA14_21560*), which are sigma and anti-

sigma factors, respectively *(16)*. Overexpression of *froR* increases *fro* expression whereas

overexpression of *froI* eliminates *fro* induction *(3)*. Flow activates FroR-dependent transcription

of *froABCD* through an unknown mechanism.

We investigated whether the *fro* system directly responds to components of the

circulatory system. Here, we show that the expression of the *fro* system is upregulated by

activated neutrophils. We determine that HOCl is a critical component of the neutrophil

activation that is responsible for the *fro* upregulation and that its upregulation reduces the

bactericidal effects of HOCl. These data suggest a novel mechanism by which a bacterial defense

system is primed to detect host immune system activity through both flow and the presence of

the immune cell effector HOCl, and responds by producing protective antioxidant proteins.

Fig. 1 Activation of *fro* transcription by flow and during infection of human tissue. (A)
RNA-seq heatmap of *P. aeruginosa* genes that are significantly upregulated at least 3-fold both
by flow and during infections in human wounds. RNA-seq datasets are from *(3, 14)*. **(B)** Top:
schematic of a *P. aeruginosa* fluorescent reporter strain that contains a transcriptional fusion of
yfp to *fro* and constitutively expresses *mCherry*. Representative phase contrast, YFP, and
mCherry images the reporter under flow (right) or non-flow (left) conditions for 30 minutes. **(C)**
Representative phase contrast, YFP, mCherry, and merged images of the *P. aeruginosa* reporter
strain co-incubated with neutrophils, macrophages, fibroblasts, or host cell-free media for 30
minutes. Dotted lines outline the boundaries of the host cells. White arrows in merged images
indicate *P. aeruginosa* with the high ratios of YFP/mCherry intensities. Images are
representative of three independent repetitions. Scale bars represent 5 μm.

RESULTS

Host cells stimulate *fro* expression in *P. aeruginosa* under non-flow conditions

We investigated if the *fro* operon is activated directly by immune cells. To quantify *fro*

expression, we used a *P. aeruginosa* strain that co-expresses the yellow fluorescent protein

(YFP) under the transcriptional control of the *fro* promoter and mCherry under the control of a

constitutive promoter (Fig. 1B). Neutrophils and macrophages are among the first immune cells

that are recruited to sites of infection. We co-incubated *P. aeruginosa* with these cells for 30

minutes to determine whether they would elicit a response in the *fro* operon. Experiments were

performed in the absence of flow in order to deconvolve the effects of flow from potential

activation of *fro* by host cells. Relative to *P. aeruginosa* that were incubated in host cell-free

media, *P. aeruginosa* in close proximity to neutrophils or that were engulfed in macrophages had

higher levels of *fro* expression (Fig. 1C). Furthermore, increases in *fro* expression were observed

in some *P. aeruginosa* that were internalized in fibroblasts (Fig. 1C). These data suggest direct

interaction or close proximity with host hosts stimulates *fro* expression in *P. aeruginosa*.

***fro* expression is induced by activated neutrophils**

90
We focused our efforts on understanding how neutrophils activate *fro* expression, as

neutrophils are one of the first immune cells that are recruited to bacterial infection sites.

P. aeruginosa triggers respiratory bursts in neutrophils *(12)*, which generates ROS and RCS. We

hypothesized that the observed *fro* activation could be a response to respiratory burst products.

To test this, we determined whether molecules released by activated neutrophils could induce *fro*

expression. We isolated cell-free conditioned medium from neutrophils that were stimulated to

induce respiratory bursts using 100 ng/mL phorbol myristate acetate (PMA) in neutrophil

isolation medium (PBGT) *(17)*. *P. aeruginosa* were cultured for 3 hours in the conditioned

medium (Fig. 2A). The expression of *fro* was quantified by computing the ratio of YFP to

mCherry fluorescence in individual *P. aeruginosa* and averaging over at least one hundred

P. aeruginosa. Conditioned medium from PMA-stimulated neutrophils increased the expression

of *fro* by 30-fold compared to medium from unstimulated neutrophils (Fig. 2B). The

upregulation in *fro* was not due to PMA, as the addition of PMA without neutrophils had no

effect on *fro* expression (Fig. 2B). The conditioned media from unstimulated neutrophils had no

significant effect on *fro* expression compared to PBGT medium alone (Fig. 2B).

Figure 2

Fig. 2 Conditioned medium from stimulated neutrophils induces *fro* expression. (A)
Representative phase-contrast and fluorescence images and **(B)** quantification of *fro* expression
levels of *P. aeruginosa* that were cultured for 3 hours in PGBT medium with or without 100
ng/mL of PMA, conditioned medium from unstimulated neutrophils, or conditioned medium
from neutrophils that were stimulated with 100 ng/mL of PMA. Bars indicate the mean and error
bars represent the standard error of the mean (SEM) of three independent experiments. P-values
were calculated using a two-tailed, unequal variance t-test. ns indicates $p > 0.05$. Scale bars
represent 5 μm.

fro expression is induced by HOCl but not by H_2O_2 or HNO_3

We assessed the ability of specific respiratory burst products to drive _fro_ activation.

Activated neutrophils generate multiple ROS and RCS oxidants, which have an inhibitory effect

on bacterial growth _(5–7, 9, 10)_. We hypothesized that these oxidants induce _fro_ expression and

tested the impact of these species on _fro_ expression at near-inhibitory concentrations (Fig. 3A).

We established that 1 μM of NaOCl or 8 μM of H_2O_2 were at the threshold of growth inhibition

for _P. aeruginosa_ (Fig. 3B), and subsequently measured _fro_ expression in cultures up to these

concentration thresholds. H_2O_2 in a range of concentrations below, at, or above 8 μM had no

significant impact on _fro_ expression (Fig. 3C-D). In contrast, 1 μM NaOCl increased _fro_

expression by up 78-fold compared to untreated cultures. This activation is not due to sodium or

chloride ions alone, as neither NaCl at the same concentration or HCl up to the highest

concentration without changing pH of the minimal medium did not induce _fro_ expression (Fig.

3D). In addition, the _fro_ response to NaOCl is not due to basic pH alone, as treatment with 6 mM

NaOH, which adjusted the medium to the same pH as 1 uM NaOCl (pH 7.1), had no significant

impact on _fro_ activation (Fig. 3D). NaOCl-induced _fro_ expression occurred at concentrations

below those associated with growth inhibition (Fig. 3B-D). The activation of _fro_ by NaOCl is

consistent with a recent transcriptional study that identified _fro_ as a gene that is upregulated by

HOCl _(18)_. We note that a higher concentration of 2 μM NaOCl did not increase _fro_ expression.

However, this concentration caused a significant decrease in cell size (Fig. S1 in Supplementary

Materials), indicative of cell stress that may have interfered with transcription.

Figure 3

Fig. 3 The expression of *fro* is induced by HOCl but not by H_2O_2 or HNO_3. (A) Schematic of
model indicating that PMA-stimulated neutrophils produce respiratory burst products that could
induce *fro* expression. **(B)** Growth profiles, measured by optical density (OD_{600}), of
P. aeruginosa that were treated with sublethal and lethal concentrations of H_2O_2, NaOCl, HNO_3,
or untreated. Data points represent the mean and error bars indicate the SEM of at least two
independent experiments. **(C)** The expression of *fro* in *P. aeruginosa* that were treated for 3
hours with sublethal concentrations of H_2O_2, NaOCl, HNO_3, or were untreated. **(D)** The
expression of *fro* in *P. aeruginosa* that were treated for 3 hours with 8 µM H_2O_2, 1 µM NaOCl,
25 mM HNO_3, 6 mM NaOH, 1 µM NaCl, 200 µM HCl, or untreated. Bars represent the mean
and error bars indicate the SEM of at least two independent experiments. Boxplots indicate the
mean (black line) and interquartile range (yellow) of three independent experiments. P-values
were calculated using a two-tailed, unequal variance t-test. ns indicates $p > 0.05$.
149

We further assessed whether *fro* expression could be activated by the strong oxidant

HNO_3. The effects of this oxidizer were measured up to inhibitory concentrations (Fig. 3B-D).

However, no impact on *fro* expression was detected. These results suggest that *fro* responds

specifically to RCS stress from HOCl and supports the hypothesis that HOCl produced by

stimulated neutrophils activates *fro* expression.

Antioxidant supplements inhibit *fro* activation

HOCl oxidizes amine and sulfur groups within amino acids *(19, 22)*. We posited that

blocking the oxidative activity of HOCl could inhibit *fro* activation. Sulfur-containing amino

acids (e.g., methionine and cysteine) and beta-mercaptoethanol (βME) are antioxidants and

major targets for HOCl-mediated oxidation *(21, 23)*. We hypothesized that supplementing 1 µM

NaOCl treatments with methionine, cysteine, or βME reduces *fro* expression. Indeed, we

observed that supplementing these antioxidants at equal or lower concentrations as NaOCl (1 µM

methionine, 0.1 µM cysteine, or 1 µM βME) significantly reduced the NaOCl-induced *fro*

expression (Fig. 4A, Fig. S2 in Supplementary Materials) indicating that the oxidative activity of

NaOCl activates *fro*. Concentrations of antioxidants that were higher than 1 µM also inhibited

fro expression (Fig. S2 in Supplementary Materials). Treatments of cysteine, methionine, and

βME alone had no significant impact on *fro* expression (Fig. S3 in Supplementary Materials).

Based on these data, we hypothesized that the activation of *fro* expression by PMA-

stimulated neutrophils is suppressed by antioxidants. Indeed, we found that supplementing

conditioned media from PMA-stimulated neutrophils with 100 µM of methionine inhibited *fro*

activation (Fig. 4B) comparable to that observed in PBGT media alone and in unstimulated

neutrophils (Figs. 2B and 4B). Together, these results suggest that stimulated neutrophils activate

fro expression through the oxidative activity of HOCl.

Fig. 4 Antioxidants inhibit *fro* activation that is induced by HOCl and conditioned media
from PMA-stimulated neutrophils. (A) The expression of *fro* in *P. aeruginosa* that are untreated,
treated with 1 µM NaOCl, or treated with 1 µM NaOCl and supplemented with 1 µM methionine,
0.1 µM cysteine, or 1 µM beta-mercaptoethanol (BME), for 3 hours. **(B)** Quantification of *fro*
expression levels in *P. aeruginosa* that were cultured in PBGT medium or in PBGT medium from
PMA-stimulated neutrophils (replotted from Fig. 2B) that was untreated or supplemented with

100 µM methionine. Bars represent the mean and error bars indicate the SEM of three independent
experiments. P-values were calculated using a two-tailed, unequal variance t-test. ns indicates
p > 0.05.
185

fro enhances _P. aeruginosa_ survival against HOCl

We investigated whether _fro_ activation has a beneficial impact on _P. aeruginosa_ survival

in response to HOCl-induced stress. We measured the growth of wild-type _P. aeruginosa_ and the

mutant strain Δ_froR_, which lacks the sigma factor that is required for _fro_ activation. In the

absence of NaOCl, both strains grew comparably. Treatment with 16 µM NaOCl significantly

inhibited the growth of the wild-type strain (Fig. 5A). This treatment resulted in even greater

growth inhibition in the Δ_froR_ strain. This result suggests that activation of the _fro_ operon

increases _P. aeruginosa_ survival against oxidative stress induced by HOCl.

We assessed the transcriptional pathways induced by _fro_ activation to understand how _fro_

activation promotes the survival of _P. aeruginosa_ during oxidative stress. We performed a Gene

Ontology (GO) analysis of RNA-seq data from the wild-type strain and Δ_froR_ mutant _(3)_. This

analysis revealed that 128 genes with oxidoreductase activity were upregulated by at least 2-fold

in the wild-type strain compared to the Δ_froR_ mutant (Supplemental Spreadsheet 1 in

Supplementary Materials). The most upregulated genes included _ahpC_, _ahpF_, and _katB_, which

encode proteins with oxidoreductase or antioxidant activity (Fig. 5B and Supplemental

Spreadsheet 2 in Supplementary Materials). These results suggest that _fro_ activation promotes _P._

aeruginosa survival against HOCl stress by activating transcriptional responses that mitigate

oxidative damage.

Figure 5

Fig. 5 FroR improves *P. aeruginosa* survival against HOCl stress and regulates the transcription of genes involved in antioxidant defense. (A) Growth profiles of wild-type (WT) *P. aeruginosa* and Δ*froR* mutants that were treated with 16 µM NaOCl or untreated. Data points represent the mean and the error bars indicate the SEM of three independent experiments. **(B)** Volcano plot indicating the fold-changes in transcripts and p-values of the wild-type (WT) strain relative to the Δ*froR* mutant using RNA-seq dataset from *(3)*. The colors reflect genes that are downregulated (blue), upregulated (red), or unchanged (grey) in WT strains compared to the Δ*froR* mutant. False discovery rate < 0.05.

Discussion

Neutrophils are activated by the presence of bacteria during the infection of host tissues. In this activated state, neutrophils produce respiratory bursts that contain high concentrations of HOCl. Whether and how bacteria respond to activated neutrophils and the RCS that the neutrophils produce has not been clear. In this study, we have demonstrated that *P. aeruginosa* responds to activated neutrophils by upregulating the transcriptional expression of the *fro* operon, which relieves HOCl-induced stress. Our data suggest that *fro* activation could function as a bacterial defense mechanism against a neutrophil attack mechanism. The *fro* response may

thus improve *P. aeruginosa* survival against neutrophils and enhance the colonization of human

tissues during infection.

One of the striking features of the *fro* system is that it is finely tuned to activate at sub-

lethal concentrations (1-2 µM) of NaOCl. Activated neutrophils produce HOCl as high as 50 µM

HOCl *(24)*. The activation of *fro* at relatively low concentrations suggests that the system

functions as a sensitive early detector of HOCl. In host environments, *fro* may activate in

environments where HOCl is present at low concentrations, such as during early stages of

neutrophil stimulation or when *P. aeruginosa* is spatially distant from activated neutrophils. The

early activation of *fro* could benefit *P. aeruginosa* by triggering a protective mechanism that

relieves the impact of HOCl stress before HOCl reaches lethal concentrations. This mechanism

would ensure that the system is already activated when *P. aeruginosa* encounters neutrophils.

Indeed, we observed that *fro* expression is strongly upregulated in *P. aeruginosa* that are in close

proximity to activated neutrophils (Fig. 1C). Additional evidence supports the function of *fro* as

a defensive early detection mechanism. The transcription of *fro* is partially activated by moving

fluids *(3)* that generate wall shear stresses in animal circulatory systems. Neutrophils move in the

fluidic networks of the circulatory system in response to tissue injury. *P. aeruginosa* that are

present in the circulatory system could prime its defenses via flowing sensing by *fro* against

potential HOCl produced by recruited neutrophils.

How does *fro* relieve HOCl stress in bacteria? The *fro* operon activates over 100 genes,

including *ahpC*, *ahpF*, and *katB,* that encode proteins that reduce the activity of ROS and are

critical for bacterial survival against oxidative stress (Fig. 5B) *(25, 26)*. While the effect of these

antioxidant proteins on RCS activity is unclear, our data suggest that *fro* activation mitigates the

toxic effects of HOCl (Fig. 5A). In addition, our results indicate that the presence of antioxidant

agents, including methionine, cysteine, or BME, reduces the activity of HOCl (Figs. 3-4).

Together, our data suggest a model in which *P. aeruginosa* senses HOCl via the *fro* system,

which upregulates the production of antioxidant proteins that suppress the toxic effects of HOCl.

This would constitute an early detection mechanism that defends against neutrophils. A recent

finding identified the RclR transcriptional regulator as a sensor of HOCl *(27)*. Further work is

necessary to determine how antioxidative proteins and antioxidative agents reduce the toxic

activity of HOCl and the extent that RclR and *fro* are co-regulated by HOCl.

In addition to promoting colonization of the lung and gastrointestinal tract *(13, 28)*, the

fro system may have additional impacts on pathogenicity, such as increasing antibiotic

resistance. Bacteria that are exposed to low doses of HOCl have higher rates of horizontal gene

transfer and can acquire antibiotic resistance genes more rapidly *(29, 30)*. The tolerance of

P. aeruginosa to HOCl via *fro* may thus provide a mechanism to facilitate antibiotic resistance.

The findings here thus suggest that the *fro* pathway could be an important defense mechanism to

target. Inhibition of this pathway could increase *P. aeruginosa* susceptibility to neutrophil-

mediated killing while minimizing antibiotic resistance.

MATERIALS AND METHODS

Strains and Growth Conditions

Experiments were performed using the *P. aeruginosa* strain AL143, which contains the *yfp* gene integrated between *froA* and *froB* and a constitutively-expressed *mCherry* gene *(3)*. For experiments with response of *P. aeruginosa* against NaOCl Δ*froR* and Δ*froI* mutant strains were used *(3)*. Strains were streaked onto LB-Miller (BD Biosciences, Franklin Lakes, NJ) petri dishes containing 2% Bacto agar (BD Biosciences, Franklin Lakes, NJ), incubated at 37 °C to obtain single colonies, and inoculated into MinA minimal medium, which contains per 1 L of water: 4.5 g KH$_2$PO4, 10.5 g K$_2$HPO$_4$, 1 g (NH$_4$)$_2$SO$_4$, 1 mM MgSO$_4$ and 0.2 % (w/v) sodium citrate *(31)*. Strains were cultured in minimal medium for 18 hours in a roller drum spinning at 200 rpm at 37 °C, diluted 1:1000 into the same medium supplemented with oxidative or antioxidative agents, conditioned medium from neutrophils, or were not supplemented, cultured for 1 hour at 37 °C, and imaged using microscopy.

Phase Contrast and Fluorescence Microscopy

Bacteria were immobilized on 1% agarose pads containing minimal medium and imaged immediately. Images were acquired using a Nikon Eclipse Ti-E microscope (Nikon, Melville, NY) containing a Nikon 100X Plan Apo (1.45 N.A.) objective, a Nikon Ph3 phase contrast condenser annulus, a Sola light engine (Lumencor, Beaverton, OR), an LED-DA/FI/TX filter set (Semrock, Rochester, NY) for visualizing the mCherry fluorescence spectrum containing a 409/493/596 dichroic and 575/25 nm filters for excitation and 641/75 nm filters for emission, an LED-CFP/YFP/MCHERRY filter set (Semrock) for visualizing YFP fluorescence containing a

459/526/596 dichroic and 509/22 nm filters for excitation and 544/24 nm filters for emission,

and a Hamamatsu Orca Flash 4.0 V2 camera (Hamamatsu, Bridgewater, NJ).

Images were acquired using Nikon NIS-Elements and analyzed using custom built

software written previously *(32)* in Matlab (Mathworks, Natick, MA). See the "Code

availability" section below to download code. Briefly, cell masks were determined from phase

contrast images using an edge-detection algorithm. The YFP to mCherry fluorescence ratios in

the masked cell areas were computed. The expression of *fro* is quantified as the ratio of YFP to

mCherry fluorescence in individual *P. aeruginosa*, which is averaged over at least one hundred

P. aeruginosa. A minimum of 100 cells were imaged and analyzed in each experiment.

Imaging *P. aeruginosa* under flow

Microfluidic devices were fabricated using standard soft lithography techniques as

described previously *(33)* with channel dimensions of 100 μm x 50 μm (width x height) and a

flow rate of approximately 5 μL/min.

Growth Curves

Growth curve experiments were performed using a Synergy HTX multi-mode plate

reader and sterile, tissue-culture treated, clear bottom, black polystyrene 96-well microplates

(Corning, Corning, NY) containing 200 μL of culture in each well. The temperature set point

was 37 °C and preheated before beginning measurements. For experiments performed with

stationary phase bacteria, overnight cultures of bacteria were grown for 18 hours at 37 °C in the

roller drum at 200 rpm to saturation, diluted into minimal medium to an optical density at 600

nm (OD_{600}) of 0.01 containing NaOCl, H_2O_2, HNO_3 or no supplement. Plates were incubated at

37 °C with continuous orbital shaking with an amplitude of 3 mm at the frequency of 180 cycles

per minute and measured for OD_{600} every 20 minutes.

Cell size analysis

P. aeruginosa were cultured for 18 hours at 37 °C in a roller drum spinning at 200 rpm,

diluted 1:1000 into fresh minimal medium, treated 1 µM NaCl, 200 µM HCl or with serial

dilutions of NaOCl, H_2O_2, HNO_3 or media only, and incubated at 37 °C in 250 mL flasks for 3

hours with shaking at 225 rpm. Cells were immobilized on 1% agarose pads containing minimal

medium, imaged using phase contrast microscopy, and analyzed using the edge detection image

analysis tools in Matlab described above. The total pixel area of each individual cell was

determined by computing the mask area and converting from pixels to μm^2 by multiplying the

mask area by a factor of 0.004225 μm^2/pixel to account for the microscope camera pixel size and

objective magnification.

Neutrophil isolation

Blood samples from healthy donors between 18 to 65 years old were collected at the

Institute for Clinical and Translational Science at the University of California, Irvine, through

approved protocols with the UCI Institutional Review Board, and informed consent was obtained

from all donors. Neutrophils were prepared essentially as described in *(34)*. Briefly, 3% dextran

(Leuconostoc spp, Mr 450,000-650,000, Sigma-Aldrich, St. Louis, MO) in PBS (Gibco,

ThermoFisher, Waltham, MA) was used to separate red blood cells (RBCs) from whole blood.

Neutrophils were purified from the remaining cells by overlaying on a Ficoll density gradient

(GE Healthcare, Chicago, IL) following centrifugation in Ficoll-Paque Centrifugation Media

(GE Healthcare, Chicago, IL) for 25 minutes at $500 \times g$.

The remaining RBCs were lysed using RBC Lysis Buffer (Fisher Scientific, Hampton,

NH), and neutrophils were resuspended in RPMI 1640 medium (ATCC, Manassas, VA) and

purity was assessed by flow cytometry using the ACEA NovoCyte Flow Cytometer and

fluorescent antibodies APC-CD11b, FITC-CD16 and PE-CD66b (eBioscience, San Diego, CA).

Neutrophils were immediately used for neutrophil stimulation or co-incubation experiments with

P. aeruginosa. This procedure routinely yielded greater than 95% CD11B-positive neutrophil

populations.

Tissue culturing

J774.1 mouse macrophages (ATCC) or primary neonatal normal human dermal

fibroblasts (Neo-NHDF) (ATCC) were cultured in 10 mL DMEM (Gibco, ThermoFisher) in T75

flasks (VWR, Radnor, PA) at 37 °C with 5% CO_2. Macrophages were passaged every three days

through scraping and Neo-NHDF cells were passaged every 5 days through trypsinization.

Co-incubation of mammalian cells with P. aeruginosa

P. aeruginosa were cultured for 18 hours in minimal medium, centrifuged at 4600 x g for

2 minutes and resuspended in DPBS (Gibco, ThermoFisher) three times, and diluted to a final

OD_{600} of 0.2. Macrophages or Neo-NHDF cells that were harvested from growth flasks, or

freshly prepared neutrophils, were washed three times by centrifuging at 300 x g for 5 minutes

and resuspending in DPBS, and 1 mL aliquots containing 4×10^5 cells/mL were transferred into

flat-bottom dishes. *P. aeruginosa* culture was added to the cells at a multiplicity of infection of

10 and incubated at 37 °C with CO_2 for 30 minutes. Dishes were aspirated until approximately

20 µL of media remained and pads consisting of 1 % agarose containing DPBS were placed on

top of the samples in the dishes to immobilize bacteria and cells. Samples were imaged

immediately using microscopy.

Preparation of neutrophil-stimulated conditioned medium

360   Approximately 2 million/mL of freshly harvested neutrophils were incubated at 37 °C in a

roller drum spinning at a speed of 4 rotations per minute in PBGT medium (10 mM phosphate

buffer pH 7.4 (2.46 mM monobasic with 7.54 mM dibasic sodium phosphate), 140 mM sodium

chloride, 10 mM potassium chloride, 0.5 mM magnesium chloride, 1 mM calcium chloride, 1

mg/mL glucose, and 5 mM taurine *(24)*). Neutrophils were supplemented with 100 ng/mL of PMA

or left untreated for 1 hour, and were separated from the supernatant by 5 minute centrifugation at

4,600 x g at 25 °C. The supernatant was isolated and used as conditioned medium.

367   *P. aeruginosa* were cultured in minimal medium for 18 hours at 37 °C in a roller drum

spinning at 200 rpm, diluted 1:1000 in conditioned media with or without the addition of 100 µM

methionine, and incubated for 3 hours. Bacterial culture from each sample was concentrated using

cellulose syringe filters with a 0.22 µm pore size (Millipore, Burlington, MA). *P. aeruginosa* cells

were immobilized on 1% agarose pads and imaged. Data analysis was performed as described in

the "cell size analysis" methods section.

Antioxidant treatments

375   *P. aeruginosa* cultured for 18 hours at 37 °C in a roller drum spinning at 200 rpm, diluted

1:1000 into fresh minimal medium were diluted 1:1000 into minimal medium alone or

supplemented with 1 µM NaOCl with an addition of methionine (1 µM, 10 µM, 100 µM), β-

mercaptoethanol (βME) (1 µM, 10 µM, 100 µM) or cysteine (0.1 µM, 1 µM, 10 µM). 50 mL of

each bacterial culture was incubated at 37 °C in 250 mL flasks for 3 hours with shaking at 225

rpm. Data analyses were performed as described in the "cell size analysis" methods section

showed that the visible decrease of *fro* response to 1 µM NaOCl was observed when using 0.1

µM cysteine, 1 µM methionine and 1 µM βME. While *fro* response was not elevated at higher

concentrations of these antioxidants.

Code Availability

The custom Matlab scripts used to process and analyze the fluorescence microscopy data

have been previously described *(32)* and are freely available from Zenodo at

https://doi.org/10.5281/zenodo.3898289. Gene Ontology enrichment analysis was performed

using GOEnrichment version 2.0.1 *(35)*. The analysis of RNA-seq data was performed using

the limma Bioconductor package in Galaxy version 3.48.0 *(36)*.

Supplementary Material

Supplemental Figures S1-S3

Supplemental Spreadsheet 1

Supplemental Spreadsheet 2

Source Data file

397
398

REFERENCES

1. Centers for Disease Control and Prevention (U.S.), *Antibiotic resistance threats in the United*
States, 2019 (Centers for Disease Control and Prevention (U.S.), 2019;
https://stacks.cdc.gov/view/cdc/82532).

2. F. C. Fang, E. R. Frawley, T. Tapscott, A. Vázquez-Torres, Bacterial Stress Responses during
Host Infection. *Cell Host Microbe* **20**, 133–143 (2016).

3. J. E. Sanfilippo, A. Lorestani, M. D. Koch, B. P. Bratton, A. Siryaporn, H. A. Stone, Z. Gitai,
Microfluidic-based transcriptomics reveal force-independent bacterial rheosensing. *Nat*
Microbiol **4**, 1274–1281 (2019).

4. E. G. Lavoie, T. Wangdi, B. I. Kazmierczak, Innate immune responses to Pseudomonas
aeruginosa infection. *Microbes Infect* **13**, 1133–1145 (2011).

5. A. W. Segal, How neutrophils kill microbes. *Annu Rev Immunol* **23**, 197–223 (2005).

6. M. B. Hampton, A. J. Kettle, C. C. Winterbourn, Inside the neutrophil phagosome: oxidants,
myeloperoxidase, and bacterial killing. *Blood* **92**, 3007–3017 (1998).

7. F. C. Fang, Antimicrobial actions of reactive oxygen species. *mBio* **2** (2011),
doi:10.1128/mBio.00141-11.

8. M. J. Gray, W.-Y. Wholey, U. Jakob, Bacterial responses to reactive chlorine species. *Annu*
Rev Microbiol **67**, 141–160 (2013).

9. C. Estrela, R. G. Ribeiro, C. R. A. Estrela, J. D. Pécora, M. D. Sousa-Neto, Antimicrobial
effect of 2% sodium hypochlorite and 2% chlorhexidine tested by different methods. *Braz Dent J*
14, 58–62 (2003).

10. J. M. Albrich, J. K. Hurst, Oxidative inactivation of Escherichia coli by hypochlorous acid.
Rates and differentiation of respiratory from other reaction sites. *FEBS Lett* **144**, 157–161
(1982).

11. J. E. Harrison, J. Schultz, Studies on the chlorinating activity of myeloperoxidase. *J Biol*
Chem **251**, 1371–1374 (1976).

12. E. T. Jensen, A. Kharazmi, K. Lam, J. W. Costerton, N. Høiby, Human polymorphonuclear
leukocyte response to Pseudomonas aeruginosa grown in biofilms. *Infect Immun* **58**, 2383–2385
(1990).

13. E. Potvin, D. E. Lehoux, I. Kukavica-Ibrulj, K. L. Richard, F. Sanschagrin, G. W. Lau, R. C.
Levesque, In vivo functional genomics of Pseudomonas aeruginosa for high-throughput
screening of new virulence factors and antibacterial targets. *Environ Microbiol* **5**, 1294–1308
(2003).

14. D. M. Cornforth, J. L. Dees, C. B. Ibberson, H. K. Huse, I. H. Mathiesen, K. Kirketerp-
Møller, R. D. Wolcott, K. P. Rumbaugh, T. Bjarnsholt, M. Whiteley, Pseudomonas aeruginosa
transcriptome during human infection. *Proc Natl Acad Sci U S A* **115**, E5125–E5134 (2018).

15. G. L. Winsor, E. J. Griffiths, R. Lo, B. K. Dhillon, J. A. Shay, F. S. L. Brinkman, Enhanced
annotations and features for comparing thousands of Pseudomonas genomes in the Pseudomonas
genome database. *Nucleic Acids Res* **44**, D646-653 (2016).

16. A. L. Boechat, G. H. Kaihami, M. J. Politi, F. Lépine, R. L. Baldini, A novel role for an ECF
sigma factor in fatty acid biosynthesis and membrane fluidity in Pseudomonas aeruginosa. *PLoS*
One **8**, e84775 (2013).

17. A. Karlsson, J. B. Nixon, L. C. McPhail, Phorbol myristate acetate induces neutrophil
NADPH-oxidase activity by two separate signal transduction pathways: dependent or
independent of phosphatidylinositol 3-kinase. *J Leukoc Biol* **67**, 396–404 (2000).

18. K. V. Farrant, L. Spiga, J. C. Davies, H. D. Williams, Response of Pseudomonas aeruginosa
to the Innate Immune System-Derived Oxidants Hypochlorous Acid and Hypothiocyanous Acid.
J Bacteriol **203**, e00300-20 (2020).

19. C. C. Winterbourn, Comparative reactivities of various biological compounds with
myeloperoxidase-hydrogen peroxide-chloride, and similarity of the oxidant to hypochlorite.
Biochim Biophys Acta **840**, 204–210 (1985).

20. D. I. Pattison, M. J. Davies, Reactions of myeloperoxidase-derived oxidants with biological
substrates: gaining chemical insight into human inflammatory diseases. *Curr Med Chem* **13**,
3271–3290 (2006).

21. G. Atmaca, Antioxidant effects of sulfur-containing amino acids. *Yonsei Med J* **45**, 776–788
(2004).

22. C. L. Hawkins, D. I. Pattison, M. J. Davies, Hypochlorite-induced oxidation of amino acids,
peptides and proteins. *Amino Acids* **25**, 259–274 (2003).

23. P. Bin, R. Huang, X. Zhou, Oxidation Resistance of the Sulfur Amino Acids: Methionine and
Cysteine. *Biomed Res Int* **2017**, 9584932 (2017).

24. J. M. Dypbukt, C. Bishop, W. M. Brooks, B. Thong, H. Eriksson, A. J. Kettle, A sensitive
and selective assay for chloramine production by myeloperoxidase. *Free Radic Biol Med* **39**,
1468–1477 (2005).

25. L. C. Seaver, J. A. Imlay, Alkyl hydroperoxide reductase is the primary scavenger of
endogenous hydrogen peroxide in Escherichia coli. *J Bacteriol* **183**, 7173–7181 (2001).

26. S. M. Brown, M. L. Howell, M. L. Vasil, A. J. Anderson, D. J. Hassett, Cloning and
characterization of the katB gene of Pseudomonas aeruginosa encoding a hydrogen peroxide-
inducible catalase: purification of KatB, cellular localization, and demonstration that it is
essential for optimal resistance to hydrogen peroxide. *J Bacteriol* **177**, 6536–6544 (1995).

27. K. V. Farrant, L. Spiga, J. C. Davies, H. D. Williams, Response of Pseudomonas aeruginosa
to the Innate Immune System-Derived Oxidants Hypochlorous Acid and Hypothiocyanous Acid.
J Bacteriol **203**, e00300-20 (2020).

28. D. Skurnik, D. Roux, H. Aschard, V. Cattoir, D. Yoder-Himes, S. Lory, G. B. Pier, A
comprehensive analysis of in vitro and in vivo genetic fitness of Pseudomonas aeruginosa using
high-throughput sequencing of transposon libraries. *PLoS Pathog* **9**, e1003582 (2013).

29. A.-M. Hou, D. Yang, J. Miao, D.-Y. Shi, J. Yin, Z.-W. Yang, Z.-Q. Shen, H.-R. Wang, Z.-G.
Qiu, W.-L. Liu, J.-W. Li, M. Jin, Chlorine injury enhances antibiotic resistance in Pseudomonas
aeruginosa through over expression of drug efflux pumps. *Water Res* **156**, 366–371 (2019).

30. M. Jin, L. Liu, D. Wang, D. Yang, W. Liu, J. Yin, Z. Yang, H. Wang, Z. Qiu, Z. Shen, D.
Shi, H. Li, J. Guo, J. Li, Chlorine disinfection promotes the exchange of antibiotic resistance
genes across bacterial genera by natural transformation. *ISME J* **14**, 1847–1856 (2020).

31. H. Malke, Jeffrey H. Miller, A Short Course in Bacterial Genetics – A Laboratory Manual
and Handbook forEscherichia coli and Related Bacteria. Cold Spring Harbor 1992. Cold Spring
Harbor Laboratory Press. ISBN: 0–87969-349–5. *J Basic Microbiol* **33**, 278–278 (1993).

32. T. Doolin, H. M. Amir, L. Duong, R. Rosenzweig, L. A. Urban, M. Bosch, A. Pol, S. P.
Gross, A. Siryaporn, Mammalian histones facilitate antimicrobial synergy by disrupting the
bacterial proton gradient and chromosome organization. *Nat Commun* **11**, 3888 (2020).

33. A. Siryaporn, M. K. Kim, Y. Shen, H. A. Stone, Z. Gitai, Colonization, competition, and
dispersal of pathogens in fluid flow networks. *Curr Biol* **25**, 1201–1207 (2015).

34. H. L. Clark, S. Abbondante, M. S. Minns, E. N. Greenberg, Y. Sun, E. Pearlman, Protein
Deiminase 4 and CR3 Regulate Aspergillus fumigatus and β-Glucan-Induced Neutrophil
Extracellular Trap Formation, but Hyphal Killing Is Dependent Only on CR3. *Front Immunol* **9**,
1182 (2018).

35. Gene Ontology Consortium, The Gene Ontology resource: enriching a GOld mine. *Nucleic
Acids Res* **49**, D325–D334 (2021).

36. C. W. Law, Y. Chen, W. Shi, G. K. Smyth, voom: Precision weights unlock linear model
analysis tools for RNA-seq read counts. *Genome Biol* **15**, R29 (2014).

Acknowledgments: The authors thank Michael Z. Zulu and Stephanie Matsuno for their

feedback on prior drafts of this article and Hew Yeng Lai for imaging of bacteria in microfluidic

devices. **Funding:** This work was supported by the Stanley Behrens Fellows in Medicine Award

to L.U, and grants from the National Institutes of Health (NIH) National Institute of Biomedical

Imaging and Bioengineering (NIBIB) (R21EB027840-01), the National Cancer Institute (NCI)

(DP2CA250382-01), the National Science Foundation (NSF) (DMS1763272), and a grant from

the Simons Foundation (594598QN) to T.L.D. **Author contributions:** I.F. and A.S. conceived

and designed the experiments. I.F. and A.H. performed the experiments. L.U. prepared

neutrophils from human samples. I.F. and L.U. analyzed data and wrote the initial draft of the

paper. I.F., L.U., T.L.D., and A.S. discussed results and edited the paper. **Competing interests:**

Authors declare that they have no competing interests. **Data and materials availability:** Raw

data for all figures that contain t-test analyses are available in the Supplementary Information

Source Data file.

Supplementary Material

Supplemental Figures S1-S3
Supplemental Spreadsheet 1 (Microsoft Excel format)
Supplemental Spreadsheet 2 (Microsoft Excel format)
Source Data file (Microsoft Excel format). Contains raw data and statistics for main and supplemental figures

Supplemental Figure S1

Fig. S1 Effects on cell size and global transcription in *P. aeruginosa* treated with NaOCl, H₂O₂, or HNO₃. Top: average area of individual *P. aeruginosa* cells, Middle: average mCherry/area, Bottom: average YFP/area in *P. aeruginosa* treated with dilutions of H2O2, NaOCl, HNO3, or untreated for 3 hours. Boxplots show the mean (black line) and the interquartile range of three independent experiments. Blue boxes indicate the datasets that were used to compute the values for Fig. 3D.

Supplemental Figure S2

Fig. S2. Dose-dependent effects of antioxidant supplements on *fro* expression induced by HOCl. Quantification of *fro* expression levels in *P. aeruginosa* cultures treated with 1 μM NaOCl and supplemented with methionine, cysteine, or beta-mercaptoethanol (BME). Plots show the mean and SEM for three independent experiments. P-values were calculated using two-tailed, unequal variance t-test. ns indicates p > 0.05.

Supplemental Figure S3

Fig. S3. Effects of antioxidants alone on *fro* expression. Quantification of *fro* expression levels in *P. aeruginosa* cultures treated with 1 µM methionine, 0.1 µM cysteine, 1 µM beta-mercaptoethanol (BME), or untreated (MinA data replotted from Fig. 3C). Plots show the mean and the standard error of the mean (SEM) for three independent experiments. P-values were calculated using two-tailed, unequal variance t-test. ns indicates $p > 0.05$.