Mutability Patterns Across the Spike Glycoprotein Reveal the Diverging and Lineage-specific Evolutionary Space of SARS-CoV-2.

Roberth A. Rojas Chávez[1], Mohammad Fili[2], Changze Han[1], Syed A. Rahman[3], Isaiah G. L. Bicar[1], Sullivan Gregory[1], Guiping Hu[2], Jishnu Das[3], Grant D. Brown[4], and Hillel Haim[1#]

[1] Department of Microbiology and Immunology, The University of Iowa, Iowa City, IA.

[2] Department of Industrial and Manufacturing Systems Engineering, Iowa State University, Ames, IA.

[3] Center for Systems Immunology, Departments of Immunology and Computational & Systems Biology, University of Pittsburgh School of Medicine, Pittsburgh, PA.

[4] Department of Biostatistics, College of Public Health, The University of Iowa, Iowa City, IA.

[#] To whom correspondence should be addressed:
Hillel Haim, MD, PhD
Department of Microbiology and Immunology
The University of Iowa
51 Newton Rd, 3-770 BSB
Iowa City, Iowa, 52242
Phone: (319) 335-9989
Email: Hillel-haim@uiowa.edu

Short Title: Evolutionary space of SARS-CoV-2 variants.

Keywords: SARS-CoV-2, COVID-19, Virus evolution, Spike protein, Mutational space.

Abstract word count: 150.

Main text word count: 4430.

ABSTRACT

Mutations in the spike glycoprotein of SARS-CoV-2 allow the virus to probe the sequence space in search of higher-fitness states. New sublineages of SARS-CoV-2 variants-of-concern (VOCs) continuously emerge with such mutations. Interestingly, the sites of mutation in these sublineages vary between the VOCs. Whether such differences reflect the random nature of mutation appearance or distinct evolutionary spaces of spike in the VOCs is unclear. Here we show that each position of spike has a lineage-specific likelihood for mutations to appear and dominate descendent sublineages. This likelihood can be accurately estimated from the lineage-specific mutational profile of spike at a protein-wide level. The mutability environment of each position, including adjacent sites on the protein structure and neighboring sites on the network of co-mutability, accurately forecast changes in descendent sublineages. Mapping of imminent changes within the VOCs can contribute to the design of immunogens and therapeutics that address future forms of SARS-CoV-2.

INTRODUCTION

Since emerging in December 2019, SARS-CoV-2 has caused devastating effects worldwide. By June 2022, more than 6 million deaths have been attributed to the infection, and estimated economic losses greater than $10 trillion are expected by the end of this year[1,2]. Several SARS-CoV-2 VOCs have appeared at different time points of the pandemic[3,4,5]. Most mutations in the VOCs that impact infection are found in the spike protein that adorns the virus surface. Spike mediates fusion with host cells and is the primary target for antibodies elicited by infection or vaccination[6]. Changes in spike can increase virus infectivity, transmissibility or resistance to vaccine-elicited antibodies and therapeutics[7]. New sublineages of the VOCs continuously appear with such mutations in spike[8,9]. Interestingly, only a minority of these changes are convergent, whereby the same substitution occurs in multiple lineages[10,11,12,13]. The latter are generally guided by positive selection pressures[14]. By contrast, most mutations that define VOC sublineages are found at distinct positions of spike and occur at evolutionarily neutral sites[15,16]. This observation raises two important questions. First, do the distinct patterns of mutations in the lineages reflect the stochastic nature of their appearance or a lineage-specific likelihood for their emergence? Second, if not driven solely by stochasticity, what clues can we identify that will inform of the evolutionary space of each lineage? Answers to these questions are critical for our ability to develop vaccines and therapeutics that can maintain their efficacy against future forms of this virus.

To address the above questions, we examined the frequency of independent mutation events at each position of spike in different SARS-CoV-2 VOCs. Lineage-specific mutational spaces were observed, as defined by the patterns of low-frequency substitutions in spike within populations infected by each lineage. We compared the mutational space of spike in each lineage with the evolutionary path of the virus within the lineage (i.e., the observed mutations that define descendant sublineages). We discovered that the sites of change in the emergent sublineages were characterized by high mutability in the ancestral lineage and a high mutability "environment", composed of adjacent positions on the protein structure and co-mutable sites across the protein. These measures of positional and environmental variability predicted remarkably well the changes observed in the new sublineages of the VOCs. Our studies of spike mutability at the protein-wide level reveal the diversifying nature of the evolutionary space of this protein and demonstrate the high predictability of the changes that give rise to new SARS-CoV-2 spike forms.

RESULTS

The mutational space of SARS-CoV-2 spike is lineage-specific and diversifying

We examined the sites of mutation in the spike protein that define sublineages of SARS-CoV-2 VOCs. Using designations of the Pango classification system[17], the lineages compared were B.1.1.7 (VOC Alpha), P.1 (Gamma), AY.4 (Delta), and lineages BA.1 and BA.2 (Omicron). Sites of mutation in descendant sublineages of the above that emerged until April 8th 2022 were compared. In addition, we calculated for these sites the synonymous and nonsynonymous mutation rates in each lineage to infer the sites under positive selection. As expected, most sites of change that define descendent sublineages did not show evidence for positive selection in their ancestral lineages (**Fig. 1a**). Importantly, limited similarity was observed between the sites of mutations in the different VOCs – only three of the 41 sublineage-defining mutation sites appeared in more than one VOC.

To investigate the basis for these distinct patterns of change, we first compared the mutational space of spike between the above lineages, as defined by the collection of sites that exhibit amino acid variability among strains that are phylogenetically closest to the lineage ancestor. In addition, as examples, we included the highly-prevalent sublineages BA.1.1 and AY4.2, which emerged from lineages BA.1 and AY.4, respectively. All other Pango-designated sublineages of the above variants were removed from the datasets. As a group representative of isolates closest to the SARS-CoV-2 ancestral strain (designated the SARS-CoV-2 "baseline"), we used sequences within 0.0015 substitutions per site from the SARS-COV-2 spike ancestral sequence (Wuhan strain, accession number NC_045512)[18]. Spike sequences that appeared at least twice in the population were aligned and "compressed" to obtain a single representative for each unique sequence. Evolutionary relationships among them were inferred, and a maximum likelihood phylogenetic tree was constructed (see Methods and **Fig. 1b**). To quantify amino acid variability at each position of spike, the lineages were partitioned into clusters of 50 sequences based on phylogeny, and the proportion of clusters that contain amino acid variability at each position was calculated (**Fig. 1c**). We designate this measure of variability "volatility", which quantifies the frequency of substitution events.

Considerable differences were observed between the volatility profiles of spike in the diverse lineages (see RBD positions in **Fig. 1d** and all spike positions in **Supplementary Fig. 1a** and **1b**). To determine the relationships between volatility patterns, we partitioned each lineage into groups of 10 clusters. The absence or presence of amino acid variability at each spike position was determined for each group, and all groups were assigned 1273-bit vectors that describe their

volatility profiles at all spike positions. Jaccard distances between the vectors were applied to
determine hierarchical relationships using the unweighted pair group method with arithmetic mean
(UPGMA) approach[19,20]. Clear clustering of group profiles from the same lineage was observed
(**Fig. 1e**). To determine statistical significance of these patterns, Euclidean distances were
measured between all vectors, and intra-lineage distances compared with inter-lineage distances
using a permutation test[21]. As shown in **Fig. 1f**, all lineages (except the smaller AY4.2) exhibited
significant specificity of their volatility patterns.

To quantify divergence of the volatility patterns, we performed pairwise comparisons
between the volatility profiles of any two lineages and the genetic distance that separates them.
Positions of the RBD, N-terminal domain of spike (NTD) and S2 subunit were analyzed
separately. The correlation coefficient for any lineage pair was then compared with the nucleotide
distance between the lineage founders. For the RBD and S2 subunits, a strong negative
relationship was observed between the genetic distance that separates any two lineage founders
and the correlation between their volatility profiles (**Fig. 1e**). For the NTD, which contains a
relatively high proportion of sites with mutations, we did not observe such a relationship.

These findings suggest that the mutational space of spike (i.e., the collection of sequence
states that are sampled) is specific for each lineage and is diversifying. As such, and assuming
that the mutational space corresponds with the evolutionary path of the virus in the population,
these findings also suggest that each lineage may have a distinct set of changes that can appear
in descendent sublineages. This possibility was explored in the studies described below.

**A high volatility state and a high volatility environment increase the likelihood of spike
positions to emerge with founder mutations in descendent lineages**

We examined the relationship between the volatility of any spike position in a population
of related strains and the emergence of mutations at this position in descendent lineages. To this
end, we first analyzed mutability profiles in the SARS-CoV-2 baseline that preceded emergence
of the VOCs. These profiles were compared with the mutations that define the emergent lineages.
For simplicity, we focused these analyses on all 615,374 spike sequences from samples collected
worldwide between December 2019 and July 2021. Evolutionary relationships among the
nucleotide sequences were inferred and a maximum likelihood tree was constructed (**Fig. 2a**).
We then partitioned the tree into discrete groups separated by a minimal distance of 0.004
substitutions per site. As expected, many groups corresponded to known SARS-CoV-2 VOCs.

The baseline groups were distinguished from the terminal emergent groups (G_{T1}-G_{T8}) using a

threshold of 0.0015 substitutions per site between the centroid of each group and the SARS-CoV-

2 spike ancestral sequence. All groups are described in **Supplementary Table 1**.

Volatility at each spike position in the baseline was compared with the absence or

presence of two types of mutations: **(i) Lineage-founder mutations (LFMs)**, which are found in

the group ancestors and in at least 50% of all sequences from that group, and **(ii) Sublineage-**

founder mutations (sLFMs), which are not found in the group ancestor and represent clonal

expansions that dominate at least one 50-sequence cluster but less than 50% of all group clusters

(see examples in **Supplementary Fig. 2a**). A total of 43 LFMs and 16 sLFMs were detected in

the baseline and terminal groups (see **Supplementary Table 1**). Most positions with high volatility

in the baseline emerged with LFMs or sLFMs (see positions of spike subunit S1 in **Fig. 2b** and of

subunit S2 in **Supplementary Fig. 2b**). Among positions with the highest volatilities, most

appeared as s/LFMs in at least one group (**Supplementary Fig. 2c**). Sites of s/LFMs were more

volatile than sites with no such mutations (**Fig. 2c**). Furthermore, non-volatile sites in the baseline

did not emerge with s/LFMs in any baseline or terminal group (**Fig. 2d**). Therefore, for any given

site, a high level of volatility (i.e., a high frequency of independent mutation events) in the baseline

group precedes (as inferred phylogenetically) the emergence of s/LFMs in the descendent

lineages.

We recently examined the within-host patterns of amino acid variability in the envelope

glycoproteins (Envs) of human immunodeficiency virus type 1 (HIV-1)[22]. We found that the

variability at many positions of the CD4-binding site can be accurately estimated by the variability

at adjacent positions on the three-dimensional structure of the protein. Analysis of the spatial

distribution patterns of volatile sites on the SARS-CoV-2 spike structure suggested a similar

clustering of volatility at multiple loci, most notably in the NTD (see **Fig. 2e** and sites with

statistically significant clustering in **Fig. 2f**). We hypothesized that if such associations are "stable"

over time, then the likelihood for future changes at any position may be associated with volatility

of its neighboring positions. To test this hypothesis, we generated a variable (designated D) that

describes for each position i the total distance-weighted "environmental" volatility:

164
$$D_i = \sum_{s=1}^{n} \frac{1}{\Delta_{is}} \cdot V_s \qquad \textbf{[1]}$$

where n is the number of positions s within 6 Å of position i, Δ_{is} is the distance between the closest

two atoms of positions i and each position s, and V_i is the volatility at each position s. Similar to

the volatility values, D values were higher for positions that emerged with s/LFMs (**Fig. 2g**).

Furthermore, positions with a non-volatile environment (i.e., a D value of zero) did not emerge

with s/LFMs (**Fig. 2h**). Therefore, high volatility at any position in the SARS-CoV-2 baseline and

high volatility at adjacent positions on the protein increase the likelihood of the site to emerge with

s/LFMs in descendent lineages.

Volatility at neighboring sites on the network of co-volatility increases the likelihood of spike positions to emerge with lineage founder mutations

We examined whether the clustering patterns of volatility at adjacent positions on spike

can be generalized to describe associations that are not dependent on physical proximity. To this

end, we used the 114 baseline clusters to determine the co-occurrence of volatility at any two

spike positions within the clusters (see schematic in **Fig. 3a**). P-values were calculated using

Fisher's exact test and used to construct the network of co-volatile sites, whereby the edges that

connect the nodes (positions) are defined by the statistical significance of the association between

their volatility patterns (see example of a network segment in **Fig. 3b** and distribution of P-values

in **Supplementary Fig. 3a**). To determine the significance threshold to apply for network

construction, we examined structural properties of the network and its robustness to random

deletion of edges. Two network topological metrics were assessed: **(i)** Degree distribution, which

describes the average number of connections each node has with other nodes, and **(ii)** Closeness

centrality, which describes for each node the sum of the path lengths to all other nodes in the

network (more central nodes have lower values)[23]. For robust scale-free networks, limited

random-edge deletions only minimally perturb their topological properties[24]. We found that

networks defined at a more stringent significance threshold (P<0.01) were more robust to edge

deletions, with minimal impact on both degree distribution and closeness centrality at the expense

of losing information (**Fig. 3c** and **Supplementary Fig. 3b**). By contrast, when less stringent

significance thresholds were used (P<0.1), the number of edges was greater (i.e., they contained

more information regarding the co-volatile positions); however, the network was less robust to

edge deletions. This suggested that an intermediate significance threshold (P<0.05) would

provide a sufficiently stable network without losing most information.

We examined whether, for any position i of spike, presence of high volatility at its network-

associated co-volatile sites (q) is associated with emergence of s/LFMs. To this end, we

generated a simple measure (R) designed to capture for each position i the total volatility of its

network-neighbors q ($q_1, q_2, q_3 \dots q_n$), using a P-value of 0.05 as the threshold:

$$R_i = \sum_{q=1}^{n} w_{iq} \cdot V_q \qquad\qquad \textbf{[2]}$$

where V_q is the volatility at each position q calculated using the baseline sequences, and w_{iq} is the

evidence for association between volatility of position i and each of its positions q (calculated as

the $-\log_{10}$(P-value) in Fisher's test). Similar to the V and D values, R values were significantly

higher for positions with s/LFMs relative to positions with no such mutations (**Fig. 3d**).

Furthermore, an R value of zero in the baseline was invariably associated with lack of s/LFM

appearance (**Fig. 3e**). Overall, V and R values for any position correlated well, and considerably

better than their correlation with D (**Fig. 3, f-h**). Nevertheless, as shown below, V and R values

exhibit different levels of predictive performance when small population sizes are tested.

We compared the performance of the three variables (V, D and R) to predict the

emergence of LFMs or sLFMs using a univariate logistic regression model. Higher classification

metrics were observed for V and R relative to D, with area under the receiver operating

characteristic curve (AUC) values higher than 0.9 for both V and R (**Fig. 3i**). In comparison,

precision of these variables was modest, at 0.3 for V and 0.24 for R, indicating a relatively high

false-positive rate. Taken together, these findings show that the emergence of s/LFMs at any

spike position is associated with a state of high volatility in the ancestral lineage, as well as high

volatility at adjacent positions on the protein and at associated sites on the co-volatility network.

218 Volatility profiles among sequences from the early pandemic capture the mutational
219 patterns of the emergent lineages

We examined whether a combination of the volatility-based variables would better capture

the observed evolutionary path of the virus than each of them separately. To this end, we indexed

all sequences by the time of sample collection and tested whether viruses that temporally

preceded emergence of SARS-CoV-2 lineages can predict the mutations they contain. For these

analyses, sequences were classified by their Pango lineage designations rather than our

phylogeny-based group definitions. We first determined the formation time of each lineage,

defined as the date by which 26 unique sequences from the lineage were detected (see **Fig. 4a**

and **Supplementary Table 2**). Based on these timelines, we divided the sequences into an "early-

phase" group that is used to predict emergence of the LFMs in the "lineage-emerging phase". The

early-phase group included one sequence from lineage B.1.1.7 and none from the other emergent

lineages. Six minor lineages emerged early in the pandemic that contained mutations at positions

614, 222 and 477 (see **Supplementary Table 3**). To avoid a potential bias, these positions were
excluded from our analyses. A total of 67 LFM sites were identified in the lineage-emerging phase.

The early-phase sequences were divided into 27 clusters of 50 sequences, which were
used to calculate V, D and R values. These values were applied to a logistic regression model
that was trained to predict the emergence of LFMs using the phylogeny-indexed baseline
sequences (see Methods section). The output of the model is the probability of each site to
emerge with LFMs in the lineage-emerging phase. For all VOCs tested, as well the non-VOC
lineages (analyzed collectively), the probabilities calculated for LFM sites were significantly higher
than probabilities assigned to the non-LFM sites (**Fig. 4b**). Predictions based on the combined
model exhibited considerably higher performance than those based on the individual variables
(**Fig. 4, c-e**).

To examine the changes in probabilities assigned to the sites of mutation during the early
stages of the pandemic, we calculated V, D and R values and the combined probability using
increasing numbers of sequences indexed by the time of sample collection. Interestingly, the
pattern of LFMs was predicted with high sensitivity and specificity by the time three clusters were
formed (150 unique sequences), corresponding to samples collected until April 1st, 2020 (**Fig. 4f**
and **Supplementary Fig. 4, a-c**)**.** Of the individual predictors, R exhibited the highest
performance, modestly lower than the combined probability, whereas performance of V gradually
increased. Further analysis of the performance of the first three clusters showed that higher
probabilities were assigned to mutation sites of lineages that emerged at earlier stages of the
pandemic (**Fig. 4g** and **Supplementary Fig. 4d**). Higher probabilities were also assigned to
convergent sites (i.e., those that emerged with LFMs in multiple lineages, **Fig. 4h** and
Supplementary Fig. 4e).

We note that while V and R values calculated using all sequences of the baseline group
correlated well (**Fig. 3f**), the performance of R was considerably higher when a small number of
sequences was available (**Fig. 4f**). For example, analysis of the two major VOCs circulating during
the lineage-emerging phase showed near-maximal R values for most LFM sites in March 2020
whereas V values of these sites gradually increased over time (**Supplementary Fig. 5**).

Taken together, these findings show that a high level of volatility at any site and at its
spatial- and network-associated sites precedes emergence of mutations in descendant lineages.
A small number of sequences is required to accurately estimate the likelihood of sites for
emergence as LFMs. Total volatility at network-associated sites exhibits a higher level of
sensitivity at earlier stages of the pandemic than volatility values of the sites.

Spike mutations in emerging SARS-CoV-2 sublineages are accurately forecasted by patterns of volatility in their ancestral lineages

We examined the patterns of volatility among sequences that preceded emergence of the VOC sublineages. For this purpose, we focused on SARS-CoV-2 lineages from the major VOCs, including B.1.1.7, P1, AY.4, BA.1 and BA.2. Sequences from the baseline of each lineage were used to forecast the mutations that define its descendent sublineages. All emergent sublineages with Pango designations and all clusters of 50 sequences that contain a non-lineage-ancestral residue as the majority variant at any site were excluded from the datasets. The remaining clusters were used to calculate V, D and R values and to assign a mutation probability to each position. Two mutation types were tested as outcomes: **(i)** Mutations that define Pango sublineages of the VOCs, and **(ii)** Mutations that are dominant in two or more 50-sequence clusters of the lineage (but are not assigned a Pango sublineage designation). As shown in **Fig. 5a**, both outcomes were predicted well using the baseline sequences of each variant (see **Supplementary Table 3** for probability values). To determine the lineage specificity of the predictions, we compared the probabilities assigned to all sites of each lineage with the mutational outcomes in all other lineages. Consistent with the above findings, the highest AUC values were observed for predictions of the changes that occurred in the homologous lineage (**Fig. 5b**).

We also investigated the changes in probabilities assigned to the sites of sublineage mutations from the time their ancestral lineage had emerged. For these tests, we focused on AY.4, which has circulated in the population for a longer time period than other lineages (global emergence in May 2021). Mutation probabilities were calculated using increasing amounts of sequences indexed by time (**Fig. 5c**). Similar to the mutations that define lineages B.1.1.7 and B.1.617.2 (**Supplementary Fig. 5**), most sites of change in the sublineages of AY.4 exhibited high mutation probabilities at early stages after AY.4 emergence. Of the 14 sites of mutation in AY.4 sublineages, nine surpassed the 0.99 probability threshold (mean 95[th] probability percentile) at least once during the first month after emergence of AY.4. We note that in these tests a positive outcome was defined as a mutation that appeared in immediately descendent sublineages (e.g., mutations that define AY.4.2 but not mutations that define AY.4.2.1). Nevertheless, several sites with high mutation probabilities were also observed in second-order lineages. For example, the second-highest probability in BA.1 was assigned to position 1081. Since this change was observed in the second-order sublineage BA.1.15.1, as designated by the Pango system, it was not considered a positive outcome site.

Finally, we examined the specific changes in amino acid occupancy at the sites of mutation

in the emergent sublineages. In most cases, the minority variant with the highest frequency in the

baseline group of each lineage also appeared as the dominant residue in the new sublineage

(see characters in red font in **Fig. 5d**). These findings further support the notion that the

evolutionary path of spike within each lineage is accurately captured by its mutational space

across all sites of the protein, as detected among early isolates of the lineage.

DISCUSSION

Since January 2022, SARS-CoV-2 variant Omicron has dominated the landscape of VOCs

circulating worldwide. Both major lineages of Omicron (BA.1 and BA.2) contain mutations in spike

that reduce virus sensitivity to immune sera and COVID-19 therapeutics[25,26]. To address these

unique antigenic properties[27,28], Omicron-specific immunogens have been developed and

tested[29,30,31]. Nevertheless, new sublineages of this VOC emerge with mutations in spike that

further impact virus transmissibility and sensitivity to COVID-19 vaccines[8,9]. Most changes that

give rise to new sublineages do not appear to be driven by positive selective pressures (**Fig. 1a**).

Instead, they occur at evolutionarily neutral positions and "hitchhike" onto the driver mutations[32].

As such, it would be expected that the evolutionary space available for spike (i.e., the sequence

states that can be occupied by expanding lineages of the virus) would be large and driven by the

stochastic nature of the hitchhiking event. Here we introduce a simple probabilistic definition of

the evolutionary space of the virus. We show that, in fact, diverse SARS-CoV-2 lineages have

vastly distinct evolutionary spaces. Each position of spike has a specific and measurable

likelihood to appear as a dominant mutation within descendants of each lineage.

The volatility of each site only partially captures its likelihood for emergence with LFMs.

More accurate estimates are provided by the mutability profile of spike at the whole-protein level,

including adjacent sites on the protein and co-volatile sites. The latter variable is more sensitive

to the changes at shorter time frames from emergence of the parental lineage. How does volatility

of the "environment" capture the mutability of each site? Clustering of volatile sites on the linear

sequence of the protein can be explained by mutational hotspots due to properties of the viral

RNA[33,34]. Clustering on the three-dimensional structure can be explained by high permissiveness

of the region for change due to their limited impact on fitness[35]. By contrast, the association

between volatility of sites separated by larger distances on the protein is less intuitive. We propose

that such associations describe the epistasis network of spike (i.e., the sites that the amino acid

occupancy of one affects the fitness of another). Indeed, the volatility of each position likely

captures its fitness profile; low volatility describes a state with a single high-fitness residue,
whereas high volatility describes the presence of multiple residues with high fitness. Accordingly,
we propose that co-volatility patterns may capture the associations between the fitness profiles
of the sites. Comparison of co-volatility network structure with structure of the epistasis network
of spike will address this question, and may allow us to identify the adaptive sites required to
facilitate changes at sites that negatively affect virus fitness[35,36].

335   We note that, despite the high predictive performance shown, these studies constitute a
relatively simple framework to understand specific factors associated with the changes in SARS-
CoV-2 spike. A more complete understanding will be generated by incorporating additional
factors, including selective pressures applied on each site, *in vitro* fitness profiles[35], and possibly
patterns of mutations within the host. Furthermore, from a computational perspective, our strategy
can be refined by applying alternative methods to define the architecture of the co-volatility
network and by using more sophisticated learners to combine the volatility-based variables. In
these studies, we have chosen to apply a simple logistic regression model to demonstrate the
predictable nature of the changes. Importantly, the use of more homogenous donor populations,
divided by their infection and vaccination status, will allow us to account for the effects of the
immune response on the evolutionary path of each variant.

346   The predictable nature of the changes in the spike protein suggest that immunogens and
therapeutics can be designed to effectively address future forms expected to dominate in the
population. Advance notice of the imminent changes in each lineage allows testing of their impact
on virus fitness and sensitivity to immune sera[37]. Knowledge of the sites that are not expected to
change is equally important. For example, several mutations in the RBD that affect virus sensitivity
to antibodies, including L452R and T478K were assigned high probabilities to occur from the
baseline group, but low probabilities to occur within lineage B.1.1.7. Similarly, the convergent
N501Y mutation[10] was assigned a high probability by the baseline group but a low probability in
AY.4 (data not shown). Accordingly, such mutations were not observed in sublineages of the
above variants. These findings further support the notion that immunogens should be tailored to
the evolutionary space that is specific to each lineages, which can be inferred at early stages after
it emerges in the population.

METHODS

Sequence alignment

Nucleotide sequences of SARS-CoV-2 isolated from humans were downloaded from the National Center for Biotechnology Information (NCBI) database, the Virus Pathogen Database and Analysis Resource (ViPR) and from the GISAID repository[38]. The following processing steps and analyses were performed within the Galaxy web platform[39]. First, excess bases were trimmed using Cutadapt, using 5'-ATGTTTGTT-3' and 3'-TACACATAA-5 "adapters" that flank the spike gene. Adapter sequences were allowed to match once with a minimum overlap of 5 bases, an error rate of 0.2 with a sequence length between 3,700 and 3,900 bases. All sequences with any nucleotide ambiguities were removed by replacing the non-standard bases with 'N' using snippy-clean_full_aln, followed by filtration of N-containing sequences using Filter FASTA. Sequences that cause frameshift mutations were excluded using Transeq. Nucleotide sequences were aligned by MAFFT, using the FFT-NS-2 method[40]. The aligned sequences were then "compressed" using Unique.seqs to obtain a single representative for each unique nucleotide sequence[41]. Nucleotide sequences were then translated with Transeq and aligned with MAFFT, FFT-NS-2[40]. The first position of each PNGS motif triplet (Asn-X-Ser/Thr, where X is any amino acid except Pro) was assigned a distinct identifier from Asn. All phylogenetic analyses were performed using the full-length spike protein, which include several sequences with amino acid insertions. To maintain consistent numbering of spike positions, all calculations described in this work were performed for the 1,273 positions of the spike protein in the SARS-CoV-2 reference strain (accession number NC_045512).

Phylogenetic tree construction and analyses

A maximum-likelihood tree was constructed for the aligned compressed nucleotide sequences using the generalized time-reversible model with CAT approximation (GTR-CAT) nucleotide evolution model with FASTTREE [42]. The tree was rooted to the sequence of the SARS-CoV-2 reference strain with MEGAX[43]. To divide the tree into "Groups" of sequences, we used an in-house code in Python (see link to GitHub repository in the Data Availability section). This tool uses the Newick file to divide the dataset into sequence groups with a user-defined genetic distance between their centroids. For all analyses we used a distance of 0.004 nucleotide substitutions per site for group partitioning. Groups that did not contain at least 50 unique sequences were excluded. To discern between baseline groups and terminal groups, we used a distance of 0.0015 nucleotide substitutions per site between each group centroid and the SARS-CoV-2 reference strain.

Calculations of volatility

To calculate volatility of spike positions, we divided all sequences of each group into clusters of 50 sequences. Sequence variability in each cluster was quantified using two approaches. To calculate volatility (V) values, we used a binary approach, whereby each position in a 50-sequence cluster was assigned a value of 1 if it contains any diversity in amino acid sequence, or a value of 0 if all sequences in the cluster contain the same amino acid. Thus, each cluster is assigned a 1,273-feature vector that describes the absence or presence of volatility at each position of spike. Volatility was then calculated by averaging values by position across all clusters. For calculations of D or R values for each position i, we used a quantitative approach to define volatility at positions associated with i (i.e., at positions s and q in **Equation 1** and **Equation 2**, respectively). Briefly, sequence variability within each cluster was measured by assigning distinct hydropathy scores to each amino acid according to a modified Black and Mould scale[44]. The Asn residue of PNGS motifs and deletions were also assigned unique values. The values assigned were: PNGS, 0; Arg, 0.167; Asp, 0.19; Glu, 0.203; His, 0.304; Asn, 0.363; Gln, 0.376; Lys, 0.403; Ser, 0.466; Thr, 0.542; Gly, 0.584; Ala, 0.68; Cys, 0.733; Pro, 0.759; Met, 0.782; Val, 0.854; Trp, 0.898; Tyr, 0.9; Leu, 0.953; Ile, 0.958; Phe, 1; deletion site, 1.5. Variability in each cluster was calculated as the standard deviation in hydropathy values among the 50 sequences, and variability values of all clusters were averaged to obtain the volatility value for each position s or q (i.e., V_s or V_q, respectively).

Lineage specificity of volatility patterns

To determine relationships between volatility profiles of spike in the diverse lineages, we partitioned each lineage into 10-cluster groups (500 sequences). All sublineages with Pango designations and all 50-sequence clusters with a dominant non-lineage-ancestral residue at any spike position were removed from the datasets. Within each group, the absence or presence of amino acid variability at each spike position was determined. All groups were thus assigned 1273-bit strings that describes the absence (0) or presence (1) of volatility at each position of the protein. Jaccard distances between the strings were calculated and all groups compared using the Unweighted Pair Group Method with Arithmetic Mean (UPGMA) method[19,20]. The output (in Newick format) was used to generate a dendrogram plot with MEGAX.

To determine the lineage specificity of the volatility patterns, we used a modification of an approach we previously described[21]. Briefly, each SARS-CoV-2 lineage was divided into groups of 10 clusters (500 sequences). Volatility in the groups was calculated for all positions of spike, and each group assigned a 1273-feature vector that describes the level of volatility at all positions

of spike. To compare the vectors, we first calculated for each lineage L the coordinates of the

centroid (C_L) among vectors from the same lineage. The mean intra-lineage distance

($d_{intralineage}$) was calculated as the average Euclidean distance between the lineage centroid C_L

and all groups from the same lineage G_L, formally $\overline{dist(C_L, G_L)}$. In addition, we calculated the mean

inter-lineage distance ($d_{interlineage}$) as the average Euclidean distance between the centroid of

lineage L and all other lineage centroids $\overline{dist(C_L, C_L)}$ ∀L'≠L. We define the ratio as:

$ratio = \frac{d_{intralineage}}{d_{interlineage}}$. The baseline ratio (S$_{base}$) was calculated as the *ratio* using the non-permuted

data. Under the null assumption concerning the evolution of volatility profiles, the intra-lineage

distances are expected to be comparable to the inter-lineage distances, while under the lineage-

specific alternative, we expect clustering of volatility profiles within each lineage even across

distinct 10-cluster groups. To test this, lineage identifiers were permuted and randomly assigned

to each group, from which the permuted ratio (S$_{rand}$) was calculated. The permutation process

was repeated 10,000 times. The P value was calculated as the fraction of the 10,000 tests that

S$_{rand}$ was smaller than S$_{base}$.

Co-volatility network construction

To determine the co-volatility of any two spike positions, we generated a matrix that

contains binary volatility values in all clusters of the tested group (rows) for all 1,273 spike

positions (columns). The co-occurrence of a volatile state between any two spike positions was

calculated using Fisher's exact test and the associated P-value determined using a custom Java

script. To construct the network of co-volatility, we used as input the matrix that describes the -

log$_{10}$(P-value) between the volatility profiles of any two spike positions, whereby nodes are the

positions of spike and the edges that connect them reflect the P-values of their association.

Network structure was visualized using the open-source software Gephi[45]. Networks were

generated using different P-value thresholds (i.e., an edge was assigned only if the P-value was

lower than 0.1, 0.05 or 0.01). To determine robustness of network structure, we randomly deleted

10, 20 or 30 percent of all edges for each of the networks, and network topological properties

were computed using the Cytoscape Network Analyzer tool[46]. Two metrics were calculated for

the complete and depleted networks: **(i)** Degree distribution, and **(ii)** Closeness centrality[23].

Calculations of positive selection

We estimated for each codon of spike the number of inferred synonymous (S) and nonsynonymous (N) substitutions using the GALAXY platform [47]. The input phylogenetic tree was constructed using FASTTREE. The dN-dS metric was used to detect codons under positive selection, where dS is the number of synonymous substitutions per site and dN is the number of nonsynonymous substitutions per site. dN-dS values were normalized using the expected number of substitutions per site. Maximum Likelihood computations of dN and dS were conducted using the HyPhy-SLAC software package [48].

Spatial clustering of volatility and calculations of the variable *D*

We performed a permutation test to determine the spatial clustering of volatile sites around each spike position. To this end, for each position i, we identified the 10 closest positions on the trimer, using coordinates of the cryo-EM structure of the cleavage-positive spike (PDB ID 6ZGI) [49]. We then calculated for each position i the statistic T_i^0:

$$T_i^0 = \sum_{j \in \varphi_i} V_i^0 * V_j^0 \qquad \qquad \textbf{[3]}$$

where V_i^0 describes the volatility at position i, V_j^0 is the volatility at the j[th] neighboring position to i, and φ_i denotes the positions numbers of the 10 closest neighbors to position i. We then permuted all positions identifiers other than *p* and calculated the statistic T_i^k:

$$T_i^k = \sum_{j \in \varphi_i} V_i^0 * V_j^k \qquad \qquad \textbf{[4]}$$

where V_j^k is the volatility at the j[th] adjacent position in the k^{th} permutation (k=*1,2, … 5,000*).

Under the null hypothesis of no spatial clustering, we would expect the neighbor labels to be arbitrary. We therefore test this null hypothesis by estimating the probability of observing a positive departure from the null distribution via:

$$P = \frac{\sum_{k=1}^{N} I_{\{T_i^k \geq T_i^0\}}}{N} \qquad \qquad \textbf{[5]}$$

where N is the total number of permutations (5,000) and I is the indicator function. Therefore, the P-value quantifies the fraction of times the volatility of the surrounding residues is larger for the permuted values relative to the non-permuted values.

To calculate D, we measured for each position i the total volatility at all sites that are within
a distance of 6 Å on the spike trimer structure. The coordinates of the cryo-electron microscopy
structure of the cleaved spike protein in the closed conformation (PDB ID 6ZGI) were used[49].
Coordinates of all atoms were included; N-acetyl-glucosamine atoms were assigned the same
position number as their associated Asn residues. We note that the 6ZGI structure is missing the
following spike residues (numbered according to the SARS-CoV-2 reference strain): 1-13, 71-75,
618-632, 677-688, 941-943 and 1146-1273. To calculate D values for these positions, we applied
the volatility values of the positions immediately adjacent on the linear sequence of spike (i.e.,
positions -1 and +1).

Combined model to predict emergence of dominant-group and subgroup-emerging mutations

To assign a probability for each position to emerge with a mutation, we used a logistic
regression model that applies V, D and R values. The model was trained using V, D and R values
calculated using the 5,700 sequences of the baseline group, with the positive outcome being the
43 GDM and 16 sGEM sites described in **Supplementary Table 1**. To this end, we first created
interaction terms between the initial predictors (i.e., V, D and R). To address the class imbalance
in our datasets (59 of the 1,273 spike positions appeared with LFMs or sLFMs) we used the
adaptive synthetic sampling approach (ADASYN)[50]. Nested cross-validation was used to tune the
model while estimating the metrics of interest. This procedure was also used to generate the
prediction probabilities for each position. Five folds were used for both the inner and outer parts
of the nested cross-validation. Grid search was utilized to optimize hyperparameters with the area
under the receiver operating characteristic curve as the objective for optimization. The model-
specific parameters that we incorporated into the hyperparameter tuning procedure are the
inverse of the regularization strength C and the penalty type. For this purpose, we used a set of
values from 0.001 to 100 for parameter C, and for penalization we used L1 norm, L2 norm, elastic
net, or no penalty in the parameter space. Since we used ADASYN to handle the class imbalance,
we also added the number of positions with similar feature values as another hyperparameter to
the search grid. The number of positions with similar feature values was set between 5 and 45.
As classification metrics, we used sensitivity, specificity, precision, recall, AUC and balanced
accuracy. The balanced accuracy metric, which is the average of sensitivity and specificity, was
used due to the relative imbalance in the datasets.

DATA AVAILABILITY STATEMENT

All IDs of the sequences used in our analyses are available on the Mendeley data repository at doi: 10.17632/wn7jwk9n22.1. Additional data related to this manuscript are available from the corresponding author upon request.

The custom code used in our studies is publicly available within the following hub repository: https://github.com/RoberthAnthonyRojasChavez/SARS2-Volatility. Source code for calculating lineage specificity can be found at https://github.com/haimlab/HIV.

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no conflicts of interest with the contents of this article.

ACKNOWLEDGEMENTS

We are grateful to Dr. Wendy Maury and Dr. Stanley Perlman for critical reading of this manuscript. We are also grateful to Dr. Benjamin Darbro for helpful discussions. We extend our thanks to the GISAID consortium, NCBI Virus, ViPR and all laboratories that publicly summited their SARS-Cov-2 sequences. This work was supported by intramural funds to HH, by grant 110028-67-RGRL to HH from the American Foundation for AIDS Research (amfAR), and by National Institutes of Health grant 1DP2AI164325 to JD.

REFERENCES

1. Cutler DM, Summers LH. The COVID-19 Pandemic and the $16 Trillion Virus. *JAMA* **324**, 1495-1496 (2020).

2. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis* **20**, 533-534 (2020).

3. Tao K*, et al.* The biological and clinical significance of emerging SARS-CoV-2 variants. *Nat Rev Genet* **22**, 757-773 (2021).

4. Tian X*, et al.* Potent binding of 2019 novel coronavirus spike protein by a SARS coronavirus-specific human monoclonal antibody. *Emerg Microbes Infect* **9**, 382-385 (2020).

5. Yi H*, et al.* The Emergence and Spread of Novel SARS-CoV-2 Variants. *Front Public Health* **9**, 696664 (2021).

6. Dai L, Gao GF. Viral targets for vaccines against COVID-19. *Nat Rev Immunol* **21**, 73-82 (2021).

7. Harvey WT*, et al.* SARS-CoV-2 variants, spike mutations and immune escape. *Nat Rev Microbiol* **19**, 409-424 (2021).

8. Focosi D, Maggi F, McConnell S, Casadevall A. Spike mutations in SARS-CoV-2 AY sublineages of the Delta variant of concern: implications for the future of the pandemic. *Future Microbiol* **17**, 219-221 (2022).

9. Saunders N*, et al.* Fusogenicity and neutralization sensitivity of the SARS-CoV-2 Delta sublineage AY.4.2. *EBioMedicine* **77**, 103934 (2022).

10. Martin DP*, et al.* The emergence and ongoing convergent evolution of the SARS-CoV-2 N501Y lineages. *Cell* **184**, 5189-5200 e5187 (2021).

11. Kistler KE, Huddleston J, Bedford T. Rapid and parallel adaptive mutations in spike S1 drive clade success in SARS-CoV-2. *bioRxiv*, (2022).

12. Meng B*, et al.* Recurrent emergence of SARS-CoV-2 spike deletion H69/V70 and its role in the Alpha variant B.1.1.7. *Cell Rep* **35**, 109292 (2021).

13. Nikolaidis M, Papakyriakou A, Chlichlia K, Markoulatos P, Oliver SG, Amoutzias GD. Comparative Analysis of SARS-CoV-2 Variants of Concern, Including Omicron, Highlights

Their Common and Distinctive Amino Acid Substitution Patterns, Especially at the Spike
ORF. *Viruses* **14**, (2022).

14.   Korber B*, et al.* Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases
Infectivity of the COVID-19 Virus. *Cell* **182**, 812-827 e819 (2020).

15.   Dearlove B*, et al.* A SARS-CoV-2 vaccine candidate would likely match all currently
circulating variants. *Proc Natl Acad Sci U S A* **117**, 23652-23662 (2020).

16.   MacLean OA*, et al.* Natural selection in the evolution of SARS-CoV-2 in bats created a
generalist virus and highly capable human pathogen. *PLoS Biol* **19**, e3001115 (2021).

17.   O'Toole A*, et al.* Assignment of epidemiological lineages in an emerging pandemic using
the pangolin tool. *Virus Evol* **7**, veab064 (2021).

18.   Wu F*, et al.* A new coronavirus associated with human respiratory disease in China.
Nature **579**, 265-269 (2020).

19.   Jaccard P. Nouvelles recherches sur la distribution florale. *Bull Soc Vaud Sci Nat* **44**, 223-
270 (1908).

20.   Sokal RR. A statistical method for evaluating systematic relationships. *Univ Kansas, Sci
Bull* **38**, 1409-1438 (1958).

21.   Han C*, et al.* Key Positions of HIV-1 Env and Signatures of Vaccine Efficacy Show Gradual
Reduction of Population Founder Effects at the Clade and Regional Levels. *mBio* **11**,
(2020).

22.   Fili M, Hu G, Han C, Kort A, Trettin J, Haim H. A New Classification Method Based on
Dynamic Ensemble Selection and its Application to Predict Variance Patterns in HIV-1
Env. *bioRxiv*, 2022.2001.2031.478521 (2022).

23.   Barabasi AL, Albert R. Emergence of scaling in random networks. *Science* **286**, 509-512
(1999).

24.   Albert R, Jeong H, Barabasi AL. Error and attack tolerance of complex networks. *Nature*
406, 378-382 (2000).

25.   Araf Y*, et al.* Omicron variant of SARS-CoV-2: Genomics, transmissibility, and responses
to current COVID-19 vaccines. *J Med Virol* **94**, 1825-1832 (2022).

26.   Cao Y*, et al.* Omicron escapes the majority of existing SARS-CoV-2 neutralizing
antibodies. *Nature* **602**, 657-663 (2022).

27.   Ai J*, et al.* Omicron variant showed lower neutralizing sensitivity than other SARS-CoV-2
variants to immune sera elicited by vaccines after boost. *Emerg Microbes Infect*, 1-24
(2021).

28.   Dejnirattisai W*, et al.* Reduced neutralisation of SARS-CoV-2 omicron B.1.1.529 variant
by post-immunisation serum. *Lancet*, (2021).

29.   Fang Z*, et al.* Omicron-specific mRNA vaccination alone and as a heterologous booster
against SARS-CoV-2. *Nat Commun* **13**, 3250 (2022).

30.   Gagne M*, et al.* mRNA-1273 or mRNA-Omicron boost in vaccinated macaques elicits
similar B cell expansion, neutralizing responses, and protection from Omicron. *Cell* **185**,
1556-1571 e1518 (2022).

31.   Qu L*, et al.* Circular RNA vaccines against SARS-CoV-2 and emerging variants. *Cell* **185**,
1728-1744 e1716 (2022).

32.   Maher MC*, et al.* Predicting the mutational drivers of future SARS-CoV-2 variants of
concern. *Sci Transl Med* **14**, eabk3445 (2022).

33.   Geller R, Domingo-Calap P, Cuevas JM, Rossolillo P, Negroni M, Sanjuan R. The external
domains of the HIV-1 envelope are a mutational cold spot. *Nat Commun* **6**, 8571 (2015).

34.   Vandelli A*, et al.* Structural analysis of SARS-CoV-2 genome and predictions of the human
interactome. *Nucleic Acids Res* **48**, 11270-11283 (2020).

35.   Starr TN*, et al.* Deep Mutational Scanning of SARS-CoV-2 Receptor Binding Domain
Reveals Constraints on Folding and ACE2 Binding. *Cell* **182**, 1295-1310 e1220 (2020).

36.   Rodriguez-Rivas J, Croce G, Muscat M, Weigt M. Epistatic models predict mutable sites
in SARS-CoV-2 proteins and epitopes. *Proc Natl Acad Sci U S A* **119**, (2022).

37.   DeGrace MM*, et al.* Defining the risk of SARS-CoV-2 variants on immune protection.
Nature **605**, 640-652 (2022).

38.   Shu Y, McCauley J. GISAID: Global initiative on sharing all influenza data - from vision to
reality. *Euro Surveill* **22**, (2017).

39.   Afgan E*, et al.* The Galaxy platform for accessible, reproducible and collaborative
biomedical analyses: 2018 update. *Nucleic Acids Res* **46**, W537-W544 (2018).

40.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**, 772-780 (2013).

41.   Schloss PD*, et al.* Introducing mothur: open-source, platform-independent, community-
supported software for describing and comparing microbial communities. *Appl Environ*
Microbiol **75**, 7537-7541 (2009).

42.   Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood trees for
large alignments. *PLoS One* **5**, e9490 (2010).

43.   Stecher G, Tamura K, Kumar S. Molecular Evolutionary Genetics Analysis (MEGA) for
macOS. *Mol Biol Evol* **37**, 1237-1239 (2020).

44.   DeLeon O*, et al.* Accurate predictions of population-level changes in sequence and
structural properties of HIV-1 Env using a volatility-controlled diffusion model. *PLoS Biol*
15, e2001549 (2017).

45.   Jacomy M, Venturini T, Heymann S, Bastian M. ForceAtlas2, a continuous graph layout
algorithm for handy network visualization designed for the Gephi software. *PLoS One* **9**,
e98679 (2014).

46.   Lotia S, Montojo J, Dong Y, Bader GD, Pico AR. Cytoscape app store. *Bioinformatics* **29**,
1350-1351 (2013).

47.   Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis
Version 7.0 for Bigger Datasets. *Mol Biol Evol* **33**, 1870-1874 (2016).

48.   Pond SL, Frost SD, Muse SV. HyPhy: hypothesis testing using phylogenies.
Bioinformatics **21**, 676-679 (2005).

49.   Wrobel AG*, et al.* SARS-CoV-2 and bat RaTG13 spike glycoprotein structures inform on
virus evolution and furin-cleavage effects. *Nat Struct Mol Biol* **27**, 763-767 (2020).

50.   He HB, Bai Y, Garcia EA, Li ST. ADASYN: Adaptive Synthetic Sampling Approach for
Imbalanced Learning. *Ieee Ijcnn*, 1322-1328 (2008).

Figure 1. The mutational space of the spike protein is specific for SARS-CoV-2 lineage. (a) Sites of mutation that define sublineages of the indicated VOCs are shown by black dots. Shaded cells indicate sites under positive selection pressure in the VOCs. **(b)** Phylogenetic tree based on 40,350 unique spike sequences from the indicated lineages. The baseline (BL) group is defined as sequences within a distance of 0.0015 nucleotide substitutions per site from the SARS-CoV-2 ancestral strain. **(c)** Schematic of our approach to calculate volatility for each position of spike. **(d)** Volatility calculated at RBD positions using sequences from the indicated VOCs or the baseline group. **(e)** Each lineage was partitioned into clusters of 500 sequences. The absence or presence of volatility in each cluster was determined, and each assigned a 1273-bit string that describes the volatility at all spike positions. Strings were compared using the UPGMA clustering method. **(f)** Euclidean distances between the strings were calculated and the centroid of each lineage determined. A permutation test was used to compare between the mean in-lineage versus between-lineage distances. P-values: *, P<0.05; **, P<0.005; ***, P<0.0005. **(g)** Volatility was calculated in each lineage at all positions of the NTD (20-286), RBD (333-527) and S2 (686-1213). The Spearman correlation coefficient between volatility values in any two lineages was calculated. Coefficients are compared with the mean genetic distance that separates any two lineages. r_S, Spearman coefficient. P-values, two-tailed test.

Figure 2. Spike positions with high volatility or a volatile environment emerge as sites of lineage-founder mutations. (a) Phylogenetic tree based on 16,808 unique spike sequences. Terminal groups are colored and labeled, with their WHO variant designations in parentheses. **(b)** Volatility values for all positions of spike subunit S1 calculated using the 114 baseline clusters (see values for subunit S2 in **Supplementary Fig. 2b**). **(c)** Comparison of volatility values between spike positions that emerged with LFMs, sLFMs or no such mutations. P-values in an unpaired T test: ***, P<0.0005; ****, P<0.00005; ns, not significant. **(d)** Number of positions that appeared with LFMs and sLFMs when volatility (V) in the baseline group was zero or larger than zero. The number of positions in each subset (n) is indicated. **(e)** Mapping of the volatility values calculated in the baseline group onto the spike trimer (PDB ID 6ZGI). **(f)** Results of a permutation test to identify positions with high volatility at their 10 closest neighbors on the trimer structure. Low P-values indicate a high-volatility environment. **(g)** The variable D describes for each position the total distance-weighted volatility at adjacent positions on the spike trimer. D values are compared between positions with LFMs, sLFMs or no such mutations. **(h)** The number of positions that emerged with LFMs or sLFMs when the D value was zero or larger than zero.

Figure 3. High volatility at co-mutable sites is associated with emergence of LFMs and sLFMs. (a) Schematic of our approach to calculate co-volatility of spike positions. For all positions, the absence (0) or presence (1) of amino acid variability was determined in each cluster of 50 sequences. The co-occurrence of a volatile state at all position pairs was determined using Fisher's test, and the P-values were used to construct the network of co-volatility between all positions. **(b)** The co-volatility network around position 614 as the root node. Edges were assigned to positions pairs if the P-value was smaller than 0.05. First- and second-degree nodes are shown. Node size corresponds to the number of triangle counts for each position. **(c)** Network robustness analyses. Networks were constructed using P-value thresholds of <0.01, <0.05 or <0.1. For each network, we randomly deleted 10%, 20% or 30% of edges and examined the effect on network stability. The degree distribution (i.e., the number of nodes associated with each position) is shown for the intact and depleted networks. **(d)** R values, which describe for each position the total weighted volatility at network-associated positions was calculated. R values for spike positions that emerged with LFMs, sLFMs or with no such mutations are shown. **(e)** Number of LFMs and sLFMs that emerged at spike positions when R in the baseline was equal to zero or greater than zero. **(f-h)** Correlations between V, D and R values calculated using the baseline sequences. r_s, Spearman coefficient. P-value, two-tailed test. **(i)** Classification metrics for evaluating performance of V, D and R to predict presence of s/LFMs. Error bars, standard errors of the means for five-fold cross validation.

Figure 4. Volatility patterns among early-pandemic viruses predict emergence of mutations during the lineage-emerging phase. (a) Timeline for emergence of SARS-CoV-2 lineages until July 2021. Lineage emergence time is defined as the date by which 26 sequences that contain all lineage mutations were identified. Lineages with WHO variant designations are indicated by their symbols (see list in Supplementary **Table 2**) and the number of LFMs in each is shown by dots. **(b)** V, D and R values were calculated for all spike positions using the early phase sequences and applied to a logistic regression model to predict emergence of LFMs. Datapoints describe probabilities assigned to all spike positions and are grouped by the lineage in which they emerged as LFMs. Values are compared between the LFM sites in the indicated VOCs (or minor lineages, labeled "Other Lin.") and the no-mutation ("No mut") sites. Significance of the differences was calculated using an unpaired T test: *, P<0.05; **, P<0.005; ***, P<0.0005; ****, P<0.00005; ns, not significant. **(c-e)** V, D or R values calculated for all spike positions using the early phase sequences are grouped as described in panel b. **(f)** V, D and R values and the combined probability were calculated using sequences from different time points of the early phase. AUC values are shown for each time point, for predicting emergence of the 67 LFMs in the lineage-emerging phase. The number of spike sequences at each time point is indicated. **(g)** LFM sites were grouped by the emergence time of the first lineage that contains them. Mutation probabilities assigned to the sites by sequences collected until April 1st 2020 are compared with the probabilities assigned to the no-mutation sites. **(h)** Probabilities assigned by the April 1st 2020 dataset are shown for LFM sites that appeared in one or more lineages. Values are compared between all groups using an unpaired T test.

Figure 5. The mutational patterns of new SARS-CoV-2 sublineages are accurately captured by volatility profiles in their ancestral lineages. (a) Sequences from the baseline of the indicated VOCs were used to calculate the combined probability for mutations at all spike positions. Values were compared with two outcomes: (i) Emergence of a mutation defines a new Pango sublineage (in red), or (ii) Presence within the lineage baseline of two or more 50-sequence clusters with a dominant non-lineage-ancestral residue (in blue). The number of events detected for each outcome (n) and the AUC values are shown. **(b)** Mutation probabilities calculated using the sequences of each lineage are compared with the outcomes observed in all lineages. **(c)** AY.4 sequences were indexed by sample collection date. The cumulative probabilities for the 14 sites that emerged as sublineages of AY.4 are shown. Circles indicate the time of sublineage emergence. Filled circles indicate that the site reached a probability of 0.99. The blue line indicates for each time point the number of sites that reached a probability of 0.99 **(d)** Weblogos describe the frequencies of minority variants at sites that appeared in the VOC sublineages by April 8th 2022. Frequencies are expressed as a percent of all sequences with a non-lineage-ancestral residue. The residues that emerged in the sublineages are shown in red font. The residue change in each lineage from the SARS-CoV-2 ancestor is shown below the axis.

Supplementary Figure 1. (a) Volatility calculated for all positions of spike using sequences from the indicated lineages or the SARS-CoV-2 baseline (BL) group. Data represent sequences from samples collected worldwide until the following dates: Baseline, until July 2021; P.1, AY.4, AY.4.2 and BA.1, until February 2022; B.1.1.7, BA.1.1 and BA.2, until March 2022. Volatility values are color-coded as indicated. **(b)** Volatilities are compared between lineages for the five positions with the highest values in each lineage.

Supplementary Figure 2. A high level of volatility at spike positions is associated with emergence of founder mutations in SARS-CoV-2 lineages. (a) Phylogenetic tree constructed from 16,808 unique spike sequences from samples collected worldwide until July 2021. Examples are shown of lineage-founder and sublineage-founder mutations in spike. (Left) Branches are colored by the amino acids that occupy spike position 501. The pattern corresponds to presence of a group-dominant mutation in $G_{T1}(\alpha)$, $G_{T6}(\gamma)$ and G_{T8}. (Right) Branches are colored by the amino acids that occupy position 1191, showing a sublineage-emerging mutation in $G_{T1}(\alpha)$. **(b)** Volatility of spike positions of the S2 subunit, as calculated using the baseline group of 5,700 sequences (114 clusters). Red bars indicate positions with lineage-founder mutations (in any terminal or baseline group). Blue bars indicate positions with sublineage-founder mutations. FP, fusion peptide; HR, heptad repeats; TM, transmembrane domain; CT, cytoplasmic tail. **(c)** Thirty spike positions with the highest volatility values. The baseline ("B") or terminal ("T") groups that contain mutations at these positions are indicated.

Supplementary Figure 3. Structural properties of the network of co-volatile sites. The co-occurrence of a volatile state at any two spike positions was determined using the 114 clusters of the baseline group. **(a)** Histogram showing the distribution of P-values calculated using Fisher's exact test for co-volatility of any two spike positions. **(b)** Networks of co-volatility between all spike positions were constructed using P-value thresholds of <0.01, <0.05 or <0.1. For each of the three networks, we randomly deleted 10%, 20% or 30% of edges and examined the effect on network stabilities. Closeness centrality values are shown for the intact and depleted networks. Higher values indicate shorter distances to all other nodes. Bars indicate the second and third quartiles and whiskers indicate minimum and maximum values.

Supplementary Figure 4. Volatility profiles among isolates from the early pandemic predict emergence of lineage-founder mutations at later stages. (a) Sequences from samples collected until the indicated time points of the early phase were used to calculate *V*, *D* and *R* values for each position. In addition, we calculated the probability for mutations at each spike position using a logistic regression model that applies *V*, *D* and *R* values. Sensitivity and specificity values are shown for prediction of the 67 LFMs that appeared during the lineage-emerging phase (September 2020 to July 2021). AUC values are shown in **Fig. 4f**. **(b)** Classification metrics for evaluating performance of the combined model to predict emergence of LFMs using sequence data collected until April 1st 2020 or all sequences of the early phase. **(c)** Precision of the combined model at different probability percentile thresholds. Probabilities were calculated for all spike positions using the April 1st 2020 dataset or all early-phase sequences. **(d)** LFM sites are grouped by the emergence time of the first lineage that contains them. Mutation probabilities assigned to the sites using sequences collected until September 19th 2020 are shown and compared with the probabilities assigned to the no-mutation sites. **(e)** Comparison of probabilities assigned to LFM sites that appeared in one or more lineages. Probabilities assigned to positions in each group are compared between all groups using an unpaired T test: *, P<0.05, ***, P<0.0005; ns, not significant.

Supplementary Figure 5. Evolution of *V*, *D* and *R* values and the combined mutation probability during the early phase of the pandemic. (a,b) Sequences from the early phase (December 30[th] 2019 to September 19[th] 2020) were divided into 36 clusters of 50 sequences. *V*, *D* and *R* values were calculated using sequences isolated from samples collected until the indicated time points. In addition, we calculated the probability for emergence of a mutation at each spike position using a logistic regression model that applies the three variables. Values for positions that emerged as LFMs in lineages B.1.1.7 and B.1.617.2 are shown. The shaded area describes probability values below the 90[th] percentile calculated for each time point.

Supplementary Table 1: Founder mutations in SARS-CoV-2 lineages and sublineages.

Group type	Group [a,b]	WHO variant	Pango lineage [c]	# of unique sequences [d]	Lineage-founder mutations [e,f]	Sublineage-founder mutations [g,h]
Terminal (T)	$G_{T1}(\alpha)$	Alpha	B.1.1.7	4,714	H69del, V70del, Y144del, N501Y, A570D, D614G, P681H, T716I, S982A, D1118H	L5F, S98F, D138H, D178H, V327I, K1191N
	$G_{T2}(\varepsilon)$	Epsilon	B.1.429	874	S13I, W152C, L452R, D614G	P26S, W258L
	$G_{T3}(\delta)$	Delta	B.1.617.2	674 (4,283)	PNGS17N, T19R, G142D, E156del, F157del, R158G, L452R,T478K, D614G, P681R, D950N	K77T, T95I, A222V, V1264L (**L5F, V70I, K97E, S112L, R214H, V289I, N1074S, V1104L**)
	$G_{T4}(\iota^3)$	Iota	B.1.526	664	L5F, T95I, D253G, E484K, D614G, A701V	S477N
	$G_{T5}(\iota^1)$	Iota	B.1.526, D2, B.1.160	598	L5F, T95I, D253G, S477N, D614G, Q957R	-
	$G_{T6}(\gamma)$	Gamma	P.1	430	L18F, T20N, P26S, D138Y, N188PNGS, R190S, K417T, E484K, N501Y, D614G, H655Y, T1027I, V1176F	-
	$G_{T7}(\iota^3)$	Iota	B.1.526	337	D80G, Y144del, F157S, L452R, D614G, T859N, D950H	T791I
	G_{T8}	-	B.1.623	53	S494P, N501Y, D614G, P681H, K854N, E1111K	-
Baseline (B)	G_{B1}	-	B.1.2	2,204	D614G	H69del, V70del, G72R, G142del, V143del, N439K, Q675H, P681H, V1228L
	G_{B2}	-	B.1.177, B.1.2	2,181	A222V, D614G	L18F, A262S, P272L, T716I
	G_{B3}	-	B.1.575, B.1.1.519	559	T478K, D614G, P681H, T732A	S494P, T716I
	G_{B4}	-	B.1.2, B.1.1.239	354	D614G, Q677H	-
	G_{B5}	-	B.1.2	173	D614G	-
	G_{B6}	-	B.1.2	166	D614G	-
	G_{B7}	-	B.1.596	125	D614G, Q677P	-
	G_{B8}	-	B.1.2	84	L5F, D614G	-
	G_{B9}	-	R.1	80	W152L, E484K, D614G, G769V	-
	G_{B10}	-	C.30, C.16, B.1.1.1	80	D614G	-
	G_{B11}	-	B.1.2	61	D614G	-
	G_{B12}	-	P.2	60	E484K, D614G, V1176F	-

[a] Grouping is based on phylogenetic analysis of 16,808 unique nucleotide sequences of spike isolated from samples collected worldwide between December 2019 and July 2021.

[b] Groups were assigned to the baseline set of sequences (G_B) if their centroid was located 0.0015 or less nucleotide substitutions per site from the reference spike sequence (accession number NC_045512).

[c] Only Pango linages that represent 10 percent or more of sequences within a group are listed.

[d] The number of $G_{T3}(\delta)$ sequences isolated from samples collected between December 2019 and September 2021 is indicated in parentheses.

[e] A mutation is defined as group-dominant if it is found in the inferred ancestral sequence of the group and in more than 50 percent of group sequences.

[f] PNGS indicates presence of Asn at the first position of a PNGS triplet where the third position is occupied by Thr or Ser and the second position is not occupied by Pro.

[g] A mutation is defined as subgroup-emerging if it is not found in the inferred group ancestor and is the dominant residue in at least one group cluster but less than 50% of all group clusters.

[h] Sublineage-founder mutation that appeared in $G_{T3}(\delta)$ between July 2021 and September 2021 are indicated in parentheses and in bold font.

Supplementary Table 2: SARS-CoV-2 lineages that emerged until July 2021 and the mutations they contain in the spike protein

Phase of pandemic	Pango Lineage [a]	Date of emergance [b]	Mutation (Probability) [c,d,e]
Early phase	A.1	03/19/2020	
	B.1	03/27/2020	614
	B.1.1	04/30/2020	614
	D.2	07/28/2020	477, 614
	B.1.177	09/02/2020	222, 614
	B.1.1.37	09/10/2020	614
Lineage-emerging phase	B.1.1.1	10/10/2020	614
	AD.2	10/12/2020	614
	B.1.2	10/14/2020	614
	B.1.1.311	10/14/2020	614
	B.1.243	10/15/2020	614
	B.1.258	10/29/2020	614, 439 (0.995)
	B.1.36.17	10/30/2020	614
	B.1.177.4	11/05/2020	614, 222
	B.1.177.57	11/05/2020	614, 222
	B.1.240	11/06/2020	614
	B.1.160	11/23/2020	477, 614
	B.1.1.7 (Alpha)	12/10/2020	69 (1.0), 70 (1.0), 144 (1.0), 501 (1.0), 570 (0.422), 614, 681 (1.0), 716 (0.986), 982 (0.299), 1118 (0.862)
	B.1.36	12/16/2020	614
	B.1.177.87	12/17/2020	222, 262 (1.0), 272 (0.879), 614
	B.1.369	12/20/2020	614
	B.1.36.8	12/29/2020	614
	B.1.177.17	12/29/2020	176 (0.999), 222, 614
	B.1.1.222	01/01/2021	732 (1.0), 614
	B.1.234	01/02/2021	614
	B.1.429 (Epsilon)	01/07/2021	13 (0.982), 152 (1.0), 452 (0.989), 614
	B.1.221	01/08/2021	98 (1.0), 614
	B.1.596	01/09/2021	614
	B.1.400	01/17/2021	614
	B.1.177.44	01/20/2021	222, 614
	B.1.311	01/25/2021	614
	B.1.595	01/29/2021	614
	B.1.427	01/31/2021	13 (0.982), 152 (1.0), 452 (0.989), 614
	B.1.526 (Iota)	02/08/2021	5 (1.0), 95 (1.0), 253 (1.0), 614
	B.1.1.519	02/10/2021	68 (0.691), 478 (0.900), 614, 732 (1.0)
	B.1.517	02/24/2021	501 (1.0), 614
	B.1.177.7	02/24/2021	18 (1.0), 222, 614
	P.2	03/04/2021	484 (0.993), 614, 1176 (0.663)
	B.1.110.3	03/04/2021	614
	B.1.575	03/08/2021	494 (0.260), 614, 681 (1.0), 716 (0.986)
	B.1.1.434	03/17/2021	614
	B.1.1.316	03/18/2021	614, 677 (1.0), 732 (1.0)
	B.1.609	03/20/2021	614
	R.1	03/23/2021	152 (1.0), 484 (0.993), 614, 769 (0.999)
	B.1.623	04/01/2021	494 (0.260), 501 (1.0), 614, 681 (1.0), 1111 (0.306)
	P.1 (Gamma)	04/07/2021	18 (1.0), 26 (1.0), 20 (0.996), 190 (0.637), 138 (1.0), 417 (0.176), 484 (0.993), 501 (1.0), 614, 655 (1.0), 1027 (0.624), 1176 (0.663)
	B.1.525	04/11/2021	52 (0.520), 67 (0.916), 69 (1.0), 70 (1.0), 144 (1.0), 484 (0.993), 614, 677 (1.0), 888 (0.244)
	B.1.351	04/20/2021	47 (0.263), 50 (0.912), 80 (1.0), 70 (1.0), 215 (0.875), 241 (0.169), 242 (0.979), 243 (0.999), 484 (0.993), 614
	B.1.241	04/22/2021	614
	B.1.617.2 (Delta)	05/05/2021	19 (1.0), 142 (0.998), 156 (0.263), 157 (0.726), 158 (0.618), 452 (0.989), 478 (0.900), 614, 681 (1.0), 950 (0.892)
	C.37	06/30/2021	75 (1.0), 76 (1.0), 246 (0.955), 247 (0.894), 248 (0.387), 249 (0.270), 250 (0.646), 251 (0.268), 252 (0.886), 253 (1.0), 452 (0.989), 490 (0.090), 614, 859 (1.0)
	AY.3	07/12/2021	19 (1.0), 156 (0.263), 157 (0.726), 158 (0.618), 452 (0.989), 478 (0.900), 614, 681 (1.0), 950 (0.892)

[a] Pango lineages associated with the sequences.

[b] The date of lineage emergence is defined as the date by which at least 26 sequences of the indicated lineage were detected.

[c] Spike mutations associated with the Pango lineage based on data published from Outbreak.info, 2021.

[d] Values in parentheses describe the probabilities assigned to each site for a mutation, based on a logistic regression model that combines V, D, and R values (see **Figure 4b**).

[e] Sites of mutations in lineages that formed before September 19th 2020 are shown in red font. These positions were excluded from our time-indexed analyses.

Supplementary Table 3: Mutations in spike that define SARS-CoV-2 sublineages

Lineage	In-lineage mutations [a]	Mutations that define Pango sublineages		
		Sublineage [b]	Mutations in first-order sublineages [c]	Mutations in second-order sublineages [d]
B.1.1.7	5 (>0.999) 98 (0.999) 138 (0.998) 1191 (>0.999)	Q.5	D138H (0.998), F490S (0.906)	
P.1	-	P.1.3	F140- (0.995), L141- (0.995), G142- (0.996), V143- (>0.999) , Y144F (0.999)	
		P.1.4	N679K (0.572)	
		P.1.5	N679K (0.572)	
		P.1.6	P681H (0.551)	
		P.1.7	P681H (0.551)	
		P.1.8	T470N (0.465), P681R (0.551), C1235F (0.168)	
		P.1.9	Q675H (0.998)	
		P.1.10	-	**P.1.10.1**: H69- (0.994), V70- (0.975)
		P.1.11	P681R (0.551)	
		P.1.12	-	**P.1.12.1**: H49Y (0.224)
AY.4	5 (>0.999) 95 (0.993) 138 (>0.999) 142 (>0.999) 1078 (>0.999) 1124 (>0.999) 1141 (>0.999) 1162 (>0.999)	AY.4.1	G446V (0.999)	
		AY.4.2	Y145H (0.978), A222V (0.953)	**AY.4.2.1**: V36F (0.420) **AY.4.2.2**:V1264L (0.973) **AY.4.2.4**: D142**G** (>0.999)
		AY.4.3	P26S (0.894), L54F (0.898)	
		AY.4.4	L5F (>0.999), S929T (0.325)	
		AY.4.5	D1259Y (0.324)	
		AY.4.10	P812S (0.841)	
		AY.4.12	P1162S (>0.999)	
		AY.4.13	V213A (0.673)	
		AY.4.14	A1078S (>0.999)	
		AY.4.15	A1078S (>0.999)	
		AY.4.16	D138H (>0.999), A845S (0.701)	
BA.1	701 (>0.999)	BA.1.1	R346K (0.925)	**BA.1.1.3**: T791I (0.232)
		BA.1.4	N417**K** (0.999), K440**N** (0.798), S446**G** (0.680), V597I (0.321)	
		BA.1.5	A831V (0.382)	
		BA.1.8	P809S (0.338)	
		BA.1.9	N417**K** (0.999), K440**N** (0.798), S446**G** (0.680)	
		BA.1.10	N417**K** (0.999), K440**N** (0.798), S446**G** (0.680)	
		BA.1.12	N417**K** (0.999), K440**N** (0.798)	
		BA.1.15	-	**BA.1.15.1**: I1081V(>0.999) **BA.1.15.2**: V320I(0.252), Q628K (0.253)
		BA.1.16	N417**K** (0.999), K440**N** (0.798), S446**G** (0.680)	
		BA.1.17	A701V (>0.999)	
		BA.1.21	-	**BA.1.21.1**: R493**Q** (0.897), S496**G** (0.887), R498**Q** (0.799), Y501**N** (0.965)
BA.2	-	BA.2.2	I1221T (0.322)	
		BA.2.3	-	**BA.2.3.1**: Q677E (0.341) **BA.2.3.4**: N641S (0.405)
		BA.2.10	-	**BA.2.10.1**: G798D (0.176)
		BA.2.11	L452R (0.657)	
		BA.2.12	S704L (0.585)	**BA.2.12.1**: L452Q (0.657)
		BA.2.13	L452M (0.657)	
		BA.2.14	W64L (0.829)	
		BA.2.15	W64L (0.829)	
		BA.2.20	N148T (0.287)	
		BA.2.22	F186S (0.479)	
		BA.2.24	D339N (0.249)	
		BA.2.35	L452R (0.657)	
		BA.2.36	I68T (0.827)	
		BA.2.38	N417T (0.667)	
		BA.2.39	L858I (0.338)	
		BA.2.40	-	**BA.2.40**: N417T(0.667)
		BA.2.41	T1231S (0.313)	

[a] In-lineage mutations are defined as positions occupied by a non-lineage ancestral residue in two or more 50-sequence clusters of the lineage but do not appear in a Pango-designated sublineage. Probability values assigned to the sites by a model that applies the V, D and R values are indicated in parentheses.

[b] Sublineages that appeared until April 8th, 2022.

[c] Reversions to the SARS-CoV-2 ancestral residues are shown in bold red font.

[d] Unique mutations in the second-order sublineages that appeared until June 17th, 2022 but do not appear in the ancestral (first-order) sublineage are shown.