Genomic signatures of host-specific selection in a parasitic plant

Authors: Emily S. Bellis[1,2,3], Clara S. von Münchow[4], Alan Kronberger[1], Calvins O. Odero[5], Elizabeth A. Kelly[3], Tian Xia[3], Xiuzhen Huang[1,2], Susann Wicke[4, 6], Steven M. Runo[5], Claude W. dePamphilis[3], Jesse R. Lasky[3]

Target journal: AJB

Corresponding Author: Emily S. Bellis, ebellis@astate.edu

[1]Department of Computer Science, Arkansas State University, State University, AR, USA
[2]Center for No-Boundary Thinking, Arkansas State University, State University, AR, USA
[3]Department of Biology, Pennsylvania State University, University Park, PA, USA
[4]Institute for Biology, Humboldt University of Berlin, Berlin, Germany
[5]Department of Biochemistry, Microbiology, and Biotechnology, Kenyatta University, Nairobi, Kenya
[6]Späth-Arboretum of the Humboldt University of Berlin, Berlin, Germany

Running head: *Striga hermonthica* population genomics

ABSTRACT

Premise—Parasitic plants and their hosts are model systems for studying genetic variation in species interactions across environments. The parasitic plant *Striga hermonthica* (witchweed) attacks a range of cereal crop hosts in Africa and exhibits substantial variation in performance on different host species. Some of this variation is due to local adaptation, but the genetic basis of specialization on certain hosts is unknown.

Methods—To identify genomic regions that are strongly differentiated between parasites attacking different host species, we present an alignment-free analysis of *S. hermonthica* population diversity using whole genome sequencing (WGS) data for 68 individuals from western Kenya. We validate our findings with germination experiments and analyses based on a *de novo* assembled draft genome.

Results—Reference-free and reference-based analyses suggest that only a small

portion of the *S. hermonthica* genome is strongly differentiated by host species in populations

from western Kenya. Analysis of host-associated *k*-mers implicated genes involved in

development of the parasite haustorium (a specialized structure used to establish vascular

connections with host roots) and a potential role of chemocyanins in molecular host-parasitic

plant interactions. Conversely, no phenotypic or genomic evidence was observed suggesting

host-specific selection on parasite response to strigolactones, hormones exuded by host roots

and required for parasite germination.

Conclusions—This study demonstrates the utility of WGS for plant species with large,

complex genomes and no available reference. Contrasting with theory emphasizing the role of

early recognition loci for host specificity, our findings support host-specific selection on later

interaction stages, recurring each generation after homogenizing gene flow.

Key words: population genomics, recurrent selective sweeps, agroecosystems,

Orobanchaceae, host-parasite coevolution, chemocyanin

INTRODUCTION

Characterizing the genomic basis of adaptation to local biotic environments is a key challenge

for evolutionary ecology (Ebert and Fields, 2020). Parasites and mutualists exert strong

influences on host fitness, and may even constitute the predominant selective pressure shaping

patterns of local adaptation in some systems (Fumagalli et al., 2011; Castellano et al., 2019).

Many host-parasite and host-mutualist systems involve a complex multi-step infection process

including many stages of interaction between host and symbiont derived molecules (Hall et al.,

2017). An outstanding question is whether adaptation to local biotic environments occurs most

often via selection on genes involved during initial infection stages, or whether genetic variation

at later stages of the interaction is also frequently maintained.

The expectation from theoretical studies is that initial recognition loci are more likely than

downstream effector loci to contribute to host genotype by parasite genotype interactions (G_H x

G_P) and correspondingly, local adaptation (Nuismer and Dybdahl, 2016). The first prediction,

that recognition loci contribute more to G_H x G_P, is supported by empirical studies of the

waterflea *Daphnia magna* and its bacterial parasite *Pasteuria ramosa* (Hall et al., 2017). In this

system, most of the genetic variance in parasite infection was associated with a single major

effect QTL linked to the early stage of parasite attachment (Hall et al., 2019). In contrast, many

different QTLs of smaller effect were associated with later stages, highlighting the potential for

independent evolution of traits involved in different stages of the infection process (Hall et al.,

2019). Supporting the second prediction that selection on recognition traits often underlies local

adaptation (Nuismer and Dybdahl, 2016), studies of plant pathosystems have revealed

reciprocal coevolutionary selection on host resistance (R) genes and parasite avirulence genes,

for example in the flax-flax rust system (Ravensdale et al., 2011; Thrall et al., 2012). However, a

high degree of genotype specificity has also been observed for many host-parasite systems

characterized by more quantitative mechanisms of resistance (Poland et al., 2009). For host-

parasite interactions characterized by quantitative genetic architectures, we still know little

regarding the genetic basis of local adaptation in natural populations and the extent to which

initial vs. later infection stages contribute to G_H x G_P.

An emerging model system for studying spatial pattern and process in coevolutionary

genomics is the parasitic plant *Striga hermonthica* (family Orobanchaceae) and its cereal hosts.

In contrast to *Striga gesnerioides,* which parasitizes cowpea via a qualitative gene-for-gene

mechanism, natural variation in host resistance to *S. hermonthica* is highly polygenic (Li and

Timko, 2009; Timko et al., 2012) with at least one large effect locus (Gobena et al., 2017).

Striga hermonthica parasitizes grass hosts including sorghum, maize, rice, and millets and is

one of the greatest biotic constraints to food security in Africa (Ejeta, 2007; Spallek et al., 2013;

Savary et al., 2019). An individual *S. hermonthica* plant can produce thousands of seeds that

are speculated survive in the soil for a decade or more under optimal conditions (Bebawi *et al.*

1984; but see Gbèhounou *et al.* 2003).

A great deal has been learned in the past decades regarding the mechanisms through

which *Striga* parasitizes diverse hosts. To germinate, parasite seeds must detect strigolactones

(SLs), hormones exuded from host roots under nutrient-deficient conditions that also stimulate

host interactions with beneficial mycorrhizal fungi (Akiyama et al., 2005). Perception of SLs is

mediated through binding to paralogs of KARRIKIN INSENSITIVE 2 (KAI2), known as

HYPOSENSITIVE TO LIGHT (HTL) receptors. SL receptors of the *KAI2d* clade rapidly

expanded and diversified during the transition to parasitism in the Orobanchaceae, a plant

family that includes thousands of mostly root parasitic species (Conn et al., 2015). For example,

the ~600 Mb genome of *Striga asiatica* contains 21 *KAI2* genes, many of which occur as

tandem duplications (Yoshida et al., 2019). *Striga hermonthica* may possess 13 or more *KAI2*

paralogs (Nelson, 2021), including 11 for which the binding affinity for diverse SLs has been

extensively characterized (Toh et al., 2015; Tsuchiya et al., 2015).

Following SL perception and parasite germination, host-derived phenolic compounds

induce formation of the haustorium, the specialized multicellular feeding structure used by

parasitic plants to infest host tissues (Cui et al., 2018). Intrusive cells of the haustoria invade

host tissues to form direct connections with host vasculature (Masumoto et al., 2021). Water,

nutrients, and other molecules including mRNA (Kim et al., 2014), small RNA (Shahid et al.,

2018), DNA (Yang et al., 2019), and proteins (Liu et al., 2020; Shen et al., 2020) can be directly

transferred across haustorial connections. In addition to low germination stimulation (Dayou et

al., 2021; Mallu et al., 2021), post-germination host resistance to *Striga hermonthica* can occur

at later interaction stages through induction of an intense hypersensitive response, formation of

a mechanical barrier, or failure of the parasite to form vascular connections (Mbuvi et al., 2017;

Mutinda et al., 2018; Kavuluko et al., 2020). Much of the genetic variation in these mechanisms

of resistance may result from host populations' local adaptation to parasitism across the range

of *Striga hermonthica* (Bellis et al., 2020).

Conversely, many parasite populations may also be adapted to their local host

communities. Across the geographic range of *S. hermonthica* in Africa, there exists

considerable variation in host crop communities, and regional abundance of a particular host

species is associated with high *Striga* performance on that crop (Bellis et al., 2021). Given the

low germination of millet- and sorghum-specific *S. hermonthica* populations in response to root

exudates from the alternate host (Parker and Reid, 1979), it is possible that at least some of this

host specialization results from natural selection on SL perception. Host-specific germination by

Striga genotypes could potentially result from duplication of different *KAI2* paralogs and

neofunctionalization to fine-tune the response to different SLs, with host community

heterogeneity maintaining presence/absence variation (Nelson, 2021). In contrast, if parasite

local adaptation involves later interaction stages, 'core parasitism genes' implicated in

haustorium development in previous studies (Yang et al., 2014) could be the targets of host-

specific selection.

Extensive genetic and germplasm resources for host species have enabled broad-scale

studies of sorghum local adaptation to *S. hermonthica* parasitism (Bellis et al., 2020), but

understanding the genetic basis of reciprocal adaptation is challenged by a paucity of genomic

data for parasites. At the population-scale, restriction-site associated DNA sequencing (RAD-

Seq) and analysis of polymorphism in transcriptomes have begun to shed light on population

level diversity in *S. hermonthica* parasites (Unachukwu *et al.* 2017; Lopez *et al.* 2019). However,

reduced representation approaches pose difficulties if only a fraction of host-associated genome

diversity is tagged by RAD-Seq markers (Lowry et al., 2017) or (in the case of transcriptomes) if

genes under selection are not expressed in sequenced tissues. Like other parasitic

angiosperms, *S. hermonthica*, a diploid species with n = 19 or 20 (Iwo et al., 1993; Aigbokhan et

al., 1998), is characterized by a larger genome than non-parasitic relatives (Lyko and Wicke,

2021), with estimated genome size of ~1 Gb (1C = 0.9 Gb; Yoshida *et al.* 2010) or greater (1C =

1.4 Gb; Estep *et al.* 2012). Several genome assembly projects are currently underway, but high

heterozygosity, obligate outcrossing, and a large genome pose substantial challenges for

assembly, and a publicly available reference for *S. hermonthica* is not currently available (but

see Qiu et al., 2022 for another genome assembly that may be available soon). Here, we

provide a reference-free analysis of population-scale diversity in *S. hermonthica,* based on

whole genome sequencing (WGS) data. Using a unique alignment-free bioinformatic approach,

we identify genetic variation associated with host-specific parasitism in natural populations and

investigate signatures of selection surrounding candidate loci. Based on these findings, we

evaluate the hypothesis that adaptation to local host populations results from selection on genes

involved in early stages of the interaction (SL perception) against the alternative that selection

on genes involved in later stages is primarily responsible.

MATERIALS AND METHODS

Sample Collection—Seeds and leaf tissue were collected from *S. hermonthica*

individuals in July 2018 from six locations in western Kenya (Fig. 1A), which are location-

matched with previously collected specimens in herbaria. Two plots with *S. hermonthica*

parasitizing different host species were sampled at each location, chosen as close as possible

and in most cases, from nearby plots on the same farm (i.e. within 15 meters). In one location

(Mumias), we were not able to find a parasite population on another host, so two maize-

parasitizing populations were sampled. Per plot, twelve *S. hermonthica* individuals were chosen

haphazardly, with effort taken to sample individuals distributed evenly throughout the plot.

Individuals were photographed before collection, and images for representative individuals at

each site were uploaded to iNaturalist. Three to four leaves per individual were sampled directly

into silica gel, before collection of the whole individual into separate paper bags. Plants were

dried in paper bags before harvesting and manual cleaning of seeds. Cleaned seeds were

shipped to Penn State for germination rate experiments and stored in individual 2 mL

microcentrifuge tubes at room temperature prior to experiments. Five voucher specimens (ESB

collection numbers 2018.1 to 2018.5) were deposited in the collection of the East African

Herbarium (EA).

Whole genome sequencing—For samples collected in 2018, whole genome

sequencing was performed for a subset of 68 *S. hermonthica* individuals. This included all 24

individuals collected from adjacent plots of finger millet and maize in Kisii and all 24 individuals

collected from adjacent plots of sorghum and maize in Homa Bay. We hypothesized that for

each of these locations, parasites sampled from different hosts would show overall high

similarity across much of the genome due to high rates of gene flow among neighboring plots.

However, we expected to identify loci under strong host-specific selection as differentiated

against this backdrop. To interpret patterns of host-specific differentiation for the Homa Bay and

Kisii populations in the context of broader population genetic diversity in western Kenya, we also

sequenced DNA from five individuals (two or three per host species) from four additional

locations. DNA was extracted from silica-dried leaf tissue in the USDA APHIS quarantine facility

at the Pennsylvania State University using the E.Z.N.A. Plant DNA DS Mini Kit (Omega Bio-tek,

Norcross, Georgia, USA) according to the manufacturer's protocol. Genomic library preparation

and paired-end 150 bp sequencing was carried out by the Texas A&M AgriLife Genomics and

Bioinformatics Service, on a single lane of a NovaSeq 6000 S4 flow cell.

Population structure—To determine whether parasites from the same location showed

the expected pattern of low genetic differentiation between different hosts, we followed a

reference-free approach to evaluate population genomic patterns among sequenced samples.

Raw reads that could be classified as plant-derived were identified using Kraken 2 (Wood et al.,

2019), based on a custom database built from the complete set of plant genomes and proteins

in the NCBI RefSeq collection, sequences from 472 Mbp of the *Striga asiatica* genome (Yoshida

et al., 2019) and *S. hermonthica* transcriptome sequences (build StHeBC4) from the Parasitic

Plant Genome Project II (Yang et al., 2014). Classified sequences were trimmed using BBduk

from BBTools (Bushnell, n.d.), removing sequence on the ends of reads with low quality

(qtrim=rl trimq=20 minlen=50) or 3' matches to adapters (k=23 mink=11 hdist=1 tpe tbo ktrim=r).

To reduce bias associated with differences in per-sample read depth, reads were downsampled

to 5.2 Gbp with BBTools Reformat.

Mash, a dimensionality reduction technique based on the MinHash algorithm, was used

to estimate genetic distance between samples based on resulting read sets (Ondov et al.,

2016). *Mash* previously showed improved performance compared to alignment-based methods

for estimating pairwise genetic distance for polyploid plant genomes using simulated and real

data (VanWallendael and Alvarez, 2020). We used a *k*-mer size of 31, removing *k*-mers with

less than 2 copies but increasing the sketch size to 1×10^7 to account for a larger volume of

input data. Principal Coordinates Analysis was performed in R version 4.0 with the pcoa function

of the 'ape' package (Paradis and Schliep, 2019). A shorter *k*-mer size (*k*=21) was also tested

but did not alter clustering patterns in PCoA. Correlation between the genetic distance matrix

and the geographic distance matrices, calculated with 'geodist', was determined using a Mantel

test (Padgham and Sumner, 2021).

If contamination is present in our Kraken2 plant genome database, our reference-free

analysis of population diversity could be biased by contaminant sequences. Therefore, we also

included an analysis of population structure based on alignment to our assembled reference

genome (see "Validation of *k*-mer based approaches"). After removing scaffolds shorter than

500 bp from the reference, high quality sequences from each *S. hermonthica* individual were

mapped using BWA-MEM v0.7.17 (Li, 2013). Sequences mapping with quality less than 20

were excluded using SAMtools v1.10 (Li et al., 2009). Pairwise genetic relationships were

calculated from genotype probabilities using *ngsDist* (Vieira et al., 2016). Genotype posterior

probabilities were assigned using ANGSD v0.935 (Korneliussen et al., 2014), ignoring sites with

a minimum coverage less than 50 or greater than 500 across all 68 individuals and contigs less

than 2.5 kb.

Host-specific differentiation—We next sought to identify particular genomic regions

differentiated between parasites growing on different hosts. This analysis targeted parasite

populations from Homa Bay or Kisii, for which we sequenced DNA from parasites for two

different hosts on the same farm from immediately adjacent plots. For the Kisii population, the

dataset included individuals from finger millet and maize, whereas for the Homa Bay population

the dataset included parasites from sorghum and maize ($n = 12$ from each host species; 48

individuals total). Counts for k-mers of length 31 were summarized across sequenced

individuals using HAWK; 31-mers were used as the longest k-mer that can be represented

efficiently on a 64-bit machine (Rahman et al., 2018). At each k-mer we considered two allelic

states (present or absent) where the k-mer was marked as present in an individual if it was

counted at least twice or absent if it was not observed at all. A single biallelic SNP in otherwise

invariant 31 bp of sequence, for example, would be represented by two distinct 31-mers. Note

that k-mers which only appear once in samples are filtered by HAWK by default (Rahman et al.,

2018). Allelic states were used to calculate the fixation index, G_{ST} (Nei and Chesser, 1983), for

each k-mer using custom Python scripts (https://github.com/em-bellis/StrigaWGS). G_{ST} is a

generalization of the widely used fixation index F_{ST} applicable to non-diploid loci (Nei, 1973); for

the purposes of our k-mer analysis, we consider loci as haploid rather than diploid, since only

presence or absence of the k-mer is scored. While we did not remove k-mers with excessively

high counts (likely derived from repetitive genomic sequences) these are not likely to be

identified as G_{ST} outliers since the k-mer would need to have a count of zero in most samples

from one population but not the other.

To gather functional information for host-associated k-mers, 31-mers with G_{ST} above 0.5

were extracted and assembled into longer contigs using ABYSS 2.0.2 (Jackman et al., 2017),

specifying a k-mer length of 29 for assembly. The k-mer length for assembly was chosen to

facilitate assembly of relatively longer contigs (minimum expected length of 59 bp, if formed by

two 29-mers overlapping with a SNP site on either side). To evaluate significance of the chosen

G_{ST} threshold, we carried out permutation analyses for Homa Bay and Kisii populations

separately. For each table (unique k-mers in rows and individuals in columns), the host label for

each individual was randomly permutated, and G_{ST} values for each k-mer recalculated. For

each permutation, we then calculated the proportion of k-mers with $G_{ST} > 0.5$, to estimate the

proportion of 'outlier' k-mers at our chosen G_{ST} threshold expected by chance. Assembled

contigs were then queried against contigs from two published *S. hermonthica* transcriptome

assemblies using BLAST optimized for short sequences (blastn-short). Transcriptome

assemblies in our BLAST database included StHeBC4 (265,694 scaffolds covering 369.7 Mb)

from the Parasitic Plant Genome Project (Westwood et al., 2012) and Sh14v2 (81,559 scaffolds

covering 83.9 Mb) from Yoshida *et al.* (2019). Annotations are based on the top hit from the

Sh14v2 transcriptome.

To further investigate genomic regions associated with host-specific differentiation, we

mapped cleaned sample reads to mRNA reference sequences, following the strategy from

Therkildsen and Palumbi (2017). Mapping to a transcriptome reference has the potential to

introduce errors in SNP calling due to intron/exon boundaries and the short length of transcripts,

so we focused this analysis on a small set of loci for which alignments could be manually

inspected including three transcripts with potential functions in haustorium development

(StHeBC4_h_c11261_g0_i1, StHeBC4_p_c12587_g2_i1, StHeBC4_u_c12903_g27039_i4; see

Results). The reference also included transcript sequences for a set of 11 previously

characterized *S. hermonthica* strigolactone receptors [GenBank accession numbers KR013121

- KR13131] (Tsuchiya et al., 2015). High quality, contaminant-filtered reads were mapped to the

reference transcriptome using BWA-MEM (Li, 2013), and alignments with low quality were

removed using SAMtools view (-q 20) (Li et al., 2009). Allele frequencies for each site in the

reference transcriptome were estimated based on genotype likelihoods using ANGSD, ignoring

low quality bases (-minQ 25), allowing reads for which only one end mapped (-

only_proper_pairs 0), and assuming known major and minor alleles but accounting for

uncertainty of the minor allele (-doMaf 3) (Kim et al., 2011; Korneliussen et al., 2014). Sites with

information for fewer than nine of twelve individuals in the population were excluded, and the

difference in estimated allele frequency between parasite populations on different hosts was

visualized with R version 4.0 (R Core Team, 2020). This strategy was used to filter false SNP

calls due to errors in mapping DNA-derived reads to a transcriptome reference, since these

SNPs should be observed as a fixed difference from the reference that occurs in both

populations. We further investigated genes with potential structural variation by extracting reads

aligned to the transcript reference and their unmapped pairs and reassembling them with

ABYSS (k=51) (Jackman et al., 2017).

Validation of k*-mer-based approaches*—To investigate patterns of selection

surrounding putative host-associated loci and validate findings from reference-free analyses, we

also mapped reads generated for *S. hermonthica* to a draft reference assembled specimen,

grown *ex situ* from seeds collected on maize in the Irimbi district of Southern Uganda

(specimens voucher deposited at MSUN). DNA was extracted from developing leaves and

inflorescences of one individual using a modified 1x CTAB-protocol with subsequent PEG-8000

precipitation and purification (Wicke et al., 2016). Genomic libraries were sequenced on an

Illumina HiSeq 2000 in 101 bp paired-end mode at Eurofins GATC Biotech GmbH (Constance,

Germany). Additional data were generated to a targeted depth of 110X using the HiSeq 2500

platform using the HiSeq SBS Kit v4 at Eurofins GATC Biotech GmbH, for which DNA from the

original extract was subjected to Φ29-polymerase based whole-genome amplification. Whole

genome-amplified DNA was size-selected for >20 kb fragments on 1% low-melting point

agarose and purified using an agarose digest-based purification with subsequent

ethanol/sodium acetate precipitation (Wicke et al., 2013). For the final assembly, we employed

Trimmomatic v0.36 (Bolger et al., 2014) to remove adapters and retain only sequences longer

than 36 bp with an average per-base quality above 15 ("ILLUMINACLIP:TruSeq3-PE.fa":

2:30:10 SLIDINGWINDOW:4:15 MINLEN:36). The quality-trimmed data were assembled using

SPAdes v3.10.1 with k-mer sizes of 21, 33, 55, and 77 (Bankevich et al., 2012). Assembly

quality was ascertained using Quast v4.5 (Gurevich et al., 2013) with default parameters for

eukaryotes. The resulting assembled contigs were contaminant-filtered, for which we ran a

nucleotide BLAST search of all contigs against the non-redundant nucleotide database (access

date: 10.07.2017) using BLAST+ v2.6 with an e-value of $1E10^{-4}$. Only contigs with the three best

hits matching to green land plants (Viridiplantae) were retained.

After removing scaffolds shorter than 500 bp from the reference, high quality sequences

from each *S. hermonthica* individual were mapped using BWA-MEM v0.7.17 (Li, 2013).

Sequences mapping with quality less than 20 were excluded using SAMtools v1.10 (Li et al.,

2009). Tajima's D was calculated in non-overlapping windows of 1-kb using ANGSD v0.935

based on the folded site frequency spectrum and including reads where only one end mapped (-

only_proper_pairs 0), to account for the highly fragmented nature of the assembly (Korneliussen

et al., 2014). F_{ST} was also calculated in non-overlapping windows with ANGSD, using the

SAMtools method for calculating genotype likelihoods. Genome-wide mean values of Tajima's D

were determined by fitting an intercept-only linear mixed model to window estimates of F_{ST} or

Tajima's D, including a random effect of 'contig' to account for increased correlation among

measurements from nearby genomic regions, with the R package lme4 (Bates et al., 2015). An

empirical *P*-value for Tajima's D for the contig containing the focal chemocyanin was calculated

based on the number of 10,000 randomly sampled windows of size matching the assembled

length of the contig (1-kb) with values more extreme than the observed value.

To validate the presence/absence polymorphism for the focal chemocyanin gene, we

performed PCR using primers designed from the reassembled 'finger millet' allele (Primer Set A:

5'-AAGATTGCGGTTACCACCAG-3' and 5'-TCTCGATCCTTTTGGAATGG-3') and the

transcript reference (Primer Set B: 5'- CAGGAGCAAGTAGAGTAGAGCA-3' and 5'-

TGGGGAAAGAGGTAGTGCAA-3'). PCR was performed with DreamTaq DNA Polymerase 2x

Mastermix (ThermoFisher, Waltham, Massachusetts, USA) under the following cycling

conditions: 95°C for 3 min; 30 cycles of 95°C (30 s), 50.3°C (30 s), 72°C (60 s); 72°C for 5 min.

Germination experiments—Seed germination was assayed in the USDA-APHIS-

permitted quarantine lab at Pennsylvania State University (permit no. P526P-21-04540). A

detailed step-by-step protocol is available from protocols.io repository (Bellis and Kelly, 2019),

following the modifications for testing seeds collected from individual plants. Briefly, seeds were

surface sterilized for 10 minutes in 1.5 mL microcentrifuge tubes with a 0.5% sodium

hypochlorite solution before preconditioning for 12 days at 30°C in separate wells of foil-

wrapped 12-well culture plates, with three replicates per unique germination stimulant and

parasite genotype combination. Germination counts were performed 3 days after addition of

germination stimulants. Tested germination stimulants included (+)5-deoxystrigol (Olchemim,

Olomouc, Czech Republic; CAS: 151716-18-6) or (±)orobanchol (Olchemim; CAS: 220493-64-1)

at 0.01 µM and (±)-GR24 (Chempep, Wellington, Florida, USA; CAS: 76974-79-3) at 0.2 µM.

GR24 is a synthetic strigolactone analog commonly used in laboratory germination studies of

parasitic plants as a positive control. 5-deoxystrigol is one of the major SLs produced by

compatible grass hosts (Awad et al., 2006) and is a potent stimulator of parasite germination

whereas orobanchol is a more dominant SL among dicot hosts (Yoneyama et al., 2008) and is a

less potent stimulator of *S. hermonthica* germination for certain genotypes (Haussmann et al.,

2004; Cardoso et al., 2014; Bellis et al., 2020).

In addition to seed collections kept separately from individual plants, we also included

tests of bulk seed collected from the Kibos population that were confirmed to have high

germinability in our previous experiments (Bellis et al., 2020). We used a generalized linear

mixed model (GLMM) with a random effect of *S. hermonthica* genotype (of the parent plant) to

compare differences in germination rate among sites, hosts, and treatments (orobanchol vs. 5-

deoxystrigol). GLMMs were implemented in R version 4.0 with the lme4 package (Bates *et al.*

2015), and significance of fixed effects was determined using likelihood ratio tests.

RESULTS

Sequencing—For the 68 Kenyan samples, on average 80% of reads per library were

classified as plant-derived using Kraken 2 (range: 71-86%). Most classified reads were

assigned specifically to *S. hermonthica* (mean: 46%) or *S. asiatica* (mean: 30%), both

represented in our database by high-quality transcriptomic or genomic resources from which

contaminant sequences (including from plant hosts) have been carefully filtered (Yang et al.,

2014; Yoshida et al., 2019). After quality and adapter trimming, on average 7.7 Gigabase pairs

of plant-derived sequence data remained per sample (range: 5.2-11.7 Gbp). Given flow

cytometry estimates of the genome size of *S. hermonthica* ranging from 1C = 0.9 Gb (Yoshida

et al., 2010) to 1C = 1.4 Gb (Estep et al., 2012), this sequencing effort corresponds to an

approximate average depth of 5.5x to 8.6x read coverage per base for each sample.

Population structure—Principal Coordinates Analysis (PCoA) based on *k*-mers

indicated high genetic diversity within populations, with only subtle differentiation of populations

from the same farm collected from different hosts (Fig. 1). Greater correlation was observed

between genetic and geographic distance than expected by chance (p = 0.001, Mantel test), but

patterns of isolation-by-distance were extremely weak (Fig. 1F). Within-population diversity was

very high at the geographic scale investigated. Many individuals exhibited a comparable range

of pairwise genetic diversity within a single field (*k*-mer distance ranging from 0.0216 - 0.0286; *n*

= 572 pairwise comparisons) as between individuals sampled more than 100 km apart (*k*-mer

distance ranging from 0.0242 - 0.0286; *n* = 336 pairwise comparisons; Fig. 1F).

371   Our alignment-free approach may not remove all contaminant sequences, and so we

also carried out a reference-based analysis using a draft reference genome assembled as part

of this study. Estimation of pairwise distances from reference-based analyses suggested that

samples were not strongly clustered by host or by sampling location, except for nine individuals

parasitizing finger millet in Kisii that clustered separately from other samples (Fig. 2). These

findings are consistent with our *k*-mer based analysis, in which the same individuals appear

close together and are also separated from other finger millet parasitizing individuals along the

first PCoA axis (Fig. 1D). Although some limited population genetic structuring may be present

within highly diverse *S. hermonthica* populations (perhaps due to multiple source populations,

for example), populations are not strongly differentiated by host at the genome-wide level.

382   ***Germination rate variation***—Previous studies have suggested that in contrast to other

locations in Africa, *S. hermonthica* populations from western Kenya demonstrate a generalist

germination response to strigolactones (Haussmann et al., 2004; Bellis et al., 2020). However,

Striga germination tests are typically conducted with bulk seeds collected from many individuals

in a field, potentially masking individual-level variation that could be segregating with respect to

parasitism on different hosts, for example if a generalist population is composed of individuals

that specialize on different resources (Bolnick et al., 2002).

389   To characterize individual-level variation in *S. hermonthica* parasitism*,* we conducted

controlled germination tests in the USDA-permitted quarantine facility at Pennsylvania State

University. Positive controls with bulk seed (collected in the Kibos region) and 0.2 µM of the

artificial strigolactone GR24 indicated good germinability for positive controls (66.5%), and no

germinated seeds were observed in wells with only sterile water. Compared to the higher

concentration of GR24, bulk seed showed slightly lower germination rates in response to 0.01

μM of the natural SLs orobanchol (51.0%) and 5-deoxystrigol (55.7%) indicating that the

concentrations of orobanchol and 5-deoxystrigol used in the individual-level experiment should

produce strong but sub-maximal germination responses.

We did not find significant host-associated germination variation among seeds collected

from individual parasites on different hosts (Table S1; Fig. 3). After accounting for treatment and

site, host-of-origin was not a statistically significant effect in our model ($P = 0.247$; likelihood

ratio test for test of full vs. reduced GLMM). The probability of germination was 30% lower for

individuals from Kisii compared to Homa Bay ($P = 0.04$; GLMM with fixed effects of 'Site' and

'Treatment') and 15% lower in response to orobanchol than to 5-deoxystrigol ($P = 0.004$;

GLMM).

Host-associated loci—Germination tests did not support host-specific differences in

strigolactone response in these populations. However, it is possible that host-specific selection

could still leave detectable signatures at the genetic level. We identified host-associated genetic

variation without a reference using a *k*-mer based approach. For Homa Bay, the total number of

31-mers generated was 2,765,562. For Kisii, the total number of *31*-mers generated was

2,123,902. Highly differentiated *k*-mers were defined as those having $G_{ST} > 0.5$. A lower

proportion of *31*-mers were highly differentiated for parasites on sorghum vs. maize hosts in

Homa Bay (0.6% of *31*-mers) compared to finger millet vs. maize hosts in Kisii (4.8% of *31*-

mers; Fig. 4A). These proportions far exceeded the proportions expected by chance, based on

G_{ST} values observed among 1000 random permutations of host labels for each of the

populations (Fig. 4B, D). In 1,000 random permutations, the maximum proportion of *k*-mers with

$G_{ST} > 0.5$ for the Homa Bay population was 0.2%, whereas the maximum for the Kisii population

was 0.5%.

Highly differentiated *31*-mers were then assembled into longer contigs for follow-up

analysis. For the Homa Bay population (maize vs. sorghum hosts), 42 contigs were assembled

ranging in length from 58 to 91 bp. Fifteen of these contigs had good BLAST hits to the

transcriptome database, with >85% identity over at least 45 bp (Table S2). For the Kisii

population (finger millet vs. maize hosts), highly differentiated *k*-mers assembled into 469

contigs with length ranging from 29 to 211 bp. Of these, 240 had good BLAST hits to transcripts

in our database (Table S3).

Of particular interest to our investigation were host-specific contigs with high levels of

similarity to known *Striga* parasitism genes including SL receptors and genes involved in

haustorium development. We first searched for similarity to a set of 11 *S. hermonthica*

strigolactone receptors that vary in their binding affinity for diverse strigolactones [GenBank

accession numbers KR013121 – KR13131] (Tsuchiya et al., 2015). We observed only one likely

spurious match to *ShHTL2* (87% similarity over 23 bp) and no hits to any of the 21 *KAI2*

paralogs from the *Striga asiatica genome* (Yoshida 2019). In follow-up analyses based on

alignments to reference *ShHTL* transcripts, only one site in *ShHTL6* had an estimated difference

in allele frequency between parasites on finger millet and maize exceeding 0.5 (Fig. S1). This

polymorphism occurred at position 457 and does not result in an amino acid change. Together,

we find little phenotypic or genomic evidence supporting host-specific differentiation for loci

involved in perception of strigolactones in these populations.

In contrast, several assembled host-specific contigs had good matches to loci implicated

in development of haustoria. Annotated transcripts included a 60 bp contig with >98% similarity

over its full length to a transcript annotated as *SUPPRESSOR OF G2 ALLELE SKP1* (*SGT1*)

(Table S2). *SGT1* was among the top upregulated genes in haustoria of the root parasitic plant

Thesium chinense and was hypothesized to be important for generating auxin response maxima

during haustorium development (Ichihashi et al., 2017). In *S. hermonthica,* it is also highly

expressed in imbibed seeds (Stage 0) and in haustoria attached to host roots (Stage 3; Fig. S2).

Parasitism on different hosts was associated with genetic structural variation in *SGT1*, with two

<100 bp regions often absent from parasites on sorghum but present for parasites on maize and

finger millet. We also identified a 59 bp contig with 96% similarity over its full length to a

transcript annotated as a pectin methylesterase; pectin methylesterases have previously well-

characterized functions in developing haustoria (Yang et al., 2014). The pectin methylesterase

transcript was not expressed in parasites grown on sorghum in most stages, but exhibited low,

non-zero expression in *S. hermonthica* seedlings after exposure to a haustorium inducing factor

(mean TPM = 0.03). Strong differentiation in transcript-aligned reads between individuals from

finger millet (frequency of the deletion allele = 0.65) and maize (frequency of the deletion allele

= 0.03) verified the signal observed from the *k*-mer association analysis.

Two additional assembled host-differentiated contigs of 113 bp and 103 bp had 99% and

97% similarity, respectively, over their full length to separate regions of a single transcript

annotated as a precursor of chemocyanin (Table S3). Chemocyanins may be of particular

interest due to the evolutionary co-option of many pollen tube genes by parasitic plants for

haustorium development (Yang et al., 2014) and the key role of chemocyanin as an attractant

for directing pollen tube growth (Kim et al., 2003). Alignment of our DNA sequences to the

chemocyanin transcript reference revealed host-associated structural variation (Fig. 5A). PCR-

based confirmation indicated 500 bp or more of genome sequence directly upstream of the 5'

end of the transcript was completely missing from parasites on maize, suggesting potential

impact on variation in gene expression levels. The fragmented nature of the draft genome

assembly precluded our ability to design PCR primers completely spanning the deletion, but

PCR banding patterns showed a characteristic absence of the region in individuals having the

allele more common on finger millet (Fig. S3), and a diversity of deletion alleles at this site (Fig.

S4). Using just sequences for individuals parasitizing finger millet, we reconstructed a 934 bp

contig, from reads that mapped to the transcript reference and their unmapped pairs.

Alignments to this 'finger millet' allele confirmed its presence in 9/12 individuals from Kisii

parasitizing finger millet and 0/12 parasites on maize (Table S3). Among all individuals

sequenced in our study, the allele is present at lowest frequency in maize parasites (14% of

individuals; $n = 35$) and intermediate frequency for parasites on sorghum (22%; $n = 18$); and

sugarcane (33%; $n = 3$)

Although patterns of allele frequency differentiation are partially correlated with overall

patterns of population structure (Fig. 2), additional lines of evidence point to an important role of

the chemocyanin in parasitic plant-host interactions. The chemocyanin transcript was present in

a previously identified list of *S. hermonthica* "core parasitism" genes, with highest expression

after exposure to haustorial initiation factors and during early post-attachment (Yang *et al.* 2014;

Fig 3B). It was also highly expressed in cells at the host-parasite interface in a study that used

laser capture microdissection to characterize gene expression in the *S. hermonthica*-sorghum

interaction (Honaas et al., 2013).

Validation of k-mer-based approaches—To further investigate signatures of selection

in the context of genome-wide patterns, we assembled a draft genome for *S. hermonthica* from

South Uganda*.* The total length of the assembled genome after filtering was 1,431 Mbp over

12,155,247 contigs, with a maximum contig length of 37.5 kb. The assembly was highly

fragmented with the largest 1.69 million contigs accounting for 50% of the assembly, and the

length of these contigs $\geq$ 110 bp (N/L50 = 1690935/110). Nevertheless, sequences of interest

were present on contigs long enough to allow for further interrogation. Specifically, the

assembled chemocyanin transcript for the finger millet allele had 97.8% identity over 918 bp to a

single contig of 1,055 bp (NODE_132909_length_1055_cov_32.632), and no other close hits.

After removing contigs shorter than 500 bp, 307026 scaffolds with a total length of 479 Mbp

remained.

To evaluate signatures of selection on the chemocyanin gene in the context of genome-

wide patterns, we estimated Tajima's D and F_{ST} for all assembled 1-kb non-overlapping

windows for maize- vs. finger millet-parasitizing individuals in the Kisii population. Consistent

with the expected signature of a selective sweep, a Tajima's D value of -1.7 in the Kisii

population indicated a significant excess of low-frequency polymorphism (empirical P-value =

0.009) for the chemocyanin contig in the *de novo* assembly compared to the genome-wide

value of 0.11 ± 0.013 (mean ± std. error; linear mixed effects model; distribution of Tajima's D

values relative to F_{ST} shown in Fig. 5C). For non-overlapping 1-kb windows with data ($n =$

237,632 windows), the median F_{ST} value was 0.038. Of these windows, 1.8% had an average

F_{ST} value greater than 0.15 and 0.5% had an average F_{ST} value greater than 0.2. The maximum

value of F_{ST} observed for the reference-based analysis was 0.57, for a 4-kb contig with similarity

to a *Phelipanche aegyptiaca* gene annotated as an ortholog of AT5G64440 (fatty acid amide

hydrolase) which is an important regulator of plant growth (Wang et al., 2006). Notably, F_{ST} for

chemocyanin contig could not be reliably calculated using the reference-based approach

because only 30 positions in the 1055-bp contig had data for a majority of individuals due to

structural variation. Together with observations from the G_{ST} analysis, our results suggest that

only a small portion of the *S. hermonthica* genome is strongly differentiated by host species in

these populations from western Kenya.

DISCUSSION

Agricultural weeds are increasingly recognized as important model systems for

addressing key questions in evolution and ecology (Vigueira et al., 2013; Baucom, 2019). In

particular, compared to many other systems, parasitic weeds offer particular advantages for the

study of coevolution including well-developed genomic and germplasm resources for their hosts,

high quality distributional data, and a rich literature describing variation in species interactions

over many decades (Bellis et al., 2020, 2021). Yet, despite recent advances in genome

sequencing for parasitic plants, evolutionary analyses particularly for species with large,

complex genomes (e.g. >1 Gb) remain a challenge (Lyko and Wicke, 2021). Consequently,

previous population-level diversity studies for *S. hermonthica*, one of the most damaging

parasitic plants in agriculture, have used reduced representation approaches (Unachukwu et al.,

2017; Lopez et al., 2019). However, reduced representation approaches such as RAD-seq may

miss signatures of selection that are highly localized in the genome (Lowry et al., 2017; Lou et

al., 2021), and transcriptomes fail to provide information regarding non-coding regions of the

genome, which also generate phenotypic diversity. As the cost of sequencing continues to

decrease, whole genome resequencing coupled with alignment-free bioinformatic approaches

can provide a promising alternate approach for surveying population genomic diversity (Voichek

and Weigel, 2020). Here, we report some of the first publicly available WGS data for field-

sampled individuals of the parasitic weed *Striga hermonthica*. Our analyses underscore high

within-population genomic variation and implicate host-specific selection on genes involved

during parasite attachment and haustorial development.

Perhaps surprisingly, we find little genomic or phenotypic evidence for host-specific

selection on strigolactone perception variation in the studied populations. Specifically, one may

expect selection on perception loci to be relatively strong, particularly since *S. hermonthica* is an

obligate parasite and so the costs of germination in the absence of a suitable host are high. This

suggests that, at least in western Kenya, most common grass hosts may be suitable, so that

costs of germinating on the wrong host are low. The genomes of *Striga* spp. include a diverse

repertoire of strigolactone receptors, each with variable affinity for different SLs (Tsuchiya et al.,

2015; Yoshida et al., 2019), providing many potential targets for selection on SL response

variation. One possibility is that these receptors are now subject to purifying selection in western

Kenya, rather than diversifying selection expected if SL perception variation is strongly linked to

fitness variation across different hosts. Environmental niche models from our previous study

(Bellis et al., 2021) predict highest suitability for *Striga* parasitism of maize across sampling

locations in our study (mean suitability: 0.92) but lower suitability for sorghum and pearl millet

(0.46 and 0.03, respectively). Another possibility is that the particular host genotypes studied

here may overlap in SL exudation profile enough that selection on parasite germination rate

variation is not strong in these natural field populations. For example, while zealactones appear

to be uniquely produced by maize (Charnikhova et al., 2017), some varieties also naturally

produce sorgomol and 5-deoxystrigol in high quantities (Yoneyama et al., 2015), strigolactones

common in sorghum root exudate that promote strong germination response in *S. hermonthica*.

A third explanation, if selection on SL perception is indeed important for local adaptation,

is that selection may only be strong in some populations across the range of *S. hermonthica.*

This explanation is consistent with the idea of coevolutionary hotspots, where there is reciprocal

selection among coevolving species (Thompson, 2005), but with the genetic targets of selection

involving different stages of the infection process in different locations. In western Kenya, for

example, coevolutionary hotspots may be 'hotter' for genes involved in parasitic interactions

post- germination than for SL perception. Kenyan *S. hermonthica* populations exhibit a more

generalist germination response compared to populations from West Africa, which show

pronounced differences in response to orobanchol vs. 5-deoxystrigol from host root exudates or

to strigolactone standards (Parker and Reid, 1979; Haussmann et al., 2004; Bellis et al., 2020).

This idea also corresponds with previous findings that East African *S. hermonthica* may have

greater average infestation success across diverse sorghum genotypes than West African

populations (Bozkurt et al., 2015). Functional G_H x G_P in strigolactone response variation may

segregate among populations in a different part of the parasite range than studied here, for

example among West African populations, or at a broader spatial scale, for example in West vs.

East African parasite populations (Haussmann et al., 2004; Bellis et al., 2020).

In contrast to general expectations (Nuismer and Dybdahl, 2016), our genomic analyses

revealed the strongest evidence for host-specific selection on genes involved in the later stages

of parasite development. The best evidence for host-specific selection on haustorium loci came

from our analyses of a transcript annotated as a chemocyanin precursor (Fig. 5). In addition to

strong differentiation between finger millet and maize hosts, the 1-kb region including the

chemocyanin exhibited an excess of rare polymorphism and multiple alleles (Fig. S3-S4),

consistent with expectations for recurrent soft sweeps from standing genetic variation (Pennings

and Hermisson, 2006). Notably, even the strongest signals of host-specific selection detected in

our study did not reveal any loci exhibiting complete or near complete differentiation between

parasites on different hosts, indicating that there may be relatively few genetic barriers to

parasitism of different cereal host species in the studied region. The 'finger millet' chemocyanin

allele, for example, was also present at low frequency in the genomes of parasites on other host

species. This suggests a neutral impact of the allele on parasitism of other hosts, and

potentially, a lack of trade-offs, given that all sequenced parasite individuals were already at an

advanced life stage. The importance of conditional neutrality for local adaptation in other

systems (Lowry et al., 2019) further highlights the complexity of selective pressures shaping

local adaptation of parasitic plants to dynamic host communities.

CONCLUSIONS

While our results emphasize the challenges of *Striga* management due to high genomic

diversity and adaptive potential, they also highlight the promise of low coverage WGS

approaches for functional genomics of non-model species. The outlier signal for two of the three

candidate loci we describe in detail here resulted from structural variation that would not have

been uncovered in an alignment-based analysis or using a RAD-Seq approach that may only

survey a small portion of the genome or be prone to allele drop-out. Reference-free approaches

continue to gain ground for studies of genomic and phenotypic variation in plants, with well-

documented advantages (VanWallendael and Alvarez, 2020; Voichek and Weigel, 2020). Our

study indicates that the utility of large WGS datasets may not be out of reach even for species

such as *Striga hermonthica* characterized by large, complex genomes. As the need to mitigate

biotic constraints on global food security becomes increasingly critical, reference-free analyses

coupled with WGS data may serve as a promising strategy for rapid characterization of alleles

involved in parasite adaptation across diverse environments.

ACKNOWLEDGMENTS

This study is based on work supported by a National Science Foundation Postdoctoral

Research Fellowship in Biology to E.S.B. under Grant 1711950, the Arkansas Biosciences

Institute (the major research component of the Arkansas Tobacco Settlement Proceeds Act of

2000), and the *Emmy Noether*-program of the German Science Foundation (WI4507/3-1 to

S.W.). A.K. and X.H. were supported by NSF EPSCoR Award 1946391. J.R.L. was supported

by NIH R35 GM138300. This work is supported by the Arkansas High Performance Computing

Center which is funded through multiple National Science Foundation grants and the Arkansas

Economic Development Commission. We also thank the Texas A&M AgriLife Research:

Genomics and Bioinformatics Service for library preparation and sequencing.

AUTHOR CONTRIBUTIONS

615 C.S.v.M., S.W., C.O.O., E.K., T.X., and E.S.B collected and processed samples and performed

616 laboratory experiments. C.S.v.M., S.W., A.K., and E.S.B. carried out analyses. X.H., S.W.,

617 S.M.R., C.W.D., J.R.L., and E.S.B. contributed to experimental design and interpretation. E.S.B.

618 and J.R.L. drafted the manuscript, with input and critical revision from all authors. All authors

619 approved the final version of the manuscript.

DATA AND CODE AVAILABILITY

Raw reads from whole genome sequencing of the 68 *S. hermonthica* individuals from the 2018

collection and the Ugandan reference have been deposited in the National Center for

Biotechnology Information (NCBI) Sequence Read Archive (SRA) database,

https://www.ncbi.nlm.nih.gov/sra (BioProject accession no. PRJNA801489). The reference

assembly of the Ugandan specimen is available for download and BLAST searches on *WARPP*

(https://warpp.app; (Kösters et al., 2021)). Images associated with the different collection sites

are available from iNaturalist. Germination rate data and code to reproduce the analyses are

available at https://github.com/em-bellis/StrigaWGS.

SUPPLEMENTARY INFORMATION

Additional supporting information may be found online in the Supporting Information Section at

the end of the article.

Appendix S1: Supplementary Figures and Tables.

LITERATURE CITED

Aigbokhan, E. I., D. K. Berner, and L. J. Musselman. 1998. Reproductive Ability of Hybrids of
Striga aspera and *Striga hermonthica*. *Phytopathology®* 88: 563–567.

Akiyama, K., K. Matsuzaki, and H. Hayashi. 2005. Plant sesquiterpenes induce hyphal
branching in arbuscular mycorrhizal fungi. *Nature* 435: 824–827.

Awad, A. A., D. Sato, D. Kusumoto, H. Kamioka, Y. Takeuchi, and K. Yoneyama. 2006.
Characterization of Strigolactones, Germination Stimulants for the Root Parasitic Plants
Striga and *Orobanche*, Produced by Maize, Millet and Sorghum. *Plant Growth*
Regulation 48: 221.

Bankevich, A., S. Nurk, D. Antipov, A. A. Gurevich, M. Dvorkin, A. S. Kulikov, V. M. Lesin, et al.
2012. SPAdes: a new genome assembly algorithm and its applications to single-cell
sequencing. *Journal of computational biology* 19: 455–477.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software* 67: 1–48.

Baucom, R. S. 2019. Evolutionary and ecological insights from herbicide-resistant weeds: what
have we learned about plant adaptation, and what is left to uncover? *New Phytologist*
223: 68–82.

Bebawi, F. F., R. E. Eplee, C. E. Harris, and R. S. Norris. 1984. Longevity of witchweed (*Striga*
asiatica) seed. *Weed Science* 32: 494–497.

Bellis, E., and E. Kelly. 2019. *Striga hermonthica* germination assay. *protocols.io*.

Bellis, E. S., E. A. Kelly, C. M. Lorts, H. Gao, V. L. DeLeo, G. Rouhan, A. Budden, et al. 2020.
Genomics of sorghum local adaptation to a parasitic plant. *Proceedings of the National*
Academy of Sciences 117: 4243.

Bellis, E. S., C. M. McLaughlin, C. W. dePamphilis, and J. R. Lasky. 2021. The geography of
parasite local adaptation to host communities. *Ecography*.

Bolger, A. M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 30: 2114–2120.

Bolnick, D. I., L. H. Yang, J. A. Fordyce, J. M. Davis, and R. Svanbäck. 2002. Measuring
Individual-Level Resource Specialization. *Ecology* 83: 2936–2941.

Bozkurt, M. L., P. Muth, H. K. Parzies, and B. I. G. Haussmann. 2015. Genetic diversity of East
and West African *Striga hermonthica* populations and virulence effects on a contrasting
set of sorghum cultivars. *Weed Research* 55: 71–81.

Bushnell, B. BBMap. Website sourceforge.net/projects/bbmap/.

Cardoso, C., T. Charnikhova, M. Jamil, P.-M. Delaux, F. Verstappen, M. Amini, D.
Lauressergues, et al. 2014. Differential Activity of *Striga hermonthica* Seed Germination
Stimulants and *Gigaspora rosea* Hyphal Branching Factors in Rice and Their
Contribution to Underground Communication. *PLOS ONE* 9: 1–11.

Castellano, D., L. H. Uricchio, K. Munch, and D. Enard. 2019. Viruses rule over adaptation in
conserved human proteins. *bioRxiv*: 555060.

Charnikhova, T. V., K. Gaus, A. Lumbroso, M. Sanders, J.-P. Vincken, A. D. Mesmaeker, C. P.
Ruyter-Spira, et al. 2017. Zealactones. Novel natural strigolactones from maize.
Phytochemistry 137: 123–131.

Conn, C. E., R. Bythell-Douglas, D. Neumann, S. Yoshida, B. Whittington, J. H. Westwood, K.
Shirasu, et al. 2015. Convergent evolution of strigolactone perception enabled host
detection in parasitic plants. *Science* 349: 540–543.

Cui, S., S. Wada, Y. Tobimatsu, Y. Takeda, S. B. Saucet, T. Takano, T. Umezawa, et al. 2018.
Host lignin composition affects haustorium induction in the parasitic plants
Phtheirospermum japonicum and *Striga hermonthica*. *New Phytologist* 218: 710–723.

Dayou, O., W. Kibet, P. Ojola, P. I. Gangashetty, S. Wicke, and S. Runo. 2021. Two-tier
witchweed (*Striga hermonthica*) resistance in wild pearl millet (*Pennisetum glaucum*)
29Aw. *Weed Science* 69: 300–306.

Ebert, D., and P. D. Fields. 2020. Host–parasite co-evolution and its genomic signature. *Nature*
Reviews Genetics 21: 754–768.

Ejeta, G. 2007. The Striga scourge in Africa: a growing pandemic. Integrating new technologies
for Striga control: towards ending the witch-hunt, 3–16. World Scientific.

Estep, M. C., B. S. Gowda, K. Huang, M. P. Timko, and J. L. Bennetzen. 2012. Genomic
characterization for parasitic weeds of the genus *Striga* by sample sequence analysis.
The Plant Genome 5.

Fumagalli, M., M. Sironi, U. Pozzoli, A. Ferrer-Admettla, L. Pattini, and R. Nielsen. 2011.
Signatures of environmental genetic adaptation pinpoint pathogens as the main selective
pressure through human evolution. *PLoS genetics* 7: e1002355.

Gbèhounou, G., A. H. Pieterse, and J. A. Verkleij. 2003. Longevity of Striga seeds reconsidered:
results of a field study on purple witchweed (*Striga hermonthica*) in Benin. *Weed*
Science 51: 940–946.

Gobena, D., M. Shimels, P. J. Rich, C. Ruyter-Spira, H. Bouwmeester, S. Kanuganti, T.
Mengiste, and G. Ejeta. 2017. Mutation in sorghum LOW GERMINATION STIMULANT 1
alters strigolactones and causes *Striga* resistance. *Proceedings of the National*
Academy of Sciences 114: 4471–4476.

Gurevich, A., V. Saveliev, N. Vyahhi, and G. Tesler. 2013. QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* 29: 1072–1075.

Hall, M. D., G. Bento, and D. Ebert. 2017. The evolutionary consequences of stepwise infection
processes. *Trends in Ecology & Evolution* 32: 612–623.

Hall, M. D., J. Routtu, and D. Ebert. 2019. Dissecting the genetic architecture of a stepwise
infection process. *Molecular ecology* 28: 3942–3957.

Haussmann, B., D. Hess, G. Omanya, R. Folkertsma, B. Reddy, M. Kayentao, H. Welz, and H.
Geiger. 2004. Genomic regions influencing resistance to the parasitic weed *Striga*
hermonthica in two recombinant inbred populations of sorghum. *Theoretical and Applied*
Genetics 109: 1005–1016.

Honaas, L. A., E. K. Wafula, Z. Yang, J. P. Der, N. J. Wickett, N. S. Altman, C. G. Taylor, et al.
2013. Functional genomics of a generalist parasitic plant: Laser microdissection of host-
parasite interface reveals host-specific patterns of parasite gene expression. *BMC Plant*
Biology 13: 9.

Ichihashi, Y., M. Kusano, M. Kobayashi, K. Suetsugu, S. Yoshida, T. Wakatake, K. Kumaishi, et
al. 2017. Transcriptomic and Metabolomic Reprogramming from Roots to Haustoria in
the Parasitic Plant, *Thesium chinense*. *Plant and Cell Physiology* 59: 729–738.

Iwo, G. A., G. O. Olaniyan, and S. W. H. Husaini. 1993. Cytological observations and
distribution of *Striga* species in central part of Nigeria With 2 Figures and one Table.
Feddes Repertorium 104: 497–501.

Jackman, S. D., B. P. Vandervalk, H. Mohamadi, J. Chu, S. Yeo, S. A. Hammond, G. Jahesh, et
al. 2017. ABySS 2.0: resource-efficient assembly of large genomes using a Bloom filter.
Genome research 27: 768–777.

Kavuluko, J., M. Kibe, I. Sugut, W. Kibet, J. Masanga, S. Mutinda, M. Wamalwa, et al. 2020.
GWAS provides biological insights into mechanisms of the parasitic plant (*Striga*)
resistance in Sorghum.

Kim, G., M. L. LeBlanc, E. K. Wafula, C. W. DePamphilis, and J. H. Westwood. 2014. Genomic-
scale exchange of mRNA between a parasitic plant and its hosts. *Science* 345: 808–
811.

Kim, S., J.-C. Mollet, J. Dong, K. Zhang, S.-Y. Park, and E. M. Lord. 2003. Chemocyanin, a
small basic protein from the lily stigma, induces pollen tube chemotropism. *Proceedings*
of the National Academy of Sciences 100: 16125–16130.

Kim, S. Y., K. E. Lohmueller, A. Albrechtsen, Y. Li, T. Korneliussen, G. Tian, N. Grarup, et al.
2011. Estimation of allele frequency and association mapping using next-generation
sequencing data. *BMC Bioinformatics* 12: 231.

Korneliussen, T. S., A. Albrechtsen, and R. Nielsen. 2014. ANGSD: Analysis of Next Generation
Sequencing Data. *BMC Bioinformatics* 15: 356.

Kösters, L. M., S. Wiechers, P. Lyko, K. F. Müller, and S. Wicke. 2021. WARPP—web
application for the research of parasitic plants. *Plant Physiology* 185: 1374–1380.

Li, H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, et al. 2009. The
Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25: 2078–2079.

Li, J., and M. P. Timko. 2009. Gene-for-gene resistance in *Striga*-cowpea associations. *Science*
325: 1094–1094.

Liu, N., G. Shen, Y. Xu, H. Liu, J. Zhang, S. Li, J. Li, et al. 2020. Extensive inter-plant protein
transfer between *Cuscuta* parasites and their host plants. *Molecular plant* 13: 573–585.

Lopez, L., E. S. Bellis, E. Wafula, S. J. Hearne, L. Honaas, P. E. Ralph, M. P. Timko, et al.
2019. Transcriptomics of host-specific interactions in natural populations of the parasitic
plant purple witchweed (*Striga hermonthica*). *Weed Science* 67: 397–411.

Lou, R., A. Jacobs, A. Wilder, and N. O. Therkildsen. 2021. A beginner's guide to low-coverage
whole genome sequencing for population genomics. *Authorea*.

Lowry, D. B., S. Hoban, J. L. Kelley, K. E. Lotterhos, L. K. Reed, M. F. Antolin, and A. Storfer.
2017. Breaking RAD: an evaluation of the utility of restriction site-associated DNA
sequencing for genome scans of adaptation. *Molecular Ecology Resources* 17: 142–
152.

Lowry, D. B., J. T. Lovell, L. Zhang, J. Bonnette, P. A. Fay, R. B. Mitchell, J. Lloyd-Reilley, et al.
2019. QTL × environment interactions underlie adaptive divergence in switchgrass
across a large latitudinal gradient. *Proceedings of the National Academy of Sciences*
116: 12933.

Lyko, P., and S. Wicke. 2021. Genomic reconfiguration in parasitic plants involves considerable
gene losses alongside global genome size inflation and gene births. *Plant Physiology*
186: 1412–1423.

Mallu, T. S., S. Mutinda, S. M. Githiri, D. Achieng Odeny, and S. Runo. 2021. New pre-
attachment *Striga* resistant sorghum adapted to African agro-ecologies. *Pest*
Management Science 77: 2894–2902.

Masumoto, N., Y. Suzuki, S. Cui, M. Wakazaki, M. Sato, K. Kumaishi, A. Shibata, et al. 2021.
Three-dimensional reconstructions of haustoria in two parasitic plant species in the
Orobanchaceae. *Plant Physiology* 185: 1429–1442.

Mbuvi, D. A., C. W. Masiga, E. K. Kuria, J. Masanga, M. Wamalwa, A. Mohamed, D. Odeny, et
al. 2017. Novel sources of witchweed (*Striga*) resistance from wild sorghum accessions.
Frontiers in plant science 8: 116.

Mutinda, S. M., J. Masanga, J. M. Mutuku, S. Runo, and A. Alakonya. 2018. KSTP 94, an open-
pollinated maize variety has postattachment resistance to purple witchweed (*Striga*
hermonthica). *Weed science* 66: 525–529.

Nei, M. 1973. Analysis of Gene Diversity in Subdivided Populations. *Proceedings of the National*
Academy of Sciences 70: 3321–3323.

Nei, M., and R. K. Chesser. 1983. Estimation of fixation indices and gene diversities. *Annals of*
Human Genetics 47: 253–259.

Nelson, D. C. 2021. The mechanism of host-induced germination in root parasitic plants. *Plant*
Physiology 185: 1353–1373.

Nuismer, S. L., and M. F. Dybdahl. 2016. Quantifying the coevolutionary potential of multistep
immune defenses. *Evolution* 70: 282–295.

Ondov, B. D., T. J. Treangen, P. Melsted, A. B. Mallonee, N. H. Bergman, S. Koren, and A. M.
Phillippy. 2016. Mash: fast genome and metagenome distance estimation using
MinHash. *Genome Biology* 17: 132.

Padgham, M., and M. D. Sumner. 2021. geodist: Fast, Dependency-Free Geodesic Distance
Calculations.

Paradis, E., and K. Schliep. 2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* 35: 526–528.

Parker, C., and D. C. Reid. 1979. Host specificity in *Striga* species: Some preliminary
observations. Proceeding of the Second International Symposium on Parasitic Weeds,
Raleigh, NC, 1979,.

Pennings, P. S., and J. Hermisson. 2006. Soft sweeps III: the signature of positive selection
from recurrent mutation. *PLoS genetics* 2: e186.

Poland, J. A., P. J. Balint-Kurti, R. J. Wisser, R. C. Pratt, and R. J. Nelson. 2009. Shades of
gray: the world of quantitative disease resistance. *Trends in plant science* 14: 21–29.

Qiu, S., J. M. Bradley, P. Zhang, R. Chaudhuri, M. Blaxter, R. K. Butlin, and J. D. Scholes.
2022. Identification of candidate virulence loci in *Striga hermonthica*, a devastating
parasite of African cereal crops. *bioRxiv*.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Rahman, A., I. Hallgrímsdóttir, M. Eisen, and L. Pachter. 2018. Association mapping from
sequencing reads using *k*-mers J. Flint [ed.],. *eLife* 7: e32920.

Ravensdale, M., A. Nemri, P. H. Thrall, J. G. Ellis, and P. N. Dodds. 2011. Co-evolutionary
interactions between host resistance and pathogen effector genes in flax rust disease.
Molecular plant pathology 12: 93–102.

Savary, S., L. Willocquet, S. J. Pethybridge, P. Esker, N. McRoberts, and A. Nelson. 2019. The
global burden of pathogens and pests on major food crops. *Nature ecology & evolution*
3: 430–439.

Shahid, S., G. Kim, N. R. Johnson, E. Wafula, F. Wang, C. Coruh, V. Bernal-Galeano, et al.
2018. MicroRNAs from the parasitic plant *Cuscuta campestris* target host messenger
RNAs. *Nature* 553: 82–85.

Shen, G., N. Liu, J. Zhang, Y. Xu, I. T. Baldwin, and J. Wu. 2020. *Cuscuta australis* (dodder)
parasite eavesdrops on the host plants' FT signals to flower. *Proceedings of the National*
Academy of Sciences 117: 23125–23130.

Spallek, T., M. Mutuku, and K. Shirasu. 2013. The genus *Striga*: a witch profile. *Molecular plant*
pathology 14: 861–869.

Therkildsen, N. O., and S. R. Palumbi. 2017. Practical low-coverage genome-wide sequencing
of hundreds of individually barcoded samples for population and evolutionary genomics
in nonmodel species. *Molecular Ecology Resources* 17: 194–208.

Thompson, J. N. 2005. The geographic mosaic of coevolution. University of Chicago Press.

Thrall, P. H., A.-L. Laine, M. Ravensdale, A. Nemri, P. N. Dodds, L. G. Barrett, and J. J. Burdon.
2012. Rapid genetic change underpins antagonistic coevolution in a natural host-
pathogen metapopulation. *Ecology letters* 15: 425–435.

Timko, M. P., K. Huang, and K. E. Lis. 2012. Host resistance and parasite virulence in *Striga*–
host plant interactions: A shifting balance of power. *Weed Science* 60: 307–315.

Toh, S., D. Holbrook-Smith, P. J. Stogios, O. Onopriyenko, S. Lumba, Y. Tsuchiya, A.
Savchenko, and P. McCourt. 2015. Structure-function analysis identifies highly sensitive
strigolactone receptors in *Striga*. *Science* 350: 203–207.

Tsuchiya, Y., M. Yoshimura, Y. Sato, K. Kuwata, S. Toh, D. Holbrook-Smith, H. Zhang, et al.
2015. Probing strigolactone receptors in *Striga hermonthica* with fluorescence. *Science*
349: 864–868.

Unachukwu, N. N., A. Menkir, I. Y. Rabbi, M. Oluoch, S. Muranaka, A. Elzein, G. Odhiambo, et
al. 2017. Genetic diversity and population structure of *Striga hermonthica* populations
from Kenya and Nigeria. *Weed Research* 57: 293–302.

VanWallendael, A., and M. Alvarez. 2020. Alignment-free methods for polyploid genomes: quick
and reliable genetic distance estimation. *bioRxiv*.

Vieira, F. G., F. Lassalle, T. S. Korneliussen, and M. Fumagalli. 2016. Improving the estimation
of genetic distances from Next-Generation Sequencing data: Genetic Distances from
NGS Data. *Biological Journal of the Linnean Society* 117: 139–149.

Vigueira, C. C., K. M. Olsen, and A. L. Caicedo. 2013. The red queen in the corn: agricultural
weeds as models of rapid adaptive evolution. *Heredity* 110: 303—311.

Voichek, Y., and D. Weigel. 2020. Identifying genetic variants underlying phenotypic variation in
plants without complete genomes. *Nature Genetics* 52: 534–540.

Wang, Y.-S., R. Shrestha, A. Kilaru, W. Wiant, B. J. Venables, K. D. Chapman, and E. B.
Blancaflor. 2006. Manipulation of *Arabidopsis* fatty acid amide hydrolase expression
modifies plant growth and sensitivity to *N* -acylethanolamines. *Proceedings of the*
National Academy of Sciences 103: 12197–12202.

Westwood, J. H., C. W. dePamphilis, M. Das, M. Fernández-Aparicio, L. A. Honaas, M. P.
Timko, E. K. Wafula, et al. 2012. The Parasitic Plant Genome Project: New Tools for
Understanding the Biology of Orobanche and Striga. *Weed Science* 60: 295–306.

Wicke, S., K. F. Müller, C. W. dePamphilis, D. Quandt, S. Bellot, and G. M. Schneeweiss. 2016.
Mechanistic model of evolutionary rate variation en route to a nonphotosynthetic lifestyle
in plants. *Proceedings of the National Academy of Sciences* 113: 9045–9050.

Wicke, S., K. F. Müller, C. W. de Pamphilis, D. Quandt, N. J. Wickett, Y. Zhang, S. S. Renner,
and G. M. Schneeweiss. 2013. Mechanisms of functional and physical genome
reduction in photosynthetic and nonphotosynthetic parasitic plants of the broomrape
family. *The Plant Cell* 25: 3711–3725.

Wood, D. E., J. Lu, and B. Langmead. 2019. Improved metagenomic analysis with Kraken 2.
Genome biology 20: 1–13.

Yang, Z., E. K. Wafula, L. A. Honaas, H. Zhang, M. Das, M. Fernandez-Aparicio, K. Huang, et
al. 2014. Comparative Transcriptome Analyses Reveal Core Parasitism Genes and
Suggest Gene Duplication and Repurposing as Sources of Structural Novelty. *Molecular*
Biology and Evolution 32: 767–790.

Yang, Z., E. K. Wafula, G. Kim, S. Shahid, J. R. McNeal, P. E. Ralph, P. R. Timilsena, et al.
2019. Convergent horizontal gene transfer and cross-talk of mobile nucleic acids in
parasitic plants. *Nature Plants* 5: 991–1001.

Yoneyama, K., R. Arakawa, K. Ishimoto, H. I. Kim, T. Kisugi, X. Xie, T. Nomura, et al. 2015.
Difference in *Striga*-susceptibility is reflected in strigolactone secretion profile, but not in
compatibility and host preference in arbuscular mycorrhizal symbiosis in two maize
cultivars. *New Phytologist* 206: 983–989.

Yoneyama, K., X. Xie, H. Sekimoto, Y. Takeuchi, S. Ogasawara, K. Akiyama, H. Hayashi, and
K. Yoneyama. 2008. Strigolactones, host recognition signals for root parasitic plants and
arbuscular mycorrhizal fungi, from Fabaceae plants. *New Phytologist* 179: 484–494.

Yoshida, S., J. K. Ishida, N. M. Kamal, A. M. Ali, S. Namba, and K. Shirasu. 2010. A full-length
enriched cDNA library and expressed sequence tag analysis of the parasitic weed,
Striga hermonthica. *BMC Plant Biology* 10: 1–10.

Yoshida, S., S. Kim, E. K. Wafula, J. Tanskanen, Y.-M. Kim, L. Honaas, Z. Yang, et al. 2019.
Genome sequence of *Striga asiatica* provides insight into the evolution of plant
parasitism. *Current Biology* 29: 3041–3052.

Appendix S1: Supplementary Figures and Tables

Figure S1. Allele frequency differences for 11 *ShHTL* receptors from Tsuchiya et al. (2015).

Allele frequencies were estimated from genotype likelihoods based on reads mapped to each

reference transcript. Allele frequency was estimated for parasites on each host species

separately ($n = 12$ per unique host and population). The difference in allele frequency estimates

between two hosts within a single population is shown.

Figure S2. (a, b) Minor allele frequency differences based on alignments to the *SGT1* transcript

sequence (StHeBC4_p_c12587_g2_i1), split by population. If the locus is not present in any of

the sequenced individuals for that population due to genetic structural variation, no data point is

shown. Frequencies were estimated based on genotype likelihoods from $n = 12$ individuals per

unique host and population. (c) Gene expression data in transcripts per million (TPM) from the

PPGPII data across the 6 stages of haustorial development from data published.

Figure S3. PCR banding results to confirm chemocyanin deletion calls. In the top panel,
binding sites for primer sets A (pink) and B (green) are shown relative to location of the
transcript sequence (solid black line), location of the deletion (grey dashed line), and conserved
plantacyanin domain (light blue rectangle). Primer set A does not yield a 676-bp band for
individuals with the deletion (lanes 11-15) whereas primer set B amplifies a region downstream
of the deletion. NTC: no template control.

911 **Figure S4. PCR banding results suggest multiple deletion variants.** PCR amplification

912 using the outer primers from set A (5'-TCTCGATCCTTTTGGAATGG-3') and B (5'-

913 TGGGGAAAGAGGTAGTGCAA-3'); see Fig. S3). NTC: no template control. Lane 1: Sh107;

914 Lane 2: Sh108, Lane 3: Sh109; Lane 4: Sh116; Lane 5: Sh117.

917   **Table S1.** GLMM for analysis of germination rate variation

	Estimate	Std. Error	z-value	P
Intercept	2.29	0.97	2.37	0.017
Site == Kisii	-2.16	0.79	-2.72	0.006
Host == Maize	-1.13	0.79	-1.43	0.153
Host == Sorghum	-1.90	1.12	-1.70	0.089
Treatment == ORO	-0.16	0.06	-2.85	0.004

921 **Table S2.** Hits to transcripts from the Parasitic Plant Genome Project (PPGPII) or Sh14v2 (from

922 Yoshida 2019) for contigs assembled from host-associated *k*-mers for *Striga hermonthica* Homa

923 Bay population (maize vs. sorghum hosts). Transcripts with greater than 85% identity over at

924 least 45 bp are shown.

PGPPII	Sh14v2	Description
StHeBC4_p_c26811_g0_i3	Sh14Contig_5850	lipid-a-disaccharide synthase-like
None	Sh14Contig_37747	None
None	Sh14Contig_44696	None
StHeBC4_p_c12903_g35328_i2	Sh14Contig_229	retrotransposon gag protein
StHeBC4_p_c12587_g2_i1	Sh14Contig_28742	disease-resistance protein SGT1
None	Sh14Contig_5301	None
None	Sh14Contig_37747	None
None	Sh14Contig_60980	hypothetical protein ASPNIDRAFT_144168
StHeBC4_u_c25350_g11_i1	Sh14Contig_78939	None
None	Sh14Contig_41198	None
None	Sh14Contig_32559	None
StHeBC4_u_c12903_g30818_i7	Sh14Contig_35967	None
None	Sh14Contig_47805	None
StHeBC4_p_c9824_g0_i1	None	None
StHeBC4_p_c12903_g42998_i1	None	None

926   **Table S3.** Hits to transcripts from the Parasitic Plant Genome Project (PPGPII) or Sh14v2 (from
927   Yoshida 2019) for contigs assembled from host-associated *k*-mers for *Striga hermonthica* Kisii
928   population (maize vs. finger millet hosts). Transcripts with greater than 85% identity over at least
929   45 bp are shown.

PGPPII	Sh14v2	Description
None	Sh14Contig_45649	None
StHeBC4_u_c12903_g32907_i4	Sh14Contig_20178	probable nicotinate-nucleotide pyrophosphorylase
StHeBC4_u_c12903_g22013_i6	Sh14Contig_24947	None
StHeBC4_u_c3941_g0_i1	Sh14Contig_42609	hypothetical protein VITISV_012016
None	Sh14Contig_70490	None
StHeBC4_u_c12903_g34431_i1	Sh14Contig_17925	upf0326 protein at4g17486-like
StHeBC4_p_c12903_g35474_i1	Sh14Contig_4481	None
StHeBC4_u_c24490_g2_i1	Sh14Contig_81114	None
StHeBC4_p_c12903_g14638_i24	Sh14Contig_23122	far-red impaired response protein
StHeBC4_u_c12903_g34431_i2	Sh14Contig_17925	upf0326 protein at4g17486-like
StHeBC4_u_c12903_g19156_i3	Sh14Contig_3012	alcohol dehydrogenase homolog
StHeBC4_p_c26043_g0_i2	Sh14Contig_13886	omega-amidase nit2
StHeBC4_p_c12903_g10585_i1	Sh14Contig_34611	None
None	Sh14Contig_20326	None
StHeBC4_p_c12903_g6756_i1	Sh14Contig_12790	None
StHeBC4_u_c12903_g15860_i5	Sh14Contig_29096	None
StHeBC4_u_c12903_g7220_i1	Sh14Contig_60050	None
None	Sh14Contig_3989	None
StHeBC4_p_c12903_g2276_i2	Sh14Contig_4881	rna polymerase ii c-terminal domain phosphatase-like 1-like
StHeBC4_u_c12903_g1840_i3	Sh14Contig_19406	e3 ubiquitin-protein ligase ring1-like
None	Sh14Contig_71030	None
StHeBC4_p_c12903_g39698_i9	Sh14Contig_45458	gag-protease polyprotein
StHeBC4_u_c12903_g32907_i12	Sh14Contig_81511	None
StHeBC4_u_c12903_g34427_i1	Sh14Contig_46360	None
StHeBC4_p_c12903_g35474_i1	Sh14Contig_4481	None
StHeBC4_p_c12903_g11444_i1	Sh14Contig_40193	polyprotein
StHeBC4_u_c19051_g0_i1	Sh14Contig_66323	None
StHeBC4_p_c12903_g10585_i1	Sh14Contig_12790	None
StHeBC4_u_c12903_g1736_i4	Sh14Contig_28260	None
None	Sh14Contig_31117	PREDICTED: uncharacterized protein LOC100256114
StHeBC4_u_c12903_g33489_i1	Sh14Contig_51019	None
StHeBC4_p_c16693_g0_i2	Sh14Contig_33691	hypothetical protein VITISV_018984
StHeBC4_p_c23323_g2_i3	Sh14Contig_3290	diphthamide biosynthesis protein 2-like
StHeBC4_p_c18975_g7_i10	Sh14Contig_7369	None
StHeBC4_p_c12903_g4626_i6	Sh14Contig_4698	retrotransposon protein
None	Sh14Contig_80867	None
StHeBC4_h_c11261_g0_i2	Sh14Contig_13520	chemocyanin precursor
StHeBC4_p_c23585_g1_i4	Sh14Contig_1911	None
StHeBC4_p_c24979_g0_i1	Sh14Contig_4792	97 kda heat shock protein
StHeBC4_h_c12903_g23955_i1	Sh14Contig_45458	gag-protease polyprotein

StHeBC4_u_c12903_g1547_i1	Sh14Contig_68469	None
None	Sh14Contig_48379	None
StHeBC4_p_c12903_g30160_i1	Sh14Contig_57886	None
None	Sh14Contig_64011	None
None	Sh14Contig_1642	retrotransposon ty1-copia subclass
StHeBC4_p_c22011_g3_i2	Sh14Contig_15531	wd repeat-containing protein 26-like
StHeBC4_p_c12903_g7195_i9	Sh14Contig_60490	unnamed protein product
StHeBC4_h_c24484_g1_i2	Sh14Contig_42033	hypothetical protein VITISV_042364
StHeBC4_p_c20621_g7_i3	Sh14Contig_4748	gag-pol precursor
StHeBC4_u_c12903_g27039_i4	Sh14Contig_4713	pectin methylesterase
StHeBC4_p_c18975_g7_i5	Sh14Contig_7369	None
None	Sh14Contig_65419	None
StHeBC4_u_c12903_g32907_i15	Sh14Contig_29096	None
None	Sh14Contig_52781	None
StHeBC4_p_c12903_g208_i2	Sh14Contig_7920	None
StHeBC4_p_c12903_g22741_i4	Sh14Contig_9532	pattern formation
StHeBC4_p_c12903_g38175_i2	Sh14Contig_17588	ring zinc finger ankyrin protein
None	Sh14Contig_4395	unnamed protein product [Vitis vinifera]
StHeBC4_u_c12903_g34427_i3	Sh14Contig_46360	None
StHeBC4_h_c11261_g0_i1	Sh14Contig_13520	chemocyanin precursor
StHeBC4_h_c12903_g40212_i2	Sh14Contig_24771	copia ltr rider
StHeBC4_p_c13584_g0_i1	Sh14Contig_33358	None
StHeBC4_u_c12903_g16409_i3	Sh14Contig_576	atp-dependent helicase brm-like
StHeBC4_h_c11483_g0_i1	Sh14Contig_55439	FAR1; Zinc finger, SWIM-type
StHeBC4_p_c12903_g20722_i11	Sh14Contig_28640	PREDICTED: uncharacterized protein LOC100854178, partial
StHeBC4_h_c12903_g12703_i2	Sh14Contig_9184	photosystem ii cp43 chlorophyll apoprotein
StHeBC4_p_c12903_g2185_i1	Sh14Contig_12790	None
StHeBC4_p_c12903_g34644_i1	Sh14Contig_4585	trehalose-phosphatase synthase 2
StHeBC4_p_c24106_g0_i1	Sh14Contig_40210	None
StHeBC4_h_c18968_g1_i5	Sh14Contig_1642	retrotransposon ty1-copia subclass
StHeBC4_u_c3088_g0_i1	Sh14Contig_31622	None
StHeBC4_u_c12903_g35782_i1	Sh14Contig_27773	None
None	Sh14Contig_23637	None
StHeBC4_u_c19051_g0_i1	Sh14Contig_63267	None
None	Sh14Contig_11046	uncharacterized protein loc100253271
StHeBC4_u_c12903_g27691_i1	Sh14Contig_35466	None
StHeBC4_u_c22214_g0_i6	Sh14Contig_43769	None
StHeBC4_u_c12903_g1736_i12	Sh14Contig_28260	None
None	Sh14Contig_58406	None
StHeBC4_u_c26755_g1_i1	Sh14Contig_68980	None
StHeBC4_p_c26987_g0_i1	Sh14Contig_48	6-4 photolyase
StHeBC4_u_c16917_g1_i1	Sh14Contig_50405	None
StHeBC4_u_c22344_g0_i6	Sh14Contig_813	thaumatin-like protein
StHeBC4_u_c12903_g699_i2	Sh14Contig_58790	None
StHeBC4_p_c12903_g14484_i1	Sh14Contig_12513	None
StHeBC4_p_c9911_g3_i3	Sh14Contig_54171	predicted protein
None	Sh14Contig_7920	None
StHeBC4_u_c12903_g34427_i8	Sh14Contig_46360	None

StHeBC4_p_c12903_g35474_i1	Sh14Contig_4481	None
None	Sh14Contig_43642	None
StHeBC4_u_c12903_g15888_i3	Sh14Contig_81511	None
StHeBC4_p_c24979_g1_i1	Sh14Contig_4792	97 kda heat shock protein
StHeBC4_p_c12903_g14483_i5	Sh14Contig_16250	tcp transcription factor 13
StHeBC4_p_c12903_g6756_i1	Sh14Contig_12790	None
StHeBC4_p_c12903_g10875_i1	Sh14Contig_20081	beta-glucan-binding protein
StHeBC4_p_c17648_g0_i3	Sh14Contig_15582	e3 ubiquitin-protein ligase upl3
StHeBC4_u_c12903_g22225_i1	Sh14Contig_48959	None
None	Sh14Contig_45784	None
None	Sh14Contig_75797	None
StHeBC4_u_c12903_g11727_i4	Sh14Contig_4041	None
StHeBC4_p_c23083_g3_i1	Sh14Contig_37240	None
None	Sh14Contig_5997	None
None	Sh14Contig_67788	None
StHeBC4_p_c12903_g6756_i1	Sh14Contig_1062	None
StHeBC4_u_c12903_g11833_i1	Sh14Contig_14842	None
StHeBC4_p_c12903_g6700_i3	Sh14Contig_43681	hypothetical protein VITISV_013115
StHeBC4_p_c20608_g0_i7	Sh14Contig_35726	PREDICTED: putative kinase-like protein TMKL1-like
None	Sh14Contig_29426	None
None	Sh14Contig_70066	None
StHeBC4_u_c12903_g29964_i1	Sh14Contig_37086	None
StHeBC4_p_c12903_g39295_i4	Sh14Contig_19126	epoxide hydrolase 2-like
StHeBC4_u_c12903_g1736_i13	Sh14Contig_29803	None
None	Sh14Contig_29803	None
StHeBC4_p_c12903_g6843_i1	Sh14Contig_4375	translation initiation factor eif-2b subunit delta-like
StHeBC4_u_c12903_g1736_i7	Sh14Contig_28260	None
StHeBC4_p_c12903_g38577_i4	Sh14Contig_21342	None
StHeBC4_p_c25304_g0_i1	Sh14Contig_5367	serine threonine-protein kinase pbs1-like
StHeBC4_u_c12903_g1736_i12	Sh14Contig_28260	None
StHeBC4_p_c12903_g7992_i1	Sh14Contig_22515	poly-specific ribonuclease parn
StHeBC4_u_c12903_g4208_i1	Sh14Contig_47575	None
StHeBC4_p_c11471_g0_i1	Sh14Contig_66224	None
StHeBC4_p_c12903_g40108_i1	Sh14Contig_17925	upf0326 protein at4g17486-like
StHeBC4_p_c26825_g3_i2	Sh14Contig_30816	unnamed protein product
StHeBC4_u_c17841_g0_i1	Sh14Contig_54135	None
StHeBC4_h_c12903_g30936_i1	Sh14Contig_28702	copia LTR rider
None	Sh14Contig_37720	None
StHeBC4_p_c12903_g10585_i1	Sh14Contig_12790	None
StHeBC4_u_c14956_g0_i2	Sh14Contig_3380	None
StHeBC4_u_c12903_g19836_i1	Sh14Contig_44128	None
StHeBC4_p_c12903_g9452_i3	Sh14Contig_68177	None
StHeBC4_p_c21038_g1_i1	Sh14Contig_46012	None
StHeBC4_u_c12903_g35474_i11	Sh14Contig_4481	None
StHeBC4_p_c12903_g15810_i1	Sh14Contig_12790	None
StHeBC4_u_c12903_g1736_i7	Sh14Contig_28260	None
StHeBC4_u_c12903_g34427_i1	Sh14Contig_61194	None

StHeBC4_p_c18569_g2_i14	Sh14Contig_67725	None
StHeBC4_u_c22214_g0_i2	Sh14Contig_43769	None
StHeBC4_p_c18975_g4_i3	Sh14Contig_77966	hypothetical protein VITISV_037041
StHeBC4_p_c22472_g0_i1	Sh14Contig_63886	predicted protein
StHeBC4_u_c24133_g4_i1	Sh14Contig_30270	None
StHeBC4_u_c12903_g12274_i1	Sh14Contig_5997	None
None	Sh14Contig_267	None
StHeBC4_u_c12903_g34427_i3	Sh14Contig_31554	None
StHeBC4_p_c12903_g33372_i2	Sh14Contig_18809	serine threonine protein kinase
StHeBC4_p_c12903_g12842_i2	Sh14Contig_58393	hypothetical protein VITISV_012059
StHeBC4_p_c12903_g13577_i4	Sh14Contig_4336	retrotransposon unclassified
StHeBC4_u_c12903_g34427_i9	Sh14Contig_46360	None
StHeBC4_u_c20739_g0_i7	Sh14Contig_30629	None
None	Sh14Contig_64863	None
StHeBC4_u_c18975_g2_i1	Sh14Contig_7369	None
StHeBC4_u_c3227_g1_i1	Sh14Contig_65419	None
StHeBC4_u_c18360_g0_i1	Sh14Contig_44903	None
StHeBC4_p_c12903_g6756_i1	Sh14Contig_1062	None
StHeBC4_p_c12903_g20268_i2	Sh14Contig_73193	None
StHeBC4_p_c12903_g15476_i8	Sh14Contig_78156	None
StHeBC4_u_c12903_g29964_i1	Sh14Contig_35346	None
None	Sh14Contig_62455	None
None	Sh14Contig_11292	None
StHeBC4_p_c12903_g43157_i2	Sh14Contig_5125	None
None	Sh14Contig_71030	None
StHeBC4_u_c12903_g18073_i1	Sh14Contig_20326	None
StHeBC4_p_c17480_g0_i1	Sh14Contig_62447	None
StHeBC4_p_c18975_g4_i6	Sh14Contig_44012	F7F22.15, related
StHeBC4_p_c12903_g32577_i4	Sh14Contig_50541	None
None	Sh14Contig_78156	None
StHeBC4_u_c12903_g32032_i1	Sh14Contig_20326	None
StHeBC4_u_c12903_g25724_i1	Sh14Contig_32318	None
StHeBC4_u_c13414_g0_i1	Sh14Contig_61194	None
None	Sh14Contig_43845	None
None	Sh14Contig_37580	None
StHeBC4_u_c12903_g1736_i12	Sh14Contig_28260	None
StHeBC4_p_c12903_g17146_i3	Sh14Contig_34580	None
StHeBC4_p_c12903_g33830_i2	Sh14Contig_22568	None
StHeBC4_p_c12903_g18700_i3	Sh14Contig_2850	None
None	Sh14Contig_61087	retrotransposon protein, putative, Ty3-gypsy subclass
StHeBC4_p_c12903_g208_i2	Sh14Contig_7920	None
StHeBC4_u_c14380_g6_i22	Sh14Contig_20089	None
StHeBC4_p_c12903_g22518_i1	Sh14Contig_39552	None
StHeBC4_u_c18188_g0_i1	Sh14Contig_30718	retrotransposon protein, putative, unclassified, expressed
StHeBC4_u_c12903_g35782_i1	Sh14Contig_27773	None
None	Sh14Contig_53940	None
None	Sh14Contig_29803	None

StHeBC4_p_c20119_g0_i1	None	None
StHeBC4_p_c12903_g26306_i42	None	None
StHeBC4_u_c12903_g15888_i8	None	None
StHeBC4_u_c12903_g34326_i2	None	None
StHeBC4_u_c22214_g0_i5	None	None
StHeBC4_u_c12903_g32907_i15	None	None
StHeBC4_p_c15477_g0_i2	None	None
StHeBC4_p_c12903_g18330_i2	None	None
StHeBC4_u_c20631_g2_i1	None	None
StHeBC4_u_c18975_g1_i1	None	None
StHeBC4_p_c12903_g33550_i2	None	None
StHeBC4_u_c20348_g0_i3	None	None
StHeBC4_u_c19161_g0_i1	None	None
StHeBC4_p_c18860_g2_i3	None	None
StHeBC4_p_c12903_g5434_i1	None	None
StHeBC4_p_c12903_g34427_i2	None	None
StHeBC4_u_c12903_g14239_i1	None	None
StHeBC4_p_c12903_g16488_i2	None	None
StHeBC4_u_c20110_g5_i1	None	None
StHeBC4_u_c17278_g0_i2	None	None
StHeBC4_p_c18569_g0_i2	None	None
StHeBC4_p_c12903_g15860_i11	None	None
StHeBC4_u_c12903_g16409_i1	None	None
StHeBC4_p_c12903_g5042_i2	None	None
StHeBC4_u_c12903_g42940_i10	None	None
StHeBC4_u_c18975_g1_i1	None	None
StHeBC4_u_c12903_g34427_i1	None	None
StHeBC4_u_c13012_g0_i1	None	None
StHeBC4_u_c20371_g0_i1	None	None
StHeBC4_u_c22472_g0_i4	None	None
StHeBC4_u_c12903_g34431_i3	None	None
StHeBC4_p_c18860_g2_i4	None	None
StHeBC4_p_c12903_g15230_i2	None	None
StHeBC4_u_c18860_g2_i48	None	None
StHeBC4_p_c12903_g8484_i1	None	None
StHeBC4_u_c18238_g1_i1	None	None
StHeBC4_u_c25350_g7_i2	None	None
StHeBC4_u_c12903_g29135_i2	None	None
StHeBC4_u_c13414_g0_i1	None	None
StHeBC4_p_c16184_g0_i1	None	None
StHeBC4_p_c12903_g27140_i4	None	None
StHeBC4_p_c12903_g18100_i3	None	None
StHeBC4_p_c26242_g21_i7	None	None
StHeBC4_u_c12903_g15860_i5	None	None
StHeBC4_p_c12903_g33449_i1	None	None
StHeBC4_u_c13012_g0_i1	None	None
StHeBC4_h_c10541_g0_i1	None	None
StHeBC4_u_c12903_g33428_i2	None	None
StHeBC4_u_c12903_g1552_i1	None	None

StHeBC4_p_c16622_g1_i2	None	None
StHeBC4_p_c12903_g8677_i10	None	None
StHeBC4_u_c10677_g0_i1	None	None
StHeBC4_p_c12903_g40108_i1	None	None
StHeBC4_u_c12903_g15860_i5	None	None
StHeBC4_p_c19033_g0_i1	None	None
StHeBC4_p_c12903_g33830_i1	None	None
StHeBC4_p_c12903_g11366_i6	None	None
StHeBC4_u_c12903_g25897_i6	None	None

931 **Table S4.** Chemocyanin variant calls for sequenced individuals. Individuals are coded as '1' if
932 they possess the 'finger millet' allele or '0' if the allele is missing.

SampleID	Site	Host	Lat	Lon	Elevation_ft	Chemocyanin
SH009	Mumias	maize	0.3342	34.47782	4306	0
SH014	Mumias	maize	0.3342	34.47782	4306	0
SH023	Mumias2	maize	0.3038	34.50713	4299	1
SH027	Mumias2	maize	0.3038	34.50713	4299	0
SH031	Mumias2	maize	0.3038	34.50713	4299	0
SH035	Kibos	sorghum	-0.0363167	34.81567	3893	1
SH039	Kibos	sorghum	-0.0363167	34.81567	3893	0
SH042	Kibos	sorghum	-0.0363167	34.81567	3893	1
SH046	Kibos	maize	-0.0341417	34.81628	3917	0
SH055	Kibos	maize	-0.0341417	34.81628	3917	0
SH065	Muhoroni2	maize	-0.15125	35.19167	4239	0
SH070	Muhoroni2	maize	-0.15125	35.19167	4239	0
SH072	Muhoroni2	sugarcane	-0.15125	35.19167	4239	0
SH074	Muhoroni2	sugarcane	-0.15125	35.19167	4239	1
SH077	Muhoroni2	sugarcane	-0.15125	35.19167	4239	0
SH079	Chemelil	maize	-0.082375	35.13167	4176	0
SH087	Chemelil	maize	-0.082375	35.13167	4176	1
SH091	Chemelil2	sorghum	-0.0941667	35.12495	4060	0
SH097	Chemelil2	sorghum	-0.0941667	35.12495	4060	1
SH101	Chemelil2	sorghum	-0.0941667	35.12495	4060	0
SH103	Kisii	finger millet	-0.6138	34.73172	4925	1
SH104	Kisii	finger millet	-0.6138	34.73172	4925	1
SH105	Kisii	finger millet	-0.6138	34.73172	4925	1
SH106	Kisii	finger millet	-0.6138	34.73172	4925	1
SH107	Kisii	finger millet	-0.6138	34.73172	4925	1
SH108	Kisii	finger millet	-0.6138	34.73172	4925	1
SH109	Kisii	finger millet	-0.6138	34.73172	4925	1
SH110	Kisii	finger millet	-0.6138	34.73172	4925	1
SH111	Kisii	finger millet	-0.6138	34.73172	4925	1
SH112	Kisii	finger millet	-0.6138	34.73172	4925	0
SH113	Kisii	finger millet	-0.6138	34.73172	4925	0
SH114	Kisii	finger millet	-0.6138	34.73172	4925	0
SH115	Kisii	maize	-0.6133167	34.7318	4930	0
SH116	Kisii	maize	-0.6133167	34.7318	4930	0
SH117	Kisii	maize	-0.6133167	34.7318	4930	0
SH118	Kisii	maize	-0.6133167	34.7318	4930	0
SH119	Kisii	maize	-0.6133167	34.7318	4930	0
SH120	Kisii	maize	-0.6133167	34.7318	4930	0
SH121	Kisii	maize	-0.6133167	34.7318	4930	0
SH122	Kisii	maize	-0.6133167	34.7318	4930	0
SH123	Kisii	maize	-0.6133167	34.7318	4930	0
SH124	Kisii	maize	-0.6133167	34.7318	4930	0
SH125	Kisii	maize	-0.6133167	34.7318	4930	0
SH126	Kisii	maize	-0.6133167	34.7318	4930	0
SH127	Homa Bay	sorghum	-0.5839667	34.4762	4173	0

SH128	Homa Bay	sorghum	-0.5839667	34.4762	4173	0
SH129	Homa Bay	sorghum	-0.5839667	34.4762	4173	0
SH130	Homa Bay	sorghum	-0.5839667	34.4762	4173	0
SH131	Homa Bay	sorghum	-0.5839667	34.4762	4173	0
SH132	Homa Bay	sorghum	-0.5839667	34.4762	4173	0
SH133	Homa Bay	sorghum	-0.5839667	34.4762	4173	1
SH134	Homa Bay	sorghum	-0.5839667	34.4762	4173	0
SH135	Homa Bay	sorghum	-0.5839667	34.4762	4173	0
SH136	Homa Bay	sorghum	-0.5839667	34.4762	4173	0
SH137	Homa Bay	sorghum	-0.5839667	34.4762	4173	0
SH138	Homa Bay	sorghum	-0.5839667	34.4762	4173	0
SH139	Homa Bay 2	maize	-0.5834	34.47635	4161	0
SH140	Homa Bay 2	maize	-0.5834	34.47635	4161	0
SH141	Homa Bay 2	maize	-0.5834	34.47635	4161	0
SH142	Homa Bay 2	maize	-0.5834	34.47635	4161	1
SH143	Homa Bay 2	maize	-0.5834	34.47635	4161	1
SH144	Homa Bay 2	maize	-0.5834	34.47635	4161	0
SH145	Homa Bay 2	maize	-0.5834	34.47635	4161	1
SH146	Homa Bay 2	maize	-0.5834	34.47635	4161	0
SH147	Homa Bay 2	maize	-0.5834	34.47635	4161	0
SH148	Homa Bay 2	maize	-0.5834	34.47635	4161	0
SH149	Homa Bay 2	maize	-0.5834	34.47635	4161	0
SH150	Homa Bay 2	maize	-0.5834	34.47635	4161	0

FIGURE LEGENDS

937
Figure 1. Population genomics of *S. hermonthica* from western Kenya. A) Map of the six

sampling locations. (B-E) Principal Coordinates Analysis (PCoA) based on *k*-mer-derived

genomic distances, performed separately for B) all sampled individuals ($n = 68$), C) five

individuals per location ($n = 30$), D) individuals from Kisii ($n = 24$), and E) individuals from Homa

Bay ($n = 24$). Relative proportion of variability explained by the first and second principal

coordinate axes is indicated for each analysis. F) Genetic vs. geographic distance. Genetic

distance was based on 31-mers. The dashed blue line indicates expectations from the best fit

line (y = 0.000013 * km + 0.0246; $R^2 = 0.05$) for $n = 4,556$ pairwise comparisons among different

individuals.

Figure 2. Population structure of Kenyan *S. hermonthica* samples from reference-based

analysis. Pairwise genetic distances were estimated from genotype probabilities, for 11,615,822

variable sites across 43,910 contigs longer than 2.5 kb. Genetic distances are colored in a heat

map-scale from 0.24 (red) to 0.33 (yellow). Individuals are ordered based on a hierarchical

clustering computed from the estimated genetic distances.

955   **Figure 3.** Germination response variation in Homa Bay and Kisii populations. Seeds from $n = 6$

956   individuals per unique site and host were tested in response to synthetic strigolactones

957   orobanchol (ORO, 0.01 µM) and 5-deoxystrigol (5DS, 0.01 µM). The dashed line indicates the

958   expectation if the percent germination in response to the two different germination stimulants is

959   identical.

Figure 4. G_{ST} and k-mer distributions. A) Observed distribution of G_{ST} values. For Homa Bay,

G_{ST} is for parasites in adjacent plots of maize vs. sorghum at 2,765,562 31-mers. For Kisii, G_{ST}

is for parasites in adjacent plots of maize vs. sorghum at 2,123,902 31-mers. B) Proportion of

31-mers with $G_{ST} > 0.5$ in the Kisii dataset. The shown distribution is based on 1,000 random

permutations of host labels. The dashed line indicates the observed proportion. C) A

representative k-mer distribution spectrum for sample SH046. Most unique 31-mers are

observed only once or twice due to the high heterozygosity of this species. D) Proportion of 31-

mers with $G_{ST} > 0.5$ in the Homa Bay dataset. The shown distribution is based on 1,000 random

permutations of host labels. The dashed line indicates the observed proportion.

Figure 5. Genetic variation in *Striga hermonthica* from Kisii, Kenya. (a) Presence/absence

variation across the chemocyanin precursor transcript (StHeBC4_h_c11261_g0_i1), for n = 12

individuals per host (finger millet or maize). If the locus is not present in any of the sequenced

individuals, no data point is shown, otherwise each point is colored according to the number of

individuals (nInd) with data for the position. Two regions in the transcript have no reads aligned

from maize-parasitizing individuals (lower panel), consistent with absence of the corresponding

k-mers and PCR-banding results that suggest a deletion. (b) Gene expression data in

transcripts per million (TPM) from the PPGPII dataset for the chemocyanin precursor transcript

across six stages of haustorial development (0: imbibed seed; 1: germinated seedling after

exposure to GR24; 2: germinated seedling after exposure to DMBQ; 3: ~48 hrs post-

attachment; 4: ~72 hrs post-attachment; 5: late post-attachment; 6.1: vegetative structures; 6.2:

reproductive structures). (c) Distribution of Tajima's D and F_{ST} values across 154,722 non-

overlapping 1-kb windows. Only windows with data for more than 50% of sites are shown,

excluding impacts of structural genetic variation as in (a). Tajima's D for the chemocyanin was -

1.7.