A dendritic mechanism for balancing synaptic flexibility and stability

Courtney E. Yaeger[1,2]**, Dimitra Vardalaki**[1,2]**, Qinrong Zhang**[3,4]**, Trang L.D. Pham**[1,2]**, Norma J.**

Brown[1,2]**, Na Ji**[3,4,5,6]**, & Mark T. Harnett**[1,2]

[1]McGovern Institute for Brain Research, MIT, Cambridge, MA, USA

[2]Department of Brain and Cognitive Sciences, MIT, Cambridge, MA, USA

[3]Department of Physics, University of California, Berkeley, CA, USA

[4]Department of Molecular and Cell Biology, University of California, Berkeley, CA, USA

[5]Helen Wills Neuroscience Institute, University of California, Berkeley, CA, USA

[6]Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National

Laboratory, Berkeley, CA, USA

To whom correspondence should be addressed: Harnett (harnett@mit.edu)

Abstract

Biological and artificial neural networks learn by modifying synaptic weights, but it is

unclear how these systems can retain previous knowledge and also acquire new

information. Here we show that cortical pyramidal neurons can solve this plasticity-

versus-stability dilemma by differentially regulating synaptic plasticity at distinct

dendritic compartments. Oblique dendrites of adult mouse layer 5b cortical pyramidal

neurons selectively received monosynaptic thalamic input, integrated linearly, and—

surprisingly—lacked synaptic potentiation. In contrast, basal dendrites, which do not

receive thalamic input, exhibited conventional NMDA receptor-mediated supralinear

integration and synaptic potentiation. Congruently, spiny synapses on oblique branches

showed low structural plasticity in vivo. A selective decline in NMDAR activity and

expression at synapses on oblique dendrites was controlled by an experience-dependent

critical period. Our results demonstrate a new biological mechanism for how single

neurons can safeguard a set of inputs from ongoing plasticity by altering synaptic

properties at distinct dendritic domains.

Main

The brain learns continuously by balancing plasticity and stability, whereby new information is

incorporated without disrupting prior knowledge. Synaptic plasticity is the key principle for

learning, but neural circuits must also exhibit sufficient synaptic stability to maintain information

storage over the long term. This process poses a major unsolved problem for both biological

and artificial neural networks, known as the plasticity-stability dilemma[1–4]. In artificial learning

systems, this problem manifests as catastrophic forgetting, where previously acquired

information is overwritten unless weight changes are limited[5–7]. While half a century of

experimental and theoretical work on biological synaptic plasticity has focused on how plasticity

is induced[8–17] and maintained within a working range[18–22], little is currently known about how

neurons sustain previously learned information throughout a lifetime of experience-dependent

plasticity.

In the mammalian sensory cortex, a classical plasticity-stability challenge occurs during

early postnatal development[23]. Initial sensory experiences must be stored and stabilized for

robust sensory processing, but cortical circuits must maintain flexibility to accommodate

subsequent learning. For example, primary sensory inputs in the visual cortex (V1) from the

dorsal lateral geniculate nucleus of the thalamus (dLGN) rapidly change at the onset of vision[24–

27]. By adulthood, these inputs are stable and no longer plastic during normal visual

experience[28–31]. However, higher-order cortical representations remain highly plastic[32–35].

Because single cortical neurons receive spatially-targeted inputs[36–39], we hypothesized that

dendritic compartmentalization could arbitrate plasticity or stability for different kinds of

information. Using complimentary physiological and imaging techniques, we identified dLGN-

recipient and non-recipient dendritic domains within layer 5b pyramidal neurons (L5b PNs) and

characterized the synaptic properties and plasticity of these domains across maturation.

Results

L5b PN oblique dendrites selectively receive thalamic input and exhibit distinct

integrative properties from basal dendrites

Visual information from dLGN projects primarily to cortical layer 4[40]. First-order thalamic inputs

are known to synapse on L5 PNs[41–43], which integrate inputs from all cortical layers, though the

location of the synaptic contacts is unclear. Physiological data from juvenile V1[38] and

anatomical data[44] suggest that dLGN inputs target the apical oblique dendrites of L5 PNs, which

can reside in layer 4. To directly test where dLGN inputs make functional synapses on L5b PNs

in adult mouse V1, we employed subcellular channelrhodopsin (ChR2)-assisted circuit

mapping[39]. After viral expression of ChR2 in dLGN (Fig. 1a), we performed whole-cell patch-

clamp recordings from V1 L5b PNs in acute slices from adult mice (postnatal day 56+). Local

photostimulation of ChR2-expressing dLGN boutons across the dendritic tree revealed

monosynaptic connections tightly restricted to apical oblique branches (Fig. 1b,c).

In light of this specific targeting of dLGN inputs to L5b PN oblique dendrites, we asked

whether these branches exhibited distinct properties compared to nearby basal dendrites (Fig.

1d). In acute slices from adult mice, basal dendrites displayed highly supralinear integration and

large local Ca^{2+} signals in response to spatiotemporally-clustered two-photon glutamate

uncaging (Fig. 1e,f,g), consistent with previous reports of NMDA spikes in juvenile cortex[45]. In

contrast, oblique dendrites integrated inputs linearly until axosomatic action potential threshold

was reached and did not show significant subthreshold Ca^{2+} influx (Fig. 1e,f,g), similar to what

has been previously described in retrosplenial cortex[36]. These differences could not be

explained by the size of uncaging events, the number of inputs, distance from the soma, action

potential threshold, or δV/δt prior to action potential initiation (Extended Data Fig. 1). Thus,

oblique dendrites of adult V1 L5b PNs selectively receive dLGN input and exhibit a specialized

linear integration mode.

Oblique dendrites lack synaptic potentiation in adult V1 L5b PNs

The lack of large NMDAR-mediated subthreshold Ca^{2+} signals in oblique dendrites led us to

hypothesize that synapses on these branches may exhibit plasticity differences compared to

basal dendrites. We tested synaptic potentiation at basal or oblique branches by pairing focal

electrical stimulation of presynaptic axons with somatic current injection. A theta glass bipolar

stimulating electrode was positioned within 10 μm of either a basal or an oblique dendritic

branch (Fig. 2a,b, Extended Data Fig. 2) to generate a 1 to 2 mV excitatory postsynaptic

potential (EPSP) in the presence of a GABA-A antagonist. EPSPs were paired with 20 ms

somatic current injections that drove bursts of 2-4 action potentials. The peak of the EPSP

preceded the peak of the first action potential by 10 ms. 5 pairings at 10 Hz were repeated 30

times, with 10 seconds between epochs (Fig. 2c). This protocol robustly potentiated synapses

on basal dendrites (Fig. 2d,f,g, Extended Data Fig. 2), similar to findings in juvenile cortex[12].

Potentiation at basal branches required pre- and post-synaptic coincidence as well as NMDARs

(Fig. 2g, Extended Data Fig. 2). In contrast, synapses on oblique dendrites did not exhibit any

significant potentiation after pairing (Fig. 2e-g, Extended Data Fig. 2). Doubling the number of

pairings did not change this outcome (Extended Data Fig. 2). These findings indicate that, unlike

basal dendrites, oblique branches of L5b PNs in adult V1 lack the capability for synaptic

potentiation under conventional plasticity induction protocols.

Synaptic plasticity in oblique dendrites is limited to a postnatal critical period

A developmental decline in synaptic potentiation has been reported for some cortical synapses
[46,47], and it is known that dLGN inputs in V1 undergo substantial refinement during early
postnatal development[25,48,49]. Therefore, we hypothesized that synapses on oblique L5b PN
dendrites may exhibit NMDAR-mediated supralinear integration and plasticity during early
postnatal development. We found that at eye-opening (postnatal days 12-14), oblique and basal
dendrites displayed similar NMDAR-dependent supralinearities and large local Ca^{2+} signals (Fig.
3a-c, Extended Data Fig. 3). These properties persisted into the canonical critical period (P18-
22)[31,50]. But by 4 weeks of age (P28-P32), oblique dendrites had developed adult-like
properties, with linear synaptic integration and minimal local Ca^{2+} signals (Fig 3c, Extended
Data Fig. 3). Synaptic plasticity at oblique dendrites followed the same developmental trajectory:
oblique and basal dendrites exhibited NMDAR-dependent synaptic potentiation at juvenile ages
(Fig. 3d, Extended Data Fig. 3) and by P28 apical integrative properties became adult-like.
These findings suggested that visual experience mediates the maturation of oblique dendrite
properties: to test this, we conducted these same experiments in mice deprived of light from
birth. In P28 dark-reared animals, oblique dendrites retained immature supralinear integration
with large local Ca^{2+} signals and synaptic potentiation (Fig. 3e, Extended Data Fig. 3).
Furthermore, dark-reared mice re-exposed to a normal light cycle for 2-4 weeks did not exhibit
adult-like integrative properties (Extended Data Fig. 3), indicating that the oblique compartment
matures only during a specific window in development. Collectively, these results reveal a
restricted developmental window in which visually-driven thalamic activity shifts the L5b PN
oblique dendritic compartment to a linearly-integrating, non-potentiating mode.
123

Oblique dendrites develop distinct postsynaptic receptor composition
One possible interpretation of our results thus far is that adult oblique dendrites lack large local
Ca^{2+} signals, integrate linearly, and do not potentiate due to a higher AMPA-to-NMDA ratio, as
adult cortical AMPA receptors (AMPARs) are significantly less Ca^{2+}-permeable than
NMDARs[51,52]. To test this, we compared AMPAR and NMDAR-mediated responses at single
spines from basal and oblique dendrites at P21 and in adults. Glutamate uncaging produced
AMPAR-mediated EPSPs within the physiological range at single spines, below the threshold
for spine Ca^{2+} influx (Fig. 4a, Extended Data Fig. 4). Mg^{2+}-free ACSF containing the AMPAR
antagonist DNQX was subsequently washed in and the uncaging protocol was repeated to
assess synaptic NMDARs (Fig. 4b). The average maximum amplitudes of EPSPs under each
condition were compared as AMPA/NMDA. At postnatal day 21, when oblique dendrites behave
similarly to basal dendrites, spines from the two branch types had comparable AMPA/NMDA. By

adulthood, this ratio increased for oblique dendrites (Fig. 4c), indicating relatively less NMDAR-

mediated synaptic conductance, consistent with linear integration, smaller local Ca^{2+} signals,

and a decreased capacity for synaptic potentiation.

To provide complementary proteomic evidence for our observed changes in functional

AMPA/NMDA, we acquired super-resolution images of oblique and basal dendrites and their

synaptic receptor content using epitope-preserving Magnified Analysis of the Proteome (eMAP,

Fig. 4d)[53,54]. Fixed slices from V1 of adult Thy1-GFP-M mice were expanded to 4x and stained

with antibodies against the presynaptic marker bassoon, the obligate NMDAR subunit GluN1,

and GluA1 and GluA2, the predominant AMPAR subunits in cortex[55]. Oblique and basal

dendrites from L5b PNs were imaged with a confocal microscope in ~40 μm-thick segments

(~180 μm expanded), and dendritic protrusions were annotated (Fig. 4e, f). Immature spines

and protrusions without bassoon were excluded post hoc (although basal and oblique branches

had similar numbers of immature and mature spines; Extended Data Fig. 4). For each spine

head, the intensity of each fluorophore was spatially integrated (Fig. 4g), and the ratio of

GluA1+GluA2/NR1 was taken. Consistent with our functional data, oblique dendrite synapses

exhibited higher AMPA/NMDA than basal dendrites (Fig. 4h, Extended Data Fig. 4). This effect

was pronounced in large spines (Fig. 4i). Taken together, these results indicate that oblique

dendrites in adults have relatively less NMDAR content and conductance, and this underlies the

differences in integration and plasticity rules between basal and oblique dendritic compartments.

Compartmentalized synaptic stability and flexibility in single neurons in vivo

Variable levels of synaptic Ca^{2+} from NMDARs trigger molecular cascades that form, enlarge,

shrink, and/or eliminate spiny synapses[56–62]. Given that we observed relatively low NMDAR

conductance and expression in apical oblique dendrites, we hypothesized that oblique dendrites

would have largely stable spiny synapses in vivo, whereas spiny synapses on basal dendrites

should exhibit dynamic structural plasticity. To investigate this, we longitudinally imaged eGFP-

expressing L5b PNs in adult mice. With extremely sparse cellular labeling and an optimized two-

photon microscopy set up (Fig. 5a), we were able to resolve spines in oblique and basal

branches in layers 4 and 5 (Fig. 5b). Across 5 or more consecutive days, we tracked spine

dynamics for both basal and oblique dendritic compartments in individual cells (n = 6 cells, 775

basal spines and 794 oblique spines, 4 animals). Spine density did not significantly differ

between basal and oblique compartments, consistent with our observations with eMAP (Fig. 5b,

Extended Data Fig. 5). The overall survival fraction of basal and oblique spines was similar to

what was previously reported for V1 apical tuft spines over a similar period of time, but spines

on oblique branches had significantly higher survival rates (Extended Data Fig. 5, see [63,64]).

When considering both the formation and elimination of all spines between days (daily turnover

ratio), oblique spines were remarkably stable, showing less than half the average turnover found

on basal dendrites in the same cell (Fig. 5d). When individual spines were categorized by

lifetime, where less than 4 days generally represents a "transient" synapse[64–68], a larger fraction

of basal dendritic spines had shorter lifetimes than oblique spines (Fig. 5a,e). Our results

demonstrate that L5b PNs in adult mice possess a dendritic domain with highly stable

synapses, in contrast to other dendritic domains with much higher rates of synaptic formation

and elimination.

Discussion

The plasticity of spiny synapses is widely considered to be the biological substrate of learning

and memory. The current model posits that smaller dendritic protrusions are involved in the

acquisition and refinement of information, whereas larger dendritic spines store acquired

information[58,63,65,67–70]. The long-term stability of specific spiny synapses is not well understood

and likely involves a combination of mechanisms, from network-level properties to molecular

regulation. Here, we show that entire dendritic domains can be highly stable and resistant to

long-term potentiation through a relative downregulation of synaptic NMDARs. These dendrites

have comparable spine densities and morphologies to neighboring, plastic dendritic

compartments but lack NMDAR-mediated supralinear integration and potentiation.

190   Our results do not rule out all possible plasticity mechanisms in oblique dendrites, but

limited NMDAR-mediated Ca^{2+} influx constrains their potential to engage conventional molecular

cascades[71]. In V1 L5b PNs, NMDAR-mediated properties decline in thalamorecipient dendrites

after a critical period of early visual experience, a process which can be prevented by delaying

the onset of vision. Thus, critical periods—the developmental window of experience required for

normal connectivity and function[50,72,73]—occur at the subcellular level, an unprecedented degree

of specificity, due to dendritic compartmentalization. Our findings establish a tractable new

model for future studies on the formation and regulation of stable dendritic domains, which likely

includes activity-dependent transcription, protein expression, and subcellular localization[74–78].

199   Protecting existing knowledge while continually acquiring new information is a long-

standing challenge for models of plasticity and biologically-inspired neural networks, which are

prone to instability and catastrophic forgetting[1,5]. To resolve this plasticity-stability dilemma,

some models decrease the plasticity of important weights[79–81]. Here we report the first biological

evidence for these theoretical limitations: after an initial strengthening in early development,

synapses on oblique dendritic spines are effectively stabilized through a domain-wide relative
decline in NMDAR expression and function. Our results are distinct from—but may interact
with—other models and observations that work to stabilize either plastic networks or individual
synapses[21,82–84]. In primary visual cortex, the stable dendritic domain receives input from the
visual thalamus, the definitive input to primary visual cortex. PNs in all layers of primary sensory
cortices receive both first-order thalamic and intracortical inputs[38,40,85,86], and this mechanism
may help sensory neurons maintain fundamental sensory representations throughout a lifetime
of learning and experience-dependent plasticity. Our results may also apply to other cortical
areas: the oblique dendritic domain in mouse association cortex similarly lacks NMDAR-
mediated supralinear integration[36]. Using this mechanism, cortical neurons may limit plasticity
within specific dendrites to maintain essential, stable representations without compromising
higher-order input plasticity. The restriction of plasticity within select dendritic compartments is a
powerful mechanism by which single neurons can solve the tradeoff between flexibility and
stability inherent to all learning systems.
218

Acknowledgements
We thank Mark Bear, Elly Nedivi, Josh Trachtenberg, and the Harnett laboratory for their
comments and suggestions and Kwanghun Chung for generously sharing equipment. This work
was funded by a Life Sciences Research Foundation postdoctoral fellowship (C.E.Y.), and a
Vallee Foundation Scholars Award, the James W. and Patricia T. Poitras Fund, and the National
Institutes of Health R01NS106031 (M.T.H.).
225

Author Contributions
C.E.Y. and M.T.H. conceived of the experiments. C.E.Y. performed patch-clamp physiology
experiments. D.V. conducted eMAP experiments. N.J.B. performed sCRACM viral injections
and histology. T.L.D.P performed viral injections and cranial window surgeries for in vivo two-
photon imaging, and Q.Z. performed deep-layer spine imaging with C.E.Y, supervised by N.J.
C.E.Y. carried out all analyses and prepared the figures. M.T.H. supervised all aspects of the
project and wrote the manuscript with C.E.Y.
233

Methods

Animals

All animal procedures were carried out in accordance with NIH guidelines for animal research

and were approved by the Massachusetts Institute of Technology Committee on Animal Care

and the Animal Care and Use Committee at the University of California, Berkeley. C57BL6 male

and female mice (Charles River Laboratories) were used in approximately equal numbers. Mice

8 weeks and older (P56+) were used for adult mouse experiments. For developmental

timepoints, mice ages P10-P14, P18-22, and P28-P32 were used. All animals were kept in

conventional social housing with unlimited food and water on a 12-hour light/dark cycle. For

dark-rearing experiments, pregnant dams were moved to a dark room at E18 and pups were

born and raised in complete darkness, with red light exposure during cage changes. Dark

reared mice were ages P28-P32 at the time of experimentation. 8-week-old Thy-1-GFP-M mice

(Jackson Laboratory) were used for eMAP experiments.

Slice preparation

Sucrose-containing artificial cerebrospinal fluid (sACSF) was used during the slicing procedure,

containing (in mM): 90 sucrose, 60 sodium chloride, 26.5 sodium bicarbonate, 2.75 potassium

chloride, 1.25 sodium phosphate, 9 glucose, 1.1 calcium chloride, and 4.1 magnesium chloride,

with an osmolality of 295-302. The sucrose solution was partially frozen to create a small

amount of slush and was kept ice-cold. Artificial cerebrospinal fluid (ACSF) was used for

recovery and recording, containing (in mM): 122 sodium chloride, 25 sodium bicarbonate, 3

potassium chloride, 1.25 sodium phosphate, 12 glucose, 1.2 calcium chloride, 1.2 magnesium

chloride, 1 ascorbate, and 3 sodium pyruvate, with an osmolality of 302-307. All solutions were

saturated with carbogen, 5% CO_2 and 95% O_2. Acute slice preparation was consistent for all

ages and experiments. Mice were put under isoflurane-induced anesthesia and decapitated.

The brain was extracted in ice-cold sucrose solution in less than one minute. The brain was

blocked at a moderate angle (approximately 20 degrees from coronal) to favor the preservation

of apical dendrites in the occipital cortex. After the brain was mounted and submerged in ice-

cold sACSF, a vibratome (Lieca VT1200S) was used to cut 300 μm-thick slices, which were

transferred to ACSF for 30-50 minutes at 36°C. Longer recovery times were used for tissue

collected from younger animals. Following recovery, slices were kept at room temperature.

Patch-clamp recording

Recordings were performed in ACSF (concentrations noted above) at 34–36 °C. Intracellular

recording solution contained (in mM): 134 potassium gluconate, 6 KCl, 10 HEPES, 4 NaCl, 4

Mg$_2$ATP, 0.3 NaGTP, and 14 phosphocreatine di(tris). Depending on the experiment, 0.05 mM

Alexa 594, 0.1 mM Alexa 488, and/or 0.1 mM OGB-1 (Invitrogen) were added to the internal

solution. An Olympus BX-61 microscope with Dodt optics and a water-immersion lens (60X, 0.9

NA; Olympus) was used to visualize cells. Whole-cell current-clamp recordings were obtained

with a Dagan BVC-700A amplifier. Patch pipettes with thin-wall glass (1.5/1.0mm OD/ID, WPI)

and resistances of 3-7 MΩ were used. Pipette capacitance was fully neutralized prior to break-

in, and series resistance was kept fully balanced throughout, ranging from 5-30 MΩ. The liquid

junction potential was not corrected. Current signals were digitized at 20 kHz and filtered at 10

kHz (Prairie View). L5b PNs were characterized by their large somas in layer 5, thick apical

dendrites, broad arborization in layer 1, low input resistance, and prominent voltage sag.

Virus injection

For sCRACM experiments, AAV2-hSyn-hChR2(H134R)-mCherry (UNC Vector Core) was

expressed in neurons in the dorsal lateral geniculate nucleus. Mice 6 weeks or older were

injected 4 weeks or more before slice preparation. Under isoflurane anesthesia, mice were

secured on a stereotaxic apparatus with a feedback-controlled heating pad (DC Temperature

Control System, FHC). Slow-release buprenorphine (1 mg/kg) was administered

subcutaneously. After scalp incision, a small burr hole was drilled over the injection site at the

following coordinates relative to bregma: anterior-posterior 2.7-2.8 mm; medio-lateral 2.3-2.4

mm. A beveled microinjection pipette containing was lowered to a depth of 2.7-2.9 mm, and

approximately 100 nL of virus was injected at a rate of 50 nL/min using a Nanoject. Following

virus injection and a five-minute rest, the pipette was removed and the incision was sutured.

Accuracy of the injection was assessed in acute slices using two-photon microscopy. For image

clarity, histology in Figure 1 was taken from a brain that was perfused, stored at 4°C overnight in

4% paraformaldehyde, transferred to PBS, and sectioned in 100 µm-thick slices. Sections were

mounted, coverslipped, and imaged under a confocal microscope (Zeiss LSM 710 with a 10x

objective, NA 0.45).

sCRACM

After 4 weeks or more of ChR2 expression in dLGN, acute slices of V1 were prepared. A two-

photon laser scanning system (Bruker), with dual galvanometers and a MaiTai DeepSee laser,

was used to confirm the presence of mCherry-expressing cell bodies in the dLGN. For all

recordings, TTX (1 µM) and 4-AP (100 µM) were added to ACSF to isolate monosynaptic

connections. Following whole-cell configuration, a 5 ms 473 nm full-field LED was used to

determine if the neuron received inputs from axons containing ChR2. If the LED drove EPSPs
or spikes, a 9 by 17 (400 x 800 µm) stimulation grid with 50 µm spacing was positioned over the
area containing the entire neuron, aligned at the pia using two-photon imaging under low
magnification (4x, 0.16 NA air objective, Olympus) and Prairie View software. The stimulation
grid controlled the location of a 473 nm laser beam (OptoEngine, LLC), and the duration of the
light pulses was kept to 1-3 ms under <2 mW power. Each point on the grid was stimulated in
order, progressing forward by column, with an interpoint delay of 1 s. 3-6 rounds of stimulation
were averaged for each neuron. Baseline voltage was defined as the 50 ms before stimulation,
and EPSP amplitudes were calculated as the baseline-subtracted maximum voltage within 50
ms after photostimulation. Population data was obtained by rotating the maps to align to the pia,
overlaying the maps to align at the soma, normalizing peak voltages, and averaging across
experiments.
318
Glutamate uncaging
Following whole-cell dialysis with a structural dye via the patch pipette, basal or apical oblique
branches were localized using two-photon imaging. A pipette containing caged glutamate (4-
methoxy-7-nitroindolinyl-caged-L-glutamate or MNI-glutamate) diluted in ACSF (10 mM) was
positioned just above the slice and over the recording site, and a picrospritzer (Picospritzer III,
Parker Hannifin) was used to puff a constant flow of caged glutamate into the region of interest.
A second Mai-Tai DeepSee laser controlled by Prairie View software was used for
photostimulation of MNI-glutamate at 720 nm. Uncaging was targeted just adjacent to spine
heads. In the imaging pathway, a linescan was used for simultaneous calcium imaging. Laser
intensity for both lasers was independently controlled (Conoptics). Linescan imaging was
performed at 1300 Hz, with a dwell time of 8 µs and a total scan time of less than 250 ms, with
baseline fluorescence kept minimal and monitored throughout. Cells with signs of photodamage
were excluded. The uncaging laser was calibrated to either a threshold calcium signal (~100%
ΔF/F) or an action potential, where the minimum power needed to drive either event was
identified and used for all points. The uncaging dwell time was 0.2 ms, and the uncaging interval
between multiple spines was 0.32 ms. Groups of 2-5 spines were stimulated independently and
then in combination with other groups up to a total of 25-40 spines. Expected values were
calculated by summing the average response of each group of spines. Ca^{2+} signals are
expressed as $(F - F_{baseline})/F_{baseline}$. For acute blockade of NMDA receptors, the competitive
antagonist D-AP5 (50 µM) was used.
339

Burst-timing dependent plasticity

For local branch stimulation, theta glass (2.0/1.4mm OD/ID) housing a bipolar stimulating

electrode (ISO-Flex, AMPI) was positioned within 10 µm of a basal or apical oblique branch.

Stimulation intensity was calibrated to generate a small EPSP of 1-3 mV, typically requiring 5-20

µA of current from the stimulating electrode. Ten minutes of baseline stimulation at 0.1 Hz was

recorded to ensure stability of the EPSP. A -50 pA hyperpolarizing step was included to

estimate input resistance. For plasticity induction, EPSPs were paired with a somatic current

injection of 400-700 pA for 20 ms duration to produce a train of 2-4 action potentials. The peak

of the first action potential was timed to be within 10 ms of the peak of the EPSP. Pairs were

done in sets of 5 at 10 Hz, and sets of 5 were repeated 30 times at 0.1 Hz. After the induction

period, EPSPs were monitored for at least 20 minutes. Inclusion criteria were as follows: resting

membrane potential could fluctuate from baseline by no more than 3 mV, input resistance could

not increase more than 20% of baseline; and series resistance had to be compensated fully

throughout (<30 MΩ). Recordings were tested for baseline stability post-hoc using either

correlation or linear regression (data not shown), and cells with significant changes in baseline

recording were not included. The necessity of pre-post pairing was shown using the same

recording set up but driving either the synaptic stimulation or the post-synaptic burst (and not

both) during the induction period. For acute blockade of NMDARs, the competitive antagonist D-

AP5 (50 µM) was used. Throughout plasticity experiments, picrotoxin (100 µM) was present in

the bath to block GABAergic transmission in acute slices from adult mice and at P28. 10 µM

picrotoxin was used for acute slices at P21, and no picrotoxin was needed at P14.

Epitope-preserving Magnified Analysis of the Proteome

Mice were perfused with cold PBS followed by cold 4% PFA while under deep anesthesia (5%

isoflurane). Brains were removed and kept in the same fixative overnight at 4°C and then

washed with PBS at 4°C for at least 1 day. 1.0 mm coronal slices of primary visual cortex were

cut on a vibratome and kept in PBS at 4°C until the day of processing. Slices were then

incubated eMAP hydrogel monomer solution (30% acrylamide [A9099, MilliporeSigma, St.

Louis, MO, USA], 10% sodium acrylate [408220, MilliporeSigma], 0.1% bis-acrylamide [161-

0142, Bio-Rad Laboratories, Hercules, CA, USA], and 0.03% VA-044 (w/v) [Wako Chemicals,

Richmond, VA, USA] in PBS]), protected from light, at 4°C overnight. For gelation, slices were

mounted between glass slides in eMAP solution and sealed in a 50 mL conical tube with

nitrogen gas at positive pressure of 10-12 psi at 37°C for 3 hours. The excess gel around the

slices was then removed. To reach a first expansion stage of 1.7x, the slices were incubated

overnight in a solution of 0.02% sodium azide (w/v) in PBS at 37°C. Slices were trimmed to

contain only parts of primary visual cortex and further sectioned with a vibratome to 75 μm

thickness (corresponding to ~40 μm thickness of the pre-expanded tissue). Slices containing

good candidate cells – L5 PNs whose apical trunk could be reconstructed at its full length in a

single slice or at most two consecutive slices – were selected during live low-resolution confocal

imaging sessions. These slices were trimmed to smallest possible samples of approximately 1.0

mm in both width and length. Slices were incubated in tissue clearing solution (6% SDS (w/v),

0.1 M phosphate buffer, 50 mM sodium sulfite, 0.02% sodium azide (w/v), pH 7.4) at 37°C for 6

hours, followed by incubation in preheated clearing solution at 95°C for 10 min. Cleared

samples were thoroughly washed with PBS + 0.1% Triton X at 37 °C. Primary antibody staining

was performed at 37°C overnight with the following antibodies: Anti-GFP (Life Technologies

A10262), Anti-NMDAR1 (SYSY 114011), Anti-AMPAR1 (SYSY 182003), Anti-AMPAR2 (SYSY

182103), and Anti-Bassoon (SYSY 141004). For secondary staining, the following fluorescent

antibodies were used: Bassoon: anti-Guinea pig-405 (AbCam ab175678); GFP: anti-Chicken-

488 (Invitrogen A11039); NMDAR1: anti-Mouse-AF+555 (Invitrogen A32727); and AMPAR1 and

AMPAR2: anti-Rabbit-AF+647 (Invitrogen A32733). Final expansion was performed just before

imaging by putting the trimmed slices in 0.1 mM tris in distilled water, and approximately 4X total

linear expansion was achieved. Slices were imaged using Leica TCS SP8 upright confocal

DM6000 microscope equipped with a 63x HC PL APO CS2/1.2 W objective, hybrid detectors,

and a white light laser. Within slices, intact L5b PNs were identified by their thick trunks and

broad apical tufts. Within single neurons, both basal and oblique dendritic branches were

imaged. To avoid photobleaching effects, no slice was imaged more than once, and basal and

oblique dendrites were imaged in alternating order for each cell. Dendritic protrusions were

analyzed using Fiji software. To draw regions of interest within spine heads in the GFP channel,

a custom-written macro code was used to apply a median blur (2 pixels), threshold the image,

and draw an ROI on the annotated protrusion. Using custom-written code in MATLAB, all color

channels were thresholded to include only signals 2 S.D. above the mean fluorescence

intensity, and signals from each channel were extracted from the ROI. The plane with peak

antibody fluorescence was identified within the spine head ROI, and each channel's

fluorescence signals within the plane were summed. Long, thin dendritic protrusions without

enlarged heads were classified as filopodia (head diameter/neck diameter < 1.3) and excluded

from the final analysis, as were spines with no detectable bassoon.

Ultrasparse eGFP expression in L5b PNs for deep layer structural imaging

Mice were approximately 8 weeks of age at the time of injection and cranial window surgery.

Prior to surgery, mice were subcutaneously injected with dexamethasone (4mg kg^{-1}) and

buprenorphine (slow-release, 1 mg kg^{-1}) and were anaesthetized with isoflurane (2% induction,

0.75-1.25% for maintenance). Mice were secured in a stereotaxic apparatus with a feedback-

controlled heating pad. Eye ointment was applied to prevent dryness (Bepanthen, Bayer), and

the scalp was shaved using hair removal cream and cleaned with iodine and ethanol. The skull

was exposed, etched, and sealed with a light-curing glue (Adhese Universal, Ivoclar). A

stainless steel headplate was secured to the skull with a light-curing flowable composite (Tetric

EvoFlow Translucent, Ivoclar). Then a 3 mm-wide craniotomy was performed over V1 (2.5 mm

lateral, 1 mm anterior to lambda). A 1:2 viral mixture of pENN.AAV.CamKII 0.4.Cre.SV40

(Addgene, diluted 1:100,000) and pAAV-hSyn-DIO-EGFP (Addgene) was injected at a volume

of 150 nl at 3 sites, approximately 200–500 µm apart, at a depth of 550 µm. Following virus

injection, the pipette was withdrawn after 5 minutes to avoid infecting superficial layers. Major

blood vessels were avoided. Cranial windows consisted of a single 3 mm coverslip attached to a

4.5 mm x 2.5 mm coverslip (170 µm thickness, custom-made, Potomac). The outer ring of the

cranial window was fixed to the skull with light-curing flowable composite (Tetric EvoFlow

Translucent, Ivoclar). Following 1-2 weeks of recovery from surgery, animals were transferred to

UC Berkeley and had 2 additional weeks of recovery.

Longitudinal spine tracking in deep cortical layers with two-photon microscopy

Ultrasparse eGFP expression (1-2 cells per injection site) allowed us to visualize L5b PN

dendritic spines 400-600 µm below the cortical surface with an optimized commercial two-

photon microscope (Thorlabs Bergamo® II Multiphoton). A femtosecond Ti:Sapphire laser

(Coherent, Chameleon Ultra II) tuned to 920 nm was used as the excitation source for all

experiments. A 25× water-immersion objective lens (Olympus, 1.05 NA, 2 mm WD) was used to

focus the excitation light into the brain and collect the emitted fluorescence. Data acquisition

and hardware were controlled using ThorImage software. Imaging started approximately 4

weeks after virus injection. Immediately prior to imaging, the mouse was anaesthetized (2%

isoflurane) and then head-fixed under the microscope with continuous anesthesia (0.5–1%

isoflurane). Body temperature was maintained using hand warmers. The headmount of the

mouse was aligned so that its cranial window was perpendicular to the excitation beam. The

correction collar of the objective lens was adjusted to minimize spherical aberration induced by

the cranial window. L5b PNs were identified by cortical depth, with somas ~500-600 µm below

the pia, and by their thick apical dendrites and broad arborization in layer 1. For each cell, z-

stacks were taken from the top-most oblique dendrite to the bottom-most basal dendrite (1 µm z

steps, 203 µm x 203 µm field of view, 0.1 µm x/y pixel^{-1}, 4 fps, 20 frames per step). The same

cells and branches were imaged daily for 5-10 days. Post-objective power ranged from 38-65

mW for different cells, and care was taken to ensure that fluorescence levels were comparable

throughout the tracking period.

Image processing and spine tracking analysis

Two-photon z-stacks were analyzed in ImageJ (v1.54f). Smaller stacks of specific basal and

oblique branches were sectioned from the original z-stack. Branches with steep changes in z

were avoided. Each substack was rigidly aligned, gamma adjusted (1.5), and gaussian filtered

(σ = 1). Branches from consecutive days were manually registered with a transparency tool

(Vitrite), and similar lookup tables were used for each timepoint. Branches were only included if

the dimmest protrusion was at least 5x higher than the standard deviation of the background

around the dendrite. Dendritic protrusions were scored on maximum intensity z-projections and

cross-checked with the original z-stack. The following criteria was used for scoring: protrusions

must extend laterally, more than 0.5 µm from the branch. Protrusions that did not reach criteria

at any point during the experiment were excluded from analysis. Daily tracking facilitated

consistent identification of structures, even with small movements of the tissue. With this

dataset, we calculated the fraction of protrusions that were present across the entire imaging

period (survival fraction, SF) and the turnover ratio (TOR), defined as the fraction of structures

that appear and/or disappear between days, as in [89] $SF(t) = N(t)/N_0$, where N_0 is the number of

structures at t = 0, and $N(t)$ is the number of structures of the original set surviving after time t.

TOR $(t_1, t_2) = (N_{gained} + N_{lost})/(N(t_1) + N(t_2))$, where $N(t_1)$ and $N(t_2)$ are the total number of

structures at the first and second time point. The lifetime of a given structure is the number of

days present within the recording window.

Data availability

All data and custom-written analysis code are available upon request.

References

1. Parisi, G.I., Kemker, R., Part, J.L., Kanan, C., and Wermter, S. (2019). Continual lifelong learning with neural networks: A review. Neural Netw *113*, 54–71. 10.1016/j.neunet.2019.01.012.

2. Mermillod, M., Bugaiska, A., and Bonin, P. (2013). The stability-plasticity dilemma: investigating the continuum from catastrophic forgetting to age-limited learning effects. Front Psychol *4*, 504. 10.3389/fpsyg.2013.00504.

3. Grossberg, S. (1980). How does a brain build a cognitive code? Psychol Rev *87*, 1–51. 10.1007/978-94-009-7758-7_1.

4. Grossberg, S. (2013). Adaptive Resonance Theory: how a brain learns to consciously attend, learn, and recognize a changing world. Neural Netw *37*, 1–47. 10.1016/j.neunet.2012.09.017.

5. Hassabis, D., Kumaran, D., Summerfield, C., and Botvinick, M. (2017). Neuroscience-Inspired Artificial Intelligence. Neuron *95*, 245–258. 10.1016/j.neuron.2017.06.011.

6. McClelland, J.L., and Rumelhart, D.E. (1985). Distributed memory and the representation of general and specific information. J Exp Psychol Gen *114*, 159–197. 10.1037/0096-3445.114.2.159.

7. McCloskey, M., and Cohen, N.J. (1989). Catastrophic Interference in Connectionist Networks: The Sequential Learning Problem. In Psychology of Learning and Motivation (Elsevier), pp. 109–165. 10.1016/S0079-7421(08)60536-8.

8. Markram, H., Lübke, J., Frotscher, M., and Sakmann, B. (1997). Regulation of synaptic efficacy by coincidence of postsynaptic APs and EPSPs. Science *275*, 213–215. 10.1126/science.275.5297.213.

9. Golding, N.L., Staff, N.P., and Spruston, N. (2002). Dendritic spikes as a mechanism for cooperative long-term potentiation. Nature *418*, 326–331. 10.1038/nature00854.

10. Froemke, R.C., Poo, M.-M., and Dan, Y. (2005). Spike-timing-dependent synaptic plasticity depends on dendritic location. Nature *434*, 221–225. 10.1038/nature03366.

11. Kampa, B.M., Letzkus, J.J., and Stuart, G.J. (2006). Requirement of dendritic calcium spikes for induction of spike-timing-dependent synaptic plasticity. J Physiol (Lond) *574*, 283–290. 10.1113/jphysiol.2006.111062.

12. Gordon, U., Polsky, A., and Schiller, J. (2006). Plasticity compartments in basal dendrites of neocortical pyramidal neurons. J Neurosci *26*, 12717–12726. 10.1523/JNEUROSCI.3502-06.2006.

13. Letzkus, J.J., Kampa, B.M., and Stuart, G.J. (2006). Learning rules for spike timing-dependent plasticity depend on dendritic synapse location. J Neurosci *26*, 10420–10429. 10.1523/JNEUROSCI.2650-06.2006.

14. Sjöström, P.J., and Häusser, M. (2006). A cooperative switch determines the sign of
synaptic plasticity in distal dendrites of neocortical pyramidal neurons. Neuron *51*, 227–238.
10.1016/j.neuron.2006.06.017.

15. Weber, J.P., Andrásfalvy, B.K., Polito, M., Magó, Á., Ujfalussy, B.B., and Makara, J.K.
(2016). Location-dependent synaptic plasticity rules by dendritic spine cooperativity. Nat
Commun *7*, 11380. 10.1038/ncomms11380.

16. Magó, Á., Weber, J.P., Ujfalussy, B.B., and Makara, J.K. (2020). Synaptic Plasticity
Depends on the Fine-Scale Input Pattern in Thin Dendrites of CA1 Pyramidal Neurons. J.
Neurosci. *40*, 2593–2605. 10.1523/JNEUROSCI.2071-19.2020.

17. Kumar, A., Barkai, E., and Schiller, J. (2021). Plasticity of olfactory bulb inputs mediated by
dendritic NMDA-spikes in rodent piriform cortex. eLife *10*. 10.7554/eLife.70383.

18. Bienenstock, E.L., Cooper, L.N., and Munro, P.W. (1982). Theory for the development of
neuron selectivity: orientation specificity and binocular interaction in visual cortex. J
Neurosci *2*, 32–48. 10.1142/9789812795885_0006.

19. Cooper, L.N., and Bear, M.F. (2012). The BCM theory of synapse modification at 30:
interaction of theory with experiment. Nat Rev Neurosci *13*, 798–810. 10.1038/nrn3353.

20. Abraham, W.C. (2008). Metaplasticity: tuning synapses and networks for plasticity. Nat Rev
Neurosci *9*, 387. 10.1038/nrn2356.

21. Turrigiano, G.G. (2008). The self-tuning neuron: synaptic scaling of excitatory synapses.
Cell *135*, 422–435. 10.1016/j.cell.2008.10.008.

22. Li, J., Park, E., Zhong, L.R., and Chen, L. (2019). Homeostatic synaptic plasticity as a
metaplasticity mechanism - a molecular and cellular perspective. Curr Opin Neurobiol *54*,
44–53. 10.1016/j.conb.2018.08.010.

23. Takesian, A.E., and Hensch, T.K. (2013). Balancing plasticity/stability across brain
development. Prog Brain Res *207*, 3–34. 10.1016/B978-0-444-63327-9.00001-1.

24. Faguet, J., Maranhao, B., Smith, S.L., and Trachtenberg, J.T. (2009). Ipsilateral eye cortical
maps are uniquely sensitive to binocular plasticity. J Neurophysiol *101*, 855–861.
10.1152/jn.90893.2008.

25. Gu, Y., and Cang, J. (2016). Binocular matching of thalamocortical and intracortical circuits
in the mouse visual cortex. eLife *5*. 10.7554/eLife.22032.

26. Leighton, A.H., and Lohmann, C. (2016). The Wiring of Developing Sensory Circuits-From
Patterned Spontaneous Activity to Synaptic Plasticity Mechanisms. Front Neural Circuits *10*,
71. 10.3389/fncir.2016.00071.

27. Wiesel, T., and Hubel, D. (1963). Single-cell responses in striate cortex of kittens deprived
of vision in one eye. J Neurophysiol *26*, 1003–1017. 10.1152/jn.1963.26.6.1003.

28. Tan, L., Tring, E., Ringach, D.L., Zipursky, S.L., and Trachtenberg, J.T. (2020). Vision
Changes the Cellular Composition of Binocular Circuitry during the Critical Period. Neuron
108, 735-747.e6. 10.1016/j.neuron.2020.09.022.

29. Wang, B.-S., Sarnaik, R., and Cang, J. (2010). Critical period plasticity matches binocular
orientation preference in the visual cortex. Neuron *65*, 246–256.
10.1016/j.neuron.2010.01.002.

30. Rose, T., Jaepel, J., Hübener, M., and Bonhoeffer, T. (2016). Cell-specific restoration of
stimulus preference after monocular deprivation in the visual cortex. Science *352*, 1319–
1322. 10.1126/science.aad3358.

31. Gordon, J.A., and Stryker, M.P. (1996). Experience-dependent plasticity of binocular
responses in the primary visual cortex of the mouse. J Neurosci *16*, 3274–3286.

32. Makino, H., and Komiyama, T. (2015). Learning enhances the relative impact of top-down
processing in the visual cortex. Nat Neurosci *18*, 1116–1122. 10.1038/nn.4061.

33. Pakan, J.M.P., Currie, S.P., Fischer, L., and Rochefort, N.L. (2018). The impact of visual
cues, reward, and motor feedback on the representation of behaviorally relevant spatial
locations in primary visual cortex. Cell Rep *24*, 2521–2528. 10.1016/j.celrep.2018.08.010.

34. Poort, J., Khan, A.G., Pachitariu, M., Nemri, A., Orsolic, I., Krupic, J., Bauza, M., Sahani, M.,
Keller, G.B., Mrsic-Flogel, T.D., et al. (2015). Learning Enhances Sensory and Multiple Non-
sensory Representations in Primary Visual Cortex. Neuron *86*, 1478–1490.
10.1016/j.neuron.2015.05.037.

35. Weskelblatt, J.B., and Niell, C.M. (2019). Distinct functional classes of excitatory neurons in
mouse V1 are differentially modulated by learning and task engagement. BioRxiv.
10.1101/533463.

36. Lafourcade, M., van der Goes, M.-S.H., Vardalaki, D., Brown, N.J., Voigts, J., Yun, D.H.,
Kim, M.E., Ku, T., and Harnett, M.T. (2022). Differential dendritic integration of long-range
inputs in association cortex via subcellular changes in synaptic AMPA-to-NMDA receptor
ratio. Neuron *110*, 1532-1546.e4. 10.1016/j.neuron.2022.01.025.

37. Young, H., Belbut, B., Baeta, M., and Petreanu, L. (2021). Laminar-specific cortico-cortical
loops in mouse visual cortex. eLife *10*. 10.7554/eLife.59551.

38. Morgenstern, N.A., Bourg, J., and Petreanu, L. (2016). Multilaminar networks of cortical
neurons integrate common inputs from sensory thalamus. Nat Neurosci *19*, 1034–1040.
10.1038/nn.4339.

39. Petreanu, L., Mao, T., Sternson, S.M., and Svoboda, K. (2009). The subcellular organization
of neocortical excitatory connections. Nature *457*, 1142–1145. 10.1038/nature07709.

40. Antonini, A., Fagiolini, M., and Stryker, M.P. (1999). Anatomical correlates of functional
plasticity in mouse visual cortex. J Neurosci *19*, 4388–4406.

41. Constantinople, C.M., and Bruno, R.M. (2013). Deep cortical layers are activated directly by
thalamus. Science *340*, 1591–1594. 10.1126/science.1236425.

42. Ferster, D., and Lindström, S. (1983). An intracellular analysis of geniculo-cortical
connectivity in area 17 of the cat. J Physiol (Lond) *342*, 181–215.
10.1113/jphysiol.1983.sp014846.

43. Kim, E.J., Juavinett, A.L., Kyubwa, E.M., Jacobs, M.W., and Callaway, E.M. (2015). Three
Types of Cortical Layer 5 Neurons That Differ in Brain-wide Connectivity and Function.
Neuron *88*, 1253–1267. 10.1016/j.neuron.2015.11.002.

44. Peters, A., and Feldman, M.L. (1977). The projection of the lateral geniculate nucleus to
area 17 of the rat cerebral cortex. IV. Terminations upon spiny dendrites. J Neurocytol *6*,
669–689. 10.1007/bf01176379.

45. Schiller, J., Major, G., Koester, H.J., and Schiller, Y. (2000). NMDA spikes in basal dendrites
of cortical pyramidal neurons. Nature *404*, 285–289. 10.1038/35005094.

46. Kirkwood, A., Lee, H.K., and Bear, M.F. (1995). Co-regulation of long-term potentiation and
experience-dependent synaptic plasticity in visual cortex by age and experience. Nature
375, 328–331. 10.1038/375328a0.

47. Crair, M.C., and Malenka, R.C. (1995). A critical period for long-term potentiation at
thalamocortical synapses. Nature *375*, 325–328. 10.1038/375325a0.

48. Sommeijer, J.-P., Ahmadlou, M., Saiepour, M.H., Seignette, K., Min, R., Heimel, J.A., and
Levelt, C.N. (2017). Thalamic inhibition regulates critical-period plasticity in visual cortex and
thalamus. Nat Neurosci *20*, 1715–1721. 10.1038/s41593-017-0002-3.

49. Cang, J., Rentería, R.C., Kaneko, M., Liu, X., Copenhagen, D.R., and Stryker, M.P. (2005).
Development of precise maps in visual cortex requires patterned spontaneous activity in the
retina. Neuron *48*, 797–809. 10.1016/j.neuron.2005.09.015.

50. Feldman, D.E. (2009). Synaptic mechanisms for plasticity in neocortex. Annu Rev Neurosci
32, 33–55. 10.1146/annurev.neuro.051508.135516.

51. Kumar, S.S., Bacci, A., Kharazia, V., and Huguenard, J.R. (2002). A developmental switch
of AMPA receptor subunits in neocortical pyramidal neurons. J Neurosci *22*, 3005–3015.

52. Hadzic, M., Jack, A., and Wahle, P. (2017). Ionotropic glutamate receptors: Which ones,
when, and where in the mammalian neocortex. J Comp Neurol *525*, 976–1033.
10.1002/cne.24103.

53. Ku, T., Swaney, J., Park, J.-Y., Albanese, A., Murray, E., Cho, J.H., Park, Y.-G., Mangena,
V., Chen, J., and Chung, K. (2016). Multiplexed and scalable super-resolution imaging of
three-dimensional protein localization in size-adjustable tissues. Nat Biotechnol *34*, 973–
981. 10.1038/nbt.3641.

54. Park, J., Khan, S., Yun, D.H., Ku, T., Villa, K.L., Lee, J.E., Zhang, Q., Park, J., Feng, G.,
Nedivi, E., et al. (2021). Epitope-preserving magnified analysis of proteome (eMAP). Sci
Adv *7*, eabf6589. 10.1126/sciadv.abf6589.

55. Cheng, D., Hoogenraad, C.C., Rush, J., Ramm, E., Schlager, M.A., Duong, D.M., Xu, P.,
Wijayawardana, S.R., Hanfelt, J., Nakagawa, T., et al. (2006). Relative and absolute

quantification of postsynaptic density proteome isolated from rat forebrain and cerebellum. Mol Cell Proteomics *5*, 1158–1170. 10.1074/mcp.D500009-MCP200.

56. Buchs, P.A., and Muller, D. (1996). Induction of long-term potentiation is associated with major ultrastructural changes of activated synapses. Proceedings of the National Academy of Sciences *93*, 8040–8045. 10.1073/pnas.93.15.8040.

57. Maletic-Savatic, M., Malinow, R., and Svoboda, K. (1999). Rapid Dendritic Morphogenesis in CA1 Hippocampal Dendrites Induced by Synaptic Activity. Science *283*, 1923–1927. 10.1126/science.283.5409.1923.

58. Matsuzaki, M., Honkura, N., Ellis-Davies, G.C.R., and Kasai, H. (2004). Structural basis of long-term potentiation in single dendritic spines. Nature *429*, 761–766. 10.1038/nature02617.

59. Zhou, Q., Homma, K.J., and Poo, M. (2004). Shrinkage of Dendritic Spines Associated with Long-Term Depression of Hippocampal Synapses. Neuron *44*, 749–757. 10.1016/j.neuron.2004.11.011.

60. Nägerl, U.V., Eberhorn, N., Cambridge, S.B., and Bonhoeffer, T. (2004). Bidirectional Activity-Dependent Morphological Plasticity in Hippocampal Neurons. Neuron *44*, 759–767. 10.1016/j.neuron.2004.11.016.

61. Kwon, H.-B., and Sabatini, B.L. (2011). Glutamate induces de novo growth of functional spines in developing cortex. Nature *474*, 100–104. 10.1038/nature09986.

62. Wiegert, J.S., and Oertner, T.G. (2013). Long-term depression triggers the selective elimination of weakly integrated synapses. Proceedings of the National Academy of Sciences *110*, E4510–E4519. 10.1073/pnas.1315926110.

63. Holtmaat, A.J.G.D., Trachtenberg, J.T., Wilbrecht, L., Shepherd, G.M., Zhang, X., Knott, G.W., and Svoboda, K. (2005). Transient and persistent dendritic spines in the neocortex in vivo. Neuron *45*, 279–291. 10.1016/j.neuron.2005.01.003.

64. Keck, T., Mrsic-Flogel, T.D., Vaz Afonso, M., Eysel, U.T., Bonhoeffer, T., and Hübener, M. (2008). Massive restructuring of neuronal circuits during functional reorganization of adult visual cortex. Nat Neurosci *11*, 1162–1167. 10.1038/nn.2181.

65. Trachtenberg, J.T., Chen, B.E., Knott, G.W., Feng, G., Sanes, J.R., Welker, E., and Svoboda, K. (2002). Long-term in vivo imaging of experience-dependent synaptic plasticity in adult cortex. Nature *420*, 788–794. 10.1038/nature01273.

66. Holtmaat, A., and Caroni, P. (2016). Functional and structural underpinnings of neuronal assembly formation in learning. Nat Neurosci *19*, 1553–1562. 10.1038/nn.4418.

67. Cane, M., Maco, B., Knott, G., and Holtmaat, A. (2014). The Relationship between PSD-95 Clustering and Spine Stability In Vivo. J. Neurosci. *34*, 2075–2086. 10.1523/JNEUROSCI.3353-13.2014.

68. Berry, K.P., and Nedivi, E. (2017). Spine Dynamics: Are They All the Same? Neuron *96*, 43–55. 10.1016/j.neuron.2017.08.008.

69. Grutzendler, J., Kasthuri, N., and Gan, W.-B. (2002). Long-term dendritic spine stability in
the adult cortex. Nature *420*, 812–816. 10.1038/nature01276.

70. Vardalaki, D., Chung, K., and Harnett, M.T. (2022). Filopodia are a structural substrate for
silent synapses in adult neocortex. Nature *612*, 323–327. 10.1038/s41586-022-05483-6.

71. Malenka, R.C., and Bear, M.F. (2004). LTP and LTD: an embarrassment of riches. Neuron
44, 5–21. 10.1016/j.neuron.2004.09.012.

72. Daw, N.W. (2009). The foundations of development and deprivation in the visual system. J
Physiol (Lond) *587*, 2769–2773. 10.1113/jphysiol.2009.170001.

73. Hooks, B.M., and Chen, C. (2020). Circuitry Underlying Experience-Dependent Plasticity in
the Mouse Visual System. Neuron *106*, 21–36. 10.1016/j.neuron.2020.01.031.

74. Liu, S.Q., and Cull-Candy, S.G. (2000). Synaptic activity at calcium-permeable AMPA
receptors induces a switch in receptor subtype. Nature *405*, 454–458. 10.1038/35013064.

75. Clem, R.L., Celikel, T., and Barth, A.L. (2008). Ongoing in vivo experience triggers synaptic
metaplasticity in the neocortex. Science *319*, 101–104. 10.1126/science.1143808.

76. Job, C., and Eberwine, J. (2001). Localization and translation of mRNA in dentrites and
axons. Nat Rev Neurosci *2*, 889–898. 10.1038/35104069.

77. Sutton, M.A., and Schuman, E.M. (2006). Dendritic Protein Synthesis, Synaptic Plasticity,
and Memory. Cell *127*, 49–58. 10.1016/j.cell.2006.09.014.

78. Holt, C.E., and Schuman, E.M. (2013). The Central Dogma Decentralized: New
Perspectives on RNA Function and Local Translation in Neurons. Neuron *80*, 648–657.
10.1016/j.neuron.2013.10.036.

79. Fusi, S., Drew, P.J., and Abbott, L.F. (2005). Cascade models of synaptically stored
memories. Neuron *45*, 599–611. 10.1016/j.neuron.2005.02.001.

80. Benna, M.K., and Fusi, S. (2016). Computational principles of synaptic memory
consolidation. Nat Neurosci *19*, 1697–1706. 10.1038/nn.4401.

81. Kirkpatrick, J., Pascanu, R., Rabinowitz, N., Veness, J., Desjardins, G., Rusu, A.A., Milan,
K., Quan, J., Ramalho, T., Grabska-Barwinska, A., et al. (2017). Overcoming catastrophic
forgetting in neural networks. Proceedings of the National Academy of Sciences *114*, 3521–
3526. 10.1073/pnas.1611835114.

82. Lee, M.-C., Yasuda, R., and Ehlers, M.D. (2010). Metaplasticity at single glutamatergic
synapses. Neuron *66*, 859–870. 10.1016/j.neuron.2010.05.015.

83. Abraham, W.C., and Bear, M.F. (1996). Metaplasticity: the plasticity of synaptic plasticity.
Trends Neurosci *19*, 126–130. 10.1016/s0166-2236(96)80018-x.

84. Lee, H.-K., and Kirkwood, A. (2019). Mechanisms of Homeostatic Synaptic Plasticity in vivo.
Front Cell Neurosci *13*, 520. 10.3389/fncel.2019.00520.

85. Harris, K.D., and Shepherd, G.M.G. (2015). The neocortical circuit: themes and variations.
Nat Neurosci *18*, 170–181. 10.1038/nn.3917.

86. Cruz-Martín, A., El-Danaf, R.N., Osakada, F., Sriram, B., Dhande, O.S., Nguyen, P.L.,
Callaway, E.M., Ghosh, A., and Huberman, A.D. (2014). A dedicated circuit links direction-
selective retinal ganglion cells to the primary visual cortex. Nature *507*, 358–361.
10.1038/nature12989.

Figure 1: Apical oblique dendrites of adult V1 L5b PNs selectively receive dLGN input and do not have NMDAR-mediated integrative properties.

a. Example confocal image of a coronal brain slice showing dLGN neurons infected with AAV2-hSyn-mCherry-ChR2. Scale bar: 500 µm.

b. Left: Example two-photon image of a V1 L5b PN whole-cell recording with overlaid photostimulation grid (blue). Scale bar, 50 µm. Middle: Average evoked EPSPs from 6 repetitions of 473 nm photostimulation for the neuron and grid shown. Soma position is indicated by the triangle. Right: Heatmap of average evoked EPSPs. Each square is 50 µm.

c. Left: Normalized heatmap of pooled L5b PNs (n = 9 neurons, 9 slices, 3 P56+ animals), aligned at the soma. Each square is 50 µm. Right: Mean normalized input as a function of depth for all neurons. Shaded region indicates the bootstrapped 95% confidence interval.

d. Example two-photon z-stack of a L5b PN with glutamate uncaging sites indicated on an oblique (purple) and a basal (green) branch, scale bar: 50 µm. Uncaging sites (stars) and linescan (yellow) are magnified in insets (scale bars: 10 µm).

e. Expected EPSPs, measured EPSPs, and local branch Ca²⁺ signals for oblique (top) and basal (bottom) dendrites shown in d. Left: Expected voltages from increasing numbers of linearly summed inputs. Middle: Measured EPSPs in response to increasing numbers of synchronous uncaging inputs. Stimulation of all inputs led to an action potential (AP, arrow). Right: Corresponding changes in local branch OGB-1 signal (ΔF/F).

f. Measured voltage (left) and branch ΔF/F (right) as a function of expected voltage for the branches shown in d. Dashed line indicates linearity (left) and threshold Ca²⁺ signal (right).

g. Adult population gain (measured/expected, left) and branch ΔF/F (right). n = 13 basal branches from 7 animals and 9 oblique branches from 6 animals. All animals were P56+. Basal APs removed for clarity. Basal vs oblique gain (AP excluded), Mann-Whitney U test: p = 7.63E-04; basal vs oblique ΔF/F (AP excluded), Mann-Whitney U test: p = 1.08E-05.

Figure 2: Oblique dendrites in adult V1 L5b PNs lack long-term synaptic potentiation, unlike basal dendrites.

a. Example two-photon z-stack of whole-cell recording with focal stimulation (green) near a basal dendrite. Scale bar: 25 µm.
b. As in a, but for an oblique dendrite (purple).
c. Plasticity induction protocol. 5 pairings of pre- and post-synaptic stimulation at 100 Hz, repeated 30 times at 0.1 Hz. Right: a 10 ms interval between the peak of the evoked EPSP and the peak of the first action potential. Scale bar: 20 mV, 50 ms; inset: 5 mV, 5 ms.
d. Example recording from the basal dendrite shown in a. After 10 minutes of baseline recording, the induction protocol is delivered (arrow). Inset shows average EPSP before (black) and after (color) pairing. Scale bar: 1 mV, 25 ms. Responses are 1 min binned averages and normalized to baseline.
e. As in d, for the oblique dendrite shown in b.
f. Normalized EPSP amplitude for neurons stimulated at basal (green) or oblique (purple) branches (2 min bins; n = 10 basal branches, 10 animals and 13 oblique branches, 12 animals. All animals were P56+.). Error bars are s.e.m. Basal vs. oblique post-induction EPSP, Mann-Whitney U test: p = 4.95E-63.
g. Change in EPSP after induction for all oblique branches, basal branches, and basal branches with DAP5 or with only presynaptic or postsynaptic activation during induction. Basal vs. oblique average change in EPSP, Mann-Whitney U test: p = 6.33E-05. All other comparisons not significant. Boxplot: median, 25^{th}, and 75^{th} percentiles with +/–2.7σ whiskers.

Figure 3: Oblique dendrites lose NMDAR-mediated properties after an experience-dependent critical period in postnatal development.

a. Expected (left), measured (middle), and ΔF/F (right) for oblique (top) and basal (bottom) dendrites from the same neuron from a P14 mouse. Note supralinear integration and large Ca^{2+} signals in both branches. Both are driven to action potential initiation (AP, arrow).

b. P14 population gain (measured/expected, left) and branch ΔF/F (right) (n = 9 basal branches and 9 oblique branches, both from 5 animals). Basal vs oblique gain*, p = 0.33; ΔF/F*, p = 0.70.

c. Maximum gain (left) and local branch ΔF/F (prior to AP initiation, right) across developmental time points (Adult, P56+: n as in Fig. 1, basal vs oblique gain*, p = 1.07E-04; ΔF/F*, p = 1.07E-04. P14: n as in b, basal vs oblique gain*, p = 0.54; ΔF/F*, p = 0.76. P21: n = 9 basal branches, 4 animals, 6 oblique branches, 2 animals, basal vs oblique gain*, p = 0.52; ΔF/F*, p = 0.68. P28: n = 10 basal branches, 5 animals; 9 oblique branches, 5 animals, basal vs oblique gain*, p = 2.17E-05; ΔF/F*: p = 4.33E-05. Boxplot parameters as in Fig. 2.

d. Plasticity (as described in Fig. 2) in basal versus oblique dendrites at P14 (n = 5 basal branches from 5 animals and 5 oblique branches from 5 animals, basal vs oblique post-induction*, p = 0.79), P21 (n= 5 basal branches from 4 animals and 7 oblique branches from 6 animals, basal vs oblique post-induction*, p = 0.87), and P28 (n = 7 basal branches, 5 animals; 7 oblique branches, 4 animals, basal vs oblique post-induction*, p = 1.25E-13).

e. P28 population gain (measured/expected, left) and branch ΔF/F (middle) in dark-reared (DR) mice (n = 9 basal branches and 10 oblique branches from 4 animals, basal vs oblique gain*, p = 0.78; ΔF/F*, p = 0.96). Plasticity for dark-reared animals (right) (n = 5 apical branches from 4 animals and 6 oblique branches from 6 animals, basal vs oblique post-induction*, p = 0.81).

*Mann-Whitney U test was used for all comparisons.

Figure 4: Changes in synaptic AMPA-to-NMDA ratio at oblique versus basal dendrites underlie differences in integration and plasticity.

a. Example two-photon image of a V1 L5b PN from a P21 mouse (scale bar: 50 μm). An oblique dendrite (circled) is shown at higher magnification (right, scale bar: 5 μm). Uncaging site indicated by star.

b. Representative average EPSPs from uncaging at single spines in P21 (left) or adult (right) mice from oblique (top) and basal (bottom) dendrites. Black traces: control, colored traces: the same spine in DNQX+Mg^{2+}-free ACSF.

c. Functional single spine AMPA/NMDA from basal and oblique dendrites in P21 and adult mice (P21: n = 72 basal spines from 4 mice and 89 oblique spines from 4 mice. Adult, P56+: n = 66 basal spines from 4 mice and 67 oblique spines from 5 mice). Outliers are not shown. P21 basal vs oblique, Mann-Whitney U test: p = 0.23; adult basal vs oblique, Mann-Whitney U test: p = 1.00E-04. Boxplot parameters as in Fig. 2.

d. Schematic of the epitope-preserving Magnified Analysis of the Proteome (eMAP) experimental procedure.

e. Confocal image of L5 PNs from a Thy1-GFP M P56+ mouse after 4x eMAP expansion. Scale bar: 300 μm.

f. Example confocal images of expanded oblique (top) and basal (bottom) dendrites from the same neuron. For each branch, immunostaining for bassoon, NR1, and GluA1+2 is shown for one spine (indicated by star). Scale bar: 10 μm, inset: 2 μm.

g. Ratio of AMPA/NMDA intensities for all spines: Mann-Whitney U test: p = 8.30E-11.

h. Ratio of AMPA/NMDA intensities for the largest 10% of basal and oblique spines (n = 131 basal dendrite spines and 171 oblique dendrite spines). Mann-Whitney U test: p = 1.08E-04.

798
Figure 5: In vivo synaptic stability and flexibility at distinct dendritic compartments.
a. 3D projection of an eGFP-expressing L5b PN, imaged in vivo with two-photon microscopy.
Scale bar: 50 µm.
b. Representative oblique (top) and basal (bottom) dendrites from the same layer 5 PN imaged
across 7 days. All transient spines (with lifetimes of 4 days or fewer) are labeled
(arrowheads) on the day prior to disappearance. Scale bar: 10 µm.
c. Average spine density per cell for basal and oblique dendrites. Wilcoxon signed-rank, p =
0.56.
d. Average daily turnover, per cell. Wilcoxon signed-rank, p = 0.03.
e. Fraction of transient (≤ 4 day lifetime) and persistent (≥ 5 day lifetime) spines on oblique
(purple) and basal (green) dendrites, per cell. Friedman's test, p = 1.07E-06.
For b-e, n = 6 cells with 775 basal spines and 794 oblique spines from 4 animals.