Dendritic compartmentalization of input-specific integration and plasticity rules across cortical development

Courtney E. Yaeger, Dimitra Vardalaki, Norma J. Brown, & Mark T. Harnett[1]

[1]McGovern Institute for Brain Research and Department of Brain and Cognitive Sciences, MIT, Cambridge, MA, USA

To whom correspondence should be addressed: Harnett (harnett@mit.edu)

Abstract

The modification of synaptic weights is critical for learning, while synaptic stability is required to maintain acquired knowledge. Single neurons have thousands of synapses within their dendritic arbors, and how the weights of specific inputs change across experience is poorly understood. Here we report that dendritic compartments receiving input from different presynaptic populations acquire distinct synaptic plasticity and integration rules across maturation. We find that apical oblique dendrites of layer 5 pyramidal neurons in adult mouse primary visual cortex receive direct monosynaptic projections from the dorsal lateral geniculate nucleus (dLGN), linearly integrate input, and lack synaptic potentiation. In contrast, basal dendrites, which do not receive dLGN input, exhibit NMDA receptor (NMDAR)-mediated supralinear integration and synaptic potentiation. Earlier in development, during thalamic input refinement, oblique and basal dendrites exhibited comparable NMDAR-dependent properties. Oblique dendrites gain mature properties with visual experience, and over the course of maturation, spines on oblique dendrites develop higher AMPA/NMDA ratios relative to basal dendrites. Our results demonstrate that cortical neurons possess dendrite-specific integration and plasticity rules that are set by the activity of their inputs. The linear, non-plastic nature of mature synapses on oblique dendrites may stabilize feedforward sensory processing while synaptic weights in other parts of the dendritic tree remain plastic, facilitating robust yet flexible cortical computation in adults.

Introduction

Adaptive behavior requires balancing the acquisition of new information with the retention of previously learned associations. Despite over half a century of work on synaptic plasticity, how neurons solve this tradeoff remains unknown. In adult visual cortex, primary sensory inputs are stable during normal visual experience[1,2–7], while cortical representations are readily remodeled[8–11]. Dendrites of cortical neurons receive both primary sensory and intracortical inputs[12–16] and play a key role in both synaptic plasticity and input compartmentalization. Convergent spatiotemporal input patterns arriving at a dendritic compartment can drive NMDAR-mediated calcium influx[17] and trigger molecular cascades for synaptic modification[18]. Afferents from specific presynaptic populations can also target particular dendritic compartments[16,19,20], where plasticity rules can be location-dependent[21–27]. However, it is unclear how distinct inputs utilize these mechanisms, and less is understood how input-specific synaptic weights are maintained across experience[28–30]. To investigate this stability-flexibility tradeoff for primary sensory cortical synapses, we first identified dLGN-recipient and non-recipient dendritic compartments in layer 5 pyramidal neurons (L5 PNs) in mouse primary visual cortex (V1). We compared synaptic integration and long-term potentiation of synapses at these two compartments in adults and across postnatal development.

Results

L5 PN oblique dendrites selectively receive thalamic input and exhibit distinct integrative properties from basal dendrites

Visual information from dLGN[13] projects primarily to V1 layer 4 cells. In an analogous sensory cortex, first-order thalamic inputs also synapse on L5 PNs[31], which integrate inputs from all cortical layers. Physiological data from juvenile V1[14] and anatomical data[32] suggest that dLGN inputs target the apical oblique dendrites of L5 PNs, which can reside in layer 4. To directly test where dLGN inputs make functional synapses on L5 PNs in mouse V1, we employed subcellular channelrhodopsin (ChR2)-assisted circuit mapping[19]. After viral expression of ChR2 in dLGN (Fig. 1a), we performed whole-cell patch-clamp recordings from V1 L5 PNs in acute slices from adult mice (P56+). Local photostimulation of ChR2-expressing dLGN boutons across the dendritic tree revealed monosynaptic connections restricted to apical oblique branches (Fig. 1b,c).

Given the specific targeting of dLGN inputs to L5 PN oblique dendrites, we asked whether these branches exhibited distinct properties compared to nearby basal dendrites (Fig. 1d). In acute slices from adult mice, basal dendrites displayed highly supralinear integration and

large local calcium influx in response to spatiotemporally-clustered two-photon glutamate

uncaging (Fig. 1e,f,g), consistent with previous reports of NMDA spikes in juvenile cortex[17]. In

contrast, oblique dendrites integrated inputs linearly until axosomatic action potential threshold

was reached and did not show significant subthreshold calcium influx (Fig. 1e,f,g). These

differences could not be explained by the size of uncaging events, the number of inputs,

distance from the soma, action potential threshold, or $\delta V/\delta t$ prior to action potential

(Supplemental Fig. 1). Thus, in adult V1 L5 PNs, oblique dendrites selectively receive dLGN

input and exhibit a specialized linear integration mode.

Oblique dendrites lack synaptic potentiation in adult V1 L5 PNs

76   Because oblique dendrites did not exhibit large local calcium signals, we hypothesized

that synapses on these branches should potentiate less than those on basal dendrites.

Therefore, we tested synaptic potentiation within dendritic compartments by pairing focal

electrical stimulation of presynaptic axons at basal or oblique dendrites with somatic current

injection. A theta glass bipolar stimulating electrode was positioned within 10 μm of either a

basal or an oblique dendritic branch (Fig. 2a,b, Supplemental Fig. 2.) to generate a 1 to 2 mV

excitatory postsynaptic potential (EPSP) in the presence of a GABA-A antagonist. EPSPs were

paired with 20 ms somatic current injections that drove bursts of 2-4 action potentials. The peak

of the EPSP and the peak of the first action potential were within 10 ms of each other, and 5

pairings at 10 Hz were repeated 30 times, with 10 seconds between epochs (Fig. 2c). This

protocol strongly potentiated synapses on basal dendrites (Fig. 2d,f, Supplemental Fig. 2),

similar to findings in juvenile cortex[21]. Potentiation at basal branches required pre- and

postsynaptic coincidence and NMDARs (Supplemental Fig. 2). In contrast, synapses on oblique

dendrites did not show any significant potentiation after pairing (Fig. 2e,f, Supplemental Fig. 2).

Increasing the numbers of pairings did not change this outcome (Supplemental Fig. 2). These

findings indicate that, unlike basal dendrites, oblique branches of L5 PNs in adult V1 lack

Hebbian synaptic potentiation.

Synapses at oblique dendrites are plastic during a postnatal critical period

95   In V1, dLGN inputs are refined during early postnatal development[33–35], after which

synaptic potentiation at some cortical synapses show a developmental decline[36–38]. Therefore,

we hypothesized that synapses on oblique L5 PN dendrites may exhibit NMDAR-mediated

supralinear integration and plasticity during early postnatal development. Indeed, we found that

at eye-opening (postnatal days 12-14), oblique and basal dendrites displayed similar NMDAR-

dependent supralinearities and large local calcium signals (Fig. 3a-c, Supplemental Fig. 3).

These properties persisted into the canonical critical period (P18-22)[39,40]. By 4 weeks of age

(P28-P32), oblique dendrites had developed adult-like properties, with linear synaptic integration

and small local calcium signals (Fig 3c, Supplemental Fig. 3). Synaptic plasticity at oblique

dendrites followed the same developmental trajectory: except for P28, when integrative

properties were adult-like, oblique and basal dendrites exhibited NMDAR-dependent synaptic

potentiation at juvenile ages (Fig. 3d, Supplemental Fig. 3). These findings suggested that

visual experience mediates the maturation of oblique dendrite properties; to test this, we

conducted these same experiments in mice deprived of light from birth. In P28 dark-reared

animals, oblique dendrites retained immature-like supralinear integration with large local calcium

signals and synaptic potentiation (Fig. 3e, Supplemental Figs. 3). Dark-reared mice re-exposed

to a normal light cycle did not develop adult-like integrative properties (Supplemental Fig. 3),

indicating that the oblique compartment developed mature properties only during a specific

window in development. Collectively, these results demonstrate a restricted developmental

window in which visually-driven thalamic activity shifts the L5 PN oblique dendritic compartment

towards a linear integration mode with synapses that do not potentiate.

Oblique dendrites develop distinct postsynaptic receptor composition

One possible interpretation of our data is that adult oblique dendrites lack large local

calcium signals, integrate linearly, and do not potentiate due to a higher AMPA/NMDA ratio, as

adult cortical AMPA receptors (AMPARs) are significantly less Ca^{2+}-permeable than

NMDARs[41,42]. To test this, we compared AMPAR and NMDAR-mediated responses at single

spines from basal and oblique dendrites at P21 and in adults. Glutamate uncaging produced

AMPAR-mediated EPSPs within the physiological range at single spines, below the threshold

for calcium influx (Fig. 4a, Supplemental Fig. 4). Then, Mg^{2+}-free ACSF containing the AMPAR

antagonist DNQX was washed in and the uncaging protocol was repeated to assess synaptic

NMDARs (Fig. 4b). The average maximum amplitudes of EPSPs under each condition were

compared as an AMPA/NMDA ratio. At postnatal day 21, when oblique dendrites behave

similarly to basal dendrites, spines from the two branch types had comparable AMPA/NMDA

ratios. By adulthood, this ratio increased for oblique dendrites (Fig. 4c), indicating relatively less

NMDAR-mediated synaptic conductance, consistent with linear integration, smaller local Ca^{2+}

signals, and a decreased capacity for synaptic potentiation.

To provide complementary proteomic evidence for our observed changes in functional

AMPA/NMDA, we acquired super-resolution images of oblique and basal dendrites and their

synaptic receptor content using epitope-preserving Magnified Analysis of the Proteome (eMAP,

Fig. 4d)[43,53]. Fixed slices from V1 of adult Thy1-GFP-M mice were expanded to 4x size and

stained with antibodies against the presynaptic marker bassoon, the obligate NMDAR subunit

GluN1, and GluA1 and GluA2, the predominant AMPAR subunits in cortex[44]. Oblique and basal

dendrites from L5 PNs were imaged with confocal microscope in ~18 μm-long segments (~75

μm expanded), and dendritic protrusions were annotated (Fig. 4e,f). Immature spines and

protrusions without bassoon were excluded post hoc (although basal and oblique branches had

similar numbers of immature and mature spines; Supplemental Fig. 4). For each spine head, the

intensity of each fluorophore was spatially integrated (Fig. 4g), and the ratio of

GluA1+GluA2/NR1 was taken. Oblique dendrite synapses exhibited higher AMPA/NMDA ratios

than basal dendrites (Fig. 4h, Supplemental Fig. 4). This effect was pronounced in large spines

(Fig. 4i). Taken together, these results indicate that oblique dendrites have relatively less

NMDAR content and conductance, and this underlies the differences in integration and plasticity

rules between basal and oblique dendritic compartments.

Discussion

Our results reveal that apical and basal dendritic compartments of adult V1 L5 PNs

exhibit different integration modes and plasticity thresholds which are regulated by visual

experience over the course of development. While basal dendrites consistently express

NMDAR-dependent supralinear integration and synaptic potentiation, apical oblique dendrites

show a developmentally-regulated transition away from NMDAR-mediated processes. Oblique

dendrites of L5 PNs receive direct dLGN input, and during early visual experience, the synaptic

AMPA/NMDA ratio on these branches increases, linearizing integration and limiting further

synaptic potentiation by adulthood. Dark rearing disrupted this process, indicating a critical

period for the maturation of synaptic properties in oblique dendrites. While critical periods have

been previously described for cortical layers and inputs[40,45], our results are the first to show that

critical periods also occur subcellularly, within dendritic compartments.

Our findings reveal that NMDAR-dependent synaptic potentiation is not universal at

spiny synapses in adult cortex, an important discovery for both biological and artificial models of

learning. The full dynamic range of plasticity in adult oblique dendrites is unclear: while our

findings cannot rule out all possible plasticity mechanisms, limited NMDAR-mediated calcium

influx constrains their potential to utilize conventional molecular cascades. Increasing AMPARs

is likely one of many mechanisms involved in regulating developing synapses. Changes in

excitatory receptor subtypes[29,41,42,46–49], kinase and phosphatase activity, as well as presynaptic

properties may also be contributing factors[28]. Our results demonstrate a naturally-occurring
example of plasticity regulation and establish a new tractable model for elucidating the
mechanisms of synapse-specific stability, an important question for both experimental and
computational neuroscience.
In contrast to what we observe in oblique dendrites, basal dendrites exhibit NMDAR-
dependent supralinear integration and synaptic potentiation throughout maturation. Basal
dendrites of L5 PNs receive intracortical input[16,20,22,50,51], and effective cortical computation may
require more flexibility and nonlinear processing of intracortical information than for feedforward
thalamic input. Following developmental refinement, thalamocortical inputs become remarkably
stable[7,39,52], while cortical representations remain plastic[8–11]. Given that PNs in all layers of
primary sensory cortex can receive both first-order thalamic and intracortical inputs[12–15], single
neurons might utilize diverse plasticity thresholds within their dendrites to maintain synapse-
specific plasticity without compromising representational fidelity. Subcellular input-specific
plasticity regulation is a powerful new mechanism by which single cortical neurons can solve the
tradeoff between flexibility and stability inherent to all learning systems.
183

Author Contributions
C.E.Y. and M.T.H. conceived of the experiments. C.E.Y. carried out all physiology experiments
and analyses. D.V. conducted the eMAP experiments and C.E.Y. analyzed the data. N.J.B.
performed viral injections and histology. M.T.H. supervised all aspects of the project and wrote
the manuscript with C.E.Y.

Acknowledgements
We thank Mark Bear, Elly Nedivi, Josh Trachtenberg, and the Harnett laboratory for their
comments and suggestions and Kwanghun Chung for generously sharing equipment. This work
was funded by a Life Sciences Research Foundation postdoctoral fellowship (C.E.Y.), and a
Vallee Foundation Scholars Award, the James W. and Patricia T. Poitras Fund, and the National
Institutes of Health R01NS106031 (M.T.H.).

Competing interests
Authors declare that they have no competing interests.

Methods
Animals

All animal procedures were carried out in accordance with NIH and Massachusetts Institute of

Technology Committee on Animal Care guidelines. C57BL6 male and female mice (Charles

River Laboratories) were used in approximately equal numbers. Mice 8 weeks and older (P56+)

were used for adult mouse experiments. For developmental timepoints, mice ages P10-P14,

P18-22, and P28-P32 were used. All animals were kept in conventional social housing with

unlimited food and water on a 12-hour light/dark cycle. For dark-rearing experiments, pregnant

dams were moved to a dark room at E18 and pups were born and raised in complete darkness,

with red light exposure during cage changes. Dark reared mice were ages P28-P32 at the time

of experimentation. 8-week-old Thy-1-GFP-M mice (Jackson Laboratory) were used for eMAP

experiments.

Slice preparation

Sucrose-containing artificial cerebrospinal fluid (sACSF) was used during the slicing procedure,

containing (in mM): 90 sucrose, 60 sodium chloride, 26.5 sodium bicarbonate, 2.75 potassium

chloride, 1.25 sodium phosphate, 9 glucose, 1.1 calcium chloride, and 4.1 magnesium chloride,

with an osmolality of 295-302. The sucrose solution was partially frozen to create a small

amount of slush and was kept ice-cold. Artificial cerebrospinal fluid (ACSF) was used for

recovery and recording, containing (in mM): 122 sodium chloride, 25 sodium bicarbonate, 3

potassium chloride, 1.25 sodium phosphate, 12 glucose, 1.2 calcium chloride, 1.2 magnesium

chloride, 1 ascorbate, and 3 sodium pyruvate, with an osmolality of 302-307. All solutions were

saturated with carbogen, 5% CO_2 and 95% O_2. Acute slice preparation was consistent for all

ages and experiments. Mice were put under isoflurane-induced anesthesia and decapitated.

The brain was extracted in ice-cold sucrose solution in less than one minute. The brain was

blocked at a moderate angle (approximately 20 degrees from coronal) to favor the preservation

of apical dendrites in the occipital cortex. After the brain was mounted and submerged in ice-

cold sACSF, a vibratome (Lieca VT1200S) was used to cut 300 µm-thick slices, which were

transferred to ACSF for 30-50 minutes at 36°C. Longer recovery times were used for tissue

collected from younger animals. Following recovery, slices were kept at room temperature.

Patch-clamp recording

Recordings were performed in ACSF (concentrations noted above) at 34–36 °C. Intracellular

recording solution contained (in mM): 134 potassium gluconate, 6 KCl, 10 HEPES, 4 NaCl, 4

Mg_2ATP, 3 NaGTP, and 14 phosphocreatine di(tris). Depending on the experiment, 0.05 mM

Alexa 594, 0.1 mM Alexa 488, and/or 0.1 mM OGB-1 (Invitrogen) were added to the internal

solution. Whole-cell current-clamp recordings were obtained with a Dagan BVC-700A amplifier.

Patch pipettes with thin-wall glass (1.5/1.0mm OD/ID, WPI) and resistances of 3-7 MΩ were

used. Pipette capacitance was fully neutralized prior to break-in, and series resistance was kept

fully balanced throughout, ranging from 5-30 MΩ. The liquid junction potential was not

corrected. Current signals were digitized at 20kHz and filtered at 10kHz (Prairie View). L5 PNs

were characterized by their large somas in layer 5, thick apical dendrites, broad arborization in

layer 1, low input resistance, and prominent voltage sag.

Virus injection

For sCRACM experiments, AAV2-hSyn-hChR2(H134R)-mCherry (UNC Vector Core) was

expressed in neurons in the dorsal lateral geniculate nucleus. Mice 6 weeks or older were

injected 4 weeks or more before slice preparation. Under isoflurane anesthesia, mice were

secured on a stereotaxic apparatus with a feedback-controlled heating pad (DC Temperature

Control System, FHC). Slow-release buprenorphine (1 mg/kg) was administered

subcutaneously. After scalp incision, a small burr hole was drilled over the injection site at the

following coordinates relative to bregma: anterior-posterior 2.7-2.8 mm; medio-lateral 2.3-2.4

mm. A beveled microinjection pipette containing was lowered to a depth of 2.7-2.9 mm, and

approximately 100 nL of virus was injected at a rate of 50 nL/min using a Nanoject. Following

virus injection and a five minute rest, the pipette was removed and the incision was sutured.

Accuracy of the injection was assessed in acute slices using two-photon microscopy. For image

clarity, histology in Figure 1 was taken from a brain that was perfused, stored at 4°C overnight in

4% paraformaldehyde, transferred to PBS, and sectioned in 100 μm-thick slices. Sections were

mounted, coverslipped, and imaged under a confocal microscope (Zeiss LSM 710 with a 10x

objective, NA 0.45).

sCRACM

After 4 weeks or more of ChR2 expression in dLGN, acute slices of V1 were prepared. A two-

photon laser scanning system (Bruker), with dual galvanometers and a MaiTai DeepSee laser,

was used to confirm the presence of mCherry-expressing cell bodies in the dLGN. For all

recordings, TTX (1 μM) and 4-AP (100 μM) were added to ACSF to isolate monosynaptic

connections. Following whole-cell configuration, a 5 ms 473 nm full-field LED was used to

determine if the neuron received inputs from axons containing ChR2. If the LED drove EPSPs

or spikes, a 9 by 17 (400 x 800 μm) stimulation grid with 50 μm spacing was positioned over the

area containing the entire neuron, aligned at the pia using two-photon imaging under low

magnification (4x, 0.16 NA air objective, Olympus) and Prairie View software. The stimulation
grid controlled the location of a 473 nm laser beam (OptoEngine, LLC), and the duration of the
light pulses was kept to 1-3 ms under <2 mW power. Each point on the grid was stimulated in
order, progressing forward by column, with an interpoint delay of 1 s. 3-6 rounds of stimulation
were averaged for each neuron. Baseline voltage was defined as the 50 ms before stimulation,
and EPSP amplitudes were calculated as the baseline-subtracted maximum voltage within 50
ms after photostimulation. Population data was obtained by rotating the maps to align to the pia,
overlaying the maps to align at the soma, normalizing peak voltages, and averaging across
experiments.
276
Glutamate uncaging
Following whole-cell dialysis with a structural dye via the patch pipette, basal or apical oblique
branches were localized using two-photon imaging. A pipette containing caged glutamate (4-
methoxy-7-nitroindolinyl-caged-L-glutamate or MNI-glutamate) diluted in ACSF (10 mM) was
positioned just above the slice and over the recording site, and a picrospritzer (Picospritzer III,
Parker Hannifin) was used to puff a constant flow of caged glutamate into the region of interest.
A second Mai-Tai DeepSee laser controlled by Prairie View software was used for
photostimulation of MNI-glutamate at 720 nm. Uncaging was targeted just adjacent to spine
heads. In the imaging pathway, a linescan was used for simultaneous calcium imaging. Laser
intensity for both lasers was independently controlled (Conoptics). Linescan imaging was
performed at 1300 Hz, with a dwell time of 8 µs and a total scan time of less than 250 ms, with
baseline fluorescence kept minimal and monitored throughout. Cells with signs of photodamage
were excluded. The uncaging laser was calibrated to either a threshold calcium signal (~100%
ΔF/F) or an action potential, where the minimum power needed to drive either event was
identified and used for all points. The uncaging dwell time was 0.2 ms, and the uncaging interval
between multiple spines was 0.32 ms. Groups of 2-5 spines were stimulated independently and
then in combination with other groups up to a total of 25-40 spines. Expected values were
calculated by summing the average response of each group of spines. Ca^{2+} signals are
expressed as $(F - F_{baseline})/F_{baseline})$. For acute blockade of NMDA receptors, the competitive
antagonist D-AP5 (50 µM) was used.
297
Spike-timing dependent plasticity
For local branch stimulation, theta glass (2.0/1.4mm OD/ID) housing a bipolar stimulating
electrode (ISO-Flex, AMPI) was positioned within 10 µm of a basal or apical oblique branch.

Stimulation intensity was calibrated to generate a small EPSP of 1-3 mV, typically requiring 5-20

µA of current from the stimulating electrode. Ten minutes of baseline stimulation at 0.1 Hz was

recorded to ensure stability of the EPSP. A -50 pA hyperpolarizing step was included to

estimate input resistance. For plasticity induction, EPSPs were paired with a somatic current

injection of 400-700 pA for 20 ms duration to produce a train of 2-4 action potentials. The peak

of the first action potential was timed to be within 10 ms of the peak of the EPSP. Pairs were

done in sets of 5 at 10 Hz, and sets of 5 were repeated 30 times at 0.1 Hz. After the induction

period, EPSPs were monitored for at least 20 minutes. Inclusion criteria were as follows: resting

membrane potential could fluctuate from baseline by no more than 3 mV, input resistance could

not increase more than 20% of baseline; and series resistance had to be compensated fully

throughout (<30 MOhms). Recordings were tested for baseline stability post-hoc using either

correlation or linear regression (data not shown), and cells with significant changes in baseline

recording were not included. The necessity of pre-post pairing was shown using the same

recording set up but driving either the synaptic stimulation or the post-synaptic burst (and not

both) during the induction period. For acute blockade of NMDARs, the competitive antagonist D-

AP5 (50 µM) was used. Throughout plasticity experiments, picrotoxin (100 µM) was present in

the bath to block GABAergic transmission in acute slices from adult mice and at P28. 10 µM

picrotoxin was used for acute slices at P21, and no picrotoxin was needed at P14.

Epitope-preserving Magnified Analysis of the Proteome

Mice were perfused with cold PBS followed by cold 4% PFA while under deep anesthesia (5%

isoflurane). Brains were removed and kept in the same fixative overnight at 4°C and then

washed with PBS at 4°C for at least 1 day. 1.0 mm coronal slices of primary visual cortex were

cut on a vibratome and kept in PBS at 4°C until the day of processing. Slices were then

incubated in eMAP hydrogel monomer solution[53] (30% acrylamide [A9099, MilliporeSigma, St.

Louis, MO, USA], 10% sodium acrylate [408220, MilliporeSigma], 0.1% bis-acrylamide [161-

0142, Bio-Rad Laboratories, Hercules, CA, USA], and 0.03% VA-044 (w/v) [Wako Chemicals,

Richmond, VA, USA] in PBS), protected from light, at 4°C overnight. For gelation, slices were

mounted between glass slides in eMAP solution and sealed in a 50 mL conical tube with

nitrogen gas at positive pressure of 10-12 psi at 37°C for 3 hours. The excess gel around the

slices was then removed. To reach a first expansion stage of 1.7x, the slices were incubated

overnight in a solution of 0.02% sodium azide (w/v) in PBS at 37°C. Slices were trimmed to

contain only parts of primary visual cortex and further sectioned with a vibratome to 75 µm

thickness (corresponding to ~40 µm thickness of the pre-expanded tissue). Slices containing

good candidate cells – layer 5 pyramidal neurons whose apical trunk could be reconstructed at

its full length in a single slice or at most two consecutive slices – were selected during live low-

resolution confocal imaging sessions. These slices were trimmed to smallest possible samples

of approximately 1.0 mm in both width and length. Slices were incubated in tissue clearing

solution (6% SDS (w/v), 0.1 M phosphate buffer, 50 mM sodium sulfite, 0.02% sodium azide

(w/v), pH 7.4) at 37°C for 6 hours, followed by incubation in preheated clearing solution at 95°C

for 10 min. Cleared samples were thoroughly washed with PBS + 0.1% Triton X at 37 °C.

Primary antibody staining was performed at 37°C overnight with the following antibodies: Anti-

GFP (Life Technologies A10262), Anti-NMDAR1 (SYSY 114011), Anti-AMPAR1 (SYSY

182003), Anti-AMPAR2 (SYSY 182103), and Anti-Bassoon (SYSY 141004). For secondary

staining, the following fluorescent antibodies were used: Bassoon: anti-Guinea pig-405 (AbCam

ab175678); GFP: anti-Chicken-488 (Invitrogen A11039); NMDAR1: anti-Mouse-AF+555

(Invitrogen A32727); and AMPAR1 and AMPAR2: anti-Rabbit-AF+647 (Invitrogen A32733).

Final expansion was performed just before imaging by putting the trimmed slices in 0.1 mM tris

in distilled water, and approximately 4X total linear expansion was achieved. Slices were

imaged using Leica TCS SP8 upright confocal DM6000 microscope equipped with a 63x HC PL

APO CS2/1.2 W objective, hybrid detectors, and a white light laser. Within slices, intact L5b

neurons were identified by their thick trunks and broad apical tufts. Within single neurons, both

basal and oblique dendritic branches were imaged. To avoid photobleaching effects, no slice

was imaged more than once, and basal and oblique dendrites were imaged in alternating order

for each cell. Dendritic protrusions were analyzed using Fiji software. To draw regions of interest

within spine heads in the GFP channel, a custom-written macro code was used to apply a

median blur (2 pixels), threshold the image, and draw an ROI on the annotated protrusion.

Using custom-written code in MATLAB, all color channels were thresholded to include only

signals 2 S.D. above the mean fluorescence intensity, and signals from each channel were

extracted from the ROI. The plane with peak antibody fluorescence was identified within the

spine head ROI, and each channel's fluorescence signals within the plane were summed.

Long, thin dendritic protrusions without enlarged heads were classified as filopodia (head

diameter/neck diameter < 1.3) and excluded from the final analysis, as were spines with no

detectable bassoon.

Data availability

All data and custom-written analysis code are available upon request.

References

1. Tan, L., Tring, E., Ringach, D. L., Zipursky, S. L. & Trachtenberg, J. T. Vision Changes the Cellular Composition of Binocular Circuitry during the Critical Period. *Neuron* **108**, 735-747.e6 (2020).
2. Wang, B.-S., Sarnaik, R. & Cang, J. Critical period plasticity matches binocular orientation preference in the visual cortex. *Neuron* **65**, 246–256 (2010).
3. LeVay, S., Wiesel, T. N. & Hubel, D. H. The development of ocular dominance columns in normal and visually deprived monkeys. *J. Comp. Neurol.* **191**, 1–51 (1980).
4. Wiesel, T. N. & Hubel, D. H. Effects of visual deprivation on morphology and physiology of cells in the cats lateral geniculate body. *J. Neurophysiol.* **26**, 978–993 (1963).
5. Wiesel, T. N. & Hubel, D. H. Single-cell responses in striate cortex of kittens deprived of vision in one eye. *J. Neurophysiol.* **26**, 1003–1017 (1963).
6. Clopath, C., Bonhoeffer, T., Hübener, M. & Rose, T. Variance and invariance of neuronal long-term representations. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **372**, (2017).
7. Rose, T., Jaepel, J., Hübener, M. & Bonhoeffer, T. Cell-specific restoration of stimulus preference after monocular deprivation in the visual cortex. *Science* **352**, 1319–1322 (2016).
8. Poort, J. *et al.* Learning Enhances Sensory and Multiple Non-sensory Representations in Primary Visual Cortex. *Neuron* **86**, 1478–1490 (2015).
9. Weskelblatt, J. B. & Niell, C. M. Distinct functional classes of excitatory neurons in mouse V1 are differentially modulated by learning and task engagement. *BioRxiv* (2019) doi:10.1101/533463.
10. Makino, H. & Komiyama, T. Learning enhances the relative impact of top-down processing in the visual cortex. *Nat. Neurosci.* **18**, 1116–1122 (2015).
11. Pakan, J. M. P., Currie, S. P., Fischer, L. & Rochefort, N. L. The impact of visual cues, reward, and motor feedback on the representation of behaviorally relevant spatial locations in primary visual cortex. *Cell Rep.* **24**, 2521–2528 (2018).
12. Harris, K. D. & Shepherd, G. M. G. The neocortical circuit: themes and variations. *Nat. Neurosci.* **18**, 170–181 (2015).
13. Antonini, A., Fagiolini, M. & Stryker, M. P. Anatomical correlates of functional plasticity in mouse visual cortex. *J. Neurosci.* **19**, 4388–4406 (1999).
14. Morgenstern, N. A., Bourg, J. & Petreanu, L. Multilaminar networks of cortical neurons integrate common inputs from sensory thalamus. *Nat. Neurosci.* **19**, 1034–1040 (2016).
15. Cruz-Martín, A. *et al.* A dedicated circuit links direction-selective retinal ganglion cells to the primary visual cortex. *Nature* **507**, 358–361 (2014).
16. Young, H., Belbut, B., Baeta, M. & Petreanu, L. Laminar-specific cortico-cortical loops in mouse visual cortex. *eLife* **10**, (2021).
17. Schiller, J., Major, G., Koester, H. J. & Schiller, Y. NMDA spikes in basal dendrites of cortical pyramidal neurons. *Nature* **404**, 285–289 (2000).
18. Malenka, R. C. & Bear, M. F. LTP and LTD: an embarrassment of riches. *Neuron* **44**, 5–21 (2004).
19. Petreanu, L., Mao, T., Sternson, S. M. & Svoboda, K. The subcellular organization of neocortical excitatory connections. *Nature* **457**, 1142–1145 (2009).
20. Gökçe, O., Bonhoeffer, T. & Scheuss, V. Clusters of synaptic inputs on dendrites of layer 5 pyramidal cells in mouse visual cortex. *eLife* **5**, (2016).
21. Gordon, U., Polsky, A. & Schiller, J. Plasticity compartments in basal dendrites of neocortical pyramidal neurons. *J. Neurosci.* **26**, 12717–12726 (2006).
22. Sjöström, P. J. & Häusser, M. A cooperative switch determines the sign of synaptic plasticity in distal dendrites of neocortical pyramidal neurons. *Neuron* **51**, 227–238 (2006).

23. Golding, N. L., Staff, N. P. & Spruston, N. Dendritic spikes as a mechanism for
cooperative long-term potentiation. *Nature* **418**, 326–331 (2002).
24. Kumar, A., Barkai, E. & Schiller, J. Plasticity of olfactory bulb inputs mediated by
dendritic NMDA-spikes in rodent piriform cortex. *eLife* **10**, (2021).
25. Weber, J. P. *et al.* Location-dependent synaptic plasticity rules by dendritic spine
cooperativity. *Nat. Commun.* **7**, 11380 (2016).
26. Froemke, R. C., Poo, M.-M. & Dan, Y. Spike-timing-dependent synaptic plasticity
depends on dendritic location. *Nature* **434**, 221–225 (2005).
27. Letzkus, J. J., Kampa, B. M. & Stuart, G. J. Learning rules for spike timing-dependent
plasticity depend on dendritic synapse location. *J. Neurosci.* **26**, 10420–10429 (2006).
28. Abraham, W. C. Metaplasticity: tuning synapses and networks for plasticity. *Nat. Rev.*
Neurosci. **9**, 387 (2008).
29. Turrigiano, G. G. The self-tuning neuron: synaptic scaling of excitatory synapses. *Cell*
135, 422–435 (2008).
30. Meyer, D., Bonhoeffer, T. & Scheuss, V. Balance and stability of synaptic structures
during synaptic plasticity. *Neuron* **82**, 430–443 (2014).
31. Constantinople, C. M. & Bruno, R. M. Deep cortical layers are activated directly by
thalamus. *Science* **340**, 1591–1594 (2013).
32. Peters, A. & Feldman, M. L. The projection of the lateral geniculate nucleus to area 17 of
the rat cerebral cortex. IV. Terminations upon spiny dendrites. *J. Neurocytol.* **6**, 669–689
(1977).
33. Gu, Y. & Cang, J. Binocular matching of thalamocortical and intracortical circuits in the
mouse visual cortex. *eLife* **5**, (2016).
34. Sommeijer, J.-P. *et al.* Thalamic inhibition regulates critical-period plasticity in visual
cortex and thalamus. *Nat. Neurosci.* **20**, 1715–1721 (2017).
35. Cang, J. *et al.* Development of precise maps in visual cortex requires patterned
spontaneous activity in the retina. *Neuron* **48**, 797–809 (2005).
36. Kirkwood, A., Lee, H. K. & Bear, M. F. Co-regulation of long-term potentiation and
experience-dependent synaptic plasticity in visual cortex by age and experience. *Nature*
375, 328–331 (1995).
37. Jiang, B., Treviño, M. & Kirkwood, A. Sequential development of long-term potentiation
and depression in different layers of the mouse visual cortex. *J. Neurosci.* **27**, 9648–9652
(2007).
38. Crair, M. C. & Malenka, R. C. A critical period for long-term potentiation at
thalamocortical synapses. *Nature* **375**, 325–328 (1995).
39. Gordon, J. A. & Stryker, M. P. Experience-dependent plasticity of binocular responses in
the primary visual cortex of the mouse. *J. Neurosci.* **16**, 3274–3286 (1996).
40. Feldman, D. E. Synaptic mechanisms for plasticity in neocortex. *Annu. Rev. Neurosci.*
32, 33–55 (2009).
41. Kumar, S. S., Bacci, A., Kharazia, V. & Huguenard, J. R. A developmental switch of
AMPA receptor subunits in neocortical pyramidal neurons. *J. Neurosci.* **22**, 3005–3015
(2002).
42. Hadzic, M., Jack, A. & Wahle, P. Ionotropic glutamate receptors: Which ones, when, and
where in the mammalian neocortex. *J. Comp. Neurol.* **525**, 976–1033 (2017).
43. Ku, T. *et al.* Multiplexed and scalable super-resolution imaging of three-dimensional
protein localization in size-adjustable tissues. *Nat. Biotechnol.* **34**, 973–981 (2016).
44. Cheng, D. *et al.* Relative and absolute quantification of postsynaptic density proteome
isolated from rat forebrain and cerebellum. *Mol. Cell. Proteomics* **5**, 1158–1170 (2006).
45. Daw, N.W. Mechanisms of plasticity in the visual cortex. in *Visual Development* 207–233
(Kluwer Academic Publishers, 2006). doi:10.1007/0-387-30484-3_11.

46. Philpot, B. D., Cho, K. K. A. & Bear, M. F. Obligatory role of NR2A for metaplasticity in
visual cortex. *Neuron* **53**, 495–502 (2007).
47. Erisir, A. & Harris, J. L. Decline of the critical period of visual plasticity is concurrent with
the reduction of NR2B subunit of the synaptic NMDA receptor in layer 4. *J. Neurosci.* **23**,
5208–5218 (2003).
48. Liu, S. Q. & Cull-Candy, S. G. Synaptic activity at calcium-permeable AMPA receptors
induces a switch in receptor subtype. *Nature* **405**, 454–458 (2000).
49. Clem, R. L., Celikel, T. & Barth, A. L. Ongoing in vivo experience triggers synaptic
metaplasticity in the neocortex. *Science* **319**, 101–104 (2008).
50. Markram, H., Lübke, J., Frotscher, M., Roth, A. & Sakmann, B. Physiology and anatomy
of synaptic connections between thick tufted pyramidal neurones in the developing rat
neocortex. *J Physiol (Lond)* **500 (Pt 2)**, 409–440 (1997).
51. Thomson, A. M. & Bannister, A. P. Postsynaptic pyramidal target selection by
descending layer III pyramidal axons: dual intracellular recordings and biocytin filling in
slices of rat neocortex. *Neuroscience* **84**, 669–683 (1998).
52. Jaepel, J., Hübener, M., Bonhoeffer, T. & Rose, T. Lateral geniculate neurons projecting
to primary visual cortex show ocular dominance plasticity in adult mice. *Nat. Neurosci.*
20, 1708–1714 (2017).
53. Park, J., Khan, S., Yun, D.H., Ku, T., Villa, K.L., Lee, J.E., Zhang, Q., Park, J., Feng, G.,
Nedivi, E., et al. Epitope-preserving magnified analysis of proteome (eMAP). *Sci. Adv.* **7**,
eabf6589. (2021).
489

Figure 1: Apical oblique dendrites in V1 L5 PNs of adult mice are selectively targeted by dLGN inputs and exhibit distinct integrative properties compared to basal dendrites.

a. Example confocal image of a coronal brain slice showing dLGN neurons infected with AAV2-hSyn-mCherry-ChR2.

b. Left: Example two-photon image of a V1 L5 PN whole-cell recording with overlaid photostimulation grid (blue). Scale bar, 50 µm. Middle: Average evoked EPSPs from 6 repetitions of 473 nm photostimulation for the neuron and grid shown. Soma position is indicated by the triangle. Right: Heatmap of average evoked EPSPs. Each square is 50 µm.

c. Left: Normalized heatmap of pooled L5 PNs (n = 9 neurons, 9 slices, 3 P56+ animals), aligned at the soma. Each square is 50 µm. Right: Mean normalized input as a function of depth for all neurons. Shaded region indicates the bootstrapped 95% confidence interval.

d. Example two-photon z-stack of a L5 PN with glutamate uncaging sites indicated on an oblique (purple) and a basal (green) branch. Uncaging sites (stars) and linescan (yellow) are magnified in insets (scale bars: 10 µm).

e. Expected EPSPs, measured EPSPs, and local branch Ca^{2+} signals for oblique (top) and basal (bottom) dendrites shown in d. Left: Expected voltages from linearly summed inputs. Middle: Measured EPSPs in response to increasing numbers of synchronous uncaging inputs. Stimulation of all inputs led to an action potential (AP, arrow). Right: Corresponding change in OGB-1 signal (ΔF/F) with input summation.

f. Measured voltage (left) and branch ΔF/F (right) as a function of expected voltage for the branches shown in d. Dashed line indicates linearity (left) and threshold Ca^{2+} signal (right).

g. Adult population gain (measured/expected, left) and branch ΔF/F (right). n = 13 basal branches from 7 animals and 9 oblique branches from 6 animals. All animals were P56+. Basal AP removed for clarity. Basal vs oblique gain (AP excluded), Mann-Whitney U test: p = 7.63E-04; basal vs oblique ΔF/F (AP excluded), Mann-Whitney U test: p = 1.08E-05.

Figure 2: Oblique dendrites in V1 L5 PNs of adult mice lack long-term potentiation, unlike basal dendrites.

a. Example two-photon z-stack of whole-cell recording with focal stimulation (green) near a basal dendrite.
b. As in a, but for an oblique dendrite (purple).
c. Plasticity induction protocol. 5 pairings of pre- and post-synaptic stimulation at 100 Hz, repeated 30 times at 0.1 Hz. Example trace below, right; inset shows the 10 ms interval between the peak of the evoked EPSP and the peak of the first action potential. Scale bar: 20 mV, 50 ms; inset: 5 mV, 5 ms.
d. Example recording from the basal dendrite shown in a. After 10 minutes of baseline recording, the induction protocol is delivered (arrows denote start and stop). Inset shows average EPSP before (black) and after (color) pairing. Scale bar: 1 mV, 25 ms. Responses are 1 min binned averages.
e. As in d, for the oblique dendrite shown in b.
f. Mean normalized peak EPSP amplitude for neurons stimulated at basal (green) or oblique (purple) branches (2 min bins; n = 10 basal branches from 10 animals and 13 oblique branches from 12 animals. All animals were P56+.). Error bars are s.e.m. Basal vs. oblique post-induction EPSP, Mann-Whitney U test: p = 4.95E-63.

Figure 3: Developmental regulation of synaptic integration and plasticity in oblique versus basal dendrites.

a. Expected (left), measured (middle), and ΔF/F (right) for oblique (top) and basal (bottom) dendrites from the same neuron from a P14 mouse. Note supralinear integration and large Ca^{2+} signals in both branches. Both are driven to action potential (AP, arrow).

b. P14 population gain (measured/expected, left) and branch ΔF/F (right) (n = 9 basal branches and 9 oblique branches, both from 5 animals). Basal vs oblique gain*, p = 0.33; ΔF/F*, p = 0.70.

c. Maximum gain (left) and local branch ΔF/F (prior to AP initiation, right) across developmental time points (Adult, P56+: n as in Fig. 1, basal vs oblique gain*, p = 1.07E-04; ΔF/F*, p = 1.07E-04. P14: n as in b, basal vs oblique gain*, p = 0.54; ΔF/F*, p = 0.76. P21: n = 9 basal branches, 4 animals, 6 oblique branches, 2 animals, basal vs oblique gain*, p = 0.52; ΔF/F*, p = 0.68. P28: n = 10 basal branches, 5 animals; 9 oblique branches, 5 animals, basal vs oblique gain*, p = 2.17E-05; ΔF/F*: p = 4.33E-05.

d. Plasticity (as described in Fig. 2) in basal versus oblique dendrites at P14 (n = 5 basal
branches from 5 animals and 5 oblique branches from 5 animals, basal vs oblique post-
induction*, p = 0.79), P21 (n= 5 basal branches from 4 animals and 7 oblique branches from
6 animals, basal vs oblique post-induction*, p = 0.87), and P28 (n = 7 basal branches, 5
animals; 7 oblique branches, 4 animals, basal vs oblique post-induction*, p = 1.25E-13).
e. P28 population gain (measured/expected, left) and branch ΔF/F (middle) in dark-reared
(DR) mice (n = 9 basal branches and 10 oblique branches from 4 animals, basal vs oblique
gain*, p = 0.78; ΔF/F*, p = 0.96). Plasticity for dark-reared animals (right) (n = 5 apical
branches from 4 animals and 6 oblique branches from 6 animals, basal vs oblique post-
induction*, p = 0.81).
*Mann-Whitney U test was used for all comparisons.
561

Figure 4: Changes in synaptic AMPA-to-NMDA ratio at oblique versus basal dendrites underlie differences in integration and plasticity.

a. Example two-photon image of a V1 L5 PN from a P21 mouse. An oblique dendrite (circled) is shown at higher magnification (right). Uncaging site indicated by star.

b. Representative average EPSPs from uncaging at single spines in P21 (left) or adult (right) mice from oblique (top) and basal (bottom) dendrites. Black traces: control, colored traces: the same spine in DNQX+Mg^{2+}-free ACSF.

c. Functional single spine AMPA/NMDA ratios from basal and oblique dendrites in P21 and adult mice (P21: n = 72 basal spines from 4 mice and 89 oblique spines from 4 mice. Adult, P56+: n = 66 basal spines from 4 mice and 67 oblique spines from 5 mice). Outliers are not shown. P21 basal vs oblique, Mann-Whitney U test: p = 0.23; adult basal vs oblique, Mann-Whitney U test: p = 1.00E-04.

d. Schematic of the Magnified Analysis of the Proteome (eMAP) experimental procedure.

e. Confocal image of L5 PNs from a Thy1-GFP M P56+ mouse after 4x eMAP expansion.

f. Example confocal images of expanded oblique (top) and basal (bottom) dendrites from the same neuron. For each branch, immunostaining for bassoon, NR1, and GluA1+2 is shown for one spine (indicated by star). Scale bar: 10 µm, spine inset: 2 µm.

581   g. Probability density of NR1 and GluA1+2 signals for all spines (n = 1,305 basal dendrite
582   spines and 1,719 oblique dendrite spines from 13 cells, 3 mice).
583   h. Ratio of AMPA/NMDA intensities for all spines: Mann-Whitney U test: p = 8.30E-11.
584   i. Ratio of AMPA/NMDA intensities for the largest 10% of basal and oblique spines (n = 131
585   basal dendrite spines and 171 oblique dendrite spines). Mann-Whitney U test: p = 1.08E-04.
586

Supplemental Figure 1. Properties of adult basal and oblique dendrites and experimental parameters of glutamate uncaging experiments.

a. Example voltage waveform and corresponding δV/δt for an AP evoked by glutamate uncaging at an oblique branch. Scale bar, top: 5 mV, 10 ms, bottom: 1 mV/s.
b. Voltage thresholds for APs evoked by uncaging at basal (green) and oblique (purple) dendrites (n = 12 basal branches, 7 animals; 9 oblique branches, 6 animals. Mann-Whitney U test: p = 0.27).
c. Maximum δV/δt prior to AP initiation for uncaging at basal and oblique branches (n = 12 basal branches, 7 animals; 9 oblique branches, 6 animals. Mann-Whitney U test: p = 0.64).
d. Maximum ΔF/F for subthreshold responses at basal dendrites and maximum ΔF/F in oblique dendrites following an AP (n = 13 basal branches, 7 animals; 8 oblique branches, 6 animals. Mann-Whitney U test: p = 0.69).
e. Uncaging site distance from soma for basal and oblique dendrites (n = 13 basal branches, 7 animals, 9 oblique branches, 6 animals. Mann-Whitney U test: p = 0.34).
f. EPSP amplitude for basal and oblique dendrites driven by 5 uncaging points (n = 13 basal branches, 7 animals; 9 oblique branches, 6 animals. Mann-Whitney U test: p = 0.64).
g. Maximum number of uncaging points used in basal or oblique experiments (n = 13 basal branches, 7 animals; 9 oblique branches, 6 animals. Mann-Whitney U test: p = 1).

Supplemental Figure 2. Experimental measurement of local electrical stimulation spatial spread and properties of synaptic potentiation in adult L5 PNs.

a. Two-photon z-stack showing experimental set up. Theta glass pipette housing a bipolar electrode is positioned close to a basal branch to produce EPSPs and local Ca^{2+} signals. The branch is then severed with a laser (yellow) and EPSPs are measured again. The damaged proximal part of branch is indicated with a dotted line for clarity.

b. A local branch Ca^{2+} signal (top) and somatically-recorded EPSP (bottom) were driven in the basal branch shown in a, prior to the laser cut (pre, green). Following severing of the branch, the EPSP is greatly reduced (post, gray).

c. Percent reduction of peak EPSP amplitude following laser cutting, n = 6 basal branches from 6 neurons and 5 mice.

d. Input resistance (top) and resting membrane potential (bottom) monitored throughout adult plasticity experiments shown in Figure 2. Neither corresponded with changes in synaptic potentiation.

e. Coordinated pre- and post-synaptic activity and NMDA receptors are required for synaptic potentiation in basal dendrites. EPSPs (top) in basal dendrites following induction protocols with presynaptic-only stimulation (black, n = 6 cells, 5 animals), postsynaptic-only (gray, n = 7 cells, 4 animals), and pre/post pairing in the presence of D-AP5 (blue, n = 5 cells, 5 animals) with corresponding input resistance (middle) and resting membrane potential (bottom). Presynaptic-only or postsynaptic-only before versus after pairing comparisons were not statistically significant (pre-only, Mann-Whitney U test: p = 0.42, post-only, Mann-

628   Whitney U test: p = 0.66,); pairing with D-AP5 showed small but statistically significant
629   depression, Mann-Whitney U test: p = 0.006.

630   f. EPSPs (2 min binned averages) from a basal and oblique dendrite before and after plasticity
631   induction with 60 pairings. Despite double the number of pairings, the oblique branch does
632   not potentiate.

Supplemental Figure 3. Dendritic input-output relationships and plasticity properties for developmental timepoints and controls.

All panels are gain (top) and branch ΔF/F (bottom) vs. number of glutamate uncaging inputs.

a. Blockade of NMDARs by D-AP5 eliminates supralinear gain and branch calcium influx for both basal (blue) and oblique (grey) dendrites in P14 mice.
b. Basal and oblique dendrites exhibit supralinear gain and large Ca^{2+} signals in P21 mice under control conditions. Basal vs oblique gain*: p = 0.29; ΔF/F*: p = 0.49.
c. In P28 mice, basal dendrites exhibit supralinear integration and large Ca^{2+} signals, in contrast to oblique dendrites, which are linear and lack Ca^{2+} signals prior to AP initiation. Basal vs oblique gain*: p = 3.14E-06; ΔF/F*: p = 7.84E-04. AP excluded in both comparisons.

d. Both basal and oblique dendrites in P28 dark-reared animals integrate supralinearly with
large Ca^{2+} signals. Basal vs oblique gain*: p = 0.78; ΔF/F*: p = 0.96, Mann-Whitney U test.
These processes are NMDAR-dependent (D-AP5 blockade: basal, blue, oblique, grey).
e. Both basal and oblique dendrites in dark-reared animals that are returned to normal light
conditions for 2-4 weeks retain supralinear integration and large Ca^{2+} signals. Basal vs
oblique gain*: p = 0.75; ΔF/F*: p = 0.65.
f. Corresponding input resistance (top) and resting membrane potential (bottom) during
plasticity experiments in Fig 3, for each developmental timepoint and dark-reared animals.
g. D-AP5 prevents synaptic potentiation in P14 basal and oblique dendrites (n = 4 oblique
dendrites in 3 animals, 2 basal dendrites in 2 animals, pre vs post induction*: p = 1.41E-06).

*Mann-Whitney U test was used for all comparisons.

Supplemental Figure 4. Amplitudes of uncaging-evoked single spine EPSPs and properties of spines in basal and oblique dendrites in expanded tissue.

a. AMPAR-mediated EPSP amplitudes for all spines used to measure functional AMPA/NMDA in P21 and adult mice (Mann-Whitney U test: p = 0.43).
b. Probability density of immature spines in basal and oblique dendrites (n = 392 immature spines in basal dendrites, 513 immature spines in oblique dendrites, 13 cells, 3 animals, Mann-Whitney U test: p = 0.34).
c. Probability density of mature spines in basal and oblique dendrites (n = 1305 spines in basal dendrites, 1719 spines in oblique dendrites, 13 cells, 3 animals, Mann-Whitney U test: p = 0.77).
d. Ratio of AMPA/NMDA fluorescence intensity normalized by spine head area for spines in basal and oblique dendrites, Mann-Whitney U test: p = 2.79E-08.