**Title**: Oncogenic protein condensates suppress growth factor perception and modulate drug tolerance

**Authors:** David Gonzalez-Martinez[1#], Lee Roth[1#], Thomas R. Mumford[1], Juan Guan[2,3], Bo Huang[4,5,6], Asmin Tulpule[7], Trever G. Bivona[8], Lukasz J. Bugaj[1,9,10,*]

**Affiliations:**
[1]Department of Bioengineering, University of Pennsylvania, Philadelphia, PA, 19104, USA
[2]Department of Physics, Department of Anatomy and Cell Biology, University of Florida, Gainesville, FL 32611
[3]Department of Anatomy and Cell Biology, University of Florida, Gainesville, FL 32611
[4]Department of Pharmaceutical Chemistry, UCSF, San Francisco CA 94143, USA
[5]Department of Biochemistry and Biophysics, UCSF, San Francisco CA 94143, USA
[6]Chan Zuckerberg Biohub, San Francisco 94158, USA
[7]Division of Pediatric Hematology/Oncology, UCSF, San Francisco, CA 94143, USA
[8]Department of Medicine, Division of Hematology and Oncology, UCSF, San Francisco, CA 94143, USA
[9]Abramson Cancer Center, University of Pennsylvania, Philadelphia, PA, 19104, USA
[10]Institute of Regenerative Medicine, University of Pennsylvania, Philadelphia, PA, 19104, USA

[#]equal contributions
[*]corresponding author

**Correspondence:** bugaj@seas.upenn.edu

**Abstract**
Drug resistance remains a central challenge towards durable cancer therapy, including for cancers driven by the EML4-ALK oncogene. EML4-ALK and related fusion oncogenes form cytoplasmic protein condensates that transmit oncogenic signals through the Ras/Erk pathway. However, whether such condensates play a role in drug response is unclear. Here, we used optogenetics to find that condensates suppress signaling through endogenous RTKs including EGFR. Notably, ALK inhibition hypersensitized RTK signals, which are known to drive resistance. Suppression of RTKs occurred because condensates sequestered downstream adapter proteins that are required for RTK signal transmission. Strikingly, EGFR hypersensitization resulted in rapid and pulsatile Erk signal reactivation, which originated from neighboring apoptotic cells. Paracrine signals promoted survival during ALK inhibition, and blockade of paracrine signals suppressed drug tolerance. Our results uncover a regulatory role for RTK fusion condensates in cancer drug response and demonstrate the potential of optogenetics for uncovering functional biomarkers of cancer cells.

## Introduction

Despite the development of potent oncogene inhibitors, drug resistance remains an unsolved challenge towards durable cancer therapies. Thus, there remains a critical need to better understand molecular interactions between host cells, oncogenes, and targeted inhibitors to identify effective treatment strategies that forestall or prevent resistance.

The oncoprotein EML4-ALK (echinoderm microtubule-associated protein-like 4-anaplastic lymphoma kinase) is a receptor tyrosine kinase (RTK) fusion oncogene that drives ~3-7% of non-small cell lung cancer (NSCLC), the leading cause of cancer-related death in the US (Siegel et al., 2021; Takeuchi et al., 2012). RTK fusions are a large class of chimeric oncogenes that share a similar composition, where the enzymatic fragment of an RTK is fused through chromosomal rearrangement to a fusion partner domain that is often multimeric (Du and Lovly, 2018; Shaw et al., 2013). Cancers driven by RTK fusions commonly exhibit oncogene addiction, where blockade of the oncogene causes cell death(Weinstein, 2002). For EML4-ALK+ cancers, multiple FDA-approved ALK inhibitors achieve initial tumor regression, but resistance and tumor relapse inevitably emerge (Gainor et al., 2016; Solomon et al., 2018).

EML4-ALK drives oncogenicity primarily through sustained Ras/Erk signaling (Hrustanovic et al., 2015). Durable treatment of cancers driven by Ras/Erk has been challenged by robust autoinhibitory feedback loops that result in pathway reactivation after treatment. For example, in the case of BRAF V600E+ cancers, BRAF inhibition suppresses oncogenic Erk signals but simultaneously relieves Erk-dependent negative feedback of RTKs, resulting in strong EGFR stimulation, cell survival, and drug resistance (Corcoran et al., 2012; Gerosa et al., 2020; Lito et al., 2012; Prahallad et al., 2012). For EML4-ALK+ cancers, re-activation of RTK signals after ALK inhibition also promotes drug tolerance and acquired resistance (Hrustanovic et al., 2015; Sasaki et al., 2011; Tani et al., 2016; Vaishnavi et al., 2017; Voena et al., 2013; Wilson et al., 2012) (**Figure 1A**), although specific feedback mechanisms that mediate this reactivation are not well understood.

Recently it was found that EML4-ALK forms large cytoplasmic protein granules—or condensates— in cancer cells(Hrustanovic et al., 2015; Tulpule et al., 2021) (**Figure 1B**). Condensate formation resulted in part from the ability of RTK adapter proteins like Grb2 to enlarge small EML4-ALK oligomers through multivalent interactions. Such higher-order assemblies of proteins may have numerous and even opposing consequences on intracellular biochemistry, for example acting as amplifiers or inhibitors of molecular reactions(Banani et al., 2017). Although condensate formation serves as an amplifier that promotes EML4-ALK activation and signaling(Tulpule et al., 2021), it is unclear whether condensates might have additional impact on a cell's ability to transmit signals. Of particular interest are functional interactions between EML4-ALK condensates and transmembrane RTKs including the epidermal growth factor receptor (EGFR), which signal through the same downstream effectors as EML4-ALK (**Figure 1C**). These interactions are potentially important because EGFR is overexpressed in over 80% of NSCLC and is correlated with poor prognosis (Rusch et al., 1993) and drug resistance (Davies et al., 2013; Lito et al., 2012; Obenauf et al., 2015; Straussman et al., 2012; Tanimoto et al., 2014; Vaishnavi et al., 2017; Wilson et al., 2012).

We recently developed a method called 'functional profiling' to detect how oncogene expression can corrupt cell signal transmission (Bugaj et al., 2017, 2018). In this approach, we apply precisely defined signaling stimuli, for example using light-activated 'optogenetic' signaling probes, and we quantify differential responses in downstream signals, transcription, and cell fate as a function of oncogene expression or drug treatment. Previously, functional profiling revealed

that certain BRAF-mutant cancer cells and drug-treated cells exhibited abnormally slow Ras/Erk
pathway activation kinetics, and that such kinetics can cause cells to misinterpret dynamic stimuli
leading to hyperproliferation(Bugaj et al., 2018).
In the present work, we applied optogenetics to functionally profile the interaction between
EML4-ALK condensates and transmembrane RTK signaling (**Figure 1D**). We discovered that
EML4-ALK expression dramatically suppressed a cell's perception of RTK signals. We found
that suppression resulted not from biochemical feedback, but rather from sequestration of RTK
adapters by EML4-ALK condensates. Suppression was rapidly reversed by ALK inhibition,
resulting in hypersensitization of RTKs to external growth factors. Restored perception of
external ligands promoted rapid EGFR/Erk reactivation after drug treatment, which suppressed
cell killing and permitted tolerance to ALK inhibitors. Our work thus uncovers an important role
for oncogenic condensates in signal regulation and drug response and suggests novel candidate
co-targets to enhance the durability of ALK inhibitor therapy.
102
**Results**
***Optogenetic profiling reveals that RTK signaling is suppressed in EML4-ALK+ cancer***
***cells.***
To determine whether EML4-ALK could alter RTK signal transmission, we expressed a light-
sensitive fibroblast growth factor receptor (optoFGFR (Kim et al., 2014)) in the STE-1 cancer
cell line, which is driven by EML4-ALK(V1) (Lovly et al., 2014) (**Figure 1E, S1A**). We observed
signal transmission by applying blue light stimuli and measuring single-cell phospho-Erk (ppErk)
levels through immunofluorescence. We found that optoFGFR could induce only minimal ppErk
signal increase (1.4-fold) above the high basal ppErk levels attributed to active EML4-ALK
(**Figure 1F, top)**. We then asked whether signaling responses might change in the presence of
ALK inhibition. Pre-treatment with the ALK inhibitor crizotinib (ALKi, 1 µM) eliminated the high
tonic ppErk signal observed in untreated cells, consistent with the ability of EML4-ALK to drive
strong signals through the Ras/Erk pathway. Strikingly, subsequent light stimulation now drove
strong Erk signaling, surpassing levels achieved in the absence of drug (**Fig. 1F, bottom)**. The
dynamic range (fold-change) of signal induction increased as a function of light intensity and
reached a maximum of 29-fold increase at the highest levels of light stimulation (**Figure 1G**).
Notably, ppErk increase was measured in response to exceedingly low levels of light (8
mW/cm$^2$), a level that did not provide measurable increase in untreated cells, suggesting that
ALK inhibition can hypersensitize cancer cells to weak RTK signals (**Figure S1B**). We note that
although the magnitude of dynamic range increase was dependent on the expression levels of
optoFGFR, the general trends held across all expression levels (**Figure S1C**). These results
suggested the existence of a strong suppressive effect of EML4-ALK activity on transmembrane
RTK signaling.
126
***EML4-ALK suppresses — and ALK inhibition restores — EGFR signaling***
To determine whether suppression of RTKs impacted signaling through endogenous
epidermal growth factor receptor (EGFR), we measured ppErk induction upon addition of
epidermal growth factor (EGF) in STE-1 and H3122 cells, two EML4-ALK+ cell lines (Lovly et
al., 2014) (**Figure 2A)**. Strong (100 ng/mL) EGF stimulation gave minimal ppErk response, but
pre-treatment with ALKi substantially increased ppErk dynamic range in both cell lines (1.8- vs
11.2-fold in STE-; 1.7- vs 6.6-fold in H3122) (**Figure 2B,C**). As before, increased dynamic range
was due to lower baseline — but also higher maximal — ppErk signaling, and the magnitude of

dynamic range increase was dose-dependent (**Figure 2C, S2A**). Thus, we concluded that
EML4-ALK activity suppresses EGFR signaling in cancer cells.
We next asked whether expression of EML4-ALK was sufficient to suppress RTK
signaling. We transiently expressed EML4-ALK(V1) in non-transformed lung epithelial Beas2B
cells and we observed responses to EGF (**Figure 2D**). EML4-ALK expression raised basal ppErk
levels (t = 0, **Figure 2E**), and EGF stimulation resulted in only a small increase in ppErk relative
to untransfected cells (**Figure 2E**). Furthermore, pre-incubation with ALKi reversed suppression
of EGFR/Erk signals in transfected cells (**Figure 2E,F**), echoing our observations in cancer cells.
Collectively, our results show that EML4-ALK expression is sufficient to strongly dampen
signaling through transmembrane EGFR, and that ALK inhibition restores and potentiates these
signals.
146
***Mapping EGFR suppression using optogenetics***. We sought to understand the molecular
mechanism by which EML4-ALK suppressed EGFR signaling. To narrow candidate
mechanisms, we first determined the duration of ALKi pre-incubation that was required to
observe enhanced EGFR/Erk signaling. We preincubated both STE-1 and H3122 cells with 0-
60 min of ALKi, and we analyzed ppErk levels in response to 15 min of EGF stimulation (**Figure
3A**). In both cell lines, an increase in Erk signal amplitude was observed with as little as 5 min
of ALKi pre-incubation (20 minutes total including stimulation), and rose to half-max with only
~15 mins of pre-incubation (**Figure 3B**). Such fast response suggested a primarily post-
translational mechanism.
To pinpoint the node in the EGFR/Erk pathway that was suppressed by EML4-ALK, we
measured ppErk responses to optogenetic stimulation at successive nodes of the pathway in
the presence or absence of drug (**Figure 3C,D**). We stimulated STE-1 cell lines that stably
expressed either optoFGFR or optoSOS, which allows activation of Ras/Erk signals through
light-induced membrane recruitment of the SOS catalytic domain(Benman et al., 2022; Toettcher
et al., 2013) (**Figure 3D, S2**). As before, light stimulation of optoFGFR yielded little increase in
ppErk, but ALKi pretreatment enhanced ppErk induction (**Figure 3E, left**), consistent with relief
of optoFGFR suppression by ALKi. By contrast, the maximal level of ppErk induced by optoSOS
was unchanged by ALKi pre-treatment (**Figure 3E, right**). These results suggest that EML4-
ALK suppresses signaling upstream of Ras stimulation, for example at the receptor level.
However, further experiments showed that EGF-stimulated phospho-EGFR (pEGFR) levels
were unchanged by ALKi pre-incubation, demonstrating that EML4-ALK does not directly
suppress RTK phosphorylation (**Figure S3**). Collectively, our results indicate that EML4-ALK
suppresses RTK signaling downstream of receptor phosphorylation but upstream of Ras
activation, implicating a role for the adapter proteins that couple these two nodes (**Figure 3F**).
Oncogene-mediated feedback suppression of RTKs has been described previously, most
notably as a result of Erk-dependent transcriptional and post-translational negative
feedback(Corcoran et al., 2012; Gerosa et al., 2020; Lito et al., 2012; Prahallad et al., 2012;
Turke et al., 2012). To test the role of Erk-dependent negative feedback in EML4-ALK+ cancer
cells, we decoupled ALK inhibition from the loss of Erk signaling using optoSOS stimulation
(**Figure 3G**). In this experiment, we again assessed EGF response after ALKi preincubation, but
we supplemented one experimental group with optogenetic Ras/Erk signaling at levels that
matched those from drug-naïve STE-1 cells, thus maintaining any Erk-dependent feedback that
would have been otherwise lost through ALK inhibition (**Figure 3G,H,I, S4A**). We found that
optoSOS stimulation during ALKi pre-incubation did not diminish the enhanced response to EGF

(**Figure 3J, S4B**). In agreement, the levels of Erk-dependent negative regulator Spry2 did not
change within the first 60 min of ALKi pre-incubation (**Figure S4C**). These results show that, in
EML4-ALK+ cancer cells, EGFR suppression is not mediated by established, Erk-dependent
mechanisms.
185
***EML4-ALK condensates suppress EGFR through sequestration of RTK effectors.*** We
suspected that condensation of EML4-ALK could play a causal role in RTK suppression. To test
the effects of such higher-order EML4-ALK organization, we examined EGF response in cells
transfected with EML4-ALK mutants that fail to form condensates due to mutations in either the
trimerization domain (ΔTD) or the kinase domain (K589M)(Tulpule et al., 2021). In contrast to
full-length EML4-ALK (V1) (**Figure 4A, left**), the condensate-deficient mutants had minimal
effect on EGF-induced Erk activity (**Figure 4A, middle and right**), suggesting that condensation
was essential for its ability to modulate EGFR sensitivity.
Protein condensation can act as a negative regulator in both natural and engineered
systems when it sequesters essential components of biochemical reactions (Garabedian et al.,
2021; Omer et al., 2018; Thedieck et al., 2013). We thus hypothesized that EML4-ALK
condensates suppressed transmembrane RTKs by sequestering the adapter proteins that are
required to transmit RTK signals. Such adapters, including Grb2 and SOS1 which link RTKs to
Ras/Erk signaling, are also required to transmit EML4-ALK signals  (Tulpule et al., 2021), and
thus represent shared resources that could implement competitive inhibition of RTKs.
To test this hypothesis, we observed the localization of the Grb2 adapter using Beas2B
cells where Grb2 was fluorescently tagged at the endogenous locus (Tulpule et al., 2021). In
normal cells, Grb2 appeared diffuse in the cytoplasm but translocated to activated EGFR within
<1 min of EGF addition (**Fig. 4B,C, Supplementary Movie 1**). However, upon transfection with
EML4-ALK, Grb2 strongly colocalized with EML4-ALK puncta (**Figure S5**), and Grb2 remained
sequestered in the puncta even upon addition of EGF (**Figure 4B,C,D, Supplementary Movie
1**). We then asked how Grb2 localization might change in the presence of ALKi (**Figure 4E**).
ALKi caused Grb2 puncta to rapidly dissolve into the cytoplasm ($t_{1/2}$ = 17 +/- 2 min) (**Figure
4F,G, Supplementary Movie 2**), in line with recent reports (Sampson et al., 2021). Finally,
sequential treatment with ALKi and EGF allowed robust membrane translocation of Grb2 even
in the presence of EML4-ALK (**Figure 4H,I**). These results suggest that EML4-ALK condensates
sequester Grb2 and suppress translocation to the membrane upon EGF stimulation, providing
a mechanism by which EML4-ALK can competitively inhibit EGFR signaling.
As a further test of our model, we directly measured the EGF-induced recruitment of
adapters in H3122 cancer cells through co-immunoprecipitation.  SOS1 is recruited by Grb2 to
activated receptors and stimulates Ras upon recruitment. We found that SOS1 co-precipitated
with EGFR upon EGF stimulation, but only when preincubated with ALKi, further suggesting that
active EML4-ALK condensates sequester RTK adapters (**Figure 4J, S6**). Finally, our conceptual
model predicts that distinct RTK fusions that form condensates would similarly repress EGFR
signaling. We thus measured EGF response in TPC-1 cells, which harbor the CCDC6-RET
fusion. Like EML4-ALK, CCDC6-RET forms condensates that colocalize with many RTK
adapters, including Grb2 and SOS1(Tulpule et al., 2021). As predicted, EGF stimulation induced
only moderate levels of Erk phosphorylation, whereas pre-treatment with RET inhibitor BLU-667
permitted strong stimulation (2.0- vs 6.0-fold stimulation) to levels beyond those achievable in
drug-naive cells (**Figure S7**). Collectively, our results suggest that competition for RTK adapters

is a key mechanism by which condensates of EML4-ALK —and potentially other RTK fusions —
suppress a cell's perception of transmembrane RTK signals (**Figure 4K**).

***RTK resensitization promotes rapid signal reactivation upon ALK inhibition.***
RTK stimulation promotes drug tolerance and acquired resistance across cancer types
(Straussman et al., 2012; Wilson et al., 2012). We thus sought to understand whether
hypersensitization of RTKs could promote RTK signaling during ALKi therapy. We monitored
RTK/Erk signaling in drug-treated populations of STE-1 cells using ErkKTR, a biosensor that
reports on Erk activity through nuclear exclusion of a fluorescent protein (Regot et al., 2014).
(**Figure 5A)**. In the absence of ALKi, Erk activity was at an intermediate level, indicated by
relatively equal distribution between the cytoplasm and nucleus (**Figure 5B, Supplementary
Movie 3**). Notably, this localization pattern did not change over the course of 22 hrs, reflecting
the tonic signaling downstream of EML4-ALK. By contrast, treatment with ALKi induced a rapid
initial decrease of Erk activity followed by a striking appearance of Erk activity pulses within ~1
hr after treatment (**Supplementary Movie 3**). Each Erk pulse lasted 10-20 minutes, and the
pulse amplitude exceeded levels observed in the absence of drug, paralleling our earlier results
with optogenetic or ligand stimulation (**Figure 5C**).
Activity pulses appeared sporadically but in a spatially coordinated manner, appearing
either simultaneously or as a traveling wave within small clusters of neighboring cells (**Figure
5B, Supplementary Movie 3**). This pattern was consistent with RTK stimulation through
paracrine signaling. To test this hypothesis, we sought to block paracrine signals through
inhibition of either the EGF receptor (EGFRi, 1 μM erlotinib) or matrix metalloproteases (MMPi,
10 μM marimastat), which release EGFR ligands from the cell surface to enable paracrine
signaling (Dong et al., 1999; Peschon et al., 1998; Sahin et al., 2004). Co-treatment with ALKi
and EGFRi suppressed Erk pulses after drug addition (ALKi vs ALKi/EGFRi: 58 ± 7% vs 12 ±
5% of cells showing ≥ 1 pulse over 22 hrs; error = 95% CI), indicating EGFR activation causes
the observed ERK pulses (**Fig. 5D,E,F, Supplementary Movie 3**). Co-treatment with ALKi and
MMPi similarly reduced Erk reactivation pulses (ALKi vs ALKi/MMPi: 58 ± 7% vs 36 ± 7% of
cells showing ≥ 1 pulse over 22 hrs; error = 95% CI), though to a lesser extent than with EGFRi,
potentially due to MMP-independent juxtacrine signals (Brachmann et al., 1989; Wong et al.,
1989) (**Fig. 5D,E,F, Supplementary Movie 3**). Thus, ALKi treatment in STE-1 cells decreases
Erk signals but is rapidly followed by RTK reactivation mediated by paracrine signals.

**Signal reactivation results from paracrine signals from dying cells.**
We next sought to determine the source of the paracrine signals. We observed that
pulsing events often appeared next to dying cells, and that co-inhibition of ALKi with EGFRi or
MMPi prevented this pulsing (**Figure 5G,H, Supplementary Movie 3)**. These observations are
consistent with paracrine ligand secretion from apoptotic cells, which has previously been
observed to promote cell survival and homeostasis within epithelial sheets (Aikin et al., 2020;
Gagliardi et al., 2021; Valon et al., 2021). To quantify this effect, we measured signal activation
in cells that neighbored a dying cell within the ~hour preceding its death (**Figure 5I).** We then
counted Erk pulses in these neighbors (N) and compared pulse counts to those from randomly
selected non-neighbors (R) over that same time interval (for more details, see **Methods).** Our
analysis revealed that, in the absence of ALKi, pulses were almost never observed in either the
N or R populations (DMSO, **Figure 5 J,K**). However, in ALKi-treated cells, N cells pulsed
significantly more than R cells (N vs R: 27 ± 4% vs 4 ± 2% cells with ≥ 1 pulse; error = 95% CI)

(**Fig. 5J,K**). Co-treatment with EGFRi eliminated Erk pulsing in neighbors (N cells in ALKi vs
ALKi/EGFRi: 27 $\pm$ 4% vs 2 $\pm$ 1%), whereas co-treatment with MMPi dramatically reduced
neighbor signaling relative to ALKi alone (N cells in ALKi vs ALKi/MMPi: 27 $\pm$ 4% vs 10 $\pm$
2%),(**Fig. 5J,K).** Together, our results demonstrate that virtually all observed Erk reactivation is
associated with paracrine signals associated with dying drug-treated cells. Importantly, because
Erk pulses were not observed in untreated cells —even in neighbors of dying cells (**Figure 5J**)—
these data suggest that ALKi-induced RTK hypersensitization is an essential first step for the
perception of paracrine ligands during ALKi therapy.
280
***Signal reactivation pulses activate gene transcription***
To understand the extent to which the short ALKi-dependent Erk pulses could impact cell
behavior, we asked whether the observed pulses could stimulate downstream transcription. Erk
activity is a strong driver of transcription, including of a class of rapidly responding immediate
early genes (IEGs) that begin transcription within minutes of Erk activity(Wilson et al., 2017).
EGR1 is an IEG that has been implicated in drug resistance to ALK inhibitors (Voena et al.,
2013). Additionally, EGR1 expression is adaptive, such that its expression peaks by ~ 1 hr but
then decays within 1-2 hours, even in the presence of constant upstream signal (Bugaj et al.,
2018; Sukhatme et al., 1987) (**Figure S8A**). Thus, accumulation of EGR1 indicates the presence
of only recent Erk activation(Davies et al., 2020). We thus examined the extent to which EGR1
accumulated in STE-1 cells upon drug treatment (**Fig. 6A**). In untreated cancer cells, EGR1
levels remained low despite high Erk signaling from EML4-ALK, consistent with EGR1
adaptation to tonic Erk signals (**Figure 6B,C**). Upon ALK inhibition, a distinct peak of EGR1-high
cells appeared and grew at 4 and 6 hrs after drug treatment, indicating that ALKi-induced Erk
pulses could indeed drive transcription. Importantly, co-inhibition of ALK and EGFR prevented
the appearance of EGR1-positive cells, consistent with transcription resulting from paracrine
signals through EGFR (**Figure 6B,C**). Thus, ALKi-induced Erk activity pulses provide sufficient
signal to drive gene expression changes that could regulate cell fate.
299
***Signal reactivation promotes acute drug tolerance and cell persistence during ALK***
***inhibition.*** Finally, we asked whether RTK hypersensitization and resultant Erk pulses could
counteract cell killing and promote drug tolerance to ALKi therapy (**Fig. 6D).** In STE-1 cells
imaged over 22 hrs, ALKi treatment induced moderate cell death (45 $\pm$ 7% of cells; error = 95%
CI). However, co-treatment of ALKi with either EGFRi or MMPi increased cell death (52 $\pm$ 7%
and 75 $\pm$ 6% of cells, respectively) (**Fig. 6E**). In both ALKi- and ALKi/MMPi-treated cells,
surviving cells showed an increased number of Erk pulses compared to cells that died (ALKi,
live vs dying: 1.8 $\pm$ 0.2 vs 0.9 $\pm$ 0.1 pulses per cell; ALKi/MMPi, live vs dying: 1.6 $\pm$ 0.2 vs 0.4 $\pm$
0.1 pulses per cell. Mean $\pm$ SE, **Figure 6F**), further drawing a link between RTK reactivation and
cell survival. We confirmed our results through an independent assay of cell death using a
fluorescent reporter of caspase-3 activity (NucView), an indicator of apoptosis, over the first 24
hr of treatment in EML4-ALK+ cells. As before, while ALK inhibition led to increased caspase-3,
co-treatment with EGFRi significantly increased the caspase-3+ cell fraction in STE-1 cells (28
$\pm$ 2 vs. 41 $\pm$ 2%). Similarly, co-treatment with MMPi also increased caspase-3 (from 28 $\pm$ 2 to 50
$\pm$ 6%) (**Figure 6G**). Similar trends were observed in H3122 cells (**Figure S8B**). Neither EGFRi
nor MMPi treatment alone showed enhanced killing over untreated cells. Notably, addition of 50
ng/mL EGF reversed enhanced killing in the ALKi/MMPi condition in STE-1 cells (61 $\pm$ 2 to 39 $\pm$
1%, respectively), further suggesting that synergistic killing in the presence of MMPi resulted

from blockade of signal reactivation (**Figure 6H**). Although EGF addition could not rescue
enhanced killing under combined ALKi/EGFRi treatment, optogenetic pulses of optoFGFR
lowered cell death in this condition (**Figure S8C,D,E**). Finally, we observed that co-treatments
of ALKi with either EGFRi or MMPi suppressed long-term drug tolerance measured at 17-days
(**Figure 6I**). Together, our results indicate that hypersensitization of EGFR and restored
perception of paracrine ligands leads to reactivation of survival signals soon after drug treatment,
which limits the cytotoxicity of ALK therapies and promotes drug tolerance, the first step towards
acquired resistance. The mechanisms that underlie these events may thus provide novel
treatment co-targets to enhance therapy in EML4-ALK+ cancers.
327
**Discussion**
Our results advance a novel function for protein condensates in cancer cells and their
response to targeted therapy (Jiang et al., 2020) (**Figure 7**). In addition to amplifying oncogenic
signaling (Tulpule et al., 2021), EML4-ALK condensates simultaneously act as a molecular
sponge for RTK adapters including Grb2 and SOS. This sequestration desensitizes the cell's
perception of external ligands by limiting the amount of adapters available to transduce the
activated RTK signal. However, upon ALK inhibition, adapters are rapidly released to the
cytoplasm, hypersensitizing cellular response to ligands in the cell's microenvironment. RTK
hypersensitization promotes cell survival and drug tolerance in response to targeted inhibitors,
due at least in part to apoptosis-induced paracrine signals.
We show that protein condensation provides a mechanism by which oncogenes can
suppress RTK signaling. Analogous RTK suppression has been observed in other, molecularly
distinct cancers. BRAF V600E+ melanoma and colorectal cancer cells suppress RTKs through
Erk-dependent transcription of negative regulators (Sproutys), and through inhibitory
phosphorylation of SOS1(Gerosa et al., 2020; Lito et al., 2012; Prahallad et al., 2012). Inhibition
of BRAF suppresses these mechanisms and leads to rapid, pulsatile Erk reactivation and drug
resistance, similar to our observations. Separately, in cancers driven by EGFR, Erk provides
suppressive phosphorylation of EGFR receptors, which is lost during MEK inhibition and leads
to reactivation of ErbB3 and PI3K(Turke et al., 2012). In contrast to these and similar examples,
our findings demonstrate that such RTK suppression can also be implemented through
biophysical, rather than biochemical, feedback. Nevertheless, the variety of mechanisms by
which oncogenes can suppress RTKs hints that such feedback might play an important role in
establishing permissive conditions for oncogenesis. It will thus be interesting to more
comprehensively understand the diverse oncogenic contexts in which such suppression and
reactivation occurs.
It has been repeatedly observed that ligand-induced signaling through EGFR promotes
survival and resistance to ALK inhibitors in EML4-ALK+ cancer cells (Obenauf et al., 2015;
Sasaki et al., 2011; Tani et al., 2016; Tanimoto et al., 2014; Vaishnavi et al., 2017; Vander Velde
et al., 2020; Wilson et al., 2012). Our work builds on these prior studies to show that 1) ALK
inhibition is a necessary first step to allow cancer cell perception of paracrine signals, 2) EGFR
ligands are sent from apoptotic neighbor cells, and 3) ALK-induced resensitization to RTK
signals happens within minutes of drug treatment. In addition to fast RTK reactivation, previous
work found that slower-timescale transcriptional amplification of KRas or downregulation of
DUSP6 contributes to resistance development (Hrustanovic et al., 2015). Our results are
complementary to these findings, which could be expected to further sensitize cells to external
RTK ligands.

Dying cancer cells send survival signals to their neighbors through a mechanism that
requires proteolytic processing of EGFR ligands, thus adding to previously identified
mechanisms whereby drug treatment induces secretion that promotes cell survival and drug
tolerance (Kurtova et al., 2015; Obenauf et al., 2015). Blockade of apoptosis-induced paracrine
signals enhanced cell killing and limited tolerance in response to ALK inhibition. Although co-
inhibition of EGFR is highly effective in promoting durable drug responses across a variety of
cancer cells, (Sasaki et al., 2011; Vaishnavi et al., 2017; Vander Velde et al., 2020; Voena et al.,
2013), a recent clinical trial of combined inhibition of crizotinib (ALKi) and erlotinib (EGFRi) in
EML4-ALK+ NSCLC failed due to frequent adverse effects and low maximum tolerated dose
(Ou et al., 2017). We show that co-inhibition of certain matrix metalloproteases may be an
alternative strategy limit RTK signaling during targeted therapy. We note, however, that previous
studies found that long-term MMP inhibition can promote resistance by causing accumulation of
transmembrane RTKs (e.g. AXL) that are also MMP targets (Miller et al., 2016). Thus, further
studies will be required to understand whether MMPi co-treatment can indeed enhance therapy,
or whether MMPi scheduling could leverage acute benefits while avoiding deleterious chronic
effects.
Our finding that the sequestration of RTK adapters can functionally desensitize RTK
signals could inspire new types of therapeutic targets that mimic this behavior, for example
through inhibition or sequestration of important RTK adapters. Of note, the recent discovery of
EML4-ALK condensates has raised the idea that disaggregation of the condensates might be
therapeutically beneficial (Cai et al., 2021; Hirai et al., 2020). Our work cautions, however, that
such disaggregation strategies will likely be subject to the same adapter redistribution and rapid
RTK reactivation observed with small molecule ALK inhibitors, potentially challenging their
efficacy.
A unique promise of functional profiling is that common signaling abnormalities may be
identified and inform therapies among genetically distinct cancers (Bugaj et al., 2017). Because
our work revealed signaling principles that result from a likely common property (condensation)
of RTK fusions (Du and Lovly, 2018; Tulpule et al., 2021), we anticipate that our findings may
be broadly relevant to other cancer cells in this class. We show that in a cancer line driven by a
CCDC6-RET fusion that also forms large aggregates (Tulpule et al., 2021), RET activity similarly
suppresses EGFR, and RET inhibition potentiates EGFR signaling (**Figure S7)**. These results
are consistent with a previous study that showed that Grb2 can change association from the
fusion to EGFR upon kinase inhibition in cancers driven by ALK, RET, NTRK1 and ROS1
fusions, including in patient samples of primary tumors, resistant tumors, and
metastases(Vaishnavi et al., 2017). These findings imply common behavior across molecularly
distinct RTK fusions and further point to the clinical relevance of these phenomena. Future work
will determine the extent to which the principles we describe in EML4-ALK+ cancers will extend
to this more diverse array of malignancies.
402

**Acknowledgements.** The authors thank Dr. Magdalena Niewiadomska-Bugaj for feedback on
statistics, data presentation and analysis. The authors also thank Dr. Alex Hughes (Penn), Dr.
Sydney Shaffer, and Dr. Yael Mossé for helpful comments on the manuscript. This work was
supported by the National Institutes of Health (R35GM138211 for L.J.B and D.G.M.;
R01CA231300 for B.H. and T.G.B; U54CA224081, R01CA204302, R01CA211052 and
R01CA169338 for T.G.B.), UCSF Marcus Program in Precision Medicine Innovation to B.H. and

T.G.B, and the UCSF Physician Scientist Scholar Program and Alex's Lemonade Stand for A.T.
B.H. is a Chan Biohub Investigator.
411

412 **Author Contributions** D.G.M. and L.J.B. conceived the study. D.G.M. and L.R. performed
413 experiments. J.G. and B.H. generated endogenously-tagged cell lines and provided guidance
414 on imaging. D.G.M., T.R.M. and L.J.B. analyzed data. A.T., T.G.B., and L.J.B. provided
415 experimental feedback and guidance. L.J.B. supervised the work. D.G.M and L.J.B. wrote the
416 manuscript and made figures, with editing from all authors.
417

418 **Conflicts of interest.** D.G.M. and L.J.B. have filed a provisional patent based on the findings in
419 this work. T.G.B. is an advisor to Novartis, Astrazeneca, Revolution Medicines, Array/Pfizer,
420 Springworks, Strategia, Relay, Jazz, Rain, Engine, Scorpion and receives research funding from
421 Strategia, Kinnate, Verastem, and Revolution Medicines. A.T. is an advisor to Faze Medicines.
422
423
424

# Figures

**Figure 1. Optogenetics functional profiling of EML4-ALK+ cancer cells reveals suppression of RTK signaling.** A) EML4-ALK+ cancer cells treated with ALK inhibitors (ALKi) can persist through therapy and acquire stable drug resistance. B) Immunofluorescence staining shows punctate expression of ALK in two EML4-ALK+ cancer cell lines. C) Although EML4-ALK and transmembrane RTKs signal through the same downstream adapters, the extent to which EML4-ALK condensates interact with RTKs, and whether interactions could impact drug responses, is not clear. D) Functional profiling allows us to observe input/output responses to RTK signaling to understand functional interactions between RTKs and EML4-ALK. E) optoFGFR allows blue-light-induced stimulation of FGFR signaling, including through the Ras/Erk pathway. F) Single-cell immunofluorescence analysis of ppErk levels in STE-1 cells stimulated with light (optoFGFR) in the presence of either DMSO (grey) or the ALK inhibitor crizotinib (ALKi, 1μM; orange). F) ppErk fold-change in response to 5 min of blue light stimulation. Blue light intensity ranged from 2% to 60% of maximum (160 mW/cm$^2$). Data points show the ratio of ppErk from stimulated and unstimulated cells. Each condition was performed in duplicate, with each replicate representing the mean of ~4000 cells.

442
443
444
445
**Figure 2. EML4-ALK suppresses EGFR signaling.** A) EGF stimulates EGFR and downstream
Ras/Erk signaling. B) ppErk levels in response to EGF (100 ng/mL) in the presence of 1µM ALKi
(orange) or DMSO (grey) in STE-1 and H3122 cancer cells. Data points and ribbons represent
mean +/- SEM of three replicates, each of which represents the mean of ~1000-3000 cells
measured through IF. C) Quantification of dynamic range from experiment in 2B over a range of
EGF concentrations. D) EML4-ALK(V1) was transiently expressed in lung epithelial Beas2B
cells. E) Time course of ppErk immunofluorescence levels in response to EGF stimulation (50
ng/mL). Data points represent mean +/- SEM of three replicates, each of 200-400 transfected or
1000-2000 untransfected cells. F) Dynamic range of ppErk in EML4-ALK-expressing Beas2B in
response to EGF in the presence or absence of ALKi (1 µM) pretreatment.

456
**Figure 3. Mapping RTK feedback suppression using optogenetics.** A) Time course of RTK
hypersensitization was obtained by pre-incubating cancer cells with 1 μM crizotinib for a variable
period before stimulation with EGF (50 ng/mL) for 15 min, followed by fixation and
immunostaining for ppErk. B) ppErk induction as a function of ALKi pre-incubation time,
quantified by immunofluorescence. Open circles = unstimulated. Closed circles = EGF
stimulated. Data points represent mean +/- SEM of triplicates, each consisting of 1000-3000
cells. Error bars are smaller than the data points for unstimulated samples. C) Pinpointing the
location of EML4-ALK interaction with RTK/Erk signaling. D) OptoFGFR and optoSOS permit
optogenetic stimulation at successive nodes of the pathway. E) Quantification of
immunofluorescence of ppErk in response to optoFGFR or optoSOS in the presence or absence
of ALKi. ALK-dependent suppression is observed only with optoFGFR, suggesting that
suppression happens upstream of Ras but downstream of RTK activation (F). G) Testing the
role of Erk-dependent negative feedback on RTK suppression. Stimulating optoSOS drives
elevated basal levels of ppErk during ALKi treatment and sustains any Erk-dependent negative
feedback that would otherwise be lost during ALK inhibition. H) STE-1 cells were treated with
either ALKi or ALKi and optoSOS, and EGF response was assessed. I) Predicted results and
implications for Erk-dependent feedback. J) Cells treated with both ALKi and optoSOS showed
comparable increase in EGF-induced ppErk as cells receiving ALKi only, suggesting that Erk-
dependent negative feedback does not suppress EGFR in EML4-ALK+ cancer cells. Data points
represent means of ~500-900 cells per condition. ** p < 0.01 by Student's t-test.

**Figure 4. EML4-ALK condensates suppress RTK signals through sequestration of RTK effectors.** A) Quantification of ppErk response to EGF (50 ng/mL) stimulation in Beas2B cells that expressed EML4-ALK (V1), EML4-ALK (ΔTD), and kinase-dead EML4-ALK (K589M). Data are means +/- SEM of triplicates, each of which represents 500-1000 cells.  Data points represent mean. B) Beas2B cells that harbored endogenously-tagged Grb2 (Grb2:mNG) were transiently transfected with EML4-ALK and stimulated with EGF (50ng/mL) to visualize Grb2 translocation in the presence and absence of EML4-ALK. C) Impaired membrane translocation of Grb2 in the presence of EML4-ALK condensates. Time in mm:sec. See **Supplementary Movie 1**. D) Line scan of Grb2 intensity distribution in the presence (red) or absence (green) of EML4-ALK expression, as depicted in C. E) Grb2 localization was visualized upon treatment

with 1 µM ALKi. F) Grb2 shifted from condensates to cytoplasm after ALKi treatment. Time in
hh:mm. See **Supplementary Movie 2**. G) Quantification of Grb2 cluster dissociation kinetics
after ALKi treatment. H) Grb2 distribution was examined upon sequential treatment with ALKi
and EGF. I) ALKi released Grb2 to the cytoplasm, which allowed its membrane translocation in
the presence of EGF. J) Immunoprecipitation of EGFR shows co-precipitation of SOS1 only in
the presence of ALKi pretreatment and EGF. K) Conceptual model of how EML4-ALK
suppresses transmembrane RTKs. EML4-ALK sequesters adapters like Grb2/SOS1 and
prohibits their translocation to activated RTKs. ALK inhibition releases adapter sequestration
and restores cellular response to RTK stimulation.
504

505
506
507

**Figure 5. ALK inhibition hypersensitizes cancer cells to paracrine growth factors secreted from dying neighbor cells.** A) The ErkKTR reporter indicates Erk activity through nuclear-cytoplasmic translocation of a fluorescent protein. B) Live cell imaging of STE-1 cells expressing ErkKTR in the presence or absence of ALKi (1 μM crizotinib). See **Supplementary Movie 3**. C) Representative single-cell traces of cytoplasmic/nuclear ErkKTR intensity ratio from conditions shown in B. D) Quantification of ErkKTR activity in the presence of ALKi or its combination with EGFRi (1 μM erlotinib) or MMPi (10 μM marimastat). E) Quantification of Erk activity pulses per

cell. F) Fraction of cells that exhibited any pulses over 21.5 hrs of imaging. Error bars indicate 95% CI. n = 200 cells per condition. G) Apoptotic cells can secrete paracrine EGFR ligands to their neighbors. Paracrine signaling can be blocked by inhibiting either EGFR or the MMPs that are required to shed certain EGFR ligands from the surface of the sender cell. H) ErkKTR activity pulses are primarily observed surrounding a dying cell during ALK inhibition, but not in the absence of drug or in the presence of EGFR or MMP inhibitors. I) Definition of neighbors and non-neighbors of a death event. J) Quantification of pulses per cell for each death event in neighbors or a randomly chosen subset of cells not near a death event (see **Methods** for more details). K) Fraction of total neighbor vs random non-neighbor cells that show any pulsing. Error bars = 95% CI.

**Flgure 6. Resensitization to paracrine signals drives transcription and promotes drug tolerance.** A) Signaling through EGFR activates Ras/Erk and stimulates transcription, including of the immediate early gene EGR1. B) Upon ALK inhibition, EGR1 expression increases, but not upon co-inhibition of ALK and EGFR. C) Comparison of EGR1 expression after 6 hrs of the indicated treatment. D) We hypothesized that restored perception of paracrine signals might promote survival, and that blocking paracrine signals would promote killing. E) Visualization of live cell imaging of Erk activity pulses and cell death. F) Quantification of pulses per cell in cells that died (D) or survived (L) through 22 hrs of imaging, in the conditions where Erk pulsing could be observed. G) Caspase-3 activation was assessed using the NucView reporter after 24 hr treatment with the indicated drugs. H) EGF (50 ng/mL) addition promotes survival during ALKi treatment and counteracts enhanced killing from ALKi/MMPi co-treatment, but not from ALKi/EGFRi co-treatment. Dotted line is cell killing from DMSO control treatment. Each data point represents the fraction of caspase-3+ cells from 2000-3000 cells.  See **Methods** for more

540    details. I) Crystal violet staining (left) and cell counts (right) of cell survival after 17 days of the
541    indicated treatments.

542
543
**Figure 7. Model of how EML4-ALK condensates interact with transmembrane RTKs and**
**its effect on drug tolerance.** We find that EML4-ALK condensates suppress perception of
EGFR signals by sequestering RTK adapters like Grb2 and SOS1. Upon ALK inhibition,
adapters are released to the cytoplasm where they can transduce signals from EGFR and other
RTKs. Although ALK inhibitors effectively suppress ALK activity, they simultaneously restore a
cell's perception to paracrine signals, including those from neighboring apoptotic cells. Such
paracrine signals promote drug tolerance, and blocking them with inhibitors of either EGFR or
MMPs enhances cell killing during ALK inhibitor therapy.
552
553

**Supplementary Figures**

**Supplementary Figure 1. OptoFGFR functional profiling of STE-1 cancer cells.** A) optoFGFR comprises the intracellular domain of FGFR1 fused to the PHR domain from Arabidopsis Cryptochrome 2, which clusters under blue light stimulation(Bugaj et al., 2013). The construct is anchored in the membrane through N-terminal myristoylation. B) Single cell distributions of ppErk intensity in STE-1 cells stimulated by the indicated intensity of blue light in the presence (orange) or absence (grey) of ALKi (2 hr pre-incubation of 1µM crizotinib). Strong ppErk response at low light intensity in the presence, but not the absence, of ALKi suggests that ALKi hypersensitizes cells to RTK stimulation. C) The magnitude of ppErk response in optoFGFR STE-1 cells is a function of both light intensity and optoFGFR expression levels. Results are presented by binning optoFGFR-mCh expression by quartiles. ( 1<25%, 25% < 2 < 50%, 50% <3 < 75%, 4 > 75%).

**Supplementary Figure 2. Schematics of the optoFGFR and optoSOS tools used in Figure 3.** optoFGFR(Kim et al., 2014) allows optogenetic stimulation of the FGFR receptor using light-induced clustering of the Cryptochrome 2 protein(Bugaj et al., 2013). OptoSOS allos Ras activation through membrane recruitment of the SOS2 catalytic domain (SOS2$_{cat}$). Membrane recruitment is achieved through blue light dimerization of sspB to the iLid protein(Guntas et al., 2015) which is anchored to the membrane through a CAAX motif.

**Supplementary Figure 3. EGFR signal suppression does not result from decreased EGFR**
**phosphorylation.** Following overnight starvation, STE-1 cells were pretreated with ALKi or
DMSO for 2 hrs. Cells were then stimulated with either EGF (50 ng/mL) for 15 min and then
were fixed and immunostained for ppErk. ALKi pretreatment did not alter pEGFR (Y1068) levels
in either the presence or absence of EGF, whereas pALK (Y1507) decreased. Thus EML4-ALK
does not modulate EGFR signaling through changes in EGFR phsophorylation. Data points
represent mean intensity of 3000-5000 cells.
589

590
**Supplementary Figure 4. Determining the role of Erk-dependent negative feedback on**
**RTK suppression.** A) Demonstration that optoSOS can maintain high Erk levels (comparable
to those from the uninhibited oncogene) during ALKi pre-treatment. Optogenetic stimulation of
Ras/Erk was used to decouple ALK inhibition from loss of Erk signaling in **Figure 3G-J**. B) ppErk
levels after 15 min of EGF stimulation in the presence or absence of optoSOS activation.
Maintenance of optoSOS activation during EGF stimulation did not result in elevated Erk
signaling in the absence of ALKi (as seen in **Figure 3J**). Data in A) and B) represent      signal
from top 25% of optoSOS-expressing cells. Data points represent mean ± SEM of 200-600 cells
per condition. C) Western blot showing levels of negative regulator Spry2 over the first hour of
ALKi treatment. Despite loss of Erk activity, Spry2 levels remain unchanged over this time period.
601

Grb2-mNG      EML4-ALK-mCh      Merge

603
604
605 **Supplementary Figure 5. EML4-ALK condensates colocalize with Grb2 puncta.** Beas2B
606 cells that harbored endogenously tagged Grb2 (Grb2:mNG) were transfected with EML4-ALK-
607 mCh. In transfected cells, EML4-ALK co-localized with Grb2. In untransfected cells, Grb2 was
608 diffuse.
609

610   **Supplementary Figure 6. Co-immunoprecipitation of EGFR with SOS1.** H3122 cells were
611   starved overnight, pretreated with crizotinib (2 hr), and were stimulated with EGF for 2 minutes
612   before lysis and immunoprecipitation. Right 4 lanes are reproduced from **Figure 4J**.

617
**Supplementary Figure 7. CCDC6-RET+ cancer cells show dynamic suppression of EGFR**
**through RTK fusion activity.** TPC-1 cells harbor CCDC6-RET, a distinct RTK fusion that forms
condensates.  We tested the extent to which CCDC6-RET could modulate EGFR dynamic range
as observed for EML4-ALK. A) Plot shows ppErk response to EGF stimulation after 2 hr pre-
treatment with DMSO (grey) or 100 nM BLU-667 (purple), a RET inhibitor. Data represent mean
± SEM of triplicates, each of which represents ~1000 cells. B) Quantification of ppErk dynamic
range shows that oncogene inhibition enhances signaling through EGFR.  Thus, multiple RTK
fusions that form condensates dynamically suppress EGFR signaling.
626
627
628

630
**Supplementary Figure 8. RTK/Erk pulses activate transcription and promote cell survival**
**during ALKi treatment.** A) Adaptive response of EGR1 expression in STE-1 cells treated with
ALKi and 50 ng/mL of EGF.  EGR1 peaks at 1 hr after stimulation and then adapts to near
baseline expression. EGR response is stronger and earlier than observed in **Figure 6B** because
here all cells were stimulated at t = 0, whereas in **6B** only individual cells were stimulated
sporadically through paracrine signals starting 1-2 hrs after ALK treatment. B) Blockade of EGFR
or MMPs accelerates cell death in H3122 in response to ALKi, as shown STE-1 cells in **Figure**
**6G**. C) We tested whether optoFGFR stimulation could reduce the enhanced cell death response
in STE-1 cells co-treated with ALKi and EGFRi.  D). A 10 min pulse of blue light (96 mW/cm$^2$)
every hour produced strong, periodic ppErk levels. Data points represent mean of three
replicates, each representing ~2000 cells.  E) Hourly pulses of optoFGFR/Erk (10 min of 96
mW/cm$^2$) reduced cell death in both ALKi and ALKi/EGFRi treated cells. Data points represents
the mean +/- SEM of three replicates, each representing ~10,000 cells.
644
645
646

**Methods**

Cell lines and cell culture

     All cell lines were maintained at 37 °C and 5% $CO_2$ using a standard cell culture incubator. STE-1, H3122, TPC-1, and Beas2B cells were cultured in RPMI-1640 growth medium (RPMI-1640 containing L-glutamine supplement with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin (P/S)). For experiments, cells were seeded in 96 or 384 well plates coated with fibronectin (MilliporeSigma, FC01010MG) diluted to 10 µg/mL in PBS. For 96 well plate experiments 10,000 STE-1 and H3122 cells were seeded in 200 µl of cell culture media per well. For 384 well plate experiments 5,000 STE-1, H3122, TPC-1 cells or 2500 Beas2B cells were seeded in 50 µl of culture media per well. LentiX- HEK 293T cells (TakaraBio, #632180) were cultured using DMEM supplemented with 10% FBS and 1% P/S.

*Plasmid design and assembly*

     All cloning was performed by PCR and DNA assembly using NEBuilder® HiFi DNA Assembly Master Mix (New England Biolabs #E2621). Constructs for transient expression were cloned into pEGFP-C1 backbones encoding a CMV promoter. Fluorescently-tagged EML4-ALK (pCMV EML4-ALK-GFP) was generated by inserting the coding sequence for EML4-ALK(V1) upstream of GFP. To generate untagged EML4-ALK for transient expression in Beas2B cells, EGFP was replaced with P2A-H2B-iRFP (pCMV EML4-ALK-P2A-H2B-iRFP). EML4-ALK (ΔTD) mutant was cloned by excluding bases 310-459 from pCMV-EML4-ALK-P2A-H2B-iRFP through PCR amplification followed by plasmid assembly using HiFi. EML4-ALK (K589M) was cloned by single point mutation encoded on the primer used to amplify pCMV-EML4-ALK-P2A-H2B-iRFP followed by blunt end ligation using NEB T4 ligase (NEB, #M0202). Constructs for generation of stable cell lines were cloned into pHR lentiviral or CLPIT retroviral plasmid backbones. For optogenetic control of FGFR signaling (Kim et al., 2014), the FGFR intracellular domain with a N-terminal myristoylation site was inserted upstream of mCh-Cry2(Bugaj et al., 2013) in a CLPIT plasmid to create CLPIT Myr-mCh-FGFR(ICD)-Cry2. For optogenetic control of SOS signaling, we generated pHR sspB-SOScat-mCh-2A-iLid-CAAX by inserting an mCherry coding sequence to replace BFP, which we described previously (Benman et al., 2022). For live-cell tracking of ppErk activity, we generated pHR ErkKTR-mRuby2 by inserting an mRuby2 coding sequence in place of BFP in a construct we described previously (Benman et al., 2022). Visualization of nuclei was achieved using pLenti PGK DEST-H2B-iRFP670 (Addgene plasmid #90237).

Transient transfection

     pCMV-EML4-ALK-P2A-H2B-iRFP, pCMV-EML4-ALK(ΔTD)-P2A-H2B-iRFP, pCMV-EML4-ALK(K589M)-P2A-H2B-iRFP and CMV-3XFLAG-mCh-EML4-ALK were transiently transfected into wt Beas2B or Beas2B mNG:Grb2 cells using Lipofectamine™ 3000 (Invitrogen, L3000001). Briefly, $2.5 \times 10^3$ cells were seeded in fibronectin-coated 384 well plates in RPMI growth medium. For an individual well: 0.25 µl of Lipofectamine 3000 reagent was diluted in 1 µl of OptiMEM (Gibco, #31985070) and incubated at room temperature for 5 min. This mix was then added to a diluted DNA mix composed of 25 ng of plasmid DNA and 0.5 µl of P3000 reagent, diluted in OptiMEM to a final volume of 1.25µl. After 15 min of incubation at room temperature, transfection mix was added to cells cultured in 50 µl culture medium. Following 6 hrs of transfection, media was exchanged with fresh growth medium. Cells were then starved

693  for at least 6 hrs using starvation medium (RPMI + 1% P/S) with or without phenol red for fixed
694  or live-cell assays, respectively.
695
<u>Generation of stable cell lines</u>
Cell lines were generated using viral transduction. CLPIT-Myr-mCh-FGFR(ICD)-Cry2,
pHR sspB(nano)-mCh-SOS2(cat)-2A-iLid-CAAX, pHR KTR-mRuby2, and PGK-DEST-H2B-
iRFP670 vectors were packaged into lentivirus using HEK 293T cells. Briefly, $7.0 \times 10^5$ cells
were seeded in each well of a 6 well plate. For pHR transfer vectors, cells were transfected with
1.5 µg of pHR transfer vector, 1.33 µg of pCMV-dR8.91 (Addgene #12263), and 0.17 µg pMD2.G
(Addgene #12259). For CLPIT transfer vectors, cells were transfected with 1.25 µg of transfer
vector, 0.5 µg of pCMV-VSVG (Addgene #8454) and 0.75 µg of pCMV-gag/pol. Transfections
were performed using the calcium phosphate method. The following day, medium was removed
and replaced with fresh growth medium.  After 24 and 48 hrs, virus-containing supernatant was
collected and stored at 4 °C. Supernatants were then centrifuged at 200 x g for 2 min to remove
cell debris, filtered through a 0.45 µm filter (Fisher Scientific, catalog number 13-1001-07), and
used immediately or stored at -80°C. For transduction, $2.0 \times 10^5$ H3122 or STE-I cells were
seeded along with 500 µl of viral supernatant. Transduced cells were expanded and sorted (BD
Influx) for appropriate expression levels. For STE-1 cells that expressed optoFGFR or optoSOS,
the top 20% or 30% of mCh+ cells, respectively were sorted, expanded, and used for
experiments. Beas2B cells expressing endogenously tagged Grb2:mNG (Beas2B Grb2:mNG)
generated previously (Tulpule et al., 2021).
714
<u>Optogenetic Stimulation</u>
Light stimulation was achieved in microwell plates using the optoPlate-96 (1-color blue
version(Bugaj and Lim, 2019)). A 20 mm tall black adaptor was used to ensure even light
diffusion across each of the 384 well plate wells. To investigate the ppErk response to optoFGFR
stimulation (**Figure 1 F,G**), cells were stimulated with 500 ms every 10 sec for 5 min (light
intensity ranging from $3.2 mW/cm^2$ (min) to $160 mW/cm^2$ (max)). To maintain basal ppErk levels
during ALK inhibition (**Figure 2 H,J**) optoSOS-expressing STE-1 cells were stimulated for 500ms
every 10 sec ($160 W/cm^2$ blue light) for 15 min, with stimulation beginning simultaneously with
drug treatment (1µM crizotinib). Cells were then stimulated with 50 ng/ml EGF (PeproTech, 315-
09) while still being exposed with blue light as previously described. For optogenetic rescue of
cell death during ALKi/ EGFRi treatment, cells were stimulated with blue light ($96 mW/cm^2$,
500msON/10s OFF) for 10 min every hour, with stimulation beginning simultaneously with drug
treatment.
728
<u>Growth Factor Stimulation Assay</u>
Plated cells were starved overnight with RPMI-1640 starvation medium. Starvation was
achieved by performing 7X 80% washes either manually or using the BioTek 405 LS microplate
washer. Cells were treated with inhibitor for 2 hours (1µM crizotinib (Sigma-Aldrich, PZ0191) or
0.1µM BLU-667 (Biovision, B2548)). Cells were then stimulated at their respective time points
with EGF following a delay that allowed for all wells to be fixed simultaneously. Cells were fixed
with 4% paraformaldehyde (PFA) (Electron Microscopy Sciences, 15710) by adding the
appropriate volume of 16% PFA to the culture medium to yield a final concentration of 4%.
Samples were then immediately prepared for immunostaining.
738

<u>Immunofluorescence</u>

740    After PFA fixation (10 mins), cells were permeabilized with 0.5% Triton-X100 in PBS for
741 10 min, followed by incubation in ice-cold 100% methanol for 10 min. Samples were then blocked
742 with blocking solution (1% bovine serum albumin (BSA) (Fisher, BP9706100) diluted in PBS) for
743 1 hour at room temperature. Samples were incubated in primary antibody diluted in blocking
744 solution for either 2 hours at room temperature (RT) or overnight at 4 °C. Primary antibodies
745 used were: phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204), Cell Signaling #4370, 1:400
746 dilution; phospho-EGF Receptor (Tyr1068), Cell Signaling #3777, 1:800; EGR1, Cell Signaling
747 # 4153, 1:800). After incubation with primary antibody, samples were washed 7X with 80%
748 washes of 0.1% Tween-20 in PBS (PBS-T) using the BioTek 405 LS microplate washer.
749 Samples were then incubated in blocking solution containing secondary antibody (IgG (H+L)
750 Cross-Adsorbed Goat anti-Rabbit, DyLight™ 488, Invitrogen #35553, 1:500; Goat anti-Rabbit
751 IgG (H+L) Cross-Adsorbed Secondary Antibody, DyLight™ 650, Invitrogen #SA510034, 1:500)
752 and 4,6-diamidino-2-phenylindole (DAPI; ThermoFisher Scientific #D1306, 300 nM) for 1 hour at
753 RT. Samples washed with PBS-T as previously described. Samples were left in a final volume
754 of 100 μL PBS-T for imaging and storage.

756 <u>Immunoblotting and co-Immunoprecipitation</u>
757 For immunoblots STE1 or H3122 cells (3 X $10^5$) were plated in each well of a 6 well plate,
758 cultured for 24 hours, and subsequently serum starved for 16 hours. Cells were treated with 1
759 μM crizotinib for the times indicated, after which cells were washed in ice cold PBS and lysed in
760 RIPA buffer (50 mM Tris pH 7.5, 150 mM NaCl, 1% NP40, 0.1% SDS, 0.5% DOC, 1 mM EDTA,
761 2 mM sodium vanadate and protease inhibitor). Protein concentration of cleared cell lysates was
762 determined by BCA protein assay kit (Thermo Scientific #23225) and 20-30 μg of lysed samples
763 were subjected to SDS–polyacrylamide gel electrophoresis (SDS-PAGE). For co-
764 immunoprecipitation, STE1 or H3122 cells (2.5 X $10^6$) were plated on 10 cm plates, cultured for
765 24 hours, and subsequently serum-starved for 16 hours. Cells were treated with 1 μM crizotinib
766 and 10 ng/mL EGF as indicated, washed with ice cold PBS, and lysed (50 mM HEPES pH7.4,
767 150mM NaCl, 1% Triton X-100, 1 mM EDTA, 1 mM EGTA, 10% glycerol, 2 mM sodium vanadate
768 and protease inhibitor (Sigma #P8340)).  Cleared cell lysates were incubated for 2 hours with
769 Protein A/G agarose beads (Santa Cruz, SC-2003) that were hybridized with EGFR antibody
770 (Thermo, clone H11). Beads were then washed 5 times with HNTG buffer (20 mM HEPES pH
771 7.4, 150 mM NaCl, 0.1% Triton X-100, 10% glycerol), and sample buffer was added to elute
772 proteins. Eluates or 25–30 μg of protein lysate were loaded in a precast 4-15% gradient SDS–
773 polyacrylamide gel for electrophoresis (mini-protean TGX precast gel, Bio-RAD, # 456-1084).
774    Protein separations were transferred onto a nitrocellulose membrane using the Trans-blot
775 Turbo RTA transfer kit (Bio-rad, #170-4270) according to manufacturer's protocol. Membranes
776 were blocked in 5% milk in Tris buffer saline with 0.5% Tween-20 (TBS-T)  for 1 hour and
777 incubated overnight at 4°C with primary antibodies against EGFR (CST #4267), SOS1 (CST
778 #5890), SPRY2 (CST #14954), pERK1/2 (CST #4370), tubulin (CST #3873). Each primary
779 antibody was used at a dilution of 1:1000 in TBS-T with 3% BSA. After washing with TBS-T,
780 membranes with incubated with secondary antibodies in TBS-T with 3% BSA for 1 hr at room
781 temperature (IRDye® 800CW Goat anti-Rabbit IgG, 1;20,000 dilution, LI-COR #926-32211;
782 IRDye® 680RD Donkey anti-Mouse IgG, 1:20,000 dilution, LI-COR, #926-68072). Membranes
783 were then imaged on the LI-COR Odyssey scanner.

 Live cell imaging

Live cell imaging was performed using a Nikon Ti2-E microscope equipped with a
Yokagawa CSU-W1 spinning disk, 405/488/561/640 nm laser lines, an sCMOS camera
(Photometrics), and a motorized stage. Cells were maintained at 37 °C and 5% $CO_2$ using an
environmental chamber (Okolabs). Transfected Beas2B Grb2:mNG cells were imaged 48 hours
after transfection with a 40X oil immersion objective. To visualize Grb2 localization during drug
response and EGF stimulation, cells were imaged for 2 hours following addition of 1 µM crizotinib
and subsequent EGF stimulation, both added manually during image acquisition.  For monitoring
Erk activity after drug treatment, STE-1 cells stably expressing ErkKTR-mRuby2 and H2B-
iRFP670 were seeded in a 96 well plate (Falcon #353072) at a density of $1.5x10^4$ cells/well. The
following day, media was replaced with  phenol-free RPMI-1640 with L-glutamine supplemented
with 2% FBS and 1% P/S. Immediately before imaging, cells were supplemented with NucView
(Biotium #10402 , 1 µM) and the indicated drugs (crizotinib (1 µM), erlotinib (1 µM, Sigma-
Aldrich, #SML2156), marimastat (10 µM, Cayman, #14869)) or DMSO (1:5000)). Cells were
imaged every 5 minutes for 22 hours using confocal microscopy using 20X magnification.
800
Caspase-3 Activation and Drug Tolerance Assay
To assay acute cell killing, STE-I and H3122 cells were seeded in a 96 well plate. The
following day, media was replaced with phenol-free RPMI-1640 supplemented with 2% FBS, 1%
P/S, and treated with 1µM NucView 488 and the indicated (1µM crizotinib, 1µM erlotinib, 10µM
marimastat). 24 hours after drug addition, cells were incubated in 5 µg/mL Hoechst 33342
Hydrochloride (Cayman 15547) for 15 minutes. Cells were then imaged under live-cell confocal
microscopy using 10X magnification. To assay drug tolerance, H3122 cells were seeded in a 96
well plate as previously described. The following day, media was replaced with RPMI with 2%
FBS, 1% P/S and treated with the indicated inhibitors (1µM crizotinib, 1µM erlotinib, 10µM
marimastat). Media with inhibitors was replaced every 2 days. Following 17 days of treatment,
cells were fixed with 4% PFA and permeabilized with 100% methanol for 10 min. Cells were then
incubated in DAPI (300 nM) for 30 min, imaged, and nuclei were counted through image analysis
813
Image processing and analysis
*Immunostaining and caspase-3 reporter*
Images of fixed-cell immunostaining and live-cell caspase-3 reporter were quantified
using CellProfiler (v 4.0.7)(Lamprecht et al., 2007). Briefly, cell nuclei were segmented using the
DAPI channel, and the cytoplasmic fluorescence was measured within a 5-pixel ring that
circumscribed the nucleus. For measuring caspase-3 activity and drug tolerance, DAPI or
Hoechst-stained nuclei were segmented using ilastik(Berg et al., 2019), quantified in CellProfiler,
and quantification was exported to R for processing and data visualization using the tidyR
package (Wickham et al., 2019).
823
*Live cell imaging of Grb2*
Time lapse imaging of Grb2 puncta was quantified with a custom MATLAB script. Briefly, cells
were segmented manually and clusters were identified in a semi-automated manner through the
following steps. First, cell intensities were normalized to their internal median. Images were then
passed through sequential Gaussian, top hat, and Laplacian filters to enhance clusters while
suppressing other features like background and high frequency noise. These transformed
intensities were used to identify bright pixels at the center of clusters with a user-defined intensity

cutoff. From each of these center points, neighboring pixels were compared to a threshold
intensity that was set by the local background of the cell and scaled by a second user-defined
parameter. Neighboring pixels above this threshold are included in the cluster, while pixels below
the threshold are excluded. This occurs iteratively, where adjacent pixels to each included pixel
were also checked to be included or excluded until all new neighbors were below the threshold.
Cluster properties of each cell were exported and processed in R for analysis and visualization.
837
*Live cell imaging of ErkKTR*
ErkKTR dynamics and cell death were tracked and quantitated in a semi-automated
manner using the p53CellCinema package (Reyes et al., 2018) in MATLAB, and data was
processed and visualized in R and RStudio using the tidyR package. Briefly, cells with low or no
ErkKTR expression were first excluded from analysis. Then, ErkKTR cytoplasmic/nuclear ratios
from remaining cells were imported into MATLAB, and peaks were identified using the
'findpeaks' function. Peak calls were then manually inspected for outliers, for example resulting
from apoptotic cell debris, and outlier cells were removed from analysis. Of note, we consistently
observed Erk pulses preceding cell division events across conditions. This phenomenon has
been previously observed and has been shown to be independent of Ras/Erk signaling,
potentially due to non-specific activation from a cyclin dependent kinase(Gerosa et al., 2020).
Thus, we disregarded all pulses that occurred within 10 frames (50 mins) preceding cell division.
To count Erk pulses in neighbors vs non-neighbors of dying cells, we first identified each cell
death event. We then identified the neighbors of the dying cell by identifying which nuclear
centroids were within a 100 pixel radius (~32 μm, or 2-3 cell diameters) of the dying cell for each
of the 10 frames preceding the death event. We then counted the total pulses of those neighbors
over the indicated time frame, and compared against pulse counts from a random subset of non-
neighbor cells over the same time frame. Importantly, for each death event, the number of non-
neighbors matched the number of neighbors, and the sampled non-neighbors were not
neighbors of other death events over that same time span. For visualization in **Figure 6E**, cells
that divided during the course of the experiment are represented as separate cells, and the Erk
history of the mother cell is reproduced for both cells. However, for quantification of Erk pulses,
mother cell pulses were counted only once.
861

**References**

Aikin, T.J., Peterson, A.F., Pokrass, M.J., Clark, H.R., and Regot, S. (2020). MAPK activity dynamics regulate non-cell autonomous effects of oncogene expression. Elife *9*.

Banani, S.F., Lee, H.O., Hyman, A.A., and Rosen, M.K. (2017). Biomolecular condensates: organizers of cellular biochemistry. Nat. Rev. Mol. Cell Biol. *18*, 285–298.

Benman, W., Berlew, E.E., Deng, H., Parker, C., Kuznetsov, I.A., Lim, B., Siekmann, A.F., Chow, B.Y., and Bugaj, L.J. (2022). Temperature-responsive optogenetic probes of cell signaling. Nat. Chem. Biol. *18*, 152–160.

Berg, S., Kutra, D., Kroeger, T., Straehle, C.N., Kausler, B.X., Haubold, C., Schiegg, M., Ales, J., Beier, T., Rudy, M., et al. (2019). ilastik: interactive machine learning for (bio)image analysis. Nat. Methods *16*, 1226–1232.

Brachmann, R., Lindquist, P.B., Nagashima, M., Kohr, W., Lipari, T., Napier, M., and Derynck, R. (1989). Transmembrane TGF-alpha precursors activate EGF/TGF-alpha receptors. Cell *56*, 691–700.

Bugaj, L.J., and Lim, W.A. (2019). High-throughput multicolor optogenetics in microwell plates. Nat. Protoc. *14*, 2205–2228.

Bugaj, L.J., Choksi, A.T., Mesuda, C.K., Kane, R.S., and Schaffer, D.V. (2013). Optogenetic protein clustering and signaling activation in mammalian cells. Nat. Methods *10*, 249–252.

Bugaj, L.J., O'Donoghue, G.P., and Lim, W.A. (2017). Interrogating cellular perception and decision making with optogenetic tools. J. Cell Biol. *216*, 25–28.

Bugaj, L.J., Sabnis, A.J., Mitchell, A., Garbarino, J.E., Toettcher, J.E., Bivona, T.G., and Lim, W.A. (2018). Cancer mutations and targeted drugs can disrupt dynamic signal encoding by the Ras-Erk pathway. Science *361*.

Cai, D., Liu, Z., and Lippincott-Schwartz, J. (2021). Biomolecular Condensates and Their Links to Cancer Progression. Trends Biochem. Sci. *46*, 535–549.

Corcoran, R.B., Ebi, H., Turke, A.B., Coffee, E.M., Nishino, M., Cogdill, A.P., Brown, R.D., Della Pelle, P., Dias-Santagata, D., Hung, K.E., et al. (2012). EGFR-mediated re-activation of MAPK signaling contributes to insensitivity of BRAF mutant colorectal cancers to RAF inhibition with vemurafenib. Cancer Discov. *2*, 227–235.

Davies, A.E., Pargett, M., Siebert, S., Gillies, T.E., Choi, Y., Tobin, S.J., Ram, A.R., Murthy, V., Juliano, C., Quon, G., et al. (2020). Systems-Level Properties of EGFR-RAS-ERK Signaling Amplify Local Signals to Generate Dynamic Gene Expression Heterogeneity. Cell Syst *11*, 161-175.e5.

Davies, K.D., Mahale, S., Astling, D.P., Aisner, D.L., Le, A.T., Hinz, T.K., Vaishnavi, A., Bunn, P.A., Jr, Heasley, L.E., Tan, A.-C., et al. (2013). Resistance to ROS1 inhibition mediated by EGFR pathway activation in non-small cell lung cancer. PLoS One *8*, e82236.

Dong, J., Opresko, L.K., Dempsey, P.J., Lauffenburger, D.A., Coffey, R.J., and Wiley, H.S.
(1999). Metalloprotease-mediated ligand release regulates autocrine signaling through the
epidermal growth factor receptor. Proc. Natl. Acad. Sci. U. S. A. *96*, 6235–6240.

Du, Z., and Lovly, C.M. (2018). Mechanisms of receptor tyrosine kinase activation in cancer.
Mol. Cancer *17*, 58.

Gagliardi, P.A., Dobrzyński, M., Jacques, M.-A., Dessauges, C., Ender, P., Blum, Y., Hughes,
R.M., Cohen, A.R., and Pertz, O. (2021). Collective ERK/Akt activity waves orchestrate epithelial
homeostasis by driving apoptosis-induced survival. Dev. Cell *56*, 1712-1726.e6.

Gainor, J.F., Dardaei, L., Yoda, S., Friboulet, L., Leshchiner, I., Katayama, R., Dagogo-Jack, I.,
Gadgeel, S., Schultz, K., Singh, M., et al. (2016). Molecular Mechanisms of Resistance to First-
and Second-Generation ALK Inhibitors in ALK-Rearranged Lung Cancer. Cancer Discov. *6*,
1118–1133.

Garabedian, M.V., Wang, W., Dabdoub, J.B., Tong, M., Caldwell, R.M., Benman, W., Schuster,
B.S., Deiters, A., and Good, M.C. (2021). Designer membraneless organelles sequester native
factors for control of cell behavior. Nat. Chem. Biol. *17*, 998–1007.

Gerosa, L., Chidley, C., Fröhlich, F., Sanchez, G., Lim, S.K., Muhlich, J., Chen, J.-Y.,
Vallabhaneni, S., Baker, G.J., Schapiro, D., et al. (2020). Receptor-Driven ERK Pulses
Reconfigure MAPK Signaling and Enable Persistence of Drug-Adapted BRAF-Mutant
Melanoma Cells. Cell Syst *11*, 478-494.e9.

Guntas, G., Hallett, R.A., Zimmerman, S.P., Williams, T., Yumerefendi, H., Bear, J.E., and
Kuhlman, B. (2015). Engineering an improved light-induced dimer (iLID) for controlling the
localization and activity of signaling proteins. Proc. Natl. Acad. Sci. U. S. A. *112*, 112–117.

Hirai, N., Sasaki, T., Okumura, S., Minami, Y., Chiba, S., and Ohsaki, Y. (2020). Monomerization
of ALK Fusion Proteins as a Therapeutic Strategy in ALK-Rearranged Non-small Cell Lung
Cancers. Front. Oncol. *10*, 419.

Hrustanovic, G., Olivas, V., Pazarentzos, E., Tulpule, A., Asthana, S., Blakely, C.M., Okimoto,
R.A., Lin, L., Neel, D.S., Sabnis, A., et al. (2015). RAS-MAPK dependence underlies a rational
polytherapy strategy in EML4-ALK–positive lung cancer. Nat. Med. *21*, 1038.

Jiang, S., Fagman, J.B., Chen, C., Alberti, S., and Liu, B. (2020). Protein phase separation and
its role in tumorigenesis. Elife *9*.

Kim, N., Kim, J.M., Lee, M., Kim, C.Y., Chang, K.-Y., and Heo, W.D. (2014). Spatiotemporal
control of fibroblast growth factor receptor signals by blue light. Chem. Biol. *21*, 903–912.

Kurtova, A.V., Xiao, J., Mo, Q., Pazhanisamy, S., Krasnow, R., Lerner, S.P., Chen, F., Roh, T.T.,
Lay, E., Ho, P.L., et al. (2015). Blocking PGE2-induced tumour repopulation abrogates bladder
cancer chemoresistance. Nature *517*, 209–213.

Lamprecht, M.R., Sabatini, D.M., and Carpenter, A.E. (2007). CellProfiler$^{TM}$: free, versatile
software for automated biological image analysis. Biotechniques *42*, 71–75.

Lito, P., Pratilas, C.A., Joseph, E.W., Tadi, M., Halilovic, E., Zubrowski, M., Huang, A., Wong,
W.L., Callahan, M.K., Merghoub, T., et al. (2012). Relief of profound feedback inhibition of
mitogenic signaling by RAF inhibitors attenuates their activity in BRAFV600E melanomas.
Cancer Cell *22*, 668–682.

Lovly, C.M., McDonald, N.T., Chen, H., Ortiz-Cuaran, S., Heukamp, L.C., Yan, Y., Florin, A.,
Ozretić, L., Lim, D., Wang, L., et al. (2014). Rationale for co-targeting IGF-1R and ALK in ALK
fusion-positive lung cancer. Nat. Med. *20*, 1027–1034.

Miller, M.A., Oudin, M.J., Sullivan, R.J., Wang, S.J., Meyer, A.S., Im, H., Frederick, D.T., Tadros,
J., Griffith, L.G., Lee, H., et al. (2016). Reduced Proteolytic Shedding of Receptor Tyrosine
Kinases Is a Post-Translational Mechanism of Kinase Inhibitor Resistance. Cancer Discov. *6*,
382–399.

Obenauf, A.C., Zou, Y., Ji, A.L., Vanharanta, S., Shu, W., Shi, H., Kong, X., Bosenberg, M.C.,
Wiesner, T., Rosen, N., et al. (2015). Therapy-induced tumour secretomes promote resistance
and tumour progression. Nature *520*, 368–372.

Omer, A., Patel, D., Lian, X.J., Sadek, J., Di Marco, S., Pause, A., Gorospe, M., and Gallouzi,
I.E. (2018). Stress granules counteract senescence by sequestration of PAI-1. EMBO Rep. *19*.

Ou, S.-H.I., Govindan, R., Eaton, K.D., Otterson, G.A., Gutierrez, M.E., Mita, A.C., Argiris, A.,
Brega, N.M., Usari, T., Tan, W., et al. (2017). Phase I Results from a Study of Crizotinib in
Combination with Erlotinib in Patients with Advanced Nonsquamous Non-Small Cell Lung
Cancer. J. Thorac. Oncol. *12*, 145–151.

Peschon, J.J., Slack, J.L., Reddy, P., Stocking, K.L., Sunnarborg, S.W., Lee, D.C., Russell,
W.E., Castner, B.J., Johnson, R.S., Fitzner, J.N., et al. (1998). An essential role for ectodomain
shedding in mammalian development. Science *282*, 1281–1284.

Prahallad, A., Sun, C., Huang, S., Di Nicolantonio, F., Salazar, R., Zecchin, D., Beijersbergen,
R.L., Bardelli, A., and Bernards, R. (2012). Unresponsiveness of colon cancer to BRAF(V600E)
inhibition through feedback activation of EGFR. Nature *483*, 100–103.

Regot, S., Hughey, J.J., Bajar, B.T., Carrasco, S., and Covert, M.W. (2014). High-Sensitivity
Measurements of Multiple Kinase Activities in Live Single Cells. Cell *157*, 1724–1734.

Reyes, J., Chen, J.-Y., Stewart-Ornstein, J., Karhohs, K.W., Mock, C.S., and Lahav, G. (2018).
Fluctuations in p53 Signaling Allow Escape from Cell-Cycle Arrest. Mol. Cell *71*, 581-591.e5.

Rusch, V., Baselga, J., Cordon-Cardo, C., Orazem, J., Zaman, M., Hoda, S., McIntosh, J., Kurie,
J., and Dmitrovsky, E. (1993). Differential expression of the epidermal growth factor receptor
and its ligands in primary non-small cell lung cancers and adjacent benign lung. Cancer Res.
*53*, 2379–2385.

Sahin, U., Weskamp, G., Kelly, K., Zhou, H.-M., Higashiyama, S., Peschon, J., Hartmann, D.,
Saftig, P., and Blobel, C.P. (2004). Distinct roles for ADAM10 and ADAM17 in ectodomain
shedding of six EGFR ligands. J. Cell Biol. *164*, 769–779.

Sampson, J., Richards, M.W., Choi, J., Fry, A.M., and Bayliss, R. (2021). Phase-separated foci
of EML4-ALK facilitate signalling and depend upon an active kinase conformation. EMBO Rep.
e53693.

Sasaki, T., Koivunen, J., Ogino, A., Yanagita, M., Nikiforow, S., Zheng, W., Lathan, C., Marcoux,
J.P., Du, J., Okuda, K., et al. (2011). A novel ALK secondary mutation and EGFR signaling cause
resistance to ALK kinase inhibitors. Cancer Res. *71*, 6051–6060.

Shaw, A.T., Hsu, P.P., Awad, M.M., and Engelman, J.A. (2013). Tyrosine kinase gene
rearrangements in epithelial malignancies. Nat. Rev. Cancer *13*, 772–787.

Siegel, R.L., Miller, K.D., Fuchs, H.E., and Jemal, A. (2021). Cancer Statistics, 2021. CA Cancer
J. Clin. *71*, 7–33.

Solomon, B.J., Besse, B., Bauer, T.M., Felip, E., Soo, R.A., Camidge, D.R., Chiari, R., Bearz,
A., Lin, C.-C., Gadgeel, S.M., et al. (2018). Lorlatinib in patients with ALK-positive non-small-cell
lung cancer: results from a global phase 2 study. Lancet Oncol. *19*, 1654–1667.

Straussman, R., Morikawa, T., Shee, K., Barzily-Rokni, M., Qian, Z.R., Du, J., Davis, A.,
Mongare, M.M., Gould, J., Frederick, D.T., et al. (2012). Tumour micro-environment elicits innate
resistance to RAF inhibitors through HGF secretion. Nature *487*, 500–504.

Sukhatme, V.P., Kartha, S., Toback, F.G., Taub, R., Hoover, R.G., and Tsai-Morris, C.H. (1987).
A novel early growth response gene rapidly induced by fibroblast, epithelial cell and lymphocyte
mitogens. Oncogene Res. *1*, 343–355.

Takeuchi, K., Soda, M., Togashi, Y., Suzuki, R., Sakata, S., Hatano, S., Asaka, R., Hamanaka,
W., Ninomiya, H., Uehara, H., et al. (2012). RET, ROS1 and ALK fusions in lung cancer. Nat.
Med. *18*, 378–381.

Tani, T., Yasuda, H., Hamamoto, J., Kuroda, A., Arai, D., Ishioka, K., Ohgino, K., Miyawaki, M.,
Kawada, I., Naoki, K., et al. (2016). Activation of EGFR Bypass Signaling by TGFα
Overexpression Induces Acquired Resistance to Alectinib in ALK-Translocated Lung Cancer
Cells. Mol. Cancer Ther. *15*, 162–171.

Tanimoto, A., Yamada, T., Nanjo, S., Takeuchi, S., Ebi, H., Kita, K., Matsumoto, K., and Yano,
S. (2014). Receptor ligand-triggered resistance to alectinib and its circumvention by Hsp90
inhibition in EML4-ALK lung cancer cells. Oncotarget *5*, 4920–4928.

Thedieck, K., Holzwarth, B., Prentzell, M.T., Boehlke, C., Kläsener, K., Ruf, S., Sonntag, A.G.,
Maerz, L., Grellscheid, S.-N., Kremmer, E., et al. (2013). Inhibition of mTORC1 by astrin and
stress granules prevents apoptosis in cancer cells. Cell *154*, 859–874.

Toettcher, J.E., Weiner, O.D., and Lim, W.A. (2013). Using optogenetics to interrogate the
dynamic control of signal transmission by the Ras/Erk module. Cell *155*, 1422–1434.

Tulpule, A., Guan, J., Neel, D.S., Allegakoen, H.R., Lin, Y.P., Brown, D., Chou, Y.-T., Heslin, A.,
Chatterjee, N., Perati, S., et al. (2021). Kinase-mediated RAS signaling via membraneless
cytoplasmic protein granules. Cell.

Turke, A.B., Song, Y., Costa, C., Cook, R., Arteaga, C.L., Asara, J.M., and Engelman, J.A. (2012). MEK inhibition leads to PI3K/AKT activation by relieving a negative feedback on ERBB receptors. Cancer Res. *72*, 3228–3237.

Vaishnavi, A., Schubert, L., Rix, U., Marek, L.A., Le, A.T., Keysar, S.B., Glogowska, M.J., Smith, M.A., Kako, S., Sumi, N.J., et al. (2017). EGFR Mediates Responses to Small-Molecule Drugs Targeting Oncogenic Fusion Kinases. Cancer Res. *77*, 3551–3563.

Valon, L., Davidović, A., Levillayer, F., Villars, A., Chouly, M., Cerqueira-Campos, F., and Levayer, R. (2021). Robustness of epithelial sealing is an emerging property of local ERK feedback driven by cell elimination. Dev. Cell *56*, 1700-1711.e8.

Vander Velde, R., Yoon, N., Marusyk, V., Durmaz, A., Dhawan, A., Miroshnychenko, D., Lozano-Peral, D., Desai, B., Balynska, O., Poleszhuk, J., et al. (2020). Resistance to targeted therapies as a multifactorial, gradual adaptation to inhibitor specific selective pressures. Nat. Commun. *11*, 2393.

Voena, C., Di Giacomo, F., Panizza, E., D'Amico, L., Boccalatte, F.E., Pellegrino, E., Todaro, M., Recupero, D., Tabbò, F., Ambrogio, C., et al. (2013). The EGFR family members sustain the neoplastic phenotype of ALK+ lung adenocarcinoma via EGR1. Oncogenesis *2*, e43.

Weinstein, I.B. (2002). Cancer. Addiction to oncogenes--the Achilles heal of cancer. Science *297*, 63–64.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., et al. (2019). Welcome to the tidyverse. J. Open Source Softw. *4*, 1686.

Wilson, M.Z., Ravindran, P.T., Lim, W.A., and Toettcher, J.E. (2017). Tracing Information Flow from Erk to Target Gene Induction Reveals Mechanisms of Dynamic and Combinatorial Control. Mol. Cell *67*, 757-769.e5.

Wilson, T.R., Fridlyand, J., Yan, Y., Penuel, E., Burton, L., Chan, E., Peng, J., Lin, E., Wang, Y., Sosman, J., et al. (2012). Widespread potential for growth-factor-driven resistance to anticancer kinase inhibitors. Nature *487*, 505–509.

Wong, S.T., Winchell, L.F., McCune, B.K., Earp, H.S., Teixidó, J., Massagué, J., Herman, B., and Lee, D.C. (1989). The TGF-alpha precursor expressed on the cell surface binds to the EGF receptor on adjacent cells, leading to signal transduction. Cell *56*, 495–506.