Title: Oncogenic protein condensates modulate cell signal perception and drug tolerance.

Authors: David Gonzalez-Martinez[1#], Lee Roth[1#], Thomas R. Mumford[1], Asmin Tulpule[2], Trever G.
Bivona[3], Lukasz J. Bugaj[1,4,5,*]

Affiliations:
[1]Department of Bioengineering, University of Pennsylvania, Philadelphia, PA, 19104, USA
[2]Division of Pediatric Hematology/Oncology, UCSF, San Francisco, CA 94143, USA
[3]Department of Medicine, Division of Hematology and Oncology, UCSF, San Francisco, CA 94143,
USA
[4]Abramson Cancer Center, University of Pennsylvania, Philadelphia, PA, 19104, USA
[5]Institute of Regenerative Medicine, University of Pennsylvania, Philadelphia, PA, 19104, USA

[#]equal contributions
[*]corresponding author

Correspondence: bugaj@seas.upenn.edu

Abstract

Drug resistance remains a central challenge towards durable cancer therapy, including for cancers
driven by the EML4-ALK oncogene. EML4-ALK and related fusion oncogenes form cytoplasmic protein
condensates that transmit oncogenic signals through the Ras/Erk pathway. However, whether such
condensates play a role in drug response or resistance development is unclear. Here, we applied
optogenetic functional profiling to examine how EML4-ALK condensates impact signal transmission
through transmembrane receptor tyrosine kinases (RTKs), a common route of resistance signaling. We
found that condensates dramatically suppress signaling through activated RTKs including EGFR.
Conversely, ALK inhibition restored and hypersensitized RTK signals. Modulation of RTK sensitivity
occurred because EML4-ALK condensates sequestered downstream adapters that are required to
transduce signals from both EML4-ALK and ligand-stimulated RTKs. Strikingly, EGFR
hypersensitization resulted in rapid and pulsatile Erk signal reactivation within 10s of minutes of drug
addition. EGFR reactivation originated from paracrine signals from neighboring apoptotic cells, and
reactivation could be blocked by inhibition of either EGFR or matrix metalloproteases. Paracrine signals
promoted survival during ALK inhibition, and blockade of paracrine signals accelerated cell killing and
suppressed drug tolerance. Our results uncover a regulatory role for protein condensates in cancer
signaling and drug response and demonstrate the potential of optogenetic profiling for drug discovery
based on functional biomarkers in cancer cells.

Introduction
Despite the development of potent oncogene inhibitors, drug resistance remains an unsolved
challenge towards durable cancer therapies. Thus, there remains a critical need to better understand
molecular interactions between host cells, oncogenes, and targeted inhibitors to identify effective
treatment strategies that forestall or prevent resistance.
The oncoprotein EML4-ALK (echinoderm microtubule-associated protein-like 4-anaplastic
lymphoma kinase) is a receptor tyrosine kinase (RTK) fusion oncogene that drives ~3-7% of non-small
cell lung cancer (NSCLC), the leading cause of cancer-related death in the US (Takeuchi et al. 2012;

Siegel et al. 2021). RTK fusions are a large class of chimeric oncogenes that share a similar
composition, where the enzymatic fragment of an RTK is fused through chromosomal rearrangement to
a fusion partner domain that is often multimeric (Shaw et al. 2013; Du and Lovly 2018). Cancers driven
by RTK fusions commonly exhibit oncogene addiction, where blockade of the oncogene causes cell
death(Weinstein 2002). For EML4-ALK+ cancers, multiple FDA-approved ALK inhibitors achieve initial
patient responses and tumor regression, although resistance and tumor relapse inevitably emerge
(Gainor et al. 2016; Solomon et al. 2018).
EML4-ALK drives oncogenicity primarily through sustained Ras/Erk signaling (Hrustanovic et al.
2015). Durable treatment of cancers driven by Ras/Erk has been challenged by robust autoinhibitory
feedback loops that result in pathway reactivation after treatment. For example, in the case of BRAF
V600E+ cancers, BRAF inhibition leads to relief of negative feedback loops that permit strong EGFR
stimulation, cell survival, and drug resistance (Lito et al. 2012a; Prahallad et al. 2012a; Gerosa et al.
2020a; Corcoran et al. 2012). For EML4-ALK+ cancers, re-activation of RTK signals after ALK inhibition
also promotes drug tolerance and acquired resistance (Vaishnavi et al. 2017; Sasaki et al. 2011;
Hrustanovic et al. 2015; T. R. Wilson et al. 2012; Voena et al. 2013; Tani et al. 2016) (**Figure 1A**),
although specific feedback mechanisms that mediate this reactivation are not well understood.
Recently it was found that EML4-ALK forms large cytoplasmic protein granules—or
condensates— in cancer cells(Hrustanovic et al. 2015; Tulpule et al. 2021) (**Figure 1B**). Condensate
formation resulted in part from the ability of RTK adapter proteins like Grb2 to enlarge small EML4-ALK
oligomers through multivalent interactions. Such higher-order assemblies of proteins may have
numerous and even opposing consequences on intracellular biochemistry, for example acting as
amplifiers or inhibitors of molecular reactions(Banani et al. 2017). Although condensate formation
serves as an amplifier that promotes EML4-ALK activation and signaling, it is unclear whether
condensates might have additional impact on a cell's ability to transmit signals. Of particular interest are
functional interactions between EML4-ALK condensates and transmembrane RTKs including the
epidermal growth factor receptor (EGFR), which signal through the same downstream effectors as
EML4-ALK (**Figure 1C**). These interactions are potentially important because EGFR is overexpressed
in over 80% of NSCLC and is correlated with poor prognosis (Rusch et al. 1993) and drug resistance
(Vaishnavi et al. 2017; Tanimoto et al. 2014; T. R. Wilson et al. 2012; Davies et al. 2013; Obenauf et al.
2015; Straussman et al. 2012; Lito et al. 2012a).
We recently developed a method called 'functional profiling' to detect how oncogene expression
alters cell signal transmission (L. J. Bugaj et al. 2018; Lukasz J. Bugaj, O'Donoghue, and Lim 2017). In
this approach, we apply precisely defined signaling stimuli, for example using light-activated
'optogenetic' signaling probes, and we quantify differential responses in downstream signals,
transcription, and cell fate as a function of oncogene expression or drug treatment. Previously,
functional profiling revealed that certain BRAF-mutant cancer cells and drug-treated cells exhibited
abnormally slow Ras/Erk pathway activation kinetics, and that such kinetics can cause cells to
misinterpret dynamic stimuli leading to hyperproliferation(L. J. Bugaj et al. 2018).
In the present work, we applied optogenetic profiling to uncover functional relationships between
EML4-ALK condensates and transmembrane RTKs (**Figure 1D**). We discovered that EML4-ALK
dramatically suppressed RTK signal transmission. We found that suppression resulted not from
biochemical feedback, but rather from sequestration of RTK adapters by EML4-ALK condensates.
Suppression was rapidly reversed by ALK inhibition, resulting in hypersensitization of RTKs to external
growth factors. Restored perception of external ligands promoted rapid EGFR/Erk reactivation after

drug treatment, which suppressed cell killing and permitted tolerance to ALK inhibitors. Our work thus
uncovers a signal regulatory role for oncogenic condensates and suggests novel candidate co-targets
to enhance the durability of ALK inhibitor therapy.
93
Results
Optogenetic profiling reveals that RTK signaling is suppressed in EML4-ALK+ cancer cells.
To determine whether EML4-ALK could alter RTK signaling properties, we expressed a light-sensitive
fibroblast growth factor receptor (optoFGFR (Kim et al. 2014)) in the STE-1 cancer cell line, which is
driven by EML4-ALK(V1) (Lovly et al. 2014) (**Figure 1E, S1A**). We observed signal transmission by
applying blue light stimuli and measuring single-cell phospho-Erk (ppErk) levels through
immunofluorescence. We found that blue light stimulation of optoFGFR could induce only minimal
ppErk signal increase (1.4-fold) above the high basal ppErk levels attributed to active EML4-ALK
(**Figure 1F, top)**. We then asked whether signaling responses might change in the presence of ALK
inhibition. Pre-treatment with the ALK inhibitor crizotinib (ALKi, 1 µM) eliminated the high tonic ppErk
signal observed in untreated cells, consistent with the ability of EML4-ALK to drive strong signals
through the Ras/Erk pathway. Strikingly, subsequent light stimulation now drove strong Erk signaling,
surpassing levels achieved in the absence of drug (**Fig. 1F, bottom)**. The dynamic range (fold-change)
of signal induction increased as a function of light intensity and reached a maximum of 29-fold increase
at the highest levels of light stimulation (**Figure 1G**). Notably, ppErk increase in ALKi-treated cells could
be measured after stimulation with exceedingly low levels of light (8 mW/cm^2), a level that did not
provide measurable increase in untreated cells, suggesting that ALK inhibition can hypersensitize
cancer cells to weak RTK signals (**Figure S1B**). We note that although the magnitude of dynamic range
increase was dependent on the expression levels of optoFGFR, the general trends described held
across all expression levels (**Figure S1C**). These results suggested the existence of a strong negative
regulatory interaction between EML4-ALK activity and transmembrane RTK signaling.
115
EML4-ALK suppresses — and ALK inhibition restores — EGFR signaling
To determine whether negative regulation of RTKs could be observed in the context of native
epidermal growth factor receptor (EGFR), we measured ppErk induction upon addition of epidermal
growth factor (EGF) in STE-1 and H3122 cells, two EML4-ALK+ cell lines (Lovly et al. 2014) (**Figure
2A)**. Strong (100 ng/mL) EGF stimulation gave minimal ppErk response, but pre-treatment with ALKi
resulted in notably increased ppErk dynamic range in both cell lines (1.7- vs 9.5-fold in STE-; 1.7- vs
7.4-fold in H3122) (**Figure 2B,C**). As before, increased dynamic range was due to lower baseline — but
also higher maximal — ppErk signaling, and the magnitude of dynamic range increase was dose-
dependent (**Figure 2C, S2A**). Thus, we concluded that EML4-ALK activity suppresses EGFR signaling
in cancer cells.
We next asked whether expression of EML4-ALK was sufficient to suppress RTK signaling. We
transiently expressed EML4-ALK(V1) in non-transformed lung epithelial Beas2B cells and we observed
responses to EGF (**Figure 2D**). As expected, EML4-ALK expression raised basal ppErk levels (t = 0,
Figure 2E). However, upon addition of EGF, ppErk signaling was weaker (stronger ppErk suppression)
than in untransfected cells (**Figure 2E**). Finally, pre-incubation with ALKi reversed suppression of
EGFR/Erk signals in transfected cells (**Figure 2E,F**). Collectively, our results show that EML4-ALK
expression is sufficient to strongly dampen signaling through transmembrane EGFR, and that ALK
inhibition restores and potentiates these signals.

Mapping EGFR suppression using optogenetics*.* We sought to understand the molecular
mechanism by which EML4-ALK suppressed EGFR signaling. To narrow candidate mechanisms, we
first determined the duration of ALKi pre-incubation that was required to observe enhanced EGFR/Erk
signaling. We preincubated both STE-1 and H3122 cells with 0-60 min of ALK inhibitor, and we
analyzed ppErk levels in response to 15 min of EGF stimulation (**Figure 3A**). In both cell lines, an
increase in Erk signal amplitude was observed with as little as 5 min of ALKi pre-incubation (20 minutes
total including stimulation), and rose to half-max with only ~15 mins of pre-incubation (**Figure 3B**).
Such fast response suggested a primarily post-translational mechanism.
To locate the node in the EGFR/Erk pathway that was suppressed by EML4-ALK, we stimulated
the pathway at successive nodes using optogenetics in the presence and absence of drug (**Figure
3C,D**). In addition to optoFGFR, we also generated STE-1 cells that expressed optoSOS, which allows
activation of Ras/Erk signals through light-induced membrane recruitment of the SOS catalytic
domain(Toettcher, Weiner, and Lim 2013; Benman et al. 2022) (**Figure 3D, S2**). As before, light
stimulation of optoFGFR yielded little increase in ppErk, but ppErk induction was enhanced with pre-
treatment with ALKi (**Figure 3E, left**), consistent with relief of optoFGFR suppression by ALKi. By
contrast, the maximal level of ppErk induced by optoSOS was unchanged by ALKi pre-treatment
(**Figure 3E, right**), suggesting that EML4-ALK suppresses signaling upstream of Ras, for example at
the receptor level. However, direct immunofluorescence showed that EGF-stimulated phospho-EGFR
(pEGFR) levels were unchanged by ALKi pre-incubation (**Figure S3**). Collectively, our results indicate
that EML4-ALK suppresses RTK signaling downstream of receptor phosphorylation but upstream of
Ras activation, implicating a role for the adapter proteins that couple these two nodes (**Figure 3F**).
Oncogene-mediated feedback suppression of RTKs has been described previously, most
notably as as a result of Erk-dependent transcriptional and post-translational negative
feedback(Prahallad et al. 2012b; Corcoran et al. 2012; Lito et al. 2012b; Gerosa et al. 2020b; Turke et
al. 2012). To test the role of Erk-dependent negative feedback in EML4-ALK+ cancer cells, we
decoupled ALK inhibition from the loss of Erk signaling using optoSOS stimulation (**Figure 3G**). In this
experiment, we again assessed EGF response in STE-1 cells with ALKi preincubation, but we
supplemented one experimental group with optogenetic levels of Erk signaling that matched those from
drug-naïve STE-1 cells, thus maintaining any Erk-dependent negative feedback that would have been
lost through ALK inhibition (**Figure 3 H,I, S4**). Our results showed that ALKi preincubation again
increased the strength of EGF induction, and optoSOS stimulation during pre-incubation did not
diminish the strength of induction. These results show that in EML4-ALK+ cancer cells, EGFR
suppression —and drug-induced relief of suppression — is not mediated by established, Erk-dependent
mechanisms (**Figure 3J**).
169

EML4-ALK condensates suppress EGFR through competitive sequestration of RTK effectors.
We suspected that condensation of EML4-ALK could play a causal role in RTK suppression. To test the
effects of such higher-order EML4-ALK organization, we examined EGF response in cells transfected
with EML4-ALK mutants that fail to form condensates due to mutations in either the trimerization
domain (ΔTD) or the kinase domain (K589M)(Tulpule et al. 2021). In contrast to full-length EML4-ALK
(V1) (**Figure 4A, left**), the condensate-deficient mutants had minimal effect EGF-induced Erk activity
(**Figure 4A, middle and right**), suggesting that condensation was essential for its ability to modulate
EGFR sensitivity.

Protein condensation can act as a negative regulator in both natural and engineered systems
when it sequesters essential components of biochemical reactions (Thedieck et al. 2013; Omer et al.
2018; Garabedian et al. 2021). We thus hypothesized that EML4-ALK condensates suppressed
transmembrane RTKs by sequestering the adapter proteins that are required to transmit RTK signals.
Such adapters, including Grb2 and SOS1 which link RTKs to Ras/Erk signaling, are also required to
transmit EML4-ALK signals (Tulpule et al. 2021), and thus represent shared resources that could
implement competitive inhibition of RTKs.
To test this hypothesis, we observed the localization of the Grb2 adapter using Beas2B cells where
Grb2 was fluorescently tagged at the endogenous locus (Tulpule et al. 2021). In normal cells, Grb2
appeared diffuse in the cytoplasm but translocated to activated EGFR within <1 min of EGF addition
(**Fig. 4D,E, Supplementary Movie 1**). However, upon transfection with EML4-ALK, Grb2 strongly
colocalized with EML4-ALK puncta (**Figure S5**), and Grb2 remained sequestered in the puncta even
upon addition of EGF (**Figure 4D,E,F, Supplementary Movie 1**). We then asked how Grb2 localization
might change in the presence of ALKi (**Figure 4G**). ALKi caused Grb2 puncta to rapidly dissolve into
the cytoplasm ($t_{1/2}$ = 17 +/- 2 min) (**Figure 4H,I, Supplementary Movie 2**), in line with recent reports
(Sampson et al. 2021). Finally, sequential treatment with ALKi and EGF allowed robust membrane
translocation of Grb2 even in the presence of EML4-ALK (**Figure 4J,K**). These results suggest that
EML4-ALK condensates sequester Grb2 and suppress translocation to the membrane upon EGF
stimulation, providing a mechanism by which EML4-ALK can competitively inhibit EGFR signaling.
As a further test of our model, we measured the EGF-induced recruitment of SOS1 in H3122
cancer cells through co-immunoprecipitation. SOS1 is recruited by Grb2 to activated receptors and
stimulates Ras upon recruitment. We found that SOS1 co-precipitated with EGFR upon EGF
stimulation, but only when preincubated with ALKi, further suggesting that active EML4-ALK
condensates sequester RTK adapters (**Figure 4J, S6**). Finally, our conceptual model predicts that
distinct RTK fusions that form condensates would similarly repress EGFR signaling. We thus measured
EGF response in TPC-1 cells, which harbor the CCDC6-RET fusion. Like EML4-ALK, CCDC6-RET
forms condensates that colocalize with many RTK adapters, including Grb2 and SOS1(Tulpule et al.
2021). As predicted, EGF stimulation induced only moderate levels of Erk phosphorylation, whereas
pre-treatment with RET inhibitor BLU-667 permitted strong stimulation (2.2- vs 7-fold stimulation) to
levels beyond those achievable in drug-naive cells (**Figure S7**). Collectively, our results suggest that
competition for RTK adapters is a key mechanism by which condensates of EML4-ALK —and
potentially other RTK fusions — suppress a cell's perception of transmembrane RTK signals (**Fig. 4K**).
210
RTK resensitization promotes rapid signal reactivation upon ALK inhibition.
RTK stimulation promotes drug tolerance and acquired resistance across cancer types (Straussman et
al. 2012; T. R. Wilson et al. 2012). We thus sought to understand whether hypersensitization of RTKs
could promote RTK signaling during ALKi therapy. We monitored RTK/Erk signaling in populations of
STE-1 cells using ErkKTR, a biosensor that reports on Erk activity through nuclear exclusion of a
fluorescent protein (Regot et al. 2014). (**Figure 5A**). In the absence of ALKi, Erk activity was at an
intermediate level, indicated by relatively equal distribution between the cytoplasm and nucleus (**Figure
5B, Supplementary Movie 3**). Notably, this localization pattern did not change over the course of 22
hrs, reflecting the tonic signaling downstream of EML4-ALK. By contrast, treatment with ALKi induced a
rapid initial decrease of Erk activity followed by a striking appearance of Erk activity pulses within as
little as 30 min after treatment (**Supplementary Movie 3**). Each Erk pulse lasted 10-20 minutes, and

the pulse amplitude exceeded levels observed in the absence of drug, paralleling our earlier results
with optogenetic or ligand stimulation (**Figure 5C**).
Activity pulses appeared sporadically but were not confined to individual cells. Rather, Erk
activity appeared in small clusters of neighboring cells either simultaneously or as a wave of activity
(**Figure 4B, Supplementary Movie 3**). This pattern was consistent with RTK stimulation through
paracrine signaling. To test this hypothesis, we sought to block paracrine signals through inhibition of
either the EGF receptor (EGFRi, 1 µM erlotinib) or matrix metalloproteases (MMPi, 10 µM marimastat),
which release EGFR ligands from the cell surface to enable paracrine signaling (Dong et al. 1999;
Peschon et al. 1998; Sahin et al. 2004). Co-treatment with ALKi and EGFRi suppressed Erk activity
pulses after drug addition (ALKi vs ALKi/EGFRi: 58 $\pm$ 7% vs 12 $\pm$ 5% of cells showing $\geq$ 1 pulse over 22
hrs; error = 95% CI), indicating EGFR activation causes the observed ERK pulses (**Fig. 4D,E,F,**
Supplementary Movie 3). Co-treatment with ALKi and MMPi similarly reduced Erk reactivation pulses
(ALKi vs ALKi/MMPi: 58 $\pm$ 7% vs 36 $\pm$ 7% of cells showing $\geq$ 1 pulse over 22 hrs; error = 95% CI),
though to a lesser extent than with EGFRi, potentially due to MMP-independent juxtacrine signals
(Brachmann et al. 1989; Wong et al. 1989) (**Fig. 4D,E,F, Supplementary Movie 3**). Thus, ALKi
treatment in STE-1 cells decreases Erk signals but is rapidly followed by RTK reactivation mediated by
paracrine signals.
239
Signal reactivation results from paracrine signals from dying cells.
We next sought to determine the source of the paracrine signals. We observed that pulsing
events often appeared next to dying cells, and that co-inhibition of ALKi with EGFRi or MMPi prevented
this pulsing (**Figure 5G,H, Supplementary Movie 3)**. These observations are consistent with paracrine
ligand secretion from apoptotic cells, which has previously been observed to promote cell survival and
homeostasis within epithelial sheets (Gagliardi et al. 2021). To quantify this effect, we observed signal
activation in cells that neighbored a dying cell within the ~hour preceding cell death (**Figure 5C).** We
then counted Erk pulses in these neighbors (N) and compared pulse counts to those from randomly
selected non-neighbors (R) over that same time interval (for more details, see **Methods).** Our analysis
revealed that, in the absence of ALKi, pulses were almost never observed in either the N or R
populations (DMSO, **Figure 5 D,E**). However, in ALKi-treated cells, N cells pulsed significantly more
than R cells (N vs R: 27 $\pm$ 4% vs 4 $\pm$ 2% cells with $\geq$ 1 pulse; error = 95% CI) (**Fig. 5D,E**). Co-treatment
with EGFRi eliminated Erk pulsing in neighbors (N cells in ALKi vs ALKi/EGFRi: 27 $\pm$ 4% vs 2 $\pm$ 1%),
whereas co-treatment with MMPi dramatically reduced neighbor signaling relative to ALKi alone (N cells
in ALKi vs ALKi/MMPi: 27 $\pm$ 4% vs 10 $\pm$ 2%),(**Fig. 5D,E).** Together, our results demonstrate that
virtually all observed Erk reactivation is associated with paracrine signals associated with dying drug-
treated cells. Importantly, because Erk pulses were not observed in untreated cells —even in neighbors
of dying cells (**Figure 5J**)—these data suggest that ALKi-induced RTK hypersensitization is an
essential first step for the perception of paracrine ligands during ALKi therapy.
259
Signal reactivation pulses activate gene transcription
To understand the extent to which the short ALKi-dependent Erk pulses could impact cell
behavior, we asked whether the observed pulses could stimulate downstream transcription. Erk activity
is a strong driver of transcription, including of a class of rapidly responding immediate early genes
(IEGs) that begin transcription within minutes of Erk activity(M. Z. Wilson et al. 2017). EGR1 is an IEG
that has been implicated in drug resistance to ALK inhibitors (Voena et al. 2013). Additionally, EGR1

expression is adaptive, such that its expression peaks by ~ 1 hr but then decays within 1-2 hours, even
in the presence of constant upstream signal (Sukhatme et al. 1987) (**Figure S8A**). Thus, accumulation
of EGR1 indicates the presence of only recent Erk activation. We thus examined the extent to which
EGR1 accumulated in STE-1 cells upon drug treatment (**Fig. 6A**). In untreated cancer cells, EGR1
levels remained low despite high Erk signaling from EML4-ALK, consistent with EGR1 adaptation to
tonic Erk signals (**Figure 6B,C**). Upon ALK inhibition, a distinct peak of EGR1-high cells appeared and
grew at 4 and 6 hrs after drug treatment, indicating that ALKi-induced Erk pulses could indeed drive
transcription. Importantly, co-inhibition of ALK and EGFR prevented the appearance of EGR1-positive
cells, consistent with transcription resulting from paracrine signals through EGFR (**Figure 6B,C**). Thus,
ALKi-induced Erk activity pulses provide sufficient signal to drive gene expression changes that could
regulate cell fate.
277
Signal reactivation promotes acute drug tolerance and cell persistence during ALK inhibition.
Finally, we asked whether RTK hypersensitization and resultant Erk activity pulses could counteract cell
killing and promote drug tolerance to ALK inhibitor therapy (**Fig. 6D).** In STE-1 cells imaged over 22
hrs, ALKi treatment induced moderate cell death ($45 \pm 7\%$ of cells; error = 95% CI). However, co-
treatment of ALKi with either EGFRi or MMPi increased cell death ($52 \pm 7\%$ and $75 \pm 6\%$ of cells,
respectively) (**Fig. 6E**). In both ALKi- and ALKi/MMPi-treated cells, surviving cells showed an increased
number of Erk pulses compared to cells that died (ALKi, live vs dead: 1.8 ± 0.2 vs 0.9 ± 0.1 pulses per
cell; ALKi/MMPi, live vs dead: 1.6 ± 0.2 vs 0.4 ± 0.1 pulses per cell. Mean $\pm$ SE, **Figure 6F**), further
drawing a link between RTK reactivation and cell survival. We confirmed our results through an
independent assay of cell death using a fluorescent reporter of caspase-3 activity (Nucview), an
indicator of apoptosis, over the first 24 hr of treatment in EML4-ALK+ cells. As before, while ALK
inhibition led to increased caspase-3, co-treatment with EGFRi significantly increased the caspase-3+
cell fraction in STE-1 cells (28 ± 2 vs. $41 \pm 2\%$). Similarly, co-treatment with MMPi also increased
caspase-3 (from 28 ± 2 to $50 \pm 6\%$) (**Figure 6G**). Similar trends were observed in H3122 cells (**Figure
S8B).** Neither EGFRi nor MMPi treatment alone showed enhanced killing over untreated cells. Notably,
addition of 50 ng/mL EGF reversed enhanced killing in the ALKi/MMPi condition in STE-1 cells (61 ± 2
to $39 \pm 1\%$, respectively), further suggesting that synergistic killing in the presence of MMPi resulted
from blockade of signal reactivation (**Figure 6H**). Although EGF addition could not rescue enhanced
killing under combined ALKi/EGFRi treatment, optogenetic pulses of optoFGFR lowered cell death in
this condition (**Figure S8C,D,E**). Finally, we observed that co-treatments of ALKi with either EGFRi or
MMPi dramatically suppressed long-term drug tolerance measured at 17-days (**Figure 6I**). Together,
our results indicate that hypersensitization of EGFR and restored perception of paracrine ligands leads
to reactivation of survival signals soon after drug treatment, which limits the cytotoxicity of ALK
therapies and promotes drug tolerance, the first step towards acquired resistance. The mechanisms
that underlie these events may thus provide novel treatment co-targets for to enhance therapy in EML4-
ALK+ cancers.
304
Discussion
Our results advance a novel function for protein condensates in cancer cells and their response
to targeted therapy (Jiang et al. 2020). In addition to amplifying oncogenic signaling (Tulpule et al.
2021), EML4-ALK condensates simultaneously act as a molecular sponge for RTK adapters including
Grb2 and SOS. This sequestration desensitizes the cell's perception of external ligands by limiting the

amount of adapters available to transduce the activated RTK signal. However, upon ALK inhibition,
adapters are rapidly released to the cytoplasm, hypersensitizing cellular response to ligands in the
cell's microenvironment. RTK hypersensitization promotes cell survival and drug tolerance in response
to targeted inhibitors, due at lease in part to apoptosis-induced paracrine signals.
314   Adapter sequestration by fusion condensates represents a novel mechanism by which
oncogenes can suppress RTK signaling. Analogous RTK suppression has been observed in other,
molecularly distinct cancers. BRAF V600E+ melanoma and colorectal cancer cells suppress RTKs
through Erk-dependent transcription of negative regulators (Sproutys), and through inhibitory
phosphorylation of SOS1(Lito et al. 2012b; Prahallad et al. 2012b; Gerosa et al. 2020b). Inhibition of
BRAF suppresses these mechanisms and leads to rapid, pulsatile Erk reactivation and drug resistance,
similar to our observations. Separately, in cancers driven by EGFR, Erk provides suppressive
phosphorylation of EGFR receptors, which is lost during MEK inhibition and leads to reactivation of
ErbB3 and PI3K(Turke et al. 2012). In contrast to these and similar examples, our findings
demonstrate that such RTK suppression can also be implemented through biophysical, rather than
biochemical, feedback.   Nevertheless, the variety of mechanisms by which oncogenes can suppress
RTKs hints that such feedback might play an important role in establishing permissive conditions for
oncogenesis. It will thus be interesting to more comprehensively understand the diverse oncogenic
contexts in which such suppression and reactivation occurs.
328   It has been repeatedly observed that ligand-induced signaling through EGFR promotes survival
and resistance to ALK inhibitors in EML4-ALK+ cancer cells (T. R. Wilson et al. 2012; Vander Velde et
al. 2020; Tani et al. 2016; Tanimoto et al. 2014; Vaishnavi et al. 2017; Sasaki et al. 2011; Obenauf et
al. 2015). Our work builds on these prior studies to show that 1) ALK inhibition is a necessary first step
to allow cancer cell perception of paracrine signals and 2) ALK-induced resensitization to RTK signals
happens within minutes of drug treatment. In addition to fast RTK reactivation, previous work found that
slower-timescale transcriptional amplification of KRas or downregulation of DUSP6 contributes to
resistance development (Hrustanovic et al. 2015). Our results are complementary to these findings,
which could be expected to further sensitize cells to external RTK ligands.
337   We found that dying cancer cells send survival signals to their neighbors through a mechanism
that requires proteolytic processing of EGFR ligands, thus adding to previously identified mechanisms
whereby drug treatment induces secretion that promotes cell survival and drug tolerance (Obenauf et
al. 2015; Kurtova et al. 2015). Blockade of apoptosis-induced paracrine signals enhanced cell killing
and limited tolerance in response to ALK inhibition. Although co-inhibition of EGFR is highly effective in
promoting durable drug responses across a variety of cancer cells, (Vaishnavi et al. 2017; Vander
Velde et al. 2020; Sasaki et al. 2011; Voena et al. 2013), a recent clinical trial of combined inhibition of
crizotinib (ALKi) and erlotinib (EGFRi) in EML4-ALK+ NSCLC failed due to frequent adverse effects and
low maximum tolerated dose (Ou et al. 2017). We show that co-inhibition of certain matrix
metalloproteases may be an alternative strategy limit RTK signaling during targeted therapy. We note,
however, that previous studies found that long-term MMP inhibition can promote resistance by causing
accumulation of transmembrane RTKs (e.g. AXL) (Miller et al. 2016). Thus, further studies will be
required to understand whether MMPi co-treatment can indeed enhance therapy, or whether MMPi
scheduling could leverage acute benefits while avoiding deleterious chronic effects.
351   Our finding that the sequestration of RTK adapters can functionally desensitize RTK signals
could inspire new types of therapeutic targets that mimic this behavior, for example through inhibition or
sequestration of important RTK adapters. Of note, the recent discovery of EML4-ALK condensates has

raised the idea that disaggregation of the condensates might be therapeutically beneficial (Hirai et al.
2020; Cai, Liu, and Lippincott-Schwartz 2021). Our work cautions, however, that such disaggregation
strategies will likely be subject to the same adapter redistribution and rapid RTK reactivation observed
with small molecule ALK inhibitors, potentially challenging their efficacy.
358   A unique promise of functional profiling is that common signaling abnormalities may be
identified and inform therapies among genetically distinct cancers (Lukasz J. Bugaj, O'Donoghue, and
Lim 2017). Because our work revealed signaling principles that result from a likely common property
(condensation) of RTK fusions (Du and Lovly 2018; Tulpule et al. 2021), we anticipate that our findings
may be broadly relevant to other cancer cells in this class. We show that in a cancer line driven by a
CCDC6-RET fusion that also forms large aggregates (Tulpule et al. 2021), RET activity similarly
suppresses EGFR, and RET inhibition potentiates EGFR signaling (**Figure S7)**. These results are
consistent with a previous study that showed that Grb2 can change association from the fusion to
EGFR upon kinase inhibition in cancers driven by ALK, RET, NTRK1 and ROS1 fusions, further
implying common behavior across molecularly distinct RTK fusions (Vaishnavi et al. 2017). Future work
will determine the extent to which the principles we describe in EML4-ALK+ cancers will extend to this
more diverse array of malignancies.
370

371   **Acknowledgements.** The authors thank Dr. Magdalena Niewiadomska-Bugaj for feedback on
372   statistics, data presentation and analysis. The authors also thank Dr. Alex Hughes (Penn) and Dr.
373   Sydney Shaffer, and Dr. Yael Mosse for helpful comments on the manuscript. The Beas2B cell line with
374   endogenously tagged Grb2 was a gift from Bo Huang, PhD (UCSF). This work was supported by the
375   National Institutes of Health (R35GM138211 for L.J.B).

376
377   **Author Contributions** D.G.M. and L.J.B. conceived the study. D.G.M. and L.R. performed
378   experiments. D.G.M., T.R.M. and L.J.B. analyzed data. L.J.B. supervised the work. A.T., T.G.B., L.R.,
379   and L.J.B. provided experimental feedback and guidance. D.G.M and L.J.B. wrote the manuscript and
380   made figures, with editing from all authors.
381
382   **Conflicts of interest.** D.G.M. and L.J.B. have filed a provisional patent based on the findings in this
383   work.

392   **Figures**

393
Figure 1. Optogenetic functional profiling of EML4-ALK+ cancer cells reveals suppression of
RTK signaling. A) EML4-ALK+ cancer cells treated with ALK inhibitors (ALKi) can persist through
therapy and acquire stable drug resistance. B) Immunofluorescence staining shows punctate
expression of ALK in two EML4-ALK+ cancer cell lines. C) Although EML4-ALK and transmembrane
RTKs signal through the same downstream adapters, the extent to which EML4-ALK condensates
interact with RTKs, and whether interactions could impact drug responses, is not clear. D) Functional
profiling allows us to observe input/output responses to RTK signaling to understand functional
interactions between RTKs and EML4-ALK. E) optoFGFR allows blue-light-induced stimulation of
FGFR signaling, including through the Ras/Erk pathway. F) Single-cell analysis of ppErk levels in STE-
1 cells stimulated with light (optoFGFR) in the presence of either DMSO (grey) or the ALK inhibitor
crizotinib (ALKi, 1μM; orange). F) ppErk fold-change in response to 5 min of blue light stimulation. Blue
light intensity ranged from 2% to 60% of maximum (1.45 W/cm^2). Data points show the ratio of ppErk
from stimulated and unstimulated cells. Each condition was performed in biological replicate, with each
replicate representing the mean of ~1000 cells. This experiment is representative of three independent
experiments.

Figure 2. EML4-ALK suppresses EGFR signaling. A) EGF stimulates EGFR and downstream Ras/Erk signaling. B) ppErk levels in response to EGF (100 ng/mL) in the presence of 1µM ALKi (orange) or DMSO (grey) in STE-1 and H3122 cancer cells. Data points and ribbons represent mean +/- SEM of three replicates, each of which represents the mean of ~1000 cells. C) Quantification of dynamic range from experiment in 2B over a range of EGF concentrations. D) EML4-ALK(V1) was transiently expressed in lung epithelial Beas2B cells. E) Time course of ppErk immunofluorescence levels in response to EGF stimulation (50 ng/mL). Data points represent mean +/- SEM of three replicates. G) Dynamic range of ppErk in EML4-ALK-expressing Beas2B in response to EGF in the presence or absence of ALKi (1 µM) pretreatment.

Figure 3. Mapping RTK feedback suppression using optogenetics. A) Time course of RTK
hypersensitization was obtained by pre-incubating cancer cells with 1 µM crizotinib for varying times
before stimulation with EGF (50 ng/mL) for 15 min, followed by fixation and immunostaining for ppErk.
B) ppErk induction as a function of ALKi pre-incubation time, quantified by immunofluorescence. Open
circles = unstimulated. Closed circles = EGF stimulated. Data points represent mean +/- SEM of
triplicates. Error bars are smaller than the data points for unstimulated samples. C)Pinpointing the
location of EML4-ALK interaction with RTK/Erk signaling. D) OptoFGFR and optoSOS permit
optogenetic stimulation at successive nodes of the pathway. E) Quantification of immunofluorescence
of ppErk in response to optoFGFR or optoSOS in the presence or absence of ALKi. ALK-dependent
suppression is observed only with optoFGFR, suggesting that suppression happens upstream of Ras
but downstream of RTK activation (F). G) Testing the role of Erk-dependent negative feedback on RTK
suppression. Stimulating optoSOS drives high levels of Erk during ALKi treatment and sustains any
Erk-dependent negative feedback that would otherwise be lost during ALK inhibition. H) STE-1 cells
were treated with either ALKi or ALKi and optoSOS, and EGF response was assessed. I) Predicted
results and implications for Erk-dependent feedback. J) Cells treated with both ALKi and optoSOS
showed the same increase in EGF-induced ppErk as cells receiving ALKi only, suggesting that Erk-
dependent negative feedback does not suppress EGFR in EML4-ALK+ cancer cells. Data points
represent means of ~600-1000 cells per condition.

448
Figure 4. EML4-ALK condensates suppress RTK signals through sequestration of RTK
effectors. A) Quantification of ppErk response to EGF (50ng/mL) stimulation in Beas2B cells that
expressed EML4-ALK (V1), EML4-ALK (ΔTD), and kinase-dead EML4-ALK (K589M). Data are means
of biological triplicates, each of which represents the mean of 500-1000 cells. Error = 1 SEM. B)
Beas2B cells that harbored endogenously-tagged Grb2 (Grb2:mNG) were transiently transfected with
EML4-ALK and stimulated with EGF (50ng/mL) to visualize Grb2 translocation in the presence and
absence of EML4-ALK. C) Impaired membrane translocation of Grb2 in the presence of EML4-ALK

condensates. Time in mm:sec. See Supplementary Movie 1. D) Line scan of Grb2 intensity distribution
in the presence (red) or absence (green) of EML4-ALK expression, as depicted in C. E) Grb2
localization was visualized upon treatment with 1 µM ALKi. F) Grb2 shifted from condensates to
cytoplasm after ALKi treatment. Time in hh:mm. See Supplementary Movie 2. G) Quantification of Grb2
cluster dissociation kinetics after ALKi treatment. H) Grb2 distribution was examined upon sequential
treatment with ALKi and EGF. I) ALKi released Grb2 to the cytoplasm, which allowed its membrane
translocation in the presence of EGF. J) Immunoprecipitation of EGFR shows co-precipitation of SOS1
only in the presence of ALKi pretreatment and EGF. K) EML4-ALK sequesters adapters like
Grb2/SOS1 and prohibits their translocation to RTKs upon ligand stimulation. ALK inhibition releases
adapter sequestration and restores cellular response to RTK stimulation.
466

467
468
469

Figure 5. ALK inhibition resensitizes cancer cells to paracrine growth factors secreted from
dying neighbor cells. A) The ErkKTR reporter indicates Erk activity through nuclear-cytoplasmic
translocation of a fluorescent protein. (B) Live cell imaging of STE-1 cells expressing ErkKTR in the
presence or absence of ALKi (1 µM crizotinib). See **Supplementary Movie 3**. C) Representative
single-cell traces of cytoplasmic/nuclear ErkKTR intensity ratio from conditions shown in B. D)
Quantification of ErkKTR activity in the presence of ALKi or its combination with EGFRi (1 µM erlotinib)
or MMPi (10 µM marimastat). E) Quantification of Erk activity pulses per cell. F) Fraction of cells that
exhibited any pulses over 21.5 hrs of imaging. Error bars indicate 95% CI. n = 200 cells per condition.

G) Apoptotic cells can secrete paracrine EGFR ligands to their neighbors. Paracrine signaling can be
blocked by inhibiting either EGFR or the MMPs that are required to shed certain EGFR ligands from the
surface of the sender cell. H) ErkKTR activity pulses are primarily observed surrounding a dying cell
during ALK inhibition, but not in the absence of drug or in the presence of EGFR or MMP inhibitors. I)
Definition of neighbors and non-neighbors of a death event. J) Quantification of pulses per cell for each
death event in neighbors or a randomly-chosen subset of cells not near a death event (see **Methods**
for more details). K) Fraction of total neighbor vs random non-neighbor cells that show any pulsing.
Error bars = 95% CI.
486

FIgure 6. Resensitization to paracrine signals drives transcription and promotes drug tolerance.
A) Signaling through EGFR activates Ras/Erk and stimulates transcription, including of the immediate
early gene EGR1. B) Upon ALK inhibition, EGR1 expression increases, but not upon co-inhibition of
ALK and EGFR. C) Comparison of EGR1 expression after 6 hrs of the indicated treatment. D) We
hypothesized that restored perception of paracrine signals might promote survival, and that blocking
paracrine signals would promote killing. E) Visualization of live cell imaging of Erk activity pulses and
cell death. F) Quantification of pulse per cell in cells that died (D) or survived (L) through 22 hrs of
imaging, in the conditions where Erk pulsing could be observed. H) Caspase-3 activation was assessed
using the NucView reporter after 24 hr treatment with the indicated drugs. I) EGF (50 ng/mL) addition
promotes survival during ALKi treatment and counteracts enhanced killing from ALKi/MMPi co-
treatment, but not from ALKi/EGFRi co-treatment. Dotted line is cell killing from DMSO control
treatment. Each data point represents the fraction of caspase-3+ cells out of ~10,000 total cells. See
Methods for more details. I) Crystal violet staining (left) and cell counts (right) of cell survival after 17
days of the indicated treatments.

502
503
Figure 7. Model of how EML4-ALK condensates interact with transmembrane RTKs and its
effect on drug tolerance. We find that EML4-ALK condensates suppress perception of EGFR signals
by sequestering RTK adapters like Grb2 and SOS1. Upon ALK inhibition, adapters are released to the
cytoplasm where they can transduce signals from EGFR and other RTKs. Although ALK inhibitors
effectively suppress ALK activity, they simultaneously restore a cell's perception to paracrine signals,
including those from neighboring apoptotic cells. Such paracrine signals promote drug tolerance, and
blocking them with inhibitors of either EGFR or MMPs enhances cell killing during ALK inhibitor therapy.
511
512

Supplementary Figures

Supplementary Figure 1. OptoFGFR functional profiling of STE-1 cancer cells. A) optoFGFR
comprises the intracellular domain of mouse FGFR1 fused to the PHR domain from Arabidopsis
Cryptochrome 2, which clusters under blue light stimulation. The construct is anchored in the membrane
through N-terminal myristoylation. B) Single cell distributions of ppErk intensity in STE-1 cells stimulated
by the indicated intensity of blue light in the presence (orange) or absence (grey) of ALKi (2 hr pre-
incubation of 1µM crizotinib). Strong ppErk response at low light intensity in the presence, but not the
absence, of ALKi suggests that ALKi hypersensitizes cells to RTK stimulation. C) The magnitude of ppErk
response in optoFGFR STE-1 cells is a function of both light intensity and optoFGFR expression levels.
Results are presented by binning optoFGFR-mCh expression by quartiles. (1<25%, 25% < 2 < 50%,
50% <3 < 75%, 4 > 75%).

Supplementary Figure 2. Schematics of the optoFGFR and optoSOS tools used in Figure 3.

531
532
Supplementary Figure 3. EGFR signal suppression does not result from decreased EGFR
phosphorylation. Following overnight starvation, STE-1 cells were pretreated with ALKi or
DMSO for 2 hrs, were stimulated with either EGF (50 ng/mL) or fresh starvation medium for 15
min, and then were fixed and immunostained. ALKi pretreatment did not alter pEGFR (Y1068)
levels in either the presence or absence of EGF, whereas pALK (Y1507) decreased. Thus
increased EGFR signaling in ALKi pretreated cells does not result from increased EGFR
phosphorylation.
540

Supplementary Figure 4. Optogenetic maintenance of high ppErk during ALKi treatment. Demonstration that optoSOS can maintain high Erk levels (comparable to those from the uninhibited oncogene) during ALKi pre-treatment. Optogenetic stimulation of Ras/Erk was used to decouple ALK inhibition from loss of Erk signaling in **Figure 3G-J**.

547
548
549 **Supplementary Figure 5. EML4-ALK condensates colocalize with Grb2 puncta.** Beas2B cells that
550 harbored endogenously tagged Grb2 (Grb2:mNG) were transfected with EML4-ALK-mCh. In transfected
551 cells, EML4-ALK co-localized with Grb2. In untransfected cells, Grb2 was diffuse.
552

553   **Supplementary Figure 6. Co-immunoprecipitation of EGFR with SOS1.** H3122 cells were starved
554   overnight, pretreated with crizotinib (2 hr), and were stimulated with EGF for 2 minutes before lysis and
555   immunoprecipitation. Right 4 lanes are reproduced from **Figure 4J**.
556
557
558
559

561
Supplementary Figure 7. CCDC6-RET+ cancer cells show dynamic suppression of EGFR through
RTK fusion activity. TPC-1 cells harbor CCDC6-RET, a distinct RTK fusion that forms condensates.
We tested the extent to which CCDC6-RET could modulate EGFR dynamic range as observed for EML4-
ALK. A) Plot shows ppErk intensity response to EGF stimulation after 2 hr pre-treatment with DMSO
(grey) or 1 μM BLU-667 (purple), a RET inhibitor. Data represents mean, and error bars represent 95%
CI of triplicates, each of which represents ~1000 cells. B) Quantification of ppErk dynamic range shows
that oncogene inhibition enhances signaling through EGFR. Thus, multiple RTK fusions that form
condensates dynamically suppress EGFR signaling.
570
571
572

Supplementary Figure 8. RTK/Erk pulses activate transcription and promote cell survival during ALKi treatment. A) Adaptive response of EGR1 expression in STE-1 cells treated with ALKi and 50 ng/mL of EGF. EGR1 peaks at 1 hr after stimulation and then adapts to near baseline expression. EGR response is stronger and earlier than observed in **Figure 6B** because here all cells were stimulated at t = 0, whereas in **6B** only individual cells were stimulated sporadically through paracrine signals starting 1-2 hrs after ALK treatment. B) Blockade of EGFR or MMPs accelerates cell death in H3122 in response to ALKi, as shown STE-1 cells in **Figure 6G**. C) We tested whether optoFGFR stimulation could reduce the enhanced cell death response in STE-1 cells co-treated with ALKi and EGFRi. D). A 10 min pulse of blue light (96 mW/cm^2) every hour produced strong, periodic ppErk levels. E) Hourly pulses of optoFGFR/Erk (10 min of 96 mW/cm^2) reduced cell death in both ALKi and ALKi/EGFRi treated cells. Data represents the mean, error bars = 95% CI caspase-3+ cells of three replicates, each representing ~10,000 cells.

Methods

Cell Lines and Cell Culture

All cell lines were maintained at 37°C and 5% CO_2 using standard cell culture incubator. STE-1, H3122, TPC-1 and Beas2B cell lines were cultured in RPMI-1640 media containing L-glutamine supplement with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin (P/S). For experiments, cells were seeded in 96 or 384 well plates coated with 10 µg/mL MilliporeSigma™ Chemicon™ Human Plasma Fibronectin Purified Protein in PBS. For 96 well plate experiments 10,000 STE-1 and H3122 cells were seeded in 200µl of cell culture media per well. For 384 well plate experiments 5,000 STE-1 cells, 5000 H3122 cells, or 2000 Beas2B cells were seeded in 50µl of culture media per well.

Constructs

EML4-ALK variant 1 (EML4ALK(V1)) cDNA was cloned into a pCMV vector with a C-terminal 2A-H2B-iRFP670 tag. Deletion of EML4 trimerization domain was achieved by deleting bases 310-459 of EML4-ALK full-length sequence GenBank: AB274722.1 and EML4-ALK kinase-dead was cloned by mutating K589M as previously described (Tulpule et al. 2021). EML4ALK(V1)-mCh was cloned by replacing the C-terminal 2A-H2B-iRFP670 tag with mCh. Erk kinase translocation reporter (KTR-mCh) was cloned into a pHR lentiviral vector with a C-terminal mCh tag. H2B-iRFP670 nuclear marker was cloned into a pHR. For optogenetic control of FGFR signaling, the cytoplasmic domain of FGFR1 receptor with an N-terminal myristylation site was fused to the N-terminal of mCh-Cry2. All cloning was performed using NEB Hifi DNA assembly reagent.

Transfection and viral transduction

EML4ALKV1, ΔTD EML4ALK, and K589M EML4ALK plasmids were transfected into Beas2B cells using Lipofectamine™ 3000 Transfection Reagent (Invitrogen, L3000001) following manufacturer recommendations. OptoFGFR, OptoSOS, KTR-mCh, and H2B-iRFP670 vectors were packaged into lentivirus using lentiX- HEK 293T cells. Briefly cells were transfected with pHR packaging vectors and supernatant was collected over two days. 300,000 STE-I cells were then seeded in 6 well plate with media containing virus. Cells were then expanded, and successfully transduced cells were sorted using fluorescence activated cell sorting (BD Influx System)

Optogenetic Stimulation

Light stimulation was achieved using a single-color blue LED optoPlate-96(L. J. Bugaj and Lim 2019). A 13 mm tall black adaptor was used for even light diffusion across each of the 384 well plate wells being illuminated. Light intensity was corrected to stimulate wells with a range of blue

629 light intensities spanning from 0 to 870 mW/cm^2. Cells were stimulated for 500ms every 10
630 seconds at the specified light intensity.
631
Growth Factor Stimulation Assay
24 hr following cell plating, cells were starved with RPMI-1640 media containing L-glutamine
and 1% P/S using the a BioTEK 405 platge washer. 24 hrs after starvation, cells were treated
with inhibitor for 2 hours (1µM Crizotinib (Sigma-Aldrich, PZ0191) or 1-100 µM BLU-667
(Biovision, B2548)). Cells were then stimulated at their respective time points with epidermal
growth factor (EGF) (PeproTech 315-09) following a delay that allows for all wells to be fixed at
the same time. Cells were then immediately prepared for immunostaining.
639
Immunostaining
Immediately after treatment, cells were fixed using 4% paraformaldehyde for 10 min and
permeabilized using 0.5% Triton-X100 in PBS for 10 min. Cells were further permeabilized using
100% methanol at -20°C for 10 min and samples were blocked with 1% bovine serum albumin
(Fisher, BP9706100) in PBS (PBS+1%BSA) for 1 hour at room temperature. Cells were then
incubated in primary antibody diluted in PBS+1%BSA for 2 hours at room temperature or
overnight at 4C (phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204), Cell Signaling, catalog
number 4370, 1:400 dilution; Phospho-EGF Receptor (Tyr1068), Cell Signaling, catalog number
3777, 1:800; EGR1, Cell Signaling, catalog number 4153, 1:800). Primary antibody was then
washed with 0.1% Tween-20 in PBS (PBS+T). Cells were then incubated in secondary antibody
diluted in PBS+1%BSA as recommended (IgG (H+L) Cross-Adsorbed Goat anti-Rabbit,
DyLight™ 488, Invitrogen, catalog number 35553, 1:500; Goat anti-Rabbit IgG (H+L) Cross-
Adsorbed Secondary Antibody, DyLight™ 650, Invitrogen, catalog number SA510034, 1:500)
and and 4,6-diamidino-2-phenylindole (DAPI; ThermoFisher Scientific, catalog number D1306,
300 nM) for 1 hour at room temperature and washed with PBS+T.
655
Live Cell Imaging
A Nikon Ti2-E microscope equipped with a Yokagawa CSU-W1 spinning disk, 405/488/561/640
nm laser lines, an sCMOS camera (Photometrics), and a motorized stage was used for live cell
imaging. Temperature was maintained at 37°C and CO2 concentration was maintained at 5%
using an environmental chamber (Okolabs). Transfected Beas2B::mNGGrb2 cells were imaged
48 hours after transfection with a 40X oil immersion lens. To visualize Grb2 localization in
response to growth factor stimulation after oncogene inhibition, cells were imaged for 2 hours
following 1µM crizotinib treatment. Cells were then stimulated with 50 ng/mL EGF (PreproTech
315-09). STE1 cells stably expressing ERKKTR-Ruby and H2B-iRFP670 were seeded in a 96
well plate (1.5x10^5 cells/well) and cultured in RPMI supplemented with 2% FBS. 48 hours post-
seeding, cells were supplemented with NucView (1:1000), Criztinib (1 µM), Erlotinib (1 µM),
Marimastat (10 µM) or DMSO (1:5000). Cells were imaged every 5 minutes for 22 hours at 20X
magnification.

<u>Cell Death Assay</u>

The day following cell plating, cells were washed with phenol free RPMI-1640 with L-glutamine supplemented with 2% FBS, 1% P/S, 1μM NucView 488, and respective inhibitor treatment (Crizotinib 1μM, Erlotinib 1μM, Marimastat 10μM). 24 hours after treatment addition, cells were treated with 5 μg/mL Hoechst 33342 Hydrochloride (Cayman 15547) for 15 minutes. Cells were then imaged using confocal microscopy as previously described.

<u>Image Processing and Analysis</u>

Immunostaining and live imaging of caspase+ cells was performed using Cell Profiler version 4.0.7. Briefly, cell nuclei were segmented using the DAPI or Hoechst 33342 Hydrochloride stain and cytoplasm was obtained by expanding a 5-pixel ring from the nuclei. Data was then exported into R for data visualization using the dplyr and ggplot2 packages.

<u>Live cell image analysis.</u>

Grb2 cluster dissociation

Grb2 clusters time lapse imaging was quantified with a custom MATLAB script. Briefly, cells were segmented manually and clusters were identified through the following steps. First, cell intensities were first normalized to their internal median, and a series of transforms were performed to enhance clusters while suppressing other features, like background and high frequency noise. These transformed intensities were used to identify bright pixels at the center of clusters, using a user-defined intensity cutoff. From each of these center points, neighboring pixels were compared to a threshold intensity that was set by the local background of the cell and scaled by a second user-defined parameter. Neighboring pixels above this threshold are included in the cluster, while pixels below the threshold are excluded. This occurs iteratively, where adjacent pixels to each included pixel were also checked to be included or excluded until all new neighbors were below the threshold.

ErkKTR dynamics

ErkKTR dynamics and cell death were tracked and quantitated in a semi-automated manner using the p53CellCinema package (Reyes et al. 2018) in MATLAB, and data was processed and visualized in R and RStudio using the tidyR(Wickham et al. 2019) package. Briefly, cells with low or no ErkKTR expression were first excluded from analysis. Then, ErkKTR cytoplasmic/nuclear ratios from remaining cells were imported into MATLAB, and peaks were identified using the 'findpeaks' function. Peak calls were then manually inspected for outliers, for example resulting from apoptotic cell debris, and outlier cells were removed from analysis. Of note, we consistently observed Erk pulses preceding cell division events across conditions. This phenomenon has been previously observed and has been shown to be independent of Ras/Erk signaling, potentially due to non-specific activation from a cyclin dependent kinase(Gerosa et al.

2020). Thus, we disregarded all pulses that occurred within 10 frames (50 mins) preceding cell
division. To count Erk pulses in neighbors vs non-neighbors of dying cells, we first identified
each cell death event. We then identified the neighbors of the dying cell by identifying which
nuclei centroids were within a 100 pixel radius of the dying cell for each of the 10 frames
preceding the death annotation. We then counted the total pulses of those neighbors over the
indicated time frame, and compared against pulse counts from a random subset of non-neighbor
cells over the same time frame. Importantly, for each death event, the number of non-neighbors
that were sampled matched the number of neighbors, and the sampled non-neighbors were not
neighbors of other death events over that same time span. For visualization in **Figure 6E**, cells
that divided during the course of the experiment are represented as separate cells, and the Erk
history of the mother cell is reproduced for both cells. However, for quantification of Erk pulses,
mother cell pulses were counted only once.

References

Banani, Salman F., Hyun O. Lee, Anthony A. Hyman, and Michael K. Rosen. 2017. "Biomolecular
Condensates: Organizers of Cellular Biochemistry." *Nature Reviews. Molecular Cell Biology* 18
(5): 285–98.
Benman, William, Erin E. Berlew, Hao Deng, Caitlyn Parker, Ivan A. Kuznetsov, Bomyi Lim, Arndt F.
Siekmann, Brian Y. Chow, and Lukasz J. Bugaj. 2022. "Temperature-Responsive Optogenetic
Probes of Cell Signaling." *Nature Chemical Biology* 18 (2): 152–60.
Brachmann, R., P. B. Lindquist, M. Nagashima, W. Kohr, T. Lipari, M. Napier, and R. Derynck. 1989.
"Transmembrane TGF-Alpha Precursors Activate EGF/TGF-Alpha Receptors." *Cell* 56 (4): 691–
700.
Bugaj, L. J., A. J. Sabnis, A. Mitchell, J. E. Garbarino, J. E. Toettcher, T. G. Bivona, and W. A. Lim.
2018. "Cancer Mutations and Targeted Drugs Can Disrupt Dynamic Signal Encoding by the
Ras-Erk Pathway." *Science* 361 (6405). https://doi.org/10.1126/science.aao3048.
Bugaj, Lukasz J., Geoff P. O'Donoghue, and Wendell A. Lim. 2017. "Interrogating Cellular Perception
and Decision Making with Optogenetic Tools." *The Journal of Cell Biology* 216 (1): 25–28.
Cai, Danfeng, Zhe Liu, and Jennifer Lippincott-Schwartz. 2021. "Biomolecular Condensates and Their
Links to Cancer Progression." *Trends in Biochemical Sciences* 46 (7): 535–49.
Corcoran, Ryan B., Hiromichi Ebi, Alexa B. Turke, Erin M. Coffee, Michiya Nishino, Alexandria P.
Cogdill, Ronald D. Brown, et al. 2012. "EGFR-Mediated Re-Activation of MAPK Signaling
Contributes to Insensitivity of BRAF Mutant Colorectal Cancers to RAF Inhibition with
Vemurafenib." *Cancer Discovery* 2 (3): 227–35.
Davies, Kurtis D., Sakshi Mahale, David P. Astling, Dara L. Aisner, Anh T. Le, Trista K. Hinz, Aria
Vaishnavi, et al. 2013. "Resistance to ROS1 Inhibition Mediated by EGFR Pathway Activation in
Non-Small Cell Lung Cancer." *PloS One* 8 (12): e82236.
Dong, J., L. K. Opresko, P. J. Dempsey, D. A. Lauffenburger, R. J. Coffey, and H. S. Wiley. 1999.
"Metalloprotease-Mediated Ligand Release Regulates Autocrine Signaling through the
Epidermal Growth Factor Receptor." *Proceedings of the National Academy of Sciences of the*

755    *United States of America* 96 (11): 6235–40.

756 Du, Zhenfang, and Christine M. Lovly. 2018. "Mechanisms of Receptor Tyrosine Kinase Activation in
757    Cancer." *Molecular Cancer* 17 (1): 58.

758 Gagliardi, Paolo Armando, Maciej Dobrzyński, Marc-Antoine Jacques, Coralie Dessauges, Pascal
759    Ender, Yannick Blum, Robert M. Hughes, Andrew R. Cohen, and Olivier Pertz. 2021. "Collective
760    ERK/Akt Activity Waves Orchestrate Epithelial Homeostasis by Driving Apoptosis-Induced
761    Survival." *Developmental Cell* 56 (12): 1712-1726.e6.

762 Gainor, Justin F., Leila Dardaei, Satoshi Yoda, Luc Friboulet, Ignaty Leshchiner, Ryohei Katayama,
763    Ibiayi Dagogo-Jack, et al. 2016. "Molecular Mechanisms of Resistance to First- and Second-
764    Generation ALK Inhibitors in ALK-Rearranged Lung Cancer." *Cancer Discovery* 6 (10): 1118–
765    33.

766 Garabedian, Mikael V., Wentao Wang, Jorge B. Dabdoub, Michelle Tong, Reese M. Caldwell, William
767    Benman, Benjamin S. Schuster, Alexander Deiters, and Matthew C. Good. 2021. "Designer
768    Membraneless Organelles Sequester Native Factors for Control of Cell Behavior." *Nature*
769    *Chemical Biology* 17 (9): 998–1007.

770 Gerosa, Luca, Christopher Chidley, Fabian Fröhlich, Gabriela Sanchez, Sang Kyun Lim, Jeremy
771    Muhlich, Jia-Yun Chen, et al. 2020a. "Receptor-Driven ERK Pulses Reconfigure MAPK
772    Signaling and Enable Persistence of Drug-Adapted BRAF-Mutant Melanoma Cells." *Cell*
773    *Systems* 11 (5): 478-494.e9.

774 Hirai, Noriko, Takaaki Sasaki, Shunsuke Okumura, Yoshinori Minami, Shinichi Chiba, and Yoshinobu
775    Ohsaki. 2020. "Monomerization of ALK Fusion Proteins as a Therapeutic Strategy in ALK-
776    Rearranged Non-Small Cell Lung Cancers." *Frontiers in Oncology* 10 (April): 419.

777 Hrustanovic, Gorjan, Victor Olivas, Evangelos Pazarentzos, Asmin Tulpule, Saurabh Asthana, Collin M.
778    Blakely, Ross A. Okimoto, et al. 2015. "RAS-MAPK Dependence Underlies a Rational
779    Polytherapy Strategy in EML4-ALK–Positive Lung Cancer." *Nature Medicine* 21 (August): 1038.

780 Jiang, Shan, Johan Bourghardt Fagman, Changyan Chen, Simon Alberti, and Beidong Liu. 2020.
781    "Protein Phase Separation and Its Role in Tumorigenesis." *ELife* 9 (November).
782    https://doi.org/10.7554/eLife.60264.

783 Kim, Nury, Jin Man Kim, Minji Lee, Cha Yeon Kim, Ki-Young Chang, and Won Do Heo. 2014.
784    "Spatiotemporal Control of Fibroblast Growth Factor Receptor Signals by Blue Light." *Chemistry*
785    *& Biology* 21 (7): 903–12.

786 Kurtova, Antonina V., Jing Xiao, Qianxing Mo, Senthil Pazhanisamy, Ross Krasnow, Seth P. Lerner,
787    Fengju Chen, et al. 2015. "Blocking PGE2-Induced Tumour Repopulation Abrogates Bladder
788    Cancer Chemoresistance." *Nature* 517 (7533): 209–13.

789 Lito, Piro, Christine A. Pratilas, Eric W. Joseph, Madhavi Tadi, Ensar Halilovic, Matthew Zubrowski,
790    Alan Huang, et al. 2012a. "Relief of Profound Feedback Inhibition of Mitogenic Signaling by RAF
791    Inhibitors Attenuates Their Activity in BRAFV600E Melanomas." *Cancer Cell* 22 (5): 668–82.

792 Lovly, Christine M., Nerina T. McDonald, Heidi Chen, Sandra Ortiz-Cuaran, Lukas C. Heukamp,
793    Yingjun Yan, Alexandra Florin, et al. 2014. "Rationale for Co-Targeting IGF-1R and ALK in ALK
794    Fusion-Positive Lung Cancer." *Nature Medicine* 20 (9): 1027–34.

795 Miller, Miles A., Madeleine J. Oudin, Ryan J. Sullivan, Stephanie J. Wang, Aaron S. Meyer, Hyungsoon
796    Im, Dennie T. Frederick, et al. 2016. "Reduced Proteolytic Shedding of Receptor Tyrosine
797    Kinases Is a Post-Translational Mechanism of Kinase Inhibitor Resistance." *Cancer Discovery* 6
798    (4): 382–99.

799 Obenauf, Anna C., Yilong Zou, Andrew L. Ji, Sakari Vanharanta, Weiping Shu, Hubing Shi, Xiangju
800    Kong, et al. 2015. "Therapy-Induced Tumour Secretomes Promote Resistance and Tumour
801    Progression." *Nature* 520 (7547): 368–72.

802 Omer, Amr, Devang Patel, Xian Jin Lian, Jason Sadek, Sergio Di Marco, Arnim Pause, Myriam
803    Gorospe, and Imed Eddine Gallouzi. 2018. "Stress Granules Counteract Senescence by
804    Sequestration of PAI-1." *EMBO Reports* 19 (5). https://doi.org/10.15252/embr.201744722.

805 Ou, Sai-Hong Ignatius, Ramaswamy Govindan, Keith D. Eaton, Gregory A. Otterson, Martin E.

Gutierrez, Alain C. Mita, Athanassios Argiris, et al. 2017. "Phase I Results from a Study of
Crizotinib in Combination with Erlotinib in Patients with Advanced Nonsquamous Non-Small Cell
Lung Cancer." *Journal of Thoracic Oncology: Official Publication of the International Association*
for the Study of Lung Cancer 12 (1): 145–51.
Peschon, J. J., J. L. Slack, P. Reddy, K. L. Stocking, S. W. Sunnarborg, D. C. Lee, W. E. Russell, et al.
1998. "An Essential Role for Ectodomain Shedding in Mammalian Development." *Science* 282
(5392): 1281–84.
Prahallad, Anirudh, Chong Sun, Sidong Huang, Federica Di Nicolantonio, Ramon Salazar, Davide
Zecchin, Roderick L. Beijersbergen, Alberto Bardelli, and René Bernards. 2012a.
"Unresponsiveness of Colon Cancer to BRAF(V600E) Inhibition through Feedback Activation of
EGFR." *Nature* 483 (7387): 100–103.
Regot, Sergi, Jacob J. Hughey, Bryce T. Bajar, Silvia Carrasco, and Markus W. Covert. 2014. "High-
Sensitivity Measurements of Multiple Kinase Activities in Live Single Cells." *Cell* 157 (7): 1724–
34.
Rusch, V., J. Baselga, C. Cordon-Cardo, J. Orazem, M. Zaman, S. Hoda, J. McIntosh, J. Kurie, and E.
Dmitrovsky. 1993. "Differential Expression of the Epidermal Growth Factor Receptor and Its
Ligands in Primary Non-Small Cell Lung Cancers and Adjacent Benign Lung." *Cancer Research*
53 (10 Suppl): 2379–85.
Sahin, Umut, Gisela Weskamp, Kristine Kelly, Hong-Ming Zhou, Shigeki Higashiyama, Jacques
Peschon, Dieter Hartmann, Paul Saftig, and Carl P. Blobel. 2004. "Distinct Roles for ADAM10
and ADAM17 in Ectodomain Shedding of Six EGFR Ligands." *The Journal of Cell Biology* 164
(5): 769–79.
Sampson, Josephina, Mark W. Richards, Jene Choi, Andrew M. Fry, and Richard Bayliss. 2021.
"Phase-Separated Foci of EML4-ALK Facilitate Signalling and Depend upon an Active Kinase
Conformation." *EMBO Reports*, October, e53693.
Sasaki, Takaaki, Jussi Koivunen, Atsuko Ogino, Masahiko Yanagita, Sarah Nikiforow, Wei Zheng,
Christopher Lathan, et al. 2011. "A Novel ALK Secondary Mutation and EGFR Signaling Cause
Resistance to ALK Kinase Inhibitors." *Cancer Research* 71 (18): 6051–60.
Shaw, Alice T., Peggy P. Hsu, Mark M. Awad, and Jeffrey A. Engelman. 2013. "Tyrosine Kinase Gene
Rearrangements in Epithelial Malignancies." *Nature Reviews. Cancer* 13 (11): 772–87.
Siegel, Rebecca L., Kimberly D. Miller, Hannah E. Fuchs, and Ahmedin Jemal. 2021. "Cancer
Statistics, 2021." *CA: A Cancer Journal for Clinicians* 71 (1): 7–33.
Solomon, Benjamin J., Benjamin Besse, Todd M. Bauer, Enriqueta Felip, Ross A. Soo, D. Ross
Camidge, Rita Chiari, et al. 2018. "Lorlatinib in Patients with ALK-Positive Non-Small-Cell Lung
Cancer: Results from a Global Phase 2 Study." *The Lancet Oncology* 19 (12): 1654–67.
Straussman, Ravid, Teppei Morikawa, Kevin Shee, Michal Barzily-Rokni, Zhi Rong Qian, Jinyan Du,
Ashli Davis, et al. 2012. "Tumour Micro-Environment Elicits Innate Resistance to RAF Inhibitors
through HGF Secretion." *Nature* 487 (7408): 500–504.
Sukhatme, V. P., S. Kartha, F. G. Toback, R. Taub, R. G. Hoover, and C. H. Tsai-Morris. 1987. "A
Novel Early Growth Response Gene Rapidly Induced by Fibroblast, Epithelial Cell and
Lymphocyte Mitogens." *Oncogene Research* 1 (4): 343–55.
Takeuchi, Kengo, Manabu Soda, Yuki Togashi, Ritsuro Suzuki, Seiji Sakata, Satoko Hatano, Reimi
Asaka, et al. 2012. "RET, ROS1 and ALK Fusions in Lung Cancer." *Nature Medicine* 18 (3):
378–81.
Tani, Tetsuo, Hiroyuki Yasuda, Junko Hamamoto, Aoi Kuroda, Daisuke Arai, Kota Ishioka, Keiko
Ohgino, et al. 2016. "Activation of EGFR Bypass Signaling by TGFα Overexpression Induces
Acquired Resistance to Alectinib in ALK-Translocated Lung Cancer Cells." *Molecular Cancer*
Therapeutics 15 (1): 162–71.
Tanimoto, Azusa, Tadaaki Yamada, Shigeki Nanjo, Shinji Takeuchi, Hiromichi Ebi, Kenji Kita, Kunio
Matsumoto, and Seiji Yano. 2014. "Receptor Ligand-Triggered Resistance to Alectinib and Its
Circumvention by Hsp90 Inhibition in EML4-ALK Lung Cancer Cells." *Oncotarget* 5 (13): 4920–

28.

Thedieck, Kathrin, Birgit Holzwarth, Mirja Tamara Prentzell, Christopher Boehlke, Kathrin Kläsener,
Stefanie Ruf, Annika Gwendolin Sonntag, et al. 2013. "Inhibition of MTORC1 by Astrin and
Stress Granules Prevents Apoptosis in Cancer Cells." *Cell* 154 (4): 859–74.

Toettcher, J. E., O. D. Weiner, and W. A. Lim. 2013. "Using Optogenetics to Interrogate the Dynamic
Control of Signal Transmission by the Ras/Erk Module." *Cell* 155 (6): 1422–34.

Tulpule, Asmin, Juan Guan, Dana S. Neel, Hannah R. Allegakoen, Yone Phar Lin, David Brown, Yu-
Ting Chou, et al. 2021. "Kinase-Mediated RAS Signaling via Membraneless Cytoplasmic Protein
Granules." *Cell*, April. https://doi.org/10.1016/j.cell.2021.03.031.

Turke, Alexa B., Youngchul Song, Carlotta Costa, Rebecca Cook, Carlos L. Arteaga, John M. Asara,
and Jeffrey A. Engelman. 2012. "MEK Inhibition Leads to PI3K/AKT Activation by Relieving a
Negative Feedback on ERBB Receptors." *Cancer Research* 72 (13): 3228–37.

Vaishnavi, Aria, Laura Schubert, Uwe Rix, Lindsay A. Marek, Anh T. Le, Stephen B. Keysar,
Magdalena J. Glogowska, et al. 2017. "EGFR Mediates Responses to Small-Molecule Drugs
Targeting Oncogenic Fusion Kinases." *Cancer Research* 77 (13): 3551–63.

Vander Velde, Robert, Nara Yoon, Viktoriya Marusyk, Arda Durmaz, Andrew Dhawan, Daria
Miroshnychenko, Diego Lozano-Peral, et al. 2020. "Resistance to Targeted Therapies as a
Multifactorial, Gradual Adaptation to Inhibitor Specific Selective Pressures." *Nature
Communications* 11 (1): 2393.

Voena, C., F. Di Giacomo, E. Panizza, L. D'Amico, F. E. Boccalatte, E. Pellegrino, M. Todaro, et al.
2013. "The EGFR Family Members Sustain the Neoplastic Phenotype of ALK+ Lung
Adenocarcinoma via EGR1." *Oncogenesis* 2 (April): e43.

Weinstein, I. Bernard. 2002. "Cancer. Addiction to Oncogenes--the Achilles Heal of Cancer." *Science*.

Wilson, Maxwell Z., Pavithran T. Ravindran, Wendell A. Lim, and Jared E. Toettcher. 2017. "Tracing
Information Flow from Erk to Target Gene Induction Reveals Mechanisms of Dynamic and
Combinatorial Control." *Molecular Cell* 67 (5): 757-769.e5.

Wilson, Timothy R., Jane Fridlyand, Yibing Yan, Elicia Penuel, Luciana Burton, Emily Chan, Jing Peng,
et al. 2012. "Widespread Potential for Growth-Factor-Driven Resistance to Anticancer Kinase
Inhibitors." *Nature* 487 (7408): 505–9.

Wong, S. T., L. F. Winchell, B. K. McCune, H. S. Earp, J. Teixidó, J. Massagué, B. Herman, and D. C.
Lee. 1989. "The TGF-Alpha Precursor Expressed on the Cell Surface Binds to the EGF
Receptor on Adjacent Cells, Leading to Signal Transduction." *Cell* 56 (3): 495–506.