Sensory target detection at local and global time scales dissociates modality-specific and

supramodal dynamics along a cortical hierarchy

Maria Niedernhuber[1,2,*], Federico Raimondo[3,5,6,*], Jacobo D. Sitt[7,**], & Tristan A. Bekinschtein[1,**]

[1] Cambridge Consciousness and Cognition Lab, Department of Psychology, University of

Cambridge, UK

[2] Body, Self, and Plasticity Lab, Department of Psychology, University of Zurich,

Switzerland

[3] Brain and Spine Institute, Paris, France

[4] National Institute of Health and Medical Research, Paris, France

[5] Institute of Neuroscience and Medicine, Brain & Behaviour (INM-7), Research Centre

Jülich, Jülich, Germany

[6] Institute of Systems Neuroscience, Medical Faculty, Heinrich Heine University Düsseldorf,

Düsseldorf, Germany

[7] Sorbonne Université, Institut du Cerveau - Paris Brain Institute - ICM, Inserm, CNRS,

APHP, Hôpital de la Pitié Salpêtrière, Paris, France. Université de Paris, Paris, France

Author Note

This project was funded by BJA/RCoA WKR0-2018-0060 to TAB. J.D.S. is supported by the Paris Brain Institute (France) and the program "Investissements d'avenir" ANR-10- IAIHU-06, by the CARNOT maturation grant, Title: Con&Heart to JDS, by the NeuroCatalyst grant, Title: NeuroEnviroTech, as well as a the PerBrain study funded by the European Union (ERA PerMed JTC2019 "PerBrain").

* Maria Niedernhuber and Federico Raimondo share first authorship.** Tristan Bekinschtein and Jacobo D. Sitt share last authorship.

The authors made the following contributions. Maria Niedernhuber: Conceptualization, Data collection, Data analysis, Writing - Original Draft Preparation, Writing - Review & Editing; Federico Raimondo: Conceptualization, Data collection, Data analysis, Writing - Review & Editing; Jacobo D. Sitt: Conceptualization, Writing - Review & Editing; Tristan A. Bekinschtein: Conceptualization, Writing - Review & Editing.

Correspondence concerning this article should be addressed to Maria Niedernhuber, Craik-Marshall Building, 20 Downing Pl, Cambridge CB2 1QW, United Kingdom. E-mail: mn473@cam.ac.uk

Abstract

How do cortical networks support the detection of irregular sensory events for different senses? Here we reconcile two views of cortical function according to which supramodal processing is either deferred to associative cortices in higher levels of a unimodal-to-supramodal cortical hierarchy or features across all cortical processing stages. In a series of hierarchical multisensory target detection studies using EEG, we applied a temporal cross-decoding approach to reveal the cortical dynamics elicited by rule-based global and feature-based local sensory deviations within and between the visual, somatosensory and auditory modality. We uncovered simple feature-based sensory deviations presented in temporal vicinity to a monotonous input stream triggered an MMN-like local negativity which decayed quickly and early in different sensory modalities. Conversely, complex rule-based targets tracked across time evoked a P3b-like global effect which generalised across a late time window. Ultimately, a series of temporal cross-modality decoding analyses demonstrated that global effects are maintained in supramodal networks whereas local effects propagate along modality-specific cortical networks and evolve into short-lived supramodal activity. Taken together, each sense implements a cortical hierarchy which orchestrates target detection responses operating on local and global time scales at successive processing stages. Converging evidence from different analyses and studies, we show that cortical organisation largely follows a gradient in which modality-specific processes dominate local responses and higher-order processes encode abstract supramodal information fed forward from modality-specific cortices. In conclusion, the cortex implements a hierarchy which hosts increasingly supramodal processes in each level rather than a strict hierarchy which delays supramodal function to associative cortices.

Keywords: attention, target detection, P300, Mismatch Negativity, predictive coding
Word count: 7552 words

Sensory target detection at local and global time scales dissociates modality-specific and supramodal dynamics along a cortical hierarchy

Introduction

The ability to detect deviant sensory events in a stream of predictable stimuli is crucial for adaptive behaviour. To enable this, each sense relies on a dedicated hierarchical system of cortices spanning from primary sensory to associative and frontal areas (Allen et al., 2016a; Chao, Takaura, Wang, Fujii, & Dehaene, 2018; Chennu et al., 2013; Cornella, Leung, Grimm, & Escera, 2012; J. Murray et al., 2014; Rao & Ballard, 1999; Wacongne et al., 2011). Are there unified principles of cortical organisation which govern target detection processes across sensory systems? A prominent proposal states each perceptual system encompasses a series of cortices arranged along a simple-to complex hierarchy according to which successive levels encode increasingly abstract sensory information (de Lange, Heilbron, & Kok, 2018; Felleman & Van Essen, 1991; Kiebel, Daunizeau, & Friston, 2008; J. Murray et al., 2014; Rao & Ballard, 1999). Indeed, there is strong evidence across different sensory modalities that neurons in primary and secondary sensory areas respond to simple features such as shapes in the primary visual cortex, pitch in the primary auditory cortex, pressure in the primary somatosensory cortex), whereas frontal regions encode more complex information such as visual landscapes, sound patterns and tactile objects (Allen et al., 2016a; Bekinschtein et al., 2009; Fardo et al., 2017; Hochstein & Ahissar, 2002; Hubel & Wiesel, 1962; Maunsell & Essen, 1983; C. Murray, 2004; Nelken & Bar-Yosef, 2008; Rao & Ballard, 1999; Riesenhuber & Poggio, 1999; Van Essen & Maunsell, 1983).

Converging evidence from target detection studies in different sensory modalities supports the notion that target detection processes are organised along a simple-to-complex gradient from modality-specific to frontal cortical regions. A rich body of literature identified the prefrontal cortex as a key site for complex rule-visual search

tasks in which simple targets are detected was found to rely on early visual regions based perceptual learning of sensory targets (Miller, 2009). In the visual system, target detection involving the spatial conjunction of multiple visual features has been shown to recruit attention-dependent frontoparietal processes (Donner et al., 2000; Kristjánsson, Vuilleumier, Schwartz, Macaluso, & Driver, 2007). Conversely, early visual cortices support visual search tasks in which simple targets are detected (Ester, Serences, & Awh, 2009). In the auditory domain, a series of studies mapping the neural basis of simple and complex auditory targets provide evidence for a hierarchical organisation of the auditory system. Higher-order frontal cortices were found to process complex targets such as auditory patterns, whereas simple pitch deviations depended on activity in early auditory cortices (Bekinschtein et al., 2009; Chennu et al., 2016; Cornella et al., 2012; Uhrig, Dehaene, & Jarraya, 2014; Wacongne et al., 2011). Taken together, these results raise the question whether the detection of complex targets relies on a common frontal activation pattern in different sensory modalities, while the detection of simple targets recruits modality-specific areas.

We propose that cortical signatures of sensory targets are arranged along a gradient of supramodality according to which responses to complex rule-based targets largely rely on sustained late supramodal activation patterns while modality-specific processing dominates responses to simple targets (pre-registered https://osf.io/3mqvy/). Our approach exploits differences in the susceptibility of electrophysiological responses to bottom-up and top-down variables to dissociate their neural dynamics in different levels of the cortical hierarchy. Sensory targets are followed by two cortical responses which can be measured using EEG: the Mismatch Negativity and the P300 complex (consisting of two components called P3a and P3b). The MMN is an early local negativity associated with sensory change detection in a temporal proximity to the expected input (Mäntysalo & Näätänen, 1987; Risto Näätänen, Tervaniemi, Sussman, Paavilainen, & Winkler, 2001; Shinozaki, Yabe, Sutoh, Hiruma, & Kaneko, 1998; Wacongne, Changeux, & Dehaene,

2012). Originally discovered in the auditory modality (R. Näätänen, Simpson, & Loveless,

1982), the MMN can be detected in different sensory modalities (Allen et al., 2016b;

Amado & Kovács, 2016; Czigler & Csibra, 1992; Czigler et al., 2006; Shen, Smyk, Meltzoff,

& Marshall, 2018). Consistent with its early appearance, the MMN has been found to

originate in primary sensory cortices dedicated to the modality in which the sensory

irregularity was generated Hamm & Yuste (2016). Canonically considered to be

pre-attentive (Tiitinen, May, Reinikainen, & Näätänen, 1994), the MMN is modulated by

attention but resilient to distraction (Auksztulewicz & Friston, 2015; Chennu et al., 2013).

The P3b is a late distributed positive potential which marks the detection of sensory

targets and can be elicited in a range of sensory modalities (Polich, 2007; Yamaguchi &

Knight, 1991b, 1991a), In contrast to the MMN, the P3b seems to require an active

memory to track the sensory context in which irregularities are embedded. The P3b also

also lasts longer (Pegado et al., 2010) and disappears without attention to the sensory

irregularity (Katayama & Polich, 1998; Polich, 2007; Squires, Petuchowski, Wickens, &

Donchin, 1977). Based on these properties, the auditory MMN and P3b can be located as

cortical responses to simple versus complex sensory targets in lower-order auditory cortical

regions and higher-order frontoparietal levels of the cortical hierarchy respectively

(Bekinschtein et al., 2009; Chao et al., 2018; Wacongne et al., 2011).

Here we use multisensory versions of a hierarchical sensory target detection paradigm

("local-global paradigm") to obtain a P3b-like global top-down response as well as a

MMN-like local bottom-up response. The local-global paradigm achieves this by

manipulating the complexity of the sensory context in which a stimulus change appears

(Bekinschtein et al., 2009). We hypothesised that the global effect as a top-down-driven

sensory novelty response to complex targets in higher-order cortices might share neural

signatures between the senses. In contrast, the local effect which can be regarded as a

bottom-up response to simple targets might be supported by early localised

modality-specific activity in primary sensory cortices. In two different multisensory

variants of the local-global paradigm, we uncovered commonalities between the cortical dynamics of responses to feature-based local and rule-based global sensory deviations using a temporal decoding approach (King & Dehaene, 2014). We show that classifiers trained to decode global effects in a specific sensory modality were able to decode other global effects in a late extended time window. In contrast, local effects shared few short-lived cortical signatures between the senses.

Results

Hierarchically nested sensory deviations elicit local and global effects in different sensory modalities

We investigated commonalities in cortical responses to rule-based global and feature-based local sensory deviations using multisensory versions of the local-global paradigm. In this task, expectations about sensory stimuli are selectively violated at two hierarchical levels: the local and global level. To achieve this, the paradigm varies the temporal extent of the sensory context in which a deviant stimulus is presented (Bekinschtein et al., 2009). Sensory deviations are introduced either at a local level (by occasionally varying the last stimulus in a stimulus group) or at a global level (by introducing rare stimulus groups in blocks of frequently appearing stimulus groups), as shown in Figure 1. Capitalising on differences in the responsiveness of cortical processes to top-down manipulation, this design enables us to extract local and global cortical markers which can be located at successive hierarchical levels in the cortex. MMN-like local effects can be extracted from feature-based sensory deviations and manifest as an early amplitude enhancement when locally standard trials (groups of five identical ipsilateral stimuli) are contrasted with locally deviant trials (groups of four identical ipsilateral stimuli followed by a contralateral input). Rule-based sensory deviations evoke P3b-like global effects which are established by contrasting ERP time courses extracted from globally deviant trials with

standard trials in a late time window between 250-600 ms after onset of the last stimulus in

a trial. As such, the global effect is associated with top-down target detection processes

which require the retention of the preceding sensory input stream in memory in different

sensory systems. To examine commonalities in local or global cortical responses between

different sensory modalities, we developed two versions of the local-global paradigm in

which sensory inputs are sampled from different sensory modalities. In the bimodal version

of the paradigm, we presented blocks of auditory or somatosensory hierarchical sensory

stimulation. The trimodal version of the paradigm contained unisensory blocks of either

visual, somatosensory or auditory sensory inputs.

In a first step, we established that hierarchical manipulations of sensory context elicit

a MMN-like local effect and a P3b-like global effect in the auditory, somatosensory and

visual modality in both experiments. To that end, we performed cluster-based permutation

of ERP amplitude time courses (Maris & Oostenveld, 2007) displayed in Figure 1.

Replicating earlier findings (Bekinschtein et al., 2009; Chennu et al., 2013), the auditory

local effect in the bimodal paradigm manifested as a frontotemporal bipolar two-peak

difference wave (cluster t = -12000, p < 0.001) and a one-peak difference wave in the

trimodal paradigm (cluster t = -5396.6, p = 0.001). A somatosensory local effect emerged

as a central negativity between ~50-150 ms in the bimodal paradigm (cluster t = -752.31, p

= 0.003) and as a two-peak negativity between ~100-350 ms the trimodal paradigm

(cluster t = -3298.4, p = 0.004). We also identified a visual local effect as a negativity in a

mid-range time window between ~100-350 ms (cluster t = -6178.9, p < 0.001) shown in

Figure 1.

At a global level, a comparison of globally deviant and standard trials revealed a

positive difference wave in a time window from ~250 ms until the end of the trial regardless

of sensory modality (with some shifts in onset of the effect). In the bimodal (cluster t =

22143, p < 0.001) and trimodal study (cluster t = 27934, p < 0.001), we revealed an

auditory global effect as a positive deflection with a posterior distribution. A

somatosensory global effect presented with a similar posterior distribution in both the bimodal (cluster t = 18351, p < 0.001) and trimodal study (cluster t = 21421, p < 0.001). Ultimately, a visual global effect emerged as a positive difference wave in a relatively late time window from ~400 ms until the end of the trial (cluster t = 12857, p = 0.001) shown in Figure 1. These results show that complex targets which require the conscious tracking of sensory patterns across time elicit a global effect in different sensory modalities. Our finding that the global effect manifests as a large, late and posterior positive deflection regardless of sensory modality complements previous studies which characterise the related P300 as a late positivity (Bekinschtein et al., 2009; Bledowski et al., 2004; Chennu et al., 2013; Walz et al., 2013), Taken together, our results show that the functional dissociation of local and global effects is a supramodal property of target detection systems in different sensory domains.

Cortical responses to rule-based but not feature-based sensory targets are sustained across time in each sensory modality

Previous work has shown that functional differences between cortical responses to auditory rule-based and feature-based targets are reflected in the extent to which they are maintained in auditory networks. Cortical activation patterns in response to auditory rule-based targets are sustained in time, whereas feature-based auditory deviations decay quickly (King & Dehaene, 2014). Is the temporal evolution of cortical target detection responses a property common to different sensory domains? We examined whether global effects are linked to a sustained cortical activation pattern whereas local effects are supported by short-lived activity regardless of sensory modality. We employed temporal decoding to characterise neural activation patterns elicited by local and global effects in different sensory modalities. In short, temporal decoding is a machine learning approach used to characterise the temporal evolution of cortical activation patterns linked to sensory events. A classifier trained at a time point t is not only tested at t but at all other

Figure 1. **Experimental design and ERP results.**   Block type design for the local-global paradigm used in the bimodal and trimodal study (A). Cluster-based permutation test results for ERP amplitude differences showing that a local and global effect can be obtained in different sensory modalities for the trimodal (B) and the bimodal study (C). On the left, each panel shows ERP amplitude time courses of the corresponding deviant-standard condition pair. Next to the ERP voltage time course plot, the corresponding topographical map is shown. Time periods in which a cluster-based permutation test identified a difference between both conditions are highlighted in red in the ERP time course plot and delineated with a black line on the topographical map. The time point at which the difference between both conditions is maximal is delineated with a dotted red line in the ERP amplitude time course plot. The electrode at which this difference was obtained is marked with an orange dot on the topographical map.

remaining time points. This leads to a temporal generalisation matrix of classification performance scores. The shape of the temporal generalisation matrix offers insights into the temporal dynamic of cognitive operations and their cortical generators (Dehaene & King, 2016).

We observed that local effects could be decoded along the diagonal in a mid-range time window for each sensory modality (Figure 2). Across different contrasts, local effects were found to decay quickly. This finding is mostly consistent with a serial activation of different cortices dedicated to the sensory modality in which the local effect was applied (King, Gramfort, Schurger, Naccache, & Dehaene, 2014). At a local level, the visual local effect was maintained briefly between 200-400 ms (mean AUC = 0.511 ± 0.029, max. AUC = 0.61 at 268 ms training time and 268 ms testing time, mean cluster t = 3.327, p < 0.05). In the trimodal study, the somatosensory local effect (mean AUC = 0.511 ± 0.024, max. AUC = 0.564 at 316 ms training time and 312 ms testing time, mean cluster t = 2.91, p < 0.05) and the auditory local effect (mean AUC = 0.509 ± 0.028, max. AUC = 0.65 at 212 ms training time and 212 ms testing time, mean cluster t = 2.599, p < 0.05) were maintained for ~100 ms from ~200 ms. In the bimodal study, the somatosensory effect was best decoded between 200-300 ms (mean AUC = 0.516 ± 0.026, max. AUC = 0.629 at 208 ms training time and 208 ms testing time, mean cluster t = 4.472 ± 7.53, p < 0.05). In this study, temporal decoding revealed a classification score matrix consistent with two distinct processes underpinning the auditory local effect. From ~100 ms, the auditory local effect can be decoded along the diagonal which suggests a serial propagation of cortical activity along the auditory cortical hierarchy. However, cortical activity is sustained for ~150 ms from ~200 ms (mean AUC = 0.507 ± 0.04, max. AUC = 0.714 at 188 ms training time and 188 ms testing time, mean cluster t = 2.11 ± 13.151, p < 0.05). Although classification scores for the auditory and somatosensory local effect differed from chance across an extended time window, classifiers performed only slightly better than chance from ~350 ms for both effects.

²⁴³   At a global level, we found that classifiers trained to distinguish deviant and standard

²⁴⁴ trials from ~200 ms generalised across other time samples in the remaining trial window

²⁴⁵ regardless of which sensory modality was tested (Figure 2). This decoding procedure led to

²⁴⁶ a rectangular classification score matrix for each global contrast. In line with earlier ERP

²⁴⁷ time course comparison results (Figure 1), the visual global effect manifested relatively late

²⁴⁸ from ~400 ms (mean AUC = 0.512 ± 0.06, max. AUC = 0.656 at 516 ms training time and

²⁴⁹ 520 ms testing time, mean cluster t = 3.19, p < 0.05). In the trimodal study, the

²⁵⁰ somatosensory global effect appeared from ~350 ms ((mean AUC = 0.52 ± 0.048, max.

²⁵¹ AUC = 0.644 at 512 ms training time and 508 ms testing time, mean cluster t = 5.561, p

²⁵² < 0.05)) and the auditory global effect from ~200 ms (mean AUC = 0.544 ± 0.069, max.

²⁵³ AUC = 0.69 at 532 ms training time and 524 ms testing time, mean cluster t = 12.502, p

²⁵⁴ < 0.05). In the bimodal study, temporal generalisation was found relatively early from

²⁵⁵ ~150 ms for both the somatosensory (mean AUC = 0.56 ± 0.057, max. AUC = 0.697 at

²⁵⁶ 372 ms training time and 344 ms testing time, mean cluster t = 19.317 ± 19.074, p < 0.05)

²⁵⁷ and auditory global effect (mean AUC = 0.599 ± 0.048, max. AUC = 0.695 at 432 ms

²⁵⁸ training time and 428 ms testing time, mean cluster t = 33.062 ± 17.556, p < 0.05).

²⁵⁹ Cluster-based permutation tests comparing differences in classification scores between the

²⁶⁰ local and global effect show that the global effect generalises in a late time window whereas

²⁶¹ the local effect does not (Figure 2). Collectively, our results indicate that cortical

²⁶² activation associated with the global effect starts no earlier than ~150 ms after onset of the

²⁶³ last stimulus in a trial and is sustained over time until the trial ends. However, we

²⁶⁴ observed some shifts in the onset latency between different global effects with the visual

²⁶⁵ global effect not appearing before 400 ms. Despite these shifts, this pattern suggests that a

²⁶⁶ single cortical system is active in that time window (King & Dehaene, 2014). This finding

²⁶⁷ leaves the question open whether this system is dedicated to a specific sensory modality or

²⁶⁸ shared between different senses.

Figure 2. **Temporal generalisation analysis within sensory modalities.** Panels display temporal generalisation results for the local (left) and global effect (right) in the trimodal (A) and bimodal dataset (B). Each panel shows results of a temporal generalisation analysis in which a classifier is trained to distinguish deviant and standard trials at each time point and then tested on all remaining time points in a trial. Classification scores are displayed on a red-to-blue gradient. On top of each matrix, adjacent classification scores different from chance (0.5) in a cluster-based permutation test are highlighted with a black line. Next to each matrix, we show a series of time courses of classification scores produced by classifiers trained at the specified time point. On top of each matrix, we plotted the classification score time course for a decoding procedure in which a classifier is trained and tested on deviant-standard pairs at the same time point (corresponding to the matrix diagonal). Classification scores different from chance are highlighted with a blue shade. On the right side, we show results of a cluster-based permutation test contrasting classification score matrices of the corresponding local and global effect. Each subplot shows the classification score time courses of a classifier trained to separate global deviant-standard pairs at the specified time point and tested across the remaining time window, and its local counterpart. Time periods in which local and global effect classification time courses differ are highlighted in red shade.

A supramodal cortical signature sustained in time supports global but not local

effects

Here we provide evidence for the hypothesis that cortical hierarchies dedicated to

each sense are organised along a gradient of supramodality. Building up on our finding that

global effects are associated with a sustained late cortical activation pattern, we further

demonstrate that this sustained pattern is shared between different sensory modalities. We

also show that local effects in different sensory modalities rely on few, if any, common

cortical signatures in comparison to global effects.

Temporal decoding was employed to examine whether cortical responses share neural

dynamics between sensory modalities. To identify common activity patterns linked to

evoked responses, temporal generalisation analysis can be applied in two ways: Classifiers

can be trained to separate a deviant-standard condition pair in a target sensory modality

at a time point t and tested on deviant-standard condition pairs in a different sensory

modality across all time points in a trial. Alternatively, a classifier can be trained to

separate deviant-standard condition pairs when they are pooled across all sensory

modalities separately at a global and a local level. We used a 5-fold stratified

cross-validation approach in which classifiers were trained on 4/5 of the data and tested on

the remaining 1/5 for both analyses (see Methods for details).

We initially trained a classifier to separate globally deviant from standard trials when

trials for each condition pair are pooled across sensory modalities. This procedure revealed

temporal generalisation in a late time window across different global contrast in both

datasets (Figure 3). We provide evidence for shared activity supporting the auditory and

visual global effect from ~400 ms regardless of whether classifiers were trained on the visual

and tested on the auditory contrast (mean AUC = 0.517 ± 0.065, max. AUC = 0.663 at

452 ms training time and 572 ms testing time, mean cluster t = 4.622, p < 0.05) or vice

versa (mean AUC = 0.508 ± 0.06, max. AUC = 0.653 at 676 ms training time and 496 ms

testing time, mean cluster t = 3.37, p < 0.05). A similar pattern was found for

somatosensory and visual effects when classifiers were trained on visual and tested on

somatosensory deviant-standard pairs (mean AUC = 0.51 ± 0.053, max. AUC = 0.636 at

540 ms training time and 464 ms testing time, mean cluster t = 3.202, p < 0.05) or vice

versa (mean AUC = 0.51 ± 0.067, max. AUC = 0.675 at 580 ms training time and 540 ms

testing time, mean cluster t = 2.945, p < 0.05). Shared activity between the auditory and

somatosensory global effect was found from ~350 ms when classifiers were trained on the

auditory and tested on the somatosensory contrast (mean AUC = 0.531 ± 0.046, max.

AUC = 0.644 at 424 ms training time and 580 ms testing time, mean cluster t = 10.207, p

< 0.05) or vice versa (mean AUC = 0.532 ± 0.064, max. AUC = 0.688 at 576 ms training

time and 432 ms testing time, mean cluster t = 10.256, p < 0.05). This finding was

replicated in the bimodal dataset in a more extensive time window from ~150 ms when

classifiers were trained on the somatosensory and tested on the auditory effect (mean AUC

= 0.568 ± 0.059, max. AUC = 0.674 at 188 ms training time and 160 ms testing time,

mean cluster t = 22.265 ± 19.849, p < 0.05) and vice versa (mean AUC = 0.572 ± 0.049,

max. AUC = 0.663 at 292 ms training time and 348 ms testing time, mean cluster t =

24.748 ± 17.699, p < 0.05). Taken together, these findings show that there is shared

activation between global effects regardless of which sensory modality is used for training

and testing. The global effect consistently manifests in a rectangular classification score

matrix which suggests that common activation is maintained in cortical networks (with

some temporal shifts in onset times).

Interestingly, temporal decoding of local effects revealed short-lived temporal

generalisation around ~200 ms in both datasets. In the trimodal dataset, short-lived shared

representations from ~200-350 ms were found to be associated with local effects across all

comparisons: auditory to somatosensory (mean AUC = 0.511 ± 0.024, max. AUC = 0.587

at 208 ms training time and 304 ms testing time, mean cluster t = 4.07, p < 0.05),

somatosensory to auditory (mean AUC = 0.509 ± 0.033, max. AUC = 0.612 at 296 ms

training time and 208 ms testing time, mean cluster t = 2.952, p < 0.05), auditory to

visual (mean AUC = 0.506 ± 0.029, max. AUC = 0.615 at 196 ms training time and 268

ms testing time, mean cluster t = 2.075, p < 0.05), visual to auditory (mean AUC = 0.52

± 0.017, max. AUC = 0.588 at 264 ms training time and 196 ms testing time, mean cluster

t = 6.587, p < 0.05), somatosensory to visual (mean AUC = 0.506 ± 0.027, max. AUC =

0.597 at 284 ms training time and 272 ms testing time, mean cluster t = 1.812, p < 0.05)

and visual to somatosensory (mean AUC = 0.497 ± 0.018, max. AUC = 0.546 at 256 ms

training time and 216 ms testing time, mean cluster t = -0.897, p < 0.05)). In the bimodal

dataset, we found evidence for shared representations from ~200 ms when classifiers were

trained on somatosensory and tested on auditory deviant-standard pairs (mean AUC =

0.515 ± 0.03, max. AUC = 0.677 at 208 ms training time and 176 ms testing time, mean

cluster t = 4.905 ± 9.025, p < 0.05) and vice versa (mean AUC = 0.508 ± 0.024, max.

AUC = 0.639 at 172 ms training time and 208 ms testing time, mean cluster t = 2.991 ±

8.745, p < 0.05).

Our result demonstrates that local effects are supported by supramodal transient

activation patterns starting from ~200 ms. Although our finding is compatible with the

idea that local effects are processed in cortical hierarchies dedicated to the sensory

modality in which the stimulation is applied, we provide evidence for some overlap in

higher-order or associative regions between these hierarchies. In sum, these results support

the hypothesis that global effects share sustained neural activation patterns between the

senses while local effects share fewer, if any, activity (Figure 3)

To corroborate these findings, we use temporal decoding to examine the temporal

evolution of cortical activity when global deviant-standard pairs and local deviant-standard

pairs are each pooled across sensory modalities. Our results show that the local effect is

associated with some temporal generalisation from ~180-250 ms in both the bimodal (mean

AUC = 0.508 ± 0.025, max. AUC = 0.649 at 204 ms training time and 204 ms testing

time, mean cluster t = 2.747 ± 8.431, p < 0.05)and the trimodal dataset (mean AUC =

Figure 3. **Temporal generalisation analysis within and between sensory modalities.** Each panel shows results from a series of temporal generalisation analyses decoding the local effect (left) and the global effect (right) in the trimodal dataset (A) and bimodal dataset (B). Each matrix shows results of a temporal generalisation analysis in which a classifier is trained to distinguish deviant and standard trials at each time point in the training modality and then tested on all remaining time points in a trial in the testing modality. For each experiment, this leads to an n-by-n matrix with n being the number of sensory modalities tested in a study (3 for the trimodal and 2 for the bimodal study). Temporal generalisation matrices placed along the diagonal of a panel show results from a decoding analysis performed within a sensory modality and are highlighted with a purple frame. The remaining temporal generalisation matrices show results from a decoding procedure in which a classifier is trained to distinguish deviant and standard trials corresponding to the sensory modality indicated next to each row and tested on the sensory modality corresponding to the column label. AUC-ROC classification scores are shown on a red-to-blue gradient. Classification score clusters which were found to be different from chance in a cluster-based permutation test are delineated with a black line. 0 ms describes the onset of the last stimulus in a trial.

0.513 ± 0.026, max. AUC = 0.574 at 220 ms training time and 220 ms testing time, mean cluster t = 3.874, p < 0.05). For local effects, our result is most consistent with the involvement of a series of modality-specific neural generators in early levels of the cortical hierarchy and a contribution of supramodal regions in later stages. Again, temporal decoding revealed sustained shared activation starting from ~150 ms until trial end for the global effect in the bimodal dataset (mean AUC = 0.579 ± 0.057, max. AUC = 0.667 at 340 ms training time and 336 ms testing time). In the trimodal dataset, shared activation between global effects extended across the complete time window. We found evidence for sustained activity from ~350 until trial end. Our results also reveal a rectangular classification score cluster which differs slightly but significantly from chance from the onset of the last stimulus in a trial to ~250 ms (mean AUC = 0.524 ± 0.057, max. AUC = 0.657 at 520 ms training time and 544 ms testing time, mean cluster t = 7.354, p < 0.05). Based on work by King et al. (King & Dehaene, 2014), these temporal generalisation results suggest that there is a common generator for global effects in different sensory modalities. In contrast, multiple neural generators support the local effect. For the local effect, our evidence shows that these generators are modality-specific in early stages and supramodal in later stages of cortical processing (Figure 4). Ultimately, we assessed whether there are differences in decoding strength between the local and global effect in a comparison of maximum classification scores in a cluster drawn from all 14 temporal decoding analyses. Peak decoding performance was found to differ between local and global effects using a Mann-Whitney U test (U = 31, p < 0.001).

Figure 4. **Temporal decoding of combined sensory modalities.** In a temporal decoding analysis, we pooled deviant trials and standard trials regardless of sensory modality and tested whether a classifier trained to discriminate deviant and standard trials at a specific time point might generalise to the remaining time points. Each panel shows the resulting matrix of AUC-ROC classification scores for the global (A) and local effect (B). In each panel, results for the global effect are shown on the right and results for the local effect are displayed on the left. In conditions in which cluster-based permutation was performed, clusters which differ from chance are highlighted with a purple horizontal line for classification performance in intervals of 100 ms and green for diagonal classification performance. 0 ms marks the onset of the last stimulus in a trial. Clusters of AUC-ROC scores which differed from chance were delineated with a dotted line. Each rain cloud plot shows the distribution of maximum classification scores in a cluster drawn from all 14 temporal decoding analyses for the local and global effect as well as corresponding box-and-whisker plots. Significant differences assessed with a Mann-Whitney U Test are highlighted with an asterisk (C).

₃₇₀   **Discussion**

₃₇₁ **Hierarchically nested sensory targets elicit local and global responses across**

₃₇₂ **sensory modalities**

₃₇₃   A long-standing debate in neuroscience revolves around which sensory and perceptual

₃₇₄ processes are supramodal or modality-specific (Cao, Summerfield, Park, Giordano, &

₃₇₅ Kayser, 2019; Driver & Noesselt, 2008; Faivre, Filevich, Solovey, Kühn, & Blanke, 2018;

₃₇₆ Walz et al., 2013). Inspired by the notion that perceptual systems in the cortex are

₃₇₇ hierarchically organised on a simple-to-complex axis (Dürschmid et al., 2016; Kiebel et al.,

₃₇₈ 2008; C. Murray, 2004; Rao & Ballard, 1999), we investigated whether cortical responses to

₃₇₉ sensory targets implemented in successive levels of the cortical hierarchy are ordered along

₃₈₀ a gradient of supramodality. We examined whether cortical responses to sensory targets

₃₈₁ rely more on supramodal activation patterns when sensory targets are complex. Across a

₃₈₂ series of two experiments combining evidence from the somatosensory, visual and auditory

₃₈₃ modality, we first established that selective hierarchically nested divergences of sensory

₃₈₄ inputs can trigger a MMN-like effect at a local level and a P3b-like effect at a global level

₃₈₅ in different sensory modalities. We applied temporal decoding to show that complex

₃₈₆ rule-based targets elicit a sustained global response in each modality which relies on a

₃₈₇ single supramodal network in higher-order cortical networks. Our results demonstrate that

₃₈₈ the prolonged maintenance of cortical activation elicited by the global effect and the serial

₃₈₉ propagation of the local effect are principles of cortical function found across sensory

₃₉₀ modalities. Extending earlier temporal decoding results showing that the auditory global

₃₉₁ effect is maintained in higher-order cortical networks while cortical activity representing

₃₉₂ the auditory local effect is serially propagated along cortical areas (King et al., 2014), we

₃₉₃ demonstrate that cortical hierarchies implement target detection processes which track

₃₉₄ sensory irregularities in hierarchically nested different timescales at successive cortical

₃₉₅ processing stages for each sense. Crucially, our finding that the P3b-like global effect relies

on sustained supramodal cortical signatures while the local effect elicits short-lived responses with few commonalities between the senses supports the notion of a gradient of supramodality underpinning cortical hierarchies but also refutes the idea that early cortical processes are strictly unimodal.

Rule-based sensory targets elicit supramodal and sustained responses in the cortex

Our finding that the P3-like global effect as a response to complex sensory targets is supported by sustained common supramodal activation patterns contributes to a debate around its putative supramodal underpinnings. Some studies investigating cortical activation linked to the auditory and visual P3 suggest a common network including the insula and frontoparietal areas between these senses (Linden et al., 1999; Walz et al., 2013). However, other studies highlight a contribution of higher-order regions in modality-specific cortical hierarchies to the P300 (Bledowski et al., 2004). Our results support the notion that the P3 is partially supported by a supramodal network while at the same time not ruling out contributions of modality-specific processes.

In the recent decade, the global effect has been proposed to be a cortical signal reflecting the conscious processing of incoming sensory stimulation (Bekinschtein et al., 2009). By this view, the global effect reflects recurrent information flow in a global neuronal workspace which maintains cortical signatures to become consciously accessible (Dehaene & Changeux, 2011). Sensory inputs from modality-specific cortices are fed forward to the global neuronal workspace which broadcasts integrated multisensory information from the top down to the levels below (Mashour, Roelfsema, Changeux, & Dehaene, 2020). Our observation that the cortical signatures supporting the global effect are supramodal are consistent with the theory that the global effect marks a supramodal top-down-driven process in which sensory information is amplified for conscious access via allocated attention (Chennu et al., 2013).

Local responses to sensory targets are linked to short-lived modality-specific activation

We discovered a short-lived supramodal contribution to the MMN-like local effect which refutes a classic view according to which the MMN relies on modality-specific mechanisms (Nyman et al., 1990; Pazo-Alvarez, Cadaveira, & Amenedo, 2003). In our study, temporal cross-decoding analyses uncovered short-lived supramodal signatures for the local effect starting from ~200 ms after onset of the last stimulus in a trial. Previous studies demonstrate that local effects are supported by a temporofrontal network in which neuronal messages are propagated forward to the inferior frontal gyrus after initial processing in primary sensory areas which raises the possibility that frontal contributions to the MMN might host supramodal signatures. Both the visual and auditory MMN were found to consist of an earlier component in modality-specific early sensory cortices followed by an attention-modulated late frontal component from ~200 ms after oddball onset (Deouell, 2007; Hedge et al., 2015). Similarly, studies of effective connectivity underpinning the MMN show that the potential is likely supported by a network spanning primary and secondary sensory cortices as well as frontal regions in different sensory modalities (Auksztulewicz & Friston, 2015; Chennu et al., 2016; Fardo et al., 2017; Garrido et al., 2009; Ostwald et al., 2012). Finally, the temporal characteristics of supramodal signatures supporting the local effect are congruent with a contribution of frontal areas linked to attention and target detection (Garrido, Kilner, Stephan, & Friston, 2009). Combined with earlier results, our results suggest that the MMN-like local effect might consist of a modality-specific and a supramodal component. Our result is consistent with the idea that successive layers in the cortical hierarchy might support increasingly supramodal processes.

A gradient of supramodality as a principle of cortical organisation

The canonical view of cortical function states that cortical hierarchies implement a strict unimodal-to-supramodal gradient. According to this view, supramodal processing is deferred to associative and frontal cortices (Felleman & Van Essen, 1991). Mounting evidence demonstrates that multisensory processes are ubiquitous in the cortical hierarchy and occur at all processing stages which refutes the idea that early cortices are strictly unimodal (Driver & Noesselt, 2008; Ghazanfar & Schroeder, 2006). Integrating both views, our findings support the view that the cortex is hierarchically organised along a gradient of supramodality. Feature-based sensory irregularities trigger quickly decaying and early modality-specific processes which dominate early stages of sensory processing. Finally, a sustained late response to rule-based sensory irregularities shared between sensory modalities might reflect a recurrent supramodal process in higher-order cortical areas. However, our finding that local responses evolve into short-lived supramodal activation patterns provide evidence for the notion that early cortical function is not strictly specific to a sensory modality.

Our results are compatible with a predictive coding view of cortical function. Predictive coding states that cortical responses to irregular sensory information reflect a prediction error resulting from a reconciliation of actual sensory inputs and their predictions (Clark, 2013; Friston, 2005; Hohwy, 2012; Rao & Ballard, 1999). From this perspective, local and global effects can be seen as manifestations of prediction errors located at temporally dissociable successive levels of a dedicated cortical hierarchy for each sense (Wacongne et al., 2011). A central idea in predictive coding is that higher-order levels of the cortical hierarchy converge information from different senses forwarded from the levels below to generate predictions about the sensory environment (Clark, 2013; de Lange et al., 2018; Friston, 2005; Hohwy, 2012). This aligns with our result that higher-order cortical responses share supramodal signatures between the senses while

lower-order responses largely rely on modality-specific activation patterns.

Most research on the MMN and P3b as cortical underpinnings of prediction error

signals concentrates on the auditory or visual domain, and comparably less is known about

the somatosensory P3b or MMN (Linden et al., 1999; Ostwald et al., 2012). Earlier studies

employed the local-global paradigm to demonstrate that the MMN-like local effect is

generated in the primary auditory cortex and the global effect relies on activity in

frontoparietal regions (Bekinschtein et al., 2009; Chao et al., 2018; Chennu et al., 2013; El

Karoui et al., 2015; Uhrig et al., 2014; Wacongne et al., 2011). Extending earlier work, we

deliver an integrative framework for cortical responses to sensory targets tracking different

timescales at successive levels of the cortical hierarchy across different sensory modalities.

Materials and methods

Participants

We developed two multisensory versions of the local-global paradigm. In the bimodal

version, separate blocks of somatosensory or auditory expectation violations were

presented. In addition to purely visual, somatosensory and auditory blocks, the trimodal

version of the paradigm encompassed blocks in which trials combined inputs from two

different sensory modalities which could either be visual, auditory or somatosensory. Only

individuals with no history of neurological or psychiatric conditions and no tactile and

auditory impairment were recruited into both studies. In addition, participants with visual

impairments were excluded from participant in the trimodal study. All participants gave

written and informed consent. Data for the bimodal local-global paradigm were collected

at the University of Cambridge and obtained ethical approval from the Department of

Psychology (CPREC 2014.25). For the bimodal study, we invited individuals aged 18-35 to

participate through the SONA participant database at the Department of Psychology. We

recruited 30 individuals (15/15 female/male, mean+-STD age is 24.5667(+-4.5157)) who

were paid £10 per hour for a duration of 3-3.5 hours. The trimodal task was performed at the Centre Nationale de la Recherche Scientifique (CNRS) in France. Participants for this study were invited through the CNRS RISC system. Only individuals aged 18-80 were asked to participate in the trimodal study. Of 54 participants (35 female, mean($\pm$STD) age: 25.20($\pm$4.1) years) who participated in the trimodal study, 7 were excluded due to a recording error. Participants in the trimodal study were paid €40 for their effort.

Materials

In the trimodal study, auditory stimulation was applied using Etymotic noise-isolating insert earphones. Eccentric Rotating Mass motors controlled by two Texas Instruments DRV2605 haptic drivers were employed to deliver vibrotactile stimulation to the wrist. Two independent 8x8 LED matrices implemented in virtual reality goggles were used to apply visual inputs isolating visual hemifields. The setup was controlled using an Arduino Zero board. The bimodal study used auditory inputs generated by mixing three sinusoidal signals of either 500, 1000, and 2000 Hz (tone type A) or 350, 700, and 1400 Hz (tone type B) in Matlab R2016 based on (Chennu et al., 2013) and applied using EARTONE 3A insert earphones. Tactile stimulation was delivered using a custom-made device which applies mechanical pins to the fingertip with a Saia-Burgess 12.7 mm stroke, 12 v, 4 W DC push-action solenoid with 0.3-0.6 N force, and no nominal delay after current onset) controlled by an Arduino Mega board.

Paradigm

In a series of two experiments. expectations about sensory inputs were violated either locally within trials or globally between trials. Trials were composed of five stimuli with a stimulus duration of 50 ms and a stimulus onset interval of 150 ms. Each trial consisted of four identical ipsilateral stimuli followed by a deviant contralateral stimulus (locally deviant trials) or another ipsilateral stimulus (locally standard trials). Contrasting locally

deviant and standard trials reveals a MMN-like amplitude difference referred to as local effect in a time window between 50-250 ms after onset of the last stimulus in a trial. Global expectation violations are achieved when a frequently presented trial type is occasionally interspersed with a different trial type. A comparison of frequent (globally standard) trials and rare (globally deviant) trials uncovers a global effect which manifests as a late distributed P3b-like positive wave (Bekinschtein et al., 2009; Chennu et al., 2013, 2016; Phillips et al., 2016; Sitt et al., 2014; Wacongne et al., 2011).

During each study, two block types were presented. The bimodal study consisted of 8 auditory and somatosensory blocks. Standard stimuli in 50% of blocks were presented on the left side and on the right side in the remaining blocks. Each block consisted of 78% globally standard trials and 22% globally deviant trials. In block type bX, locally standard trials were occasionally interrupted by 22 % locally deviant trials which could equally likely be a locally deviant trial in which only the laterality or both laterality and stimulus type were varied. In block type BY, a stream of locally deviant trials in which the last stimulus was applied to the contralateral body hemisphere was occasionally interrupted by locally standard trials or locally deviant trials in which the last stimulus differed in laterality and stimulus type. In the trimodal study, blocks were composed of 80% globally standard and 20% globally deviant trials. Trials randomly started on the left or right side within each block with 50% probability. In blocktype tX, locally standard trials consisting of five identical stimuli were interspersed with locally deviant trials in which the last stimulus was applied to the contralateral hemisphere. Blocktype bY consisted of a sequence of locally deviant trials occasionally interrupted by locally standard trials.

In the bimodal study, somatosensory locally standard trials consisted of five touches ipsilaterally applied to the index finger. We introduced two types of locally deviant trials in which the last stimulus in a trial was applied to the contralateral index finger or contralateral middle finger. In auditory blocks, locally standard trials presented as five identical sounds. Local deviations were introduced by varying either only the laterality of

the ear which received the last sound in a trial or both the laterality and pitch of the last sound in a trial. The latter locally deviant trial type was globally deviant type in each block type and the former locally deviant trial type was globally standard in block type Y and deviant in block type X. The bimodal study consisted of 8 auditory and 8 somatosensory blocks. The block order was pseudo-randomised so that the experiment started with somatosensory block type Y followed by somatosensory block type X, no more than two consecutive blocks were presented in the same sensory modality, and each half of the experiment contained equal proportions of somatosensory and auditory blocks. Each block consisted of 158-160 trials and lasted ~4.5 min. Inter-trial intervals were randomly sampled from a uniform distribution between 800-1000 ms in steps of 50 ms. In each block, 30-34 globally deviant trials were embedded in a sequence of 112 globally standard trials and both deviant types occurred in equal proportions. Each globally deviant trial was preceded either by 2, 3, 4 or 5 globally standard trials in equal proportions. Participants were exposed to white noise during somatosensory blocks to avoid auditory cues from the tactile stimulation device.

The trimodal study contained blocks with exclusively auditory, visual or somatosensory stimulation. Participants in this study also underwent multisensory blocks in which expectations violations require to converge inputs from two of these sensory modalities and which were not further analysed. Local deviations were introduced by varying the laterality of the last stimulus in a trial. 50% of trials applied standard stimuli to the left hemisphere (ear, visual hemifield, or wrist) and the last stimulus to the right hemisphere (and vice versa). Participants underwent three experimental sessions with a total duration of 20 min (4.5 min per session). The remaining two sessions applied crossmodal stimulation and analyses were not included in this paper. 16 participants were presented with somatosensory blocks, 15 participants with auditory blocks and 16 participants with visual blocks. Each block type was presented twice per session in a fixed order: X-Y-X-Y. 31 (~20%) trials included in each block were globally deviant. Each

globally deviant trial was preceded by 3, 4, or 5 globally standard trials.

EEG data analyses

EEG data acquisition was performed using a Net Amps 300 amplifier with an

Electrical Geodesics 256-channel high density EEG net at the ICM in Paris for the bimodal

study and a 128-channel high density EEG net at the Department of Psychology,

University of Cambridge for the trimodal study. We performed EEG data preprocessing

using EEGLAB 2019 in Matlab 2019b. In a first step, data were down-sampled to 250 Hz,

filtered between 0.5-30 Hz and epoched with respect to the onset of the last stimulus in a

trial. Habituation trials at the start of each block were removed. Having removed

electrodes placed on the neck and cheek which record mostly muscle artefacts, we retained

91 electrodes in the bimodal data set and 175 electrodes in the trimodal data set for

further analysis. We performed baseline removal using a window of 100 ms before epoch

onset. Noisy trials (with a variance of >350) and channels (with a variance of >500) were

temporarily removed using a semi-automated procedure. We removed artefacts resulting

from sweat, eye and muscle movements using independent component analysis. Ultimately,

we removed the remaining artefacts using trial-wise interpolation.

We performed cluster-based permutation analyses to test for differences in ERP

amplitude time courses, We used the Common Average as a reference and performed a

baseline correction in a time window of 100 ms before onset of the last stimulus in a trial.

For each condition pair with unequal trial numbers, trials in the condition with a higher

trial number were randomly deselected until the number of trials in both conditions was

equal. Cluster-based permutation uses Monte Carlo partitioning to obtain a cluster-level

t-statistic. To perform Monte Carlo partitioning, data are pooled and randomly divided

into two new data sets of equal size. We performed two-sided t-tests on the subject

averages time-channel pairs and retained only t-values with p < 0.05. Spatiotemporally

adjacent t-values were summarised and the largest cluster-level summarised t-value was

identified. Having performed this procedure 3000 times, we determined the p-value corresponding to the proportion of maximal cluster-level t-values larger than the observed t-value in the original comparison. Conditions were deemed to be different if $p < 0.05$ (Oostenveld, Fries, Maris, & Schoffelen, 2011).

We applied temporal decoding to examine whether two contrasts rely on similar cortical signatures. Temporal decoding is a machine learning procedure which assesses whether a classifier trained to discriminate two trial types at one time point will generalise to the remaining time points in a sample. We applied a bootstrapping procedure in which 5 trials were randomly sampled from each dataset until we reached 540 epoch averages for each trial type (deviant/standard) respectively. This procedure was repeated 50 times. Classification performance was assessed on each of the resulting datasets. To perform temporal decoding within a sensory modality, we used a 5-fold stratified cross-validation procedure in which linear support vector machines were trained to optimally separate standard and deviant trials on 4/5 of the data set and tested on the remaining data. For temporal decoding between sensory modalities, we fitted classifiers using training and testing datasets from two different sensory modalities. For each condition pair, we trained and tested classifiers on every time point in a time window of 600 ms after onset of the last stimulus in a trial. Classification performance was assessed using the Area Under the Curve Receiver Operating Characteristic (AUC-ROC) which is a non-parametric criterion-free measure of separability. This procedure results in a training time vs testing time temporal generalisation matrix with AUC-ROC classification scores as cells. To identify adjacent AUC-ROC scores which differ from chance, we performed a Monte Carlo cluster-based permutation analysis with 1024 random partitions on classification score averages from each bootstrapped dataset and applied two-tailed paired t-tests to identify clusters of AUC-ROC values ($p < 0.05$) which differ from chance (King & Dehaene, 2014; King et al., 2014).

References

Allen, M., Fardo, F., Dietz, M. J., Hillebrandt, H., Friston, K. J., Rees, G., & Roepstorff, A. (2016a). Anterior insula coordinates hierarchical processing of tactile mismatch responses. *NeuroImage*, *127*, 34–43. https://doi.org/10.1016/j.neuroimage.2015.11.030

Allen, M., Fardo, F., Dietz, M. J., Hillebrandt, H., Friston, K. J., Rees, G., & Roepstorff, A. (2016b). Anterior insula coordinates hierarchical processing of tactile mismatch responses. *NeuroImage*, *127*, 34–43. https://doi.org/10.1016/j.neuroimage.2015.11.030

Amado, C., & Kovács, G. (2016). Does surprise enhancement or repetition suppression explain visual mismatch negativity? *European Journal of Neuroscience*, *43*(12), 1590–1600. https://doi.org/10.1111/ejn.13263

Auksztulewicz, R., & Friston, K. (2015). Attentional Enhancement of Auditory Mismatch Responses: A DCM/MEG Study. *Cerebral Cortex*, bhu323. https://doi.org/10.1093/cercor/bhu323

Bekinschtein, T. A., Dehaene, S., Rohaut, B., Tadel, F., Cohen, L., & Naccache, L. (2009). Neural signature of the conscious processing of auditory regularities. *Proceedings of the National Academy of Sciences*, *106*(5), 1672–1677. https://doi.org/10.1073/pnas.0809667106

Bledowski, C., Prvulovic, D., Hoechstetter, K., Scherg, M., Wibral, M., Goebel, R., & Linden, D. E. J. (2004). Localizing P300 Generators in Visual Target and Distractor Processing: A Combined Event-Related Potential and Functional Magnetic Resonance Imaging Study. *Journal of Neuroscience*, *24*(42), 9353–9360. https://doi.org/10.1523/JNEUROSCI.1897-04.2004

Cao, Y., Summerfield, C., Park, H., Giordano, B. L., & Kayser, C. (2019). Causal

Inference in the Multisensory Brain. *Neuron, 102*(5), 1076–1087.e8.

https://doi.org/10.1016/j.neuron.2019.03.043

Chao, Z. C., Takaura, K., Wang, L., Fujii, N., & Dehaene, S. (2018). Large-Scale

Cortical Networks for Hierarchical Prediction and Prediction Error in the

Primate Brain. *Neuron, 100*(5), 1252–1266.e3.

https://doi.org/10.1016/j.neuron.2018.10.004

Chennu, S., Noreika, V., Gueorguiev, D., Blenkmann, A., Kochen, S., Ibanez, A.,

. . . Bekinschtein, T. A. (2013). Expectation and Attention in Hierarchical

Auditory Prediction. *Journal of Neuroscience, 33*(27), 11194–11205.

https://doi.org/10.1523/JNEUROSCI.0114-13.2013

Chennu, S., Noreika, Gueorguiev, Shtyrov, Bekinschtein, & Henson, R. (2016).

Silent Expectations: Dynamic Causal Modeling of Cortical Prediction and

Attention to Sounds That Weren't. *Journal of Neuroscience, 36*(32), 8305–8316.

https://doi.org/10.1523/JNEUROSCI.1125-16.2016

Clark, A. (2013). Whatever next? Predictive brains, situated agents, and the future

of cognitive science. *Behavioral and Brain Sciences, 36*(03), 181–204.

Cornella, M., Leung, S., Grimm, S., & Escera, C. (2012). Detection of simple and

pattern regularity violations occurs at different levels of the auditory hierarchy.

PloS One, 7(8), e43604. https://doi.org/10.1371/journal.pone.0043604

Czigler, I., & Csibra, G. (1992). Event-Related Potentials and the Identification of

Deviant Visual Stimuli. *Psychophysiology, 29*(4), 471–485.

https://doi.org/10.1111/j.1469-8986.1992.tb01722.x

Czigler, I., Winkler, I., Pató, L., Várnagy, A., Weisz, J., & Balázs, L. (2006). Visual

temporal window of integration as revealed by the visual mismatch negativity

event-related potential to stimulus omissions. *Brain Research, 1104*(1), 129–140.

https://doi.org/10.1016/j.brainres.2006.05.034

de Lange, F. P., Heilbron, M., & Kok, P. (2018). How Do Expectations Shape
Perception? *Trends in Cognitive Sciences*, *22*(9), 764–779.
https://doi.org/10.1016/j.tics.2018.06.002

Dehaene, S., & Changeux, J.-P. (2011). Experimental and Theoretical Approaches
to Conscious Processing. *Neuron*, *70*(2), 200–227.
https://doi.org/10.1016/j.neuron.2011.03.018

Dehaene, S., & King, J.-R. (2016). Decoding the Dynamics of Conscious Perception:
The Temporal Generalization Method. In *Micro-, Meso-and Macro-Dynamics of
the Brain* (pp. 85–97). Springer.

Deouell, L. Y. (2007). The Frontal Generator of the Mismatch Negativity Revisited.
Journal of Psychophysiology, *21*(3-4), 188–203.
https://doi.org/10.1027/0269-8803.21.34.188

Donner, T., Kettermann, A., Diesch, E., Ostendorf, F., Villringer, A., & Brandt, S.
A. (2000). Involvement of the human frontal eye field and multiple parietal areas
in covert visual selection during conjunction search. *European Journal of
Neuroscience*, *12*(9), 3407–3414.
https://doi.org/10.1046/j.1460-9568.2000.00223.x

Driver, J., & Noesselt, T. (2008). Multisensory Interplay Reveals Crossmodal
Influences on 'Sensory-Specific' Brain Regions, Neural Responses, and
Judgments. *Neuron*, *57*(1), 11–23. https://doi.org/10.1016/j.neuron.2007.12.013

Dürschmid, S., Edwards, E., Reichert, C., Dewar, C., Hinrichs, H., Heinze, H.-J., . . .
Knight, R. T. (2016). Hierarchy of prediction errors for auditory events in
human temporal and frontal cortex. *Proceedings of the National Academy of
Sciences*, *113*(24), 6755–6760. https://doi.org/10.1073/pnas.1525030113

El Karoui, I., King, J.-R., Sitt, J., Meyniel, F., Van Gaal, S., Hasboun, D., . . .
Naccache, L. (2015). Event-Related Potential, Time-frequency, and Functional

Connectivity Facets of Local and Global Auditory Novelty Processing: An

Intracranial Study in Humans. *Cerebral Cortex*, *25*(11), 4203–4212.

https://doi.org/10.1093/cercor/bhu143

Ester, E. F., Serences, J. T., & Awh, E. (2009). Spatially Global Representations in

Human Primary Visual Cortex during Working Memory Maintenance. *Journal

of Neuroscience*, *29*(48), 15258–15265.

https://doi.org/10.1523/JNEUROSCI.4388-09.2009

Faivre, N., Filevich, E., Solovey, G., Kühn, S., & Blanke, O. (2018). Behavioral,

Modeling, and Electrophysiological Evidence for Supramodality in Human

Metacognition. *Journal of Neuroscience*, *38*(2), 263–277.

https://doi.org/10.1523/JNEUROSCI.0322-17.2017

Fardo, F., Auksztulewicz, R., Allen, M., Dietz, M. J., Roepstorff, A., & Friston, K.

J. (2017). Expectation violation and attention to pain jointly modulate neural

gain in somatosensory cortex. *Neuroimage*, *153*.

Felleman, D. J., & Van Essen, D. (1991). Distributed hierarchical processing in the

primate cerebral cortex. *Cerebral Cortex (New York, N.Y. : 1991)*, *1*(1), 1–47.

https://doi.org/10.1093/cercor/1.1.1

Friston, K. (2005). A theory of cortical responses. *Philosophical Transactions of the

Royal Society B: Biological Sciences*, *360*(1456), 815–836.

https://doi.org/10.1098/rstb.2005.1622

Garrido, M. I., Friston, K. J., Kiebel, S. J., Stephan, K. E., Baldeweg, T., & Kilner,

J. M. (2008). The functional anatomy of the MMN: A DCM study of the roving

paradigm. *NeuroImage*, *42*(2), 936–944.

https://doi.org/10.1016/j.neuroimage.2008.05.018

Garrido, M. I., Kilner, J. M., Kiebel, S. J., & Friston, K. J. (2009). Dynamic Causal

Modeling of the Response to Frequency Deviants. *Journal of Neurophysiology*,

101(5), 2620–2631. https://doi.org/10.1152/jn.90291.2008

Garrido, M. I., Kilner, J. M., Stephan, K. E., & Friston, K. J. (2009). The

mismatch negativity: A review of underlying mechanisms. *Clinical*

Neurophysiology, 120(3), 453–463.

Ghazanfar, A., & Schroeder, C. (2006). Is neocortex essentially multisensory?

Trends in Cognitive Sciences, 10(6), 278–285.

https://doi.org/10.1016/j.tics.2006.04.008

Hamm, J. P., & Yuste, R. (2016). Somatostatin Interneurons Control a Key

Component of Mismatch Negativity in Mouse Visual Cortex. *Cell Reports,*

16(3), 597–604. https://doi.org/10.1016/j.celrep.2016.06.037

Hedge, C., Stothart, G., Todd Jones, J., Rojas Frías, P., Magee, K. L., & Brooks, J.

C. W. (2015). A frontal attention mechanism in the visual mismatch negativity.

Behavioural Brain Research, 293, 173–181.

https://doi.org/10.1016/j.bbr.2015.07.022

Hochstein, S., & Ahissar, M. (2002). View from the Top: Hierarchies and Reverse

Hierarchies in the Visual System. *Neuron, 36*(5), 791–804.

https://doi.org/10.1016/S0896-6273(02)01091-7

Hohwy, J. (2012). Attention and conscious perception in the hypothesis testing

brain. *Attention and Consciousness in Different Senses*, 74.

Hubel, D. H., & Wiesel, T. N. (1962). Receptive fields, binocular interaction and

functional architecture in the cat's visual cortex. *The Journal of Physiology,*

160(1), 106–154. https://doi.org/10.1113/jphysiol.1962.sp006837

Katayama, J., & Polich, J. (1998). Stimulus context determines P3a and P3b.

Psychophysiology, 35(1), 23–33.

Kiebel, S. J., Daunizeau, J., & Friston, K. J. (2008). A Hierarchy of Time-Scales

754 and the Brain. *PLOS Computational Biology*, *4*(11), e1000209.

755 https://doi.org/10.1371/journal.pcbi.1000209

756 King, J.-R., & Dehaene, S. (2014). Characterizing the dynamics of mental

757 representations: The temporal generalization method. *Trends in Cognitive*

758 *Sciences*, *18*(4), 203–210. https://doi.org/10.1016/j.tics.2014.01.002

759 King, J.-R., Gramfort, A., Schurger, A., Naccache, L., & Dehaene, S. (2014). Two

760 Distinct Dynamic Modes Subtend the Detection of Unexpected Sounds. *PLOS*

761 *ONE*, *9*(1), e85791. https://doi.org/10.1371/journal.pone.0085791

762 Kristjánsson, Á., Vuilleumier, P., Schwartz, S., Macaluso, E., & Driver, J. (2007).

763 Neural Basis for Priming of Pop-Out during Visual Search Revealed with fMRI.

764 *Cerebral Cortex*, *17*(7), 1612–1624. https://doi.org/10.1093/cercor/bhl072

765 Linden, D. E. J., Prvulovic, D., Formisano, E., Völlinger, M., Zanella, F. E.,

766 Goebel, R., & Dierks, T. (1999). The Functional Neuroanatomy of Target

767 Detection: An fMRI Study of Visual and Auditory Oddball Tasks. *Cerebral*

768 *Cortex*, *9*(8), 815–823. https://doi.org/10.1093/cercor/9.8.815

769 Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and

770 MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190.

771 https://doi.org/10.1016/j.jneumeth.2007.03.024

772 Mashour, G. A., Roelfsema, P., Changeux, J.-P., & Dehaene, S. (2020). Conscious

773 Processing and the Global Neuronal Workspace Hypothesis. *Neuron*, *105*(5),

774 776–798. https://doi.org/10.1016/j.neuron.2020.01.026

775 Maunsell, J. H., & Essen, D. van. (1983). The connections of the middle temporal

776 visual area (MT) and their relationship to a cortical hierarchy in the macaque

777 monkey. *Journal of Neuroscience*, *3*(12), 2563–2586.

778 https://doi.org/10.1523/JNEUROSCI.03-12-02563.1983

Mäntysalo, S., & Näätänen, R. (1987). The duration of a neuronal trace of an auditory stimulus as indicated by event-related potentials. *Biological Psychology*, *24*(3), 183–195.

Miller, E. K. (2009). Executive Function and Higher-Order Cognition: Definition and Neural Substrates. *Encyclopedia of Neuroscience*, *4*, 7.

Murray, C. (2004). An interpretative phenomenological analysis of the embodiment of artificial limbs. *Disability and Rehabilitation*, *26*(16), 963–973.

Murray, J., Bernacchia, A., Freedman, D. J., Romo, R., Wallis, J. D., Cai, X., . . . Wang, X.-J. (2014). A hierarchy of intrinsic timescales across primate cortex. *Nature Neuroscience*, *17*(12), 1661–1663. https://doi.org/10.1038/nn.3862

Näätänen, R., Simpson, M., & Loveless, N. E. (1982). Stimulus deviance and evoked potentials. *Biological Psychology*, *14*(1), 53–98. https://doi.org/10.1016/0301-0511(82)90017-5

Näätänen, Risto, Tervaniemi, M., Sussman, E., Paavilainen, P., & Winkler, I. (2001). 'Primitive intelligence' in the auditory cortex. *Trends in Neurosciences*, *24*(5), 283–288. https://doi.org/10.1016/S0166-2236(00)01790-2

Nelken, I., & Bar-Yosef, O. (2008). Neurons and objects: The case of auditory cortex. *Frontiers in Neuroscience*, *2*. https://doi.org/10.3389/neuro.01.009.2008

Nyman, G., Alho, K., Laurinen, P., Paavilainen, P., Radil, T., Reinikainen, K., . . . Näätänen, R. (1990). Mismatch negativity (MMN) for sequences of auditory and visual stimuli: Evidence for a mechanism specific to the auditory modality. *Electroencephalography and Clinical Neurophysiology/Evoked Potentials Section*, *77*(6), 436–444. https://doi.org/10.1016/0168-5597(90)90004-W

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive

Electrophysiological Data [Research Article].

https://www.hindawi.com/journals/cin/2011/156869/.

https://doi.org/10.1155/2011/156869

Ostwald, D., Spitzer, B., Guggenmos, M., Schmidt, T. T., Kiebel, S. J., &

Blankenburg, F. (2012). Evidence for neural encoding of Bayesian surprise in

human somatosensation. *NeuroImage*, *62*(1), 177–188.

Pazo-Alvarez, P., Cadaveira, F., & Amenedo, E. (2003). MMN in the visual

modality: A review. *Biological Psychology*, *63*(3), 199–236.

https://doi.org/10.1016/S0301-0511(03)00049-8

Pegado, F., Bekinschtein, T., Chausson, N., Dehaene, S., Cohen, L., & Naccache, L.

(2010). Probing the lifetimes of auditory novelty detection processes.

Neuropsychologia, *48*(10), 3145–3154.

https://doi.org/10.1016/j.neuropsychologia.2010.06.030

Phillips, H. N., Blenkmann, A., Hughes, L. E., Kochen, S., Bekinschtein, T. A.,

Rowe, J. B., & others. (2016). Convergent evidence for hierarchical prediction

networks from human electrocorticography and magnetoencephalography.

Cortex, *82*, 192–205.

Polich, J. (2007). Updating P300: An integrative theory of P3a and P3b. *Clinical

Neurophysiology*, *118*(10), 2128–2148.

https://doi.org/10.1016/j.clinph.2007.04.019

Rao, R. P., & Ballard, D. H. (1999). Predictive coding in the visual cortex: A

functional interpretation of some extra-classical receptive-field effects. *Nature

Neuroscience*, *2*(1), 79–87.

Riesenhuber, M., & Poggio, T. (1999). Hierarchical models of object recognition in

cortex. *Nature Neuroscience*, *2*(11), 1019–1025. https://doi.org/10.1038/14819

Shen, G., Smyk, N. J., Meltzoff, A. N., & Marshall, P. J. (2018). Using somatosensory mismatch responses as a window into somatotopic processing of tactile stimulation. *Psychophysiology*, *55*(5), e13030. https://doi.org/10.1111/psyp.13030

Shinozaki, N., Yabe, H., Sutoh, T., Hiruma, T., & Kaneko, S. (1998). Somatosensory automatic responses to deviant stimuli. *Cognitive Brain Research*, *7*(2), 165–171. https://doi.org/10.1016/S0926-6410(98)00020-2

Sitt, J. D., King, J.-R., El Karoui, I., Rohaut, B., Faugeras, F., Gramfort, A., … Naccache, L. (2014). Large scale screening of neural signatures of consciousness in patients in a vegetative or minimally conscious state. *Brain*, *137*(8), 2258–2270. https://doi.org/10.1093/brain/awu141

Squires, K., Petuchowski, S., Wickens, C., & Donchin, E. (1977). The effects of stimulus sequence on event related potentials: A comparison of visual and auditory sequences. *Perception & Psychophysics*, *22*(1), 31–40. https://doi.org/10.3758/BF03206077

Tiitinen, H., May, P., Reinikainen, K., & Näätänen, R. (1994). Attentive novelty detection in humans is governed by pre-attentive sensory memory. *Nature*, *372*(6501), 90.

Uhrig, L., Dehaene, S., & Jarraya, B. (2014). A Hierarchy of Responses to Auditory Regularities in the Macaque Brain. *Journal of Neuroscience*, *34*(4), 1127–1132. https://doi.org/10.1523/JNEUROSCI.3165-13.2014

Van Essen, D. C., & Maunsell, J. H. R. (1983). Hierarchical organization and functional streams in the visual cortex. *Trends in Neurosciences*, *6*, 370–375. https://doi.org/10.1016/0166-2236(83)90167-4

Wacongne, C., Changeux, J.-P., & Dehaene, S. (2012). A Neuronal Model of Predictive Coding Accounting for the Mismatch Negativity. *Journal of*

Neuroscience, *32*(11), 3665–3678.

https://doi.org/10.1523/JNEUROSCI.5003-11.2012

Wacongne, C., Labyt, E., van Wassenhove, V., Bekinschtein, T., Naccache, L., & Dehaene, S. (2011). Evidence for a hierarchy of predictions and prediction errors in human cortex. *Proceedings of the National Academy of Sciences*, *108*(51), 20754–20759. https://doi.org/10.1073/pnas.1117807108

Walz, J. M., Goldman, R. I., Carapezza, M., Muraskin, J., Brown, T. R., & Sajda, P. (2013). Simultaneous EEG-fMRI Reveals Temporal Evolution of Coupling between Supramodal Cortical Attention Networks and the Brainstem. *Journal of Neuroscience*, *33*(49), 19212–19222. https://doi.org/10.1523/JNEUROSCI.2649-13.2013

Yamaguchi, S., & Knight, R. T. (1991a). Anterior and posterior association cortex contributions to the somatosensory P300. *Journal of Neuroscience*, *11*(7), 2039–2054.

Yamaguchi, S., & Knight, R. T. (1991b). P300 generation by novel somatosensory stimuli. *Electroencephalography and Clinical Neurophysiology*, *78*(1), 50–55.