Constitutively active STING causes neuroinflammation and degeneration

of dopaminergic neurons in mice

Eva M Szegö[1*], Laura Malz[2*], Nadine Bernhardt[3], Angela Rösen-Wolff[4], Björn H.

Falkenburger[5#], Hella Luksch[6#]

[1] Department of Neurology, TU Dresden, Dresden, Germany; e-mail:

Eva.Szego@uniklinikum-dresden.de

[2] Departments of Neurology & Pediatrics, TU Dresden, Dresden, Germany; e-mail:

Laura.Malz@uniklinikum-dresden.de

[3] Department of Psychiatry, TU Dresden, Dresden, Germany; e-mail:

Nadine.Bernhardt@uniklinikum-dresden.de

[4] Department of Pediatrics, TU Dresden, Dresden, Germany; e-mail: Angela.Roesen-

Wolff@uniklinikum-dresden.de

[5] Department of Neurology, TU Dresden, Dresden, Germany; Deutsches Zentrum für

Neurodegenerative Erkrankungen, Dresden, Germany; e-mail: bfalken@ukdd.de

[6] Department of Pediatrics, TU Dresden, Dresden, Germany; e-mail:

Hella.Luksch@uniklinikum-dresden.de

* EMSz and LM contributed equally

# BHF and HL contributed equally

Abstract

The innate immune system can protect against certain aspects of neurodegenerative diseases, but also contribute to disease progression. Stimulator of interferon genes (STING) is activated after detection of cytoplasmic dsDNA by cGAS (cyclic GMP-AMP synthase) as part of the defense against viral pathogens, activating type I interferon and NF-kB/inflammasome signaling. In order to specifically test the relevance of this pathway for the degeneration of dopaminergic neurons in Parkinson's disease, we studied a mouse model with heterozygous expression of the constitutively active STING variant N153S.

In adult mice expressing N153S STING, the number of dopaminergic neurons was smaller than in controls, as was the density of dopaminergic axon terminals and the concentration of dopamine in the striatum. We also observed alpha-synuclein pathology and a lower density of synaptic puncta. Neuroinflammation was quantified by staining astroglia and microglia, by measuring mRNAs, proteins and nuclear translocation of transcription factors. Neuroinflammatory markers were already elevated in juvenile mice, thus preceding the degeneration of dopaminergic neurons. Inflammation and neurodegeneration were blunted in mice deficient for signaling by type I interferons or inflammasomes, but not suppressed completely.

Collectively, these findings demonstrate that chronic activation of the STING innate immunity pathway is sufficient to cause degeneration of dopaminergic neurons. This pathway could be targeted therapeutically.

Introduction

Inflammation contributes significantly to the pathogenesis of neurodegenerative diseases, including Parkinson's disease (PD) (Harms et al., 2021; Hirsch and Standaert, 2021). Inflammatory serum markers are associated with more severe PD symptoms and with a more rapid progression of cognitive decline (Hall et al., 2018; Mollenhauer et al., 2019). Anti-inflammatory drugs like aspirin have been associated with a lower risk of developing PD in epidemiological studies (Chen et al., 2003). The pathological hallmarks of PD include the degeneration of dopaminergic neurons in the substantia nigra and the cytoplasmic inclusions termed Lewy bodies (Obeso et al., 2017). Aggregates of alpha-synuclein (aSyn), one of the main constituents of Lewy bodies, stimulate monocytes and microglia (Grozdanov et al., 2019) and astroglia (Chou et al., 2021), suggesting that inflammatory changes in PD respond to aSyn pathology and contribute to disease progression (Harms et al., 2021; Hirsch and Standaert, 2021).

The stimulator of interferon genes (STING) responds to cytoplasmic dsDNA as part of the innate immunity defense against viral pathogens (Paul et al., 2021). STING is activated by cyclic GMP-AMP synthase (cGAS), which binds dsDNA and catalyzes synthesis of the second messenger cyclic GMP-AMP (Motwani et al., 2019). This pathway can be activated by viral nucleic acids but also by self-DNA (Chen et al., 2016; Li and Chen, 2018; Motwani et al., 2019). The cGAS-STING pathway has been implicated in the pathogenesis of PD (Beyer et al., 2020), but also Alzheimer's disease and amyotrophic lateral sclerosis (Chen et al., 2021; Li et al., 2021; Paul et al., 2021). For instance, mice deficient for the PD-associated proteins PINK and parkin fail to degrade damaged mitochondria, accumulate mitochondrial DNA in the cytosol and show a STING-mediated inflammatory phenotype after exhaustive exercise (Sliter et al., 2018).

The two main transcription factors activated by STING are the interferon regulatory factor 3 (IRF3) (Chen et al., 2021) and nuclear factor 'kappa-light-chain-enhancer' of activated B-cells (NF-κB) (Liu et al., 2014). Consequently, type I interferons (IFN) and pro-inflammatory cytokines are produced, triggering a secondary inflammation through the activation of inflammasomes (Hopfner and Hornung, 2020). Additionally, STING can activate inflammasomes directly (Wang et al., 2020). Inflammasomes are major signaling hubs that activate caspase-1 and control the bioactivity of pro-inflammatory cytokines of the interleukin (IL)-1 family (Gaidt et al., 2017; Schroder and Tschopp, 2010). Activation of inflammasomes formed by the NOD-, LRR- and pyrin domain-containing protein 3 (NLRP3)

has been linked to the progression of several neurodegenerative diseases (Heneka et al.,

2018). NLRP3 immunoreactivity is increased in mesencephalic neurons of PD patients and

NLRP3 variants are associated with the risk to develop PD (von Herrmann et al., 2018).

Conversely, inhibiting NLRP3 mitigates degeneration of dopaminergic neurons and aSyn

pathology in mouse models (Gordon et al., 2018). NRLP3 has also been associated with

Alzheimer's disease (Heneka et al., 2018), prion disease (Nazmi et al., 2019), and traumatic

brain injury (Sen et al., 2020).

Neuroinflammation receives growing attention in neurodegenerative diseases because it

represents a promising therapeutic target. Designing such therapies requires, however, to

determine the specific effects of individual components of this highly interconnected

signaling network, which responds to diverse stimuli and involves many different cell types.

Given the role of the STING pathway in PINK/parkin related damage (Sliter et al., 2018), we

wanted to learn more about the role of STING for PD pathogenesis and determine whether

specific activation of STING is sufficient to cause degeneration of dopaminergic neurons. In

order to test this, we used a mouse model with heterozygous expression of the STING genetic

variant N153S (Luksch et al., 2019). Constitutively active STING mutants cause an

autoinflammatory disease in humans termed STING-associated vasculopathy with onset in

infancy (SAVI) (Crow and Casanova, 2014; Liu et al., 2014). SAVI is characterized by

systemic inflammation with acral vasculitis, T cell lymphopenia, and interstitial pulmonary

disease. Major features of systemic inflammation in SAVI are recapitulated in STING N153S

knockin mice (Luksch et al., 2019, Siedel et al., 2020). For simplicity, we refer to these as

STING ki mice here and use the term STING WT for the corresponding wild type littermate

controls. In this work, we determined the extent of neuroinflammation in the brains of young

and adult mice, the extent of dopamine neuron degeneration, and aSyn pathology.

Furthermore, we used additional knockout mice to determine which of the known

downstream signaling pathways contribute to STING-induced neurodegeneration.

Materials and Methods

Source of chemicals, antibodies, composition of buffers, equipment and software used in this study are listed in supplemental Table S1.

Animals

All animal experiments were carried out in accordance with the European Communities Council Directive of November 24, 1986 (86/609/EEC) and approved by the Landesdirektion Dresden, Germany. Mice of both sexes were housed under a 12-hour light and dark cycle with free access to pelleted food and tap water in the Experimental Center, Technische Universität, Dresden, Germany. Heterozygous STING N153S/WT ki mice (STING ki) were previously described (Luksch et al., 2019). STING ki or STING WT mice were crossed to $Ifnar1^{-/-}$ mice (a gift from Axel Roers, Dresden, Germany) and $Casp1^{-/-}$ mice (a gift from Stefan Winkler, Dresden, Germany).

For all genotypes, five-week-old (from here referred as juvenile) or 20-23-week-old (referred as adult) animals were sacrificed with an overdose of isoflurane (Baxter, Lessines, Belgium). For western blot analysis and for gene expression analysis, brains were rapidly removed from the skull and washed in ice-cold Tris-buffered saline (TBS, pH 7.4). Cortex and striatum were dissected, snap-frozen in liquid nitrogen and stored at -80°C until use. For histology, mice were perfused transcardially with 4 % paraformaldehyde (PFA) in TBS. After post-fixation (4 % PFA, overnight) and cryoprotection (30 % sucrose in TBS), 30 µm-thick coronal brain sections were cut in a cryostat (Leica, Germany).

Immunofluorescence stainings of mouse brain sections

To quantify the number of dopaminergic neurons in the *substantia nigra* (SN), every fifth section throughout the entire SN was stained for tyrosine hydroxylase (TH) as previously (Szegő et al., 2021). In brief, after blocking (2 % bovine serum albumin, 0,3 % Triton X-100 in TBS; 1 h RT), sections were incubated with the primary antibody in blocking solution (two overnights), followed by the fluorescently labelled secondary antibody (Alexa 488 conjugated donkey anti-sheep, 1:2000, overnight). Sections were counterstained with Hoechst and mounted with Fluoromount-G.

To quantify the density of dopaminergic axon terminals (fibers) and neuroinflammation in the striatum, every sixth section throughout the entire striatum was stained with a cocktail of primary antibodies: TH (Pel Freeze, P40101, 1:1000), Iba1 (Wako, 019-19741, 1:1000)

and GFAP (abcam, ab4674, 1:2000). As fluorescently labelled secondary antibodies, Alexa

488 conjugated donkey anti-sheep, Alexa 555 conjugated donkey anti-rabbit, Alexa 647

conjugated donkey anti-chicken were used (1:2000, overnight). Sections were counterstained

with Hoechst and mounted with Fluoromount-G.

Quantification of dopaminergic neuron number, striatal fiber density and gliosis

The number of dopaminergic somata in the SN was determined by supervised manual

counting by an investigator blinded to the experimental groups. For each animal, every fifth

section throughout the rostro-caudal extent of the SN (2.54 to -3.88 mm posterior to Bregma

based on Paxinos and Franklin, 2001) was incorporated into the counting procedure. In each

section, z stacks were acquired (step size: 2 µm, 5 slices in total) from both hemispheres with

a 20x objective (N.A 0.8, Axio Imager 2, Zeiss). Stacks were stitched to reconstruct the entire

SN. After adjusting the threshold and carefully marking the borders of the SN, only TH-

positive cell bodies with a visible nucleus in the blue channel were counted by ImageJ (1,53c;

Cell Counter plugin).The total number of neurons per SN was estimated by multiplying the

counted cell number by five (every fifth section was used for this analysis). For quantification

of gliosis, five fluorescent images were acquired from every sixth striatal section stained for

GFAP and Iba1 using a 20x objective. After adjusting the threshold and noise removal

(Background subtraction, rolling ball radius 50) from the individual images (separately for

GFAP and Iba1 channels), the area fraction was determined by ImageJ from ten regions of

interest per image. Results were analyzed using a generalized linear mixed model (glm) in

RStudio with a hierarchically nested design (expressed as percent area) as previously (Szegő

et al., 2021).

From the same striatal sections, the density of the dopaminergic axon terminals (fibers)

was determined as described previously (Szegő et al., 2012). In brief, z-stack images were

acquired (five planes, 0.5 µm step size, 100x objective, N.A. 1.4; Axio Imager 2, Zeiss). TH-

positive fibers were delineated from the maximal intensity projection (ImageJ) after adjusting

the threshold, noise removal and binarization, and density was expressed as percent area.

Every sixth section per animal, five images per section and ten boxes per image were

analyzed in a hierarchically nested design as above.

Protein analyses

To detect protein changes, cortical and striatal tissue were mechanically lysed in a buffer

containing 250 mM sucrose, 50 mM TRIS (pH 7,5), 1 mM EDTA, 5 mM $MgCl_2$, 1 % Triton

X-100 in the presence of protease and phosphatase inhibitors (MedChem Express) as

previously (Szegő et al., 2012). Samples were centrifuged (14000 g, 30 min, 4 °C) and

protein concentration in the supernatant was determined with the BCA method

(ThermoFisher, Germany). After boiling with 4x Laemmli buffer (1 M Tris pH 6.8, 0.8 %

SDS, 40 % glycerol, 5 % β-mercaptoethanol, traces of bromophenol blue, 5 min, 95 ºC), 5 µg

protein was loaded onto a 4-20 % Tris/glycine SDS gel for western blot analysis. Membranes

were fixed in paraformaldehyde (10 min, RT) and blocked with 1 % bovine serum albumin,

0,05 % Tween 20 in TBS. Membranes were incubated first in the presence of antibodies

against phosphorylated aSyn and aSyn, (Cell Signaling), then with βIII-tubulin as loading

control (overnight, 4 ºC). Following washing, membranes were incubated in the presence of

horseradish peroxidase-conjugated secondary antibodies (donkey anti-mouse or donkey anti-

rabbit). Signal was detected using chemiluminescent substrate and a camera-based system.

ImageJ was used to determine the optical density of protein bands and all data were analyzed

for each group (n = 5 animals/group) based on 3 independent blots. Optical densities were

normalized to the expression of the density of the tubulin loading control of the same sample,

and then expressed relative to the WT animals.

Gene expression analyses

Total RNA was extracted from snap frozen dissected prefrontal cortex tissue by using the

RNeasy Mini Kit (Qiagen, Germany) according to the manufacturer's instructions. cDNA

was generated by MMLV reverse transcription (Promega Germany). Quantitative Real Time

PCR assays were carried out by using QuantStudio 5 (Thermo Fisher Scientific, Germany)

and GoTaq®aPCR Master Mix with SYBR green fluorescence (Promega, Germany). PCR

primer sequences were retrieved from the Primer Bank database (Spandidos et al., 2009).

Expression of genes was normalized to the expression of the housekeeping genes (*Hprt1*,

Rpl13a, *Eef2*) and to the STING WT by using the ΔΔCt method. Sequences of primers are

listed in supplemental Table S2.

Statistical analyses

Sample numbers for each analysis are listed in supplemental Table S4. In graphs, markers

represent individual animals; lines represent mean and standard deviation (SD) of all animals.

Data normality was tested by the Kolmogorov-Smirnov test and graphically by QQ plot (R,

version 2.8.0; R Development Core Team 2008). Grubbs test was used to identify outliers. t-

test, Mann-Whitney test or two-way ANOVA were performed using GraphPad Prism

(Versions 5.01 and 9.0.0). Linear regression was performed using R. For generalized linear

mixed-effects model (Szegő et al., 2021), animal, section and image were used as random

effects nested within each other (R package: lme4). p values are indicated in the graphs by

symbols with * representing $p<0.05$, ** representing $p<0.01$, *** representing $p<0.001$.

Exact p values are given in the Figure legends.

Results

Neuroinflammation and degeneration of dopaminergic neurons in mice with constitutive STING activation

To characterize the neuronal phenotype in adult STING ki mice, we assessed neuroinflammation and the integrity of the dopaminergic nigrostriatal system. First, we stained striatal sections for Iba1 to determine the activation of microglia. The Iba1 positive area fraction was 9-fold higher in STING ki mice than in STING WT (Figure 1A and E). Activation of astroglia, as determined by GFAP staining, was 26-fold higher in STING ki mice than in STING WT (Figure 1B and F). Thus, STING ki mice exhibit a strong neuroinflammatory phenotype, consistent with increased systemic inflammation (Luksch et al., 2019).

Next, we asked whether the chronic neuroinflammation in the STING ki mice is associated with the degeneration of dopaminergic neurons. Somata of dopaminergic neurons in the *substantia nigra* (SN, Figure 1C) and dopaminergic axon terminals in the striatum (Figure 1D) were identified by staining for tyrosine hydroxylase (TH). The SN of STING ki mice contained significantly fewer TH-positive neurons than the SN of STING WT (Figure 1G and figure supplement 1A). Similarly, the density of dopaminergic axon terminals (fibers) in the striatum was lower in STING ki mice than in STING WT (Figure 1H). Accordingly, the concentration of dopamine in the striatum was lower in STING ki mice than in STING WT mice (Figure 1I). The concentration of dopamine metabolites was higher in STING ki mice (supplemental Figure 1B), suggesting increased dopamine turnover as commonly observed with dopamine depletion. Taken together, these findings demonstrate that the integrity of nigrostriatal dopaminergic neurons is compromised in STING ki mice.

Neuroinflammation without obvious degeneration of the dopaminergic neurons in juvenile mice with constitutive STING activation

In order to explore whether the compromised integrity of dopaminergic neurons is a consequence of a prolonged neuroinflammation, we next analyzed brain sections of juvenile (5-week-old) STING ki and STING WT mice (Figure 2). Microglia was already significantly activated in juvenile STING ki mice. The area fraction of the Iba1 staining was 2-fold higher in STING ki mice than in STING WT (Figure 2A and E). Similarly, the area fraction of GFAP signal was 14-fold higher in STING ki mice than in STING WT (Figure 2B and F), suggesting activation of astroglia in juvenile STING ki mice. However, the number of TH-

positive neurons in the *substantia nigra* was not different between juvenile STING ki mice

and STING WT (Figure 2C and G, figure supplement 1C). Similarly, striatal axon terminals

(Figure 2D and H) and the concentrations of striatal dopamine and its metabolites (Figure 2I,

figure supplement 1D) were not different between STING ki mice and STING WT.

Taken together, these findings demonstrate that the compromised integrity of the

nigrostriatal system in adult STING ki mice (Figure 1) represents an adult-onset

neurodegeneration and not a developmental defect. Given that activation of microglia and

astroglia in STING ki mice precedes degeneration of dopaminergic neurons, STING-induced

neuroinflammation could contribute to the neurodegeneration.

aSyn pathology and synaptic defects in the striatum of STING ki mice

To analyze aSyn pathology in the striatum of STING ki mice, we measured the amount of

aSyn protein phosphorylated at serine 129 (paSyn), which is considered one of the major

pathological forms of aSyn (Anderson et al., 2006; Fujiwara et al., 2002; Samuel et al., 2016).

Lysates of adult STING ki mice contained a substantial amount of paSyn (Figure 3A), which

was barely detectable in STING WT. The amount of total aSyn was lower in STING ki mice

(Figure 3C) and the ratio of paSyn to total aSyn increased (Figure 3D) – as commonly

observed in synucleinopathy models and PD patients (Anderson et al., 2006; Chatterjee et al.,

2020; Fujiwara et al., 2002; Szegö et al., 2022).

Thioflavin S (ThioS) binds to the characteristic β-sheet conformation of amyloid-

containing proteins, including aSyn (Froula et al., 2019; Neumann et al., 2002). The number

of cells with ThioS-positive inclusions was higher in the striatum of adult STING ki mice

than in STING WT (Figure 3 B and E), consistent with the findings from the aSyn

immunoblots.

Since dopamine depletion and aSyn pathology can compromise synaptic integrity, we next

quantified the density of synapses in the striatum. Presynaptic puncta were detected by

staining against synapsin; post-synaptic puncta were detected by staining against homer

(Figure 3F and G). The density of synapsin puncta was 18% lower in adult STING ki than in

STING WT (Figure 3H), the density of post-synaptic puncta was 9% lower in STING ki than

in STING WT mice (Figure 3I). In summary, we observed aSyn pathology and a reduced

density of synapses in the striatum of adult STING ki mice.

Type I IFN signaling and NF-κB/inflammasome dependent signaling are activated in the brain of STING ki mice

In order to analyze the signaling pathways by which constitutively active STING causes degradation of dopaminergic neurons and aSyn pathology, we examined the expression of selected interferon-stimulated genes (ISGs) by quantitative real time PCR in juvenile and adult STING ki and STING WT mice (Figure 4A-D). In STING WT mice, we observed an age-dependent increase for interferon-induced GTP-binding protein Mx1 (*Mx-1*, 10-fold, p= 0,0000006, comparison not depicted in Figure 4. for clarity), interferon-gamma induced protein 10 kD (*Ip-10*, 22-fold, p= 0,000105) and *Sting1* (3,7-fold, p=0,0008399), consistent with previous findings (Harris et al., 2020).

In STING ki mice, the expression of ISGs was generally higher than in STING WT, both comparing juvenile mice and comparing adult mice (Figures 4A-D, p-values in Figure legend). The expression of *Sting1* was not significantly different between adult STING ki and STING WT, confirming that the expression of the STING N153S mutant is not changing.

cGAS/STING activation also leads to the activation of NF-κB/inflammasome pathway (Balka et al., 2020; Balka and De Nardo, 2021; Wang et al., 2020). We therefore analyzed induction of the NF-κB/inflammasome pathway by quantitative real time PCR of the downstream mediators, tumor necrosis factor alpha (*Tnfα*), interleukin 1 beta (*Il-1β*), and caspase 1 (*Casp1*) (Figure 4E-G). In STING WT mice, the expression of *Tnfα*, *Il-1β*, *Casp1* increased with age (*Tnfα*: 5-fold, p= 0,0238678; *Il-1β*: 21-fold, p= 0,0000038; *Casp1*: 150-fold, p<0,00001, comparisons not depicted in Figure 4. for clarity), consistent with earlier reports (Mejias et al., 2018). In juvenile STING ki mice, the expression of *Tnfα* was 13-fold higher than in juvenile STING WT (Figure 4E; p= 0,0001448), *Il-1β* was 33-fold higher (Figure 4F, p= 0,00005) and *Casp1* was 9-fold higher (Figure 4G; p= 0,0000369). In adult mice, expression of *Tnfa* and *Casp1* was not different between STING WT and STING ki whereas expression of *Il-1β* was 6-fold higher in adult STING ki than in adult STING WT (Figure 4F, p= 0,0389).

STAT3 and NF-κB translocate to the nucleus in STING ki mice

Nuclear trafficking is critical for the function of transcription factors such as STAT3 and NF-κB (Balka and De Nardo, 2021; Noguchi et al., 2013). To further investigate the activation of the type I IFN and NF-kB-dependent pathways in juvenile and adult STING ki mice, we quantified the number of nuclei positive for phosphorylated STAT3 (pSTAT3) and NF-κB in the striatum (Figure 5). There were only very few pSTAT3-positive nuclei in juvenile and

adult STING WT mice (Figure 5A and C). In juvenile STING ki mice, the average number of

pSTAT3-positive nuclei was 4-fold higher than in juvenile STING WT mice (Figure 5C, p=

0,05678, 2-way ANOVA). In adult STING ki mice, the number of pSTAT3-positive nuclei

was 15-fold higher than in STING WT (p= 0,00004). Similarly, the number of NF-kB-

positive nuclei was 3-fold higher in juvenile STING ki mice than in STING WT (Figure 5B

and, p= 0,009). In adult STING ki mice, the number of NF-kB-positive nuclei was 6-fold

higher than in STING WT (Figure 5D, p= 0,007).

Taken together, our results show nuclear translocation of pSTAT3 and NF-kB in the

striatum of STING ki mice, consistent with a robust activation of type I IFN and NF-

kB/inflammasome dependent signaling in STING ki mice.

318   Type I IFN and NF-kB/inflammasome signaling contribute to neuroinflammation in

319   STING ki mice

We next sought to investigate the contribution of the type I IFN and NF-kB/inflammasome

signaling pathways to neuroinflammation in STING ki mice. In order to test the involvement

of the IFN dependent pathway, we crossed STING ki mice with mice deficient for type I IFN

receptor (*Ifnar1*$^{-/-}$); in order to test the involvement of the NF-kB/inflammasome pathway, we

crossed STING ki mice with mice deficient for caspase-1 (*Casp1*$^{-/-}$).

We first analyzed ISGs and NF-kB dependent gene expression by quantitative real time

PCR in adult *Ifnar1*$^{-/-}$ and *Casp1*$^{-/-}$ mice (Figure 6A-E). The N153S STING-induced increase

in the interferon dependent genes *Ifi44 and Mx-1* was abrogated in *Ifnar1*$^{-/-}$ mice (Figure 6A

and B), but unaltered in *Casp1*$^{-/-}$ mice. This is consistent with the dependence of *Ifi44 and*

Mx-1 on type I interferon signaling. The N153S STING-induced increase in expression of *Il-*

1β (Figure 6C) was abrogated in *Casp1*$^{-/-}$ mice, but not in *Ifnar1*$^{-/-}$ mice, consistent with the

fact that *Il-1β* is NF-κB dependent. The N153S STING-induced increase in *Ip-10* expression

was 17-fold in WT (Figure 6D, p= 0,0030844), 12-fold in *Ifnar1*$^{-/-}$ (p= 0,002589) and 6-fold

in *Casp 1*$^{-/-}$ (p= 0,0041598), i.e. not statistically different between WT and *Ifnar1*$^{-/-}$ mice (p=

0,4398) or *Casp 1*$^{-/-}$ mice (p= 0,0623). Expression of *Tnfa* (Figure 6E) was not different

between STING ki and STING WT in this age group, consistent with Figure 5A, and

unaltered in *Ifnar1*$^{-/-}$ and *Casp1*$^{-/-}$ mice. Expression of *Sting1* was also not altered in any on

the genotypes (figure supplement 2A), confirming that alteration in transgene expression did

not underlie the observed effects. Collectively, these analyses confirm the activation of both

IFN signaling and NF-κB/inflammasome signaling in the brain of STING ki mice. *Ifnar1*$^{-/-}$

selectively interferes with the interferon dependent pathway whereas $Casp1^{-/-}$ interferes with

the NF-kB/inflammasome pathway.

We next measured glial activation in $Ifnar1^{-/-}$ and $Casp1^{-/-}$ mice. N153S STING-

induced activation of microglia was still observed in adult $Ifnar1^{-/-}$ and $Casp1^{-/-}$ mice (Figure

6F and H, fold increase and p values in legend). In order to determine whether the extent of

N153S STING-induced microglia activation was significantly different between $Ifnar1^{-/-}$

mice and Ctrl., we calculated the interaction between the factors 'STING genotype' and

'$Ifnar1$ genotype' in two-way ANOVA. The extent of microglia activation was significantly

smaller in adult $Ifnar1^{-/-}$ mice (p= 0,004724) than in Ctrl. Similarly, the extent of N153S

STING-induced microglia activation was significantly smaller in $Casp1^{-/-}$ mice (p=

0,000021). The extent of N153S STING-induced microglia activation did not differ between

$Ifnar1^{-/-}$ and $Casp1^{-/-}$ (p= 0,0805731). Collectively, these findings suggest that N153S

STING-induced microglia activation depends both on type 1 IFN signaling and on NF-

kB/inflammasome signaling.

The extent of N153S STING-induced astroglia activation (Figure 6G and I) was

significantly reduced from 26-fold in Ctrl. to 16-fold in $Ifnar1^{-/-}$ (p=0,0158378 for interaction

of 2-way ANOVA) and to 4-fold in $Casp1^{-/-}$ (p<0.00001 for interaction of 2-way ANOVA),

suggesting that astroglia activation might depend more on NF-kB/inflammasome signaling

than on interferon dependent signaling.

Type I IFN and NF-kB/inflammasome signaling contribute to neurodegeneration in

STING ki mice

Next, we asked whether the N153S STING-induced degeneration of dopaminergic axon

terminals in the striatum is abrogated in adult $Ifnar1^{-/-}$ and $Casp 1^{-/-}$ mice (Figure 7).

Interestingly, the density of dopaminergic axon terminals in the striatum was already lower in

$Ifnar1^{-/-}$ mice and $Casp 1^{-/-}$ mice with WT STING than in control $Ifnar1^{+/+}$ and $Casp1^{+/+}$ mice

(p= 0,0143071 and p= 0,0000248, two-way ANOVA). These findings suggest that $Ifnar1$ and

$Casp1$ are required for proper proliferation, maturation and/or maintenance of dopaminergic

neurons and their axon terminals. Indeed, the Wnt-β-catenin pathway is regulated by

interferons (Kovács et al., 2019) and regulates the differentiation of midbrain dopaminergic

neurons (Szegő et al., 2017). Furthermore, IL-1β induces the differentiation of dopaminergic

neurons (Ling et al., 1998; Rodriguez-Pallares et al., 2005), and we found marginally reduced

$Il-1b$ expression in adult $Casp1^{-/-}$ mice (Figure 6C STING WT).

Both in *Ifnar1*[-/-] and in *Casp 1*[-/-] mice, the N153S STING-induced degeneration of

dopaminergic axon terminals in the striatum was less pronounced than in the control

Ifnar1[+/+-] and *Casp1*[+/+] mice (p= 0,00157 and p= 0,007326 for interaction in two-way

ANOVA). N153S STING-induced fiber loss was not statistically different between *Ifnar1*[-/-]

and *Casp1*[-/-] mice (p= 0,468 for interaction). Consistent with the less pronounced

degeneration of dopaminergic axon terminals, the N153S STING-induced reduction in

striatal dopamine and increase in its metabolism was not observed in *Ifnar1*[-/-] and *Casp1*[-/-]

mice (Figure 7C, figure supplement 2B). These findings suggest that both pathways

contribute to the degeneration of dopaminergic axon terminals, and blocking either pathway

could reduce neurodegeneration.

Discussion

In this work, we demonstrated that the expression of the constitutively active STING variant N153S causes neuroinflammation, followed by degeneration of dopaminergic neurons and aSyn pathology. N153S STING-induced microglia activation and degeneration of dopaminergic neurons involve both type I IFN-dependent and NF-κB/inflammasome-dependend signaling, while astroglia activation might depend predominantly on the NF-κB/inflammasome pathway.

Constitutive STING activation causes neuroinflammation

STING is expressed in the central nervous system. Its expression is highest in microglia, but it can also be detected in astroglia and neurons (Jeffries and Marriott, 2017). Microglia are the primary immune cells of the central nervous system (Wolf et al., 2017) and highly activated upon pathogen invasion or tissue damage. Microglia activation in STING ki mice (Figure 1 and 2) is therefore in line with the increased inflammatory phenotype found in the lungs and spleen of STING ki mice (Luksch et al., 2019; Siedel et al., 2020) and in patients with STING-associated vasculopathy with onset in infancy syndrome (Liu et al. 2014).

Consistent with this morphologically defined neuroinflammatory phenotype, we observed increased expression of the ISGs *Ifi44* and *Mx-1* (Figure 4). These effects are associated with interferon signaling, the main pathway downstream of STING (Decout et al., 2021), demonstrating that expressing the N153S mutant of STING indeed activates interferon dependent pathways. Microglia activation was reduced when N153S STING was expressed in mice deficient for Ifnar1 (*Ifnar1*[-/-], Figure 6H), confirming that the type I IFN pathway is involved in STING-dependent microglia activation as previously demonstrated by others (Warner et al., 2017). However, microglia activation by N153S STING was not completely blocked in *Ifnar1*[-/-] mice, suggesting that additional pathways are involved.

Indeed, next to type I IFN signaling, we also observed increased expression of *Il-1b* and *Tnfa* (Figure 4), confirming the activation of the NF-κB/inflammasome pathway in STING ki mice (Balka et al., 2020; Balka and De Nardo, 2021; B. C. Liu et al., 2018). Consistently, in STING ki mice, nuclear translocation of NF-κB was observed significantly more often (Figure 5). We also observed increased expression of *Casp1*, indicating that the inflammasome pathways are activated by constitutively active STING, consistent with previous work about systemic inflammation (Luksch et al., 2019). Accordingly, the extent of microglia activation by N153S STING was reduced in *Casp1*[-/-] mice (Figure 6H). Therefore,

our results suggest that STING-dependent microglia activation involves inflammasome

activation, in addition to type I IFN-dependent pathways.

Next to the activation of microglia, we also observed activation of astroglia (Figures 1

and 2). STING-induced astroglia activation was only blunted in *Ifnar1*$^{-/-}$ mice, but it was

completely blocked in *Casp1*$^{-/-}$ mice (Figure 6I). This finding suggests that chronic STING-

induced astroglia activation might depend mainly on NF-κB/inflammasome signaling.

Astroglia activation can result directly from the expression of N153S STING in astroglia and

indirectly through the activation of microglia (Kwon and Koh, 2020; Wolf et al., 2017). The

differential effect in *Ifnar1*$^{-/-}$ and *Casp1*$^{-/-}$ mice suggests that astroglia activation is not only a

downstream consequence of microglia activation.

Chronic STING activation leads to the degeneration of dopaminergic neurons

We observed a reduced number of TH-positive neurons in the *substantia nigra* of STING ki

mice, a reduced density of dopaminergic axon terminals and a reduced concentration of

striatal dopamine (Figure 1G-I). These changes occurred in adult mice and were not present

in juvenile mice (Figure 2G-I). The degeneration of dopaminergic neurons thus is a

consequence of the inflammatory changes in our model.

On the *Ifnar1*$^{-/-}$ and *Casp1*$^{-/-}$ backgrounds, the effect of STING ki was difficult to

assess. Yet, the relative reduction in TH fiber density in STING ki mice, as compared to

STING WT mice, was smaller on the *Ifnar1*$^{-/-}$ and *Casp1*$^{-/-}$ backgrounds than in controls

(Figure 7B), as was the extent of dopamine depletion F (Figure 7C). These findings are

consistent with the partial rescue of microglia activation in *Ifnar1*$^{-/-}$ and *Casp1*$^{-/-}$ mice (Figure

6F and H). They suggest that both pathways contribute to the degeneration of dopaminergic

neurons. In further studies, the developmental effects of *Ifnar1* and *Casp1* deficiency could

be circumvented by using conditional knockout mice or pharmacological inhibitors.

In our model, degeneration of dopaminergic neurons and their axon terminals in STING

ki mice is likely a secondary effect of glia activation and secretion of inflammatory cytokines

– in line with previous findings demonstrating a role for inflammation in the pathogenesis of

PD (Hall et al., 2018; Mollenhauer et al., 2019). In addition, constant STING activity within

dopaminergic neurons could contribute to their degeneration – both in our STING ki mice

and in the pathogenesis of PD. Indeed, dopaminergic neurons accumulate oxidative damage

as a consequence of dopamine synthesis and electrical pacemaking activity (Guzman et al.,

2010), and even moderate oxidative stress can stimulate the STING pathway (Sliter et al.,

2018; West et al., 2015). Further work is required to determine the importance of STING

activation in dopaminergic neurons for their degeneration in this model.

Accumulation of protein aggregates in STING ki mice

We observed an accumulation of pathological aSyn and an increased number of Thioflavin S

positive cells in STING ki mice (Figure 3A-E). These findings suggest that chronic STING

activation induces aSyn pathology. They are consistent with the recent observation that

priming rats with a mimic of viral dsDNA precipitates aSyn pathology (Olsen et al., 2019)

and with the aSyn aggregation following viral encephalitis (Bantle et al., 2021).

Inflammatory signals are therefore active promotors of aSyn pathology and not only

responsive to aSyn pathology.

The mechanism by which prolonged neuroinflammation leads to aSyn pathology is still

unknown. Both inflammation and IFN induce the expression of the double-stranded (ds)

RNA-dependent protein kinase (PKR) (Gal-Ben-Ari et al., 2019). PKR can phosphorylate

aSyn at serine 129, resulting in aSyn pathology (Reimer et al., 2018). Moreover, clearance of

aSyn aggregates occurs primarily through autophagy (Ebrahimi-Fakhari et al., 2011), and

INFα increases expression of mammalian target of rapamycin (mTOR) (Liu et al., 2016),

which is expected to reduce autophagy initiation. On the other hand, acute STING activation

can induce autophagy (Hopfner and Hornung, 2020; Y. Liu et al., 2018; Moretti et al., 2017).

aSyn pathology in our model therefore could be explained by an exhaustion of the autophagy

machinery, as it was suggested recently (Bido et al., 2021), and the overall effect of

inflammatory pathways on autophagy and aSyn pathology could be bimodal. Accordingly,

degeneration of dopaminergic neurons and accumulation of aSyn aggregates was also

observed in mice deficient for IFNβ (Ejlerskov et al., 2015; Magalhaes et al., 2021).

Conclusions

In this work, we observed the degeneration of dopaminergic neurons in mice expressing a

constitutively active STING variant. These findings support the hypothesis that chronic

neuroinflammation is sufficient to trigger degeneration of dopaminergic neurons and aSyn

pathology. They indicate that neuroinflammation plays an active role in the pathogenesis of

PD and could be a promising therapeutic target. Further work is required to determine the

precise signaling pathway by which STING activation causes aSyn pathology and dopamine

neuron degeneration.

Declarations

Ethical Approval and Consent to participate

Animals were approved by local authorities (Landesdirektion Dresden)and conducted in accordance with guidelines of the Federation for European Laboratory Animal Science Associations (FELASA).

Consent for publication

All authors approved the manuscript.

Availability of data and materials

The dataset supporting the conclusions of this article is included within the article and its additional files.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the German Research Foundation (DFG) SFB TRR237, B18 and FA-658-3-1.

Authors' contributions

EMSz, ARW, BHF and HL conceived research. EMSz, LM, NB and HL performed research and analyzed data. EMSz, ARW, BHF and HL wrote the first draft of the manuscript. All authors contributed and approved the manuscript.

Acknowledgements

We thank Min Ae Lee-Kirsch (TU Dresden) for very helpful comments on the manuscript and Andrea Kempe, Annett Böhme, Katrin Höhne and Kristin Wogan for excellent technical assistance.

511 ## List of Abbreviations

aSyn	alpha-synuclein
Casp1	Caspase1
cGAS	cyclic GMP-AMP synthase
dsDNA	double strand DNA
GFAP	glial fibrillary acidic protein
Iba1	ionized calcium-binding adapter molecule 1
Ifi44	interferon induced protein 44
IFN	interferon
Ifnar1	interferon alpha receptor1
Il-1β	interleukin 1 beta
Ip-10	interferon-gamma induced protein 10 kD
IRF3	interferon regulatory factor 3
ISG	interferon-stimulated gene
ki	knock in
Mx1	interferon-induced GTP-binding protein
NF-kB	nuclear factor 'kappa-light-chain-enhancer' of activated B-cells
NRLP3	the nucleotide-binding oligomerization domain (NOD), leucine-rich repeat (LRR)-containing protein 3
PD	Parkinson's disease
SAVI	STING-associated vasculopathy with onset in infancy
SN	substantia nigra
STAT3	signal transducer and activator of transcription 3
STING	stimulator of interferon genes
TH	tyrosine hydroxylase
TNFβ	tumor necrosis factor beta
WT	wild type

References

Anderson JP, Walker DE, Goldstein JM, De Laat R, Banducci K, Caccavello RJ, Barbour R, Huang J, Kling K, Lee M, Diep L, Keim PS, Shen X, Chataway T, Schlossmacher MG, Seubert P, Schenk D, Sinha S, Gai WP, Chilcote TJ. 2006. Phosphorylation of Ser-129 is the dominant pathological modification of α-synuclein in familial and sporadic lewy body disease. *J Biol Chem* **281**:29739–29752. doi:10.1074/JBC.M600933200

Balka KR, De Nardo D. 2021. Molecular and spatial mechanisms governing STING signalling. *FEBS J* **288**:5504–5529. doi:10.1111/FEBS.15640

Balka KR, Louis C, Saunders TL, Smith AM, Calleja DJ, D'Silva DB, Moghaddas F, Tailler M, Lawlor KE, Zhan Y, Burns CJ, Wicks IP, Miner JJ, Kile BT, Masters SL, De Nardo D. 2020. TBK1 and IKKε Act Redundantly to Mediate STING-Induced NF-κB Responses in Myeloid Cells. *Cell Rep* **31**. doi:10.1016/J.CELREP.2020.03.056

Bantle CM, Rocha SM, French CT, Phillips AT, Tran K, Olson KE, Bass TA, Aboellail T, Smeyne RJ, Tjalkens RB. 2021. Astrocyte inflammatory signaling mediates α-synuclein aggregation and dopaminergic neuronal loss following viral encephalitis. *Exp Neurol* **346**. doi:10.1016/j.expneurol.2021.113845

Beyer MMSS, Lonnemann N, Remus A, Latz E, Heneka MT, Korte M. 2020. Enduring Changes in Neuronal Function upon Systemic Inflammation Are NLRP3 Inflammasome Dependent. *J Neurosci* **40**:5480–5494. doi:10.1523/JNEUROSCI.0200-20.2020

Bido S, Muggeo S, Massimino L, Marzi MJ, Giannelli SG, Melacini E, Nannoni M, Gambarè D, Bellini E, Ordazzo G, Rossi G, Maffezzini C, Iannelli A, Luoni M, Bagicaluppi M, Gregori S, Nicassio F, Broccoli V. 2021. Microglia-specific overexpression of α-synuclein leads to severe dopaminergic neurodegeneration by phagocytic exhaustion and oxidative toxicity. *Nat Commun 2021 121* **12**:1–15. doi:10.1038/s41467-021-26519-x

Chatterjee K, Roy A, Banerjee R, Choudhury S, Mondal B, Halder S, Basu P, Shubham S, Dey S, Kumar H. 2020. Inflammasome and α-synuclein in Parkinson's disease: A cross-sectional study. *J Neuroimmunol* **338**. doi:10.1016/J.JNEUROIM.2019.577089

Chen H, Zhang SM, Hernán MA, Schwarzschild MA, Willett WC, Colditz GA, Speizer FE, Ascherio A. 2003. Nonsteroidal Anti-inflammatory Drugs and the Risk of Parkinson Disease. *Arch Neurol* **60**:1059. doi:10.1001/archneur.60.8.1059

Chen K, Lai C, Su Y, Bao WD, Yang LN, Xu P-P, Zhu L-Q. 2021. cGAS-STING-mediated IFN-I response in host defense and neuro-inflammatory diseases. *Curr Neuropharmacol* **19**. doi:10.2174/1570159X19666210924110144

Chen Q, Sun L, Chen ZJ. 2016. Regulation and function of the cGAS-STING pathway of
cytosolic DNA sensing. *Nat Immunol* **17**:1142–1149. doi:10.1038/ni.3558

Chou TW, Chang NP, Krishnagiri M, Patel AP, Lindman M, Angel JP, Kung PL, Atkins C,
Daniels BP. 2021. Fibrillar α-synuclein induces neurotoxic astrocyte activation via RIP
kinase signaling and NF-κB. *Cell Death Dis* **12**. doi:10.1038/s41419-021-04049-0

Crow YJ, Casanova J-L. 2014. STING-Associated Vasculopathy with Onset in Infancy — A
New Interferonopathy. *N Engl J Med* **371**:568–571. doi:10.1056/nejme1407246

Decout A, Katz JD, Venkatraman S, Ablasser A. 2021. The cGAS–STING pathway as a
therapeutic target in inflammatory diseases. *Nat Rev Immunol* **21**:548–569.
doi:10.1038/s41577-021-00524-z

Ebrahimi-Fakhari D, Cantuti-Castelvetri I, Fan Z, Rockenstein E, Masliah E, Hyman BT,
McLean PJ, Unni VK. 2011. Distinct Roles In Vivo for the Ubiquitin–Proteasome
System and the Autophagy–Lysosomal Pathway in the Degradation of α-Synuclein. *J*
Neurosci **31**:14508. doi:10.1523/JNEUROSCI.1560-11.2011

Ejlerskov P, Hultberg JG, Wang JY, Carlsson R, Ambjørn M, Kuss M, Liu Y, Porcu G,
Kolkova K, Friis Rundsten C, Ruscher K, Pakkenberg B, Goldmann T, Loreth D, Prinz
M, Rubinsztein DC, Issazadeh-Navikas S. 2015. Lack of Neuronal IFN-β-IFNAR
Causes Lewy Body- and Parkinson's Disease-like Dementia. *Cell* **163**:324–339.

Froula JM, Castellana-Cruz M, Anabtawi NM, Camino JD, Chen SW, Thrasher DR, Freire J,
Yazdi AA, Fleming S, Dobson CM, Kumita JR, Cremades N, Volpicelli-Daley LA.
2019. Defining α-synuclein species responsible for Parkinson's disease phenotypes in
mice. *J Biol Chem* **294**:10392–10406. doi:10.1074/JBC.RA119.007743

Fujiwara H, Hasegawa M, Dohmae N, Kawashima A, Masliah E, Goldberg MS, Shen J,
Takio K, Iwatsubo T. 2002. α-synuclein is phosphorylated in synucleinopathy lesions.
Nat Cell Biol **4**:160–164. doi:10.1038/ncb748

Gaidt MM, Ebert TS, Chauhan D, Ramshorn K, Pinci F, Zuber S, O'Duill F, Schmid-Burgk
JL, Hoss F, Buhmann R, Wittmann G, Latz E, Subklewe M, Hornung V. 2017. The
DNA Inflammasome in Human Myeloid Cells Is Initiated by a STING-Cell Death
Program Upstream of NLRP3. *Cell* **171**:1110-1124.e18.
doi:10.1016/J.CELL.2017.09.039

Gal-Ben-Ari S, Barrera I, Ehrlich M, Rosenblum K. 2019. PKR: A kinase to remember.
Front Mol Neurosci **11**:480. doi:10.3389/FNMOL.2018.00480/BIBTEX

Gordon R, Albornoz EA, Christie DC, Langley MR, Kumar V, Mantovani S, Robertson
AAB, Butler MS, Rowe DB, O'Neill LA, Kanthasamy AG, Schroder K, Cooper MA,

Woodruff TM. 2018. Inflammasome inhibition prevents -synuclein pathology and

581   dopaminergic neurodegeneration in mice. *Sci Transl Med* **10**.

582   doi:10.1126/scitranslmed.aah4066

583   Grozdanov V, Bousset L, Hoffmeister M, Bliederhaeuser C, Meier C, Madiona K, Pieri L,

584   Kiechle M, McLean PJ, Kassubek J, Behrends C, Ludolph AC, Weishaupt JH, Melki R,

585   Danzer KM. 2019. Increased Immune Activation by Pathologic α-Synuclein in

586   Parkinson's Disease. *Ann Neurol* **86**:593–606. doi:10.1002/ana.25557

587   Guzman JN, Sanchez-Padilla J, Wokosin D, Kondapalli J, Ilijic E, Schumacker PT, Surmeier

588   DJ. 2010. Oxidant stress evoked by pacemaking in dopaminergic neurons is attenuated

589   by DJ-1. *Nature* **468**:696–700. doi:10.1038/nature09536

590   Hall S, Janelidze S, Surova Y, Widner H, Zetterberg H, Hansson O. 2018. Cerebrospinal

591   fluid concentrations of inflammatory markers in Parkinson's disease and atypical

592   parkinsonian disorders. *Sci Rep* **8**. doi:10.1038/S41598-018-31517-Z

593   Harms AS, Kordower JH, Sette A, Lindestam Arlehamn CS, Sulzer D, Mach RH. 2021.

594   Inflammation in Experimental Models of α -Synucleinopathies. *Mov Disord* **36**:37–49.

595   doi:10.1002/mds.28264

596   Harris NM, Roy-O'Reilly M, Ritzel RM, Holmes A, Sansing LH, O'Keefe LM, McCullough

597   LD, Chauhan A. 2020. Depletion of CD4 T cells provides therapeutic benefits in aged

598   mice after ischemic stroke. *Exp Neurol* **326**. doi:10.1016/J.EXPNEUROL.2020.113202

599   Heneka MT, McManus RM, Latz E. 2018. Inflammasome signalling in brain function and

600   neurodegenerative disease. *Nat Rev Neurosci* **19**:610–621. doi:10.1038/s41583-018-

601   0055-7

602   Hirsch EC, Standaert DG. 2021. Ten Unsolved Questions About Neuroinflammationnin

603   Parkinson's Disease. *Mov Disord* **36**:16–24. doi:10.1002/mds.28075

604   Hopfner KP, Hornung V. 2020. Molecular mechanisms and cellular functions of cGAS–

605   STING signalling. *Nat Rev Mol Cell Biol* **21**:501–521. doi:10.1038/s41580-020-0244-x

606   Jeffries AM, Marriott I. 2017. Human microglia and astrocytes express cGAS-STING viral

607   sensing components. *Neurosci Lett* **658**:53–56. doi:10.1016/j.neulet.2017.08.039

608   Kovács B, Vajda E, Nagy EE. 2019. Regulatory Effects and Interactions of the Wnt and

609   OPG-RANKL-RANK Signaling at the Bone-Cartilage Interface in Osteoarthritis. *Int J*

610   *Mol Sci* **20**. doi:10.3390/IJMS20184653

611   Kwon HS, Koh SH. 2020. Neuroinflammation in neurodegenerative disorders: the roles of

612   microglia and astrocytes. *Transl Neurodegener* **9**. doi:10.1186/S40035-020-00221-2

Li F, Wang N, Zheng Y, Luo Y, Zhang Y. 2021. cGAS- Stimulator of Interferon Genes

Signaling in Central Nervous System Disorders. *Aging Dis* **12**:1658–1674.

doi:10.14336/AD.2021.0304

Li T, Chen ZJ. 2018. The cGAS-cGAMP-STI NG pathway connects DNA damage to

inflammation, senescence, and cancer. *J Exp Med* **215**:1287–1299.

doi:10.1084/jem.20180139

Ling ZD, Potter ED, Lipton JW, Carvey PM. 1998. Differentiation of mesencephalic

progenitor cells into dopaminergic neurons by cytokines. *Exp Neurol* **149**:411–423.

doi:10.1006/EXNR.1998.6715

Liu BC, Sarhan J, Panda A, Muendlein HI, Ilyukha V, Coers J, Yamamoto M, Isberg RR,

Poltorak A. 2018. Constitutive Interferon Maintains GBP Expression Required for

Release of Bacterial Components Upstream of Pyroptosis and Anti-DNA Responses.

Cell Rep **24**:155-168.e5. doi:10.1016/J.CELREP.2018.06.012

Liu WL, Yang HC, Hsu CS, Wang CC, Wang TS, Kao JH, Chen DS. 2016. Pegylated IFN-α

suppresses hepatitis C virus by promoting the DAPK-mTOR pathway. *Proc Natl Acad*

Sci U S A **113**:14799–14804. doi:10.1073/pnas.1618517114

Liu Y, Gordesky-Gold B, Leney-Greene M, Weinbren NL, Tudor M, Cherry S. 2018.

Inflammation-Induced, STING-Dependent Autophagy Restricts Zika Virus Infection in

the Drosophila Brain. *Cell Host Microbe* **24**:57-68.e3. doi:10.1016/j.chom.2018.05.022

Liu Y, Jesus AA, Marrero B, Yang D, Ramsey SE, Montealegre Sanchez GA, Tenbrock K,

Wittkowski H, Jones OY, Kuehn HS, Lee C-CR, DiMattia MA, Cowen EW, Gonzalez

B, Palmer I, DiGiovanna JJ, Biancotto A, Kim H, Tsai WL, Trier AM, Huang Y, Stone

DL, Hill S, Kim HJ, St. Hilaire C, Gurprasad S, Plass N, Chapelle D, Horkayne-Szakaly

I, Foell D, Barysenka A, Candotti F, Holland SM, Hughes JD, Mehmet H, Issekutz AC,

Raffeld M, McElwee J, Fontana JR, Minniti CP, Moir S, Kastner DL, Gadina M, Steven

AC, Wingfield PT, Brooks SR, Rosenzweig SD, Fleisher TA, Deng Z, Boehm M, Paller

AS, Goldbach-Mansky R. 2014. Activated STING in a vascular and pulmonary

syndrome. *N Engl J Med* **371**:507–518. doi:10.1056/NEJMOA1312625

Luksch H, Stinson WA, Platt DJ, Qian W, Kalugotla G, Miner CA, Bennion BG, Gerbaulet

A, Rösen-Wolff A, Miner JJ. 2019. STING-associated lung disease in mice relies on T

cells but not type I interferon. *J Allergy Clin Immunol* **144**:254-266.e8.

doi:10.1016/j.jaci.2019.01.044

Magalhaes J, Tresse E, Ejlerskov P, Hu E, Liu Y, Marin A, Montalant A, Satriano L,

Rundsten CF, Carlsen EMM, Rydbirk R, Sharifi-Zarchi A, Andersen JB, Aznar S,

Brudek T, Khodosevich K, Prinz M, Perrier JFM, Sharma M, Gasser T, Issazadeh-

Navikas S. 2021. PIAS2-mediated blockade of IFN-β signaling: a basis for sporadic
649   Parkinson disease dementia. *Mol Psychiatry 2021* 1–17. doi:10.1038/s41380-021-
650   01207-w

651 Mejias NH, Martinez CC, Stephens ME, De Rivero Vaccari JP. 2018. Contribution of the
652   inflammasome to inflammaging. *J Inflamm (United Kingdom)* **15**:1–10.
653   doi:10.1186/S12950-018-0198-3/FIGURES/6

654 Mollenhauer B, Zimmermann J, Sixel-Döring F, Focke NK, Wicke T, Ebentheuer J,
655   Schaumburg M, Lang E, Friede T, Trenkwalder C. 2019. Baseline predictors for
656   progression 4 years after Parkinson's disease diagnosis in the De Novo Parkinson
657   Cohort (DeNoPa). *Mov Disord* **34**:67–77. doi:10.1002/mds.27492

658 Moretti J, Roy S, Bozec D, Martinez J, Chapman JR, Ueberheide B, Lamming DW, Chen ZJ,
659   Horng T, Yeretssian G, Green DR, Blander JM. 2017. STING Senses Microbial
660   Viability to Orchestrate Stress-Mediated Autophagy of the Endoplasmic Reticulum. *Cell*
661   **171**:809-823.e13. doi:10.1016/j.cell.2017.09.034

662 Motwani M, Pesiridis S, Fitzgerald KA. 2019. DNA sensing by the cGAS–STING pathway
663   in health and disease. *Nat Rev Genet* **20**:657–674. doi:10.1038/s41576-019-0151-1

664 Nazmi A, Field RH, Griffin EW, Haugh O, Hennessy E, Cox D, Reis R, Tortorelli L, Murray
665   CL, Lopez-Rodriguez AB, Jin L, Lavelle EC, Dunne A, Cunningham C. 2019. Chronic
666   neurodegeneration induces type I interferon synthesis via STING, shaping microglial
667   phenotype and accelerating disease progression. *Glia* **67**:1254–1276.
668   doi:10.1002/glia.23592

669 Neumann M, Kahle PJ, Giasson BI, Ozmen L, Borroni E, Spooren W, Müller V, Odoy S,
670   Fujiwara H, Hasegawa M, Iwatsubo T, Trojanowski JQ, Kretzschmar HA, Haass C.
671   2002. Misfolded proteinase K-resistant hyperphosphorylated alpha-synuclein in aged
672   transgenic mice with locomotor deterioration and in human alpha-synucleinopathies. *J*
673   *Clin Invest* **110**:1429–1439. doi:10.1172/JCI15777

674 Noguchi S, Yamada N, Kumazaki M, Yasui Y, Iwasaki J, Naito S, Akao Y. 2013. socs7, a
675   target gene of microRNA-145, regulates interferon-β induction through STAT3 nuclear
676   translocation in bladder cancer cells. *Cell Death Dis* **4**:e482.
677   doi:10.1038/CDDIS.2013.11

678 Obeso JA, Stamelou M, Goetz CG, Poewe W, Lang AE, Weintraub D, Burn D, Halliday GM,
679   Bezard E, Przedborski S, Lehericy S, Brooks DJ, Rothwell JC, Hallett M, DeLong MR,
680   Marras C, Tanner CM, Ross GW, Langston JW, Klein C, Bonifati V, Jankovic J,
681   Lozano AM, Deuschl G, Bergman H, Tolosa E, Rodriguez-Violante M, Fahn S,

Postuma RB, Berg D, Marek K, Standaert DG, Surmeier DJ, Olanow CW, Kordower

JH, Calabresi P, Schapira AHV, Stoessl AJ. 2017. Past, present, and future of

Parkinson's disease: A special essay on the 200th Anniversary of the Shaking Palsy.

Mov Disord. doi:10.1002/mds.27115

Olsen LK, Cairns AG, Ådén J, Moriarty N, Cabre S, Alamilla VR, Almqvist F, Dowd E,

McKernan DP. 2019. Viral mimetic priming enhances α-synuclein-induced

degeneration: Implications for Parkinson's disease. *Brain Behav Immun* **80**:525–535.

doi:10.1016/j.bbi.2019.04.036

Paul BD, Snyder SH, Bohr VA. 2021. Signaling by cGAS–STING in Neurodegeneration,

Neuroinflammation, and Aging. *Trends Neurosci* **44**:83–96.

doi:10.1016/j.tins.2020.10.008

Paxinos G, Franklin KBJ. 2001. Paxinos and Franklin's the Mouse Brain in Stereotaxic

Coordinates, Academic Press.

Reimer L, Vesterager LB, Betzer C, Zheng J, Nielsen LD, Kofoed RH, Lassen LB, Bølcho U,

Paludan SR, Fog K, Jensen PH. 2018. Inflammation kinase PKR phosphorylates α-

synuclein and causes α-synuclein-dependent cell death. *Neurobiol Dis* **115**:17–28.

doi:10.1016/J.NBD.2018.03.001

Rodriguez-Pallares J, Guerra MJ, Labandeira-Garcia JL. 2005. Angiotensin II and

interleukin-1 interact to increase generation of dopaminergic neurons from neurospheres

of mesencephalic precursors. *Brain Res Dev Brain Res* **158**:120–122.

doi:10.1016/J.DEVBRAINRES.2005.06.009

Samuel F, Flavin WP, Iqbal S, Pacelli C, Renganathan SDS, Trudeau LE, Campbell EM,

Fraser PE, Tandon A. 2016. Effects of Serine 129 Phosphorylation on α-Synuclein

Aggregation, Membrane Association, and Internalization. *J Biol Chem* **291**:4374.

doi:10.1074/JBC.M115.705095

Schroder K, Tschopp J. 2010. The Inflammasomes. *Cell* **140**:821–832.

doi:10.1016/j.cell.2010.01.040

Sen T, Saha P, Gupta R, Foley LM, Jiang T, Abakumova OS, Hitchens TK, Sen N. 2020.

Aberrant ER Stress-Induced Neuronal-IFNβ Elicits White Matter Injury Due to

Microglial Activation and T-Cell Infiltration after TBI. *J Neurosci Off J Soc Neurosci*

40:424–446. doi:10.1523/JNEUROSCI.0718-19.2019

Siedel H, Roers A, Rösen-Wolff A, Luksch H. 2020. Type I interferon-independent T cell

impairment in a Tmem173 N153S/WT mouse model of STING associated vasculopathy

with onset in infancy (SAVI). *Clin Immunol* **216**:108466.

doi:10.1016/j.clim.2020.108466

Sliter DA, Martinez J, Hao L, Chen X, Sun N, Fischer TD, Burman JL, Li Y, Zhang Z,

Narendra DP, Cai H, Borsche M, Klein C, Youle RJ. 2018. Parkin and PINK1 mitigate

STING-induced inflammation. *Nature*. doi:10.1038/s41586-018-0448-9

Spandidos A, Wang X, Wang H, Seed B. 2009. PrimerBank: A resource of human and mouse

PCR primer pairs for gene expression detection and quantification. *Nucleic Acids Res*

38:D792–D799.

Szegő ÉM, Boß F, Komnig D, Gärtner C, Höfs L, Shaykhalishahi H, Woerdehoff M, Saridaki

T, Schulz JB, Hoyer W, Falkenburger BH. 2021. A β-wrapin targeting the N-terminus of

α-synuclein monomers reduces fibril-induced aggregation in neurons. *Front Neurosci*

15:751.

Szegő ÉM, Gerhardt E, Kermer P, Schulz JB. 2012. A30P α-synuclein impairs dopaminergic

fiber regeneration and interacts with L-DOPA replacement in MPTP-treated mice.

Neurobiol Dis **45**:591–600.

Szegő ÉM, Gerhardt E, Outeiro TF. 2017. Sirtuin 2 enhances dopaminergic differentiation

via the AKT/GSK-3β/β-catenin pathway. *Neurobiol Aging* **56**:7–16.

Szegö EM, Van den Haute C, Höfs L, Baekelandt V, Van der Perren A, Falkenburger BH.

2022. Rab7 reduces α-synuclein toxicity in rats and primary neurons. *Exp Neurol*

347:113900. doi:10.1016/J.EXPNEUROL.2021.113900

von Herrmann KM, Salas LA, Martinez EM, Young AL, Howard JM, Feldman MS,

Christensen BC, Wilkins OM, Lee SL, Hickey WF, Havrda MC. 2018. NLRP3

expression in mesencephalic neurons and characterization of a rare NLRP3

polymorphism associated with decreased risk of Parkinson's disease. *npj Park Dis* **4**:24.

doi:10.1038/s41531-018-0061-5

Wang W, Hu D, Wu C, Feng Y, Li A, Liu W, Wang Y, Chen K, Tian M, Xiao F, Zhang Q,

Shereen MA, Chen W, Pan P, Wan P, Wu K, Wu J. 2020. STING promotes NLRP3

localization in ER and facilitates NLRP3 deubiquitination to activate the inflammasome

upon HSV-1 infection. *PLoS Pathog* **16**. doi:10.1371/JOURNAL.PPAT.1008335

Warner JD, Irizarry-Caro RA, Bennion BG, Ai TL, Smith AM, Miner CA, Sakai T,

Gonugunta VK, Wu J, Platt DJ, Yan N, Miner JJ. 2017. STING-associated vasculopathy

develops independently of IRF3 in mice. *J Exp Med* **214**:3279–3292.

doi:10.1084/JEM.20171351

West AP, Khoury-Hanold W, Staron M, Tal MC, Pineda CM, Lang SM, Bestwick M,

Duguay BA, Raimundo N, MacDuff DA, Kaech SM, Smiley JR, Means RE, Iwasaki A,

Shadel GS. 2015. Mitochondrial DNA stress primes the antiviral innate immune

response. *Nature* **520**:553–557. doi:10.1038/nature14156

Wolf SA, Boddeke HWGM, Kettenmann H. 2017. Microglia in Physiology and Disease.

Annu Rev Physiol **79**:619–643. doi:10.1146/ANNUREV-PHYSIOL-022516-034406

Figure Legends

Figure 1. Constitutive STING activation induces neuroinflammation and neurodegeneration in adult mice.

(A) Representative images of striatal sections from STING WT and STING ki mice stained for the microglia marker Iba1. Scale bar: 50 μm. (B) Representative images of striatal sections stained for the astroglia marker GFAP. Scale bar: 50 μm. (C) Representative images of midbrain sections containing the *substantia nigra* (SN) from STING WT and STING ki mice (stitched from two microscopy fields) stained for tyrosine hydroxylase (TH). Scale bar: 100 μm. (D) Representative images of striatal sections stained for TH from STING WT and STING ki mice. Scale bar: 10 μm. (E) Area fraction positive for Iba1, normalized to the mean of STING WT mice. Markers represent individual animals (black: STING WT animals, red: STING ki animals). Lines represent mean ± SD. Comparison by t-test (***: p=0,0007). (F) Area fraction positive for GFAP, normalized to the mean of STING WT mice (**: p=0,0011; t-test). (G) Number of TH-positive neurons (*: p=0,0257; t-test). (H) Area fraction positive for TH (**: p=0,0081; t-test). (I) Concentration of dopamine (*: p=0,0448; t-test) in striatal lysates from STING WT and STING ki animals, normalized to the mean concentration in STING WT. Graphs showing quantification of the dopamine metabolites are in figure supplement 1B.

Figure 2. Neuroinflammation without neurodegeneration in juvenile mice with constitutive STING activation.

(A) Representative images of striatal sections stained for the microglia marker Iba1 from 5 week-old STING WT and STING ki mice. Scale bar: 50 μm (B) Representative images of striatal sections stained for the astroglia marker GFAP from 5 week-old STING WT and STING ki mice. Scale bar: 50 μm (C) Representative images of midbrain sections containing the *substantia nigra* (SN, stitched from two microscopy fields) stained for tyrosine hydroxylase (TH) from 5 week-old STING WT and STING ki mice. Scale bar: 100 μm (D) Representative images of striatal sections stained for TH from 5 week-old STING WT and STING ki mice. Scale bar: 10 μm. (E) Area fraction positive for Iba1, normalized to the mean of STING WT (***: p=0,0009; t-test). (F) Area fraction positive for GFAP, normalized to the mean of STING WT brains (***: p=0,0007; t-test). (G) Number of TH-positive neurons (mean ± SD; t-test). (H) Area fraction positive for TH (mean ± SD, t-test). (I) Dopamine concentration in striatal lysates from 5 week-old STING WT and STING ki mice, measured by HPLC and normalized

to the mean of STING WT (mean ± SD, t-test). Dopamine metabolites are in figure supplement

1D.

Figure 3. Constitutive STING activation induces alpha-synuclein pathology and synapse

loss in adult mice.

(A) Representative Western blot images showing phosphorylated alpha-synuclein (S129;

paSyn, upper panel), total alpha-synuclein (aSyn, middle panel) and the loading control βIII-

tubulin (lower panel). (B) Representative images of striatal sections from 20 week-old STING

WT and STING ki mice stained with Thioflavin S. Scale bar: 10 μm (C) Ratio of total alpha-

synuclein and loading control, expressed relative to the mean of STING WT (p=0,0491; mean

± SD; t-test). (D) Immunoreactivity to paSyn and aSyn, expressed relative to the mean of

STING WT (p=0,0059; mean ± SD; t-test). (E) Number of cells with inclusions positive for

Thioflavin S (ThioS) per mm^2 (*: p=0,0141; t-test). (F-G) Representative images of striatal

sections from 20 week-old STING WT and STING ki mice stained for the presynaptic marker

synapsin (F) or for the post-synaptic marker homer (G). Scale bar: 10 μm. (H-I) Area fraction

positive for synapsin (H, p=0,0053) or homer (I, p=0,0408) (mean ± SD; t-test).

Figure 4. Activation of IFN and NF-κB/inflammasome dependent signaling in juvenile

and adult STING ki mice.

(A-D) Expression of ISGs in the frontal cortex of STING WT and STING ki mice. (A) *Ifi44*

(***: p=0,0002277; *: p= 0,044987), (B) *Mx-1* (***: p=0,0000003; *: p= 0,016835; for

comparison between age groups ***: p= 0,000602), (C) *Ip-10* (***: p= 0,000001; **: p=

0,0017215; for comparison between age groups **: p= 0,001483), (D) *Sting1* (*: p=

0,0184042). (E-G) Expression of NF-κB/inflammasome related genes in the frontal cortex of

STING WT and STING ki mice. (E) *Tnfα* (***: p= 0,0001448, for comparison between age

groups *: p= 0,03952). (F) *Il-1b* (***: p= 0,00005, *: p= 0,0389, for comparison between age

groups **: p= 0,00241). (G) *Casp1* (***: p= 0,0000369, for comparison between age groups

***: p= 0,000064). Markers represent individual animals, bars represent mean ± SD. Analysis

was two-way ANOVA with Tukey HSD post-hoc test.

Figure 5. Nuclear translocation of pSTAT3 and NF-κB in the striatum of 5 and 20 week-

old STING WT and STING ki mice.

(A) Representative images of striatal sections from 5 week-old (upper images) and 20 week-

old (lower images) STING WT and STING ki mice stained for Iba1 (green), GFAP (white) and

phosphorylated-STAT3 (pSTAT3; red). Images show color coded merged channels (center)

and in addition pSTAT3 staining in grayscale (left and right). Scale bar: 10 μm. (B)

Representative images of striatal sections from 4 week-old (upper images) and 20 week-old

(lower images) STING WT and STING ki mice stained for Iba1 (green), GFAP (white) and

NF-κB (red) . NF-kB staining is shown in gray in separate images. Scale bar: 10 μm. (C)

Number of pSTAT3-positive nuclei/mm^3 (***: p=0,00004; **: p=0,0025 for the interaction;

two-way ANOVA, Bonferroni post-hoc test). (D) Number of NF-kB-positive nuclei/mm^3 (**:

p=0,009; ***: p=0,0007; mean ± SD; two-way ANOVA, Bonferroni post-hoc test).

Figure 6. Neuroinflammation in adult double transgenic mice with STING ki and knock-

out for Ifnar1 or Caspase-1.

(A-E) Expression of IFN related genes and NF-κB/inflammasome related genes in the frontal

cortex of adult STING WT and STING ki mice (ANOVA with Tukey HSD post-hoc test). (A)

Ifi44 (*: p= 0,01414; **: p= 0,0037655; for interaction between Ctrl. and and *Ifnar1$^{-/-}$*, **: p=

0,005101). (B) *Mx-1* (*: p= 0,02823; **: p= 0,00573). (C) *Il-1b* (Ctrl. background *: p=

0,04534; **: p= 0, 005405; *Casp1$^{-/-}$ background* *: p= 0,0107096; for interaction between Ctrl.

and *Casp1$^{-/-}$* , *: p= 0,01298). (D) *Ip-10* (Ctrl. background **: p= 0,0030844; *Ifnar1$^{-/-}$*

background: p= 0,025893; *Casp1$^{-/-}$ background* **: p= 0,0041598). (E) *Tnfα* (all differences

n.s.). *Sting1* expression is shown on figure supplement 2A. (F) Representative images of striatal

sections stained for the microglia marker Iba1. Sections were obtained from adult STING WT

(upper images) or STING ki (lower images) mice on a background of interferon a receptor

knockout (*Ifnar1$^{-/-}$*), caspase 1 knockout (*Casp1$^{-/-}$*) or *Ifnar1$^{+/+}$*, *Casp1$^{+/+}$* (Ctrl.). Scale bar: 50

μm. (G) Representative images of striatal sections stained for the astroglia marker GFAP from

STING WT (upper images) or STING ki (lower images) mice on a background of interferon a

receptor knockout (*Ifnar1$^{-/-}$*), caspase 1 knockout (*Casp1$^{-/-}$*) or *Ifnar1$^{+/+}$*, *Casp1$^{+/+}$* (Ctrl.). Scale

bar: 50 μm. (H) Area fraction positive for Iba1, normalized to the mean of STING WT brains

(differences in $^{+/+}$ mice ***: p= 0,0000001; for *Ifnar1$^{-/-}$* ***: p=0,000003; for *Casp1$^{-/-}$* ***: p=

0,0029374; two-way ANOVA with Bonferroni post-hoc test). (I) Area fraction positive for

GFAP, normalized to STING WT on Ctrl. Background (***: p= 0,0000 for STING WT vs

STING ki on Ctrl.; ***: p= 0,0000 on *Ifnar1$^{-/-}$*; background, ***: p=xyz on Casp1-/-

background; two-way ANOVA with Bonferroni post-hoc test).

Figure 7. Degeneration of dopaminergic neurons in double transgenic mice with STING

N153S/WT ki and knock-out for Ifnar1 or Caspase-1.

(A) Representative images of striatal sections stained for tyrosine hydroxylase (TH) from STING WT (upper images) or STING ki (lower images) mice on a background of interferon a receptor knockout (*Ifnar1*$^{-/-}$), caspase-1 knockout (*Casp1*$^{-/-}$) or *Ifnar1*$^{+/+}$, *Casp1*$^{+/+}$ (Ctrl.). Scale bar: 10 μm. (B) Area fraction positive for TH, normalized to STING WT on Ctrl. background (***: p=0,0000 for Ctrl. background; *: p= 0,043 for *Ifnar1*$^{-/-}$; **: p=0,0126845 for *Casp1*$^{-/-}$; for interaction between Ctrl. background and *Ifnar1*$^{-/-}$: p= 0,00157; between Ctrl. and *Casp1*$^{-/-}$: p= 0,007326; two-way ANOVA with Bonferroni post-hoc test). (C) Concentration of dopamine in striatal lysates of STING WT and STING ki mice, normalized to STING WT on Ctrl. background. (***: p= 0,0005; t-test). Dopamine metabolism is shown on figure supplement 2B.

<u>Supplemental Information</u>

Figure supplement 1, related to Figures 1 and 2. Number of TH-positive neurons in the

***substantia nigra* and dopamine metabolism in the striatum of STING WT and STING ki**

mice

(A) Number of TH-positive neurons in the substantia nigra of adult mice (p=0,0257; t-test). (B)

Dopamine metabolism (concentration of dopamine metabolites DOPAC + HVA) / dopamine

in adult mice (p=0,0179; t-test). (C) Number of TH-positive neurons in the substantia nigra of

juvenile mice (p=0,5188; t-test). (D) Dopamine metabolism in juvenile mice (p=0,9545; t-test).

Figure supplement 2, related to Figures 6 and 7. *Sting1* expression and dopamine

metabolism in KO animals.

(A) Expression of *Sting1* in the frontal cortex of STING WT and STING ki mice on a

background of interferon a receptor knockout (*Ifnar1$^{-/-}$*), caspase 1 knockout (*Casp1$^{-/-}$*) or

Ifnar1$^{+/+}$, *Casp1$^{+/+}$* (Ctrl.). (all differences n.s., 2-way ANOVA). (B) Dopamine metabolism

in STING WT and STING ki mice. (**: p= 0,0073; 2-way ANOVA).

Supplemental Table S1.– List of materials and antibody dilutions

Antibody	Dilution	Source	ID
Anti-Iba1, rabbit	1:1000	Fujifilm Wako Chemicals	Cat# 019-19741
Anti-GFAP, chicken	1:2000	Abcam	Cat# ab4674
Anti-NFκB p65 (D14E12), rabbit	1:500	Cell Signaling Technology	Cat# 8242
Anti-Tubulin βIII, rabbit	1:5000	Covance	Cat# PRB-435P
Anti-phospho-a-synuclein, rabbit	1:1000	Abcam	Cat# ab51253
Anti-a-synuclein, mouse	1:2000	BD Biosciences	Cat# 610787
Anti-TH, sheep	1:2000	Pel Freez	Cat# P40101
Anti-homer, rabbit	1.1000	Synaptic Systems	Cat# 160002
Anti-Synapsin, chicken	1:1000	Synaptic System	Cat# 106006
HRP conjugated donkey anti-mouse IgG	1:5000	Jackson ImmunoResearch	Cat# 715-035-150
HRP conjugated donkey anti-rabbit IgG	1:5000	Jackson ImmunoResearch	Cat# 711-035-152
Alexa 555 conjugated donkey anti-rabbit	1:2000	Invitrogen	Cat# A31572
Alexa 647 conjugated donkey anti-chicken	1:2000	Jackson ImmunoResearch	Cat# 703-605-155
Alexa 488 conjugated donkey anti-sheep	1:2000	Invitrogen	Cat# A11015
Alexa 488 conjugated donkey anti-rabbit	1:2000	Invitrogen	Cat# A21206
Reagent			
PhosSTOP phosphatase inhibitor		Sigma-Aldrich	Cat# 4906845001
Bromphenol blue		Honeywell Fluka	Cat# 32712
Mini-PROTEAN TGX Precast Gels 4-20%		BioRad	Cat# 4561094
Precision Plus Protein WesternC Protein Standards		BioRad	Cat# 1610376
WesternBright Chemilumineszenz Substrat Sirius		Biozym Scientific GmbH	Cat# 541019
M-MLV Reverse Transriptase		Promega	Cat# M1701
RNasin Ribonuclease Inhibitor		Promega	Cat# N2511
GoTaq® qPCR Master Mix		Promega	Cat# A6002
Deoxynucleotide Triphosphates (dNTPs)		Promega	Cat# U1205
M-MLV 5x Reaction Buffer		Promega	Cat# M1701
Pierce Micro BCA Protein-Assay-Kit		Thermo Fisher Scientific Inc.	Cat# 23235
SV Total RNA Isolation System		Promega	Cat# Z3101
Fluoromount-G		Southern Biotech	Cat# 0100-01
Equipment			
Infinite 200M Multimode-Plate-Reader		Tecan Group	
QuantStudio 5 Real-Time PCR System		Thermo Fisher Scientific Inc.	
CM3050S Kryostat		Leica Mikrosysteme Vertrieb GmbH	
Zeiss Axio Observer Z1		Carl Zeiss	
Zeiss Spinning Disc		Carl Zeiss	
Software			
GraphPad Prism 5. 01 and 9.0.0		GraphPad Software	
ImageJ Fiji		Wayne Rasband (NIH)	
QuantStudio5 qPCR Data Analysis Software		Thermo Fisher Scientific Inc.	

Supplemental Table S2. – List of qRT PCR Primers

Gene	Primer forward (5' - 3')	Primer revers (5' - 3')
Ifi44	*AACTGACTGCTCGCAATAATGT*	*GTAACACAGCAATGCCTCTTGT*
Mx-1	AACCCTGCTACCTTTCAA	AAGCATCGTTTTCTCTATTTC
Sting1	CTGCTGACATATACCTCAGTTG	GAGCATGTTGTTATGTAGCTG
Ip-10	CCAAGTGCTGCCGTCATTTTC	GGCTCGCAGGGATGATTTCAA
Tnfα	CCTGTAGCCCACGTCGTAG	GGGAGTAGACAAGGTACAACCC
Casp1	GCTGCCTGCCCAGAGCACAAG	CTCTTCAGAGTCTCTTACTG
Il-1β	GAAATGCCACCTTTTGACAGTG	TGGATGCTCTCATCAGGACAG
Hprt1	TCAGTCAACGGGGGACATAAA	GGGGCTGTACTGCTTAACCAG
Rpl13a	AGCCTACCAGAAAGTTTGCTTAC	GCTTCTTCTTCCGATAGTGCATC
Eef2	CCGACTCCCTTGTGTGCAA	AGTTCAGGTCGTTCTCAGAGAG

Supplemental Table S4. – Sample numbers in each analysis

Assay	Group -- number of animals used/excluded						Reason*
Western blot	**STING WT adult**	**STING ki adult**					
	5/0	5/0					
qPCR	**STING WT adult**	**STING ki adult**	**STING WT juvenile**	**STING ki juvenile**			
Ifi44	5/0	5/0	5/0	5/0			
Ip-10	4/1	4/1	5/0	5/0			Outlier
Mx-1	4/1	4/1	5/0	5/0			Outlier
Tnfa1	5/0	5/0	5/0	5/0			
Casp1	5/0	4/1	5/0	5/0			Outlier
Il-1b	4/1	5/0	4/1	5/0			Outlier
Sting1	5/0	5/0	5/0	4/1			Outlier
	STING WT	**STING ki**	**STING WT Ifnar1⁻/⁻**	**STING ki Ifnar1⁻/⁻**	**STING WT Casp1⁻/⁻**	**STING ki Casp1⁻/⁻**	
Ifi44	5/0	5/0	5/0	5/0	4/1	5/0	Outlier
Ip-10	4/1	4/1	4/1	4/1	4/1	4/1	Outlier
Mx-1	4/1	4/1	4/1	5/0	5/0	5/0	Outlier
Tnfa1	5/0	5/0	5/0	4/1	5/0	5/0	Outlier
Il6	4/1	5/0	5/0	4/1	4/1	4/1	Outlier
Casp1	5/0	5/0	5/0	5/0	5/0	5/0	Outlier
Il-1b	5/0	5/0	4/1	5/0	4/1	5/0	Outlier
Histology	**STING WT adult**	**STING ki adult**	**STING WT juvenile**	**STING ki juvenile**			
	5/0	5/0	5/0	6/0			
	STING WT	**STING ki**	**STING WT, Ifnar1⁻/⁻**	**STING ki, Ifnar1⁻/⁻**	**STING WT, Casp1⁻/⁻**	**STING ki, Casp1⁻/⁻**	
	5/0	5/0	5/0	6/0	5/0	5/0	
HPLC	**STING WT adult**	**STING ki adult**	**STING WT juvenile**	**STING ki juvenile**			
	5/0	5/0	5/0	5/0			
	STING WT	**STING ki**	**STING WT, Ifnar1⁻/⁻**	**STING ki, Ifnar1⁻/⁻**	**STING WT, Casp1⁻/⁻**	**STING ki, Casp1⁻/⁻**	
	8/0	7/0	3/0	3/0	9/0	6/0	

*Grubbs`test was used to detect outliers (Prism).

Figure 1

Figure 2

A STING WT   STING ki

kDa

14 —   paSyn

14 —   aSyn

55 —   βIII tubulin

B STING WT   STING ki

Thio S striatum

10 μm

● STING WT   ● STING ki

C
aSyn/βIIItubulin
(fold STING WT)
*

D
paSyn/aSyn
(fold STING WT)
**

E
Cells with ThioS
(fold STING WT)
*

F STING WT   STING ki

synapsin striatum

G

homer striatum

10 μm

H
Synapsin density
(fold STING WT)
**

I
Homer density
(fold STING WT)
*

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

A

TH striatum

STING WT

STING ki

Ctrl. *Ifnar1*^{-/-} *Casp1*^{-/-}

10 μm

B

TH fiber density
(fold STING WT)

Ctrl. *Ifnar1*^{-/-} *Casp1*^{-/-}

C

Dopamine
(fold STING WT)

Ctrl. *Ifnar1*^{-/-} *Casp1*^{-/-}

● STING WT ● STING ki

Supplemental Figure 1

Supplemental Figure 2

A *Sting1*

B

STING WT   STING ki