A Split-GAL4 screen identifies novel sleep-promoting neurons in the Ventral Nerve Cord of *Drosophila*

Joseph Jones[1], Brandon Holder[1], Kiran Eiken[1], Alex Vogt[1], Adriana Velarde[1], Ally Elder[1], Jennifer McEllin[1] and Stephane Dissel[1*]

[1]Department of Genetics, Developmental and Evolutionary Biology, School of Biological and Chemical Sciences, University of Missouri-Kansas City, 5007 Rockhill Rd, Kansas City, MO 64110

*Correspondence to: dissels@umkc.edu

Abstract

As in the mammalian system, sleep in *Drosophila* is regulated by multiple brain regions. Among them, neurons projecting to the dorsal Fan-Shaped Body (dFB) have been intensively studied and the data suggest they play a critical role for in sleep regulation. The 23E10-GAL4 driver is the most widely used tool to label and manipulate dFB neurons. Multiple studies have reported that activation of 23E10-GAL4 neurons promotes sleep. However, anatomical analysis revealed that 23E10-GAL4 labels 23-30 dFB neurons in the *Drosophila* brain and many non-dFB neurons in the brain and in the

Ventral Nerve Cord (VNC), the fly equivalent of the spinal cord. To better understand

the role of individual dFB neurons in sleep regulation, we undertook a Split-GAL4

screen to gain access to subsets of 23E10-GAL expressing cells. In this study, we

report the discovery of two VNC cholinergic sleep-promoting neurons labeled by the

23E10-GAL4 driver.

Introduction

Understanding the neural basis of behavior is a major aspect of neurobiology. However,

unequivocally assigning a behavior to a specific neuron or group of neurons is not a

trivial task. The extreme complexity, diversity and connectivity of the mammalian

nervous system renders this task even more daunting. To study the neural basis of a

behavior, an investigator must be able to specifically manipulate a distinct group of cells

and monitor the behavior of interest. Such approaches require precise genetic tools that

allow for the manipulation of discrete neurons or groups of neurons. While such tools

exist in mammalian systems (1), they may not be available for all the diverse types of

neurons that underlie a specific behavior. It is therefore not surprising that animal

models have been extensively used to untangle the neural basis of many different

behaviors (2). One such model, the fruit fly *Drosophila melanogaster*, with its simpler

nervous system and unparalleled genetic toolbox, has become a workhorse in the study

of complex behaviors, such as courtship (3, 4), feeding (5, 6), and sleep (7). Multiple

binary systems have been developed to access and manipulate specific groups of

neurons in the fly nervous system: GAL4/UAS (8); LexA/LexAop (9) and; QF/QUAS

(10). Of these 3 systems, the GAL4/UAS system has been by far the most extensively

used. However, the expression patterns of GAL4 drivers are often not restricted enough

to clearly link a behavior to specific neurons. Importantly, refinement of GAL4

expression pattern can be achieved by employing the intersectional Split-GAL4

technology (11).

Sleep is a behavior that has been described in a variety of species ranging from jellyfish

to humans (12). Sleep is regulated by two processes, the circadian clock which gates

the occurrence of sleep, and the sleep homeostat which controls the intensity and

duration of sleep, in response to prior wakefulness (13). Although the precise function of

sleep still remains unknown, there is ample evidence supporting the notion that sleep is

required for maintaining optimal physiological and behavioral performance (14). A priori,

sleep could appear to be a detrimental activity as it competes with other motivated

behaviors such as feeding, mating, or parenting, and renders organisms defenseless

against potential predators. However, despite these negative outcomes, sleep has been

maintained throughout evolution, emphasizing its essential value (15).

Over the last 20 years, multiple studies have clearly demonstrated that the mechanisms

and regulation of sleep are largely conserved from flies to mammals (16). Like in the

mammalian system, sleep-regulating centers are found in many areas in the *Drosophila*

brain (16, 17). One such sleep-regulating center that has gathered a lot of attention

comprises neurons that project to the dorsal Fan-Shaped body (dFB). The Fan-Shaped

Body (FB) is part of the central complex in the *Drosophila* brain, a region organized into

multiple layers that plays a role in many functions, including locomotion control (18),

courtship behavior (19) and memory (20, 21), nociception (22), visual feature

recognition (23) and processing (24), social behaviors (25) and, feeding decision-

making (26).

Previous studies demonstrated that increasing the activity of neurons contained in the

C5-GAL4, 104y-GAL4, and C205-GAL4 drivers strongly promotes sleep (27, 28). While

the expression patterns of these 3 independent drivers are broad, they show prominent

overlap in neurons that project to the dFB (27). Based on these observations, the

authors concluded that it is likely that the dFB plays a role in regulating sleep but could

not rule out a role for neurons outside the dFB (27). In addition, further studies

established that reducing the excitability of 104y-GAL4 neurons reduces sleep (29, 30).

Highlighting the strong interaction between sleep and memory, activation of 104y-GAL4

neurons consolidates Short-Term Memory (STM) to Long-Term Memory (LTM) (27, 31)

and restores STM to the classical memory mutant *rutabaga* and *dunce* (32). Taken

together, these data point at dFB neurons as important modulator of sleep.

Further work supported a role for the dFB in sleep homeostasis by demonstrating that

sleep deprivation increases the excitability of 104y-GAL4 dFB neurons (29). A follow up

study proposed that increasing sleep pressure switches dFB neurons from an

electrically silent to an electrically active state and that this process is regulated by

dopaminergic signaling to the dFB (33) and the accumulation of mitochondrial reactive

oxygen species in dFB neurons (34). The physiological properties of dFB neurons have

led to the suggestion that dFB neurons are functionally analogous to the ventrolateral

preoptic nucleus (VLPO), a key sleep-regulating center in the mammalian brain (29, 35).

Importantly, recent studies have used 23E10-GAL4, a more restrictive driver to

manipulate and monitor dFB neurons. (33, 36). Several groups have demonstrated that

chronic and acute activation (using thermogenetic or optogenetic methods) of 23E10-

GAL4 neurons increases sleep (31, 37-39). Because of its more restricted expression

pattern and strong capacity to modulate sleep, 23E10-GAL4 (as it relates to sleep) is

seen as a dFB specific driver by most in the scientific community. Remarkably, a recent

study described an electron microscopy-based connectome of the central complex

including the dFB and demonstrated that the 23E10-GAL4 driver contains 31 neurons in

the brain that project to the dFB (40). Whether these 31 dFB neurons can be considered

as a functionally homogeneous group is still unclear. Interestingly, Qian *et al*.

demonstrated that only a small subset of 23E10-GAL4 neurons express the 5HT2b

serotonin receptor and that these cells regulate sleep and sleep homeostasis (37).

Unfortunately, this study did not investigate the role of non-5HT2b expressing 23E10-

Gal4 neurons in sleep regulation. In addition, whether there are non-dFB 5HT2b

expressing 23E10-GAL4 neurons that could potentially play a role in sleep modulation is

not clearly addressed in this study either. Taken together, these data suggest that

23E10-GAL dFB neurons probably constitute a functionally heterogenous group of cells.

Surprisingly, activation of 23E10-GAL4 expressing neurons, unlike 104y-GAL4, does

not consolidate STM to LTM (31). In this study, the authors demonstrated that in

addition to the dFB, 104y-GAL4 expresses in neurons projecting to the ventral FB (vFB)

and that these neurons also regulate sleep and are responsible for memory

consolidation (31). These data emphasize the need to use genetic tools that are as

specific as possible to clearly assign a role in sleep regulation and sleep function to

specific neurons/groups of neurons. Often, GAL4 drivers that are extensively used to

manipulate a specific neuronal group also express in regions outside the area of

interest. This makes it difficult to clearly identify the neural basis of sleep behavior.

With this in mind, we wanted to know which of the 31 dFB 23E10-GAL4 neurons are

important for sleep regulation. To do so, we employed a targeted, intersectional Split-

GAL4 strategy (11) focused on 23E10-GAL4 dFB neurons. Surprisingly, our data

demonstrates that in the 23E10-GAL driver, there are non-dFB sleep-promoting

neurons. Importantly, these cholinergic neurons are located in the Ventral Nerve Cord

(VNC), the fly equivalent to the spinal cord. Nevertheless, we confirm that 23E10-GAL4

dFB neurons regulate sleep homeostasis while 23E10-GAL4 VNC neurons do not.

Thus, our work reveals that there are multiple groups (at least two) of sleep-regulating

neurons in an extensively used "dFB-specific" GAL4 driver. These data clearly highlight

the need to use extremely specific tools to manipulate limited number of neurons when

trying to unequivocally assign a behavior to a group of cells.

Results

The 23E10-GAL4 driver promotes sleep and contains many non dFB neurons in

the central nervous system

Previous studies have reported that increasing the activity of dFB neurons using the

23E10-GAL driver reliably increases sleep (31, 37-39). We first sought to confirm these

observations in our laboratory. Thus, we expressed the thermogenetic TrpA1 cation

channel, which is activated by transferring flies to 31°C (41), in 23E10-GAL4 expressing

neurons. Individual flies were monitored for two days at 22°C (baseline days, no

activation), then temperature was raised to 31°C for 24h on day 3 (activation day). A

schematic of the experimental setup is provided in Figure 1A. As seen in Figure 1B,C,

raising the temperature changes the sleep profile of both parental controls and results in

a loss of sleep (Figure 1E). These data are in agreement with previous reports on the

effect of high temperature on sleep (42, 43). On the other hand, activating 23E10-GAL4

neurons strongly increases sleep (Figure 1D,E). Importantly, activation of 23E10-GAL4

neurons does not affect the amount of locomotor activity when the flies are awake

(Figure 1F). Thus, we confirmed that activating 23E10-GAL4 neurons promotes sleep

and that the increase in sleep is not caused by a general motor defect, in agreement

with previous reports using the same GAL4 driver (31, 37-39) .

23E10-GAL4 has become the driver of choice to manipulate dFB neurons and many in

the field see it as a "dFB-specific" tool. However, GAL4 lines often are not that specific

and commonly express in neurons outside the region of interest. To clarify the

expression pattern of 23E10-GAL4, we expressed GFP and carefully quantified the

number of neurons that are GFP positive. As seen in Figure 1G,H, there are many

neurons labelled by 23E10-GAL4 in the brain and in the VNC (See also Video 1 and

Video 2). Importantly, many of these neurons are not dFB neurons. Our analysis

revealed that there are more than 50 neurons in the brain labelled by 23E10-GAL4, only

half of which are dFB neurons (Table1). In addition, 23E10-GAL labels about 18

neurons in the VNC (Table1). Interestingly, we note that while the electron microscopy

study discussed above identified 31 dFB 23E10-GAL4 neurons (40), we do not

systematically observe 31 neurons in every brain. This discrepancy is probably due to

the different methodologies used in the two studies. We suspect that some 23E10 dFB

neurons may be only weakly expressing GAL4 and are therefore not always visible

using confocal microscopy. However, based on this expression pattern, we conclude

that it is impossible to unequivocally assign a role in sleep promotion to 23E10-GAL4

dFB neurons.

The 23E10-GAL4 driver contains sleep-promoting neurons that are not dFB

neurons

To shed light on the role of individual 23E10-GAL4 expressing neurons in sleep

regulation, we employed a Split-GAL4 strategy (11) to access a reduced number of

23E10-GAL4 expressing cells. In designing this screen, our primary region of interest in

the brain was the dFB, a choice dictated by the number of studies supporting a role for

this structure in sleep regulation (27-30, 33, 34, 37, 44, 45). Central to the Split-GAL4

technology is the fact that the functional GAL4 transcription factor can be separated into

two non-functional fragments, a GAL4 DNA-binding domain (DBD) and an activation

domain (AD). Different enhancers are used to express these two fragments and the

functional GAL4 transcription factor is reconstituted only in the cells in which both

enhancers are active (11).

We obtained 22 different AD lines based on their associated GAL4 line expressing in the dFB, as observed using the Janelia Flylight website (36, 46). Since the goal of our screen was to identify the contribution of individual 23E10 dFB neurons in sleep regulation, we designed a targeted approach by combining these individual AD lines to a 23E10-DBD line (46, 47), thus creating 22 new Split-GAL4 lines named FBS for Fan-Shaped Body Splits lines (See Table2 for a description of these lines). These newly created lines were screened behaviorally for their ability to modulate sleep and anatomically to identify their expression patterns. To do so, each individual Split-GAL4 FBS line was crossed to a line expressing both the thermogenetic TrpA1 cation channel and a mCD8GFP construct. Flies were maintained at 22°C for 2 days before raising the temperature to 31°C at the beginning of day 3 for a duration of 24h. Temperature was brought down to 22°C on day 4 (Figure 2A). For each individual fly, we calculated the percentage of total sleep change between activation day (day3) and baseline day (day2) (Figure 2A). As a control for this screen, we used an enhancerless AD construct (created in the vector used to make all the AD lines) combined with 23E10-DBD, since it is the common element in all the Split-GAL4 lines screened in this experiment. As seen in Figure 2B,D, control female flies show a reduction in sleep upon transferring to 31°C, an effect particularly evident during the night. These data are in agreement with previous studies documenting that changes in temperature modulate sleep architecture and that high temperature disturbs sleep at night (42, 43, 48, 49) . However, acute activation of the neurons contained in 9 of the 22 FBS lines led to significant increases in percentage sleep change in female flies (FBS4, FBS5, FBS42, FBS45, FBS53, FBS68, FBS72, FBS81 and FBS84), when compared with controls. Individual sleep

traces for all 22 FBS lines are shown in Figure 2E,F,G,H and Figure 2-figure

supplement 1. Importantly, analysis of activity counts during awake time reveals that

increases in sleep are not caused by a reduction in locomotor activity (Figure 2C). On

the contrary, 3 of the strongly sleep-inducing lines even show an increase in waking

activity upon neuronal activation (Figure 2C). This increase is likely explained by the

need for a fly to perform tasks that are mutually exclusive to sleep in a reduced amount

of waking time. While increases in total sleep are indicative of increase sleep quantity,

this measurement does not provide information about sleep quality or sleep depth. To

assess whether sleep quality is modulated by FBS lines, we analyzed sleep

consolidation during the day and night in these flies. As seen in Figure 2-figure

supplement 2A, upon neuronal activation, daytime sleep bout duration is significantly

increased in 8 out of the 9 sleep-promoting lines (FBS4, FBS5, FBS42, FBS45, FBS53,

FBS68, FBS72 and FBS84), while it is unaffected in FBS81. Notably, increased sleep

bout duration is believed to be an indication of increased sleep depth (50). Thus, these

data indicate that activating neurons contained in these 8 FBS lines not only increases

sleep level but probably also increases sleep depth. Interestingly, one of the lines that

did not show an increase in total sleep, has a significant increase in daytime sleep bout

duration (FBS28). During the nighttime, the effect of high temperature on sleep is

obvious as most lines, including the control, display a significant decrease in sleep bout

duration when raising the temperature to 31°C (Figure 2-figure supplement 2B). Only 6

of the 9 sleep-promoting lines (FBS4, FBS5, FBS42, FBS45, FBS53 and FBS68)

maintain similar nighttime sleep bout durations before and during neuronal activation.

Importantly, in these 6 lines nighttime sleep bout duration is significantly increased at

31°C when compared to control flies (Figure 2-figure supplement 2B). These data

indicate that most sleep-promoting Split-GAL4 lines increase daytime sleep

consolidation and suggest that they could also increase nighttime sleep consolidation.

However, because of the clear effect of high temperature on nighttime sleep, this claim

requires further validation.

In addition, since sleep in *Drosophila* is sexually dimorphic (51-53), we systematically

assessed male flies in our experiments. As seen in Figure 2-figure supplement 3, we

obtained almost identical behavioral data when analyzing male flies using our

thermogenetic activation approach. Thus, we conclude that we have identified 9 FBS

lines that promote sleep when thermogenetically activated in males and females.

Having identified 9 different Split-GAL4 lines that significantly increased sleep when

activated, we sought to identify the neurons that are contained within these lines.

Because the goal of our screen was to identify sleep-regulating dFB neurons in the

23E10-GAL4 pattern of expression, we expected that most of our FBS lines, sleep-

promoting or not, would express in dFB cells. This is indeed the case as 20 out of the

22 FBS lines express in some dFB neurons (Figure 2, Figure 2-figure supplement 1,

Table 2). The two exceptions to this are FBS25 (expresses in no neurons at all) and

FBS5 as discussed below. Regarding sleep-promoting FBS lines, 8 out of 9 have some

expression in the dFB. However, and surprisingly, one of the strongly sleep-promoting

lines, FBS5, does not express in any dFB neurons at all (Figure 2F, Video 3, Video 4).

Rather, FBS5 strongly expresses in two clusters of 4-5 neurons located in the anterior

ventrolateral protocerebrum and in 4 VNC cells located in the metathoracic ganglion. A

close examination of the anatomy of these 4 VNC neurons indicate that they have

processes in the brain. Two of these neurons show a very specific pattern of expression

with brain processes located in the superior medial protocerebrum. Because of their

characteristic look, we named these processes "bowtie". These processes are

highlighted in the right panel in Figure 2F (see also Video 3). FBS5 additionally labels

two neurons in the VNC whose cell bodies are located very close to the "bowtie"

neurons. These neurons have characteristic processes in each leg ganglion (yellow

arrows in Figure 2F-VNC image). Based on their anatomy, we hypothesize that these

cells could be Taste Projection Neurons 1 (TPN1) that receive inputs from sweet

gustatory receptor leg neurons (GRNs) and convey this information to the brain in the

subesophageal zone (SEZ) (54). We observed these putative TPN1 neurons in 14 out

of the 22 FBS lines. Importantly, since all the neurons in FBS5 are part of the 23E10-

GAL4 expression pattern, these data indicate that this driver contains non-dFB sleep-

promoting cells.

Most other sleep-promoting line expression patterns are very discrete with only two

major areas of expression; dFB neurons, and the metathoracic ganglion region

containing the two "bowtie" neurons and TPN1-like cells (FBS42, FBS45, FBS53,

FBS68, FBS72, FBS81 and FBS84) (Figure 2G,H; Figure 2-figure supplement 1;

Table2). Some of these FBS lines show minimal amount of expression elsewhere in the

brain, but there is no consistency in these additionally labeled cells between different

lines. Of note, FBS4 expresses in 18-23 dFB neurons (Figure 2E, Table 2) and no other

cells in the brain or VNC. Considering that 23E10-GAL4 reliably labels 23-30 dFB

neurons in our confocal microscopy experiments (Table1), we conclude that FBS4

expresses in 75-80% of 23E10 dFB neurons. Because of its dFB specific expression

pattern and its ability to increase sleep when thermogenetically activated, we suspect

that FBS4 probably contains neuron(s) that are regulating sleep. However, we also note

that the sleep increase triggered by FBS4 thermogenetic activation is not as striking as

when activating other strongly sleep-promoting lines like FBS5, FBS53 or FBS68

(Figure 2B for females and Figure 2-figure supplement 3A for males).

One particularly interesting line is FBS45. FBS45 is strongly sleep-promoting and is

expressed in 21-32 dFB neurons, indicating that most if not all 23E10 dFB neurons are

present in FBS45 (Figure 2G, Table2). However, FBS 45 also expresses in the two

"bowtie" neurons. Importantly, FBS45 does not label any other neurons in the brain and

VNC.

When combining behavioral and anatomical data, it appears that the numbers of dFB

neurons contained within a specific FBS line cannot be predictive of whether the line

can promote sleep upon neuronal activation. For example, FBS68 strongly promotes

sleep and expresses in only 3-6 dFB neurons (Figure 2H, Table 2), while FBS60

contains 22-27 dFB neurons and does not promote sleep (Figure 2-figure supplement

1L, Table 2).

Remarkably, a careful examination of expression patterns reveals that many FBS lines

appear to express in the "bowtie" VNC neurons. Note that the brain processes of these

cells are located extremely close to the axonal projections of dFB neurons, making the

visualization of these VNC projections sometimes difficult, depending on the strength of

dFB projection's staining. Nevertheless, these VNC "bowtie" neurons are present in 17

out of 22 FBS lines (See Figure 2 and Figure 2-figure supplement 1 VNC "bowtie"

processes). Importantly, we can see these neurons and their processes in all sleep-

promoting lines with the notable exception of FBS4 (Figure 2, Figure 2-figure

supplement 1, Table 2), as discussed above. Upon examination of the literature, we

found that neurons with similar projections are frequently seen in many Janelia GAL4

lines (55). This study reports that these neurons (referred to as "sparse T" in this work)

and their processes are seen in up to 60% of all Janelia GAL4 lines. The fact that these

"bowtie" neurons are frequently present in many GAL4 lines probably explains why we

see them in the vast majority of our FBS lines (18 out of 22 or 82% of them).

In conclusion, our thermogenetic screen and anatomical analysis identified 9 sleep-

promoting lines with diverse patterns of expression. While the identity of the neurons

responsible for sleep promotion in these different lines is not clear at this stage, our

analysis indicates that there are multiple groups of sleep-promoting neurons in the

23E10-GAL4 pattern of expression; one of which appears to be part of the dFB (FBS4)

while the other is non-dFB based (FBS5).

Optogenetic confirmation of sleep promoting FBS lines

Because of the strong effect that temperature has on sleep, we sought to confirm our

thermogenetic findings using an alternative way to manipulate FBS lines. The logic

behind our thinking is that it may be difficult to fully describe and characterize the sleep

behaviors of our FBS lines when temperature itself has such a profound effect on sleep.

Furthermore, a recent study demonstrated that some of the effects of high temperature

on sleep are mediated by GABAergic transmission on dFB neurons (49). Taken

together, these data suggest that thermogenetic manipulations may not be optimal to

conduct our screen. We thus undertook an optogenetic screen using CsChrimson (56).

Our optogenetic experimental setup is described in Figure 3A. Notably, for neurons to

be activated when expressing CsChrimson, flies need to be fed all trans-retinal and the

neurons must be stimulated with a 627nm LED light source (56). Thus, optogenetic

approaches provide a much better control on the timing and parameters of activation.

Nonetheless, optogenetic activation gave results that are mostly identical to the

thermogenetic approach. That is, most sleep-promoting lines identified using TrpA1 also

induce sleep when activated with CsChrimson in retinal-fed flies (FBS5, FBS42, FBS45,

FBS53, FBS68 and FBS72) (Figure 3B for female data). Notable differences are FBS4,

FBS81 and FBS84 which do not promote sleep when optogenetically activated in

female flies. However, activating FBS81 and FBS84 significantly increases sleep in

retinal-fed male flies (Figure 3-figure supplement 2A). Importantly, unlike FBS81 and

FBS84, activation of FBS4 neurons does not increase sleep in retinal-fed males (Figure

3-figure supplement 2A). Thus, there is a clear discrepancy for FBS4 comparing

thermogenetic and optogenetic activations. Another difference between the two

activation approaches is FBS70 that is sleep-promoting using CsChrimson in retinal-fed

female flies while it was not using TrpA1 (Figure 3B). Surprisingly, retinal-fed

FBS70>UAS-CsChrimson males do not increase sleep upon LED stimulation,

potentially revealing a sexually dimorphic phenomenon (Figure 3-figure supplement 2A).

Importantly, assessment of locomotor activity during waking period reveals that

optogenetic activation does not decrease waking activity in long sleeping retinal-fed

females (Figure 3C) and males (Figure 3-figure supplement 2B), ruling out the

possibility that locomotor defects are the underlying cause of the sleep phenotypes

observed. Examination of sleep bout duration in retinal-fed flies indicates that most

sleep-promoting FBS lines increase sleep consolidation during the day (Figure 3D for

females and Figure 3-figure supplement 2C for males) and during the night (Figure 3E

for females and Figure 3-figure supplement 2D for males). These data indicate that

optogenetic activation increases sleep quantity and probably sleep quality (sleep depth)

in many of the FBS lines first identified using our thermogenetic approaches.

Interestingly, some FBS lines that did not increase total sleep upon optogenetic

activation increased daytime sleep bout duration: FBS57 and FBS60 females (Figure

3D) and FBS33, FBS35, FBS64 and FBS87 males (Figure 3-figure supplement 2C).

These data suggest that these lines may also express in sleep-regulating neurons.

Importantly, analysis of sleep parameters in vehicle-fed flies demonstrate that the sleep

phenotypes we observed are specific to retinal-fed and LED stimulated flies (Figure 3-

figure supplement 1 for females and Figure 3-figure supplement 3 for males).

In conclusion, our thermogenetic and optogenetic neuronal activation revealed several

strongly sleep-promoting lines. When looking at their expression patterns, our data

clearly demonstrate that 23E10 GAL4 contains at least two different sleep-promoting

regions, one that is part of the dFB and one that is not.

The 23E10-GAL4 driver contains sleep-promoting neurons that are not located in

the brain

To reveal the identity of the non-dFB 23E10 sleep-promoting neurons, we undertook a

second screen creating additional Split-GAL4 lines focusing on neurons that are present

in FBS5. To facilitate this task, we used Color Depth MIP mask search, a software plug-

in for Fiji (57). This approach identified several AD lines (containing neurons of interest)

that we combined with our 23E10-DBD stock. An example of such a line is shown in

Figure 4A-D. This Split-GAL4 line (VT020742-AD;23E10-DBD) expresses only in 4 VNC

neurons (Figure 4C). Projections of 2 of these are clearly "bowtie" like (gray arrows in

Figure 4A-B). Projections of the other two appear to be identical to TPN1 neurons that

have been previously described (yellow arrows in Figure 4A-D). Remarkably,

VT020742-AD;23E10-DBD does not express in any other neurons in the brain and

VNC. To confirm the presence of TPN1 neurons in this Split-GAL4 line we expressed

GFP using a TPN1 specific driver, 30A08-LexA (54) while simultaneously expressing

RFP using VT020742-AD;23E10-DBD. As seen in Figure 4-figure supplement 4A, there

is clear GFP and RFP co-expression in two out of the four VT020742-AD;23E10-DBD

neurons (yellow arrows). In addition, we also see co-expression of GFP and RFP in the

processes that go into each leg ganglion, but only RFP expression in the "bowtie"

processes. From these data we conclude that VT020742-AD;23E10-DBD labels 4

neurons in the metathoracic ganglion of the VNC: 2 "bowtie" neurons and 2 TPN1 cells.

Optogenetic activation of these 4 neurons strongly increases sleep in retinal-fed

females (Figure 4K) and males (Figure 4-figure supplement 2A). In addition, daytime

and nighttime sleep bout duration are significantly increased in retinal-fed VT020742-

AD;23E10-DBD>UAS-CsChrimson flies (Figure 4M-N for females and Figure 4-figure

supplement 2C-D for males). Importantly, analysis of waking activity reveals that the

sleep increases are not caused by a deficit in locomotor activity (Figure 4L for females

and Figure 4-figure supplement 2B for males).

We then wanted to find out whether TPN1 neurons are the cells involved in sleep

promotion in VT020742-AD;23E10-DBD. We thus created a 30A08-AD;23E10-DBD

Split-GAL4 line. This line labels only the two TPN1 neurons and no other cells in the

brain and VNC (Figure 4E-F). Importantly, optogenetic activation of 30A08-AD;23E10-

DBD neurons has no effect on sleep (Figure 4K-N and Figure 4-figure supplement 2A-

D), indicating that the sleep-promoting capacity of VT020742-AD;23E10-DBD most

likely lies within the two "bowtie" neurons. To further rule out an involvement of TPN1

neurons in sleep promotion, we adopted the Split-GAL4 repressor Killer Zipper (KZip+)

technology which inhibits an AD and DBD from forming a functional GAL4 (58).

Expressing KZip+ with 30A08-LexA while expressing GFP with VT020742-AD;23E10-

DBD, reduces the number of GFP positive cells from 4 to 2 (Figure 4-figure supplement

4B). Only the two "bowtie" neurons can be seen in this combination. Thus, we can

effectively subtract the TPN1 neurons from the VT020742-AD;23E10-DBD pattern of

expression. Optogenetic activation of the remaining "bowtie" neurons (in retinal-fed

VT020742-AD;23E10-DBD>UAS-CsChrimson, 30A08-LexA>LexAop2-KZip+ flies) still

promotes sleep (Figure 4-figure supplement 4C). Taken together these data indicate

that the two "bowtie" neurons in VT020742-AD;23E10-DBD (and 23E10-GAL4) increase

sleep when activated and that TPN1 neurons are not involved in sleep promotion.

To further refine our analysis, we constructed an additional Split-GAL4 line (VT013602-

AD;23E10-DBD) that only expresses in the two "bowtie" like VNC neurons (gray arrows

in Figure 4G-J, Video 5, Video 6). Optogenetic activation of retinal-fed VT013602-

AD;23E10-DBD neurons strongly promotes sleep (Figure 4K and Figure 4-figure

supplement 2A for male data) and increases daytime (Figure 4M and Figure 4-figure

supplement 2C) and nighttime bout duration (Figure 4N and Figure 4-figure supplement

2D). Again, increases in sleep are not caused by abnormal locomotor activity (Figure 4L

and Figure 4-figure supplement 2B). Importantly, analysis of sleep parameters in

vehicle-fed flies reveal sleep phenotypes observed are specific to activation of

VT013602-AD; 23E10-DBD neurons (Figure 4-figure supplement 1 and Figure 4-figure

supplement 3). Since activating the two 'bowtie" neurons increases daytime and

nighttime sleep bout duration, we hypothesize that this manipulation increases sleep

quality, namely sleep depth. To investigate this possibility, we measured arousal

thresholds in VT013602-AD;23E10-DBD>UAS-CsChrimson flies. As seen in Figure 4-

figure supplement 5, the proportion of retinal-fed flies awakened by mild mechanical

perturbations (1 and 2) is significantly reduced in LED stimulated (LED ON, activated)

individuals compared with non-activated flies (LED OFF). Thus, these data demonstrate

that when activated, the two "bowtie" neurons increase sleep quantity and sleep depth

as indicated by an increase in arousal threshold. Importantly, at the highest perturbation

level (4), retinal-fed LED activated VT013602-AD;23E10-DBD>UAS-CsChrimson flies

respond to the stimulus similarly to controls, indicating that the sleep that is induced by

activation of these neurons is reversible.

To assess whether the "bowtie" neurons seen in VT020742-AD;23E10-DBD and

VT013602-AD;23E10-DBD are the same, we constructed a line that carries both

VT020742-AD and VT013602-AD as well as 23E10-DBD. Expressing GFP with this line

labels only 4 VNC neurons, 2 "bowtie" cells and 2 TPN1 (Figure 4-figure supplement

4D). If the "bowtie" neurons were different in VT020742-AD;23E10-DBD and VT013602-

AD;23E10-DBD we would expect to see 6 cells. Since this is not the case, we can

conclude that the 2 'bowtie" neurons observed in VT020742-AD;23E10-DBD and

VT013602-AD;23E10-DBD are the same. To provide functional support to this finding,

we created an additional Split-GAL4 line, VT013602-AD;VT020742-DBD. This Split-

GAL4 line labels 2 "bowtie" cells in the VNC with typical processes in the brain (gray

arrows in Figure 4-figure supplement 4E) and promotes sleep when activated

thermogenetically (Figure 4-figure supplement 4F-H). Taken together, these data

demonstrate that we have unequivocally identified two novel sleep-promoting neurons

located in the VNC. Importantly, these cells are part of the 23E10-GAL4 expression

pattern. We will from now on refer to these neurons as VNC-SP (for VNC Sleep-

Promoting).

Silencing VNC-SP neurons reduces sleep

Having demonstrated that activation of VNC-SP neurons increases sleep duration and sleep depth, we wondered whether these cells could modulate sleep bi-directionally. Thus we performed experiments to silence the activity of VNC-SP neurons either chronically, by expressing the inward rectifying potassium channel Kir2.1 (59) or acutely by using the temperature sensitive Shi[ts1] construct (60). As seen in Figure 5A-B, total sleep is significantly reduced when expressing Kir2.1 in VNC-SP neurons in female flies. This reduction in sleep is accompanied by a reduction of both daytime (Figure 5C) and nighttime sleep bout duration (Figure 5D), suggesting a defect in sleep consolidation. Importantly, analysis of activity counts during waking time reveals that the decrease in sleep is not due to hyperactivity (Figure 5E). We obtained identical behavioral results when expressing Kir2.1 in VNC-SP neurons of male flies (Figure 5F-J). Thus, chronic silencing of VNC-SP neurons decreases sleep and disrupts sleep consolidation.

To confirm the findings obtained using chronic silencing we decided to employ an acute silencing protocol by expressing Shi[ts1] (60) in VNC-SP neurons. Notably, this approach relies on increasing temperature to silence neurons. Because of the previously reported effect of temperature on sleep, it is not surprising to see that the shift in temperature reduces sleep in control flies (Figure 5K for females and Figure 5N for males). Acute silencing of VNC-SP neurons also reduces sleep, an effect particularly striking at night (Figure 5L and Figure 5M). Importantly, nighttime sleep is significantly more reduced in VNC-SP> Shi[ts1] flies than in controls (Figure 5M and Figure 5P).

Taken together, our chronic and acute silencing data demonstrate that VNC-SP

neurons positively regulate sleep. Importantly, these results further reinforce our

findings using optogenetic and thermogenetic activation experiments.

VNC-SP neurons express and use acetylcholine to modulate sleep

Having demonstrated that VNC-SP neurons modulate sleep bi-directionally, we sought

to identify the neurotransmitter(s) used by these cells. A previous study suggested that

most neurons in the VNC use one of three neurotransmitters: GABA, Glutamate or

Acetylcholine (61). With this in mind, we expressed GFP in VNC-SP neurons and

incubated them with antibodies to GABA, choline acetyltransferase (ChAT), the enzyme

necessary to produce Acetylcholine and the vesicular glutamate transporter (VGlut). As

seen in Figure 6A, we observed clear co-labelling of GFP and ChAT in VNC-SP

neurons. To further confirm that VNC-SP neurons express ChAT, we employed a

combinatorial approach expressing RFP in VNC-SP cells and GFP driven by a ChAT-

LexA (62) construct that is inserted within the endogenous ChAT locus. This construct

ensures that the expression of LexA is controlled by ChAT regulatory sequences and

should reflect transcriptional activity at the endogenous ChAT locus. As seen in Figure

6-figure supplement 1A, we observe clear co-labelling of GFP and RFP in VNC-SP

neurons (gray arrow). In addition, we employed a similar approach using 23E10-GAL to

drive expression of RFP. As seen in Figure 6-figure supplement 1B, we observe clear

co-labelling of GFP and RFP in two neurons in the metathoracic ganglion of the VNC

(gray arrows). The evidence clearly suggests that these two cells are the VNC-SP

neurons. Interestingly, in the process of this experiment, we found that 2-3 dFB neurons

per brain hemisphere are co-labelled by 23E10-GAL4 and ChAT-LexA construct (Figure

6-figure supplement 1B, bottom panels, red arrows). Taken together, these data

indicate that VNC-SP neurons and some 23E10-GAL4 dFB neurons are cholinergic.

Importantly, we did not detect any co-labelling between GFP and VGlut (Figure 6B) or

GABA (Figure 6C) in these experiments. To find out whether VNC-SP neurons express

any additional neurotransmitters we also performed experiments using antibodies to

serotonin (5HT, Figure 6D), tyrosine hydroxylase (TH, Figure 6E), the key enzyme in

the synthesis of dopamine, and tyrosine decarboxylase 2 (Tdc2, Figure 6F), an enzyme

involved in the synthesis of octopamine. We did not observe any co-labelling with any of

these 3 antibodies.

These data indicate that VNC-SP neurons are cholinergic but they do not provide any

information as to whether cholinergic transmission is needed for the sleep modulating

role of VNC-SP neurons. To address this question, we sought to disrupt cholinergic

transmission in VNC-SP neurons using an RNAi approach. Thus, we obtained two

independent RNAi lines for ChAT. We first sought to validate them by crossing them to

the pan-neuronal elav-GAL4 driver. One of the lines (60028) led to lethality when

crossed to elav-GAL4, indicative of a significative reduction in ChAT levels. The second

line (25856) gave rise to progeny suggesting it is not very efficient. Note that a similar

approach was used to validate ChAT RNAi constructs in other studies (63). However, to

further confirm the efficiency of these ChAT RNAi constructs we performed a qPCR

analysis on flies expressing ChAT RNAi (60028) specifically in the eyes using a GMR-

GAL4 driver. Figure 6-figure supplement 1C demonstrates that ChAT mRNA levels are

significantly reduced (about 20%) in the eyes of GMR-GAL4>ChAT RNAi (60028) flies

compared with controls. Notably, ChAT mRNA levels are unaffected in the body of

these flies, illustrating the efficiency and specificity of the ChAT RNAi line. Importantly,

qPCR analysis of the second ChAT RNAi line (25856) showed no significant reduction

in ChAT levels, further demonstrating its inefficiency (data not shown). Based on this

data, we decided to only employ one ChAT RNAi (60028) in our behavioral studies.

As seen in Figure 6-figure supplement 1D-E, ChAT levels as measured by

immunocytochemistry are significantly reduced when expressing ChAT RNAi in VNC-

SP neurons. Behaviorally, reducing ChAT levels in VNC-SP neurons significantly

reduces sleep (Figure 6G-H) and, daytime (Figure 6I) and nighttime (Figure 6J) sleep

bout duration in female flies. Importantly, the reduction in sleep is not due to

hyperactivity (Figure 6K). We obtained similar behavioral data when looking at males

(Figure 6-figure supplement 1F-I). Taken together, our data indicate that cholinergic

transmission is involved in regulating the sleep modulating effect of VNC-SP neurons.

We also note that the behavioral effects of reducing ChAT levels and silencing VNC-SP

neurons are consistent with each other, suggesting that in the absence of ChAT, the

ability of VNC-SP neurons to positively regulate sleep is seriously diminished. To further

interrogate the necessity of cholinergic transmission for VNC-SP neurons to promote

sleep, we activated VNC-SP neurons optogenetically, while simultaneously expressing

ChAT RNAi in these cells. As seen in Figure 6L, expression of CsChrimson and a RNAi

line targeting GFP (control RNAi) does not prevent sleep increase in retinal-fed flies.

However, expressing ChAT RNAi completely abolishes the sleep promotion triggered by

activation of VNC-SP neurons. In conclusion, our data clearly demonstrate that VNC-SP

neurons are cholinergic and that disruption of cholinergic transmission abolishes the

sleep-promoting role of these cells.

To further confirm that the sleep phenotypes we observe are due to VNC-SP neurons,

we optogenetically activated VNC-SP cells while expressing the Split-GAL4 repressor

KZip+ (58) under the control of a ChAT-LexA driver. As seen in Figure 6-figure

supplement 1J, expressing the KZip+ repressor in ChAT expressing neurons blocks the

sleep induction caused by activation of VNC-SP neurons. These data strongly suggest

that the KZip+ is expressed in VNC-SP neurons and actively represses the binding of

the GAL4-AD and GAL4-DBD constructs. To visually confirm repression of GAL4

activity, we expressed GFP in VNC-SP neurons and the KZip+ repressor in ChAT-LexA

cells. As seen in Figure 6-figure supplement 1K, expressing the KZip+ repressor in

ChAT expressing neurons strongly reduces the amount of GFP staining seen in VNC-

SP neurons. Taken together, these data confirm that the sleep phenotypes we observe

are due to activation of VNC-SP neurons and further demonstrate that these cells are

cholinergic.

Since VNC-SP neurons are observed in 18 of our 22 FBS lines, and in most sleep-

promoting lines (except for FBS4), we wondered whether VNC-SP neurons are the cells

responsible for the sleep increase in all our lines. We decided to focus on FBS45, since

it expresses in most if not all 23E10-GAL4 dFB neurons (Table 2) and in the two VNC-

SP neurons. Importantly, FBS45 is one of the strongest sleep-promoting lines we have

studied in this work. Our question was the following: Is the sleep increase caused by

activation of FBS45 due to VNC-SP neurons, dFB neurons, or both group of cells? To

answer this question, we expressed UAS-CsChrimson in FBS45 neurons while

simultaneously driving the expression of the KZip+ repressor in ChAT-LexA cells. We

hypothesize that this manipulation should repress GAL4 formation specifically in VNC-

SP neurons. However, since we have reported that ChAT-LexA also expresses in some

23E10-GAL4 dFB neurons (Figure 6-figure supplement 1B), our manipulation may

repress GAL4 in a few dFB neurons. As seen in Figure 6-figure supplement 1L,

expressing the KZip+ repressor in ChAT-LexA prevents retinal-fed FBS45>UAS-

CsChrimson flies from increasing sleep when activated. These data indicate that the

neurons responsible for the sleep increase when activating FBS45 cells are cholinergic

and that expressing CsChrimson only in FBS45 neurons that are not targeted by ChAT-

LexA (non-cholinergic) cannot promote sleep. We then visually assessed the extent of

ChAT-LexA driven KZip+ repression in FBS45 neurons by expressing UAS-GFP. As

seen in Figure 6-figure supplement 1M, expressing the KZip+ repressor in ChAT-LexA

completely abolishes GFP expression in VNC-SP neurons in *FBS45>UAS-GFP; ChAT-*

LexA>LexAop2-KZip+ flies. However, as we expected based on our previous data,

ChAT-LexA>LexAop2-KZip+ consistently removes GFP expression in 4-6 dFB neurons

as well. Taken together, these data suggest that the sleep-promoting capacity of FBS45

relies on cholinergic neurons. While we strongly believe that the cells responsible are

VNC-SP neurons, we cannot completely exclude a role for one of the ChAT expressing

23E10-GAL4 dFB neurons. Further work will help clarify the role of cholinergic dFB

neurons in sleep regulation.

Since the VNC receives and integrates sensory inputs from the periphery and sends

this information to the brain (64-66), we wondered whether VNC-SP neurons could

similarly send sleep-relevant signals to the brain. The location of VNC-SP neurons in

the VNC, and their processes in the brain suggests that this is a likely possibility. To find

out, we expressed the dendritic marker DenMark (67) in conjunction with the

presynaptic marker synaptotagmin-GFP (syt.eGFP) (68) in VNC-SP neurons. Our data

indicate that VNC-SP neurons have mostly postsynaptic sites in the VNC, as expected if

these cells receive sensory information from the periphery (Figure 6-figure supplement

2, middle panels). However, and surprisingly, the brain processes of these cells are

labeled with both the presynaptic and postsynaptic markers (Figure 6-figure supplement

2, top panels). This suggests that VNC-SP neurons are well positioned to receive

sensory inputs from the periphery in the VNC and that they are sending this information

to the brain. However, since the brain processes of VNC-SP neurons are also harboring

postsynaptic sites, it is highly likely that they also get inputs in the brain. Nevertheless,

our data suggest that VNC-SP neurons send sleep-relevant signals to the brain.

Multiple sleep-promoting regions in 23E10-GAL4 with functional specificity

Our data unequivocally demonstrate that VNC-SP neurons, which are part of the

23E10-GAL4 pattern of expression, are sleep-promoting. Since 23E10-GAL4 is widely

seen as dFB specific driver when it comes to sleep regulation, the identification of VNC-

SP neurons is a major finding that impacts our current views about the role of the dFB in

sleep regulation. Our data shows that all but one FBS sleep-promoting lines contain

VNC-SP neurons. The only sleep-promoting line that is dFB specific (FBS4) only mildly

promotes sleep (when compared with other lines) in our thermogenetic screen but does

not using optogenetic. These data raise important questions about the role of the dFB in

sleep regulation and sleep-promotion.

To try to address these questions we went back to the 23E10-GAL4 driver and its

23E10-LexA counterpart (69). First, we wanted to check whether 23E10-GAL4 and

23E10-LexA express in the same number of neurons. This is an important control since

it was reported that often for a given enhancer sequence, the LexA expression pattern

does not fully recapitulate the GAL4 expression pattern (69). We created a line

expressing GFP under the control of 23E10-LexA and RFP under the control of 23E10-

GAL4. As seen in Figure 7A-B, the 23E10-LexA driver does not fully recapitulate the

23E10-GAL4 pattern of expression in the dFB. Some cells are clearly co-labelled by

GFP and RFP (yellow arrows) but some other dFB neurons only express RFP (gray

arrow). Importantly, we did not observe any dFB cells (or any other cell in the brain) that

are only labeled by 23E10-LexA. These data indicate that, like many other LexA drivers,

23E10-LexA only expresses in a subset of the 23E10-GAL4 pattern of expression. We

then turned our attention to the metathoracic ganglion of the VNC. As seen in Figure

7C, 23E10-LexA labels the two TPN1 neurons (yellow arrows) while 23E10-GAL4 labels

the two TPN1 cells as well as the two VNC-SP neurons (gray arrows). The processes of

TPN1 neurons are highlighted in Figure 7E and those from VNC-SP in Figure 7D. We

see clear co-labelling of TPN1 processes by 23E10-LexA and 23E10-GAL4, but VNC-

SP processes are only labeled by 23E10-GAL4. Taken together, we conclude that

23E10-LexA only expresses in a subset of 23E10-GAL4 neurons and that 23E10-LexA

does not express in VNC-SP neurons.

We then wondered whether 23E10-LexA could promote sleep when activated

optogenetically. As seen in Figure 7F, optogenetic activation of retinal-fed 23E10-

GAL4>UAS-CsChrimson strongly increases sleep as expected. Controls show no

changes in sleeping levels. However, optogenetic activation of retinal-fed 23E10-

LexA>LexAop2-CsChrimson has no effect on sleep (Figure 7G). The stark difference in

behavior seen when activating 23E10-GAL4 and 23E10-LexA could be explained by the

fact that 23E10-GAL4 expresses in more neurons than 23E10-LexA. In this scenario the

absence of VNC-SP neurons in the 23E10-LexA pattern of expression must account for

some if not all this difference. However, since 23E10-LexA does not fully recapitulate

the expression pattern of 23E10-GAL4 in the dFB, we cannot rule out that some of

these missing dFB neurons may be involved in sleep promotion. An alternative

explanation is that 23E10-LexA may not be as strong a driver as 23E10-GAL4 and

therefore would not express sufficient levels of the CsChrimson construct. In our

experience this is very unlikely as we have optogenetically activated other pairs of

Gal4/LexA drivers and obtained identical and strong sleep activation with both the LexA

and GAL4 counterparts (data not shown).

Does our data mean that the dFB plays no role at all in sleep regulation? Based on the

number of studies supporting a role for this structure in sleep regulation (27-30, 33, 34,

37, 44, 45) we think that this is very unlikely. Most of the previous work supports a role

for dFB neurons in sleep homeostasis. In particular, sleep deprivation increases the

excitability of dFB neurons (29) and augmented sleep pressure switches dFB neurons

from an electrically silent to an electrically active state (33). Furthermore, reducing the

excitability of dFB neurons by reducing levels of the Rho-GTPase-activating protein

encoded by the crossveinless-c (cv-c) gene leads to a defect in sleep homeostasis (29).

With this in mind, we wanted to assess whether silencing all 23E10-Gal4 neurons, or

just 23E10-GAL4 dFB and VNC-SP neurons separately would affect sleep homeostasis

(Figure 8A). As seen in Figure 8B, chronic silencing of all 23E10-GAL4 neurons by

expression of Kir2.1 effectively blocks homeostatic sleep rebound when compared with

both parental controls. Interestingly, chronic silencing of VNC-SP cells does not disrupt

homeostatic sleep rebound (Figure 8C) while silencing of most 23E10-GAL4 dFB

neurons (contained in the FBS4 line) does (Figure 8D). These data support a role for

23E10-GAL4 dFB neurons in sleep homeostasis, in agreement with previous studies.

Taken together, our data indicate that the 23E10-GAL4 driver contains at least two

groups of sleep-regulating neurons, one located in the dFB that is mostly involved in

sleep homeostasis and VNC-SP neurons that are dispensable for sleep homeostasis.

Discussion

The strength of the fly model lies in its strong genetic techniques, allowing researchers to selectively manipulate discrete populations of neurons and monitor how specific behaviors are affected. To do so, an impressive collection of binary expression systems have been developed. In particular, the GAL4/UAS system (8) has been the keystone of *Drosophila* neurobiological studies. However, GAL4 drivers are often not specific enough and express in cells outside the region of interest. This can make the task of unequivocally assigning a given behavior to a specific neuron or group of neurons particularly difficult. Importantly, refinement of GAL4 expression pattern can be achieved by employing the intersectional Split-GAL4 technology (11).

Like mammalian sleep, sleep in *Drosophila* is regulated by multiple areas in the brain (16, 17). Particularly, an expanding amount of evidence supports a role for dFB neurons in sleep regulation (27-30, 33, 34, 37, 44, 45). The most widely used tool to manipulate dFB neurons is the 23E10-GAL4 driver (33, 36), and thermogenetic or optogenetic activation of 23E10-GAL4 neurons results in increased sleep (31, 37-39). However, whether 23E10-GAL4 dFB neurons can be considered as a functionally homogeneous group is still unclear. Also unclear is whether the sleep-promotion caused by activation of 23E10-GAL4 neurons is due to dFB neurons. This concern is reinforced by the fact that 23E10-GAL4 expresses in many neurons outside the dFB in the brain and in the VNC (Figure 1, Table 1). In this study, we sought to identify the role of individual 23E10-GAL4 dFB neurons in sleep regulation.

To identify sleep-promoting neurons within the 23E10-GAL4 pattern of expression we

adopted a targeted Split-GAL4 strategy combining 22 individual AD lines (selected for

the strong dFB expression pattern of their GAL4 counterparts) with a 23E10-DBD line.

We screened these 22 novel Split-GAL4 FBS lines behaviorally and anatomically. First,

we employed a thermogenetic approach to activate the neurons contained in these 22

lines. These experiments identified 9 FBS lines that promote sleep when activated in

females and males (Figure 2). An anatomical assessment of each line revealed that

most sleep-promoting FBS lines express in some dFB neurons, as expected based on

our targeted approach. However, they also express in other areas in the brain and

especially in the metathoracic ganglion of the VNC. In fact, we identified two very clear

and characteristic processes associated with 4 different VNC neurons in many FBS

lines, sleep-promoting or not. The first set of two VNC neurons have typical processes

in each leg ganglion and send processes in the brain that terminate in the SEZ. Our

data demonstrated that these cells are TPN1 neurons (Figure 4) (54). We observed

these neurons in 14 out of the 22 FBS lines. However, we demonstrated that TPN1

neurons are not promoting sleep when activated (Figure 4). The second set of two VNC

neurons send processes to the superior medial protocerebrum. Due to their

characteristic anatomical features, we named these processes "bowtie". Importantly,

these processes are seen in 18 out of 22 FBS lines (and in 8 out of the 9 sleep-

promoting lines). The high frequency of observation of these processes in our FBS lines

suggest that "bowtie" neurons are commonly expressed in many GAL4 lines. In fact, a

MultiColor FlpOut (MCFO) study reported that these neurons are observed in more than

60% of all Janelia GAL4 lines (55). Only one line, FBS4, promotes sleep when

thermogenetically activated and expresses only in dFB cells, containing about 80% of

23E10-GAL4 dFB neurons. FBS4 can thus be considered as a dFB specific driver. We

note that the amount of sleep induction triggered by thermogenetic activation of FBS4 is

lower than in most other sleep-promoting FBS lines. However, the most surprising

finding from our anatomical analysis is FBS5, a line that has no expression in the dFB,

and still promotes sleep when activated. This clearly demonstrates that 23E10-GAL4

contains sleep-promoting neurons that are not dFB cells.

Because temperature has a strong impact on sleep (42, 43, 48, 49), we confirmed our

findings using an optogenetic activation method (Figure 3). Most sleep-promoting FBS

lines increased sleep when activated using either methods. The notable exception is the

dFB specific FBS4 line, which did not affect sleep at all upon optogenetic activation.

What is the source of this discrepancy? We can propose several scenarios. The first

possible explanation is that the threshold of activation needed for the neurons contained

in FBS4 to fire and promote sleep is high. This would imply that thermogenetic

activation, which is constant during the 24h activation day, provides stronger activation

than the fragmented optogenetic protocol. Considering that dFB neurons are subject to

dopaminergic regulation (33), we hypothesize that under baseline conditions, dFB

neurons are under severe dopaminergic inhibition and that our optogenetic activation

protocol is not sufficient to bring these cells above firing threshold. A second possibility

is that FBS4 expresses in neurons underlying the effect of high temperature on sleep.

This hypothesis is supported by a recent study demonstrating that some of the effects of

high temperature on sleep are mediated by GABAergic transmission on dFB neurons

(49). It is thus conceivable, that activating FBS4 neurons thermogenetically may

counteract the effect of increasing temperature on sleep. This would explain why we

only see a sleep-promoting effect when activating FBS4 using thermogenetic and not

optogenetic.

A further screen demonstrated that the VNC 'bowtie" neurons are sleep-promoting

(Figure 4). Activating these cells (renamed VNC-SP neurons) increases sleep, sleep

consolidation and sleep depth by increasing arousal thresholds. Moreover, chronic, and

acute silencing of VNC-SP neurons reduce sleep and sleep consolidation (Figure 5)

confirming that VNC-SP neurons positively regulate sleep. We then sought to discover

the neurochemical identity of VNC-SP neurons and identified acetylcholine as a key

modulator of sleep in VNC-SP neurons (Figure 6). Reducing cholinergic transmission in

VNC-SP neurons reduces sleep and sleep consolidation, an effect mirroring the

silencing experiments described earlier. Moreover, we demonstrate that VNC-SP

neurons use cholinergic transmission to promote sleep when optogenetically activated.

Since we observed 'bowtie" processes in most FBS lines, we assume that most of these

lines express in VNC-SP neurons. One could therefore wonder why we do not observe

total sleep changes when activating these lines? Importantly, our analysis of sleep

parameters including day and night sleep bout duration reveals that sleep consolidation

is significantly increased using at least one activation protocol with 7 of these "non

sleep-promoting" "bowtie" containing lines (FBS28, FBS33, FBS35, FBS57, FBS60,

FBS64 and FBS87). Thus, out of the 18 FBS lines that express in VNC-SP neurons, 16

of them increase at least one sleep parameter when activated. The only lines that

express in VNC-SP neurons and do not affect any sleep parameters when activated are

FBS1 and FBS58. Interestingly, we observed that the GFP signal in the VNC-SP

"bowtie" processes of these two lines is weak (Figure 2-figure supplement 1), indicating

reduced GAL4 expression in these cells. This observation may provide an explanation

for the lack of sleep changes when activating these lines. We also suspect that levels of

GAL4 expression in VNC-SP neurons could explain why some FBS lines are promoting

sleep more strikingly than others.

Where in the nervous system are these VNC-SP cholinergic signals received? We show

that VNC-SP neurons receive synaptic inputs in the VNC and have presynaptic sites in

the brain (Figure 6-figure supplement 2). This suggests that VNC-SP neurons are well

positioned to receive and integrate sleep-relevant sensory inputs from the periphery in

the VNC and that they are sending this information to the brain. The nature of these

inputs will require additional investigations that will further uncover the function and role

of VNC-SP neurons in sleep regulation. Known modulators of sleep include the immune

system and metabolic functions. Based on their location, it is possible that VNC-SP

neurons integrate signals relevant to these two sleep-modulating systems.

Our data raised important questions about the role of dFB neurons in sleep regulation.

Importantly, we show that 23E10-GAL4 dFB neurons are involved in sleep homeostasis

as silencing them completely abrogates sleep rebound following a night of sleep

deprivation (Figure 8). Our data is in agreement with previous work supporting a role for

dFB neurons in sleep homeostasis (29, 33). Interestingly, we also demonstrate that

VNC-SP neurons are not part of the homeostatic sleep circuit. These findings suggest

that dFB neurons and VNC-SP neurons regulate different aspects and functions of

sleep. Here again, further work will determine the role of VNC-SP neurons.

Finally, our data clearly highlight the need to obtain tools that are as specific as possible

when attempting to link a given behavior with a specific neuron or group of neurons. In

particular, the frequency at which VNC-SP neurons are observed in existing GAL4 lines

(in up to 60% of all Janelia GAL4 lines) (55) raises some important methodological

questions. For example, are VNC-SP neurons present in the expression pattern of a

GAL4 line that is known to modulate sleep and is seen as a tool specific for a given

structure in the brain? Future studies using GAL4 lines in sleep studies will need to take

account of the potential presence of VNC-SP neurons in their expression pattern.

Perhaps more importantly, existing data may need to be reinterpreted in the light of our

finding.

Conclusions

Our work identified a novel group of cholinergic sleep-promoting neurons located in the

VNC in *Drosophila*. We show that these cells are perfectly positioned to receive and

integrate sensory inputs from the periphery and send this information to the brain. We

demonstrate that these cells regulate sleep quantity and sleep quality. We further

demonstrate that they are not involved in sleep homeostasis whereas 23E10-GAL4 dFB

neurons are. Further work will need to determine the nature of the sleep-relevant

information that is relayed by VNC-SP neurons.

Materials and Methods

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (*D. melanogaster*)	P{GMR23E10-GAL4}attP2	Bloomington Drosophila Stock Center	RRID:BDSC_49032	
Genetic reagent (*D. melanogaster*)	w[*]; P{y[+t7.7] w[+mC]=UAS-TrpA1(B).K}attP16	Bloomington Drosophila Stock Center	RRID:BDSC_26263	
Genetic reagent (*D. melanogaster*)	Canton-S	Laboratory of Paul Shaw		
Genetic reagent (*D. melanogaster*)	w[*]; P{y[+t7.7] w[+mC]=10XUAS-IVS-mCD8::GFP}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_32186	
Genetic reagent (*D. melanogaster*)	w[*]; P{y[+t7.7] w[+mC]=10XUAS-IVS-mCD8::GFP}attP2	Bloomington Drosophila Stock Center	RRID:BDSC_32185	
Genetic reagent (*D. melanogaster*)	w[1118]; P{y[+t7.7] w[+mC]=23E10-GAL4.DBD}attP2/TM3	Bloomington Drosophila Stock Center	RRID:BDSC_69269	
Genetic reagent (*D. melanogaster*)	w[1118]; P{y[+t7.7] w[+mC]=p65.AD.Uw}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_71210	Empty-AD
Genetic reagent (*D. melanogaster*)	P{R84C10-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_70833	AD for FBS 4
Genetic reagent (*D. melanogaster*)	P{R23E12-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_70602	AD for FBS 5

Genetic reagent (*D. melanogaster*)	w[1118]; P{y[+t7.7] w[+mC]=VT050238-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_72962	AD for FBS 45
Genetic reagent (*D. melanogaster*)	P{VT023823-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_86629	AD for FBS 68
Genetic reagent (*D. melanogaster*)	P{R12D12-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_70539	AD for FBS 1
Genetic reagent (*D. melanogaster*)	P{R42H01-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_70687	AD for FBS 6
Genetic reagent (*D. melanogaster*)	P{R22A12-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_70120	AD for FBS 25
Genetic reagent (*D. melanogaster*)	P{R54G07-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_70737	AD for FBS 28
Genetic reagent (*D. melanogaster*)	P{R49C04-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_71075	AD for FBS 33
Genetic reagent (*D. melanogaster*)	P{R65C03-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_71005	AD for FBS 35
Genetic reagent (*D. melanogaster*)	P{VT058545-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_74090	AD for FBS 41
Genetic reagent (*D. melanogaster*)	P{R12G09-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_68826	AD for FBS 42
Genetic reagent (*D. melanogaster*)	P{VT043400-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_75874	AD for FBS 53
Genetic reagent (*D. melanogaster*)	P{VT025720-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_73432	AD for FBS 57
Genetic reagent (*D. melanogaster*)	P{VT064566-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_71451	AD for FBS 58
Genetic reagent (*D. melanogaster*)	P{VT023818-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_73262	AD for FBS 60
Genetic reagent (*D. melanogaster*)	P{VT027956-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_73061	AD for FBS 64

Genetic reagent (*D. melanogaster*)	P{VT034811-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_71337	AD for FBS 70
Genetic reagent (*D. melanogaster*)	P{VT026953-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_74413	AD for FBS 72
Genetic reagent (*D. melanogaster*)	P{R74G10-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_71131	AD for FBS 81
Genetic reagent (*D. melanogaster*)	P{VT007754-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_74138	AD for FBS 84
Genetic reagent (*D. melanogaster*)	P{VT045793-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_71441	AD for FBS 87
Genetic reagent (*D. melanogaster*)	w[1118]; P{y[+t7.7] w[+mC]=20XUAS-IVS-CsChrimson.mVenus}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_55135	
Genetic reagent (*D. melanogaster*)	P{VT020742-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_73460	
Genetic reagent (*D. melanogaster*)	P{R30A08-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_71007	
Genetic reagent (*D. melanogaster*)	P{VT013602-p65.AD}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_74209	
Genetic reagent (*D. melanogaster*)	P{GMR30A08-lexA}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_54756	
Genetic reagent (*D. melanogaster*)	y[1] w[*]; P{y[+t7.7] w[+mC]=10XUAS-IVS-mCD8::RFP}attP18 P{y[+t7.7] w[+mC]=13XLexAop2-mCD8::GFP}attP8	Bloomington Drosophila Stock Center	RRID:BDSC_32229	
Genetic reagent (*D. melanogaster*)	w[*]; P{y[+t7.7] w[+mC]=13XLexAop2-KZip+.3XHA}su(Hw)attP5/CyO; TM6B, Tb[1]/MKRS	Bloomington Drosophila Stock Center	RRID:BDSC_76253	
Genetic reagent (*D. melanogaster*)	w[*]; P{y[+t7.7] w[+mC]=13XLexAop2-KZip+.3XHA}attP2	Bloomington Drosophila Stock Center	RRID:BDSC_76254	
Genetic reagent (*D. melanogaster*)	P{VT020742-GAL4.DBD}attP2	Bloomington Drosophila Stock Center	RRID:BDSC_72912	

Genetic reagent (*D. melanogaster*)	w[1118]; P{y[+t7.7] w[+mC]=GAL4.DBD.Uw}attP2	Bloomington Drosophila Stock Center	RRID:BDSC_71209	Empty-DBD
Genetic reagent (*D. melanogaster*)	w[*]; P{w[+mC]=UAS-Hsap\KCNJ2.EGFP}7	Bloomington Drosophila Stock Center	RRID:BDSC_6595	UAS-Kir2.1
Genetic reagent (*D. melanogaster*)	pJFRC100-20XUAS-TTS-Shibire-ts1-p10 in VK00005	Laboratory of Gerry Rubin	(70)	UAS-Shi[ts1]
Genetic reagent (*D. melanogaster*)	GMR-GAL4	Laboratory of Jeff Price	(71)	
Genetic reagent (*D. melanogaster*)	P{GawB}elav[C155]	Bloomington Drosophila Stock Center	RRID:BDSC_458	elav-GAL4
Genetic reagent (*D. melanogaster*)	P{TRiP.HMC05021}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_60028	Expresses dsRNA for RNAi of ChAT under UAS control
Genetic reagent (*D. melanogaster*)	y[1] v[1]; P{y[+t7.7] v[+t1.8]=UAS-GFP.VALIUM10}attP2	Bloomington Drosophila Stock Center	RRID:BDSC_35786	Control RNAi line
Genetic reagent (*D. melanogaster*)	w[1118]; P{w[+mC]=UAS-Dcr-2.D}2	Bloomington Drosophila Stock Center	RRID:BDSC_24650	UAS-Dicer2 on chromosome 2
Genetic reagent (*D. melanogaster*)	w[1118]; P{w[+mC]=UAS-Dcr-2.D}10	Bloomington Drosophila Stock Center	RRID:BDSC_24651	UAS-Dicer2 on chromosome 3
Genetic reagent (*D. melanogaster*)	w[1118]; P{y[+t7.7] w[+mC]=20XUAS-IVS-CsChrimson.mVenus}attP2	Bloomington Drosophila Stock Center	RRID:BDSC_55136	
Genetic reagent (*D. melanogaster*)	w[1118]; P{y[+t7.7] w[+mC]=20XUAS-IVS-CsChrimson.mVenus}attP18	Bloomington Drosophila Stock Center	RRID:BDSC_55134	
Genetic reagent (*D. melanogaster*)	w[*]; Mi{Trojan-lexA:QFAD.0}ChAT[MI04508-TlexA:QFAD.0] CG7715[MI04508-TlexA:QFAD.0-X]/TM6B, Tb[1]	Bloomington Drosophila Stock Center	RRID:BDSC_60319	ChAT-LexA
Genetic reagent (*D. melanogaster*)	P{GMR23E10-lexA}attP40	Bloomington Drosophila Stock Center	RRID:BDSC_52693	

Genetic reagent (*D. melanogaster*)	w[1118]; P{y[+t7.7] w[+mC]=13XLexAop2-IVS-CsChrimson.mVenus}attP18	Bloomington Drosophila Stock Center	RRID:BDSC_55137	
Genetic reagent (*D. melanogaster*)	w[1118]; P{w[+mC]=UAS-syt.eGFP}3	Bloomington Drosophila Stock Center	RRID:BDSC_62138	
Genetic reagent (*D. melanogaster*)	w[*]; P{y[+t7.7] w[+mC]=5XUAS-DenMark::smGdP-V5}su(Hw)attP5	Bloomington Drosophila Stock Center	RRID:BDSC_6926	
Genetic reagent (*D. melanogaster*)	P{TRiP.JF01877}attP2 (RNAi against ChAT)	Bloomington Drosophila Stock Center	RRID:BDSC_25856	Expresses dsRNA for RNAi of ChAT under UAS control
Antibody	Anti-GFP (Chicken polyclonal)	Aves Labs, Inc	GFP-1020	dilution 1:1000
Antibody	anti-bruchpilot (mouse monoclonal)	Developmental Studies Hybridoma Bank	nc82-s	dilution 1:50
Antibody	Anti-DsRed (rabbit polyclonal)	Takara	632496	dilution 1:250
Antibody	Anti-ChAT (mouse monoclonal)	Developmental Studies Hybridoma Bank	Chat4B1-c	dilution 1:100
Antibody	Anti-VGlut (rabbit)	Aaron DiAntonio (Washington University)	(72)	dilution 1:500
Antibody	Anti-GABA (rabbit polyclonal)	Sigma-Aldrich	A2052	dilution 1:1000
Antibody	Anti-5-HT (rabbit polyclonal)	Sigma-Aldrich	S5545	dilution 1:500
Antibody	Anti-TH (mouse monoclonal)	Immunostar	22941	dilution 1:100
Antibody	Anti-Tdc2 (rabbit polyclonal)	Covalab	pab0822-P	dilution 1:200
Antibody	Anti-V5 (mouse monoclonal)	Sigma-Aldrich	V8012	dilution 1:400
Antibody	Goat anti-chicken Alexafluor 488	Invitrogen	A32931	dilution 1:800
Antibody	Goat anti-mouse Alexafluor 555	Invitrogen	A32727	dilution 1:400

Antibody	Goat anti-rabbit Alexafluor 555	Invitrogen	A32732	dilution 1:400
Antibody	Goat anti-mouse Alexafluor 633	Invitrogen	A21052	dilution 1:400
Antibody	Goat anti-rabbit Alexafluor 633	Invitrogen	A21071	dilution 1:400
Chemical compound, drug	retinal	Sigma	R2500	400µM
Chemical compound, drug	Vectashield	Vector Laboratories	H-1000	
Chemical compound, drug	20% paraformaldehyde	Electron Microscopy Sciences	15713	4% dilution
Media	Drosophila Schneider's Medium	Gibco	21720024	
Media	Normal Goat Serum	MP Biomedicals	191356	5% dilution

Flies

831   Flies were cultured at 25˚C with 50% humidity under a 12-hour light: 12-hour dark cycle.

832   Flies were kept on a standard yeast and molasses diet (per 1L: 50g yeast, 15g sucrose,

833   28g corn syrup, 33.3mL molasses, 9g agar).

Sleep

Sleep was assessed as previously described (32, 73). 4-10 day old virgin females or

males were used as described in figure legends. Briefly, flies were placed into individual

65 mm tubes and all activity was continuously measured through the Trikinetics

Drosophila Activity Monitoring System (www.Trikinetics.com, Waltham, Ma). Locomotor

activity was measured in 1-minute bins and sleep was defined as periods of quiescence

lasting at least 5 minutes.

Optogenetic activation

For optogenetic activation, flies expressing CsChrimson were loaded into individual

65mm tubes with food supplemented with 400μM all-trans retinal (Sigma, #R2500) and

kept in the dark for 48 hours. Sleep was then monitored under low-light conditions for 48

hours of baseline measurements. To activate CSChrimson, flies were put under 627nm

LEDs (LuxeonStar LXM2-PD01-0040) set to a pulse cycle of (5ms on, 95ms off) x 20

with a 4 second delay between pulse cycles. The light intensity of this protocol is

$0.024mW/mm^2$

Sleep deprivation

Sleep deprivation was performed as previously described (32, 74). Briefly, flies were

placed into individual 65 mm tubes and the sleep-nullifying apparatus (SNAP) was used

to sleep deprive flies for 12 hours during the dark phase. Sleep homeostasis was

calculated for each individual fly as the ratio of the minutes of sleep gained above

baseline during 24h of recovery sleep divided by the total min of sleep lost during 12 h

of sleep deprivation.

Arousal threshold

Retinal or vehicle-fed VNC-SP>UAS-CsChrimson flies were placed into individual 65

mm tubes in Trikinetics monitors. These were then loaded in the SNAP sleep

deprivation apparatus. The SNAP tilts asymmetrically from −60° to +60°, such that

sleeping flies are disturbed during each downward movement (75). Each individual

SNAP movement takes 15 seconds to complete. To assess arousal threshold, we

modified the SNAP parameters so that flies were subjected to only 1, 2, or 4 downward

movements in the SNAP every hour for 24h and their locomotor responses were

analyzed. We then averaged each individual hour response to obtain the percentage of

flies awakened by a given strength of perturbation.

QPCR

QPCR were performed as previously described (32, 76). Briefly, total RNA was isolated

from the bodies and eyes of ~10 flies per replicate with Trizol (Invitrogen, Carlsbad, CA)

and DNAse I digested. cDNA synthesis was performed using Superscript IV VILO

(Invitrogen) according to manufacturer protocol. cDNA samples were loaded in

duplicate and quantitatively analyzed using SYBR Green Master Mix (Applied

Biosystems) and gene specific primers using the Applied Biosystems 7500 system.

Expression values for RP49 were used to normalize results between groups. The

following primer sets were used: rp49: fw aagaagcgcaccaagcacttcatc, rev

tctgttgtcgatacccttgggctt and ChAT: fw caccgagcgatacaggatgg, rev

ggcaccttgggtagagtgtc.

Immunocytochemistry

Flies were cold anesthetized, brains and VNC were dissected in ice cold Schneider's

Drosophila Medium (Gibco, 21720024) and blocked in 5% normal goat serum.

Following blocking flies were incubated overnight at 4˚C in primary antibody, washed in

PBST and incubated overnight at 4˚C in secondary antibody. Description of antibodies

used can be found in the resources table. Brains and VNCs were mounted on polylysine

treated slides in Vectashield H-1000 mounting medium. Imaging was performed on a

Zeiss 510 meta confocal microscope using a Plan-Apochromat 20x or Plan-Neofluar

40x objective. Z-series images were acquired with a 1 uM slice size using the same

settings (laser power, gain, offset) to allow for comparison across genotypes. Images

were processed and analyzed using ImageJ.

Statistics

Statistical analyses were performed with Prism9 software (GraphPad). Normal

distribution was assessed with the D'Agostino-Pearson test. Normally distributed data

were analyzed with parametric statistics: t-test, one-way analysis of variance or two-way

ANOVA followed by the planned pairwise multiple comparisons as described in the

legends. For data that significantly differed from the normal distribution, non-parametric

statistics were applied, Mann-Whitney U test or Kruskal–Wallis test followed by Dunn's

multiple test. Some non-normally distributed data were subjected to log transformation

or Box-Cox transformation before two-way ANOVA followed by planned pairwise

multiple comparisons as described in the legends. All statistically different groups are

defined as $*P < 0.05$, $**P<0.01$, $***P<0.001$ and $****P<0.0001$.

Author Contributions

S.D., J.J. and J.M.E. designed the experiments and wrote the paper. J.J., B.H., K.E.,

A.V., A.V., A.E., J.M.E and S.D. performed the experiments. J.J., J.M.E and S.D.

analyzed the data.

Acknowledgments

912 We thank Aaron DiAntonio, Gerry Rubin, Paul Shaw and Jeff Price for sharing reagents.

Competing Interests

915 The authors have no competing interests.

References

1.   Park HG, Carmel JB. Selective Manipulation of Neural Circuits. Neurotherapeutics.
2016;13(2):311-24.
2.   Moulin TC, Covill LE, Itskov PM, Williams MJ, Schioth HB. Rodent and fly models in behavioral
neuroscience: An evaluation of methodological advances, comparative research, and future
perspectives. Neurosci Biobehav Rev. 2021;120:1-12.
3.   Yamamoto D, Koganezawa M. Genes and circuits of courtship behaviour in Drosophila males.
Nat Rev Neurosci. 2013;14(10):681-92.
4.   Anderson DJ. Circuit modules linking internal states and social behaviour in flies and mice. Nat
Rev Neurosci. 2016;17(11):692-704.
5.   Bhumika, Singh AK. Regulation of feeding behavior in Drosophila through the interplay of
gustation, physiology and neuromodulation. Front Biosci (Landmark Ed). 2018;23:2016-27.
6.   Scott K. Gustatory Processing in Drosophila melanogaster. Annu Rev Entomol. 2018;63:15-30.
7.   Ly S, Pack AI, Naidoo N. The neurobiological basis of sleep: Insights from Drosophila. Neurosci
Biobehav Rev. 2018;87:67-86.
8.   Brand AH, Perrimon N. Targeted gene expression as a means of altering cell fates and generating
dominant phenotypes. Development. 1993;118(2):401-15.
9.   Lai SL, Lee T. Genetic mosaic with dual binary transcriptional systems in Drosophila. Nat
Neurosci. 2006;9(5):703-9.
10.   Potter CJ, Tasic B, Russler EV, Liang L, Luo L. The Q system: a repressible binary system for
transgene expression, lineage tracing, and mosaic analysis. Cell. 2010;141(3):536-48.
11.   Luan H, Peabody NC, Vinson CR, White BH. Refined spatial manipulation of neuronal function by
combinatorial restriction of transgene expression. Neuron. 2006;52(3):425-36.

12.   Keene AC, Duboue ER. The origins and evolution of sleep. J Exp Biol. 2018;221(Pt 11).

13.   Borbely AA. A two process model of sleep regulation. Hum Neurobiol. 1982;1(3):195-204.

14.   Krause AJ, Simon EB, Mander BA, Greer SM, Saletin JM, Goldstein-Piekarski AN, et al. The sleep-
deprived human brain. Nat Rev Neurosci. 2017;18(7):404-18.

15.   Miyazaki S, Liu CY, Hayashi Y. Sleep in vertebrate and invertebrate animals, and insights into the
function and evolution of sleep. Neurosci Res. 2017;118:3-12.

16.   Shafer OT, Keene AC. The Regulation of Drosophila Sleep. Curr Biol. 2021;31(1):R38-R49.

17.   Dissel S. Drosophila as a Model to Study the Relationship Between Sleep, Plasticity, and
Memory. Front Physiol. 2020;11:533.

18.   Strauss R. The central complex and the genetic dissection of locomotor behaviour. Curr Opin
Neurobiol. 2002;12(6):633-8.

19.   Sakai T, Kitamoto T. Differential roles of two major brain structures, mushroom bodies and
central complex, for Drosophila male courtship behavior. J Neurobiol. 2006;66(8):821-34.

20.   Joiner MA, Griffith LC. Mapping of the anatomical circuit of CaM kinase-dependent courtship
conditioning in Drosophila. Learn Mem. 1999;6(2):177-92.

21.   Sakai T, Inami S, Sato S, Kitamoto T. Fan-shaped body neurons are involved in period-dependent
regulation of long-term courtship memory in Drosophila. Learn Mem. 2012;19(12):571-4.

22.   Hu W, Peng Y, Sun J, Zhang F, Zhang X, Wang L, et al. Fan-Shaped Body Neurons in the
Drosophila Brain Regulate Both Innate and Conditioned Nociceptive Avoidance. Cell Rep.
2018;24(6):1573-84.

23.   Liu G, Seiler H, Wen A, Zars T, Ito K, Wolf R, et al. Distinct memory traces for two visual features
in the Drosophila brain. Nature. 2006;439(7076):551-6.

24.   Weir PT, Dickinson MH. Functional divisions for visual processing in the central brain of flying
Drosophila. Proc Natl Acad Sci U S A. 2015;112(40):E5523-32.

25.   Kacsoh BZ, Bozler J, Hodge S, Bosco G. Neural circuitry of social learning in Drosophila requires
multiple inputs to facilitate inter-species communication. Commun Biol. 2019;2:309.

26.   Sareen PF, McCurdy LY, Nitabach MN. A neuronal ensemble encoding adaptive choice during
sensory conflict in Drosophila. Nat Commun. 2021;12(1):4131.

27.   Donlea JM, Thimgan MS, Suzuki Y, Gottschalk L, Shaw PJ. Inducing sleep by remote control
facilitates memory consolidation in Drosophila. Science. 2011;332(6037):1571-6.

28.   Ueno T, Tomita J, Tanimoto H, Endo K, Ito K, Kume S, et al. Identification of a dopamine pathway
that regulates sleep and arousal in Drosophila. Nat Neurosci. 2012;15(11):1516-23.

29.   Donlea JM, Pimentel D, Miesenbock G. Neuronal Machinery of Sleep Homeostasis in Drosophila.
Neuron. 2014;81(6):1442.

30.   Liu Q, Liu S, Kodama L, Driscoll MR, Wu MN. Two dopaminergic neurons signal to the dorsal fan-
shaped body to promote wakefulness in Drosophila. Curr Biol. 2012;22(22):2114-23.

31.   Dag U, Lei Z, Le JQ, Wong A, Bushey D, Keleman K. Neuronal reactivation during post-learning
sleep consolidates long-term memory in Drosophila. Elife. 2019;8.

32.   Dissel S, Angadi V, Kirszenblat L, Suzuki Y, Donlea J, Klose M, et al. Sleep restores behavioral
plasticity to Drosophila mutants. Curr Biol. 2015;25(10):1270-81.

33.   Pimentel D, Donlea JM, Talbot CB, Song SM, Thurston AJF, Miesenbock G. Operation of a
homeostatic sleep switch. Nature. 2016;536(7616):333-7.

34.   Kempf A, Song SM, Talbot CB, Miesenbock G. A potassium channel beta-subunit couples
mitochondrial electron transport to sleep. Nature. 2019;568(7751):230-4.

35.   Alam MA, Kumar S, McGinty D, Alam MN, Szymusiak R. Neuronal activity in the preoptic
hypothalamus during sleep deprivation and recovery sleep. J Neurophysiol. 2014;111(2):287-99.

36.   Jenett A, Rubin GM, Ngo TT, Shepherd D, Murphy C, Dionne H, et al. A GAL4-driver line resource
for Drosophila neurobiology. Cell Rep. 2012;2(4):991-1001.

37.   Qian Y, Cao Y, Deng B, Yang G, Li J, Xu R, et al. Sleep homeostasis regulated by 5HT2b receptor in
a small subset of neurons in the dorsal fan-shaped body of drosophila. Elife. 2017;6.
38.   Yap MHW, Grabowska MJ, Rohrscheib C, Jeans R, Troup M, Paulk AC, et al. Oscillatory brain
activity in spontaneous and induced sleep stages in flies. Nat Commun. 2017;8(1):1815.
39.   Berry JA, Cervantes-Sandoval I, Chakraborty M, Davis RL. Sleep Facilitates Memory by Blocking
Dopamine Neuron-Mediated Forgetting. Cell. 2015;161(7):1656-67.
40.   Hulse BK, Haberkern H, Franconville R, Turner-Evans DB, Takemura SY, Wolff T, et al. A
connectome of the Drosophila central complex reveals network motifs suitable for flexible navigation
and context-dependent action selection. Elife. 2021;10.
41.   Hamada FN, Rosenzweig M, Kang K, Pulver SR, Ghezzi A, Jegla TJ, et al. An internal thermal
sensor controlling temperature preference in Drosophila. Nature. 2008;454(7201):217-20.
42.   Ishimoto H, Lark A, Kitamoto T. Factors that Differentially Affect Daytime and Nighttime Sleep in
Drosophila melanogaster. Front Neurol. 2012;3:24.
43.   Lamaze A, Ozturk-Colak A, Fischer R, Peschel N, Koh K, Jepson JE. Regulation of sleep plasticity
by a thermo-sensitive circuit in Drosophila. Sci Rep. 2017;7:40304.
44.   Donlea JM, Pimentel D, Talbot CB, Kempf A, Omoto JJ, Hartenstein V, et al. Recurrent Circuitry
for Balancing Sleep Need and Sleep. Neuron. 2018;97(2):378-89 e4.
45.   Tainton-Heap LAL, Kirszenblat LC, Notaras ET, Grabowska MJ, Jeans R, Feng K, et al. A
Paradoxical Kind of Sleep in Drosophila melanogaster. Curr Biol. 2021;31(3):578-90 e6.
46.   Tirian L, Dickson BJ. The VT GAL4, LexA, and split-GAL4 driver line collections for targeted
expression in the *Drosophila* nervous system. BioRxiv 198648 [Preprint] October 5, 2017 Available from:
https://doiorg/101101/198648. 2017.
47.   Dionne H, Hibbard KL, Cavallaro A, Kao JC, Rubin GM. Genetic Reagents for Making Split-GAL4
Lines in Drosophila. Genetics. 2018;209(1):31-5.
48.   Parisky KM, Agosto Rivera JL, Donelson NC, Kotecha S, Griffith LC. Reorganization of Sleep by
Temperature in Drosophila Requires Light, the Homeostat, and the Circadian Clock. Curr Biol.
2016;26(7):882-92.
49.   Kim JH, Ki Y, Lee H, Hur MS, Baik B, Hur JH, et al. The voltage-gated potassium channel Shaker
promotes sleep via thermosensitive GABA transmission. Commun Biol. 2020;3(1):174.
50.   van Alphen B, Yap MH, Kirszenblat L, Kottler B, van Swinderen B. A dynamic deep sleep stage in
Drosophila. J Neurosci. 2013;33(16):6917-27.
51.   Isaac RE, Li C, Leedale AE, Shirras AD. Drosophila male sex peptide inhibits siesta sleep and
promotes locomotor activity in the post-mated female. Proc Biol Sci. 2010;277(1678):65-70.
52.   Khericha M, Kolenchery JB, Tauber E. Neural and non-neural contributions to sexual dimorphism
of mid-day sleep in Drosophila melanogaster: a pilot study. Physiol Entomol. 2016;41(4):327-34.
53.   Wu B, Ma L, Zhang E, Du J, Liu S, Price J, et al. Sexual dimorphism of sleep regulated by juvenile
hormone signaling in Drosophila. PLoS Genet. 2018;14(4):e1007318.
54.   Kim H, Kirkhart C, Scott K. Long-range projection neurons in the taste circuit of Drosophila. Elife.
2017;6.
55.   Meissner GW, Dorman Z, Nern A, Forster K, Gibney T, Jeter J, et al. An image resource of
subdivided Drosophila GAL4-driver expression patterns for neuron-level searches. bioRxiv.
2020:2020.05.29.080473.
56.   Klapoetke NC, Murata Y, Kim SS, Pulver SR, Birdsey-Benson A, Cho YK, et al. Independent optical
excitation of distinct neural populations. Nat Methods. 2014;11(3):338-46.
57.   Otsuna H, Ito M, Kawase T. Color depth MIP mask search: a new tool to expedite Split-GAL4
creation. BioRxiv 318006 [Preprint] May 9, 2018 Available from: https://doiorg/101101/318006v1.
2018.

58.   Dolan MJ, Luan H, Shropshire WC, Sutcliffe B, Cocanougher B, Scott RL, et al. Facilitating Neuron-
Specific Genetic Manipulations in Drosophila melanogaster Using a Split GAL4 Repressor. Genetics.
2017;206(2):775-84.
59.   Baines RA, Uhler JP, Thompson A, Sweeney ST, Bate M. Altered electrical properties in
Drosophila neurons developing without synaptic transmission. J Neurosci. 2001;21(5):1523-31.
60.   Kitamoto T. Conditional modification of behavior in Drosophila by targeted expression of a
temperature-sensitive shibire allele in defined neurons. J Neurobiol. 2001;47(2):81-92.
61.   Lacin H, Chen HM, Long X, Singer RH, Lee T, Truman JW. Neurotransmitter identity is acquired in
a lineage-restricted manner in the Drosophila CNS. Elife. 2019;8.
62.   Diao F, Ironfield H, Luan H, Diao F, Shropshire WC, Ewer J, et al. Plug-and-play genetic access to
drosophila cell types using exchangeable exon cassettes. Cell Rep. 2015;10(8):1410-21.
63.   Ni JD, Gurav AS, Liu W, Ogunmowo TH, Hackbart H, Elsheikh A, et al. Differential regulation of
the Drosophila sleep homeostat by circadian and arousal inputs. Elife. 2019;8.
64.   Allen AM, Neville MC, Birtles S, Croset V, Treiber CD, Waddell S, et al. A single-cell transcriptomic
atlas of the adult Drosophila ventral nerve cord. Elife. 2020;9.
65.   Court R, Namiki S, Armstrong JD, Borner J, Card G, Costa M, et al. A Systematic Nomenclature for
the Drosophila Ventral Nerve Cord. Neuron. 2020;107(6):1071-9 e2.
66.   Tsubouchi A, Yano T, Yokoyama TK, Murtin C, Otsuna H, Ito K. Topological and modality-specific
representation of somatosensory information in the fly brain. Science. 2017;358(6363):615-23.
67.   Nicolai LJ, Ramaekers A, Raemaekers T, Drozdzecki A, Mauss AS, Yan J, et al. Genetically encoded
dendritic marker sheds light on neuronal connectivity in Drosophila. Proc Natl Acad Sci U S A.
2010;107(47):20553-8.
68.   Zhang YQ, Rodesch CK, Broadie K. Living synaptic vesicle marker: synaptotagmin-GFP. Genesis.
2002;34(1-2):142-5.
69.   Pfeiffer BD, Ngo TT, Hibbard KL, Murphy C, Jenett A, Truman JW, et al. Refinement of tools for
targeted gene expression in Drosophila. Genetics. 2010;186(2):735-55.
70.   Pfeiffer BD, Truman JW, Rubin GM. Using translational enhancers to increase transgene
expression in Drosophila. Proc Natl Acad Sci U S A. 2012;109(17):6626-31.
71.   Takahashi Y, Hirose F, Matsukage A, Yamaguchi M. Identification of three conserved regions in
the DREF transcription factors from Drosophila melanogaster and Drosophila virilis. Nucleic Acids Res.
1999;27(2):510-6.
72.   Daniels RW, Collins CA, Chen K, Gelfand MV, Featherstone DE, DiAntonio A. A single vesicular
glutamate transporter is sufficient to fill a synaptic vesicle. Neuron. 2006;49(1):11-6.
73.   Shaw PJ, Cirelli C, Greenspan RJ, Tononi G. Correlates of sleep and waking in Drosophila
melanogaster. Science. 2000;287(5459):1834-7.
74.   Seugnet L, Suzuki Y, Vine L, Gottschalk L, Shaw PJ. D1 receptor activation in the mushroom
bodies rescues sleep-loss-induced learning impairments in Drosophila. Curr Biol. 2008;18(15):1110-7.
75.   Shaw PJ, Tononi G, Greenspan RJ, Robinson DF. Stress response genes protect against lethal
effects of sleep deprivation in Drosophila. Nature. 2002;417(6886):287-91.
76.   Dissel S, Seugnet L, Thimgan MS, Silverman N, Angadi V, Thacher PV, et al. Differential activation
of immune factors in neurons and glia contribute to individual differences in resilience/vulnerability to
sleep disruption. Brain Behav Immun. 2015;47:75-85.

1077

1078

 Figure Legends

 Figure 1. The 23E10-GAL4 driver contains many neurons outside the dFB.

A) Diagram of the experimental assay. Sleep was measured at 22°C for 2 days to
establish baseline sleep profile. Flies were then shifted to 31°C for 24h at the start of
day3 to increase activity of the targeted cells by activating the TrpA1 channel, and then
returned to 22°C. White bars (L) represent the 12h of light and black bars (D) represent
the 12h of dark that are oscillating daily.

B-C) Sleep profile in minutes of sleep per hour for day 2 (22°C, blue line) and day 3
(31°C, red line) for parental control female flies: *23E10-Gal4/+* (B) and *UAS-TrpA1/+*
(C).

D) Sleep profile in minutes of sleep per hour for day 2 (22°C, blue line) and day 3 (31°C,
red line) for *23E10-GAL4>UAS-TrpA1* female flies.

E) Box plots of total sleep change in % ((total sleep on day3-total sleep on day2/total
sleep on day2) X 100) for data presented in B-D. The bottom and top of each box
represents the first and third quartile, and the horizontal line dividing the box is the
median. The whiskers represent the 10th and 90th percentiles. Flies expressing UAS-
TrpA1 in 23E10-GAL4 significantly increase sleep when switched to 31°C compared
with parental controls, Kruskal-Wallis ANOVA followed by Dunn's multiple comparisons.
****$P<0.0001$, n=25-31 flies per genotype.

F) Box plots of locomotor activity counts per minute awake for flies presented in B-D.
The bottom and top of each box represents the first and third quartile, and the horizontal
line dividing the box is the median. The whiskers represent the 10th and 90th percentiles.
Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test
found no differences in locomotor activity between 22°C and 31°C. n.s.= not significant.
n=25-31 flies per genotype.

G-H) Representative confocal stacks of a female *23E10-GAL4>UAS-mCD8GFP* brain
(G) and ventral nerve cord (VNC, H). Green, anti-GFP; magenta, anti-nc82 (neuropile
marker).

Figure 2: Thermogenetic screen of FBS lines.

A) Diagram of the experimental assay. Sleep was measured at 22°C for 2 days to establish baseline sleep profile. Flies were then shifted to 31°C for 24h at the start of day3 to increase activity of the targeted cells by activating the TrpA1 channel, and then returned to 22°C. White bars (L) represent the 12h of light and black bars (D) represent the 12h of dark that are oscillating daily.

B) Box plots of total sleep change in % ((total sleep on day3-total sleep on day2/total sleep on day2) X 100) for female control (Empty-AD; 23E10-DBD) and 22 FBS lines expressing UAS-TrpA1; UAS-GFP. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10^{th} and 90^{th} percentiles. The gray rectangle spanning the horizontal axis indicates the interquartile range of the control. Kruskal-Wallis ANOVA followed by Dunn's multiple comparisons revealed that 9 FBS lines increase sleep significantly more than control flies when transferred to 31°C. **$P<0.01$, ***$P<0.001$, ****$P<0.0001$, n=30-51 flies per genotype.

C) Box plots of locomotor activity counts per minute awake for flies presented in B. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10^{th} and 90^{th} percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test found that for 3 sleep-promoting FBS lines (FBS5, FBS45 and FBS53) locomotor activity per awake time is increased while no differences are seen for the other 6 sleep-promoting lines between 22°C and 31°C. **$P<0.01$, ***$P<0.001$, ****$P<0.0001$, n=30-51 flies per genotype.

D-H) Left, sleep profile in minutes of sleep per hour for day 2 (22°C, blue line) and day 3 (31°C, red line) for empty control (Empty-AD; 23E10-DBD, D), FBS4 (E), FBS5 (F), FBS45 (G) and FBS68 (H) female flies expressing TrpA1 and GFP. Right, Representative confocal stacks for each *FBS>UAS-TrpA1; UAS-mCD8GFP* line of a female brain (left panel), VNC (middle panel) as well as a magnified view of VNC "bowtie" processes as indicated by the gray rectangle in F. Yellow arrows in F indicate TPN1-like processes in FBS5. Green, anti-GFP; magenta, anti-nc82 (neuropile marker).

Figure 3: Optogenetic screen of FBS lines.

A) Diagram of the experimental assay. Sleep was measured in retinal-fed and vehicle-fed flies for 2 days without 627nm LED stimulation to establish baseline sleep profile. LEDs were were then turned on for 24h at the start of day3 to increase activity of the targeted cells by activating the CsChrimson channel, and then turned off on day4. White bars (L) represent the 12h of light and black bars (D) represent the 12h of dark that are oscillating daily.

B) Box plots of total sleep change in % ((total sleep on day3-total sleep on day2/total sleep on day2) X 100) for control (Empty) and 22 experimental vehicle-fed and retinal-fed female flies expressing CsChrimson upon 627nm LED stimulation. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10^{th} and 90^{th} percentiles. Two-way ANOVA followed by Sidak's multiple comparisons revealed that 7 retinal-fed FBS lines increase sleep significantly when stimulated with 627nm LEDs when compared with vehicle-fed flies. **$P<0.01$, ***$P<0.001$, ****$P<0.0001$, n=21-39 flies per genotype and condition.

C) Box plots of locomotor activity counts per minute awake for retinal-fed flies presented in B. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10^{th} and 90^{th} percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test found that for most sleep-promoting FBS lines, locomotor activity per awake time is not affected when the flies are stimulated with 627nm LEDs while it is increased in *FBS45>UAS-CsChrimson* flies. ****$P<0.0001$, n=21-38 flies per genotype.

D) Box plots of daytime sleep bout duration (in minutes) for retinal-fed flies presented in B. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10^{th} and 90^{th} percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons indicate that daytime sleep bout duration is increased in 10 FBS lines expressing CsChrimson when stimulated with 627nm LEDs. *$P<0.05$, **$P<0.01$, ****$P<0.0001$, n=21-38 flies per genotype.

E) Box plots of nighttime sleep bout duration (in minutes) for retinal-fed flies presented in B. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons indicate that nighttime sleep bout duration is increased in 6 FBS lines expressing CsChrimson when stimulated with 627nm LEDs. ****$P<0.0001$, n=21-38 flies per genotype.

Figure 4: The 23E10-GAL driver contains sleep-promoting neurons that are located in the VNC.

A-D) Representative confocal stacks of a female *VT020742-AD; 23E10-DBD>UAS-GFP* brain (A), brain sagittal view (B), VNC (C) and VNC sagittal view (D). The cell bodies panel in C is a magnified view of the metathoracic area of the VNC. Yellow arrows indicate TPN1 like processes and gray arrows "bowtie" processes. Green, anti-GFP; magenta, anti-nc82 (neuropile marker). A= anterior, P= posterior, V= ventral and D= dorsal.

E-F) Representative confocal stacks of a female *30A08-AD; 23E10-DBD>UAS-GFP* brain (E) and VNC (F). The cell bodies panel in F is a magnified view of the metathoracic area of the VNC. Yellow arrows indicate TPN1 processes in the brain and VNC. Green, anti-GFP.

G-J) Representative confocal stacks of a female *VT013602-AD; 23E10-DBD>UAS-GFP* brain (G), brain sagittal view (H), VNC (I) and VNC sagittal view (J). The cell bodies panel in I is a magnified view of the metathoracic area of the VNC. Gray arrows indicate "bowtie" processes and cell bodies in I and J. Green, anti-GFP; magenta, anti-nc82 (neuropile marker). A= anterior, P= posterior, V= ventral and D= dorsal.

K) Box plots of total sleep change in % for control (*Empty-AD; 23E10-DBD>UAS-CsChrimson*), *VT020742-AD; 23E10-DBD>UAS-CsChrimson*, *30A08-AD; 23E10-DBD>UAS-CsChrimson* and *VT013602-AD; 23E10-DBD>UAS-CsChrimson* vehicle-fed and retinal-fed female flies upon 627nm LED stimulation. The bottom and top of each

box represents the first and third quartile, and the horizontal line dividing the box is the

median. The whiskers represent the 10th and 90th percentiles. Two-way ANOVA

followed by Sidak's multiple comparisons revealed that retinal-fed *VT020742-AD;*

23E10-DBD>UAS-CsChrimson and *VT013602-AD; 23E10-DBD>UAS-CsChrimson* flies

increase sleep significantly when stimulated with 627nm LEDs when compared with

vehicle-fed flies. **$P<0.01$, ****$P<0.0001$, n.s.= not significant. n=13-34 flies per

genotype and condition.

L) Box plots of locomotor activity counts per minute awake for retinal-fed flies presented

in K. The bottom and top of each box represents the first and third quartile, and the

horizontal line dividing the box is the median. The whiskers represent the 10th and

90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple

comparisons test shows that locomotor activity per awake time is not affected when the

flies are stimulated with 627nm LEDs. n.s.= not significant. n=15-34 flies per genotype.

M) Box plots of daytime sleep bout duration (in minutes) for retinal-fed flies presented in

K. The bottom and top of each box represents the first and third quartile, and the

horizontal line dividing the box is the median. The whiskers represent the 10th and

90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple

comparisons indicates that daytime sleep bout duration is increased in retinal-fed

VT020742-AD; 23E10-DBD>UAS-CsChrimson and *VT013602-AD; 23E10-DBD>UAS-*

CsChrimson flies when stimulated with 627nm LEDs. **$P<0.01$, ****$P<0.0001$, n.s.= not

significant. n=15-34 flies per genotype.

N) Box plots of nighttime sleep bout duration (in minutes) for retinal-fed flies presented

in K. The bottom and top of each box represents the first and third quartile, and the

horizontal line dividing the box is the median. The whiskers represent the 10th and

90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple

comparisons indicates that nighttime sleep bout duration is increased in retinal-fed

VT020742-AD; 23E10-DBD>UAS-CsChrimson and *VT013602-AD; 23E10-DBD>UAS-*

CsChrimson flies when stimulated with 627nm LEDs. ***$P<0.001$, ****$P<0.0001$, n.s.=

not significant. n=15-34 flies per genotype.

1225

Figure 5: Silencing VNC-SP neurons reduces sleep.

A) and **F)** Sleep profile in minutes of sleep per hour for control (*Empty-AD; 23E10-DBD>UAS-Kir2.1,* blue line) and *VNC-SP>Kir2.1 (VT013602-AD; 23E10-DBD>UAS-Kir2.1,* red line) female (A) and male (F) flies.

B) and **G)** Box plots of total sleep time (in minutes) for flies presented in A and F, respectively. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. A two-tailed unpaired t test revealed that total sleep is significantly reduced in *VNC-SP>Kir2.1* female flies compared to controls (B). A two-tailed Mann-Whitney U test revealed that total sleep is significantly reduced in *VNC-SP>Kir2.1* male flies compared to controls (G). ****P<0.0001. n=58-60 flies per genotype for females and n=48 flies per genotype for males.

C) and **H)** Box plots of daytime sleep bout duration (in minutes) for flies presented in A and F, respectively. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-tailed Mann-Whitney U tests revealed that daytime sleep bout duration is significantly reduced in *VNC-SP>Kir2.1* female (C) and male (H) flies compared to controls. *P<0.05, ***P<0.0001. n=58-60 flies per genotype for females and n=48 flies per genotype for males.

D) and **I)** Box plots of nighttime sleep bout duration (in minutes) for flies presented in A and F, respectively. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-tailed Mann-Whitney U tests revealed that nighttime sleep bout duration is significantly reduced in *VNC-SP>Kir2.1* female (D) and male (I) flies compared to controls. ***P<0.001. n=58-60 flies per genotype for females and n=48 flies per genotype for males.

E) and **J)** Box plots of locomotor activity counts per minute awake for flies presented in A and F, respectively. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent

the 10th and 90th percentiles. Two-tailed Mann-Whitney U tests revealed that activity per

minute awake is significantly reduced in *VNC-SP>Kir2.1* female (E) while it is

unchanged in male (J) flies compared to controls. **P<0.001, n.s.= not significant. n=58-

60 flies per genotype for females and n=48 flies per genotype for males.

K) and **N)** Sleep profile in minutes of sleep per hour for control (*Empty-AD; 23E10-*

DBD>UAS-Shi^{ts1}) female (K) and male (N) flies at 22°C (blue line) and 32°C (red line).

L) and **O)** Sleep profile in minutes of sleep per hour for *VNC-SP>UAS-Shi^{ts1}* female (L)

and male (O) flies at 22°C (blue line) and 32°C (red line).

M) and **P)** Box plots of nighttime sleep change in % for female flies presented in K and L

(M) and male flies presented in N and O. The bottom and top of each box represents

the first and third quartile, and the horizontal line dividing the box is the median. The

whiskers represent the 10th and 90th percentiles. Two-tailed unpaired t tests revealed

that *VNC-SP>Shi^{ts1}* flies lose significantly more sleep when transferred to 32°C

compared with controls in both males and females. *P<0.05. n=27-31 flies per

genotype for females and n=31 flies per genotype for males.

1270

Figure 6: VNC-SP neurons are cholinergic.

A-F) Representative confocal stacks focusing on the metathoracic ganglion of the VNC

of female *VT013602-AD; 23E10-DBD>UAS-GFP* flies stained with antibodies to ChAT

(A), VGlut (B), GABA (C), 5HT (D), TH (E) and Tdc2 (F). Gray arrows in A indicate

colocalization of GFP and ChAT staining in VNC-SP neurons. Green, anti-GFP;

magenta, anti-ChAT (A), anti-VGlut (B), anti-GABA (C), anti-5HT (D), anti-TH (E) and

anti-Tdc2 (F).

G) Sleep profile in minutes of sleep per hour for control (*Empty-AD; 23E10-DBD>UAS-*

ChAT^{RNAi 60028}, blue line) and *VNC-SP>ChAT^{RNAi}* (*VT013602-AD; 23E10-DBD> UAS-*

ChAT^{RNAi 60028}, red line) female flies.

H) Box plots of total sleep time (in minutes) for flies presented in G. The bottom and top

of each box represents the first and third quartile, and the horizontal line dividing the

box is the median. The whiskers represent the 10th and 90th percentiles. A two-tailed

unpaired t test revealed that *VNC-SP> ChAT^RNAi* flies sleep significantly less than

controls. ****P<0.0001. n=22-30 flies per genotype.

I) Box plots of daytime sleep bout duration (in minutes) for flies presented in G. The

bottom and top of each box represents the first and third quartile, and the horizontal line

dividing the box is the median. The whiskers represent the 10th and 90th percentiles. A

two-tailed Mann-Whitney U test revealed that *VNC-SP> ChAT^RNAi* flies daytime sleep

bout duration is significantly reduced compared with controls. ****P<0.0001. n=22-30

flies per genotype.

J) Box plots of nighttime sleep bout duration (in minutes) for flies presented in G. The

bottom and top of each box represents the first and third quartile, and the horizontal line

dividing the box is the median. The whiskers represent the 10th and 90th percentiles. A

two-tailed Mann-Whitney U test revealed that *VNC-SP> ChAT^RNAi* flies nighttime sleep

bout duration is significantly reduced compared with controls. ****P<0.0001. n=22-30

flies per genotype.

K) Box plots of locomotor activity counts per minute awake for flies presented in G. The

bottom and top of each box represents the first and third quartile, and the horizontal line

dividing the box is the median. The whiskers represent the 10th and 90th percentiles. A

two-tailed Mann-Whitney U test revealed that there is no difference in waking activity

between *VNC-SP> ChAT^RNAi* flies and controls. n.s= not significant. n=22-30 flies per

genotype.

L) Box plots of total sleep change in % for vehicle-fed and retinal-fed *VT013602-AD;*

23E10-DBD>UAS-Cs Chrimson; UAS-GFP.VALIUM10 (control) and *VT013602-AD;*

23E10-DBD>UAS-Cs Chrimson; UAS-ChAT^RNAi 60028 flies stimulated by 627nm LEDs.

The bottom and top of each box represents the first and third quartile, and the horizontal

line dividing the box is the median. The whiskers represent the 10th and 90th percentiles.

A two-way ANOVA followed by Sidak's multiple comparisons shows that sleep is

significantly increased in control flies and that expressing ChAT RNAi in VNC-SP

neurons completely abolishes the sleep-promoting effect of VNC-SP neurons activated

by 627nm LEDs. ****P<0.0001, n.s= not significant. n=31-36 flies per genotype and
condition.

1314

Figure 7: The 23E10-LexA driver does not express in VNC-SP neurons.

A-E) Representative confocal stacks of a female *23E10-LexA>LexAop2-GFP; 23E10-*
GAL4>UAS-RFP fly showing the brain (A), a magnified view of dFB neurons (B), the
VNC (C), VNC-SP neurons projections in the brain (D) and TPN1 neurons processes in
the VNC (E). Yellow arrows in B show dFB neurons that are labeled by both the 23E10-
LexA and 23E10-GAL4 drivers, while the gray arrow shows a neuron that is only labeled
by 23E10-GAL4. Our analysis revealed that in these co-labelling experiments, 23E10-
LexA expresses in 13-20 dFB neurons while 23E10-GAL4 labels 19-26 dFB cells per
brain. Importantly, all the neurons labeled by 23E10-LexA are part of the 23E10-GAL4
expression pattern. Yellow arrows in C show the TPN1 neurons and gray arrows the
VNC-SP neurons. Green, anti-GFP; magenta, anti-RFP.

F) Box plots of total sleep change in % for vehicle-fed and retinal-fed *23E10-GAL4/+*
and *UAS-CsChrimson/+* parental controls and *23E10-GAL4>UAS-CsChrimson* female
flies stimulated by 627nm LEDs. The bottom and top of each box represents the first
and third quartile, and the horizontal line dividing the box is the median. The whiskers
represent the 10th and 90th percentiles. Two-way ANOVA followed by Sidak's multiple
comparisons revealed that retinal-fed *23E10-GAL4>UAS-CsChrimson* flies significantly
increase sleep when stimulated with 627nm LEDs. Both parental controls show no
difference. ****P<0.0001, n.s= not significant. n=24-42 flies per genotype and condition.

G) Box plots of total sleep change in % for vehicle-fed and retinal-fed *23E10-LexA/+*
and *LexAop2-CsChrimson/+* parental controls and *23E10-LexA>LexAop2-CsChrimson*
female flies stimulated by 627nm LEDs. The bottom and top of each box represents the
first and third quartile, and the horizontal line dividing the box is the median. The
whiskers represent the 10th and 90th percentiles. A two-way ANOVA followed by Sidak's
multiple comparisons revealed that no differences in sleep following stimulation. n.s=
not significant. n=23-31 flies per genotype and condition.

Figure 8: The 23E10-GAL4 contains two groups of sleep regulating neurons.

A) Representative confocal stacks of a female *23E10-GAL4>UAS-GFP* fly showing dFB neurons and VNC-SP neurons.

B) Box plots of total sleep recovered in % during the first 24h following 12h of sleep deprivation at night for *23E10-GAL4/+* and *UAS-Kir2.1/+* parental controls and *23E10-GAL4>UAS-Kir2.1* female flies. A one-way ANOVA followed by Tukey's multiple comparisons demonstrate that both parental controls have a bigger homeostatic sleep rebound than *23E10-GAL4>UAS-Kir2.1* female flies. $*P<0.05$, n.s= not significant. n=47-62 flies per genotype.

C) Box plots of total sleep recovered in % during the first 24h following 12h of sleep deprivation at night for *Empty-AD; 23E10-DBD>UAS-Kir2.1* and *VT013602-AD; 23E10-DBD>UAS-Kir2.1* female flies. A two-tailed unpaired t test revealed that there is no difference in homeostatic sleep rebound. n.s= not significant. n=24-29 flies per genotype.

D) Box plots of total sleep recovered in % during the first 24h following 12h of sleep deprivation at night for *Empty-AD; 23E10-DBD>UAS-Kir2.1* and *FBS4>UAS-Kir2.1* female flies. A two-tailed unpaired t test revealed that *FBS4>UAS-Kir2.1* flies have a significantly reduced homeostatic sleep rebound compared to controls. $**P<0.01$. n=15-28 flies per genotype.

Table Legends

Table 1: Quantification of the number of cells labeled by the 23E10-Gal4 driver.

Average number ± SEM and range of dFB neurons, all brain neurons and VNC neurons labeled in *23E10-GAL4>UAS-GFP* female flies.

Table 2: Description of FBS lines.

Identification of the AD construct combined with 23E10-DBD in each FBS line. Average

number ± SEM and range of dFB neurons labeled by each line. Bowtie (Y/N) indicates

whether "bowtie" processes are seen in the brain, Y= yes, N= no. Range of additional

(non-dFB) neurons labeled in the brain of each line. Average number ± SEM of VNC

metathoracic cells labeled by each FBS line. TPN1 (Y/N) indicates whether TPN1

neurons are present in the expression pattern, Y= yes, N= no. Range of additional cells

in the VNC include any VNC neurons that is not TPN1 or "bowtie" like (VNC-SP).

1374

Video Legends

Video 1: Brain of a 23E10-GAL4>UAS-GFP female.

Video 2: VNC of a 23E10-GAL4>UAS-GFP female.

Video 3: Brain of a FBS5>UAS-GFP female.

Video 4: VNC of a FBS5>UAS-GFP female.

Video 5: Brain of a VT013602AD-23E10-DBD (VNC-SP)>UAS-GFP female.

Video 6: VNC of a VT013602AD-23E10-DBD (VNC-SP)>UAS-GFP female.

1382

Supplemental Figure Legends

Figure 2-figure supplement 1: Additional sleep profiles and confocal images of
FBS lines.

A-R) Sleep profile (left) and representative confocal stacks (right) for brain and VNC of

female *FBS>UAS-TrpA1; UAS-GFP* for each FBS line not presented in Figure 2.

Figure 2-figure supplement 2: Additional sleep parameters for FBS female flies
thermogenetically activated.

A) Box plots of daytime sleep bout duration in minutes for flies presented in Figure 2B.

The bottom and top of each box represents the first and third quartile, and the horizontal

line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test revealed that 9 FBS line show a significant increase in daytime sleep bout duration between 22°C and 31°C. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, n=30-51 flies per genotype.

B) Box plots of nighttime sleep bout duration in minutes for flies presented in Figure 2B. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test revealed that control and most FBS line show a significant decrease in nighttime sleep bout duration between 22°C and 31°C. Only 6 sleep-promoting FBS lines show no difference between 22°C and 31°C. Dunnett's multiple comparisons reveal that for 5 of them, nighttime sleep bout duration at 31°C is significantly increased compared with Empty control flies (# on figure). *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, n.s= not significant. n=30-51 flies per genotype.

Figure 2-figure supplement 3: Thermogenetic activation in male flies.

A) Box plots of total sleep change in % ((total sleep on day3-total sleep on day2/total sleep on day2) X 100) for male control (Empty-AD; 23E10-DBD) and 22 FBS lines expressing UAS-TrpA1; UAS-GFP. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. The gray rectangle spanning the horizontal axis indicates the interquartile range of the control. Kruskal-Wallis ANOVA followed by Dunn's multiple comparisons revealed that 9 FBS lines increase sleep significantly more than control flies when transferred to 31°C. *P<0.05, **P<0.01, ****P<0.0001, n=26-46 flies per genotype.

B) Box plots of locomotor activity counts per minute awake for flies presented in A. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles.

Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test found that for 4 sleep-promoting FBS lines (FBS5, FBS45, FBS53 and FBS68) locomotor activity per awake time is increased while no differences are seen for the other 5 sleep-promoting lines between 22°C and 31°C. *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$, n=26-46 flies per genotype.

C) Box plots of daytime sleep bout duration for flies presented in A. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test found that for 7 out of the 9 sleep-promoting FBS lines, daytime sleep bout duration is significantly increased at 31°C compared with 22°C. *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$, n=26-46 flies per genotype.

D) Box plots of nighttime sleep bout duration for flies presented in A. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons revealed that control and most FBS line show a significant decrease in nighttime sleep bout duration between 22°C and 31°C. 5 sleep-promoting FBS lines show no difference between 22°C and 31°C (FBS5, FBS42, FBS53, FBS81 and FBS84) while FBS45 and FBS68 show an increase in nighttime sleep bout duration at 31°C. Dunnett's multiple comparisons reveal that for FBS5, FBS42, FBS45, FBS53, FBS68,FBS81 and FBS84, nighttime sleep bout duration at 31°C is significantly increased compared with Empty control flies (# on figure). **$P<0.01$, ***$P<0.001$, ****$P<0.0001$, n=26-46 flies per genotype.

Figure 3-figure supplement 1: Vehicle-fed sleep data for females in optogenetic experiments.

A) Box plots of locomotor activity counts per minute awake for vehicle-fed flies presented in Figure 3B. The bottom and top of each box represents the first and third

quartile, and the horizontal line dividing the box is the median. The whiskers represent

the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's

multiple comparisons test found that all lines except FBS42 show no difference in

locomotor activity per awake time when the flies are stimulated with 627nm LEDs.

***P<0.001. n=20-39 flies per genotype.

B) Box plots of daytime sleep bout duration for vehicle-fed flies presented in Figure 3B.

The bottom and top of each box represents the first and third quartile, and the horizontal

line dividing the box is the median. The whiskers represent the 10th and 90th percentiles.

Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test

found that most vehicle-fed sleep-promoting lines show no difference in daytime sleep

bout duration when the flies are stimulated with 627nm LEDs. *P<0.05, **P<0.01,

***P<0.001. n=20-39 flies per genotype.

C) Box plots of nighttime sleep bout duration for vehicle-fed flies presented in Figure 3B.

The bottom and top of each box represents the first and third quartile, and the horizontal

line dividing the box is the median. The whiskers represent the 10th and 90th percentiles.

Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test

show no difference in nighttime sleep bout duration when vehicle-fed flies are stimulated

with 627nm LEDs. n=20-39 flies per genotype.

1468

Figure 3-figure supplement 2: Optogenetic activation in males.

A) Box plots of total sleep change in % ((total sleep on day3-total sleep on day2/total

sleep on day2) X 100) for control (Empty) and 22 experimental vehicle-fed and retinal-

fed male flies expressing CsChrimson upon 627nm LED stimulation. The bottom and

top of each box represents the first and third quartile, and the horizontal line dividing the

box is the median. The whiskers represent the 10th and 90th percentiles. Two-way

ANOVA followed by Sidak's multiple comparisons revealed that 9 retinal-fed FBS lines

increase sleep significantly when stimulated with 627nm LEDs when compared with

vehicle-fed flies. *P<0.05, **P<0.01, ****P<0.0001. n=20-44 flies per genotype and

condition.

B) Box plots of locomotor activity counts per minute awake for retinal-fed flies presented in A. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test found that for most sleep-promoting FBS lines, locomotor activity per awake time is not affected when the flies are stimulated with 627nm LEDs while it is increased in *FBS72>UAS-CsChrimson* flies. *P<0.05, n=24-44 flies per genotype.

C) Box plots of daytime sleep bout duration (in minutes) for retinal-fed flies presented in A. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons indicate that daytime sleep bout duration is increased in 12 FBS lines expressing CsChrimson when stimulated with 627nm LEDs. *P<0.05, ***P<0.001, ****P<0.0001, n=24-44 flies per genotype.

D) Box plots of nighttime sleep bout duration (in minutes) for retinal-fed flies presented in A. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons indicate that nighttime sleep bout duration is increased in 7 FBS lines expressing CsChrimson when stimulated with 627nm LEDs. *P<0.05, **P<0.01, ****P<0.0001, n=24-44 flies per genotype.

Figure 3-figure supplement 3: Vehicle-fed sleep data for males in optogenetic experiments.

A) Box plots of locomotor activity counts per minute awake for vehicle-fed flies presented in Figure 3-figure supplement 2A. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test found no difference in locomotor activity

per awake time when the flies are stimulated with 627nm LEDs. n=20-40 flies per genotype.

B) Box plots of daytime sleep bout duration for vehicle-fed flies presented in Figure 3-figure supplement 2A. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test found that most vehicle-fed sleep-promoting lines show no difference in daytime sleep bout duration when the flies are stimulated with 627nm LEDs. $*P<0.05$, $**P<0.01$, $***P<0.001$. n=20-40 flies per genotype.

C) Box plots of nighttime sleep bout duration for vehicle-fed flies presented in Figure 3-figure supplement 2A. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test show no difference in nighttime sleep bout duration when vehicle-fed flies are stimulated with 627nm LEDs. n=20-40 flies per genotype.

Figure 4-figure supplement 1: Vehicle-fed sleep data for females in optogenetic activation of lines containing only VNC neurons.

A) Box plots of locomotor activity counts per minute awake for vehicle-fed flies presented in Figure 4K. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test found that locomotor activity per awake time is reduced in vehicle-fed *VT020742-AD; 23E10-DBD>UAS-CsChrimson* and *VT013602-AD; 23E10-DBD>UAS-CsChrimson* flies that are stimulated with 627nm LEDs. $**P<0.01$, $***P<0.001$. n.s.= not significant. n=13-32 flies per genotype.

B) Box plots of daytime sleep bout duration for vehicle-fed flies presented in Figure 4K. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles.

Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test found no difference in daytime sleep bout duration when the flies are stimulated with 627nm LEDs. n.s= not significant. n=13-32 flies per genotype.

C) Box plots of nighttime sleep bout duration for vehicle-fed flies presented in Figure 4K. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test show that nighttime sleep bout duration is not different when vehicle-fed flies are stimulated with 627nm LEDs. n.s.= not significant. n=13-32 flies per genotype.

Figure 4-figure supplement 2: Optogenetic activation of lines containing only VNC neurons. Male data.

A) Box plots of total sleep change in % for control (*Empty-AD; 23E10-DBD>UAS-CsChrimson*), *VT020742-AD; 23E10-DBD>UAS-CsChrimson*, *30A08-AD; 23E10-DBD>UAS-CsChrimson* and *VT013602-AD; 23E10-DBD>UAS-CsChrimson* vehicle-fed and retinal-fed male flies upon 627nm LED stimulation. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way ANOVA followed by Sidak's multiple comparisons revealed that retinal-fed *VT020742-AD; 23E10-DBD>UAS-CsChrimson* and *VT013602-AD; 23E10-DBD>UAS-CsChrimson* flies increase sleep significantly when stimulated with 627nm LEDs when compared with vehicle-fed flies. ****P<0.0001, n.s.= not significant. n=18-40 flies per genotype and condition.

B) Box plots of locomotor activity counts per minute awake for retinal-fed flies presented in A. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test show that locomotor activity per awake time is increased in *VT020742-AD; 23E10-DBD>UAS-CsChrimson* flies and is not affected in the other

genotypes when the flies are stimulated with 627nm LEDs. **P<0.05, n.s.= not
significant. n=21-40 flies per genotype.

C) Box plots of daytime sleep bout duration (in minutes) for retinal-fed flies presented in
A. The bottom and top of each box represents the first and third quartile, and the
horizontal line dividing the box is the median. The whiskers represent the 10th and
90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple
comparisons indicates that daytime sleep bout duration is increased in retinal-fed
VT020742-AD; 23E10-DBD>UAS-CsChrimson and *VT013602-AD; 23E10-DBD>UAS-*
CsChrimson flies when stimulated with 627nm LEDs. ****P<0.0001, n.s.= not
significant. n=21-40 flies per genotype.

D) Box plots of nighttime sleep bout duration (in minutes) for retinal-fed flies presented
in A. The bottom and top of each box represents the first and third quartile, and the
horizontal line dividing the box is the median. The whiskers represent the 10th and
90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple
comparisons indicates that nighttime sleep bout duration is increased in retinal-fed
VT020742-AD; 23E10-DBD>UAS-CsChrimson and *VT013602-AD; 23E10-DBD>UAS-*
CsChrimson flies when stimulated with 627nm LEDs. ****P<0.0001, n.s.= not
significant. n=21-40 flies per genotype.

1584

Figure 4-figure supplement 3: Vehicle-fed sleep data for males in optogenetic
activation of lines containing only VNC neurons.

A) Box plots of locomotor activity counts per minute awake for vehicle-fed flies
presented in Figure 4-figure supplement 2A. The bottom and top of each box represents
the first and third quartile, and the horizontal line dividing the box is the median. The
whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA
followed by Sidak's multiple comparisons test found that locomotor activity per awake
time is reduced in vehicle-fed *VT020742-AD; 23E10-DBD>UAS-CsChrimson* and
VT013602-AD; 23E10-DBD>UAS-CsChrimson flies that are stimulated with 627nm
LEDs. **P<0.01, n.s.= not significant. n=18-32 flies per genotype.

B) Box plots of daytime sleep bout duration for vehicle-fed flies presented in Figure 4-figure supplement 2A. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test found no difference in daytime sleep bout duration when the flies are stimulated with 627nm LEDs. n.s= not significant. n=18-32 flies per genotype.

C) Box plots of nighttime sleep bout duration for vehicle-fed flies presented in Figure 4-figure supplement 2A. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. Two-way repeated measures ANOVA followed by Sidak's multiple comparisons test show that nighttime sleep bout duration is not increased when vehicle-fed flies are stimulated with 627nm LEDs. n.s= not significant. n=18-32 flies per genotype.

Figure 4-figure supplement 4: Additional data for TPN1 and VNC-SP neurons.

A) Representative confocal stacks of a female *30A08-LexA>LexAop2-GFP; VT020742-AD; 23E10-DBD>UAS-RFP* fly showing the brain, the VNC and a magnified view of the cell bodies in the metathoracic ganglion. Yellow arrows show the TPN1 neurons and their processes in the VNC and in the brain and gray arrows the VNC-SP neurons and their processes. TPN1 neurons are labeled by 30A08-LexA and VT020742-AD; 23E10-DBD while VNC-SP neurons are only present in VT020742-AD; 23E10-DBD. Green, anti-GFP; magenta, anti-RFP.

B) Representative confocal stacks of a female *30A08-LexA>LexAop2-KZip+; VT020742-AD; 23E10-DBD>UAS-GFP* fly showing the brain and the VNC. The KZip+ repressor effectively remove expression in the TPN1 neurons leaving only the VNC-SP neurons. Green, anti-GFP; magenta, anti-nc82 (neuropile marker).

C) Box plots of total sleep change in % for *Empty-AD; 23E10-DBD>UAS-CsChrimson*, *VT020742-AD; 23E10-DBD>UAS-CsChrimson*, *Empty-AD; 23E10-DBD>UAS-CsChrimson; 30A08-LexA> LexAop2-KZip+* and *VT020742-AD; 23E10-DBD>UAS-*

CsChrimson; 30A08-LexA> LexAop2-KZip⁺ vehicle-fed and retinal-fed female flies upon 627nm LED stimulation. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10^{th} and 90^{th} percentiles. Two-way ANOVA followed by Sidak's multiple comparisons revealed that retinal-fed *VT020742-AD; 23E10-DBD>UAS-CsChrimson* and *VT020742-AD; 23E10-DBD>UAS-CsChrimson; 30A08-LexA> LexAop2-KZip⁺* flies increase sleep significantly when stimulated with 627nm LEDs when compared with vehicle-fed flies. Tukey's multiple comparisons demonstrate that there is no difference in total sleep change between retinal-fed *VT020742-AD; 23E10-DBD>UAS-CsChrimson* and *VT020742-AD; 23E10-DBD>UAS-CsChrimson; 30A08-LexA> LexAop2-KZip⁺* flies. **P<0.01, ****P<0.0001, n.s.= not significant. n=17-31 flies per genotype and condition.

D) Representative confocal stacks of a female *VT020742-AD/ VT013602-AD; 23E10-DBD>UAS-GFP* fly showing the brain and the VNC. This line labels the two TPN1 cells and the two VNC-SP neurons. Green, anti-GFP; magenta, anti-nc82 (neuropile marker).

E) Representative confocal stacks of a female *VT013602-AD; VT020742-DBD>UAS-GFP* fly showing the brain and the VNC. This line labels the two VNC-SP neurons and their processes in the brain (gray arrows). Green, anti-GFP; magenta, anti-nc82 (neuropile marker).

F) Sleep profile of control female (*Empty-AD; Empty-DBD>UAS-TrpA1)* at 22°C (blue line) and 31°C (red line).

G) Sleep profile of *VT013602-AD; VT020742-DBD>UAS-TrpA1* female flies at 22°C (blue line) and 31°C (red line).

H) Box plots of total sleep change in % for flies presented in F and G. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10^{th} and 90^{th} percentiles. A two-tailed unpaired t test revealed that activating VT013602-AD; VT020742-DBD neurons significantly increases sleep compared with controls. ****P<0.0001. n=13-14 flies per genotype.

Figure 4-figure supplement 5: Arousal threshold for VNC-SP>UAS-Cs Chrimson flies

Arousal threshold in vehicle-fed and retinal-fed *VT013602-AD; 23E10-DBD>UAS-CsChrimson* female flies. Percentage of flies awakened by a stimulus of increasing strength (1, 2 and 4 downward movements in the SNAP apparatus) with and without 627nm LEDs stimulation. Two-way ANOVA followed by Sidak's multiple comparisons indicates that in retinal-fed flies activation of VT013602-AD; 23E10-DBD neurons reduce the responsiveness to the 1 and 2 stimulus strength when compared with non-activated flies. No difference in responsiveness is seen at the strongest stimulus (4). Two-way ANOVA followed by Sidak's multiple comparisons indicates that in vehicle-fed flies no difference in responsiveness is seen between LED stimulated and non-stimulated flies. ****$P<0.0001$, n.s.= not significant. n=16 flies per genotype and condition.

Figure 6-figure supplement 1: Additional data demonstrating that VNC-SP neurons are cholinergic.

A) Representative confocal stacks of a female *ChAT-LexA>LexAop2-GFP; VT013602-AD; 23E10-DBD (VNC-SP)>UAS-RFP* focusing on the cell bodies in the metathoracic ganglion of the VNC. Gray arrows show the VNC-SP neurons that are co-labelled by the ChAT-LexA driver. Green, anti-GFP; magenta, anti-RFP.

B) Representative confocal stacks of a female *ChAT-LexA>LexAop2-GFP; 23E10-GAL4>UAS-RFP.* Top panels, focusing on the cell bodies in the metathoracic ganglion of the VNC. Gray arrows show the VNC-SP neurons that are co-labelled by the ChAT-LexA driver. Bottom panels, focusing on dFB neurons. Red arrows show 23E10-GAL4 dFB neurons that are also labeled by ChAT-LexA. Green, anti-GFP; magenta, anti-RFP.

C) Quantification of ChAT knockdown efficiency by RNAi. qPCR was performed on the body and eyes of control flies (*GMR-GAL4>+*) and flies expressing ChAT RNAi (line 60028) in the eyes driven by GMR-GAL4 (*GMR-GAL4>ChAT RNAi*). Expression levels were normalized to control levels. Two-way ANOVA followed by Sidak's multiple

comparisons revealed that in *GMR-GAL4>ChAT RNAi* flies ChAT levels are significantly reduced in the eyes but not in the body. **P<0.01. n.s.= not significant. n=3-4 replicates per genotype.

D) ChAT immunostaining in *VNC-SP>UAS-GFP* (top panels) and *VNC-SP>UAS-GFP; UAS-ChAT RNAi* flies (bottom panels). Gray arrows point to neurons positive for GFP and ChAT. Green, anti-GFP; magenta, anti-ChAT.

E) Quantification of data presented in D. A one-tailed unpaired t test revealed that expressing ChAT RNAi in VNC-SP neurons significantly reduce ChAT levels as measured with ChAT antibody staining. *P<0.05. n=14-17 VNC analyzed per genotype.

F) Box plots of total sleep time (in minutes) for control (*Empty-AD; 23E10-DBD>ChAT 60028*) and *VNC-SP>ChAT 60028* male flies. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. A two-tailed Mann-Whitney U test revealed that *VNC-SP> ChAT^{RNAi}* male flies sleep significantly less than controls. ****P<0.0001. n=24-30 flies per genotype.

G) Box plots of daytime sleep bout duration (in minutes) for control (*Empty-AD; 23E10-DBD>ChAT 60028*) and *VNC-SP>ChAT 60028* male flies. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. A two-tailed Mann-Whitney U test revealed that *VNC-SP> ChAT^{RNAi}* male flies daytime sleep bout duration is significantly reduced compared with controls. ***P<0.001. n=24-30 flies per genotype.

H) Box plots of nighttime sleep bout duration (in minutes) for control (*Empty-AD; 23E10-DBD>ChAT 60028*) and *VNC-SP>ChAT 60028* male flies. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10th and 90th percentiles. A two-tailed Mann-Whitney U test revealed that *VNC-SP> ChAT^{RNAi}* male flies nighttime sleep bout duration is significantly reduced compared with controls. ***P<0.001. n=24-30 flies per genotype.

I) Box plots of locomotor activity counts per minute awake for control (*Empty-AD; 23E10-DBD>ChAT 60028*) and *VNC-SP>ChAT 60028* male flies. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10^{th} and 90^{th} percentiles. A two-tailed unpaired t test revealed that there is no difference in waking activity between *VNC-SP> ChAT^{RNAi}* male flies and controls. n.s= not significant. n=24-30 flies per genotype.

J) Box plots of total sleep change in % for *VT013602-AD; 23E10-DBD>UAS-CsChrimson; 2x LexAop2KZip^{+}* (control, no LexA driver) and for *VT013602-AD; 23E10-DBD>UAS-CsChrimson; ChAT-LexA>2x LexAop2KZip^{+}* vehicle-fed and retinal-fed female flies upon 627nm LED stimulation. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10^{th} and 90^{th} percentiles. Two-way ANOVA followed by Sidak's multiple comparisons revealed that retinal-fed *VT013602-AD; 23E10-DBD>UAS-CsChrimson; ChAT-LexA>2x LexAop2KZip^{+}* flies do not increase sleep compared with vehicle-fed flies when stimulated with 627nm LEDs. Control flies increase sleep significantly. ***P<0.001, n.s.= not significant. n=11-16 flies per genotype and condition.

K) Representative confocal stacks of *VT013602-AD; 23E10-DBD>UAS-GFP* (left) and *VT013602-AD; 23E10-DBD>UAS-GFP; ChAT-LexA>LexAop2KZip^{+}* (right) female flies. Expression of GFP in VNC-SP neurons is greatly reduced by the expression of the KZip^{+} repressor. Gray arrows show VNC-SP neurons. Green, anti-GFP; magenta, anti-RFP.

L) Box plots of total sleep change in % for *Empty-AD; 23E10-DBD>UAS-CsChrimson; ChAT-LexA> LexAop2KZip^{+}* and for *FBS45>UAS-CsChrimson; ChAT-LexA> LexAop2KZip^{+}* vehicle-fed and retinal-fed female flies upon 627nm LED stimulation. The bottom and top of each box represents the first and third quartile, and the horizontal line dividing the box is the median. The whiskers represent the 10^{th} and 90^{th} percentiles. Two-way ANOVA followed by Sidak's multiple comparisons revealed no differences in sleep when flies are stimulated with 627nm LEDs. n.s.= not significant. n=22-29 flies per genotype and condition.

M) Representative confocal stacks of *FBS45>UAS-GFP* (left) and *FBS45>UAS-GFP;*

ChAT-LexA>LexAop2KZip⁺ (right) female flies. Expression of GFP in VNC-SP neurons

is completely abolished by the expression of the KZip⁺ repressor. Gray arrows show

VNC-SP neurons. In addition, the KZip⁺ repressor also consistently remove expression

in 4-6 dFB neurons. On average 19.33 ± 0.62 dFB neurons are labeled in *FBS45>UAS-*

GFP; ChAT-LexA>LexAop2KZip⁺ while it is more than 25 without the KZip⁺ repressor

(Table 2). Green, anti-GFP; magenta, anti-RFP.

1746

Figure 6-figure supplement 2: VNC-SP neurons are well positioned to integrate

sensory inputs in the VNC and send this information to the brain.

Representative confocal stacks of a female *VT013602-AD; 23E10-DBD> UAS-*

5xDenMark-V5; UAS-syt.eGFP. Top panels, focusing on the brain "bowtie" processes.

Middle panels, the VNC and bottom panels, a magnified view of the cell bodies area.

Green, anti-GFP; magenta, anti-V5.

1753

1754

1755

Figure 1

A

	22°				31°		22°	
	L	D	L	D	L	D	L	D
Day		1		2		3		4

Total Sleep Change (%)=

$$\frac{\text{Total Sleep on Day3- Day2}}{\text{Total Sleep on Day2}} \times 100$$

B *23E10-GAL4/+*

C *UAS-TrpA1/+*

D *23E10-GAL4>UAS-TrpA1*

E

F

23E10-GAL4>UAS-mCD8GFP

G Brain — 100 μm

H Ventral Nerve Cord — 50 μm

Figure 2

Figure 3

Figure 4

Chronic silencing-Kir2.1

Females

Males

Acute silencing-Shibire[ts1]

Figure 6

VT013602-AD;23E10-DBD>UAS-GFP

Figure 7

Figure 8

Table 1

	Average (n)	Range
dFB neurons	25.68 ± 0.50 (19)	23-30
All brain neurons	51.79 ± 1.05 (19)	44-62
VNC neurons	17.56 ± 0.99 (9)	12-22

Table 2

FBS line	AD component	Average whole brain dFB neurons ± SEM (n)	Range dFB neurons	Bowtie (Y/N)	Range of additional cells in brain	Average VNC cells in metathoracic ganglion ± SEM (n)	TPN1 (Y/N)	Range of additional cells in VNC
FBS 1	12D12-AD	4.64 ± 0.40 (n=14)	2-7	Y	0	2.00 ± 0.32 (n=11)	N	0
FBS 4	84C10-AD	20.80 ± 0.35 (n=15)	18-23	N	0	0.00 ± 0.50 (n=15)	N	0
FBS 5	23E12-AD	0 ± 0 (n=26)	0	Y	5-10	3.89 ± 0.11 (n=26)	Y	0
FBS 6	42H01-AD	3.85 ± 0.52 (n=13)	1-4	N	0	0.00 ± 0.00 (n=8)	N	0
FBS 25	22A12-AD	0 ± 0 (n=5)	0	N	0	0.00 ± 0.00 (n=5)	N	0
FBS 28	54G07-AD	13.08 ± 0.63 (n=12)	10-18	Y	0-1	3.09 ± 0.17 (n=22)	N	0
FBS 33	49C04-AD	26.50 ± 1.15 (n=10)	23-31	Y	0	3.17 ± 0.32 (n=12)	Y	0
FBS 35	65C03-AD	5.00 ± 1.06 (n=6)	2-8	Y	1-2	3.60 ± 0.24 (n=5)	Y	0
FBS 41	VT058545-AD	11.50 ± 0.43 (n=10)	10-14	N	0-2	3.00 ± 0.41 (n=4)	Y	0
FBS 42	12G09-AD	21.56 ± 0.80 (=10)	18-25	Y	2-4	3.00 ± 0.37 (n=6)	Y	0-2
FBS 45	VT050238-AD	25.11 ± 1.41 (n=9)	21-32	Y	0	2.40 ± 0.16 (n=15)	N	0
FBS 53	VT043400-AD	23.86 ± 0.51 (n=7)	22-26	Y	0	2.91 ± 0.31 (n=11)	N	0-1
FBS 57	VT025720-AD	7.40 ± 0.68 (n=5)	5-9	Y	0	3.00 ± 0.31 (n=7)	Y	0
FBS 58	VT064566-AD	21.50 ± 1.28 (n=6)	18-27	Y	0-2	3.10 ± 0.28 (n=10)	Y	0-1
FBS 60	VT023818-AD	24.56 ± 0.50 (n=9)	22-27	Y	0	3.27 ± 0.38 (n=11)	Y	0-2
FBS 64	VT027956-AD	15.00 ± 2.12 (n=7)	11-19	Y	0	3.40 ± 0.49 (n=10)	Y	0
FBS 68	VT023823-AD	4.45 ± 0.34 (n=11)	3-6	Y	0	4.08 ± 0.49 (n=13)	Y	0-4
FBS 70	VT034811-AD	4.86 ± 0.40 (n=7)	3-6	Y	0-7	3.90 ± 0.38 (n=10)	Y	0-4
FBS 72	VT026953-AD	24.40 ± 1.50 (n=7)	20-28	Y	0-1	3.27 ± 0.25 (n=10)	Y	0
FBS 81	74G10-AD	22.17 ± 0.70 (n=6)	19-24	Y	0	2.80 ± 0.58 (n=5)	N	0-2
FBS 84	VT007754-AD	23.80 ± 0.58 (n=8)	22-25	Y	0-2	3.36 ± 0.31 (n=12)	Y	0-4
FBS 87	VT045793-AD	18.55 ± 0.53 (n=13)	17-22	Y	0-2	3.62 ± 0.29 (n=13)	Y	0-6

Figure 2-S1

Figure 2-S2

Figure 2-S3

Figure 3-S1

Figure 3-S2

Figure 3-S3

A

B

C

Figure 4-S2

Figure 4-S4

A   *VT020742-AD; 23E10-DBD>UAS-mCD8RFP; 30A08-LexA> LexAop2-mCD8GFP*

B   *VT020742-AD; 23E10-DBD>UAS-GFP; 30A08-LexA> LexAop2-KZip+*

C

D   *VT020742-AD; VT013602-AD; 23E10-DBD>UAS-GFP*

E   *VT013602-AD; VT020742-DBD; >UAS-GFP*

F   *Empty> UAS-TrpA1*

G   *VT013602-AD; VT020742-DBD> UAS-TrpA1*

H

Figure 4-S5

Figure 6-S2