Genome-wide identification, characterization, and comparative analysis of NLR resistance genes in *Coffea spp.*

Mariana de Lima Santos [1*], Mário Lúcio Vilela de Resende [1], Gabriel Sérgio Costa Alves[2],
Jose Carlos Huguet-Tapia[3], Márcio Fernando Ribeiro de Resende Júnior[4] [†], Jeremy Todd
Brawner[3* †]

[1]Laboratório de Fisiologia do Parasitismo, Faculdade de Ciências Agrárias, Departamento de
Fitopatologia, Universidade Federal de Lavras, Lavras, Minas Gerais, Brazil

[2]Laboratório de Processos Biológicos e Produtos Biotecnológicos, Instituto de Ciências
Biológicas, Departamento de Biologia Celular, Universidade de Brasília, Brasília, Distrito
Federal, Brazil

[3]Institute of Food and Agricultural Sciences, Department of Plant Pathology, University of
Florida, Gainesville, Florida, FL, United States

[4] Institute of Food and Agricultural Sciences, Horticultural Sciences Department, University of
Florida, Gainesville, Florida, FL, United States

* Correspondence:
Corresponding Authors
santos-ml@outlook.com
jeremybrawner@ufl.edu
17
†These authors have contributed equally to this work and share last authorship

Keywords: resistance genes, Coffea, nucleotide-binding site leucine-rich repeat, Genome-
wide, NLR-annotator.

Abstract

The largest family of disease resistance genes in plants are nucleotide-binding site leucine-rich
repeat genes (NLRs). The products of these genes are responsible for recognizing avirulence
proteins (Avr) of phytopathogens and triggering specific defense responses. Identifying NLRs in
plant genomes with standard gene annotation software is challenging due to their multidomain
nature, sequence diversity, and clustered genomic distribution. We present the results of a
genome-wide scan and comparative analysis of NLR loci in three coffee species (*Coffea
canephora*, *Coffea eugenioides* and their interspecific hybrid *Coffea arabica*). A total of 1311
non-redundant NLR loci were identified in *C. arabica*, 927 in *C. canephora*, and 1079 in *C.
eugenioides*, of which 809, 562, and 695 are complete loci, respectively. The NLR-annotator tool
used in this study showed extremely high sensitivities and specificities (over 99%) and increased
the detection of putative NLRs in the reference coffee genomes. The NLRs loci in coffee are
distributed among all chromosomes and are organized mostly in clusters. The *C. arabica* genome
presented a smaller number of NLR loci when compared to the sum of the parental genomes (*C.
canephora*, and *C. eugenioides*). There are orthologous NLRs (orthogroups) shared between
coffee, tomato, potato, and reference NLRs and those that are shared only among coffee species,
which provides clues about the functionality and evolutionary history of these orthogroups.

Phylogenetic analysis demonstrated orthologous NLRs shared between *C. arabica* and the
parental genomes and those that were possibly lost. The NLR family members in coffee are
subdivided into two main groups: TIR-NLR (TNL) and non-TNL. The non-TNLs seem to
represent a repertoire of resistance genes that are important in coffee. These results will support
functional studies and contribute to a more precise use of these genes for breeding disease-
resistant coffee cultivars.

1   Introduction

Coffee is a globally important agricultural commodity that plays a significant economic role in
producing and consuming countries (Krishnan, 2017). The genus *Coffea* consists of more than
100 botanical species (Davis et al., 2006), however, the most cultivated species are *Coffea*
canephora and *Coffea arabica*. *C. canephora* is diploid (2n = 2x = 22 chromosomes) (Denoeud
et al., 2014), while *C. arabica* is a allotetraploid (2n = 4x = 44 chromosomes) (Tran et al., 2018)
originated from natural hybridization between *C. canephora* and *C. eugenidoides* (Lashermes et
al., 1999; Bawin et al., 2020). Among the more than 50 coffee-producing countries, Brazil,
Vietnam, Colombia, and Indonesia are major producers, with Brazil being the largest producer
by volume. Currently, coffee diseases are the main factor affecting productivity (Cerda et al.,
2017). Examples of diseases associated with coffee include cercosporiosis (*Cercospora*
coffeicola), bacterial blight (*Pseudomonas syringae* pv. *Garcae*), anthracnose (*Colletotrichum*
coffeanum), root-knot nematodes (*Meloidogyne spp.*), coffee berry disease – CBD
(*Colletotrichum kahawae*), and coffee leaf rust – CLR (*Hemileia vastatrix*) (Cabral et al., 2016;
Krishnan, 2017). CLR is one of the most devastating diseases found in coffee and is present in
all regions of the world where coffee is grown (McCook and Vandermeer, 2015; Cabral et al.,
2016). Currently, 95% of *C. arabica* varieties cultivated in Brazil are susceptible to CLR due to
the emergence of variants of the pathogen (Cabral et al., 2016). Given the increasing problem of
plant pathogens in coffee production, a greater understanding of the set of receptors regulating
the plant immune system of coffee is needed.

Throughout evolution, plants have developed sophisticated systems to defend themselves from
pathogens. The plant immune system involves both broad-spectrum and specific recognition of
pathogens. Broad-spectrum recognition is related to the detection of pathogen□associated
molecular patterns (PAMP), such as fungal chitin or bacterial flagella, by pattern recognition
receptors (PRR) that are anchored to the plasma membrane and trigger the PAMP-triggered
immunity (PTI) (Boutrot and Zipfel, 2017). Specific recognition, on the other hand, primarily
involves receptors encoded by resistance genes (R genes) that detect the presence of pathogen
effector proteins and trigger effector-triggered immunity (ETI) (Jones and Dangl, 2006). Both
types of recognition occur dynamically and continuously, converging into signaling pathways
that activate essential mechanisms for downstream responses to pathogen recognition (Lu and
Tsuda, 2021; Yuan et al., 2021).

The R genes have been extensively studied in several crops to facilitate their greater use in plant
breeding (Jupe et al., 2013; Wan et al., 2013; Lozano et al., 2015; Inturrisi et al., 2020;
Steuernagel et al., 2020). The protein products of these genes recognize directly or indirectly
effector proteins that are secreted by pathogens (Kourelis and Van Der Hoorn, 2018) and trigger
a series of signaling steps that lead to the hypersensitive response (HR) that activates cell death
and potentially leads to systemic acquired resistance (SAR) (Kachroo and Robin, 2013; Jones et

al., 2016). The largest and most diverse group of R genes found in plants belong to the
nucleotide-binding site leucine-rich repeat family (NLR or NBS-LRR) (Jones et al., 2016). The
proteins encoded by these genes are typically modular, and many have a variable N-terminal
domain-containing Toll/interleucina-1 (TIR) or coiled-coil (CC) receptors. As well, the
nucleotide-binding domain (NB-ARC or NBS) is a canonical feature of NRLs, that is shared
with human apoptotic protease-activating factor-1 (*APAF-1*) and *Caenorhabditis elegans* death-4
(*CED-4*) proteins. A C-terminal region comprising a variable number of leucine-rich repeats
(LRRs) is another common feature in NLR genes (Jones et al., 2016; Shao et al., 2019). The NB-
ARC domain is highly conserved and is involved in the active and inactive state of the NLS
protein (Bonardi et al., 2012; Jones et al., 2016). This domain presents motifs that are
characteristic of the ATPase family, such as p-loop, kinase 2, and the RNBS (Resistance
Nucleotide Binding Site) A, RNBS-C, and RNBS-D motifs (Van Ghelder et al., 2019).
Mutations in specific residues within these motifs may cause the loss of protein function or self-
activation and interfere with the regulation or activation of defense mechanisms. (Monteiro and
Nishimura, 2018; Bezerra-Neto et al., 2020).

The domains described above provide a more refined classification of NLRs proteins, which may
be characterized into two main groups: TNLs (TIR-NLRs) or non-TNL (which include CNLs -
CC-NLRs). The truncation of a single domain, such as LRR (CN or TN), TIR or CC (NL), in
both C and N terminal domains (N), may also be used to classify NLR genes (Monteiro and
Nishimura, 2018). Additionally, atypical or non-canonical integrated domains (IDs) that act as
decoys and play roles in oligomerization or downstream signaling may be present, demonstrating
the structural diversity of this NLR family (Kroj et al., 2016; Wang et al., 2021). The number of
NLRs in plant genomes varies greatly and is often organized in tandem, which facilitates
duplication, contraction, and transposition and provides a reservoir of genetic variation that
allows plant evolutionary dynamics to respond to changes phytopathogen populations (Barragan
and Weigel, 2021). These genes are often under selection pressure, resulting in a large number of
pseudogenes and variable loci content within the same species, among species, and across plant
populations (Schatz et al., 2014; Steuernagel et al., 2015; Sun et al., 2020; Hufford et al., 2021).

The knowledge of how NLRs are distributed throughout the genome and their diversity is of
great interest as it may reveal new sources of resistance that may be used to develop new
cultivars (Monteiro and Nishimura, 2018). The growing number of sequenced plant genomes
facilitates the search for novel NLR and has led to the genome-wide analysis of NLR genes
(Denoeud et al., 2014; Song et al., 2015; Scott et al., 2020; Wang et al., 2021). However, its large
number, frequently clustered genomic distribution, and low expression in uninfected tissues
make cataloging NLR genes challenging and often underestimates the number of NLRs in
genomes (Jupe et al., 2013; Steuernagel et al., 2015, 2020). To mitigate this problem, some
specific gene/loci NLR annotation pipelines have been developed to augment standard gene
annotation software and improve our ability to identify and locate genes belonging to this family.
Some examples of these pipelines are NBSPred (Kushwaha et al., 2016), NLGenomeSweeper
(Toda et al., 2020), and NLR-annotator, a new version of NLR-parser (Steuernagel et al., 2015,
2020). The NLR-Annotator is a tool used to annotate NLR loci that use 20 highly curated motifs
present in NLR proteins and does not depend on the support of transcript data (Jupe et al., 2012;
Steuernagel et al., 2020). Since it was published, this tool has been applied in several studies to
prospect and annotate R genes in recently sequenced genomes (Muliyar et al., 2020; Read et al.,

2020; Scott et al., 2020), to check the completeness of previous annotations (Muliyar et al.,
2020), and for studies of the resistance-related locus (Jost et al., 2020).

The genome of *C. canephora* was published in 2014, which allowed the first genome-wide NLR
study in coffee (Denoeud et al., 2014). In 2018, the *C. arabica* and *C. eugenioides* genomes were
deposited at the NCBI, providing an essential resource for studying the structure and evolution of
NLRs in arabica coffee and the contribution of the genomes that gave rise to this species. For
coffee production to continue advancing in producing regions worldwide, adequate disease
management is of great importance. A range of strategies must be used to control the main
phytosanitary problems associated with coffee production. Using these genomic resources is
essential for informing breeding strategies aimed at developing resistance to disease in coffee.
Given the above, this study aimed to: (i) identify NLR loci in *C. arabica*, *C. canephora,* and *C.*
eugenioides genomes using the NLR-annotator tool and discuss the improvements in annotation
derived from the use of a specific pipeline for NLR genes in coffee, (ii) catalog, classify and
characterize the distribution of NLRs loci in the *coffee spp.* genomes, and (iii) understand the
contribution of *C. canephora* and *C. eugenioides* to the NLR repertoire of *C. arabica*.

2   Materials and Methods

2.1   Coffee Genomic Resources

Three genomes were used in this study. The *C. arabica* (Caturra red - Cara_1.0, GenBank
assembly accession: GCA_003713225.1) and *C. eugenioides* (Ceug_1.0, GenBank assembly
accession: GCA_003713205.1) genomes are available from the NCBI (National Center for
Biotechnology Information) database (https://www.ncbi.nlm.nih.gov/) and the *C. canephora*
genome is available at Coffee genome hub (https://coffee-genome.org/) (Denoeud et al., 2014).
For the three species, the genome files, sets of predicted proteins, predicted genes, and GFF
(General Feature Format) were used.

2.2   Identification of loci NLR in *Coffea* spp. genomes

The identification of NLR loci in *Coffea spp.* was accomplished by the NLR-Annotator,
(Steuernagel et al., 2020) using the default parameters. The tool uses combinations of short
motifs of 15 to 50 amino acids to classify a genomic locus as an NLR. These motifs were defined
using domains of known NLR proteins used as a training set in a study carried out by Jupe et al.
2012 (Supplementary Table 1).

In summary, the pipeline is divided into three steps: 1) dissection of genomic input sequence into
20-kb fragments overlapping by 5 kb; 2) translating each fragment into all six reading frames
and searching for the motifs associated with NLR by NLR-Parser to create an xml-based report
file. The NLR-Parser searches for combinations of doublets or triplets of motifs and records their
nucleotide positions, disregarding motifs that occur randomly. Finally in step 3, the NLR-
Annotator uses the xml file as input, integrates data from all fragments, evaluates positions and
combinations of motifs. In this step, the NB-ARC motifs are used as the principal seed to
annotate NLR loci, generate output files (.gff, .bed - Browser Extensible Data, .txt and file of the
NB-ARC motifs as multiple alignments to complete loci) based on coordinates and orientation
the initial input genomic sequence (Steuernagel et al., 2020).

Each section of the genomic sequence associated with a single NLR is called an 'NLR locus' and
this refers to an NB-ARC domain (or associated motif) followed or not by one or more leucine-
rich repeats (LRRs). From the sets of motifs that are identified, these loci are classified as
complete (containing the P-loop, at least three consecutive NB-ARC motifs, and at least one
LRR), complete (pseudogenes), partial and partial (pseudogenes) (Supplementary Table 1).
Therefore, the NLR-Annotator identifies the NLR loci that are either active genes or
pseudogenes. The number of NLR loci and their classification is described in the output file.txt
(Steuernagel et al., 2020 and https://github.com/steuernb/NLR-Annotator).

2.3　Validation of the NLR-annotator sensitivity and specificity in coffee genomes

To validate the sensitivity and specificity of the NLR-annotator in the coffee genomes, we
initially classified the protein sequences of the three genomes using PfamScan
(https://pfam.xfam.org/) version 1.5 with an e-value of less than 1E- 5 and models from Pfam-A.
Subsequently, proteins that had the NB-ARC domain (PF00931) were filtered, and from this
process, we obtained the ID of the genes corresponding to each protein. With the list of gene
model IDs of the NLR family, it was then possible to filter the GFF files and obtain the positions
of the genes that had already been annotated in each genome.

We identified overlapping intervals to compare the NLR loci detected by NLR-Annotator and
the NLR genes that had already been annotated in the genomes. We used the information from
.gff files from both annotations for an overlay analysis using bedtools intersect (version 2.29.2).
An overlap was only considered if both, the locus, and gene, were on the same strand. This
analysis made it possible to distinguish the loci identified by NLR-annotator that were or were
not overlapping with the gene models from the reference genomes.

For NLR genes already annotated in the genomes and not identified by NLR-annotator, a search
for motifs by NLR-Parser was performed to obtain the xml and txt output (options -c and -o) as
well the detection of conserved domains using the NCBI Conserved Domain Database
(https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi) for nucleotides sequences. Standard
parameters were used for the conserved domains analysis, except for the threshold (E-value),
which was set to 1E-5. The Graphical summary was set to provide a concise view of the results.
To characterize the NLR loci only found by NLR-annotator and to make sure they were
homologous with NLRs already annotated in plants, we aligned the sequences for these loci with
NCBI's non-redundant protein database (nr) (https://blast.ncbi.nlm.nih.gov) using BLASTx
(BLAST - version 2.10.1 with the max_target_seqs option set to 5). For loci that did not have
homology with NLRs proteins, a conserved domain analysis was also performed as previously
described.

The sensitivity of the pipeline was calculated as the ratio of the number of loci identified by
NLR-annotator (including motifs detected by NLR-Parser in the second step of the pipeline) to
the number of NLRs genes already annotated in the genomes. The specificity was calculated as
the ratio of the number of loci identified by NLR-annotator that are related to NLRs genes or
have characteristic domains of that family to the total number of loci identified. Characteristic
domains were defined as domains overlapping with the annotations already described in the
studied genomes, homology with NLR proteins by BLASTx or NB-ARC domains identified
with conserved domains analysis.

2.4   Distribution of NLR loci in Coffee's chromosomes

In order to visualize the distribution of NLR loci on chromosomes of the three analyzed coffee species, the annotation files from NLR annotator (.txt) were used to extract the genomic position and classifications of the loci. The chromosome size information in Mb was obtained from the NCBI (for *C. arabica* and *C. eugenioides*) and Coffee genome hub (for *C. canephora*) and the visualization was created using the R software with the chromoMap package (Anand and Lopez, 2020). ChromoMap, divides the chromosomes as a continuous composition of loci. Each locus, consist of a specific genomic range determined algorithmically based on chromosome length and then the annotations are inserted. The detailed annotation information on each locus NLRs (complete, complete pseudogene, partial and partial pseudogene) is displayed in an HTML file.

2.5   Prediction of genes in the complete loci found only by the NLR-annotator

Gene prediction was performed using the AUGUSTUS program version 3.3.3 (http://bioinf.uni-greifswald.de/augustus/) (Stanke et al., 2006) using gene models from *Solanum lycopersicum* and allowing for the prediction of only complete genes.

2.6   Orthologous groups and Phylogenetic Analyses

The complete loci identified in the coffee genomes by the NLR-annotator, being those loci that overlap with gene models of the reference genomes and loci that were annotated by AUGUSTUS as putative genes were the focus of ortholog and phylogenetic analysis. In order to make a comparison with the set of coffee NLRs, 326 NLR loci identified in tomato (*Solanum licopersicum* - Heinz 1706) by Andolfo et al., 2014, 755 loci identified in potato (*Solanum tuberosum*) by Jupe et al., 2013, 67 NLR reference genes (functionally characterized protein) obtained from The Plant Resistance Genes database - PRGDB (http://prgdb.org/prgdb/, S2 table) (Osuna-Cruz et al., 2018) and the *CED-4* gene from *Caenorhabditis elegans* (outgroup) were also added. All these sequences were classified according to the rules of motifs established by the NLR-annotator and only those considered as complete NLR were used for these analyses. This criterion was used to standardize the methodology for classifying loci as complete or not.

The amino acid sequences of the NB-ARC domain were extracted from the set of complete NLR loci for the 5 species (*C. arabica*, *C. canephora*, *C. eugenioides*, *S. licopersicum* and *S. tuberosum*) along with the reference genes by NLR -annotator (parameter -a). All NB-ARC domain of these complete loci (hereafter called NLRs) were compared with each other using BLASTP, all-by-all (E value <1e-10). The markov clustering algorithm was performed with inflation value of 1.5 and then NLRs in the same cluster were classified as orthologous subgroups by OrthoMCL version 1.4 (standard parameters) (Li et al., 2003). In order to analyze and visualize the number of orthogroups shared between the species and the ones that are unique to a single species, we used the UpSetR package in R (Conway et al., 2017).

The NLRs clustered into single-copy orthogroups (orthogroups that have one copy of each NLR present once in each of the 5 genomes or reference NLRs) by OrthoMCL were used to construct a phylogenetic tree. The sequences were aligned using MAFFT version 6.903 (Katoh et al., 2002), with the --auto parameter to select the best alignment strategy. The tree was inferred using RAxML version 8.2.10 (Stamatakis, 2014) with the PROTGAMMAAUTO model (the JTT model was selected as having the highest likelihood) with 100 bootstrap replicates. A second

247 phylogenetic tree classifying the coffee NLRs was also constructed with the above-mentioned
248 parameters using the entire set of complete NLRs. Coffee NLRs were classified in the tree using
249 the previous classification describing 67 reference NLRs (Supplementary Table 2) and the
250 tomato and potato NLRs (Jupe et al., 2012, 2013; Andolfo et al., 2014). The trees were
251 visualized and edited using the Interactive Tree of Life (iTOL) tool (Letunic and Bork, 2021).

3   Results

3.1   NLRs Identification, validation of the sensitivity and specificity of NLR-annotator in coffee genomes

A total of 1318 loci were identified for *C. arabica*, 932 for *C. canephora*, and 1081 for *C.
eugenioides* (Supplementary Table 3). In each species, we identified some loci that are in the
same position but were separated by the NLR-annotator as two distinct NLRs. We found 7, 5,
and 2 repeated loci in the aforementioned species, respectively (Supplementary Table 3
highlighted in blue). Considering these repeated loci when counting the number and distribution
of the NLRs loci on the chromosomes, the most complete classification was considered. After
the identification of these regions, it was found that there were 1311 non-redundant loci for *C.
arabica* (627 from the *C. canephora* subgenome - CaC, 650 from the *C. eugenioides* subgenome
- CaE and 34 Unassigned - Un), 927 for *C. canephora* (559 mapped on chromosomes and 367
Un) and 1079 for *C. eugenioides* (944 mapped on chromosomes and 135 Un). The number of
complete loci found in each species was 809 (*C. arabica*), 562 (*C. canehora*), and 695
(*C.eugenioides*).

To examine whether there was a consensus between the gene models for NLRs that have
previously been annotated in the genomes and loci identified by NLR-annotator, an overlap
analysis was performed. PfamScan analyses were conducted on the set of predicted proteins and
the subsequent selection of NLR proteins annotated in each genome showed that 1015, 709, and
869 genes encoded proteins (including isoforms) containing the NB-ARC domain in the *C.
arabica*, *C. canephora*, and *C. eugenioides* genomes, respectively (Figure 1, Supplementary
Table 4). The overlap between the genomic positions of these genes and the positions of loci
from NLR-annotator showed that of 1311, 927, and 1079 loci identified by NLR-annotator for *C.
arabica*, *C. canephora*, and *C.eugenioides* respectively, 1013 (99,80%), 687 (96,90%), and 857
(98,62%) overlap with the genes already annotated in the reference genomes. A total of 298, 240
and 222 non overlapping NLRs were found, respectively (Figure 1, Table 1). We also noticed
that there are cases in which more than one NLR loci overlapped with a single NLR gene, and
the opposite was also found in all three genomes. A Venn diagram representing these data is
shown in figure 1 as the intersection and data are highlighted in Supplementary Table 5.

The overlap analysis also made it possible to identify genes annotated in the reference genomes
that did not overlap with any locus from NLR-annotator. To examine these genes, an NLR-parser
analysis, with options -c (file.xml) and -o (file.txt), was performed on this set. Among the genes
not identified by NLR-annotator for *C. arabica* (18), *C. canephora* (25), and *C. eugenioides* (24),
7, 3, and 4, respectively, were below the standard threshold (1E-5) for the MAST based motif
search used by NLR-Parser. Additionally, 9, 17, and 16 genes present motifs that were detectable
using the standard threshold but did not contain at least three consecutive motifs belonging to the
NB-ARC domain or presented as motifs in random order. These loci were therefore not

288 annotated in the third step of the NLR-annotator (Supplementary Table 6). After this analysis, we
289 also identified and confirmed genes that were not found by NLR-annotator. Two genes were not
290 found in *C. arabica* (LOC113737176 and LOC113735982), five genes in *C. canephora*,
291 (Cc02_g12220, Cc03_g10360, Cc07_g18800, Cc00_g21910 and Cc00_g35420) and four genes
292 in *C. eugenioides* (LOC113766771, LOC113766774, LOC113766615 and LOC113777141).
293 Supplementary text 1, supplementary table 6 and supplementary figure 1 present additional
294 details from this analysis. After these analyses, it was possible to verify that the NLR annotator
295 showed a sensitivity of 99.8%, 99.4%, and 99.7% for *C. arabica*, *C. canephora* and *C.
296 eugenioides*, respectively.

297 As stated above, the overlap analysis also made it possible to detect that the NLR-annotator
298 identified loci that were complete, complete (pseudogenes), partial and partial (pseudogenes) that
299 did not overlap with genes already annotated in reference genomes (Figure 1, Table 1). To
300 further investigate these loci and ensure that they were indeed related to genes encoding NLR
301 proteins, a BLASTx analysis was performed. This analysis showed that of the 298, 240, and 222
302 loci in *C. arabica*, *C canephora*, and *C. eugenioides*, only 7, 4, and 6 did not show homology
303 with resistance proteins being found among the five best hits, respectively. (Supplementary
304 Table 7, highlighted in orange).

305 To describe the sequences that did not show homology to NLRs proteins by BLASTx, a
306 conserved domains analysis was performed (Supplementary Figure 2). Many of these loci do not
307 show homology with NLRs proteins because most of the sequence contains domains related to
308 the family of proteins involved in the activity of transposable elements such as ribonuclease H
309 (RNase H) and reverse transcriptases (RTs). However, it was also possible to identify
310 characteristic domains of NLR proteins such as NB-ARC, Toll / interleukin-1 receptor (TIR),
311 RX-CC_like, and Rx_N, suggesting that these loci cannot be considered false positives. Only
312 three loci did not present characteristic domains, Chr11c_nlr_73_Ca, chr0_nlr_300_Cc and
313 Chr8_nlr_67_Ce and all these loci were partial (pseudogenes). These loci were removed from
314 further analysis. From these results, it was possible to verify that the specificity of the NLR-
315 annotator was 99.9% in all three genomes.

3.2 Distribution of NLR loci in the Coffea spp. Genome

317 Considering all detected loci, in *C. canephora*, chromosomes 3 and 11 have the most significant
318 number of identified loci, including complete, complete (pseudogene), partial and partial
319 (pseudogene). For *C. eugenioides*, chromosomes 3 and 11 also contain many NLR and large
320 numbers of loci were also found on chromosomes 5 and 8. The total number of loci found on
321 chromosome 8 is almost the same as the number identified on 11, however, the number of
322 complete NLR on chromosome 8 is higher, with 16 more loci identified. For *C. arabica*,
323 chromosomes 3 and 11 from the *C. canephora* and *C. eugenioides* subgenomes, respectively,
324 also have the largest number of NLR. This what was also found on chromosome 8 from the *C.
325 eugenioides* subgenoma. For *C. arabica*, the *C. eugenoides* subgenome generally has a slightly
326 higher number of NLRs loci as reported above. The number of loci of this subgenome on
327 chromosomes 8 and 11 stand out in comparison to *C. canephora* subgenome, with 34 and 30
328 more loci, respectively. The chromosomes with the fewest loci for the three species are 9 and 10,
329 and chromosome 4 specifically for *C. eugenioides*. The number of loci in unmapped sequences

(Unassigned) for the *C. canephora* reference genome represent 39.7%, which was much higher
than the number of loci found in *C. eugenioides* (12.5%) and *C. arabica* (2,6%) (Figure 2A).

The chromosomal location of these loci in the three species demonstrated that most loci are
organized in clusters and are unevenly distributed across the entire chromosome. In addition,
there are clusters that have the four different types of loci or at least two types, presenting a
stretch of complete, complete (pseudogene), partial and/or partial (pseudogene). Not all loci were
clustered, we also found loci of the four types that were physically isolated in chromosomes
(Figure 2B).

3.3   Gene prediction for complete loci found only by NLR-annotator

Since the NLR-annotator is not a gene predictor but is a tool to annotate loci associated with
NLRs, the gene-finding program AUGUSTUS was used to characterize the loci found only by
NLR-annotator and that were classified as complete (S3 Table highlighted in orange). This
analysis aimed to verify whether these complete loci could be considered potential gene models.
This analysis showed that of the 70 and 67 complete loci for *C. arabica* and *C. canephora*, 64
and 66, are potential gene models, respectively. For *C. eugenioides*, all 71 loci were predicted as
potential genes. The loci that were not identified as potential genes are in red in supplementary
table 3.

3.4   Ortholog Groups and Phylogenetic Analysis

From the ortholog group analysis conducted using OrthoMCL, 803 complete loci of *C. arabica*,
561 of *C. canephora* and 695 of *C. eugenioides* were used. Six and 1 loci of *C. arabica* and *C.*
canephora, respectively, were removed from analysis because they are complete loci that are not
overlapping gene models or were not identified as putative genes by AUGUSTUS analysis.
Additionally, 151 tomato loci (out of 326) and 403 potato loci (out of 755) that were classified as
complete loci by NLR-annotator as well as 67 reference NLRs and *CED-4* were used. Out of a
total of 2681 NLRs, 2038 (76%) were grouped into 593 orthologous groups, hereinafter referred
to as orthogroups (Figure 3, Supplementary Table 8). The number of coffee NLRs present within
these orthogroups were 591, 427, 584, which represents 73.6%, 76.1% and 84% of the total
NLRs found for *C. arabica*, *C. canephora* and *C. eugenioides*, respectively. Two hundred and
seventy-two orthogroups were in single-copies, containing 647 NLRs, of which only 7 are
reference NLRs.

There were 10 orthogroups shared by all species and reference NLRs and 11 were shared only
among species and not the reference NLRs. As expected, the greatest number of orthogroups
were shared among coffee NLRs, 200 orthogroups containing 783 NLRs were shared only
among *C. arabica* (296: 163 CaE, 130 CaC e 3 un), *C. canephora* (215) and *C. eugenioides*
(272), respectively. The comparison between *C. arabica* NLRs with only one of the diploid
species showed that *C. eugenioides* shares a slightly higher number of orthogroups (78) than *C.*
canephora (71) and also of NLRs within these orthogroups (orthogroup Ca/Ce = 87/96 NLRs,
orthogroup Ca/Cc = 86/74 NLRs). When the comparison was only between the two diploid
species, it was observed that 34 orthogroups are shared only between them. The number of
orthogroups shared between NLRs of the same coffee species was 31 in *C. eugenioides*, 24 in *C.*
arabica and 9 in *C. canephora*.

The number of NLR orthogroups shared only between the three coffee species and one solanum
specie was higher among potato (9) than tomato (3) NLRs, however it should be noted that the
number of potato NLR in the analysis was almost 3 times larger than the number of tomato NLR.
Forty-six orthogroups contain NLRs from at least one coffee species and one solanum species.
Fifteen orthogroups were shared between reference NLRs, and at least one coffee and a solanum
species, and grouped 21 reference NLRs (Figure 3, Supplementary Table 8, highlighted in light
blue). Of these, *C. canephora* and/or *C. eugenioides* are present in three orthogroups with
reference genes in which *C. arabica* is absent (ORTHOMCL16: Cc, Soly, Soltu e *Hero*;
ORTHOMCL17: Ce, Soly, Soltu e *Rpiblb1*; ORTHOMCL24: Cc, Ce, Soly, Soltu e *VAT*),
indicating these orthogroups are not present in the hybrid. Four orthogroups were clustered in
only the three coffee species and reference NLRs (ORTHOMCL1, ORTHOMCL19,
ORTHOMCL119 and ORTHOMCL199, Figure 3 and Supplementary Table 8, highlighted in
dark blue), which contained *Lr10*, *MLA1*, *MLA10*, *MLA13*, *Mla12*, *Mla6*, *Pi36*, *Pikm2TS*, *FOM-*
2, *Rdg2a* e *Pm3*. The percentage of orphans (i.e., NLRs not assigned to any ortholog group by
OrthoMCL) among coffee NLRs was highest in *C. arabica* (26.4% -212) followed by *C.*
canephora (23.9% - 134) and *C. eugenioides* (16% -111) (Supplementary Table 9).

The phylogenetic tree of single-copy orthologous NLRs showed that most clades are shared only
among coffee species (Figure 4), but it was also possible to observe clades that clustered NLRs
from solanum, coffee, and reference. Among the clades that clustered coffee NLRs, 71 presented
groupings of orthologs between *C. arabica*, *C. canephora* and *C. eugenioides*, and most of these
are located within the same chromosome. One of these clades, in addition to grouping NLRs of
the three coffee species, includes the reference NLR *RPS2* (RESISTANCE to P. SYRINGAE 2)
(Figure 4). This clade was supported by a high bootstrap value (100%) and was grouped in the
ORTHOMCL45 (Supplementary Table 8). All loci in this cluster were found on chromosome 6
for the three coffee species. Clades that contained NLRs of *C. arabica* and *C. canephora*, *C.*
arabica and *C. eugenioides* and a few *C. canephora* and *C. eugenioides* were also observed.
These are within the same chromosome or on different chromosomes.

Phylogenetic analysis for coffee NLRs classification revealed that members of the NLR
superfamily are grouped into 2 main groups: TIR-NLR (including TNL and NLs) and non-TNLs
(including CNLs and NLs) (Figure 5). NLRs belonging to the non-TNL group outnumbered
those in the TNL group in coffee genomes. We also found that the non-TNLs group is divided
into 13 subgroups and that all subgroups had NLRs from all studied coffee genomes. The same
pattern occurred in the TNL group. Within non-TNLs subgroups it was possible to observe
clades with a greater number of NLRs from *C. arabica* that are shared with *C. eugenioides*
(bands on the outer ring of the tree with a predominance of green and blue colors). There were
exclusive coffee clades as well as clades that contained NLRs that were shared with potato,
tomato, and reference NLRs.

4   Discussion

4.1   NLRs Identification and use of NLR-annotator in coffee genomes

In this study, we investigated loci related to genes of the NLR family in three coffee genomes
and compared these loci with NLR from other plants. The annotation of genes in this family is a
high priority in plant genome sequencing and annotation projects because losses from pathogens

are among the main problems for sustainable agriculture (Steuernagel et al., 2015). A better
understanding of disease resistance in crops will provide plant breeders with tools that may be
used to produce long-term solutions for dealing with future environmental change. A catalog of
NLRs loci, whether complete genes or pseudogenes, within and between species, provides a
toolbox for exploring NLRs that has not previously been described (Jones et al., 2016). Given the
importance of coffee and the availability of the recent *C. arabica* and *C. eugenioides* genomes,
the study of NLRs loci in these species represents an essential source of information for the
development of new disease-resistant cultivars.

As NLR-annotator has already been validated in *C. canephora* (Steuernagel et al., 2020), we
initially used this genome to ensure the reproducibility of the tool in our study and subsequently
applied it with the *C. arabica* and *C. eugenioides* genomes. The tool identified 932 loci NLR for
C. canephora, which is in accordance with Steuernagel et al., 2020, and also predicted of two
distinct NLRs loci within the same genomic position that was previously reported. This
repeatable annotation is the result of a complete NB-ARC domain preceded by a truncated NB-
ARC domain, which makes the tool use the two NB-ARC domains as distinct seeds to identify
two NLRs for the same locus. This has also been reported when this tool was used on the wheat
genome (Steuernagel et al., 2020). The sensitivities and specificities of this tool in coffee
genomes were extremely high (above 99%). In the *Arabidopsis thaliana* genome, which
represents a well-annotated model plant genome, this tool achieved 95% sensitivity, and all loci
that were found were validated to be associated with NLRs (specificity of 100%) (Steuernagel et
al., 2020). In the Nipponbare reference genome of rice, the detection success rate was 99.2%
(Read et al., 2020).

As NLR genes have repeated and clustered genomic distributions in plants, their representation
in genomes using standard gene callers can be underestimated (Jupe et al., 2013; Steuernagel et
al., 2015, 2020). In addition to the high rate of specificity and sensitivity, the NLR-annotator
allowed for the identification of complete loci for coffee in regions distinct from the gene models
already annotated in the reference genomes. This study, therefore, increased the number of
putative NLR genes detected in the reference genomes of coffee species. The complete loci
identified by NLR-annotator that did not overlap the reference genome annotation have also been
reported in rice (Read et al., 2020). It is also relevant to highlight that as this tool is not limited to
searching for functional genes, the complete (pseudogenes) loci that did not overlap with
annotations of the reference genome were also identified for *C. arabica* (90), *C. canephora* (56),
and *C eugenioides* (65). The location of these loci also represent an important resource, since
non-functional alleles identified in sequenced accessions may represent functional NLRs in other
individuals of the same species (Steuernagel et al., 2020). The caturra cultivar (*C. arabica*)
sequenced and used in this study, for example, is used as a susceptible control to differentiate
Hemileia vastatrix races among differential clones (Zambolim and Caixeta, 2021). Pseudogene
regions in this genome may indicate functional genes present in other coffee cultivars.

Our data showed that 18 of the 20 loci found only by NLR-annotator, that did not present
homology to NLRs proteins by BLASTx analysis, have protein domains involved in the activity
of transposable elements (TE). It is known that TE are abundant in plant genomes and that they
play an important role in adaptive evolution and contribute to the evolutionary dynamics of
plant-pathogen interactions (Malacarne et al., 2012; Zhang et al., 2014; Kim et al., 2017). Many
R genes are flanked by TE, which in addition to being sources of genetic variability, are involved

in suppressing or increasing the expression of these genes (Seidl and Thomma, 2017). The Ty3-
gypsy-like TE performance has been reported in a region around the S_H3 locus associated with
CLR resistance. This TE has been described in *C. arabica* subgenomes, replacing the
orthologous counterpart of *C. canephora* with that of *C. eugenioides* (homoeologous non-
reciprocal transposition - HNRT) (Cenci et al., 2012). Moreover, there is evidence of functional
R genes that have evolved through TE-mediated duplications (Seidl and Thomma, 2017), which
demonstrates their importance in the evolutionary changes and expansion of NLR receptors and
justifies the presence of domains related to TE in the studied loci (Zhang et al., 2014; Kim et al.,
2017).

4.2   Distribution of NLR loci in the *Coffea spp.* Genome

Although *C. arabica* results from a natural interspecific hybridization event between *C.*
canephora and *C. eugenioides*, the number of loci found was not proportional to the sum of the
two subgenomes, showing that the hybrid has a relatively smaller number of NLRs loci. The *C.*
canephora genome size is about 690 Mbp, and the *C. eugenioides* is 665 Mbp (Noirot et al.,
2003; Clarindo and Carvalho, 2011; Hamon et al., 2015). The *C. arabica* genome, on the other
hand, is slightly smaller than the sum of its two combined parental genomes (about 1276 Mbp)
(Hamon et al., 2015). This may explain the smaller number of NLRs in this species. Genome
contraction is common in amphidiploids, which may be related to chromosomal rearrangements,
including duplication, insertions, and deletions after initial hybridization (Hamon et al., 2015).
An example of the number of NLRs being smaller than the sum of the corresponding parents was
reported in *Brassica juncea* (Indian mustard), a species formed by hybridization between the
diploid Brassica species of *B. rapa,* and *B. nigra* (Inturrisi et al., 2020). Moreover, differences in
the genome assembly quality may also have interfered with the identification of NLR loci.

Among the three species analyzed, the only one with a genome-wide NLR study already reported
is *C. canephora* (Denoeud et al., 2014). The NLR gene data from the previous study agrees with
much of our findings. A large number of NLR loci in unanchored scaffolds for this species has
also been described. Here 210 complete NLR loci were identified in unanchored scaffolds for *C.*
canephora, while in the first description of the manually curated genes, 213 were not mapped.
The number of mapped NLR genes was 348, while in our study, there were 352 complete loci. In
C. canephora, it has also been reported that NLRs genes are located on all chromosomes, but
with an increased number found on chromosomes 1, 3, 5, 8, 11, which together represented
70.1% of the mapped NLR genes. Although we have highlighted chromosomes 3 and 11 as
having a greater number of NLR, chromosomes 1, 5, and 8 also contain large numbers of NLR
loci in the three species studied here. Together, all these chromosomes represent 68.2, 71.4 and
70.0% of the total of NLR loci mapped for *C. arabica, C. canephora,* and *C. eugenioides*,
respectively. Moreover, the low number of NLR genes on chromosomes 9 and 10 had been
previously reported was confirmed in this study (Denoeud et al., 2014). These comparisons show
that the three species display a conserved pattern with regards to the chromosomal distribution of
NLR loci.

The NLR loci found in the three studied coffee species are arranged in clusters that group
complete loci, pseudogenes and partial. These genes tend to group together and provide birth-
and-death events for functional NLRs (Ling et al., 2021). In these clusters it is possible to find
tandem gene duplications, recombination hotspots or active transposon elements functioning as a

reservoir of genetic variation to generate specificity for new pathogen variants (Michelmore and
Meyers, 1998; Zhang et al., 2014). Within plant genomes many R genes have been found to
reside in clusters (Jupe et al., 2012; Andolfo et al., 2014, 2021; Seo et al., 2016; Zheng et al.,
2016; Read et al., 2020). The S_H3 locus in coffee, for example, corresponds to a complex cluster
of multiple genes, including CNL-like NLR genes (Ribas et al., 2011; Cenci et al., 2012). The
number of complete or functional loci in plants represents a fraction of the total number of loci
found (Jupe et al., 2012; Seo et al., 2016). This happens precisely because the evolutionary
dynamics within these clusters favor the coexistence of functional genes, pseudogenes, and
partial genes, which differ between plants in consequence of evolutionary routes for certain
pathosystems.

Recent discoveries show that NLRs can be multi domain receptors, that is they present domains
integrated to the canonical form NLR or TNL/CNL (Bailey et al., 2018; Wang et al., 2021).
Knowing regions of the genome that have this canonical form can facilitate the description of
non-canonical integrated domains that are upstream or downstream from the more conserved
region (Monteiro and Nishimura, 2018). Other relevant information is that activation of NLRs
often happens in complexes and there is evidence that truncated NLRs can form
heterocomplexes with complete NLRs, or may act as the main receptors in defense activation
(Monteiro and Nishimura, 2018). NLRs truncated as *CbCN* (*Capsicum baccatum* – CC-NB-
ARC), and *TN2* (TIR-NB-ARC) act in resistance to pathogens (Zhao et al., 2015; Son et al.,
2021). This evidence reinforces the importance of knowing loci related to R genes, whether
complete, pseudogenes or partial.

4.3   Ortholog groups and Phylogenetic Analyses

The genus *Coffea* belongs to the Rubiaceae family, which is in the asterid clade that also
contains the Solanaceae family. Many studies have used species of the genus *Solanum* to obtain
insights into the genomic and evolutionary architecture of coffee (Lin et al., 2005; Lefebvre-
Pautigny et al., 2010; Denoeud et al., 2014). Species of the genus *Solanum* have also been used
as models for understanding the molecular processes related to plant-pathogen interaction. This
supports their use for comparative approaches to lead to discoveries of NLR loci or functionally
important gene families in coffee (Andolfo et al., 2021). Our results showed that of the 17
reference NLRs that have been cloned and characterized in species of the genus *Solanum* and
that were also used in this study, 8 of them are present in shared orthogroups with coffee
(Supplementary Table 2 and Supplementary Table 8), being 1 TNL (*Gro1-4)* and 7 CNL (*Hero*,
Prf, *Rpi-blb1*, *Rx2*, *Sw-5*, *Tm-2a*, *Tm-2*). These genes have been found to be involved in
resistance to a diverse group of pathogens including viruses, oomycetes, bacteria and nematodes
(Bendahmane et al., 2000; Van Der Vossen et al., 2003; Paal et al., 2004; Andolfo et al., 2021).
In total, 46 orthogroups were shared between coffee and solanum indicating that these orthologs
were probably present before the speciation of these two groups. Reference NLRs characterized
in species such as *Hordeum vulgare*, *Oryza sativa*, *Triticum aestivum*, *Glycine max*, *Arabidopsis*
thaliana e *Cucumis melo* also share othogroups with coffee NLRs and all of these NLR belong to
the CNL class (Supplementary Table 2 and Supplementary Table 8). These orthogroups are
important as they indicate roles that may be inferred and further investigated in coffee. An
interesting orthogroup that obtained high support in the phylogenetic tree was the one that
clustered the *RPS2* reference NLR as well as NLRs present in all three coffee species. *RPS2 i*s a

resistance gene of *Arabidopsis thaliana* that confers resistance to *Pseudomonas syringae* bacteria
that express the *avrRpt2* avirulence gene (Bent et al., 1994; Mindrinos et al., 1994).

In general, plant species exhibit differences in the number of NLR genes contained within their
genomes. Amplification of certain groups has been detected (Seo et al., 2016). This diversity has
not been associated with genome size or phylogenetic relationships, but is related to the
specialization of each particular host (Wan et al., 2013; Lozano et al., 2015; Seo et al., 2016;
Zheng et al., 2016; Steuernagel et al., 2020). In all three coffee species, a set of orphan genes and
orthogroups that share NLRs within the same species were detected. In tomato, 45 of ~320 NLRs
sequences are more similar to each other than to any other sequences compared (Andolfo et al.,
2021) and orthogroups that share NLRs within the same species of *Solanum* are attributed to
duplication events that generate different gene repertoires and result in species-specific
subfamilies (Seo et al., 2016).

The single-copy ortrogroups provide more reliable results for interpreting evolutionary processes
between groups of evaluated genes by allowing for the identification of true orthologs between
different groups of plants (Zimmer et al., 2007; Duarte et al., 2010). The results from the single-
copy ortrogroups tree showed that some orthologous NLRs seem to have a common ancestor
only among coffee species. The S_H3 locus, for example, was described as being shared only
among coffee species suggesting that the ancestral copy S_H3-CNL was inserted into the S_H3
locus after the divergence of the *Solanum* and *Coffea* lineages (Ribas et al., 2011). The clades
that clustered orthologous NLRs from *C. arabica*, *C. canephora* and *C. eugenioides* probably
represent NLR present in both ancestral diploids genomes, which were passed to *C. arabica*
genome. On the other hand, clades that clustered only *C. canephora* and *C. eugenioides* loci may
represent ancestral NLRs that were lost in *C. arabica* or that underwent so many modifications in
this species that makes it difficult to find homology between these NLRs. These NLR may
provide valuable resistance mechanisms that are not present in the *C. arabica* hybrid. Nucleotide
level changes, such as deletions, insertions and rearrangements have been observed in coffee
RGA (Resistance gene analogues) (Noir et al., 2001; Hendre et al., 2011). It is also known that it
is very likely that the sequenced genotypes of *C. canephora* and *C. eugenioides* present
significant differences from the ancestral donors of *C. arabica* subgenomes, which may explain
the lack of homology in certain NLR groups (Cenci et al., 2012).

The NB-ARC is the most conserved domain in the NLR gene family. Despite the conservation of
this domain, it is possible to distinguish the TIR (TNL) and non-TIR classes based on different
residues inside the motifs present in this region (Jones et al., 2016; Shao et al., 2019; Van
Ghelder et al., 2019). Therefore, this domain has been used to describe the phylogenetic
relationships between the sequences of this group and classify them (Andolfo et al., 2014). The
classification of NLRs in coffee revealed that the non-TNL class were present in greater numbers
than those of the TNL group in each of the three analyzed coffee genomes. It is known that non-
TNL genes that include many CNL are widely distributed in monocots and dicots, while TNL are
mainly found in dicots (McHale et al., 2006; Zheng et al., 2016). The low frequency of TNLs in
coffee agrees with results found for species of the solanum group, such as pepper, tomato and
potato (Andolfo et al., 2014; Seo et al., 2016). Our results are also consistent with the low
frequency of TNLs found in previous studies of coffee (Hendre et al., 2011; Denoeud et al.,
2014). Thus, it is possible to suggest that, as in solanum, non-TNLs represent an important
repertoire of resistance genes in coffee. Additionally, the TNL group and the non-TNLs

subgroups contained NLRs from *C.arabica*, *C. canephora* and *C. eugenioides*, indicating
conservation of the NLR classes across coffee genomes and suggesting that all subgroups were
present in a common ancestor, similar to what has been described for comparisons of species
within the solanum group (Seo et al., 2016).

In the two phylogenetic trees analyzed, the clades group coffee NLRs that are mostly present in
the same chromosomes but groupings of NLRs present on different chromosomes were also
detected. Genes located on the same chromosome tend to group into subclades in the
phylogenetic tree. However, rearrangement events of the chromosomes can affect NLR loci and
modify their genomic order or location (Cenci et al., 2012; Zheng et al., 2016; Ling et al., 2021).

Considering the relevance of coffee, few studies have been conducted addressing the
identification of NLR in genomes of this crop. RGA studies using degenerate primers for NB-
ARC region have already been performed (Noir et al., 2001; Hendre et al., 2011; Kumar, 2012),
in addition to studies in S_H3 locus (Cenci et al., 2010, 2012; Lashermes et al., 2010), but very
little is known about the NLR family in cultivated (*C.arabica* e *C. canephora*) and uncultivated
coffee species (such as *C. eugenioides*). This is the first study focused on genome wide
identification of NLRs in the *C. arabica* genome, and also adds information to the existing report
for the *C. canephora* genome (Denoeud et al., 2014). The Genome-wide identification of coffee
NLRs allows for more in-depth molecular studies provides further information for identifying
candidate genes for cloning and subsequent functional validation of NLR genes while also
expanding the range of NLR that are available for breeding of this crop (Seo et al., 2016).

5   Conclusion

Our analysis showed that the use of a specific pipeline for resistance genes was efficient in
detecting NLR loci in the studied coffee genomes and increased the information available for the
location of these loci in *C. arabica*, *C. canephora* and *C. eugenioides*. The NLR loci in the three
coffee species are unevenly distributed across all chromosomes and are mostly arranged in
clusters. The number of loci in *C. arabica* is less than the sum of the NLR loci from the parents
of this hybrid. Single-copy NLR orthogroups investigated in a phylogenetic tree identified
orthologous NLRs that are shared between *C. arabica* and the parental genomes as well as NLR
that were possibly lost. Coffee NLRs and some functionally characterized NLRs share common
orthogroups, which provides clues to the functionality of coffee NLRs, and paves the way for
further investigation. Loci from the NLR family are subdivided into two main groups in coffee:
TIR-NLR (TNL) and non-TNL, but non-TNLs are present in greater number and seem to
represent an important repertoire of resistance genes in coffee.

6   Conflict of Interest

618 The authors declare that the research was conducted in the absence of any commercial or
619 financial relationships that could be construed as a potential conflict of interest.

7   Author Contributions

MLi, GS, and MF: conceptualization. MLi, MLu, GS, JC, MF, and JT: methodology. MLi, JC:
software, formal analysis, investigation. MLi writing – original draft preparation. MLi, MLu,
GS, JC, MF, and JT: visualization. MLi, MLu, GS, JC, MF, and JT: validation. MLu, and MF:
resources, supervision, project administration, and funding acquisition. MLi, MLu, GS, JC, MF,
and JT: data curation; writing – review and editing. All authors have read and agreed to the
published version of the manuscript.

8   Funding

This research was funded by Coordination for the Improvement of Higher Education Personnel
(CAPES) and University of Florida to MLS, and the USDA McIntire-Stennis project FLA-PLP-
005931 "Developing and scaling up the next generation of healthy forests" to JTB.

9   Acknowledgments

We are grateful to the Universidade Federal de Lavras, the Universidade de Brasília, the
University of Florida, U.S. Department of Agriculture's Agriculture and Food Research Initiative
grant FLA-PLP006039, University of Florida's Institute for Food and Agricultural Science, and
Corn Breeding and Genomics program for supporting this collaborative research project.

References

Anand, L., and Lopez, C. M. R. (2020). chromoMap: An R package for Interactive Visualization
and Annotation of Chromosomes.

Andolfo, G., D'agostino, N., Frusciante, L., and Ercolano, M. R. (2021). The tomato
interspecific NB-LRR gene arsenal and its impact on breeding strategies. *Genes.* 12, 1–12.
doi:10.3390/genes12020184.

Andolfo, G., Jupe, F., and Witek, K. (2014). Defining the full tomato NB-LRR resistance gene
repertoire using genomic and cDNA RenSeq. *BMC plant* 14, 120, 2014. doi:10.1186/1471-
2229-14-120.

Bailey, P. C., Schudoma, C., Jackson, W., Baggs, E., Dagdas, G., Haerty, W., et al. (2018).
Dominant integration locus drives continuous diversification of plant immune receptors
with exogenous domain fusions. *Genome Biol.* 19, 1–18. doi:10.1186/s13059-018-1392-6.

Barragan, A. C., and Weigel, D. (2021). Plant NLR Diversity: The Known Unknowns of Pan-
NLRomes. *Plant Cell* 33, 814–831. doi:/10.1093/plcell/koaa002.

Bawin, Y., Ruttink, T., Staelens, A., Haegeman, A., Stoffelen, P., Mwanga Mwanga, J. C. I., et
al. (2020). Phylogenomic analysis clarifies the evolutionary origin of *Coffea arabica*. *J.*

653  *Syst. Evol.* doi:10.1111/jse.12694.

654 Bendahmane, A., Querci, M., Kanyuka, K., and Baulcombe, D. C. (2000). Agrobacterium
655  transient expression system as a tool for the isolation of disease resistance genes:
656  Application to the *Rx2* locus in potato. *Plant J.* 21, 73–81. doi:/10.1046/j.1365-
657  313x.2000.00654.x.

658 Bent, A. F., Kunkel, B. N., Dahlbeck, D., Brown, K. L., Schmidt, R., Giraudat, J., et al. (1994).
659  RPS2 of Arabidopsis thaliana: A leucine-rich repeat class of plant disease resistance genes.
660  *Science* 265, 1856–1860. doi:10.1126/science.8091210.

661 Bezerra-Neto, J. P., Araújo, F. C., Ferreira-Neto, J. R. C., Silva, R. L. O., Borges, A. N. C.,
662  Matos, M. K. S., et al. (2020). "NBS-LRR genes-Plant health sentinels: Structure, roles,
663  evolution and biotechnological applications," in *Applied Plant Biotechnology for Improving*
664  *Resistance to Biotic Stress* (Elsevier), 63–120. doi:10.1016/B978-0-12-816030-5.00004-5.

665 Bonardi, V., Cherkis, K., Nishimura, M. T., and Dangl, J. L. (2012). A new eye on NLR
666  proteins: Focused on clarity or diffused by complexity? *Curr. Opin. Immunol.* 24, 41–50.
667  doi:10.1016/j.coi.2011.12.006.

668 Boutrot, F., and Zipfel, C. (2017). Function, discovery, and exploitation of plant pattern
669  recognition receptors for broad-spectrum disease resistance. *Annu. Rev. Phytopathol.* 55,
670  257–286. doi:10.1146/annurev-phyto-080614-120106.

671 Cabral, P. G. C., Maciel-Zambolim, E., Oliveira, S. A. S., Caixeta, E. T., and Zambolim, L.
672  (2016). Genetic diversity and structure of *Hemileia vastatrix* populations on *Coffea spp.*
673  *Plant Pathol.* 65, 196–204. doi:10.1111/ppa.12411.

674 Cenci, A., Combes, M. C., and Lashermes, P. (2010). Comparative sequence analyses indicate
675  that Coffea (Asterids) and Vitis (Rosids) derive from the same paleo-hexaploid ancestral
676  genome. *Mol. Genet. Genomics* 283, 493–501. doi:10.1007/s00438-010-0534-7.

677 Cenci, A., Combes, M. C., and Lashermes, P. (2012). Genome evolution in diploid and tetraploid
678  Coffea species as revealed by comparative analysis of orthologous genome segments. *Plant*
679  *Mol. Biol.* 78, 135–145. doi:10.1007/s11103-011-9852-3.

680 Cerda, R., Avelino, J., Gary, C., Tixier, P., Lechevallier, E., and Allinne, C. (2017). Primary and
681  secondary yield losses caused by pests and diseases: Assessment and modeling in coffee.
682  *PLoS One* 12, 1–17. doi:10.1371/journal.pone.0169133.

683 Clarindo, W. R., and Carvalho, C. R. (2011). Flow cytometric analysis using SYBR Green I for
684  genome size estimation in coffee. *Acta Histochem.* 113, 221–225.
685  doi:10.1016/j.acthis.2009.10.005.

686 Conway, J. R., Lex, A., and Gehlenborg, N. (2017). UpSetR: An R package for the visualization
687  of intersecting sets and their properties. *Bioinformatics* 33, 2938–2940.
688  doi:10.1093/bioinformatics/btx364.

Davis, A. P., Govaerts, R., Bridson, D. M., and Stoffelen, P. (2006). An annotatated taxonomic
of the genus coffea (Rubiaceae). *Bot. J. Linn. Soc.* 152, 465–512. doi:10.1111/j.1095-
8339.2006.00584.x.

Denoeud, F., Carretero-Paulet, L., Dereeper, A., Droc, G., Guyot, R., Pietrella, M., et al. (2014).
The coffee genome provides insight into the convergent evolution of caffeine biosynthesis.
Science 345, 1181–1184. doi:10.1126/science.1255274.

Duarte, J. M., Wall, P. K., Edger, P. P., Landherr, L. L., Ma, H., Pires, J. C., et al. (2010).
Identification of shared single copy nuclear genes in Arabidopsis, Populus, Vitis and Oryza
and their phylogenetic utility across various taxonomic levels. *BMC Evol. Biol.* 10, 1–18.
doi:10.1186/1471-2148-10-61.

Hamon, P., Hamon, S., Razafinarivo, N. J., Guyot, R., Siljak-Yakovlev, S., Couturon, E., et al.
(2015). "Coffea Genome Organization and Evolution," in *Coffee in Health and Disease*
Prevention, 29–37. doi:10.1016/B978-0-12-409517-5.00004-8.

Hendre, P. S., Bhat, P. R., Krishnakumar, V., Aggarwal, R. K., and Donini, P. (2011). Isolation
and characterization of resistance gene analogues from Psilanthus species that represent
wild relatives of cultivated coffee endemic to India. *Genome* 54, 377–390. doi:10.1139/g11-
004.

Hufford, M. B., Seetharam, A. S., Woodhouse, M. R., Chougule, K. M., Coletta, D., Tittes, S., et
al. (2021). De novo assembly, annotation, and comparative analysis of 26 diverse maize
genomes. *Science.* doi:10.1126/science.abg5289.

Inturrisi, F., Bayer, P. E., Yang, H., Tirnaz, S., Edwards, D., and Batley, J. (2020). Genome-wide
identification and comparative analysis of resistance genes in *Brassica juncea. Mol. Breed.*
40, 1–14. doi:10.1101/2020.12.15.422814.

Jones, J. D. G., and Dangl, J. L. (2006). The plant immune system. *Nature* 444, 323–329.
doi:10.1038/nature05286.

Jones, J. D. G., Vance, R. E., and Dangl, J. L. (2016). Intracellular innate immune surveillance
devices in plants and animals. *Science* 354. doi:10.1126/science.aaf6395.

Jost, M., Singh, D., Lagudah, E., Park, R. F., and Dracatos, P. (2020). Fine mapping of leaf rust
resistance gene Rph13 from wild barley. *Theor. Appl. Genet.* 133, 1887–1895.
doi:10.1007/s00122-020-03564-6.

Jupe, F., Pritchard, L., Etherington, G. J., MacKenzie, K., Cock, P. J. A., Wright, F., et al.
(2012). Identification and localisation of the NB-LRR gene family within the potato
genome. *BMC Genomics* 13, 1–14. doi:10.1186/1471-2164-13-75.

Jupe, F., Witek, K., Verweij, W., Śliwka, J., Pritchard, L., Etherington, G. J., et al. (2013).
Resistance gene enrichment sequencing (RenSeq) enables reannotation of the NB-LRR
gene family from sequenced plant genomes and rapid mapping of resistance loci in
segregating populations. *Plant J.* 76, 530–544. doi:10.1111/tpj.12307.

Kachroo, A., and Robin, G. P. (2013). Systemic signaling during plant defense. *Curr. Opin.*
Plant Biol. 16, 527–533. doi:10.1016/j.pbi.2013.06.019.

Katoh, K., Misawa, K., Kuma, K., and Miyata, T. (2002). MAFFT: a novel method for rapid
multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* 30, 3059–
66. doi:/10.1093/nar/gkf436.

Kim, S., Park, J., Yeom, S. I., Kim, Y. M., Seo, E., Kim, K. T., et al. (2017). New reference
genome sequences of hot pepper reveal the massive evolution of plant disease-resistance
genes by retroduplication. *Genome Biol.* 18, 1–11. doi:10.1186/s13059-017-1341-9.

Kourelis, J., and Van Der Hoorn, R. A. (2018). Defended to the Nines: 25 years of Resistance
Gene Cloning Identifies Nine Mechanisms for R Protein Function. *Plant Cell* 30, 285–299.
doi:10.1105/tpc.17.00579.

Krishnan, S. (2017). Sustainable Coffee Production. *Oxford Res. Encycl. Environ. Sci.*
doi:10.1093/acrefore/9780199389414.013.224.

Kroj, T., Chanclud, E., Michel-Romiti, C., Grand, X., and Morel, J. B. (2016). Integration of
decoy domains derived from protein targets of pathogen effectors into plant immune
receptors is widespread. *New Phytol.* 210, 618–626. doi:10.1111/nph.13869.

Kumar, D. (2012). Isolation of Nucleotide Binding Site (NBS)-Leucine Rich Repeat (LRR)
Resistant Gene Analogs (Rgas) In Arabica Coffee (*Coffea arabica* L. Cv S.288). *J.*
Biotechnol. Biomater. 02. doi:10.4172/2155-952x.1000146.

Kushwaha, S. K., Chauhan, P., Hedlund, K., and Ahren, D. (2016). NBSPred: A support vector
machine-based high-throughput pipeline for plant resistance protein NBSLRR prediction.
Bioinformatics 32, 1223–1225. doi:10.1093/bioinformatics/btv714.

Lashermes, P., Combes, M. C., Ribas, A., Cenci, A., Mahé, L., and Etienne, H. (2010). Genetic
and physical mapping of the SH3 region that confers resistance to leaf rust in coffee tree
(*Coffea arabica* L.). *Tree Genet. Genomes* 6, 973–980. doi:10.1007/s11295-010-0306-x.

Lashermes, P., Combes, M. C., Robert, J., Trouslot, P., D'Hont, A., Anthony, F., et al. (1999).
Molecular characterization and origin of the Coffea arabica L. genome. *Mol. Gen. Genet.*
261, 259–266. doi:10.1007/s004380050965.

Lefebvre-Pautigny, F., Wu, F., Philippot, M., Rigoreau, M., Priyono, Zouine, M., et al. (2010).
High resolution synteny maps allowing direct comparisons between the coffee and tomato
genomes. *Tree Genet. Genomes* 6, 565–577. doi:10.1007/s11295-010-0272-3.

Letunic, I., and Bork, P. (2021). Interactive tree of life (iTOL) v3: an online tool for the display
and annotation of phylogenetic and other trees. *Nucleic Acids Res.* 49, W293–W296.
doi:10.1093/nar/gkw290.

Li, L., Stoeckert, C. J. J., and Roos, D. S. (2003). OrthoMCL: Identification of Ortholog Groups
for Eukaryotic Genomes. *Genome Res.* 13, 2178–2189. doi:10.1101/gr.1224503.candidates.

Lin, C., Mueller, L. A., Carthy, J. M., Crouzillat, D., Pétiard, V., and Tanksley, S. D. (2005).
Coffee and tomato share common gene repertoires as revealed by deep sequencing of seed
and cherry transcripts. *Theor. Appl. Genet.* 112, 114–130. doi:10.1007/s00122-005-0112-2.

Ling, J., Xie, X., Gu, X., Zhao, J., Ping, X., Li, Y., et al. (2021). High-quality chromosome-level
genomes of *Cucumis metuliferus* and *Cucumis melo* provide insight into Cucumis genome
evolution. *Plant J.* 107, 136–148. doi:10.1111/tpj.15279.

Lozano, R., Hamblin, M. T., Prochnik, S., and Jannink, J. L. (2015). Identification and
distribution of the NBS-LRR gene family in the Cassava genome. *BMC Genomics* 16, 1–14.
doi:10.1186/s12864-015-1554-9.

Lu, Y., and Tsuda, K. (2021). Intimate Association of PRR- and NLR-Mediated Signaling in
Plant Immunity. *Mol. Plant-Microbe Interact.* 34, 3–14. doi:10.1094/mpmi-08-20-0239-ia.

Malacarne, G., Perazzolli, M., Cestaro, A., Sterck, L., Fontana, P., van de Peer, Y., et al. (2012).
Deconstruction of the (paleo)polyploid grapevine genome based on the analysis of
transposition events involving NBS resistance genes. *PLoS One* 7.
doi:10.1371/journal.pone.0029762.

McCook, S., and Vandermeer, J. (2015). The Big Rust and the Red Queen: Long-Term
Perspectives on Coffee Rust Research. *Phytopathology* 105, 1164–1173.
doi:10.1094/PHYTO-04-15-0085-RVW.

McHale, L., Tan, X., Koehl, P., and Michelmore, R. W. (2006). Plant NBS-LRR proteins:
Adaptable guards. *Genome Biol.* 7. doi:10.1186/gb-2006-7-4-212.

Michelmore, R. W., and Meyers, B. C. (1998). Clusters of resistance genes in plants evolve by
divergent selection and a birth-and-death process. *Genome Res.* 8, 1113–1130.
doi:10.1101/gr.8.11.1113.

Mindrinos, M., Katagiri, F., Yu, G. L., and Ausubel, F. M. (1994). The A. thaliana disease
resistance gene RPS2 encodes a protein containing a nucleotide-binding site and leucine-
rich repeats. *Cell* 78, 1089–1099. doi:10.1016/0092-8674(94)90282-8.

Monteiro, F., and Nishimura, M. T. (2018). Structural, functional, and genomic diversity of plant
NLR proteins: An evolved resource for rational engineering of plant immunity. *Annu. Rev.*
Phytopathol. 56, 243–267. doi:10.1146/annurev-phyto-080417-045817.

Muliyar, R. K., Chowdappa, P., Behera, S. K., Kasaragod, S., Gangaraj, K. P., Kotimoole, C. N.,
et al. (2020). Assembly and Annotation of the Nuclear and Organellar Genomes of a Dwarf
Coconut (Chowghat Green Dwarf) Possessing Enhanced Disease Resistance. *Omi. A J.*
Integr. Biol. 24, 726–742. doi:10.1089/omi.2020.0147.

Noir, S., Combes, M.-C., Anthony, F., and Lashermes, P. (2001). Origin, diversity and evolution
of NBS-type disease-resistance gene homologues in coffee trees (Coffea L.). *Mol. Genet.*
Genomics 265, 654–662. doi:10.1007/s004380100459.

798 Noirot, M., Poncet, V., Barre, P., Hamon, P., Hamon, S., and De Kochko, A. (2003). Genome
799   size variations in diploid African Coffea species. *Ann. Bot.* 92, 709–714.
800   doi:10.1093/aob/mcg183.

801 Osuna-Cruz, C. M., Paytuvi-Gallart, A., Di Donato, A., Sundesha, V., Andolfo, G., Cigliano, R.
802   A., et al. (2018). PRGdb 3.0: A comprehensive platform for prediction and analysis of plant
803   disease resistance genes. *Nucleic Acids Res.* 46, D1197–D1201. doi:10.1093/nar/gkx1119.

804 Paal, J., Henselewski, H., Muth, J., Meksem, K., Menéndez, C. M., Salamini, F., et al. (2004).
805   Molecular cloning of the potato Gro1-4 gene conferring resistance to pathotype Ro1 of the
806   root cyst nematode Globodera rostochiensis, based on a candidate gene approach. *Plant J.*
807   38, 285–297. doi:10.1111/j.1365-313X.2004.02047.x.

808 Read, A. C., Moscou, M. J., Zimin, A. V., Pertea, G., Meyer, R. S., Purugganan, M. D., et al.
809   (2020). Genome assembly and characterization of a complex zfBED-NLR gene-containing
810   disease resistance locus in Carolina Gold Select rice with Nanopore sequencing. *PLoS*
811   *Genet.* 16, 1–24. doi:10.1371/journal.pgen.1008571.

812 Ribas, A. F., Cenci, A., Combes, M. C., Etienne, H., and Lashermes, P. (2011). Organization and
813   molecular evolution of a disease-resistance gene cluster in coffee trees. *BMC Genomics* 12.
814   doi:10.1186/1471-2164-12-240.

815 Schatz, M. C., Maron, L. G., Stein, J. C., Hernandez Wences, A., Gurtowski, J., Biggers, E., et
816   al. (2014). Whole genome de novo assemblies of three divergent strains of rice, Oryza
817   sativa, document novel gene space of aus and indica. *Genome Biol.* 15, 506.
818   doi:10.1186/s13059-014-0506-z.

819 Scott, A. D., Zimin, A. V, Puiu, D., Workman, R., Britton, M., Zaman, S., et al. (2020). A
820   Reference Genome Sequence for Giant Sequoia. *G3: Genes, Genomes, Genetics*. 10, 3907–
821   3919. doi: 10.1534/g3.120.401612.

822 Seidl, M. F., and Thomma, B. P. H. J. (2017). Transposable Elements Direct The Coevolution
823   between Plants and Microbes. *Trends Genet.* 33, 842–851. doi:10.1016/j.tig.2017.07.003.

824 Seo, E., Kim, S., Yeom, S. I., and Choi, D. (2016). Genome-wide comparative analyses reveal
825   the dynamic evolution of nucleotide-binding leucine-rich repeat gene family among
826   solanaceae plants. *Front. Plant Sci.* 7, 1–13. doi:10.3389/fpls.2016.01205.

827 Shao, Z. Q., Xue, J. Y., Wang, Q., Wang, B., and Chen, J. Q. (2019). Revisiting the Origin of
828   Plant NBS-LRR Genes. *Trends Plant Sci.* 24, 9–12. doi:10.1016/j.tplants.2018.10.015.

829 Son, S., Kim, S., Lee, K., Oh, J., Choi, I., Do, J., et al. (2021). The *Capsicum baccatum*-Specific
830   Truncated NLR Protein CbCN Enhances the Innate Immunity against Colletotrichum
831   acutatum. *Int. J. Mol. Sci.* 22. doi:10.3390/ijms22147672.

832 Song, W., Wang, B., Li, X., Wei, J., Chen, L., Zhang, D., et al. (2015). Identification of Immune
833   Related LRR-Containing Genes in Maize (*Zea mays L.*) by Genome-Wide Sequence
834   Analysis. *Int. J. Genomics* 2015. doi:10.1155/2015/231358.

Stamatakis, A. (2014). RAxML version 8: A tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* 30, 1312–1313. doi:10.1093/bioinformatics/btu033.

Stanke, M., Keller, O., Gunduz, I., Hayes, A., Waack, S., and Morgenstern, B. (2006).
AUGUSTUS: ab initio prediction of alternative transcripts. *Nucleic Acids Res.* 34, W435–
W439. doi:10.1093/nar/gkl200.

Steuernagel, B., Jupe, F., Witek, K., Jones, J. D. G., and Wulff, B. B. H. (2015). NLR-parser:
Rapid annotation of plant NLR complements. *Bioinformatics* 31, 1665–1667.
doi:10.1093/bioinformatics/btv005.

Steuernagel, B., Witek, K., Krattinger, S. G., Ramirez-Gonzalez, R. H., Schoonbeek, H. J., Yu,
G., et al. (2020). The NLR-annotator tool enables annotation of the intracellular immune
receptor repertoire. *Plant Physiol.* 183, 468–482. doi:10.1104/pp.19.01273.

Sun, X., Jiao, C., Schwaninger, H., Chao, C. T., Ma, Y., Duan, N., et al. (2020). Phased diploid
genome assemblies and pan-genomes provide insights into the genetic history of apple
domestication. *Nat. Genet.* 52, 1423–1432. doi:10.1038/s41588-020-00723-9.

Toda, N., Rustenholz, C., Baud, A., Le Paslier, M. C., Amselem, J., Merdinoglu, D., et al.
(2020). NLGenomeSweeper: A tool for genome-wide NBS-LRR resistance gene
identification. *Genes.* 11, 333. doi:10.3390/genes11030333.

Tran, H. T. M., Ramaraj, T., Furtado, A., Lee, L. S., and Henry, R. J. (2018). Use of a draft
genome of coffee (*Coffea arabica*) to identify SNPs associated with caffeine content. *Plant
Biotechnol. J.* 9, 1756–1766. doi:10.1111/pbi.12912.

Van Der Vossen, E., Sikkema, A., Te Lintel Hekkert, B., Gros, J., Stevens, P., Muskens, M., et
al. (2003). An ancient R gene from the wild potato species *Solanum bulbocastanum* confers
broad-spectrum resistance to *Phytophthora infestans* in cultivated potato and tomato. *Plant
J.* 36, 867–882. doi:10.1046/j.1365-313X.2003.01934.x.

Van Ghelder, C., Parent, G. J., Rigault, P., Prunier, J., Giguère, I., Caron, S., et al. (2019). The
large repertoire of conifer NLR resistance genes includes drought responsive and highly
diversified RNLs. *Sci. Rep.* 9, 1–13. doi:10.1038/s41598-019-47950-7.

Wan, H., Yuan, W., Bo, K., Shen, J., Pang, X., and Chen, J. (2013). Genome-wide analysis of
NBS-encoding disease resistance genes in *Cucumis sativus* and phylogenetic study of NBS-
encoding genes in Cucurbitaceae crops. *BMC Genomics* 14, 109. doi:10.1186/1471-2164-
14-109.

Wang, Z., Ren, H., Xu, F., Lu, G., Cheng, W., Que, Y., et al. (2021). Genome-Wide
Characterization of NLRs in *Saccharum spontaneum* L. and Their Responses to Leaf Blight
in Saccharum. *Agronomy* 11, 153. doi:10.3390/plants10020322.

Yuan, M., Jiang, Z., Bi, G., Nomura, K., Liu, M., He, S. Y., et al. (2021). Pattern-recognition
receptors are required for NLR-mediated plant immunity. *Nature*, 1–5.
doi:10.1101/2020.04.10.031294.

Zambolim, L., and Caixeta, E. T. (2021). An overview of physiological specialization ofcoffee
leaf rust – new designation of Pathotypes. *Int. J. Curr. Res.* 13, 15564–15575.

Zhang, R., Murat, F., Pont, C., Langin, T., and Salse, J. (2014). Paleo-evolutionary plasticity of
plant disease resistance genes. *BMC Genomics* 15. doi:10.1186/1471-2164-15-187.

Zhao, T., Rui, L., Li, J., Nishimura, M. T., Vogel, J. P., Liu, N., et al. (2015). A Truncated NLR
Protein, TIR-NBS2, Is Required for Activated Defense Responses in the exo70B1 Mutant.
PLoS Genet. 11, 1–28. doi:10.1371/journal.pgen.1004945.

Zheng, F., Wu, H., Zhang, R., Li, S., He, W., Wong, F. L., et al. (2016). Molecular phylogeny
and dynamic evolution of disease resistance genes in the legume family. *BMC Genomics*
17, 1–13. doi:10.1186/s12864-016-2736-9.

Zimmer, A., Lang, D., Richardt, S., Frank, W., Reski, R., and Rensing, S. A. (2007). Dating the
early evolution of plants: Detection and molecular clock analyses of orthologs. *Mol. Genet.*
Genomics 278, 393–402. doi:10.1007/s00438-007-0257-6.

Figure 1. Venn diagrams representing the overlap between the loci from NLR-annotator
and NLR genes annotated in the *C. arabica*, *C. canephora,* and *C. eugenioides* reference
genomes. The colors represent the origin of the annotation, with blue indicating those annotated
by NLR-annotator and green indicating those found in the reference genome. The intersection
refers to the overlaps that occurred once or more than once.

Figure 2. Number and chromosomal distribution of NLR loci in *C. arabica*, *C. canephora* e
C. eugenioides. **(A)** The chromosomes with the highest number of NLR loci are highlighted in
dark blue, and those with the smallest number of NRL are highlighted in light blue. CPL
identifies the completeness of NLR as: C= complete, Cps= complete (pseudogene), P= partial,
Pps= partial pseudogene and Un= unassigned. **(B)** The chromosomes represented in *C. arabica*
refer to the two subgenomes with the first originating from *C. canephopr*a and the second
originating from *C. eugenioides*. The chromosomal distribution represented in this figure does
not show all loci for each region but identifies all regions that contain NLRs loci. A more
detailed view of these chromosomes with the detail of all regions may be found at:
https://1drv.ms/u/s!As084N7WlXAIhMZfxevH93zPgU-YhQ?e=0KQ8Iy. Access the link and
download the HTML file.

Figure 3. **Upset plot of orthologous NLR groups (orthogroups) among five species, *C.***
arabica* (Ca) *C. canephora* (Cc), *C. eugenioides* (Ce), *S. tuberosum* (Soltu) and *S.
***lycopersicum* (Soly) and NLRs de referência (Ref).** The orthogroups that cluster combinations
of species/Ref NLRs is shown by the interconnected blue dots on the bottom panel. Unconnected
blue dots show orthogroups that are present within the same species. The 'Set size' represents the
total number of orthogroups per species/Ref. The 'intersection size' shows the number of
orthogroups shared between species/Ref or within the same species/Ref. The orthogroups were
clustered using OrthoMCL.

Figure 4. **Phylogenetic tree of single-copy NLR orthogroups.** The phylogenetic tree was
constructed using RAxML and was based on 647 NLRs (domain NB-ARC) that were single-
copy orthologs. The colored ring indicates coffee NLR clades, the green color represents *C.*
arabica (Ca), red represents *C. canephora* (Cc) and blue represents *C. eugenioides* (Ce). Labels
in black are coffee NLRs, green is used for *S. lycopersicum* (Soly), blue is used for *S. tuberosum*
(Soltu) and pink indicates reference proteins (Ref). Bootstrap values above 70% are indicated on
each branch with a brown circle. The pink background identifies clades that group orthologs of
Ca, Cc and Ce. The clade highlighted in purple shows the coffee NLRs and RPS2 grouping.

Figure 5. **Phylogenetic tree for coffee NLRs classification**. NB-ARC domains from 2681
NLRs clustering *C. arabica* (*Ca*), *C. canephora (Cc), C. eugenioides* (Ce), *S. lycopersicum*, *S.*
tuberosum, reference NLRs (pink balls) and CED-4 (root nematode outlier) were used to
construct the tree. The tree was constructed using RAxML. The classifications of the reference
NLRs and some *S. lycopersicum*, *S. tuberosum* NLRs were used to classify the coffee NLRs into
TNLs and Non TNLs groups (inner ring - TNL = yellow, CNL = gray and NL = Purple).
Subgroups in Non TNLs are indicated from I to XIII and alternating colors (green and purple).
Gray and yellow background highlight coffee NLRs and the outer ring separates the NLRs for
Ca, Cc and Ce in green, red and blue respectively.

Table 1. Loci identified using NLR-annotator that did not overlap with annotations of NLR
genes from coffee reference genomes.

Species	Complete	Complete (pseudogene)	Partial	Partial (pseudogene)	Total
C. arabica	70	90	65	73	298
C. canephora	67	56	73	44	240
C. eugenioides	71	65	37	49	222

Figure 1

A

Coffea arabica

CPL	Chr1c	Chr1e	Chr2c	Chr2e	Chr3c	Chr3e	Chr4c	Chr4e	Chr5c	Chr5e	Chr6c	Chr6e	Chr7c	Chr7e	Chr8c	Chr8e	Chr9c	Chr9e	Chr10c	Chr10e	Chr11c	Chr11e	Un	Total
C	42	48	17	22	71	65	25	21	42	49	23	15	44	43	35	54	11	7	16	9	54	73	23	809
Cps	21	13	7	8	38	24	5	6	28	19	3	6	14	15	12	23	1	4	1	1	12	23	5	289
P	5	4	4	6	13	12	1	4	4	5	10	2	5	6	4	3	1	0	5	8	10	7	6	125
Pps	5	2	2	3	8	4	2	1	1	2	2	4	2	4	4	9	2	0	4	3	10	13	0	88
Total	73	67	30	39	130	105	33	32	75	75	38	27	65	68	55	89	15	11	26	21	86	116	34	1311

Coffea canephora

CPL	Chr1	Chr2	Chr3	Chr4	Chr5	Chr6	Chr7	Chr8	Chr9	Chr10	Chr11	Un	Total
C	49	19	70	23	42	16	29	37	1	11	55	210	562
Cps	14	9	25	7	14	4	7	12	1	1	14	66	174
P	3	7	12	3	5	2	5	2	3	3	8	68	121
Pps	6	2	13	1	2	2	2	3	1	1	13	23	70
Total	72	37	120	34	63	24	43	54	6	16	90	367	927

Coffea eugenioides

CPL	Chr1	Chr2	Chr3	Chr4	Chr5	Chr6	Chr7	Chr8	Chr9	Chr10	Chr11	Un	Total
C	56	30	98	17	86	40	56	106	18	10	90	88	695
Cps	14	17	33	7	34	16	18	27	5	3	29	36	239
P	3	10	13	2	9	4	8	9	1	3	16	5	83
Pps	2	3	11	1	5	3	3	6	3	4	14	6	62
Total	75	60	155	27	134	63	85	148	27	20	149	135	1079

B

C. canephora

C. arabica

C. eugenioides

Legend: complete (green), complete (pseudogene) (orange), partial (yellow), partial (pseudogene) (red)

Length (bp)

Figure 2

Figure 3

Figure 4

Labels
- Coffee (black)
- Soltu (blue)
- Soly (yellow-green)
- Ref (magenta)

Ring
- Ca (green)
- Cc (red)
- Ce (blue)

Background
- Ca / Ce / Cc clades (pink)

Bootstrap
- ≥ 70% (brown circle)

Clade
- RPS2 (purple)

Figure 5

Inner ring
- TNL
- CNL
- NL

Outer ring
- Ca
- Cc
- Ce

Backgrounds
- Coffee NLRs - TNLs
- Coffee NLRs - Non-TNLs

Reference NLRs