Requirement of cholesterol for calcium-dependent vesicle fusion by stabilizing synaptotagmin-1-induced membrane bending

Houda Yasmine Ali Moussa[1*], Kyung Chul Shin[1*], Yongfeng Tong[2], Soo Jin Kim[3], Je-Kyung
Ryu[4], Said Mansour[2], Yongsoo Park[1,5*†]

[1]Neurological Disorders Research Center, Qatar Biomedical Research Institute (QBRI), Hamad Bin
Khalifa University (HBKU), Qatar Foundation, Doha, Qatar
[2]Qatar Environment and Energy Research Institute (QEERI), Hamad Bin Khalifa University (HBKU),
Doha, Qatar
[3]Division of Molecular and Life Sciences, Pohang University of Science and Technology, Pohang, 790-
784, Republic of Korea
[4]Department of Biological Sciences, KAIST, Daejeon 305-701, South Korea
[5]College of Health & Life Sciences (CHLS), Hamad Bin Khalifa University (HBKU), Qatar Foundation,
Doha, Qatar
12
13 *These authors contributed equally to this work.
14 †Corresponding authors;
15 Dr. Yongsoo Park, Neurological Disorders Research Center, Qatar Biomedical Research Institute (QBRI),
16 Hamad Bin Khalifa University (HBKU), Qatar Foundation, Doha, Qatar
17 E-mail: ypark@hbku.edu.qa
18

Competing Interests: The authors declare no competing interests.
Acknowledgements
We thank Dr. Reinhard Jahn for constructs and samples. We are deeply indebted to Dr. Kyong-Tai Kim
for technical assistanace. We thank Janarthanan Ponraj, Akshath Raghu Shetty from the Core Labs of
the Qatar Environment and Energy Research Institute (QEERI) for technical support. Thanks to Dr.
Ahmed Elalawy and Sarra Karrar from Widam Food Company for the arrangement of adrenal glands.
This work was supported by the grant from Qatar Biomedical Research Institute (Project Number SF
2019 004 to Y.P.).
27

28   **SUMMARY**
29
Cholesterol is essential for neuronal activity and function. Cholesterol depletion in the plasma membrane
impairs synaptic transmission. However, the molecular mechanisms by which cholesterol deficiency leads
to defects in vesicle fusion remain poorly understood. Here we show that cholesterol is required for Ca^{2+}-
dependent fusion using the *in-vitro* reconstitution of vesicle fusion, atomic force microscopy (AFM), and
amperometry to monitor exocytosis in chromaffin cells. Purified native vesicles were crucial for the
complete reconstitution of physiological Ca^{2+}-dependent fusion, whereas vesicle-mimicking liposomes
failed to reproduce the cholesterol effect. Intriguingly, cholesterol had no effect on membrane insertion of
synaptotagmin-1, a Ca^{2+} sensor for ultrafast fusion. Cholesterol stabilizes local membrane bending
induced by synaptotagmin-1 insertion, thereby lowering the energy barrier for Ca^{2+}-dependent fusion to
occur. Our data provide evidence that cholesterol depletion abolishes Ca^{2+}-dependent vesicle fusion by
disrupting synaptotagmin-1-induced membrane bending, and suggests that cholesterol is a master
regulator for Ca^{2+}-dependent fusion.
42

Cholesterol is a major component in cell membrane bilayers, and is essential for membrane structure and
fluidity. Brain is the most cholesterol-enriched organ and a human brain contains about 20–25% of the
body's cholesterol[1, 2]; this high density suggests that cholesterol has a critical function in the brain. Age-
related cholesterol reduction in the frontal and temporal cortices[3] results in loss of synaptic contacts,
changes in neuronal morphology, and reduced synaptic plasticity[4]. Cholesterol is associated with
neurogenesis, neurodevelopment, and synaptogenesis[5, 6]. Age-related cholesterol deficiency in the plasma
membrane leads to deficits in synaptic plasticity in mouse hippocampal neurons[7], and cholesterol
depletion by methyl-β-cyclodextrin (MCD) from the plasma membrane impairs neurotransmission and
neuronal activity, and thereby leads to synapse degeneration[8, 9].

The plasma membrane contains ~80% of total cellular cholesterol[10]. Cholesterol is capable of clustering
syntaxin-1A[11] so that soluble *N*-ethylmaleimide-sensitive factor attachment protein receptor (SNARE)
proteins become concentrated in cholesterol-enriched domains in the plasma membrane[12]. Depletion and
reduction of cholesterol level in the plasma membrane inhibit Ca^{2+}-dependent exocytosis of large dense-
core vesicles (LDCVs)[13] and cortical secretory vesicles from sea urchins[14], as well as synaptic vesicles in
hippocampal neurons[8, 15], cortical synaptosomes[16], ribbon synapses[17], and motor nerve terminals[18].
However, the molecular mechanisms by which cholesterol deficiency disrupts synaptic transmission and
induces neurodegeneration remain elusive and controversial.

Exocytosis is the process of neurotransmitter release through merging two lipid bilayers. It is mediated by
SNARE proteins[19, 20]. Neuronal SNARE proteins consist of Q-SNARE in the plasma membrane (syntaxin-
1 and SNAP-25) and R-SNARE in the vesicle membrane (synaptobrevin-2 or vesicle-associated
membrane protein-2 (VAMP-2))[19]. Synaptotagmin-1 is responsible for ultrafast Ca^{2+}-dependent
exocytosis. Synaptotagmin-1 contains two tandem C2-domains that coordinates Ca^{2+} binding; the Ca^{2+}-
bound C2 domain can be inserted to negatively charged anionic phospholipids by electrostatic
interaction[20]. In spite of intense investigation of synaptotagmin-1, the molecular mechanisms by which
synaptotagmin-1 mediates Ca^{2+}-dependent vesicle are still under debate[21].

Here we propose novel mechanisms by which cholesterol in the plasma membrane mediates
synaptotagmin-1-induced vesicle fusion. We used amperometry to monitor exocytosis in real time, and
performed *in-vitro* reconstitution of vesicle fusion including large dense-core vesicles (LDCVs) and
synaptic vesicles (SVs), and showed that cholesterol is required for Ca^{2+}-dependent vesicle fusion.
Importantly, vesicle-mimicking liposomes fail to reproduce the cholesterol effect, but purified native
vesicles, i.e., LDCVs and SVs, are crucial for the complete reconstitution of physiological Ca^{2+}-dependent
fusion. Membrane insertion of synaptotagmin-1 occurs regardless of cholesterol. Atomic force
microscopy (AFM) revealed that cholesterol stabilizes membrane bending and curvature induced by the
insertion of synaptotagmin-1, and therefore the membrane bending energy can lower the energy barrier to
fusion.

Results

Cholesterol depletion causes deficits in exocytosis.

We investigated whether local membrane curvature is observable *in-vivo* using primary chromaffin cells. Using electron microscopy (**Fig. 1a-f**) we observed the invagination of the plasma membrane into LDCVs, suggesting that the plasma membrane could be already curved before Ca^{2+} triggering. Given that the curved and invaginated membranes are enriched with cholesterol[22] and ~67% vesicles are docked to the invaginated plasma membranes (**Fig. 1e**), the membrane invagination to LDCVs we observed is likely to be cholesterol-enriched regions, where SNARE proteins are present[23, 24] (**Fig. 1a-e**).

To test the cholesterol effect on vesicle fusion, we used amperometry to monitor LDCV exocytosis in real time. Methyl-beta-cyclodextrin (MCD) depleted cholesterol from the plasma membrane, so treatment with MCD inhibited Ca^{2+}-dependent LDCV fusion (**Fig. 1g-i**).

Without cholesterol, no Ca^{2+}-dependent vesicle fusion.
To further study the molecular mechanisms by which cholesterol deficiency affects vesicle fusion, we applied a reconstitution system of vesicle fusion by using purified native vesicles such as LDCVs and SVs, as reported previously[25, 26, 27]. The plasma membrane-mimicking liposomes (PM-liposomes) contain the stabilized Q-SNARE complex (syntaxin-1A and SNAP-25A in a 1:1 molar ratio[28])(**Online Methods**).

We first tested the effect of cholesterol on basal fusion without Ca^{2+}. LDCV fusion with the PM-liposomes was readily observed (**Fig. 2a**), but the absence of cholesterol in the PM-liposomes reduced LDCV fusion by ~50% (**Fig. 2a,d**); the PM-liposomes contain either 25% or 0% cholesterol (Chol). The efficiency of the SNARE assembly and SNARE complex formation was examined using the light chain of tetanus toxin (TeNT), a protease specific for free VAMP-2; i.e. VAMP-2 in the assembled SNARE complexes is resistant to the cleavage by TeNT so that TeNT-resistant VAMP-2 represents the ternary SNARE complex formation[25]. After LDCV fusion with the PM-liposomes as a fusion assay in **Fig. 2a**, TeNT was added to cleave free VAMP-2 and quantify VAMP-2 engaged in the SNARE complex formation. Absence of cholesterol (0% Chol) in liposomes slightly reduced the ternary SNARE complex formation compared to liposomes containing cholesterol (25% Chol) (**Fig. 2b**); this result suggests that cholesterol facilitates basal fusion of LDCVs by increasing the efficiency of SNARE assembly. Cholesterol can enhance SNARE-complex formation, by inducing clustering of SNARE proteins in the plasma membrane[11, 12, 24].

Then we investigated the cholesterol effect on Ca^{2+}-dependent fusion. Although cholesterol controls Ca^{2+}-dependent neurotransmission and exocytosis in diverse cell types[8, 13, 14, 15, 16, 17, 18], the step at which vesicle fusion is impaired has not been determined. We used native vesicles to completely reconstitute vesicle fusion that reproduces Ca^{2+}-dependent vesicle fusion, correlating with the *in-vivo* data in a physiological ionic environment[25, 26, 27]. Addition of 100 µM free Ca^{2+} accelerated vesicle fusion in the presence of Mg^{2+}/ATP (**Fig. 2c,d**), where synaptotagmin-1 interacts only with PIP_2-containing membranes, but not SNARE proteins[25, 26]. VAMP-2_{1-96}, the soluble cytoplasmic region of VAMP-2, blocked vesicle fusion; this result supports SNARE-dependent vesicle fusion. Surprisingly, Ca^{2+}-evoked LDCV fusion was

completely abolished when cholesterol in the PM-liposomes was absent (**Fig. 2c,d**). The Q-SNARE
proteins were incorporated in the PM-liposomes independently of cholesterol (**Supplementary Fig. 1**).
We also confirmed that Ca^{2+}-dependent LDCV fusion did not occur without cholesterol (0% Chol) in the
PM-liposomes when the full-length syntaxin-1A and SNAP-25A binary acceptor complex (see **Online**
Methods) was included (**Supplementary Fig. 2**). High concentration of free Ca^{2+} failed to increase
vesicle fusion when the PM-liposomes contained no cholesterol (**Fig. 2e**) and cholesterol accelerated Ca^{2+}-
dependent LDCV fusion in a dose-dependent manner (**Fig. 2f**). Altogether, our results confirm that
cholesterol is required for Ca^{2+}-dependent vesicle fusion and that this *in-vitro* reconstitution reproduces
physiological exocytosis.

125 Cholesterol is not required for liposome-liposome fusion.
We further tested SVs purified from mice brain to determine whether cholesterol is essential for Ca^{2+}-
dependent SV fusion. Ca^{2+} failed to accelerate SV fusion in the absence of cholesterol in the PM-
liposomes (0% Chol) (**Fig. 3a,b**); this result is consistent with the cholesterol requirement for Ca^{2+}-
dependent LDCV fusion (**Fig. 2d**), however, Ca^{2+}-independent basal SV fusion was not affected. A unique
and major advantage of native vesicles for a fusion assay is that they maintain the native lipid and protein
diversity as well as the structural integrity of vesicles to mimic endogenous vesicle fusion. Instead of
purified native vesicles, vesicle-mimicking liposomes (V-liposomes) have been used for a fusion assay to
study the molecular mechanisms. We therefore tested V-liposomes that incorporate full length VAMP-2
and synaptotagmin-1 to determine whether the dependence on cholesterol for Ca^{2+}-dependent fusion can
be reproduced (**Fig. 3c,d**). Intriguingly, Ca^{2+}-dependent liposome fusion was slightly reduced but still
observable even in the absence of cholesterol in the PM-liposomes (**Fig. 3d**). These results indicate that
cholesterol is required for Ca^{2+}-dependent fusion of native vesicles, i.e., LDCV and SV, but not for
liposome-liposome fusion.

139 Membrane insertion of synaptotagmin-1 independently of cholesterol.
Cholesterol in the PM-liposomes is indispensable for Ca^{2+}-dependent vesicle fusion, so we investigated
whether the binding and insertion of synaptotagmin-1 to the PIP_2-containing membrane is affected by
cholesterol. The C2AB domain of synaptotagmin-1 (Syt_{97-421}) was labelled with Alexa Fluor 488 at S342C
as a donor, and the PM-liposomes (Lip., protein-free) were labelled with Rhodamine (Rho)-PE as an
acceptor (**Online Methods**). The C2AB binding to liposomes was monitored by FRET between the C2AB
domain (Alexa Fluor 488) and Rhodamine-labelled PM-liposomes (**Fig. 4a**). The C2AB domain of
synaptotagmin-1 could bind to both cholesterol-containing and cholesterol-free liposomes in response to
Ca^{2+}, and the Ca^{2+} titration for C2AB binding to the PM-liposomes (0% Chol) was comparable to that of
25% Chol-containing PM-liposomes; this result demonstrates Ca^{2+}-dependent C2AB binding to anionic
phospholipids regardless of cholesterol in liposomes (**Fig. 4a-c**). The PM-liposomes contain anionic
phospholipids including 10% PS, 4% PI, and 1% PIP_2, which provide complete coordination sites for the
Ca^{2+}-bound C2AB domain to interact with the membranes. (**Fig. 4a-c**).

To further assess membrane insertion of the C2AB domain, a fluorescence anisotropy measurement was
conducted to monitor the rotational mobility of the C2AB domain, labelled with Alexa Fluor 488 at 342C;
the PM-liposomes are label-free and protein-free (**Fig. 4d**). The C2AB insertion to the PM-liposomes

results in the increase of fluorescence anisotropy due to a reduction in the rotational mobility of the
membrane-bound C2AB domain (**Fig. 4e**). Ca^{2+} titration for membrane insertion of the C2AB domain
showed no difference between the two sets of PM-liposomes (0% vs 25% Chol) (**Fig. 4d-f**). The increases
of PIP_2 concentration in the PM-liposomes shifted Ca^{2+} titration curves to the left, which indicates an
increase in the Ca^{2+}-sensitivity of C2AB membrane binding (**Fig. 4g, Supplementary Fig. 3**). High PIP_2
concentration increases Ca^{2+} sensitivity for vesicle fusion[25]; this interaction implies that increase in the
magnitude of the negative electrostatic potential in the plasma membranes increases its attraction of Ca^{2+}-
bound synaptotagmin-1. Even 5% PIP_2 in the the PM-liposomes Ca^{2+} still failed to increase LDCV fusion
without cholesterol (**Fig. 4h**).

We also confirmed that the C2A and C2B domains bind to V-liposomes (0% PIP_2, 15% PS included), but
this interaction was completely disrupted at physiological ionic strength due to the charge-shielding effect
of Mg^{2+} and ATP (**Fig. 4i**)[25, 26]. Both the C2A and C2B domains are inserted to the plasma membrane to
drive vesicle fusion in a physiological ionic environment [25, 26]. Taken together, C2AB insertion into the
plasma membrane directly provides the driving force to lower the energy barrier for Ca^{2+}-triggered vesicle
fusion to occur. Cholesterol is only required for Ca^{2+}-dependent vesicle fusion, but is not essential for
Ca^{2+}-dependent C2AB membrane binding.

Membrane curvature is critical for Ca^{2+}-dependent fusion.
Despite the complete inhibition of Ca^{2+}-dependent vesicle fusion (**Fig. 2 and 3a,b**), membrane insertion
of the C2AB domain was still observable in the absence of cholesterol in liposomes (**Fig. 4**); this
observation strongly suggests that the downstream of membrane insertion of synaptotagmin-1 is disrupted.
Hydrophobic residues in the Ca^{2+}-binding loops of synaptotagmin-1 penetrate the inner leaflet of the
plasma membrane[29], and probably lead to local membrane bending and deformation[30, 31]. This local
membrane deformation might accelerate vesicle fusion by lowering the energy barrier[32]. Cholesterol also
regulates local membrane bending and deformation[33], so in the next experiments we examined whether
cholesterol has a critical function in membrane bending to trigger Ca^{2+}-dependent vesicle fusion.

To this end we used different PM-liposomes of different sizes: large (LUV) with 110-nm diameter and
small unilamellar vesicles (SUV) with 60-nm diameter (**Online Methods**, **Supplementary Fig. 5**). The
average diameter of LDCVs is 150 nm, ranging from 100 nm to 300 nm[27].We expected that the
synaptotagmin-1-induced changes in local membrane curvature and tension would be minimized in small
liposomes (**Fig. 5a,b**), because they are already highly curved[31]. Indeed, Ca^{2+}-dependent LDCV fusion
was dramatically lower when SUVs were used compared to LUVs (**Fig. 5c-e**). The curvature effect on
vesicle fusion was reproduced when the concentration of liposomes and the ratio of vesicle to liposome
were changed (**Supplementary Fig. 4**).

To further assess the effect of membrane curvature on vesicle fusion, we used SVs that had an average
diameter of 45 nm[34]. As expected, Ca^{2+}-dependent SV fusion with SUVs was completely impaired (**Fig.
5f,g**), whereas basal SV fusion was already saturated and augmented (**Fig. 5h**). High membrane tension
and curvature elevate basal fusion, but decrease Ca^{2+}-dependent vesicle fusion, because highly curved-

membranes are likely to fuse without Ca^{2+} and show less change in membrane tension by membrane
insertion of synaptotagmin-1.

Cholesterol stabilizes membrane bending and deformation.

Next, we performed AFM to show that cholesterol regulates local membrane bending induced by
synaptotagmin-1 (**Fig. 6**). AFM allows direct observation of membrane deformation at nanometer
resolution. The supported lipid bilayer (SLB) prepared by the PM-liposomes were planar and continuous
with 3~4 Å of background noise from the Z-position (**Fig. 6a**). The C2AB domain alone had no effect on
SLB (**Fig. 6b**), but Ca^{2+}-bound C2AB domain caused significant changes in membrane deformation with
15~20 Å invagination when 25% cholesterol was present (**Fig. 6c**). However, Ca^{2+}-bound C2AB domain
showed a weak effect on membrane deformation in the absence of cholesterol (**Fig. 6d-g**). These AFM
data provide direct evidence that cholesterol stabilizes membrane bending caused by membrane insertion
of synaptotagmin-1.

Discussion

Cholesterol regulates vesicle fusion as follows. First, it causes clustering of SNARE proteins in the plasma membrane. This clustering could increase the efficiency of membrane fusion[23, 24]. Cholesterol in the PM-liposomes slightly increased the formation of SNARE complexes, and thus increased LDCV fusion (**Fig. 2a,b,d**). Second, cholesterol regulates the physical structure, fluidity, tension, and thickness of lipid membranes[35]. Cholesterol causes negative membrane curvature that might facilitate vesicle fusion[13, 35]. Cholesterol also contributes to vesicle fusion by stabilizing fusion pores[33, 35, 36, 37]. The plasma membrane deformations occur prior to vesicle fusion, and pre-fusion membrane curvature changes can be observed[38]. Here, our data provide a novel model in which cholesterol mediates Ca^{2+}-dependent vesicle fusion by stabilizing local membrane bending caused by synaptotagmin-1 insertion. The curved plasma membrane has high bending energy, which can be released to drive fusion with vesicle membranes[30, 31, 39]. Thus, high membrane-bending energy stabilized by cholesterol lowers the energy barrier for vesicle fusion, thereby triggering Ca^{2+}-dependent vesicle fusion (**Fig. 7a,b**).

The C2AB domain of synaptotagmin-1 is inserted into the plasma membrane[29] and the membrane-binding energy of the C2AB domain is ~18 k_BT;[40] the binding leads to local membrane bending and deformation[30, 31]. Membrane bending by synaptotagmin-1 could accelerate fusion by reducing the energy barrier. However, our data indicate that membrane bending by synaptotagmin-1 seems not enough to overcome the threshold of the energy barrier for fusion (**Fig. 6d-g**), when cholesterol is absent in the plasma membrane. Cholesterol augments and potentiates the plasma membrane deformation and bending caused by synaptotagmin-1 insertion (**Fig. 6g**). This significant membrane bending energy can drive the membrane fusion (**Fig. 7a,b**).

Our *in-vitro* reconstitution of vesicle fusion has an advantage of using purified native vesicles, i.e., LDCVs and SVs, and reproduces the physiological cholesterol effect on vesicle fusion, whereas V-liposomes fail (**Fig. 3c,d**). V-liposomes are independent of cholesterol for Ca^{2+}-dependent fusion, in contrast to native vesicles (**Fig. 2,3**). V-liposomes show less efficiency of Ca^{2+}-dependent vesicle fusion and have limitations to replace native vesicles for a fusion assay. Native vesicles differ from V-liposomes in lipid composition, protein density, contents diversity, and physical property; therefore the structural integrity of native vesicles allows them to mimic endogenous vesicle fusion in the *in-vitro* reconstitution. Native vesicles have high membrane rigidity compared to V-liposomes, and therefore the bending is likely to be enough to induce membrane fusion with V-liposomes, whereas native vesicles seem to require higher membrane bending energy than V-liposomes do, in order to overcome the barrier for fusion due to membrane fluidity and rigidity. The cause of the difference between native vesicles and liposomes for fusion efficiency, and the reproduction of physiological vesicle fusion remain topics for further study.

The molecular mechanisms of synaptotagmin-1 to trigger Ca^{2+}-dependent fusion remain controversial; at least six different competing models have been proposed[21]. Synaptotagmin-1 mediates Ca^{2+}-dependent fusion by the electrostatic interaction, so several different synaptotagmin-1 models have been proposed, depending on the ionic environment[26]. Here we confirmed the cholesterol effect on vesicle fusion in a physiological ionic environment, i.e., normal ionic strength with Mg^{2+}/ATP. Both the C2A and C2B

domains of synaptotagmin-1 are inserted into the plasma membrane (**Fig. 4i**) without interacting with
SNARE proteins[26], and therefore lead to membrane bending (**Fig. 6**). Ca^{2+} fails to trigger fusion in the
absence of cholesterol despite the proteins, both SNARE assembly and synaptotagmin-1 insertion, being
fully active and functional (**Fig. 2, 4**). Cholesterol has an important function in Ca^{2+}-dependent fusion, as
a master regulator by stabilizing membrane deformation and curvature caused by synaptotagmin-1
insertion.

Age-related cholesterol reduction is linked to reduced synaptic activity, and defects in synaptic
transmission by cholesterol deficiency could result in neurodegeneration[4]. Our data explains the molecular
mechanisms how cholesterol contributes to synaptic transmission and neuronal function and may pave the
way for development of studies to explore to treatment of neurodegenerative and neurodevelopmental
disorders by optimizing cholesterol levels in the plasma membrane.

 Figure legends

Figure 1
Cholesterol depletion inhibits LDCV exocytosis in chromaffin cells. (**a**) Transmission electron
microscope (TEM) image of chromaffin cells showing invagination of the plasma membrane into LDCVs.
Scale, 500 nm. (**b-e**) Magnified TEM images of LDCVs. (**f**) LDCVs docking to the plasma membranes
either flat or curved is presented as a percentage (total 36 LDCVs docked to the plasma membranes from
three independent experiments). (**g-i**) LDCV exocytosis in chromaffin cells measured by amperometry.
MCD (10 mM, 2 h, 37°C) reduced LDCV exocytosis of chromaffin cells (**h**). (**g,h**) Shown are typical
amperometric traces upon DMPP stimulations finally evoking voltage-gated calcium channel for 20 sec.
(**i**) Data are means ± SD from 5 independent experiments.

Figure 2
Cholesterol is required for Ca^{2+}-triggered vesicle fusion. (**a**) *In-vitro* reconstitution of large dense-core
vesicle (LDCV) fusion using a lipid-mixing assay. Purified LDCVs were incubated with PM-liposomes
that incorporate the stabilized Q-SNARE complex of syntaxin-1A/SNAP-25A (Syx/SN25) in a 1:1 molar
ratio (**Online Methods**). Cholesterol (Chol) was included either 25% or 0% in the PM-liposomes. For
clarity, only endogenous VAMP-2 and synaptotagmin-1 of native LDCVs are shown. (**b**) Formation of
the ternary SNARE complex after vesicle fusion regardless of cholesterol in the PM-liposomess. LDCVs
were incubated with liposomes for 20 min, then treated with Tetanus neurotoxin (TeNT). TeNT-resistant
VAMP-2 indicates the ternary SNARE complex formation in SDS-PAGE (**Online Methods**). Boiling at
95°C disrupts the ternary SNARE complex so that VAMP-2 migrates to its size. Data are mean ± SD from
3 independent experiments (n = 3). (**c,d**) Dependence on cholesterol for Ca^{2+}-dependent vesicle fusion.
Addition of 100 µM free Ca^{2+} provoked fusion of LDCV with the PM-liposomes. Preincubation of the
PM-liposomes with VAMP-2$_{1-96}$ caused competitive inhibition that inhibited SNARE-mediated fusion.
Data are mean ± SD (n = 6). (**e**) Free Ca^{2+} was increased to 300 µM, when LDCVs fused with the PM-
liposomes (0% Chol). (**f**) Liposomes contained either 15% or 25% of Chol. Lipid composition of the PM-
liposomess: 45% PC, 15% PE, 10% PS, 25% Chol, 4% PI, and 1% PI(4,5)P$_2$. When Chol was reduced,
PC contents were adjusted accordingly. Physiological ionic strength and 1 mM MgCl$_2$/3 mM ATP were
included in all experiments.
285

Figure 3

Cholesterol is not required for liposome-liposome fusion. (**a,b**) Synaptic vesicle (SV) fusion with the

PM-liposomes that contain either 25% (**a**) or 0% (**b**) Chol. Only endogenous VAMP-2 and

synaptotagmin-1 of native SVs are presented. (**c,d**) Instead of native LDCVs or SVs, V-liposomes that

contain the full-length synaptotagmin-1 and VAMP-2 were incubated with PM-liposomes that contained

either 25% (**c**) or 0% (**d**) Chol. Lipid composition of the PM-liposomes is described in **Figure 2**. Lipid

composition of V-liposomes: 55% PC, 20% PE, 15% PS, and 10% Chol.

Figure 4

Membrane insertion of the C2AB domain of synaptotagmin-1 in the presence and absence of cholesterol. (**a-c**) Insertion of the C2AB domain of synaptotagmin-1 was monitored using FRET in which the C2AB domain (Syt-1_{97-421}) was labelled with Alexa Fluor 488 at S342C (green dots) as a donor, and the PM-liposomes (Lip.) were labelled with Rhodamine (Rho)-PE (red) as an acceptor (**Online Methods**). The PM-liposomes (protein-free) contained either 25% (**b**) or 0% Chol. (**c**) C2AB binding is presented as a percentage of maximum C2AB binding induced by 1 mM free Ca^{2+}. Data are mean $\pm$ SD from 3~5 independent experiments. (**d-f**) Binding of the C2AB domain to the PM-liposomes (protein-free; lipid composition as in **a-c** without Rho-PE) was monitored using fluorescence anisotropy. (**f**) Dose-response curve of Ca^{2+}-dependent C2AB binding to the PM-liposomes that contain either 25% or 0% Chol. Data are mean $\pm$ SD (n = 3~4). (**g**) FRET was conducted to monitor C2AB binding to liposomes as in **Fig. 4a-c**. PIP_2 was incorporated in the PM-liposomes (25% cholesterol included, protein-free). (**g**) High PIP_2 concentration increased Ca^{2+} sensitivity for C2AB binding to membranes. Data are mean $\pm$ SD (n = 3~5). (**h**) LDCV fusion with the PM-liposomes that contain 5% PIP_2 concentration and no Chol. (**i**) C2AB binding to V-liposomes was monitored using a tryptophan-dansyl FRET pair. Neither the C2A nor C2B domain binds to V-liposomes (no PIP_2) by 1 mM Ca^{2+} in the presence of 1 mM $MgCl_2$/3 mM ATP. Data are mean $\pm$ SD (n = 3).

Figure 5

High curvature strain of membrane increases basal fusion but decreases Ca^{2+}-triggered fusion. (**a,b**)

Schematic diagram showing membrane bending by membrane insertion of the C2AB domain. Large

unilamellar vesicles (LUVs), 110 nm in diameter; small unilamellar vesicles (SUVs), 60 nm in diameter.

The SUV is already highly curved. (**c,d**) Ca^{2+}-dependent LDCV fusion with the PM-liposomes, either

LUV or SUV. (**e**) Dose-response curve of Ca^{2+} on LDCV fusion with either LUV or SUV. (**f-h**) SV fusion

with the PM-liposomes, either LUV or SUV. Mouse SVs are ~45 nm in diameter. Data in **e,h** are mean ±

SD (n = 3 ~ 7).

Figure 6
Cholesterol stabilizes membrane bending induced by the C2AB domain. AFM tapping mode images
of supported lipid bilayer (SLB) prepared by the PM-liposomes. (**a-c**) AFM images and corresponding
height profiles of membrane deformation and bending for black lines across sections of the images in the
presence of 25% Chol. (**b**) C2AB domain of synaptotagmin-1 (0.5 µM) incubated with SLB. (**c**) Free 500
µM Ca^{2+} applied. (**d-f**) No cholesterol in SLB. (**g**) The height of membrane deformation measured by
AFM. Data are mean ± SEM (n = 18 ~ 37).
328
329

Figure 7
A schematic diagram summarizing synaptotagmin-1-induced membrane bending for Ca^{2+}-
dependent vesicle fusion. (**a**) Cholesterol stabilizes the plasma membrane deformation and bending
caused by synaptotagmin-1 insertion. (**b**) Energy landscapes of vesicle fusion by synaptotagmin-1.
Synaptotagmin-1-induced membrane bending enhanced by cholesterol generates the significant
membrane bending energy that can drive the membrane fusion by lowering the energy barriers to promote
the fusion intermediate.

Supplementary Figure 1
Incorporation of the Q-SNARE complex in liposomes. (**a,b**) Incorporation of the Q-SNARE complex
in liposomes in the presence and absence of cholesterol was tested using a flotation assay. (**a**) Schematic
diagram showing that liposomes float up through the gradient due to their buoyancy, whereas free proteins
remain in the bottom of the gradient. (**b**) SNARE proteins stained by coomassie blue dyes. The Q-SNARE
proteins SNAP-25A (no cysteine, cysteines are replaced by alanines) and syntaxin-1A (aa 183–288) in a
1:1 ratio by the C-terminal VAMP-2 fragment (aa 49–96), are incorporated in liposomes independently
of cholesterol.

Supplementary Figure 2

Cholesterol requirement for Ca^{2+}-evoked LDCV fusion. (**a**) No Ca^{2+}-dependent LDCV fusion was

observed when the Q-SNARE complex consisting of the full-length syntaxin-1A (1-288) and SNAP-25A

(no cysteine, cysteines are replaced by alanines) was included in liposomes with (25% Chol) and without

cholesterol (0% Chol). The binary Q-SNARE complex shows slow fusion rate and relatively low fusion

activity, as expected[25].

Supplementary Figure 3
Monitoring C2AB binding to liposomes using FRET measurement. (**a**) A schematic diagram of C2AB
domain binding to liposomes. Insertion of the C2AB domain of synaptotagmin-1 was monitored using
FRET in the C2AB domain (Syt-1$_{97\text{-}421}$) was labelled with Alexa Fluor 488 at S342C (green dots) as a
donor, and the PM-liposomes (Lip.) were labelled with Rhodamine (Rho)-PE (red) as an acceptor (**Online**
Methods). Lipid composition of the PM-liposomes: 45% PC, 13.5% PE, 10% PS, 25% Chol, 4% PI, 1%
PI(4,5)P$_2$, and 1.5% Rho-PE. When PI(4,5)P$_2$ was changed, PI contents were adjusted accordingly. (**b**)
PI(4,5)P$_2$ contents were increased to 5% in liposomes and C2AB binding was saturated with 0.5%
PI(4,5)P$_2$ in liposomes. Maximum C2AB binding induced by 1 mM free Ca^{2+} in the presence of 1 mM
MgCl$_2$/3 mM ATP.
364

Supplementary Figure 4
Ca^{2+}-dependent vesicle fusion depends on liposome membrane curvature. Large unilamellar vesicles
(LUVs), 110 nm in diameter; small unilamellar vesicles (SUVs), 60 nm in diameter. SUV is already highly
curved. A lipid-mixing assay to monitor Ca^{2+}-dependent LDCV fusion with the PM-liposomes, either
LUV or SUV. Equal amount of LDCVs (5 µL) was incubated to induce fusion with either 5 µL (**a,b**) or
20 µL (**c,d**) LUV or SUV in 1 mL fusion buffer; 120 mM K-glutamate, 20 mM K-acetate, 20 mM HEPES-
KOH (pH 7.4), 1 mM $MgCl_2$, and 3 mM ATP. 100 µM free Ca^{2+} in the presence of 1 mM $MgCl_2$/3 mM
ATP.

Supplementary Figure 5
The size distribution of proteoliposomes. The size distribution of proteoliposomes was determined using
dynamic light scattering (DLS). LUVs (110 nm in diameter) and SUVs (60 nm in diameter) were prepared
by extrusion (**Online Methods**).

References

1. Dietschy JM, Turley SD. Cholesterol metabolism in the brain. *Curr Opin Lipidol* **12**, 105-112 (2001).

2. Bjorkhem I, Meaney S. Brain cholesterol: long secret life behind a barrier. *Arterioscler Thromb Vasc Biol* **24**, 806-815 (2004).

3. Svennerholm L, Bostrom K, Jungbjer B, Olsson L. Membrane lipids of adult human brain: lipid composition of frontal and temporal lobe in subjects of age 20 to 100 years. *Journal of neurochemistry* **63**, 1802-1811 (1994).

4. Martin M, Dotti CG, Ledesma MD. Brain cholesterol in normal and pathological aging. *Biochim Biophys Acta* **1801**, 934-944 (2010).

5. Pfrieger FW. Cholesterol homeostasis and function in neurons of the central nervous system. *Cell Mol Life Sci* **60**, 1158-1171 (2003).

6. Cartocci V, Servadio M, Trezza V, Pallottini V. Can Cholesterol Metabolism Modulation Affect Brain Function and Behavior? *J Cell Physiol* **232**, 281-286 (2017).

7. Martin MG*, et al.* Constitutive hippocampal cholesterol loss underlies poor cognition in old rodents. *EMBO Mol Med* **6**, 902-917 (2014).

8. Linetti A*, et al.* Cholesterol reduction impairs exocytosis of synaptic vesicles. *J Cell Sci* **123**, 595-605 (2010).

9. Liu Q*, et al.* Neuronal LRP1 knockout in adult mice leads to impaired brain lipid metabolism and progressive, age-dependent synapse loss and neurodegeneration. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **30**, 17068-17078 (2010).

10. Lange Y, Steck TL. Active membrane cholesterol as a physiological effector. *Chem Phys Lipids* **199**, 74-93 (2016).

11. Murray DH, Tamm LK. Clustering of syntaxin-1A in model membranes is modulated by phosphatidylinositol 4,5-bisphosphate and cholesterol. *Biochemistry* **48**, 4617-4625 (2009).

12. Sieber JJ*, et al.* Anatomy and dynamics of a supramolecular membrane protein cluster. *Science* **317**, 1072-1076 (2007).

13. Zhang J, Xue R, Ong WY, Chen P. Roles of cholesterol in vesicle fusion and motion. *Biophys J* **97**, 1371-1380 (2009).

424
425   14.   Churchward MA, Rogasevskaia T, Hofgen J, Bau J, Coorssen JR. Cholesterol facilitates the
426   native mechanism of Ca2+-triggered membrane fusion. *J Cell Sci* **118**, 4833-4848 (2005).
427
428   15.   Mailman T, Hariharan M, Karten B. Inhibition of neuronal cholesterol biosynthesis with
429   lovastatin leads to impaired synaptic vesicle release even in the presence of lipoproteins or
430   geranylgeraniol. *Journal of neurochemistry* **119**, 1002-1015 (2011).
431
432   16.   Teixeira G, Vieira LB, Gomez MV, Guatimosim C. Cholesterol as a key player in the balance of
433   evoked and spontaneous glutamate release in rat brain cortical synaptosomes. *Neurochemistry*
434   *international* **61**, 1151-1159 (2012).
435
436   17.   Mercer AJ, Szalewski RJ, Jackman SL, Van Hook MJ, Thoreson WB. Regulation of presynaptic
437   strength by controlling Ca2+ channel mobility: effects of cholesterol depletion on release at the
438   cone ribbon synapse. *Journal of neurophysiology* **107**, 3468-3478 (2012).
439
440   18.   Tarakanova OI, Petrov AM, Zefirov AL. The role of membrane cholesterol in neurotransmitter
441   release from motor nerve terminals. *Doklady biological sciences : proceedings of the Academy of*
442   *Sciences of the USSR, Biological sciences sections / translated from Russian* **438**, 138-140
443   (2011).
444
445   19.   Jahn R, Scheller RH. SNAREs--engines for membrane fusion. *Nat Rev Mol Cell Biol* **7**, 631-643
446   (2006).
447
448   20.   Brunger AT, Choi UB, Lai Y, Leitz J, Zhou Q. Molecular Mechanisms of Fast Neurotransmitter
449   Release. *Annu Rev Biophys* **47**, 469-497 (2018).
450
451   21.   Park Y, Ryu JK. Models of synaptotagmin-1 to trigger Ca(2+) -dependent vesicle fusion. *FEBS*
452   *Lett* **592**, 3480-3492 (2018).
453
454   22.   Anderson RG. The caveolae membrane system. *Annual review of biochemistry* **67**, 199-225
455   (1998).
456
457   23.   Chamberlain LH, Burgoyne RD, Gould GW. SNARE proteins are highly enriched in lipid rafts
458   in PC12 cells: implications for the spatial control of exocytosis. *Proceedings of the National*
459   *Academy of Sciences of the United States of America* **98**, 5619-5624 (2001).
460
461   24.   Lang T, *et al.* SNAREs are concentrated in cholesterol-dependent clusters that define docking
462   and fusion sites for exocytosis. *EMBO J* **20**, 2202-2213 (2001).
463
464   25.   Park Y, *et al.* Controlling synaptotagmin activity by electrostatic screening. *Nature structural &*
465   *molecular biology* **19**, 991-997 (2012).
466
467   26.   Park Y, *et al.* Synaptotagmin-1 binds to PIP(2)-containing membrane but not to SNAREs at
468   physiological ionic strength. *Nature structural & molecular biology* **22**, 815-823 (2015).
469

470 27. Birinci Y, Preobraschenski J, Ganzella M, Jahn R, Park Y. Isolation of large dense-core vesicles
471   from bovine adrenal medulla for functional studies. *Sci Rep* **10**, 7540 (2020).
472
473 28. Pobbati AV, Stein A, Fasshauer D. N- to C-terminal SNARE complex assembly promotes rapid
474   membrane fusion. *Science* **313**, 673-676 (2006).
475
476 29. Herrick DZ, Sterbling S, Rasch KA, Hinderliter A, Cafiso DS. Position of synaptotagmin I at the
477   membrane interface: cooperative interactions of tandem C2 domains. *Biochemistry* **45**, 9668-
478   9674 (2006).
479
480 30. Martens S, Kozlov MM, McMahon HT. How synaptotagmin promotes membrane fusion.
481   *Science* **316**, 1205-1208 (2007).
482
483 31. Hui E, Johnson CP, Yao J, Dunning FM, Chapman ER. Synaptotagmin-mediated bending of the
484   target membrane is a critical step in Ca(2+)-regulated fusion. *Cell* **138**, 709-721 (2009).
485
486 32. Kozlov MM, Chernomordik LV. The protein coat in membrane fusion: lessons from fission.
487   *Traffic* **3**, 256-267 (2002).
488
489 33. Ivankin A, Kuzmenko I, Gidalevitz D. Cholesterol mediates membrane curvature during fusion
490   events. *Phys Rev Lett* **108**, 238103 (2012).
491
492 34. Takamori S*, et al.* Molecular anatomy of a trafficking organelle. *Cell* **127**, 831-846 (2006).
493
494 35. Yang ST, Kreutzberger AJB, Lee J, Kiessling V, Tamm LK. The role of cholesterol in
495   membrane fusion. *Chem Phys Lipids* **199**, 136-143 (2016).
496
497 36. Kreutzberger AJ, Kiessling V, Tamm LK. High cholesterol obviates a prolonged hemifusion
498   intermediate in fast SNARE-mediated membrane fusion. *Biophys J* **109**, 319-329 (2015).
499
500 37. Wu L, Courtney KC, Chapman ER. Cholesterol stabilizes recombinant exocytic fusion pores by
501   altering membrane bending rigidity. *Biophys J* **120**, 1367-1377 (2021).
502
503 38. Anantharam A, Axelrod D, Holz RW. Real-time imaging of plasma membrane deformations
504   reveals pre-fusion membrane curvature changes and a role for dynamin in the regulation of
505   fusion pore expansion. *Journal of neurochemistry* **122**, 661-671 (2012).
506
507 39. Zhang Z, Jackson MB. Membrane bending energy and fusion pore kinetics in Ca(2+)-triggered
508   exocytosis. *Biophys J* **98**, 2524-2534 (2010).
509
510 40. Gruget C*, et al.* Synaptotagmin-1 membrane binding is driven by the C2B domain and assisted
511   cooperatively by the C2A domain. *Sci Rep* **10**, 18011 (2020).
512
513 41. Ahmed S, Holt M, Riedel D, Jahn R. Small-scale isolation of synaptic vesicles from mammalian
514   brain. *Nature protocols* **8**, 998-1009 (2013).
515

516 42. Radhakrishnan A, Stein A, Jahn R, Fasshauer D. The Ca2+ affinity of synaptotagmin 1 is
517   markedly increased by a specific interaction of its C2B domain with phosphatidylinositol 4,5-
518   bisphosphate. *J Biol Chem* **284**, 25749-25760 (2009).
519
520 43. Nalefski EA, Falke JJ. Use of fluorescence resonance energy transfer to monitor Ca(2+)-
521   triggered membrane docking of C2 domains. *Methods in molecular biology* **172**, 295-303 (2002).
522
523 44. Park YS, *et al.* Involvement of protein kinase C-epsilon in activity-dependent potentiation of
524   large dense-core vesicle exocytosis in chromaffin cells. *The Journal of neuroscience : the official*
525   *journal of the Society for Neuroscience* **26**, 8999-9005 (2006).
526
527 45. Park Y, *et al.* alpha-SNAP interferes with the zippering of the SNARE protein membrane fusion
528   machinery. *J Biol Chem* **289**, 16326-16335 (2014).
529
530
531

Online Methods

Purification of large dense-core vesicles (LDCVs) and synaptic vesicles (SVs). LDCVs, also known
as chromaffin granules, were purified from bovine adrenal medullae using continuous sucrose gradient
and resuspended with fusion buffer containing 120 mM K-glutamate, 20 mM K-acetate, and 20 mM
HEPES.KOH, pH 7.4, as elsewhere[27]. SV from mouse brains were purified as described elsewhere[41].
Briefly, mice brains were homogenized in homogenization buffer supplemented with protease inhibitors,
using a glass-Teflon homogenizer. The homogenate was centrifuged for 10 min at 1,000g and the resulting
supernatant was further centrifuged for 15 min at 15,000g. The synaptosome pellet was lysed by adding
ice-cold water, followed by centrifugation for 25 min at 48,000g. The resulting supernatant was overlaid
onto a 0.7 M sucrose cushion and centrifuged for 1 h at 133,000g. The pellet was resuspended in fusion
buffer (120 mM K-glutamate, 20 mM K-acetate, 20 mM HEPES.KOH, pH 7.4).

Protein purification. All SNARE and the C2AB domain of synaptotagmin-1 constructs based on rat
sequences were expressed in *E. coli* strain BL21 (DE3) and purified by Ni^{2+}-NTA affinity chromatography
followed by ion-exchange chromatography as described elsewhere[25, 26]. The stabilized Q-SNARE
complex consisting syntaxin-1A (aa 183–288), SNAP-25A (no cysteine, cysteines replaced by alanines)
in a 1:1 ratio by the C-terminal VAMP-2 fragment (aa 49–96) was purified as described earlier[28]. The
binary Q-SNARE complex containing the full-length syntaxin-1A (1-288) and SNAP-25A (no cysteine,
cysteines replaced by alanines) was expressed using co-transformation[25]. The full-length VAMP-2,
soluble cytoplasmic region of VAMP-2 (VAMP-2_{1-96}), full-length synaptotagmin-1, C2AB domain of
synaptotagmin-1 (aa 97-421), C2A domain (aa 96-262), C2B domain (aa 248-421), and C2ab mutant
(D178A, D230A, D232A, D309A, D363A, D365A) were purified by Mono S column (GE Healthcare,
Piscataway, NJ) as described previously[42]. The stabilized Q-SNARE complex and the syntaxin-1A/SNAP-
25A binary SNARE complex were purified by Ni^{2+}-NTA affinity chromatography followed by ion-
exchange chromatography on a Mono Q column (GE Healthcare, Piscataway, NJ) in the presence of 50
mM n-octyl-β-D-glucoside (OG)[25]. The point mutated C2AB domain (S342C) was labelled with Alexa
Fluor 488 C5 maleimide (C2AB^{A488})[42].

Lipid composition of liposomes. Lipid composition (molar percentages) of the PM-liposomes that
contain the Q-SNARE complex consists of 45% PC (L-α-phosphatidylcholine), 15% PE (L-α-
phosphatidylethanolamine), 10% PS (L-α-phosphatidylserine), 25% Chol (cholesterol), 4% PI (L-α-
phosphatidylinositol), and 1% PI(4,5)P$_2$. When cholesterol was excluded (0% Chol), PC contents were
accordingly adjusted. In case of changing PI(4,5)P$_2$ concentration, PI contents were accordingly adjusted.
VAMP-2/synaptotagmin-1-containing V-liposomes are composed of 55% PC, 20% PE, 15% PS, and 10%
Chol. For FRET-based lipid-mixing assays, 1.5% 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-(7-
nitrobenz-2-oxa-1,3-diazol-4-yl (NBD-DOPE) and 1.5% 1,2-dioleoyl-sn-glycero-3-
phosphoethanolamine-N-lissamine rhodamine B sulfonyl ammonium salt (Rhodamine-DOPE) were
incorporated in the PM-liposomes (accordingly 12% unlabeled PE) as a donor and an acceptor dye,
respectively. For FRET measurement using C2AB^{A488}, 1.5% Rhodamine-DOPE was included in the PM-

liposomes (protein-free). In the case of FRET for tryptophan, 5% N-(5-dimethylaminonaphthalene-1-
dulfonyl)-1,2-dihexadecanoyl-sn-glycero-3-phosphoethanolamine, triethylammonium salt (dansyl-
DHPE) was incorporated in the PM-liposomes (protein-free). All lipids were from Avanti Polar lipids
except dansyl-DHPE (Invitrogen).
576
Preparation of proteoliposomes. Incorporation of the stabilized and binary Q-SNARE complex into
large unilamellar vesicles (LUVs, 110 nm in diameter) was achieved by OG-mediated reconstitution,
called the direct method, i.e. incorporation of proteins into preformed liposomes[25, 26]. Briefly, lipids
dissolved in a 2:1 chloroform-methanol solvent were mixed according to lipid composition. The solvent
was removed using a rotary evaporator (which generated lipid film on a glass flask), then lipids were
resuspended in 1.5 mL diethyl ether and 0.5 mL buffer containing 150 mM KCl and 20 mM HEPES/KOH
pH 7.4. After sonication on ice (3 x 45 s), multilamellar vesicles were prepared by reverse-phase
evaporation using a rotary evaporator as diethyl ether was removed. Multilamellar vesicles (0.5 mL) were
then extruded using polycarbonate membranes of pore size 100 nm (Avanti Polar lipids) to give uniformly-
distributed LUVs with the average diameter of 110 nm (**Supplementary Fig. 5**). After the preformed
LUVs had been prepared, SNARE proteins or the full-length VAMP-2/synaptotagmin-1 were
incorporated into them using OG, a mild non-ionic detergent, then the OG was removed by dialysis
overnight in 1 L buffer containing 150 mM KCl and 20 mM HEPES/KOH pH 7.4 together with 2 g SM-
2 adsorbent beads.
To make small unilamellar vesicles (SUVs) using the direct method, 110-nm LUVs produced as described
above extruded through polycarbonate membranes with 50-nm pore size (yielding SUV that had average
diameter of 60 nm, **Supplementary Fig. 5**). After preparing preformed SUVs, protein incorporation was
completed by OG as described for LUV. The size distribution of proteoliposomes was determined using
dynamic light scattering (DLS) (**Supplementary Fig. 5**). Proteoliposomes have protein-to-lipid molar
ratio of 1:500 (n/n).
597
Vesicle fusion assay. A FRET-based lipid-mixing assay was applied to monitor vesicle fusion *in vitro*[25,
26]. LDCV or SV fusion reactions were performed at 37°C in 1 mL fusion buffer containing 120 mM K-
glutamate, 20 mM K-acetate, 20 mM HEPES-KOH (pH 7.4), 1 mM $MgCl_2$, and 3 mM ATP. ATP should
be made freshly before experiments, because ATP is easily destroyed by freezing and thawing. Free Ca^{2+}
concentration in the presence of ATP and Mg^{2+} was calibrated using Maxchelator simulation program.
The fluorescence dequenching signal was measured using Fluoromax (Horiba Jobin Yvon) with
wavelengths of 460 nm for excitation and 538 nm for emission. Fluorescence values were normalized as
a percentage of maximum donor fluorescence (total fluorescence) after addition of 0.1% Triton X-100 at
the end of experiments.
607
Fluorescence resonance energy transfer (FRET). The C2AB domain of synaptotagmin-1 (30 nM,
S342C) was labeled with Alexa Fluor 488, a donor dye. C2AB fragment was engineered to contain a
single Cys residue (S342C) and labelled with Alexa Fluor 488. 1.5% Rhodamine-DOPE (Rho-PE),
incorporated in the PM-liposomes (protein-free), was used as an acceptor. Unless otherwise stated,

liposomes were LUVs prepared by the direct method. Donor fluorescence signal was measured at 37°C
using Fluoromax (Horiba Jobin Yvon) with wavelengths of 488 nm for excitation and 516 nm for emission
in 1 mL fusion buffer containing 120 mM K-glutamate, 20 mM K-acetate, 20 mM HEPES-KOH (pH 7.4),
1 mM $MgCl_2$, and 3 mM ATP. FRET was normalized as net changes of donor fluorescence intensity and
C2AB binding was presented as percentage of maximum C2AB binding induced by 1 mM Ca^{2+}. In
Supplementary Figure 3, FRET was normalized as F/F_0, where F_0 represents the initial value of the
donor fluorescence intensity.

Liposome binding of the C2AB domain was also monitored using the tryptophan-dansyl FRET
pair as a donor-acceptor dye in which dansyl-DHPE incorporated in liposomes leads to quenching of
fluorescence emitted from tryptophan of the C2AB domain[43]. Then1 μM C2AB, 3 μM C2A, or 3 μM C2B
was incubated with V-liposomes (protein-free). Donor fluorescence signal was measured at 37°C using
Fluoromax (Horiba Jobin Yvon) with wavelengths of 295 nm for excitation and 350 nm for emission in 1
mL fusion buffer containing 120 mM K-glutamate, 20 mM K-acetate, 20 mM HEPES-KOH (pH 7.4), 1
mM $MgCl_2$, and 3 mM ATP. FRET monitoring C2AB binding was normalized as a percentage of $(F_0-$
$F)/F_0$.

Fluorescence anisotropy measurements. The C2AB fragment (20 nM, S342C) were labelled with Alexa
Fluor 488[42]. Anisotropy was measured in a Fluorolog (Horiba Jobin Yvon) at 37°C in 1 ml of buffer
containing 120 mM K-glutamate, 20 mM K-acetate, and 20 mM HEPES-KOH (pH 7.4), 1 mM $MgCl_2$,
and 3 mM ATP. Excitation wavelength was 495 nm, and emission was measured at 520 nm. Lipid
composition of the PM-liposomes (protein-free) was identical to those used in a fusion assay except
labelled PE (45% PC, 15% PE, 10% PS, 25% Chol, 4% PI, and 1% PIP2). In the case of 0% Chol, PC
contents were adjusted accordingly (70% PC).

Ternary SNARE complex formation assay. Tetanus neurotoxin (TeNT) degrades free VAMP-2
whereas VAMP-2, assembled in the ternary SNARE complex, is resistant to TeNT[25]. After incubation of
LDCVs with the PM-liposomes (0% or 25% Chol) that contain the stabilized Q-SNARE complex for 20
min at 37°C, the sample was subjected to TeNT treatment (200 nM, 30 min, 37°C) then boiled for 5 min
at 95°C and analysed by immunoblotting with antibody against VAMP-2 (clone number 69.1, Synaptic
Systems (Göttingen, Germany)).

Preparation of bovine chromaffin cells. Chromaffin cells were isolated from the bovine adrenal gland
medulla by two-step collagenase digestion as previously described[44]. The cells were grown on poly-D-
lysine-coated glass coverslips in Dulbecco's modified Eagle medium/F-12 (Invitrogen, CA) containing
10% fetal bovine serum (Hyclone Laboratories, UT) and 1% antibiotics (Invitrogen, CA).

Amperometric measurement. Recordings of LDCV exocytosis from chromaffin cells were performed
as described previously[44]. Carbon-fiber electrodes were fabricated from 8 μm diameter carbon fibers and
back-filled with 3 M KCl. The amperometric current, generated by oxidation of catecholamine, was

measured using an axopatch 200B amplifier (Axon Instruments Inc., CA), which was operated in voltage-

clamp mode at a holding potential of + 650 mV.

Electron microscopy. Chromaffin cells, grown on Vitrogen collagen matrix (Cohesion, Palo Alto, CA),

were washed out with Locke's solution containing 157.4 mM NaCl, 5.6 mM KCl, 2.2 mM $CaCl_2$, 1.2 mM

$MgCl_2$, 5.6 mM D-glucose, 5 mM HEPES, and 3.6 mM $NaHCO_3$, pH 7.4 titrated by NaOH. As described

previously[44], cells were fixed with 2% paraformaldehyde and 2% glutaraldehyde in 0.05 M sodium

cacodylate buffer at pH 7.4 for 20 min at room temperature. Cells were post-fixed with 0.5% osmium

tetroxide in 0.05 M sodium cacodylate buffer at pH 7.4 for 30 min at room temperature. Cells were further

dehydrated in graded ethanol solutions and embedded in LR White resin (London Resin Co., Berkshire,

UK). Silver-gold thin sections were stained with uranyl acetate and lead citrate. The thin sections were

examined under JEOL 1200 EX2 transmission electron microscope at 80 kV.

Liposome co-flotation assay. Liposomes float up through the gradient due to their buoyancy and free

proteins remain in the bottom of the gradient, whereas proteins incorporated in liposomes co-float to the

buoyant density of the liposomes[45]. First, 30 µL of liposomes that incorporate the stabilized Q-SNARE

complex were mixed with Nycodenz (Axis Shield, 80 %, 30 µL) and a second Nycodenz layer (30 %,

40 µL) was gently applied followed by another layer of buffer (40 µL). The density gradient was

centrifuged using a Beckman TL-100 ultracentrifuge (TLS55 rotor, 100,000g, 4°C, 1 h). The 20-µL

aliquots were carefully taken from the top of the gradient and analysed by coomassie blue staining.

Atomic force microscope (AFM). Supported lipid bilayers (SLBs) was formed from LUVs by adsorption

on the surface and rupture to form lipid bilayers. LUVs prepared as described above were placed on the

surface of PDL-coated cover slips for adsorption. Then the adsorbed LUVs rupture to form SLBs on the

surface of PDL-coated cover slips. Electrostatic interactions between LUVs and the surface of the support

enhance the formation of SLBs. C2AB domain and Ca^{2+} were incubated for 1 h, then the coverslips were

washed twice with double-distilled H_2O and air-dried before imaging. AFM imaging was conducted using

the Bruker Dimension Icon platform (Bruker, USA) and performed in the 'Tapping in air mode' together

with an OTESPA type cantilever. Data were analysed using nanoscope analysis 2.0.

Figure 1

Figure 2

Figure 3

Figure 4

a C2AB
Alexa 488
Ca²⁺ Ca²⁺
Lip. (Rho-PE, PS/PIP₂)

b Ca²⁺ ↓   ↓ EGTA
FRET (F/F_0) vs Time (min)

c C2AB binding (normalized) vs Free Ca^{2+} (µM)
● Chol
● No Chol

d C2AB
Lip. (PS/PIP₂)

e Ca²⁺ ↓   ↓ EGTA
Anisotropy (a.u.) vs Time (min)

f C2AB binding (normalized) vs Free Ca^{2+} (µM)
● Chol
● No Chol

g C2AB binding (normalized) vs Free Ca^{2+} (µM)
● 0.5% PIP2
● 1% PIP2
● 5% PIP2

h LDCV ➕ LUV
No Chol
Fusion (% total fluorescence) vs Time (min)
No Ca²⁺
50 µM Ca²⁺

i Vesicle-mimicking liposome
C2 binding (normalized)
C2A   C2B
No Mg²⁺/ATP
Mg²⁺/ATP

Figure 5

Figure 6

Figure 7

a

Vesicle

Synaptobrevin-2

Syntaxin-1/
SNAP-25

Synaptotagmin-1

▼ Cholesterol

Plasma membrane

Fusion pore

b

ΔG

Membrane
bending

Bending
energy

Docked
state

Fusion

— Cholesterol
+ Cholesterol

Reaction coordinate