High rates of spontaneous chromosomal duplications unravel dosage compensation by translational regulation

Marc Krasovec[1*], Remy Merret[2,3,*], Frédéric Sanchez[1], Sophie Sanchez-Brosseau[1], Gwenaël Piganeau[1*]

[1] Sorbonne Universités, UPMC Univ Paris 06, CNRS, Biologie Intégrative des Organismes Marins (BIOM), Observatoire Océanologique, F-66650 Banyuls/Mer, France

[2] Centre National de la Recherche Scientifique, Laboratoire Génome et Développement des Plantes, UMR5096, 66860 Perpignan, France

[3] Université de Perpignan Via Domitia, Laboratoire Génome et Développement des Plantes, UMR5096, 66860 Perpignan, France

[*] Corresponding authors: krasovec@obs-banyuls.fr ; remy.merret@univ-perp.fr; gwenael.piganeau@obs-banyuls.fr

Keywords: chromosome duplication, mutation accumulation, aneuploidy, dosage compensation, translation efficiency, phytoplankton.

Running title: Spontaneous chromosome duplication rates.

ABSTRACT (203 words)

While duplications have long been recognized as a fundamental process driving major evolutionary innovations, direct estimations of spontaneous chromosome duplication rates, leading to aneuploid karyotypes, are scarce. Here, we provide the first estimations of spontaneous chromosome duplication rates in six unicellular eukaryotic species from mutation accumulation (MA) experiments. The spontaneous chromosome duplication rates reach 1×10^{-4} to 1×10^{-3} per genome per generation, which is ~4 to ~50 times less frequent than spontaneous point mutations per genome, whereas chromosome duplication events can affect 1 to 7% of the total genome size. Comparative transcriptomics between MA lines with different chromosome duplications reveals a strong positive correlation between RNA expression rate and DNA copy number. However, comparative analyses of the translation rate of mRNAs estimated by polysome profiling unravel a chromosome specific dosage compensation mechanism. In particular, one chromosome with a gene average of 2.1 excess of mRNAs is compensated by an average of ~0.7 decrease in translation rates. Altogether, our results are consistent with previous observations of a chromosome dependent effect of dosage compensation and provide evidence that it may occur during translation. These results support the existence of a yet unknown post-transcriptional mechanism orchestrating the modification of translation of hundreds of transcripts from genes located on duplicated regions in eukaryotes.

INTRODUCTION

Complete or partial chromosome duplications leading to aneuploidy karyotypes are known to be at the origin of genetic diseases, such as Trisomy 21 in humans, and is a near universal feature of tumor cells (Rajagopalan and Lengauer 2004). When a single chromosome is duplicated, an immediate issue arises: the imbalance of ploidy and gene dose within a same karyotype. This is expected to have deleterious effects because it creates an imbalance of the transcript and protein productions, which is costly and may disrupt the function of a pathway and protein interactions (Dephoure et al. 2014; Veitia and Potier 2015). The deleterious effects of aneuploidy have been documented in many different biological model systems such as yeast (Torres et al. 2007), mouse and human (Gearhart et al. 1987). In *Caenorhabditis elegans*, mutation accumulation experiments provided evidence of purifying selection against gene duplications causing excess of transcripts, as compared to gene duplications associated to invariant transcript levels (Konrad et al. 2018). Moreover, the consequences of aneuploidy on gene expression are complex and while gene expression may increase with chromosome copy number, gene expression changes may also spread outside the duplicated regions in *Arabidopsis* (Hou et al. 2018; Song et al. 2020), *Drosophila* (Devlin et al. 1988), and human cells (FitzPatrick et al. 2002). Contrasting with the before mentioned studies, aneuploidy may confer a selective advantage as a response to stress, such as resistance to drugs in *Candida albicans* (Selmecki et al. 2006) or *Saccharomyces cerevisiae* (Chen et al. 2012). Recent evidence about a lack of deleterious effect of aneuploidy is the high aneuploidy prevalence and tolerance across *Saccharomyces cerevisiae* lineages (Scopel et al. 2021), a fifth of the sequenced strains harboring atypical aneuploidy karyotypes (Peter et al. 2018). Indeed, different dosage compensation mechanisms have evolved and may restore the ancestral gene dose leading to aneuploidy tolerance. The three targets of dosage compensation mechanisms of duplicated genes are the modification of the transcription rate, the translation rate, or the protein degradation rate of duplicated genes. The most studied mechanisms are those involved in modifying the transcription rate during the evolution of heterogametic sex chromosomes, known both in plants (Muyle et al. 2012; Muyle et al. 2017; Charlesworth 2019) and animals (Disteche 2012; Graves 2016). Different mechanisms evolved either by simulating the ancestral ploidy by doubling the expression of the genes of the single copy male X (Baker et al. 1994); or by equalizing the ploidy between the two sexes by halving the

expression – silencing of one X in female (Heard et al. 1997). Although dosage
compensation is well studied in sex chromosome evolution, whether and how it
evolves after a chromosome ploidy variation is unclear, particularly for autosomes
(Kojima and Cimini 2019). Previous studies did not report consistent results, with a
significant increase of expression for the duplicated genes in *C. albicans* (Selmecki et
al. 2006), *Drosophila* (Loehlin and Carroll 2016), *S. cerevisiae* (Torres et al. 2007),
mammalian cells (Williams et al. 2008); and a decrease of the expression of the two
copies in yeast and mammals, suggesting a compensation at the transcriptional level
(Henrichsen et al. 2009; Qian et al. 2010). In the case of chromosome duplication,
evidence for dosage compensation at the transcriptional level is scarce (Stenberg et
al. 2009; Hose et al. 2020), and scaling of gene expression with gene copy number
has been reported in disomic yeasts (Kaya et al. 2020). Interestingly, dosage
compensation has been reported to occur at the post-transcriptional level via the
modification of translation efficiency in *Drosophila* (Zhang and Presgraves 2017) or at
the post-translational level for 20% of the proteome in yeast via an increased
degradation rate of proteins (Dephoure et al. 2014). Increase of protein degradation
has also been reported to be involved in compensation mechanism in human Downs
syndrome for proteins encoded on the triplicated chromosome 21 and include known
stable heteromeric protein complexes (Liu et al. 2017).
Although the consequences of chromosome duplication on transcription rates - and
to a lesser extent their consequences on translation rates and protein abundance -
have been investigated in many model organisms, our knowledge of the rate of
chromosome duplication events in eukaryotes have been yet restricted to
Saccharomyces cerevisiae (Lynch et al. 2008; Zhu et al. 2014; Liu and Zhang 2019)
and humans (Nagaoka et al. 2012; Loane et al. 2013). Here, we investigate the
spontaneous chromosome duplication rate in six unicellular photosynthetic species by
analyzing mutation accumulation (MA) experiment genomes. The principle of MA
experiments is to follow the descendants originated from a single cell under minimal
selection insured by serial bottlenecks during dozens to thousands cell divisions
(Halligan and Keightley 2009). Here, MA experiments have been performed by
maintaining 12 to 40 MA lines for a total of 1,595 to 17,250 generations depending of
the species (Krasovec et al. 2017; Krasovec, Sanchez-Brosseau, et al. 2018;
Krasovec et al. 2019). The six species include five Chlorophyta (one
Trebouxiophyceae et four Mamiellophyceae) and one Bacillariophyta, ecological

relevant primary producers in the sunlit ocean (de Vargas et al. 2015), with a large

phylogenetic spread encompassing 1.5 billion years of divergence (Yoon et al. 2004).

We further recovered cryopreserved MA lines to investigate the consequence of

chromosome duplication on the transcription and translation rates.

RESULTS

Whole chromosome duplication rate

In this study, we analyzed whole genome resequencing data from previous MA experiments (Krasovec et al. 2016; Krasovec et al. 2017; Krasovec, Sanchez-Brosseau, et al. 2018) in five haploid green algae species (Chlorophyta, four Mamiellophyceae and one Trebouxiophyceae) and one diploid diatom (Bacillariophyta) : *Picochlorum costavermella* RCC4223 (Krasovec, Vancaester, et al. 2018), *Ostreococcus tauri* RCC4221 (Blanc-Mathieu et al. 2014), *O. mediterraneus* RCC2590 (Subirana et al. 2013), *Bathycoccus prasinos* RCC1105 (Moreau et al. 2012), *Micromonas commoda* RCC299 (Worden et al. 2009) and *Phaeodactylum tricornutum* RCC2967 (Bowler et al. 2008; Giguere et al. 2022). Coverage was used as a proxy of copy number (Figure 1A and Supplemental figures S1 to S16), and unveiled whole chromosome duplication events in four of the six species (Table 1 and Table S1). Four whole chromosome duplications were observed in *M. commoda* (chromosomes C05, C12, C16 and C17), four in *B. prasinos* (chromosomes C04, C05, C06 and C19), one in *O. mediterraneus* (chromosome C14) and four in *P. tricornutum* (chromosomes C02, C14 and C23). The chromosome duplication events were mapped onto the genealogies of the MA lines to identify all independent chromosome duplication events (Figure 1B for *B. prasinos*). Genealogy analysis provided evidence that several chromosomes were duplicated two times independently: the MA lines Bp25c and Bp28b carry two independent duplications of chromosome C19; Mc08 and Mc09 carry two independent duplications of chromosome C17; and Pt11 and Pt10c carry two independent duplications of chromosomes C23. The probability of observing two independent whole chromosome duplication of the same chromosome in *B. prasinos*, *M. commoda* and *P. trichornutum* are 0.46, 0.47 and 0.34 respectively (see methods), and these probabilities are thus consistent with an equal probability of duplication across chromosomes. All independent duplication events inferred from coverage analysis and genealogies are summarized in Table 1 and S1. Unexpectedly, the analyses revealed that the ancestral line of the MA experiment in *B. prasinos* carried two copies of the chromosome C01. One of the copies had subsequently been lost 11 times independently over 4,145 generations, corresponding to a spontaneous chromosome loss of a duplicated chromosome of 0.006 per duplicated chromosome per generation in *B. prasinos*.

A. Whole genome read coverage for different mutation accumulation lines

B. Genealogy of the mutation accumulation experiment in *B. prasinos* RCC4222

Figure 1. A. Normalized raw genomic coverage of the mother line (T0 of mutation accumulation experiment), Bp20c and Bp28 mutation accumulation lines of *Bathycoccus prasinos*. Grey lines are chromosome separators. In blue are the chromosomes in double copies. Raw coverage of all lines from all species are provided in Figures S1 to S16. **B.** Pedigree of the mutation accumulation lines from the *B. prasinos* experiment. Chromosome C01 is duplicated in the T0 line (named mother line) of the experiment. This duplication is then lost several times, and five other duplications of chromosomes C04, C05, C06 and C19 occurred.

Table 1. Spontaneous whole chromosome duplication rate in six species. N_{lines} : number of MA lines,
Gen : average number of generations per MA line, Tot_{Gen} : total number of generations, N_{Chrom} : number
of chromosomes in the ancestral karyotype, N_{WCD} is the number of independent whole chromosome
duplications, U_{WCDI} is the whole chromosome duplication rate per chromosome per cell division, and
U_{Cell} is the whole chromosome duplication rate per cell division, U_{bs} is the base substitution mutation
rate per cell division. The estimation of the upper limit of N_{WCD} and U_{cell} in *O. tauri* and *P. costavermella*
relies on the assumption of one duplication event.

Species	N_{lines}	Gen	Tot_{Gen}	N_{Chrom}	N_{WCD}	Duplicated Chrom	U_{WCD}	U_{cell}	U_{bs}
Bathycoccus prasinos	35	265	4,145	19	5	C04, C05, C06, C19	0.000063	0.00121	0.0046
Micromonas commoda	37	272	4,994	17	5	C05, C12, C16, C17	0.000059	0.00100	0.0171
Ostreococcus tauri	40	512	17,250	20	0	-	<0.000003	<0.00006	0.0054
Ostreococcus mediterraneus	33	272	8,380	19	1	C14	0.000006	0.00012	0.0064
Picochlorum costavermella	12	133	1,596	10	0	-	<0.00006	<0.00063	0.0119
Phaeodactylum tricornutum	36	181	6,516	25	4	C02, C14, C23	0.000025	0.00061	0.0132

Consequences of chromosome duplication on transcription

To explore the consequences of the whole or partial chromosome duplication on
transcription, we estimated the level of transcription of a control strain (*B. prasinos*
RCC4222) and one MA line of *B. prasinos* by recovering one 4-year-old cryopreserved
MA line. This cryopreserved culture originated from MA line Bp37, and the recovered
culture is hereafter named Bp37B. Chromosome copy number of Bp37B and the
control line were estimated by whole genome resequencing. This confirmed the two
copies of chromosome C04 in Bp37B as in the original Bp37 (Figure 2A and S7), and
also revealed additional karyotypic changes: an increase in chromosome C01 copy
number (Figure 2A, Figure S17) in Bp37B. The heterogeneous coverage of
chromosome C01 led us to divide it into two regions for subsequent analyses; region
C01a (1.98 fold coverage) and C01b (1.35 fold coverage). The control line also
contained duplicated regions (Figure 2A, Figure S17), chromosome C10 and a region
of chromosome C02 named C02b, while C02a is in single copy. Limits of sub-regions
of chromosomes C01 and C02 are provided in Table S2. We interpreted coverage
values smaller than 2 (Figure S17), *e.g.* C01b in Bp37B, C10 and C02b as duplications
that are carried by a subpopulation of cells. Cultures were grown to up to 10 million
cells per ml prior to extraction so that polymorphism is not unexpected.
Comparative transcriptome analyses of Bp37B and the control line revealed that there
are on average twice the number of transcripts for genes located on the duplicated
chromosomes as compared to the control line - C04 (transcription rate *tr(i)*

average=2.13, median=1.95, estimated from TPM, Figure 2B, raw data in Table S3)

and C01a (transcription rate *tr(i)* average=2.29, median=2.02). The transcription rate

of genes on chromosomes C10 and C02b is also affected. Altogether, the transcript

production scales up with the chromosome copy number in the Bp37B and control

lines (Pearson correlation, rho=0.87, p-value<0.001, Figure 3A).

Gene-by-gene variation of the consequences of DNA copy number

Dosage invariant genes (Antonarakis et al. 2004; Lyle et al. 2004) are genes for which

the transcription is not affected by copy number, and they may be involved in

aneuploidy tolerance. We investigated differential gene expression at the individual

gene scale with Deseq2 (Love et al. 2014) to identify candidate invariant genes. We

found that the transcription rate of 158 out of the 1776 genes in two copies (9%)

located on duplicated chromosomes or regions were not significantly higher than for

genes in single copy (Table S4). Two Gene Ontology categories (RNA processing,

GO:0006396 and DNA metabolic process GO:0006259) were over-represented in this

transcriptomic invariant gene set. First, genes involved in RNA processing (3.8 times

more frequent, p-value <0.01) including members of heteromeric protein complexes

such as subunits 7 and 2 of the U6 snRNA-associated Sm-like protein (Table S5,

invariant 158 annotations). Second, the genes involved in DNA metabolic process (2.2

times more frequent in the invariant gene subset, p-value <0.04) including the DNA

Polymerase A and the DNA primase large subunit. Notable protein complex members

of the invariant data set are Histone 3 and the subunit E of the translation initiation

factor 3. There are in total 12 genes annotated as subunits in the subset of invariant

genes, this is significantly more than the frequency of protein coding genes annotated

as subunits in the complete proteome (Fisher exact test p=0.0006). This suggests that

a higher proportion of these genes coding for protein forming complexes have a gene-

specific tailored regulated transcription that is not affected by gene copy number.

Figure 2. A. Genomic coverage of 1 kbp windows along the genome of the MA line *Bathycoccus prasinos* Bp37B. **B.** Distribution of genes transcription rate *tr(i)* of the 19 chromosomes. **C.** Distribution of genes translation efficiency *ta(i)* of the 19 chromosomes. **B** and **C**. The transcription rate and translation efficiency ratio averages are different between chromosomes (ANOVA, p-value<10⁻¹⁵). **D.** Distribution of genes protein homeostasis (transcription rate x translation rate) of the 19 chromosomes. Y axis are in log2. Blue: duplications in Bp37B. Yellow: duplications in control RCC4222.

Consequences of chromosome duplication on translation

The compelling evidence of overexpression of the majority of duplicated genes prompted us to investigate whether post-transcriptional processes may temperate this excess of transcripts. This second hypothesis was tested in the same *B. prasinos* MA line by sequencing mRNAs associated with ribosomes (polysomes) in order to compare the translation efficiency *te(i)* of genes located on duplicated and non-duplicated lines, by estimating the relative proportion of mRNA binded to a ribosome in single versus duplicated genes. We found that the average translation efficiency of the genes located on the duplicated chromosome C04 was 0.71 (median=0.54) as compared to the genes on this chromosome in the control line (Figure 2C). At chromosome scale (Table S7), the translation rate is 0.67 for the chromosome C04. However, the translation efficiency of part C01a of the chromosome C01 was 0.93 and 1.54 for the part C01b. The translation efficiency of the two parts of the chromosome C02 are 1.04 and 0.76 for C02a and C02b, respectively, and 0.99 for chromosome C10. Translation rate modification is thus chromosome dependent (Figure 3): genes located on the C04 and C01b display an important variation in translation efficiency between the two lines. Last, we estimated the expected protein production in Bp37B as compared to the control line by going back to the absolute translation rate for each gene: that is the ratio of mRNA in polysomes in Bp37B as compared to the control (Figure 2D). This predicted that genes on C04 have a similar protein production rate in Bp37B and in the control line, despite a higher transcription rate as a consequence of the chromosome duplication, as well for the two parts of the chromosome C02 (Figure 2D). However, the two parts of the chromosome C01, despite different transcription and translation efficiency rates both leads to a protein level that is the double of the control line. The chromosome C10 also exhibits a lower protein content in Bp37B compared to control, meaning there is an excess of proteins from genes linked to this chromosome in the control line because of the duplication. Altogether, we observed there is no dosage compensation at the transcription level for any of the duplicated regions, but that dosage compensation may occur at the translation level on some duplicated regions and seems chromosome dependent (Figure 3).

Figure 3. Raw relative genomic coverage average (Bp37B/Control) related to average transcription rate *tr(i)* (**A**), translation efficiency *te(i)* (**B**), and average protein synthesis (average transcription rate x average translation efficiency) per chromosome (**C**). Blue: chromosomes duplicated in Bp37B; Yellow: chromosomes duplicated the control RCCC4222; Black: non-duplicated chromosomes; Grey: outlier chromosome C19. Data are provided in Table S6 and S7.

 Searching for independent molecular signatures of translation modulation

The observed dosage compensation at the translation level on chromosome C04 suggests that a post-transcriptional mechanism occurs on duplicated transcripts. As the length of the poly(A) tail is a key feature of many cytoplasmic mRNAs and is known to regulate translation (Subtelny et al. 2014; Eichhorn et al. 2016; Lim et al. 2016), we tested if DC observed for the translation of transcripts linked to duplicated chromosome C04 is associated with a variation of poly(A) tail length. Using 3'RACE experiment, we measured the poly(A) tail length in Bp37B and control (Figure 4 and S18) of the transcripts of three genes located on chromosome C04 (04g00840, 04g01730 and 04g04360), one gene located on chromosome C08 (08g03470) and C01 gene located on chromosome C01a (01g01300). We systematically observed that the poly(A) tail length is reduced in Bp37B line compared to control line for transcripts located on chromosome C04 (Figure 4), whereas no difference is observed control (Figure S18).

Figure 4. Transcripts from genes located on duplicated chromosome C04 present shorter poly(A) tail. Poly(A) tail measurement was performed using modified 3'RACE. PCR amplification was performed using primers flanking poly(A) tail (**A**) or primers anchored in 3'UTR just before poly(A) tail as internal control (**B**). Experiments was performed using total RNA from RCC4222 line (control line) and BP37B line (with duplicated chromosome C04). Illustrations representing PCR amplification are present on the right panel. Black line represents the messenger RNA and the orange line represent the ligated adapter used for reverse transcription and PCR amplification. FW. Forward primer. RV. reverse primer.

DISCUSSION

Rates of whole chromosome duplication in eukaryotes

The rates of spontaneous whole chromosome duplication reported here vary between 6×10^{-5} to 1×10^{-3} events per haploid genome per generation and are thus one to two orders of magnitude lower than spontaneous point mutations per genome per generation. Theoretically, there are two possible mechanisms at the origin of chromosome duplication i) the supernumerary replication of one chromosome before cell division, leading to one cell with one chromosome and one cell with two chromosomes; or ii) the unequal segregation of chromosomes during cell division, leading to one cell with two copies and one cell without any copy. The latter scenario has been intensively experimentally studied in human cells (Ford and Correll 1992; Cimini et al. 2001), and seems all the more likely in the extremely small-sized Mamiellophyceae cells (1 µm cell diameter) as it has been reported that they contain less kinetochore microtubules than chromosomes (Gan et al. 2011), which may induce high error rate in the chromosome segregation process. However, the whole chromosome duplication rates reported in Mamiellophyceae are in line with those found in the diatom *P. tricornutum* and in yeast (9.7×10^{-5}) (Zhu et al. 2014), the latter containing approximately as many kinetochore microtubules as chromosomes (Peterson and Ris 1976). Altogether, the spontaneous whole chromosome duplication rates reported previously in yeast (Zhu et al. 2014) and in the species from evolutionary distant eukaryotic lineages reported here (Yoon et al. 2004) suggest a high rate of spontaneous aneuploidy across the eukaryotic tree of life. To investigate whether spontaneous whole chromosome duplication had been overlooked in other lineages, we screened publicly available resequencing data of MA on the freshwater green algae *Chlamydomonas reinhardtii* (Ness et al. 2015), and identified one event for chromosome 14 in the strain CC1373. We also provided evidence of spontaneous whole chromosome duplication (four chromosomes in a single individual from a pedigree) in the brown algae *Ectocarpus* (Krasovec et al. 2022).

Interestingly, we were also able to estimate the rate of chromosome duplication loss during one experiment to be about three orders of magnitude higher than the spontaneous duplication rate. This is consistent with previous observation in *S. cerevisiae*, and our observations that cells with duplicated chromosomes are ephemeral and unlikely to be maintained for a long time in batch culture. Indeed, the coverage of the large duplication events on C01b, C02b, C04 and C10 were 1.4,1.6,

1.8 and 1.4 times the coverage of single copy chromosomes respectively, suggesting

that the proportion of cells carrying a supernumerary chromosome within the

population is 0.4, 0.6 and 0.8. In addition, we observed the loss of the chromosome

C04 duplication in one of the Bp37B lines over 4 months of sub-culturing.

As opposed to the aneuploidy generated during meiotic divisions, which has been

intensively studied in mammalian cells (Nagaoka et al. 2012), the chromosomal

duplication events reported here are generated during mitotic division. Since the

molecular mechanisms involved in chromosomal segregation in mitosis, meiosis 1 and

meiosis 2, are different, the associated aneuploidy rates are not expected to be equal.

The aneuploidy rates in mitotically dividing human cells, such as HCT116 lines, has

been estimated to be 7×10^{-2} (Thompson and Compton 2008), more than one order of

magnitude higher than the highest estimation of 1.2×10^{-3} reported here in *B. prasinos*.

The observed difference of the spontaneous point mutation rate per genome per

generation is expected to vary between species as the consequence of different

genome sizes, effective population sizes (Lynch et al. 2016), and the difference

between the observed and expected GC content (Krasovec et al. 2017). However, the

only eight spontaneous aneuploidy rates yet available are not sufficient to investigate

the reasons of chromosome duplication rate variations, such as the impact of

chromosome number or effective population size.

Fitness effect of whole chromosome duplication and dosage compensation

The high spontaneous rates of spontaneous chromosome duplication challenge the

common idea that whole chromosome duplications are highly deleterious for individual

cells. Using the number of cell divisions per days as a proxy of fitness during the MA

experiment (Krasovec et al. 2016), only one out of the 15 MA lines with a whole

chromosome duplication displayed a significant fitness decrease (Bp28b, Pearson

correlation, p-value=0.016, ρ=-0.846, Table S8). Recent studies in yeast are

inconsistent with a highly deleterious fitness effect of aneuploidy and point towards

slightly deleterious effects, as well as a significant effect of the genetic background on

aneuploidy tolerance (Scopel et al. 2021). Compensating point mutations have been

previously linked to aneuploidy tolerance in yeast (Torres et al. 2010), notably in the

gene encoding the deubiquitinating enzyme UBP6. Point mutations are unlikely to

impact the aneuploidy tolerance of the different MA lines reported here, as some MA

lines with chromosome duplication do not carry any point mutation (Mc3, Bp28b, Bp26

and Bp25), or carry only synonymous mutations and mutations located in intergenic

region (*i.e.* Om3 or Mc28, Table S9). The deleterious effect of aneuploidy could be

mitigated by a dosage compensation mechanism preserving a relative protein

homeostasis. Here in *B. prasinos*, despite the important variation in transcription and

translation between genes, we observed that transcription rates overwhelmingly

scaled up with chromosome copy number (Figure 3A) and that translation rates

differed significantly between chromosomes (Figure 3B). However, chromosome C19

displays a unique pattern. This chromosome is a small hypervariable outlier

chromosome found in all Mamiellophyceae species sequenced so far (Blanc-Mathieu

et al. 2017; Yau et al. 2020). Previous studies reported a strong variation in

transcription rates of genes on this chromosome, which is associated with resistance

to viruses (Yau et al. 2016) and characterized by lower GC content than other

chromosomes.

For the duplicated chromosomes or regions, we observed three different patterns of

translation efficiency. First, we observed no relative difference in translation rates for

genes on duplicated chromosome C10, which is duplicated in 40% percent of cells,

and region C01a, which is duplicated in 100% of cells. The inferred excess of protein

synthesis of duplicated genes on this chromosome is thus 40% (Figure 3C). Second,

we observed a decrease of the translation rate for genes linked to chromosome C04

and region C02b. More precisely, the estimated percent of mRNA linked to ribosomes

is inversely proportional to the excess of mRNA produced for duplicated genes on

these two chromosomes, predicting approximately the same protein production rate

as in cells without duplicated chromosomes (Figure 3C). Third, and very surprisingly,

we observed an increase of the translation level of genes on C01b, that are in two

copies in 40% of cells. As a consequence, the rate of protein synthesis on genes

located on region C01b reach the same relative protein synthesis than genes located

on region C01a, that is twice the protein synthesis predicted in cells with a single copy

of this chromosome. We speculate the absence of dosage compensation on region

C01a and the excess of translation on region C01b simulate the ancestral protein

homeostasis in the mother line, which contained two entirely duplicated chromosome

C01.

Chromosome-wide regulation of gene translation is relatively unexplored and poorly
understood. Indeed, while location on a chromosome may predict a gene's
transcription rate as a result of epigenetic marks on the DNA, epitranscriptomics
chromosome wide mRNA modifications mechanisms are yet to be discovered.
Notwithstanding, chromosome wide translational dosage compensation has been
previously observed in *Drosophila* (Zhang and Presgraves 2016), and at the gene
scale in several species (Zhang and Presgraves 2017; Chang and Liao 2020).
400

Poly(A) tail length
The role of poly(A) tail length in the translation efficiency is starting to be better
understood (Weill et al. 2012; Subtelny et al. 2014), and alternative polyadenylation is
indeed implicated in several processes: transcription termination by RNAP II, mRNA
stability, mRNA export and translation efficiency (Zhang et al. 2010; Di Giammartino
et al. 2011). In the cytoplasm, the poly(A) tail plays important roles in mRNA translation
and stability and the modulation of its length has an important impact on translation
efficiency (Subtelny et al. 2014; Eichhorn et al. 2016; Lim et al. 2016). As example,
during oocyte maturation and early embryonic development, an increase of poly(A) tail
length occurs for particular mRNAs resulting in an increase of translation (Subtelny et
al. 2014; Eichhorn et al. 2016; Lim et al. 2016). This modulation of poly(A) tail length
has also been found to activate some neuronal transcripts (Udagawa et al. 2012).
Recently, it has been suggested that poly(A) tails can be modulated to balance mRNA
levels and adjust translation efficiency (Slobodin et al. 2020). In this last case, the
CCR4-Not complex shortened the poly(A) tails that reduces the stability of mRNAs.
Here, our data suggest that, in a context of chromosome duplication, modulation of
poly(A) tail length could be a key post-transcriptional mechanism necessary for
dosage compensation at translation level. Future work to explore genome wide
analysis of poly(A) tail variation and post-transcriptional RNA modifications such as
adenosine methylation (Miao et al. 2022) are poised to clarify the translation
compensation mechanisms.

CONCLUSION

In conclusion, the high prevalence of whole chromosome duplication in five unicellular photosynthetic eukaryotes suggest spontaneous whole chromosome duplication is pervasive in eukaryotes. In one species, *B. prasinos*, we provide evidence that a whole duplication chromosome event is associated with dosage compensation at the post-transcriptional level which might involve the adjustment of poly(A) tail length. These results point to yet unknown post-transcriptional regulation mechanisms in DC of aneuploid karyotypes.

MATERIALS AND METHODS

Sequencing data from mutation accumulation (MA) experiments

Mutation accumulation (MA) experiments of the six species *Picochlorum costavermella* RCC4223 (Krasovec, Vancaester, et al. 2018), *Ostreococcus tauri* RCC4221 (Blanc-Mathieu et al. 2014), *O. mediterraneus* RCC2590 (Subirana et al. 2013; Yau et al. 2020), *Bathycoccus prasinos* RCC1105 (synonym to RCC4222) (Moreau et al. 2012), *Micromonas commoda* RCC299 (Worden et al. 2009) and *Phaeodactylum tricornutum* RCC2967 (Bowler et al. 2008; Giguere et al. 2022) were conducted with a flow cytometry protocol described previously for phytoplankton species in liquid medium (Krasovec et al. 2016). Briefly, a mutation accumulation experiment consists in following of MA lines that have evolved from a same cell (the ancestral line) during hundreds of generations. Relaxed selection pressure on spontaneous mutations is ensured by maintaining all MA lines at very low effective population sizes ($6 < Ne < 8.5$) throughout the experiment (Krasovec et al. 2016). As a consequence, MA experiments enable to estimate spontaneous mutation rates, excluding lethal mutations. Coupled with whole genome sequencing of ancestral and MA lines, this experiment enables the direct estimation of the spontaneous mutation rate of a species. Here, MA lines came from a single cell obtained by dilution serving as T_0 culture (named the mother line, ML) and were maintained in 24-wells plates in L1 medium at 20 °C with a 16h-dark 8h-light life cycle. Single cell bottleneck by dilution were done each 14 days to have a low effective population size and limit selection. DNA of initial line (ML line) and final time of MA lines were extracted with chloroform protocol and sequencing done with Illumina HiSeq or MiSeq by GATC biotech (Germany). To detect duplications, raw reads were mapped against the reference genomes for each strain with bwa mem v0.7.12 (Li and Durbin 2010). Then, bam files were treated with samtools v1.3.1 (Li et al. 2009) and bedtools v 2-2.18.0 (Quinlan and Hall 2010) to extract the coverage.

Statistical analyses

The probability of drawing one chromosome twice out of *k* WCD events chosen from *n* chromosomes, *P*(*j*,*n*), is equal to the ratio of the number of combinations of (*k*-1) out of *n* without replacement multiplied by *k*-1, which corresponds to the number of possible *k*-1 different chromosomes with one in two copies, out of *n*, divided by the

number of combinations of *k* out of *n* with replacement (*k*-1)*C(*k*-1,*n*)/(*n*+*k*-1), which

corresponds to the total number of possible subsamples of k chromosomes out of *n*.

$$P(k,n) = \frac{C(k-1,n) * (k-1)}{(n+k-1)!/k!\,(n-1)!}$$

All statistical analyses were performed with R (R Core Team 2022).

Dosage compensation analysis

This study was started three years after the end of the MA experiments, for which

some MA lines had been cryopreserved which allowed us to re-start a culture of one

B. prasinos MA line with a duplicated chromosome C04 (Bp37) to investigate possible

dosage compensation. The restarted culture from the cryopreserved Bp37 was

renamed Bp37B. As a control, we used the reference culture of the strain *B. prasinos*

RCC4222, that is the derived from the RCC1105 used for the MA experiment. All

cultures (Bp37B and RCC4222) were maintained under a 12:12 h light:dark regime

under 50 µmol photon m^{-2} s^{-1} white light at 20 °C. The karyotype of the defrozen

cultures Bp37B and the control RCC4222 were checked by DNAseq resequencing.

The karyotype of Bp37B contained on additional chromosome copy number as

compared to Bp37, and the karyotype of RCC4222 contained one additional copy

numbers and one less chromosome copy number as compared to the mother line (ML)

(Figure S17). The relative transcription rate was corrected by the number of DNA copy

in each line for further analysis below.

For expression analysis, total RNA was extracted using the Direct-zol RNA MiniPrep

Kit (Zymo Research, Californie, USA) from pooling flasks of cultures (100 ml cultures

with 200 million cells per ml) taken 6h before and 1h before the light on, in triplicates

for the control RCC4222 and duplicates for Bp37B. Then for translation efficiency

analysis, polysome extraction was performed for Ribo-seq as described previously

(Carpentier et al. 2020) with few modifications. Briefly, 600 mL of *B. prasinos* culture

were centrifuged at 8,000 g for 20 minutes. After centrifugation, pellets were

resuspended in 2.4 mL of polysome extraction buffer. After 10 minutes of incubation

on ice and centrifugation, 2 mL of supernatant was loaded on a 9 mL 15–60% sucrose

gradient and centrifuged for 3h at 38 000 rpm with rotor SW41 Ti. Fractions

corresponding to polysomes were pooled and polysomal RNA was extracted as

previously described (Carpentier et al. 2020). RNA library preparation was performed

on total or polysomal RNA using a NEBNext Poly(A) mRNA Magnetic Isolation Module

and a NEBNext Ultra II Directional RNA Library Prep Kit (New England Biolabs)
according to the manufacturer's instructions with 1 µg of RNA as a starting point.
Libraries were multiplexed and sequenced on a NextSeq 550. Raw reads were
mapped against the reference transcriptome of *B. prasinos* with RSEM with standard
parameters (Li and Dewey 2011). We obtained the TPM average of total RNA and
polysomes linked RNA that we compared between *B. prasinos* RCC4222 and the MA
line Bp37B. At the gene scale, we first used the total RNA TPM values to estimate the
expression difference ratio, r_{RNA} for each gene *i*, between two copies and single copy
genes as:

536
$$r_{RNA}(i) = \frac{TPM_{TWOCOPIES}(i)}{TPM_{SINGLECOPY}(i)}$$

These relative expression rates $r_{RNA}(i)$ were normalized by the expression rate median
of genes located on non-duplicated chromosomes in Bp37B (C03, C05, C06, C07,
C08, C09, C011, C012, C013, C014, C015, C016, C017 and C018), in order to
estimate the transcription rate *tr(i)* of duplicated chromosomes related to non-
duplicated chromosomes. Chromosome C19 was not considered because it is a
particularly outlier chromosome (see Discussion).

Second, we used the total RNA and polysomes linked RNA TPM values for each gene
i to calculate the translation rates of each gene, *r(i)*, as:

545
$$r(i) = \frac{PolysomeTPM(i)}{TotalTPM(i)}$$

546   The translation efficiency between a gene *i* in the two lines in single versus two copies,
547   *te(i)* was estimated as:

548
$$te(i) = \frac{r_{TWOCOPIES}(i)}{r_{SINGLECOPY}(i)}$$

The *te(i)* ratio was then normalized by the median translation efficiency of all genes
on non-duplicated chromosomes in strain BP37B (5,267 genes).

Then, to estimate the dosage compensation at the scale of the chromosomes, the
values were normalized by the average TPM of all single copy chromosomes prior to
the calculation of the ratios (Table S7).

Poly(A) tail analysis was performed as previously described with slight modifications
(Sement and Gagliardi 2014). PCR products were resolved on a 2.5% agarose gel.
Primers used in this study are available in Table S10.

Differential gene expression analyses

The statistical significance of the genes differential expression levels was further

estimated with Deseq2 (Love et al. 2014). Transcriptional invariant genes were

defined as genes present on duplicated regions and for which the relative transcription

rate $r_{RNA}(i)$ was comprised between 0.9 and 1.1. The over-representation of a certain

GO term in the transcriptional invariant gene set was compared to the genome-wide

GO term background frequency using the GO enrichment analysis with default values

implemented in pico-PLAZA workbench (Vandepoele et al. 2013).

AUTHOR CONTRIBUTIONS

MK, SSB and GP performed the mutation accumulation experiments. MK performed

the bioinformatics analysis. FS performed the cell cultures, RNA extractions, PCR and

gel migration experiments. RM designed and performed the polysome analyses,

poly(A) tail tests and RNAseq preparation. GP performed statistical analyses and

coordinated the project. MK drafted the first version of the manuscript and all authors

participated to writing the final version.

ACKNOWLEDGEMENTS

We are grateful to Claire Hemon and Elodie Desgranges for technical assistance with

the MA experiments. We acknowledge the GenoToul Bioinformatics platform

(Toulouse, France) for bioinformatics analysis support and cluster availability, the

BIOPIC platform for support with the cytometry, and the sequencing facility of the

Université de Perpignan Via Domitia BioEnvironnement platform. This work was

funded by ANRJCJC-SVSE6-2013-0005 to GP and SSB and ANR PHYTOMICS

(ANR-21-CE02-0026). This study is set within the framework of the "Laboratoires

d'Excellences (LABEX)" TULIP (ANR-10-LABX-41) and of the "École Universitaire de

Recherche (EUR)" TULIP-GS (ANR-18-EURE-0019).

COMPETING INTERESTS

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

All genomic raw reads of ML and mutation accumulation lines are available under the

bioprojects PRJNA531882 (*Ostreoccocus tauri*, *O. mediterraneus*, *Micromonas*

commoda, *Bathycoccus prasinos*), PRJNA453760 and PRJNA389600 (*Picochlorum*
costavermella), and PRJNA478011 (*Phaeodactylum tricornutum*). Transcriptomic raw
reads of the *Bathycoccus prasinos* MA lines 37 (Bp37B) and control line RCC4222 are
available under the bioproject PRJNA715163. A summary is provided in Table S11.
SUPPLEMENTARY MATERIAL
597
Figure S1 to S16. Raw coverage by 1 kb windows of all mutation accumulation lines
from *Ostreoccocus tauri*, *O. mediterraneus*, *Bathycoccus prasinos*, *Micromonas*
commoda, *Picochlorum costavermella* and *Phaeodactylum tricornutum*. The
horizontal grey lines indicate the chromosome separation, and the duplicated
chromosome are in blue. ML is the T0 genome of mutation accumulation experiments.
In *M. commoda*, a duplication of a fraction of chromosome 2 in the mother line (ML) is
maintained in all MA lines, suggesting there might be a miss-assembly due to a
duplicated region at this location.
Figure S17. Coverage analysis of the Bp37B and control RCC4222.
Figure S18. Transcripts from duplicated chromosome C01 and non-duplicated
chromosome C08 present similar poly(A) tail.
609
Table S1. List of duplicated chromosomes in the MA lines.
Table S2. Limits of sub-regions of chromosomes C01 and C02.
Table S3. Raw transcript per million (TPM) table for all 10 samples.
Table S4. Estimation of the number of dosage insensitive genes per chromosome.
Table S5. List of the dosage insensitive genes with annotations.
Table S6. Relative copy numbers of chromosomes in the two lines of *Bathycoccus*
prasinos.
Table S7. Relative TPM average per chromosome used for Figure 2.
Table S8. Fitness of MA lines with duplicated chromosomes during MA experiments.
Table S9. Point mutations previously identified in the MA lines with whole
chromosome duplications.
Table S10. Sequences of primers used for poly(A) tail length analysis.
Table S11. Summary of all accessions of the data used in this study with Bioprojects
and biosamples.
624
625

BIBLIOGRAPHY

Antonarakis SE, Lyle R, Dermitzakis ET, Reymond A, Deutsch S. 2004. Chromosome 21 and down syndrome: from genomics to pathophysiology. *Nat. Rev. Genet.* 5:725–738.

Baker BS, Gorman M, Marín I. 1994. Dosage compensation in Drosophila. *Annu. Rev. Genet.* 28:491–521.

Blanc-Mathieu R, Krasovec M, Hebrard M, Yau S, Desgranges E, Martin J, Schackwitz W, Kuo A, Salin G, Donnadieu C, et al. 2017. Population genomics of picophytoplankton unveils novel chromosome hypervariability. *Sci. Adv.* 3:e1700239.

Blanc-Mathieu R, Verhelst B, Derelle E, Rombauts S, Bouget F-Y, Carré I, Château A, Eyre-Walker A, Grimsley N, Moreau H, et al. 2014. An improved genome of the model marine alga Ostreococcus tauri unfolds by assessing Illumina de novo assemblies. *BMC Genomics* 15:1103.

Bowler C, Allen AE, Badger JH, Grimwood J, Jabbari K, Kuo A, Maheswari U, Martens C, Maumus F, Otillar RP, et al. 2008. The Phaeodactylum genome reveals the evolutionary history of diatom genomes. *Nature* 456:239–244.

Carpentier M-C, Deragon J-M, Jean V, Be SHV, Bousquet-Antonelli C, Merret R. 2020. Monitoring of XRN4 Targets Reveals the Importance of Cotranslational Decay during Arabidopsis Development. *Plant Physiol.* 184:1251–1262.

Chang AY-F, Liao B-Y. 2020. Reduced Translational Efficiency of Eukaryotic Genes after Duplication Events. *Mol. Biol. Evol.* 37:1452–1461.

Charlesworth D. 2019. Young sex chromosomes in plants and animals. *New Phytol.* 224:1095–1107.

Chen G, Bradford WD, Seidel CW, Li R. 2012. Hsp90 stress potentiates rapid cellular adaptation through induction of aneuploidy. *Nature* 482:246–250.

Cimini D, Howell B, Maddox P, Khodjakov A, Degrassi F, Salmon ED. 2001. Merotelic Kinetochore Orientation Is a Major Mechanism of Aneuploidy in Mitotic Mammalian Tissue Cells. *J. Cell Biol.* 153:517–528.

Dephoure N, Hwang S, O'Sullivan C, Dodgson SE, Gygi SP, Amon A, Torres EM. 2014. Quantitative proteomic analysis reveals posttranslational responses to aneuploidy in yeast.Dikic I, editor. *eLife* 3:e03023.

Devlin RH, Holm DG, Grigliatti TA. 1988. The influence of whole-arm trisomy on gene expression in Drosophila. *Genetics* 118:87–101.

Di Giammartino DC, Nishida K, Manley JL. 2011. Mechanisms and consequences of alternative polyadenylation. *Mol. Cell* 43:853–866.

Disteche CM. 2012. Dosage Compensation of the Sex Chromosomes. *Annu. Rev. Genet.* 46:537–560.

Eichhorn SW, Subtelny AO, Kronja I, Kwasnieski JC, Orr-Weaver TL, Bartel DP.
2016. mRNA poly(A)-tail changes specified by deadenylation broadly reshape
translation in Drosophila oocytes and early embryos.Izaurralde E, editor. *eLife*
5:e16955.

FitzPatrick DR, Ramsay J, McGill NI, Shade M, Carothers AD, Hastie ND. 2002.
Transcriptome analysis of human autosomal trisomy. *Hum. Mol. Genet.* 11:3249–
3256.

Ford JH, Correll AT. 1992. Chromosome errors at mitotic anaphase. *Genome*
35:702–705.

Gan L, Ladinsky MS, Jensen GJ. 2011. Organization of the Smallest Eukaryotic
Spindle. *Curr. Biol.* 21:1578–1583.

Gearhart JD, Oster-Granite ML, Reeves RH, Coyle JT. 1987. Developmental
consequences of autosomal aneuploidy in mammals. *Dev. Genet.* 8:249–265.

Giguere DJ, Bahcheli AT, Slattery SS, Patel RR, Browne TS, Flatley M, Karas BJ,
Edgell DR, Gloor GB. 2022. Telomere-to-telomere genome assembly of
Phaeodactylum tricornutum. *PeerJ* 10:e13607.

Graves JAM. 2016. Evolution of vertebrate sex chromosomes and dosage
compensation. *Nat. Rev. Genet.* 17:33–46.

Halligan DL, Keightley PD. 2009. Spontaneous Mutation Accumulation Studies in
Evolutionary Genetics. *Annu. Rev. Ecol. Evol. Syst.* 40:151–172.

Heard E, Clerc P, Avner P. 1997. X-chromosome inactivation in mammals. *Annu.*
Rev. Genet. 31:571–610.

Henrichsen CN, Vinckenbosch N, Zöllner S, Chaignat E, Pradervand S, Schütz F,
Ruedi M, Kaessmann H, Reymond A. 2009. Segmental copy number variation
shapes tissue transcriptomes. *Nat. Genet.* 41:424–429.

Hose J, Escalante LE, Clowers KJ, Dutcher HA, Robinson D, Bouriakov V, Coon JJ,
Shishkova E, Gasch AP. 2020. The genetic basis of aneuploidy tolerance in wild
yeast.Klein H, Tyler JK, Gresham D, Argueso JL, Bloom K, editors. *eLife* 9:e52063.

Hou J, Shi X, Chen C, Islam MS, Johnson AF, Kanno T, Huettel B, Yen M-R, Hsu F-
M, Ji T, et al. 2018. Global impacts of chromosomal imbalance on gene expression
in Arabidopsis and other taxa. *Proc. Natl. Acad. Sci.* 115:E11321–E11330.

Kaya A, Mariotti M, Tyshkovskiy A, Zhou X, Hulke ML, Ma S, Gerashchenko MV,
Koren A, Gladyshev VN. 2020. Molecular signatures of aneuploidy-driven adaptive
evolution. *Nat. Commun.* 11:588.

Kojima S, Cimini D. 2019. Aneuploidy and gene expression: is there dosage
compensation? *Epigenomics* 11:1827–1837.

Konrad A, Flibotte S, Taylor J, Waterston RH, Moerman DG, Bergthorsson U, Katju
V. 2018. Mutational and transcriptional landscape of spontaneous gene duplications
and deletions in Caenorhabditis elegans. *Proc. Natl. Acad. Sci.* 115:7386–7391.

Krasovec M, Eyre-Walker A, Grimsley N, Salmeron C, Pecqueur D, Piganeau G,
Sanchez-Ferandin S. 2016. Fitness Effects of Spontaneous Mutations in
Picoeukaryotic Marine Green Algae. *G3 Genes Genomes Genet.* 6:2063–2071.

Krasovec M, Eyre-Walker A, Sanchez-Ferandin S, Piganeau G. 2017. Spontaneous

Mutation Rate in the Smallest Photosynthetic Eukaryotes. *Mol. Biol. Evol.* 34:1770–1779.

Krasovec M, Lipinska AP, Coelho SM. 2022. Low spontaneous mutation rate in a complex multicellular eukaryote with a haploid-diploid life cycle. :2022.05.13.491831. Available from: https://www.biorxiv.org/content/10.1101/2022.05.13.491831v1

Krasovec M, Sanchez-Brosseau S, Grimsley N, Piganeau G. 2018. Spontaneous mutation rate as a source of diversity for improving desirable traits in cultured microalgae. *Algal Res.* 35:85–90.

Krasovec M, Sanchez-Brosseau S, Piganeau G. 2019. First Estimation of the Spontaneous Mutation Rate in Diatoms. *Genome Biol. Evol.* 11:1829–1837.

Krasovec M, Vancaester E, Rombauts S, Bucchini F, Yau S, Hemon C, Lebredonchel H, Grimsley N, Moreau H, Sanchez-Brosseau S, et al. 2018. Genome Analyses of the Microalga Picochlorum Provide Insights into the Evolution of Thermotolerance in the Green Lineage. *Genome Biol. Evol.* 10:2347–2365.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* 12:323.

Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows–Wheeler transform. *Bioinformatics* 26:589–595.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinforma. Oxf. Engl.* 25:2078–2079.

Lim J, Lee M, Son A, Chang H, Kim VN. 2016. mTAIL-seq reveals dynamic poly(A) tail regulation in oocyte-to-embryo development. *Genes Dev.* [Internet]. Available from: http://genesdev.cshlp.org/content/early/2016/07/14/gad.284802.116

Liu H, Zhang J. 2019. Yeast Spontaneous Mutation Rate and Spectrum Vary with Environment. *Curr. Biol. CB* 29:1584-1591.e3.

Liu Y, Borel C, Li L, Müller T, Williams EG, Germain P-L, Buljan M, Sajic T, Boersema PJ, Shao W, et al. 2017. Systematic proteome and proteostasis profiling in human Trisomy 21 fibroblast cells. *Nat. Commun.* 8:1212.

Loane M, Morris JK, Addor M-C, Arriola L, Budd J, Doray B, Garne E, Gatt M, Haeusler M, Khoshnood B, et al. 2013. Twenty-year trends in the prevalence of Down syndrome and other trisomies in Europe: impact of maternal age and prenatal screening. *Eur. J. Hum. Genet. EJHG* 21:27–33.

Loehlin DW, Carroll SB. 2016. Expression of tandem gene duplicates is often greater than twofold. *Proc. Natl. Acad. Sci.* 113:5988–5992.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15:550.

Lyle R, Gehrig C, Neergaard-Henrichsen C, Deutsch S, Antonarakis SE. 2004. Gene expression from the aneuploid chromosome in a trisomy mouse model of down syndrome. *Genome Res.* 14:1268–1274.

Lynch M, Ackerman MS, Gout J-F, Long H, Sung W, Thomas WK, Foster PL. 2016. Genetic drift, selection and the evolution of the mutation rate. *Nat. Rev. Genet.* 17:704–714.

Lynch M, Sung W, Morris K, Coffey N, Landry CR, Dopman EB, Dickinson WJ, Okamoto K, Kulkarni S, Hartl DL, et al. 2008. A genome-wide view of the spectrum of spontaneous mutations in yeast. *Proc. Natl. Acad. Sci.* 105:9272–9277.

Miao Z, Zhang T, Xie B, Qi Y, Ma C. 2022. Evolutionary Implications of the RNA N6-Methyladenosine Methylome in Plants. *Mol. Biol. Evol.* 39:msab299.

Moreau H, Verhelst B, Couloux A, Derelle E, Rombauts S, Grimsley N, Van Bel M, Poulain J, Katinka M, Hohmann-Marriott MF, et al. 2012. Gene functionalities and genome structure in Bathycoccus prasinos reflect cellular specializations at the base of the green lineage. *Genome Biol.* 13:R74.

Muyle A, Shearn R, Marais GA. 2017. The Evolution of Sex Chromosomes and Dosage Compensation in Plants. *Genome Biol. Evol.* 9:627–645.

Muyle A, Zemp N, Deschamps C, Mousset S, Widmer A, Marais GAB. 2012. Rapid De Novo Evolution of X Chromosome Dosage Compensation in Silene latifolia, a Plant with Young Sex Chromosomes. *PLOS Biol.* 10:e1001308.

Nagaoka SI, Hassold TJ, Hunt PA. 2012. Human aneuploidy: mechanisms and new insights into an age-old problem. *Nat. Rev. Genet.* 13:493–504.

Ness RW, Morgan AD, Vasanthakrishnan RB, Colegrave N, Keightley PD. 2015. Extensive de novo mutation rate variation between individuals and across the genome of Chlamydomonas reinhardtii. *Genome Res.* 25:1739–1749.

Peter J, De Chiara M, Friedrich A, Yue J-X, Pflieger D, Bergström A, Sigwalt A, Barre B, Freel K, Llored A, et al. 2018. Genome evolution across 1,011 Saccharomyces cerevisiae isolates. *Nature* 556:339–344.

Peterson JB, Ris H. 1976. Electron-microscopic study of the spindle and chromosome movement in the yeast Saccharomyces cerevisiae. *J. Cell Sci.* 22:219–242.

Qian W, Liao B-Y, Chang AY-F, Zhang J. 2010. Maintenance of duplicate genes and their functional redundancy by reduced expression. *Trends Genet. TIG* 26:425–430.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* 26:841–842.

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R project.org/.

Rajagopalan H, Lengauer C. 2004. Aneuploidy and cancer. *Nature* 432:338–341.

Scopel EFC, Hose J, Bensasson D, Gasch AP. 2021. Genetic variation in aneuploidy prevalence and tolerance across Saccharomyces cerevisiae lineages. *Genetics* 217:iyab015.

Selmecki A, Forche A, Berman J. 2006. Aneuploidy and isochromosome formation in drug-resistant Candida albicans. *Science* 313:367–370.

Sement FM, Gagliardi D. 2014. Detection of uridylated mRNAs. *Methods Mol. Biol. Clifton NJ* 1125:43–51.

Slobodin B, Bahat A, Sehrawat U, Becker-Herman S, Zuckerman B, Weiss AN, Han R, Elkon R, Agami R, Ulitsky I, et al. 2020. Transcription Dynamics Regulate Poly(A) Tails and Expression of the RNA Degradation Machinery to Balance mRNA Levels. *Mol. Cell* 78:434-444.e5.

Song MJ, Potter BI, Doyle JJ, Coate JE. 2020. Gene Balance Predicts
Transcriptional Responses Immediately Following Ploidy Change in Arabidopsis
thaliana. *Plant Cell* 32:1434–1448.

Stenberg P, Lundberg LE, Johansson A-M, Rydén P, Svensson MJ, Larsson J.
2009. Buffering of Segmental and Chromosomal Aneuploidies in Drosophila
melanogaster. *PLOS Genet.* 5:e1000465.

Subirana L, Péquin B, Michely S, Escande M-L, Meilland J, Derelle E, Marin B,
Piganeau G, Desdevises Y, Moreau H, et al. 2013. Morphology, Genome Plasticity,
and Phylogeny in the Genus Ostreococcus Reveal a Cryptic Species, O.
mediterraneus sp. nov. (Mamiellales, Mamiellophyceae). *Protist* 164:643–659.

Subtelny AO, Eichhorn SW, Chen GR, Sive H, Bartel DP. 2014. Poly(A)-tail profiling
reveals an embryonic switch in translational control. *Nature* 508:66–71.

Thompson SL, Compton DA. 2008. Examining the link between chromosomal
instability and aneuploidy in human cells. *J. Cell Biol.* 180:665–672.

Torres EM, Dephoure N, Panneerselvam A, Tucker CM, Whittaker CA, Gygi SP,
Dunham MJ, Amon A. 2010. Identification of aneuploidy-tolerating mutations. *Cell*
143:71–83.

Torres EM, Sokolsky T, Tucker CM, Chan LY, Boselli M, Dunham MJ, Amon A.
2007. Effects of aneuploidy on cellular physiology and cell division in haploid yeast.
Science 317:916–924.

Udagawa T, Swanger SA, Takeuchi K, Kim JH, Nalavadi V, Shin J, Lorenz LJ, Zukin
RS, Bassell GJ, Richter JD. 2012. Bidirectional control of mRNA translation and
synaptic plasticity by the cytoplasmic polyadenylation complex. *Mol. Cell* 47:253–
266.

Vandepoele K, Van Bel M, Richard G, Van Landeghem S, Verhelst B, Moreau H,
Van de Peer Y, Grimsley N, Piganeau G. 2013. pico-PLAZA, a genome database of
microbial photosynthetic eukaryotes. *Environ. Microbiol.* 15:2147–2153.

de Vargas C, Audic S, Henry N, Decelle J, Mahé F, Logares R, Lara E, Berney C, Le
Bescot N, Probert I, et al. 2015. Ocean plankton. Eukaryotic plankton diversity in the
sunlit ocean. *Science* 348:1261605.

Veitia RA, Potier MC. 2015. Gene dosage imbalances: action, reaction, and models.
Trends Biochem. Sci. 40:309–317.

Weill L, Belloc E, Bava F-A, Méndez R. 2012. Translational control by changes in
poly(A) tail length: recycling mRNAs. *Nat. Struct. Mol. Biol.* 19:577–585.

Williams BR, Prabhu VR, Hunter KE, Glazier CM, Whittaker CA, Housman DE,
Amon A. 2008. Aneuploidy affects proliferation and spontaneous immortalization in
mammalian cells. *Science* 322:703–709.

Worden AZ, Lee J-H, Mock T, Rouzé P, Simmons MP, Aerts AL, Allen AE, Cuvelier
ML, Derelle E, Everett MV, et al. 2009. Green evolution and dynamic adaptations
revealed by genomes of the marine picoeukaryotes Micromonas. *Science* 324:268–
272.

Yau S, Hemon C, Derelle E, Moreau H, Piganeau G, Grimsley N. 2016. A Viral
Immunity Chromosome in the Marine Picoeukaryote, Ostreococcus tauri. *PLoS*
Pathog. 12:e1005965.

Yau S, Krasovec M, Benites LF, Rombauts S, Groussin M, Vancaester E, Aury J-M,
Derelle E, Desdevises Y, Escande M-L, et al. 2020. Virus-host coexistence in
phytoplankton through the genomic lens. *Sci. Adv.* 6:eaay2587.

Yoon HS, Hackett JD, Ciniglia C, Pinto G, Bhattacharya D. 2004. A molecular
timeline for the origin of photosynthetic eukaryotes. *Mol. Biol. Evol.* 21:809–818.

Zhang X, Virtanen A, Kleiman FE. 2010. To polyadenylate or to deadenylate: that is
the question. *Cell Cycle Georget. Tex* 9:4437–4449.

Zhang Z, Presgraves DC. 2016. Drosophila X-Linked Genes Have Lower Translation
Rates than Autosomal Genes. *Mol. Biol. Evol.* 33:413–428.

Zhang Z, Presgraves DC. 2017. Translational compensation of gene copy number
alterations by aneuploidy in Drosophila melanogaster. *Nucleic Acids Res.* 45:2986–
2993.

Zhu YO, Siegal ML, Hall DW, Petrov DA. 2014. Precise estimates of mutation rate
and spectrum in yeast. *Proc. Natl. Acad. Sci. U. S. A.* 111:E2310-2318.

873
874 **Figure S17.** Coverage by 1 kb windows of the *Bathycoccus prasinos* MA line Bp37 and control
875 RCC4222.

884
Figure S18. Transcripts from duplicated chromosome C01 and non-duplicated chromosome C08
present similar poly(A) tail. Poly(A) tail measurement was performed using modified 3'RACE. PCR
amplification was performed using primers flanking poly(A) tail. Experiments was performed using total
RNA from RCC4222 line (control line) and Bp37B line (chromosome C04 duplicated). Illustrations
representing PCR amplification are present on the left panel. Black line represents the mRNA and the
orange line represent the ligated adapter used for reverse transcription and PCR amplification.