Mesenchymal-epithelial transition in lymph node metastases of oral squamous cell carcinoma is accompanied by ZEB1 expression

Kai Horny[1,2], Christoph Sproll[3], Lukas Peiffer[1,2], Frauke Furtmann[1,2,4], Patricia Gerhardt[1,2], Jan Gravemeyer[1,2], Nikolas H. Stoecklein[5], Ivelina Spassova[1,2,4], Jürgen C. Becker[1,2,4,*]

[1] Translational Skin Cancer Research, German Cancer Consortium (DKTK), 45141 Essen, Germany

[2] German Cancer Research Center (DKFZ), 69120 Heidelberg, Germany

[3] Department of Oral- and Maxillofacial Surgery, Medical Faculty, University Hospital of the Heinrich-Heine-University Düsseldorf, Germany

[4] Department of Dermatology, University Medicine Essen, 45141 Essen, Germany

[5] Department of General, Visceral and Pediatric Surgery, Medical Faculty, University Hospital of the Heinrich-Heine-University Düsseldorf, Düsseldorf, Germany

* Correspondence: j.becker@dkfz.de

Competing Interests

JCB is receiving speaker's bureau honoraria from Amgen, Pfizer, MerckSerono, Recordati and Sanofi, is a paid consultant/advisory board member/DSMB member for Boehringer Ingelheim, InProTher, MerckSerono, Pfizer, 4SC, and Sanofi/Regeneron. His group receives research grants from Bristol-Myers Squibb, Merck Serono, HTG, IQVIA, and Alcedis. None of these activities are related to the present manuscript. The other authors including K.H., C.S., L.P., F.F., P.G., J.G., N.S., I.S. declare no conflict of interest.

Abstract

Oral squamous cell carcinoma (OSCC) frequently metastasizes to the regional lymph nodes but only occasionally beyond. Initial phases of metastasis are associated with an epithelial-mesenchymal transition (EMT), the consolidation phase is associated with mesenchymal-epithelial transition (MET). This dynamic is referred to as epithelial-mesenchymal plasticity (EMP).

To capture EMP-associated heterogeneity in OSCC, we performed single-cell RNA sequencing of 5 primary tumors and 9 lymph node metastases and analyzed publicly available data of 9 additional primary tumors for a total of 7,263 carcinoma cells. Focusing initially on one metastasis to avoid inter-patient heterogeneity as confounder, we identified OSCC cells expressing genes characteristic of different epithelial and partial EMT stages. RNA velocity information together with the increase in inferred copy number variations indicated a progressive trajectory towards epithelial differentiation. Extension to all samples revealed a less stringent but essentially similar pattern. Interestingly, the epithelial differentiation in OSCC metastases is associated with higher activity of EMT-activator ZEB1, which was confirmed by co-expression of ZEB1 and cornifin-B by immunohistochemistry.

In summary, we show that EMP enables different EMT and epithelial phenotypes of OSCC cells, which are endowed with capabilities essential for the different stages of the metastatic process, including maintenance of cellular integrity.

Frequently used abbreviations

CNV   Copy number variation

DC   Dendritic cell

DEG   Differentially expressed genes

EC   Endothelial cell

ECM   Extracellular matrix

EMP   Epithelial-mesenchymal plasticity

EMT   Epithelial-mesenchymal transition

FFPE   Formalin-fixed, paraffin-embedded

GSEA   Gene set enrichment analysis

GSVA   Gene set variation analysis

HNSCC   Head and neck squamous cell carcinomas

HPV   Human papillomavirus

HTO   Hashtag oligo

IHC   Immunohistochemistry

MET   Mesenchymal-epithelial transition

MSigDB   Molecular signature database

OSCC   Oral cavity squamous cell carcinoma

PC   Principal component

pEMT   Partial EMT

scRNAseq   Single-cell RNA sequencing

SNN   Shared-nearest neighbor

UMAP   Uniform manifold approximation and projection

Introduction

Head and neck squamous cell carcinoma (HNSCC) is the sixth most common cancer worldwide, with 890,000 new cases and 450,000 deaths in 2018 [1]. The survival for HNSCC patients has improved modestly over the past decades; however, this improvement is partially attributable to the emergence of human papillomavirus (HPV)-associated HNSCC which has a better prognosis than HPV-negative tumors. One of the HPV-negative HNSCC subtypes is oral cavity squamous cell carcinoma (OSCC) which is mainly associated with tobacco and alcohol abuse [1]. OSCCs are often diagnosed at an early stage owing to the patient's self-identification of the mass lesion and symptoms, still regional lymph node metastases are frequent; thus, surgical removal of primary tumor is accompanied by neck dissection and radiotherapy [2]. Given the morbidity associated with this combined intervention, there is a need to identify molecular biomarkers to predict the presence of lymph node metastases and to prognosticate survival.

Invasion and metastasis becomes possible in many epithelial tumors through an epithelial-mesenchymal transition (EMT), i.e., a reactivation of an embryonic developmental program in which cells acquire migratory and invasive properties [3]. In EMT, epigenetic, transcriptional, and post-translational changes cause epithelial cells to break down the strong homotypic cell-cell junctions and adopt a mesenchymal morphology [4]. It is important to note that EMT should not be understood as a clearly defined process, but rather as many dynamic and complex processes, which may vary depending on tumor entity, stage, and microenvironment [4-6]. Thus, expression of EMT-related genes and their regulating transcription factors is highly heterogeneous even within one cancer entity, between patients, different lesions from one patient and between individual cancer cells within one lesion [4, 7]. Since it is a continuous, dynamic, and reversible process cancer cells can adopt a multitude of intermediate or partial states, e.g., epithelial to more mesenchymal or partial EMT (pEMT) states [5, 8-11]. Therefore, it has recently been recommended that this EMT continuum should rather referred to as epithelial-mesenchymal plasticity (EMP) [4, 10].

To capture the EMP-associated heterogeneity of cancers, single-cell analyses appear to be

the most appropriate approach, however, to date most EMP-related single-cell studies are

based on controlled *in vitro* and *in vivo* experiments [5, 6, 9, 12, 13]. Particularly in HNSCC

only few studies scrutinized EMP within freshly isolated tumor samples [8, 14, 15]. Especially

noteworthy is the landmark paper by Puram *et al.* They characterized 2,215 malignant cells

from 18 patients and discovered several pEMT states with high variability in EMP-related gene

expression [8].

In the work presented here, we investigated the phenotypic heterogeneity of 5 primary and 9

regionally metastatic OSCC lesions isolated from 7 patients using multiplexed single-cell RNA

sequencing (scRNAseq). In addition, we used a recently published series of scRNAseq data

from primary HNSCC that included 9 OSCC tumors to put our observations on an even broader

data base [15]. Our results not only confirm the EMP-associated heterogeneity of cancer cells

in primary and metastatic OSCC, but also demonstrate mesenchymal-epithelial transition

(MET) in established lymph node metastases. Surprisingly, we observed a high activity of

EMT-activator ZEB1 in metastatic OSCC cells with epithelial differentiation, a notion which was

confirmed by co-expression of ZEB1 and cornifin-B protein in individual tumor cells.

Results

Single-cell gene expression signatures of tumor cells from a single metastasis show several predominant, but not necessarily exclusive, functional phenotypes.

To avoid inter-patient heterogeneity as a confounding factor, we first focused on the analysis of a single OSCC metastasis to develop hypotheses which would be subsequently tested in the entire cohort. For this, we chose a metachronous lymph node metastasis that was removed one year after the primary tumor, because we assumed that consolidation processes are particularly pronounced in this longer existing metastasis. Multiplexed scRNAseq recovered 4,121 cells that could be assigned to the following cell types: 1,906 (46.8 %) tumor cells, 1,186 (29.1 %) fibroblasts, 507 (12.4 %) dendritic cells (DCs), 375 (9.2 %) macrophages and 102 (2.5 %) endothelial cells (ECs) (Figure 1A). The absence of T- or B-cells was in line with the histology showing completely disrupted lymph node structures and only occasional tumor-infiltrating lymphocytes (Supplementary Figure 1). OSCC cells were identified both by the presence of copy number variations (CNVs) inferred from scRNAseq data as well as the expression of epithelial markers including *S100A2, S100A14,* cytokeratins (*KRT5*, *KRT14*, *KRT17)* and stratifin (*SFN*) (Supplementary Figure 2).

Detailed phenotyping of the cancer cells identified several clusters to which we could assign the following predominant functional phenotypes that differ in their EMT state, immunomodulatory capacity, as well as their response to hypoxia, stress, and metabolic constraints (Figure 1B-D). However, the predominance of a functional phenotype does not preclude additional traits. Specifically, 515 cells exhibited a pEMT phenotype by expressing a mixture of epithelial and mesenchymal genes such as matrix metallopeptidases (*MMPs*), caveolin-1 (*CAV1*) and *CXCL14* that were previously described in pEMT signatures and are enriched in the EMT hallmark gene set from the molecular signatures database (MSigDB) (Supplementary Figure 3A) [8, 16]. In contrast, 385 cells showed higher expression of genes associated with epithelial differentiation such as *S100A7/A8/A9,* the keratinocyte envelope protein cornifin-B (*SPRR1B*), and various cytokeratins (e.g., *KRT6B* and *KRT16*). The EMP-

related gene expression patterns are correlating with established signatures from the EMTome

database (Supplementary Figure 3B-E) [7]. Interestingly, both gene expression signatures

were present in a mixed cluster with 184 cells characterized by the high expression of cell-

cycle related genes (despite the fact that we applied cell cycle regression).

With respect to cell clusters whose gene expression was not predominately associated with

EMP, 268 cells exhibited an immune-regulatory phenotype enriched for genes associated with

cytokine-mediated responses and higher expression of the chemokines *CXCL1/2/3/8* and

CCL20 and for 554 cells the gene expression pattern suggests adaptation to environmental

factors. In detail, 51 cells had higher expression of transcription factors *FOS* and *JUN*

suggesting a stress response. 310 cells can be assumed to respond to hypoxic conditions in

the tumor based on the higher expression of *NDRG1* and *EGLN3* both of which are regulated

by oxygen levels [17, 18]: *NDRG1* regulates stress response and p53-mediate caspase

activation [17], and *EGLN3* has an important role in hypoxia-inducible factor 1 alpha (HIF1α)

regulation through prolyl hydroxylation [18]. 193 cells expressed genes associated with amino

acid metabolism, starvation response and mTORC1 signaling, i.e., a regulator of mitochondrial

metabolism [19]. The upregulated genes *ASNS*, *PSAT1* and *PHGDH* integrate the metabolites

of serine and glycine metabolism into glycolysis and therefore fuel glycolysis with amino acids

[20]; hence, these cells appear to be adapted to low-glucose conditions.

OSCC cells in lymph node metastases undergo mesenchymal-epithelial transition

We next focused on possible dynamics within the predominantly EMP-related cancer cell

clusters using a higher resolved shared-nearest neighbor (SNN) clustering. We defined 4

pEMT clusters (pEMT-1 to 4), 4 clusters of more epithelial differentiated cells (epi-1 to 4) and

one cluster with mixed phenotypes (mix) (Figure 2A). pEMT-1 is enriched for genes involved

in coagulation and play a role in angiogenesis as *THBS1*, *CYR61* and *F3* (Figure 2B,

Supplementary Figure 4A). pEMT-2 and 3 showed higher expression of extracellular matrix

(ECM) remodeling genes, but pEMT-2 also higher expressed cytokeratin *KRT15* and

chemokine *CXCL14* while pEMT-3 higher expressed the serine protease inhibitor *SERPINA1*

and podoplanin (*PDPN*) which mediates efficient ECM degradation by controlling invadopodia

[21]. Of the more epithelial differentiated cell clusters, epi-2's expression profile is closest to

the pEMT cluster: higher expression of *MMP1* and lower expression of *SPRR1B* and

S100A8/A9 than epi-1, 3 and 4 (Figure 2B, E). Epi-3 showed higher expression of *S100A7* and

KRTDAP, whereas epi-4 has higher expression of kallikreins (*KLK6/7*), prostate stem cell

antigen (*PSCA*) and adipogenesis regulatory factor (*ADIRF*). Both *ADIRF* and *PSCA* play a

role in prostate cancer and *PSCA* is also reported as highly expressed in mucosal tissue, but

less in HNSCC [22-24].

To gain a better understanding on the gene expression dynamics we estimated RNA velocity,

which predicts the short-term future development in gene expression of individual cells using

the ratio of spliced and non-spliced mRNA counts (Figure 2C). This analysis revealed that

epithelial differentiated cells were strongly developing towards cluster epi-4, while most other

cells did not show a strong uniform developmental direction. Tracking the developmental

pathway within the metastasis by trajectory analysis across all EMP-related clusters revealed

a major developmental axis between pEMT and epithelial differentiated cells that is diversifying

within each end (Figure 2D, E, Supplementary Figure 4B). To confirm that we the found

progressive epithelial differentiation of the metastatic cells represents a MET, we inferred

CNVs from the scRNAseq data. This demonstrated an increased number of copy number gains

on chromosome 1, 8, 17, and 19 within epithelial differentiated cell clusters epi-1, epi-3, and

epi-4 (Figure 2F, Supplementary Figure 4C). It should be noted, however, that CNVs on

chromosomes 1 and 17 are associated with upregulated epithelial genes in close genomic

proximity; thus, these two copy number gains may not represent true genomic CNVs, but rather

reflect the high expression of these genes in epithelial differentiated cells (Supplementary

Figure 4D, E).

Intra-tumoral heterogeneity of OSCC is driven by EMP

To test whether our observation that OSCC cells in one lymph node metastasis undergo a

mesenchymal-epithelial transition is generally valid we extended our analyses by adding 5

primary tumors and 8 matched lymph node metastases from 6 patients (Table 1). From the publicly available scRNAseq data set on primary HNSCC tumors published by Kürten *et al*. [15] we chose to include the 9 HPV-negative primary tumors, which were all but one originating from the oral cavity (HN07 originated from the larynx) in our analysis. In total, we analyzed 7,263 cancer cells from 16 different patients (Figure 3A). Importantly, the frequency of cancer cells was unevenly distributed across samples, which could not be explained by differences in tumor cell content across samples as determined by histopathology (Supplementary Figure 1). Since the recovery of cells by scRNAseq is also low, the stability of epithelial tumor cell assemblies, which may not be sufficiently broken up by dissociation protocols, likely interfered with the generation of OSCC single-cell suspensions. In addition, as expected from the inter-patient heterogeneity, cancer cells were clustered based on their gene expression by patient rather than functional phenotype (Figure 3A, B). Thus, we accounted for the patient-specific effects with batch-corrected principal components (PCs) using the harmony R package which indeed resulted in a clustering by functional phenotypes [25] (Figure 3C, Supplementary Figure 5A-C). For initial annotation of the phenotype of each cluster, we were guided by the gene signatures previously identified in the indicator sample, with additional predominantly immunoregulatory phenotypes found. EMP-related phenotypes were present in all, but one tumor sample with only one cell (Figure 3D). Nevertheless, direct comparison of the frequencies of the predominant phenotypes in the pooled scRNAseq dataset proved difficult to interpret due to differences in cell numbers, and patient-specific gene expression patterns. To confirm the validity of our scRNAseq data, we examined the bulk transcriptome and deconvoluted the respective cell types for our 18 OSCC samples (Supplementary Figure 6). Using this approach, we observed a higher tumor and stroma content and less lymphocytes, which was expected as scRNAseq is known to be biased towards leukocytes. Therefore, we chose an alternative approach: to compare the EMP-related intrapatient heterogeneity of all analyzed tumor cells between patients, we calculated the similarities of the differential gene expression within each patient of all respective clusters of all patients. We first considered each patient individually and performed clustering, annotation, inferCNV and differential expression

analysis. As exemplified for the primary OSCC of patient HN01, tumor cell clusters had the same inferred CNVs and were annotated based on their phenotype using again the indicator sample as a guide (Figure 3E, Supplementary Figure 5D, E). The cosine similarity between patient-specific clusters demonstrated that within each patient, the heterogeneity in EMP is most prominent and epithelial differentiated phenotypes are profoundly different from most other, especially pEMT clusters (Figure 3F, Supplementary Figure 5F). Indeed, all pEMT phenotypes are very similar to each other and show a large overlap of the gene expression patterns with predominantly immune- and metabolic-related clusters. In epithelial differentiated cells, the most upregulated genes are *S100A8* and *S100A9*, encoding calprotectin) and *SPRR1B*, all members of the epidermal differentiation complex [27]. Of note, hypoxia- and stress-related heterogeneity is similar between patients suggesting a reactive response rather than an aspect of tumor evolution.

EMT-related transcription factor ZEB1 is highly active in metastatic epithelial differentiated OSCC cells

The transcription factors ZEB1/2, TWIST1/2, Snail (SNAI1) and Slug (SNAI2) are key for regulation of EMP [7, 27]. While mRNA expression of *SNAI2* within single OSCC cells was recently reported, the others were not detected [8]. Here, we confirm this observation as we detected *SNAI2* mRNA in almost half of the OSCC cells (Figure 4A). However, detection of low expressed genes such as transcription factors by scRNAseq, especially in 10X genomics technology, becomes unreliable due to dropout effects. Also, the activity of transcription factors is often not reflected by the dynamics of their mRNA expression alone, as their activity depends additionally on protein stability and posttranslational modifications; for example, ZEB1 protein is more stable than Snail [28]. To circumvent this problem, we inferred the activity of these transcription factors based on the mRNA expression profile of their target genes using the algorithm VIPER with regulons defined by DoRothEA database [29-31]. Using this approach, we were able to detect high activities of ZEB1, ZEB2, Snail and Slug in OSCC cells of different patients with varying EMP phenotype (Figure 4B). Since, on the one hand, the observation that epithelial differentiation in OSCC metastases is associated with higher activity of the EMT

activator ZEB1 was unexpected and, on the other hand, the method used to derive transcription factor activities appears to maybe overestimate the activity for transcriptional repressors, we addressed the plausibility of this observation (Figure 4C, Supplementary Figure 7). Consistent with the fact that one of the main functions of ZEB1 is the downregulation of E-cadherin [32] (*CDH1*), OSCC cells generally showed low expression of *CDH1* (Figure 4B). Next, we investigated the expression of ZEB1 protein by immunohistochemistry (IHC) in all 14 tumor lesions of our cohort. Since we observed nuclear ZEB1 expression in similar tumor areas as high cytoplasmic cornifin-B expression, which served as a marker of epithelial differentiation (Figure 4D, E), we validated the co-expression of ZEB-1 and cornifin-B in the same cell by immunofluorescence double-staining (Figure 4F). In line with our scRNAseq data, colocalization of both proteins was observed in a fraction of cancer cells in 9/14 (64 %) samples, double positive cells were more frequently observed in lymph node metastases (7/9, 78 %; Table 1).

Discussion

EMT represents the reactivation of an embryonic developmental program in which cells acquire migratory and invasive properties, i.e., prerequisites for invasion and metastasis of cancer [3, 33, 34]. Thus, in early stages of metastasis, tumor cells undergo EMT, whereas in established metastases the reverse process *aka* MET is also observed [35, 36]. To assess the EMP-associated heterogeneity among OSCC cells and gain some insight into the dynamics of this process, we examined the transcriptomes of 7,263 individual carcinoma cells isolated from primary or metastatic OSCC. Our results not only confirm the EMP-associated heterogeneity of cancer cells in primary and metastatic OSCC, but also demonstrate progressive MET in established lymph node metastases. Interestingly, the epithelial differentiation in OSCC metastases is associated with higher activity of EMT-activator ZEB1, which was confirmed on protein level by co-expression of ZEB1 and cornifin-B in individual tumor cells by immunofluorescence staining.

EMP appears to be the main driver of cellular heterogeneity within OSCC: detailed phenotyping

of cancer cells identified several clusters whose predominant functional phenotypes

corresponded to different EMP states, ranging from a pEMT to an epithelial differentiated state.

However, particularly the pEMT phenotypes could be overlaid by features related to

angiogenesis, ECM remodeling, metabolic adaptations, stress, and interaction with the

immune system. Metabolic adaptations include response to environmental limitations as

hypoxia and low glucose. Low glucose conditions are counteracted with upregulation of genes

related to amino acid metabolism that fuel into glycolysis [20]. While previous studies

suggested that the activity of specific metabolic pathways in OSCC varies widely among

patients [37], we observed that hypoxia- and stress-related gene expression patterns are

rather similar between patients supporting the notion of an reactive response rather than an

aspect of individual tumor evolution.

In terms of EMP dynamics, it is assumed that cells may transit from one EMP state to another

along a continuous spectrum of changes. Currently, however, it is also discussed if long-lived

phenotypes representing discrete EMP states prevail [3, 5, 6, 8, 9, 11-13]. Most studies

supporting continuous transitions are based on *in vitro* or preclinical *in vivo* models that may

not fully reflect the complexity of the tumor and its microenvironments [5, 6, 9]. Indeed, human

in situ or *ex vivo* studies suggested distinct EMP states; however, these approaches do not

fully capture cellular dynamics [8, 12, 34]. Comparing the intratumoral heterogeneity we

demonstrated that within each patient, the EMP-driven heterogeneity is most prominent and

epithelial differentiated phenotypes are profoundly different from pEMT clusters suggesting

that these states may be rather static. Moreover, gene expression dynamics estimated by RNA

velocity demonstrated epithelial differentiated cells were strongly developing towards a more

pronounced epithelial differentiation with an increasing expression of genes of the epidermal

differentiation complex [27]. Of note, OSCC cells with a pEMT phenotype did not show such a

uniform developmental direction. The assumption that epithelial differentiated metastatic cells

developed later than pEMT cells, i.e., undergoing MET, is supported by an increasing number

of inferred copy number gains towards increasing epithelial differentiation even if accounting

for the limitations of this approach.

However, we cannot conclude whether MET happened within the metastasis or primary

malignancy, as our data reflects the tumor heterogeneity within a specific timepoint of tumor

evolution. Multiple disseminated tumor cells reflecting the observed EMP heterogeneity might

have migrated collectively from the primary tumor that could be crucial for metastatic

consolidation, as we observed a similar EMP heterogeneity also within primary tumors.

The high transcriptional activity of EMT-activator ZEB1 in epithelial differentiated OSCC cells

in both primary and metastatic tumor lesions was unexpected, even considering that the

scRNAseq data inferred transcription factor activities are biased towards transcriptional

repressors. In the case of primary tumors this may be interpreted as the incipient EMT, but this

would not work for the metastatic lesions in which ZEB1 activity was associated with a

progressive epithelial differentiation. Indeed, in previous studies depletion of ZEB1 has driven

tumor cells from pEMT towards an epithelial phenotype [38, 39]. However, depletion of Zeb1

also reduces phenotypic variability of cancer cells particular their phenotypic/metabolic

plasticity [38]. While it is well established that ZEB1 together with microRNAs stabilizes EMT

through a feedforward loop, this loop could also induce epithelial differentiation based on

environmental factors [40]. In addition to the transcriptional repressor activity, ZEB1 has been

demonstrated to induce the epithelial differentiation marker cornifin-B in response to IL-1β and

IFN-γ [41]. We not only demonstrate the coincidence of *SPRR1B* mRNA expression and ZEB1

activity, but also the co-localization of cornifin-B and ZEB1 protein expression in OSCC lymph

node metastases. Thus, although ZEB1 activity is crucial for the induction of the pEMT state,

it does not seem to completely prevent partial epithelial differentiation. Remarkably, no relevant

differences in *CDH1* expression were detected between the different EMP states in the

metastatic OSCC lesions. Therefore, the epithelial differentiated phenotypes we observed

most closely correspond to a partial epithelial differentiation analogous with the observed

pEMT phenotypes. We speculate that the driving force behind this EMP-associated

heterogeneity of OSCC cells is to maintain cellular integrity. For example, ZEB1 is an ATM-
substrate linking ATM and CHK1, promoting homologous recombination-dependent DNA
repair and thereby protecting cells from genotoxic stress whereas expression of keratin
intermediate filaments helps to protect cells from stress associated apoptosis [42, 43].

In summary, heterogeneity within OSCC is dominated by EMP differences resulting in distinct
pEMT and epithelial differentiated phenotypes, but particularly the pEMT phenotypes can be
accompanied by features related to metabolic adaptations, stress, and interaction with the
immune system. These phenotypes are likely to endow capabilities essential for the different
stages of the metastatic process, including maintenance of cellular integrity. Thus, further
studies are required and already arising to unravel the factors driving these evolutionary
processes in tumor cells [44].

Patients, Material and Methods

Tissue samples

Twenty tissue samples that included primary tumors, clinically apparent lymph node metastases, and potentially affected lymph nodes from 7 OSCC patients treated at the Department of Oral-, Maxillo- and Plastic Facial Surgery, University Hospital of the Heinrich-Heine-University Düsseldorf, Germany, were used for histology, IHC, immune fluorescence as well as bulk and single-cell transcriptome analyses. The clinical details are provided in table 1. Written informed consent was obtained and the Ethics Committee of the Medical Faculty of the Heinrich-Heine-University Düsseldorf (#3090) approved the study. All procedures involving human participants are in accordance with the ethical standards of the institutional and research committee and with the Helsinki Declaration.

Histology and IHC

Hematoxylin and Eosin (H&E) and IHC were performed on 4 μm formalin-fixed, paraffin-embedded (FFPE) sections. H&E staining was performed using standard protocols (Supplementary Figure 1). Whole-slide imaging was performed using Zeiss Axioscan 7 and 10x magnification (Carl Zeiss Microscopy Deutschland GmbH, Oberkochen, Germany).

IHC using the rabbit polyclonal antibodies anti-SPRR1B (Cat. No.: SAB1301567-400UL, Sigma Aldrich, Darmstadt, Germany) anti-ZEB1 antibody (Cat. No. HPA027524-25UL, Sigma Aldrich) was performed as previously described [45]. Briefly, after sections were deparaffinized for 60 min at 60°C and rehydrated, sections were incubated for 15 min in an inverter microwave oven with antigen retrieval buffer pH 9 for anti-SPRR1B and pH 6 for anti-ZEB1. After 3x2 min washes with Tris-buffered saline with 0.1% Tween (TBST) sections were incubated for 8 min with 3% peroxidase and after an additional washing step for 30 min with 3% bovine serum albumin (BSA) in TBST. For single staining sections were subjected for 1h to anti-SPRR1B in 1:600 dilution or overnight to anti-ZEB1 in a 1:500 dilution, both at room temperature. Afterwards, the secondary anti-rabbit HRP Polymer was applied for 30 min, followed by 1:20 diluted 3,3'-Diaminobenzidin (DAB) for 10 min and 1:10 diluted hematoxylin for 3 min with

washing steps with TBST in between before a final washing with tap water for 3 min before

fixation. For multiplexed antigen detection, the OpalTM chemistry system (Akoya Biosciences,

Marlborough, MA, USA, Cat. No.: OP7TL4001KT) was used according to the manufacture's

description. Briefly, after deparaffinization and fixation, we processed the sections for 15 min

with retrieval buffers in an inverter microwave oven. Then, we incubated them with antibody

diluent for 10 min at room temperature, followed by incubation with the anti-SPRR1B antibody

for 30 min. Next, Opal Polymer horseradish peroxidase (HRP) secondary antibody solution

with the respective chromogen was applied for 10 min, antibodies were removed by microwave

treatment and the staining with anti-ZEB1 antibody was performed. Slides were lastly

incubated with 4′,6-diamidino-2-phenylindole (DAPI) for 5 min.

Single-cell RNA sequencing

Samples were processed immediately after surgery and temporarily stored for transport at 4°C

in tissue storage solution (Miltenyi Biotec, Bergisch Gladbach, Germany) as previously

described. Briefly, samples were dissociated into single-cell suspensions using the

gentleMACS Dissociator (Cat. No. 130-093-235, Miltenyi Biotec, Bergisch Gladbach,

Germany) with program "h_tumor_01", followed by 2x program "h_tumor_02" in 4.7 ml RPMI

1640 (Cat. No. P04-16500, PAN-Biotech) and an enzyme mix consisting of 200µl Enzyme H,

100 µl Enzyme R and 25 µl Enzyme A (Cat. No. 130-095-929 Miltenyi Biotec). Afterwards,

single-cell suspensions were reconstituted and washed thrice with 0.05 % BSA phosphate-

buffered saline (PBS) and filtered through a 100 µl cell strainer.

In cases multiple samples of one patient had to be analyzed (Table 1), antibody hashing for

multiplexing of samples was performed according to manufacturer's protocol. Briefly, 1 µg of

the respective TotalSeq anti-human hashtag antibody was used to incubate a maximum of ca.

2 million cells for 30 min at 4 °C (Cat. No. 394601, 394603, 394605 and 394661, 394663,

394665, respectively, Biolegend, San Diego, CA, USA). After 3 washes with PBS with 0.05 %

BSA respective cell suspensions were mixed prior to single-cell RNA library preparation. In

short, both unhashed and hashed single-cell suspensions were barcoded and processed with

the microfluidic system of 10x Genomics Chromium v2.0 platform as described in the manufacturer's protocols (10x Genomics, Leiden, Netherlands). Due to a change of system, both the 3' technology including Chromium Single Cell 3' Library & Gel Bead Kit version 2 (Cat. No. 120237), Chromium Single Cell A Chip Kit (Cat. No. 120236), Chromium i7 Multiplex Kit (Cat. No. 120262) as well as the 5' technology including Chromium Single Cell 5' Library & Gel Bead Kits version 2 (Cat. No. 1000263), Chromium Next GEM Chip K Single Cell Kit (Cat. No. 1000286), Dual Index Kit TT set A (Cat. No. 1000215) were used; for library construction the Chromium Single Cell 3'/5' Library Construction Kit (Cat. No. 1000020) was applied. After library preparation, the library from patient one was sequenced with an Illumina HiSeq 4000 (Illumina, Berlin, Germany) at the DKFZ Genomics and Proteomics Core Facility in Heidelberg and all other libraries were sequenced on an Illumina NovaSeq 6000 (Illumina, Berlin, Germany) in three runs (Run 1: patient 2,4,5; Run2: patient 3, Run3: Patient 6 and 7) at the West German Genome Center in Cologne.

Bulk transcriptome analysis

The bulk transcriptome was analyzed using a quantitative nuclease protection assay from the HTG Transcriptome Panel (HTP) according to the manufacturers protocol (Cat. No. HTG-001-224, HTG Molecular Diagnostics, Tucson, AZ, USA). Briefly, the tumor areas were macro-dissected as depicted in Supplementary Figure 1 from 4 µm FFPE sections and subjected to Proteinase K and DNase digestion. Next, the quantitative nuclease protection assay was performed using the HTG EdgeSeq Processors before adapters and sample tags were added in a PCR amplification. The resulting libraries were sequenced using on an Illumina NextSeq 500/550 High Output Kit v2.5 (75 cycles) (Cat. No. 20024906, Illumina, Berlin, Germany).

The resulting FASTQ files were processed towards a gene expression count matrix using the HTG EdgeSeq Reveal Software version 4.0.1. Quality Control, normalization, and principal component analysis (PCA) were performed using R version 4.0.5. Sample 5 failed QC due to low number of sequenced reads and was removed from the analysis. Deconvolution was performed with web application of CIBERSORTx (https://cibersortx.stanford.edu/) using a

signature matrix derived from the gene expression count matrix of combined scRNAseq data

of the samples analyzed with HTP [46]. We filtered for genes that are expressed less than 5%

within the given tumor phenotypes and randomly included only 75% of T- and B-Cells for better

performance. The resulting signature matrix was used for imputing cellular fractions from the

counts-per-million normalized HTP data without any batch correction or quantile normalization

and 500 permutations.

Bioinformatic analysis of scRNAseq data

Preprocessing

Processing from FASTQ files towards the unfiltered count matrix (barcodes x genes) was

performed using Cellranger Software Suite version 3.1.0 and the human reference genome

build GRCh38, downloaded from 10x Genomics in version 3.0.0.

Cells were identified by evaluating quality criteria inspired by Luecken *et al.*, also see

Supplementary File 1 [47]. Cells were defined by having more than 500 UMIs, less than 10 %

mitochondrial gene expression and additionally for patient 3 and 5 having more than 30

housekeeper genes expressed. The filtered count matrices (cells x genes) were further

processed using Seurat version 4.0.1 and R version 4.0.5 [48]. Demultiplexing of hashed

libraries was performed choosing manual threshold of hashtag oligo (HTO) expression based

on quality assessments described in Supplementary File 1. We removed doublets that were

identified by demultiplexing of HTO expression matrices from all analyses.

Normalization and dimensionality reduction

Every time we performed analysis of a specific set of cells, e.g., only tumor cells or only a

specific patient, the respective set of cells is normalized using the SCtransform algorithm and

the 3000 most variable genes were selected for PCA [49]. During normalization, we regressed

for cell cycle scores and percentage of mitochondrial gene expression. Cell cycle scores and

phases were determined in Seurat using log-normalized RNA counts and S- and G2M-Phase

genes defined by Tirosh *et al*. [50]. When generating a uniform manifold approximation and

projection (UMAP) without patient-specific batch effect we used corrected PCs derived with

the harmony R package version 0.1.0 [54]. Based on the variance explained by each PC and

the respective ranked elbow plot we choose an appropriate number of PCs for UMAP

visualization and SNN clustering as implemented in Seurat. For deriving patient-specific

clusters for calculating the intratumoral cosine similarity we used the same resolution

parameter for comparability. UMAPs colored by specific gene expression were ordered by

expression values.

Tumor cell identification and phenotyping

We used a combination of SNN clustering, differential gene expression, gene set enrichment

analysis (GSEA) and expression of literature-based marker genes for identification and

description of tumor cell phenotypes. Malignant cells were first identified by high epithelial gene

expression as for example cytokeratins and inferred CNVs (Supplementary Figure 2). CNVs

were inferred using the R package inferCNV version 1.6.0 with the not normalized, filtered

count matrix including all cells as input and algorithm run in "samples" mode . Inferred CNVs

of mitochondrial genes were excluded. Further, we excluded cells with ambiguous cell type

marker expression. For example, within the tumor cells of the lymph node metastasis from

patient 1, we observed few cells expressing genes typical for fibroblasts and DCs that were

subsequently excluded from tumor-specific analysis. Similarly, we removed T-Cells, B-Cells,

mast cells, fibroblasts, muscle cells, melanocytes, and other Immune cells from the tumor cell

subsets. Differential gene expression was performed by calculating the log2 foldchanges

between one cluster and all other cells from the subset based on log-normalized data using

NormalizeData function and scale factor of 10,000. We filtered for genes with log2 foldchange

greater than 0.25 and a minimum percentual expression of at least 10 % within the cluster or

all other cells. For calculating the cosine similarity, we did not filter the log2 foldchanges. GSEA

was performed using the "fgsea" R package version 1.16.0, log2 foldchanges from differential

gene expression and gene ontology biological processes (GO:BP,C5 v7.1) as well as hallmark

gene sets (H, v7.1) downloaded from MSigDB database [16, 51, 52]. Gene sets were included

if they have at least 15 genes or at maximum 500 genes within the gene set using 10,000

permutations.

For deriving epithelial differentiating and pEMT gene signatures of patient 1 we calculate

foldchanges between the "epi" and "pEMT" cluster and included genes with log2 foldchanges

greater or lesser than 1, respectively, and with at least 10 % of either epi or pEMT cells

expressing that gene. Gene set variation analysis (GSVA) was performed using the R package

GSVA version 1.38.2 with default settings, i.e., gaussian kernel [53]. As input, EMTome

signatures, the pEMT and epithelial differentiation 1 and 2 signature from Puram *et al.* [8] and

the three EMT and the epithelial senescence signatures from Kinker *et al.* were used [12].

Trajectories were inferred using SlingShot version 1.8.0 with malignant cell clusters as shown

in Figure 2A of patient 1 and the first 20 PCs [54]. RNA velocity was inferred using the VeloCyto

python and R package (version 0.6) [55]. Creation of the loom file was done using default

options and gene annotations as used for Cellranger processing. Genes were filtered based

on the minimum average expression magnitude with a threshold of 0.05 for spliced and 0.02

for unspliced reads. Velocity estimates were calculated using as distance the inverse

correlation coefficient of the PC embedding correlation matrix, the top/bottom 2 % quantiles for

gamma fit, 50 neighboring cells projecting 1 deltaT into the future and projected on the UMAP

using 200 neighbors and 50 grid points.

Transcription factor activity was inferred using the VIPER algorithm (version 1.24.0) and

regulons from the DoRothEA database (version 1.2.2) [29-31]. Hierarchical clustering in the

heatmaps was performed using Euclidean distance and ward.D2 method unless otherwise

noted. Visualization was performed using ggplot2 version 3.3.3 and ComplexHeatmap version

2.6.2 [56, 57].

Reanalysis of HNSCC dataset from Kürten *et al*.

From the publicly available scRNAseq data set on primary HNSCC tumors published the

FASTQ files of CD45 negative and HPV-negative libraries from Kürten *et al.* were downloaded

from the sequencing read archive (SRA) accessed through gene expression omnibus (GEO)

accession ID GSE164690 in BioProject PRJNA691564 [15]. The data sets were analyzed the

same as described above. HPV-negative samples were chosen for comparability to the OSCC

dataset. However, the HN07 tumor originated from the larynx, while all other samples indeed

originated from the oral cavity. We adjusted the cell identification thresholds based on our

evaluation criteria pooled for all libraries: cells were defined by having more than 175 genes

expressed, less than 10 % mitochondrial gene expression and more than 60 housekeeper

genes expressed (see Supplementary File 1).

Supplementary Information

Supplementary Figure 1: Histology of OSCC primary and metastatic tumors.

Supplementary Figure 2: | Cell type identification by differential expression and inferred CNVs for a metachronous lymph node metastasis.

Supplementary Figure 3: PEMT and epithelial differentiating gene expression signatures are comparable to previously published EMT signatures.

Supplementary Figure 4: Extended analysis of the tumor phenotype characterization for the lymph node metastasis of patient 1.

Supplementary Figure 5: Malignant phenotypes characterized across all analyzed patients.

Supplementary Figure 6: Bulk transcriptomes reveal the cellular composition of OSCC.

Supplementary Figure 7: Inferred transcription factor activity might be biased by activator or repressor function.

Table 1: Clinical and sequencing information from OSCC patients.

File 1: Supplementary Materials and Methods

Data availability

The processed datasets generated and analyzed during the current study are available in the GEO database with accession id GSE195655. The raw FASTQ files are available upon request due to privacy reasons. The high-resolution H&E Images are available through figshare under the DOI: https://doi.org/10.6084/m9.figshare.20905837.v1.

Code availability

Code used for analysis is available at https://github.com/sci-kai/single_cell_EMP.

Acknowledgements

We thank Önder Bozdogan, Translational Skin Cancer Research, German Cancer Consortium (DKTK), Essen, Germany, for his histological expertise and Linda Kubat, Department of Dermatology, University Medicine Essen, Germany, for helping with microscopy. We thank Christina H. Scheel, Skin Cancer Center, Department of Dermatology, Ruhr-University Bochum, Germany, for her helpful remarks to the manuscript. We further thank Yvonne Krause and Sophia Berger, both Department of Pathology, University Medicine Essen, Germany, for their help and guidance with HTG EdgeSeq. We further thank the DKFZ Genomics and Proteomics Core Facility and the West German Genome Center in Cologne for providing Illumina sequencing and related services.

Author Contributions

All authors approved the final version and agreed to be accountable for all aspects of the work. KH was responsible for data analysis, data curation, visualization, interpretation, conceptualization, drafting the manuscript and updating revisions. IS, LP and PG contributed by performing wet lab experiments including single-cell dissociation, scRNAseq and library preparation. FF established and performed histological, IHC and bulk transcriptome analysis. CS contributed with sample acquisition, interpretation and revising the manuscript. NS contributed by manuscript reviewing and interpretation. JG further contributed to data analysis, interpretation, and visualization. JCB was contributing by interpretation, analysis guidance, immunofluorescence examination and manuscript editing. Further, he was responsible for conceptualization and project administration including supervision and funding acquisition.

Competing Interests (COI)

JCB is receiving speaker's bureau honoraria from Amgen, Pfizer, MerckSerono, Recordati and Sanofi, is a paid consultant/advisory board member/DSMB member for Boehringer Ingelheim, InProTher, MerckSerono, Pfizer, 4SC, and Sanofi/Regeneron. His group receives research grants from Bristol-Myers Squibb, Merck Serono, HTG, IQVIA, and Alcedis. None of these

551 activities are related to the present manuscript. The other authors including K.H., C.S., L.P.,

552 F.F., P.G., J.G., N.S., I.S. declare no conflict of interest.

Funding

554 The project was funded by the German Cancer Consortium (DKTK) site budget OE 0460

555 ED003 and the federal ministry of education and research (03VP01062).

References

Johnson DE, Burtness B, Leemans CR, Lui VWY, Bauman JE, Grandis JR. Head and neck squamous cell carcinoma. *Nat Rev Dis Primers* 2020; 6: 92; https://doi.org/10.1038/s41572-020-00224-3.

Cramer JD, Burtness B, Le QT, Ferris RL. The changing therapeutic landscape of head and neck cancer. *Nat Rev Clin Oncol* 2019; 16: 669-683; https://doi.org/10.1038/s41571-019-0227-z.

Nieto MA, Huang Ruby Y-J, Jackson Rebecca A, Thiery Jean P. EMT: 2016. *Cell* 2016; 166: 21-45; https://doi.org/10.1016/j.cell.2016.06.028.

Yang J, Antin P, Berx G, Blanpain C, Brabletz T, Bronner M *et al*. Guidelines and definitions for research on epithelial–mesenchymal transition. *Nat Rev Mol Cell* 2020; 21: 341-352; https://doi.org/10.1038/s41580-020-0237-9.

McFaline-Figueroa JL, Hill AJ, Qiu X, Jackson D, Shendure J, Trapnell C. A pooled single-cell genetic screen identifies regulatory checkpoints in the continuum of the epithelial-to-mesenchymal transition. *Nat Genet* 2019; 51: 1389-1398; https://doi.org/10.1038/s41588-019-0489-5.

Cook DP, Vanderhyden BC. Context specificity of the EMT transcriptional response. *Nat Commun* 2020; 11: 2142; https://doi.org/10.1038/s41467-020-16066-2.

Vasaikar SV, Deshmukh AP, den Hollander P, Addanki S, Kuburich NA, Kudaravalli S *et al*. EMTome: a resource for pan-cancer analysis of epithelial-mesenchymal transition genes and signatures. *Br J Cancer* 2021; 124: 259-269; https://doi.org/10.1038/s41416-020-01178-9.

Puram SV, Tirosh I, Parikh AS, Patel AP, Yizhak K, Gillespie S *et al*. Single-Cell Transcriptomic Analysis of Primary and Metastatic Tumor Ecosystems in Head and Neck Cancer. *Cell* 2017; 171: 1611-1624.e1624; https://doi.org/10.1016/j.cell.2017.10.044.

Pastushenko I, Brisebarre A, Sifrim A, Fioramonti M, Revenco T, Boumahdi S *et al*. Identification of the tumour transition states occurring during EMT. *Nature* 2018; 556: 463-468; https://doi.org/10.1038/s41586-018-0040-3.

Pal A, Barrett TF, Paolini R, Parikh A, Puram SV. Partial EMT in head and neck cancer biology: a spectrum instead of a switch. *Oncogene* 2021; 40: 5049-5065; https://doi.org/10.1038/s41388-021-01868-5.

Jolly MK, Tripathi SC, Jia D, Mooney SM, Celiktas M, Hanash SM *et al*. Stability of the hybrid epithelial/mesenchymal phenotype. *Oncotarget* 2016; 7; https://doi.org/10.18632/oncotarget.8166.

12   Kinker GS, Greenwald AC, Tal R, Orlova Z, Cuoco MS, McFarland JM *et al*. Pan-cancer single-
cell RNA-seq identifies recurring programs of cellular heterogeneity. *Nat Genet* 2020; 52: 1208-
1218; https://doi.org/10.1038/s41588-020-00726-6.

13   Karacosta LG, Anchang B, Ignatiadis N, Kimmey SC, Benson JA, Shrager JB *et al*. Mapping lung
cancer epithelial-mesenchymal transition states and trajectories with single-cell resolution.
Nat Commun 2019; 10: 5587; https://doi.org/10.1038/s41467-019-13441-6.

14   Ji AL, Rubin AJ, Thrane K, Jiang S, Reynolds DL, Meyers RM *et al*. Multimodal Analysis of
Composition and Spatial Architecture in Human Squamous Cell Carcinoma. *Cell* 2020; 182: 497-
514.e422; https://doi.org/10.1016/j.cell.2020.05.039.

15   Kürten CHL, Kulkarni A, Cillo AR, Santos PM, Roble AK, Onkar S *et al*. Investigating immune and
non-immune cell interactions in head and neck tumors by single-cell RNA sequencing. *Nat
Commun* 2021; 12: 7338; https://doi.org/10.1038/s41467-021-27619-4.

16   Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The Molecular
Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* 2015; 1: 417-425;
https://doi.org/10.1016/j.cels.2015.12.004.

17   Guo D-D, Xie K-F, Luo X-J. Hypoxia-induced elevated NDRG1 mediates apoptosis through
reprograming   mitochondrial   fission   in   HCC.   *Gene*   2020;   741:   144552;
https://doi.org/10.1016/j.gene.2020.144552.

18   Ivan M, Kaelin WG. The EGLN-HIF O2-Sensing System: Multiple Inputs and Feedbacks. *Mol Cell*
2017; 66: 772-779; https://doi.org/10.1016/j.molcel.2017.06.002.

19   Laplante M, Sabatini DM. mTOR signaling at a glance. *J Cell Sci* 2009; 122: 3589-3594;
https://doi.org/10.1242/jcs.051011.

20   DeBerardinis Ralph J. Serine Metabolism: Some Tumors Take the Road Less Traveled. *Cell
Metab* 2011; 14: 285-286; https://doi.org/10.1016/j.cmet.2011.08.004.

21   Martín-Villar E, Borda-d'Agua B, Carrasco-Ramirez P, Renart J, Parsons M, Quintanilla M *et al*.
Podoplanin mediates ECM degradation by squamous carcinoma cells through control of
invadopodia stability. *Oncogene* 2015; 34: 4531-4544; https://doi.org/10.1038/onc.2014.388.

22   de Nooij-van Dalen AG, van Dongen GAMS, Smeets SJ, Nieuwenhuis EJC, Stigter-van Walsum
M, Snow GB *et al*. Characterization of the human Ly-6 antigens, the newly annotated member
Ly-6K included, as molecular markers for head-and-neck squamous cell carcinoma. *Int J Cancer*
2003; 103: 768-774; https://doi.org/10.1002/ijc.10903.

23   Reiter RE, Gu Z, Watabe T, Thomas G, Szigeti K, Davis E *et al*. Prostate stem cell antigen: A cell
surface marker overexpressed in prostate cancer. *Proc Natl Acad Sci USA* 1998; 95: 1735-1740;
https://doi.org/10.1073/pnas.95.4.1735.
644

24 Zhou K, Arslanturk S, Craig DB, Heath E, Draghici S. Discovery of primary prostate cancer
biomarkers using cross cancer learning. *Sci Rep* 2021; 11: 10433;
https://doi.org/10.1038/s41598-021-89789-x.

25 Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K *et al*. Fast, sensitive and accurate
integration of single-cell data with Harmony. *Nat Methods* 2019; 16: 1289-1296;
https://doi.org/10.1038/s41592-019-0619-0.

26 Tyler M, Tirosh I. Decoupling epithelial-mesenchymal transitions from stromal profiles by
integrative expression analysis. *Nat Commun* 2021; 12: 2592; https://doi.org/10.1038/s41467-
021-22800-1.

27 Oh IY, de Guzman Strong C. The Molecular Revolution in Cutaneous Biology: EDC and Locus
Control. *J Invest Dermatol* 2017; 137: e101-e104; https://doi.org/10.1016/j.jid.2016.03.046.

28 Stemmler MP, Eccles RL, Brabletz S, Brabletz T. Non-redundant functions of EMT transcription
factors. *Nat Cell Biol* 2019; 21: 102-112; https://doi.org/10.1038/s41556-018-0196-y.

29 Garcia-Alonso L, Holland CH, Ibrahim MM, Turei D, Saez-Rodriguez J. Benchmark and
integration of resources for the estimation of human transcription factor activities. *Genome
Res* 2019; 29: 1363-1375; https://doi.org/10.1101/gr.240663.118.

30 Holland CH, Tanevski J, Perales-Patón J, Gleixner J, Kumar MP, Mereu E *et al*. Robustness and
applicability of transcription factor and pathway analysis tools on single-cell RNA-seq data.
Genome Biol 2020; 21: 36; https://doi.org/10.1186/s13059-020-1949-z.

31 Alvarez MJ, Shen Y, Giorgi FM, Lachmann A, Ding BB, Ye BH *et al*. Functional characterization
of somatic mutations in cancer using network-based inference of protein activity. *Nat Genet*
2016; 48: 838-847; https://doi.org/10.1038/ng.3593.

32 Sánchez-Tilló E, Lázaro A, Torrent R, Cuatrecasas M, Vaquero EC, Castells A *et al*. ZEB1
represses E-cadherin and induces an EMT by recruiting the SWI/SNF chromatin-remodeling
protein BRG1. *Oncogene* 2010; 29: 3490-3500; https://doi.org/10.1038/onc.2010.102.

33 Williams ED, Gao D, Redfern A, Thompson EW. Controversies around epithelial–mesenchymal
plasticity in cancer metastasis. *Nat Rev Cancer* 2019; 19: 716-732;
https://doi.org/10.1038/s41568-019-0213-x.

34 Derynck R, Weinberg RA. EMT and Cancer: More Than Meets the Eye. *Dev Cell* 2019; 49: 313-
316; https://doi.org/10.1016/j.devcel.2019.04.026.

35 Dongre A, Weinberg RA. New insights into the mechanisms of epithelial–mesenchymal
transition and implications for cancer. *Nat Rev Mol Cell* 2019; 20: 69-84;
https://doi.org/10.1038/s41580-018-0080-4.

36 Tsai JH, Donaher JL, Murphy DA, Chau S, Yang J. Spatiotemporal regulation of epithelial-
mesenchymal transition is essential for squamous cell carcinoma metastasis. *Cancer Cell* 2012;
22: 725-736; https://doi.org/10.1016/j.ccr.2012.09.022.

37 Xiao Z, Dai Z, Locasale JW. Metabolic landscape of the tumor microenvironment at single cell
resolution. *Nat Commun* 2019; 10: 3763; https://doi.org/10.1038/s41467-019-11738-0.

38 Krebs AM, Mitschke J, Lasierra Losada M, Schmalhofer O, Boerries M, Busch H *et al*. The EMT-
activator Zeb1 is a key factor for cell plasticity and promotes metastasis in pancreatic cancer.
Nat Cell Biol 2017; 19: 518-529; https://doi.org/10.1038/ncb3513.

39 Carstens JL, Yang S, Correa de Sampaio P, Zheng X, Barua S, McAndrews KM *et al*. Stabilized
epithelial phenotype of cancer cells in primary tumors leads to increased colonization of liver
metastasis in pancreatic cancer. *Cell Rep* 2021; 35: 108990;
https://doi.org/10.1016/j.celrep.2021.108990.

40 Burk U, Schubert J, Wellner U, Schmalhofer O, Vincan E, Spaderna S *et al*. A reciprocal
repression between ZEB1 and members of the miR-200 family promotes EMT and invasion in
cancer cells. *EMBO reports* 2008; 9: 582-589; https://doi.org/10.1038/embor.2008.74.

41 Li S, Gallup M, Chen YT, McNamara NA. Molecular mechanism of proinflammatory cytokine-
mediated squamous metaplasia in human corneal epithelial cells. *Invest Ophthalmol Vis Sci*
2010; 51: 2466-2475; https://doi.org/10.1167/iovs.09-4677.

42 Zhang P, Wei Y, Wang L, Debeb BG, Yuan Y, Zhang J *et al*. ATM-mediated stabilization of ZEB1
promotes DNA damage response and radioresistance through CHK1. *Nat Cell Biol* 2014; 16:
864-875; https://doi.org/10.1038/ncb3013.

43 Oshima RG. Apoptosis and keratin intermediate filaments. *Cell Death Differ* 2002; 9: 486-492;
https://doi.org/10.1038/sj.cdd.4400988.

44 Fennell KA, Vassiliadis D, Lam EYN, Martelotto LG, Balic JJ, Hollizeck S *et al*. Non-genetic
determinants of malignant clonal fitness at single-cell resolution. *Nature* 2022; 601: 125-131;
https://doi.org/10.1038/s41586-021-04206-7.

45 Spassova I, Ugurel S, Terheyden P, Sucker A, Hassel JC, Ritter C *et al*. Predominance of Central
Memory T Cells with High T-Cell Receptor Repertoire Diversity is Associated with Response to
PD-1/PD-L1 Inhibition in Merkel Cell Carcinoma. *Clinical Cancer Research* 2020; 26: 2257-2267;
https://doi.org/10.1158/1078-0432.Ccr-19-2244.

46 Newman AM, Steen CB, Liu CL, Gentles AJ, Chaudhuri AA, Scherer F *et al*. Determining cell type
abundance and expression from bulk tissues with digital cytometry. *Nat Biotechnol* 2019; 37:
773-782; https://doi.org/10.1038/s41587-019-0114-2.

47 Luecken MD, Theis FJ. Current best practices in single-cell RNA-seq analysis: a tutorial. *Mol Syst*
Biol 2019; 15: e8746; https://doi.org/10.15252/msb.20188746.

48 Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A *et al*. Integrated analysis of
multimodal single-cell data. *Cell* 2021; 184: 3573-3587.e3529;
https://doi.org/10.1016/j.cell.2021.04.048.

49 Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq data
using regularized negative binomial regression. *Genome Biol* 2019; 20: 296;
https://doi.org/10.1186/s13059-019-1874-1.

50 Tirosh I, Izar B, Prakadan SM, Wadsworth MH, Treacy D, Trombetta JJ *et al*. Dissecting the
multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science* 2016; 352:
189; https://doi.org/10.1126/science.aad0501.

51 Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A. Fast gene set
enrichment analysis. *bioRxiv* 2021: 060012; https://doi.org/10.1101/060012.

52 Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA *et al*. Gene set
enrichment analysis: A knowledge-based approach for interpreting genome-wide expression
profiles. *Proc Natl Acad Sci USA* 2005; 102: 15545; https://doi.org/10.1073/pnas.0506580102.

53 Hänzelmann S, Castelo R, Guinney J. GSVA: gene set variation analysis for microarray and RNA-
Seq data. *BMC Bioinform* 2013; 14: 7; https://doi.org/10.1186/1471-2105-14-7.

54 Street K, Risso D, Fletcher RB, Das D, Ngai J, Yosef N *et al*. Slingshot: cell lineage and pseudotime
inference for single-cell transcriptomics. *BMC Genom* 2018; 19: 477;
https://doi.org/10.1186/s12864-018-4772-0.

55 La Manno G, Soldatov R, Zeisel A, Braun E, Hochgerner H, Petukhov V *et al*. RNA velocity of
single cells. *Nature* 2018; 560: 494-498; https://doi.org/10.1038/s41586-018-0414-6.

56 Wickham H. ggplot2: Elegant Graphics for Data Analysis. *Springer-Verlag New York* 2016;

57 Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* 2016; 32: 2847-2849;
https://doi.org/10.1093/bioinformatics/btw313.

Figure 1 | Single-cell gene expression signatures in OSCC cells from a single metastasis reveal several predominant, but not necessarily exclusive, functional phenotypes. (A) UMAP based on scRNAseq data of 4,076 cells isolated from a metachronous lymph node metastasis. Cells are annotated and summarized according to the presumed cell type. **(B)** UMAP of 1906 OSCC cells depicted in A. Cells are annotated according to predominant functional phenotype. **(C)** Heatmap for scaled, log-normalized gene expression of tumor cells (columns) split by respective phenotype and the top 10 differentially expressed genes (DEGs) (rows) of the respective phenotype against all other tumor cells. DEGs are sorted from highest to lowest log2 foldchange. Row sections are ordered like column sections. **(D)** Top 5 enriched gene sets from log2 foldchanges of respective tumor phenotypes by normalized enrichment scores (x axis). Gene sets of respective phenotypes are sorted from highest to lowest enrichment. Bars are colored by the negative decadic logarithm of the Benjamini-Hochberg adjusted p-value (padj). DCs: dendritic cells. ECs: endothelial cells.

Figure 2 | Progressive epithelial differentiation, but no strong uniform direction of development in pEMT clusters. (A) UMAP of 1906 OSCC cells annotated based on SNN clustering, defining 4 pEMT (pEMT-1 to 4), 4 epithelial differentiated (epi-1 to 4) and one mixed (mix) cluster; clusters are numbered by size. **(B)** Heatmap for scaled, log-normalized gene expression in EMP-associated tumor cell phenotypes (columns) split by EMP cluster and their top 5 DEGs (rows) against all other EMP-related tumor cell phenotypes. DEGs are sorted from highest to lowest log2 foldchange. Row sections are ordered like column sections. **(C)** Projection of RNA velocity on the UMAP depicted in A. Arrows indicate the extrapolated direction of development and their length the strength of this future development. **(D)** First two principal components of OSCC cells with the three EMP-related principal curves that are derived from trajectory inference. Graph on top visualizes the relationship between EMP clusters described by the three principal curves forming a branching trajectory. **(E)** Log-normalized expression (y-axis) of MMP1, VIM, SPRR1B and KLK7 across pseudotime values (x-axis) of curve 2, color-coded by clusters. Red lines indicate smoothed expression values over the trajectory generated with a general additive model; 95 % confidence intervals are shaded gray. **(F)** Inferred CNVs across EMP-related tumor cells (rows) for all chromosomes (columns). Red indicates copy number gains, white diploid copy number and blue copy number loss. Columns show genes categorized in chromosomes and ordered by genome position; hence the size of the chromosome reflects the number of detected genes and not its nucleotide length. Mitochondrial genes were excluded.

Figure 3 | Intra-tumoral heterogeneity of OSCC is driven by EMP. (A) UMAP based on scRNAseq data of 7,263 cancer cells from 16 different patients annotated by patient. **(B)** Heatmap for scaled, log-normalized gene expression of tumor cells split by patients and their top 5 DEGs (rows) against all other tumor cells. All patients with less than 100 cells are summarized in the 'other' column. DEGs are sorted from highest to lowest log2 foldchange. Row sections are ordered like column sections. **(C)** UMAP based on scRNAseq data depicted in A with PCs corrected for patient-specific effects using harmony. Cells are annotated according to their predominant phenotype. **(D)** Relative distribution of tumor cell phenotypes (left) and cancer cell abundance (right) across patients. The label on the y-axis shows the sample identification and tumor localization (primary tumor [PT] or lymph node metastasis [MET]). **(E)** UMAP based on scRNAseq data of 2,948 OSCC cells from patient HN01. Cells were annotated based on SNN clustering and the predominant phenotype. **(F)** Triangle heatmap of cosine similarity comparing the intratumoral heterogeneity across all patients. Cosine similarity is calculated between log2 fold changes from patient-specific clusters against all other tumor cells within the respective patient. Left side annotated are patient-specific clusters from patient 1 depicted in Figure 2A and right side from patient HN01 depicted in E. We included only patients with more than 50 tumor cells.

Figure 4 | ZEB1 is highly active in metastatic epithelial differentiated OSCC cells. (A) Percentage of tumor cells with detectable mRNA expression from scRNAseq (more than one UMI) encoding the indicted EMP-related transcription factors. **(B)** Mean inferred activity based on the target genes of the indicated transcription factors across tumor phenotypes from EMP-related patient-specific clusters. On top the log-normalized expression of *CDH1* and cornifin-B (*SPRR1B*) is shown, on bottom the localization (primary tumor [PT] or lymph node metastasis [MET]) and respective EMP phenotype of the cluster. **(C)** Mean activity of ZEB1 for epithelial differentiated and pEMT clusters of each patient, respectively. Connecting lines show dots belonging to the same patient. **(D, E)** ZEB1 (D) and cornifin-B (E) protein expression detected in serial sections by IHC of the primary tumor from patient 2; comparable areas are depicted. Scale bars equal 200 μm in overview and 100 μm in zoomed image. **(F)** Colocalized expression of ZEB1 (green) and cornifin-B (red) detected by double staining in the lymph node metastasis of patient 1 nuclei are stained in blue (DAPI), Scale bars equal 10 μm.