1   Recurrent circuits amplify corticofugal signals and drive feed-forward inhibition in the
2   inferior colliculus
3

Hannah M. Oberle[1,2#], Alexander N. Ford[2#], Pierre F. Apostolides[2,3*]
1) Neuroscience Graduate Program
2) Kresge Hearing Research Institute, Department of Otolaryngology – Head & Neck
Surgery
3) Molecular and Integrative Physiology, University of Michigan Medical School
#Equal Contribution
*corresponding author: piaposto@med.umich.edu

Abstract
The inferior colliculus (IC) is a midbrain hub critical for perceiving complex, time-varying
sounds such as speech. In addition to processing ascending inputs from most auditory
brainstem nuclei, the IC receives descending inputs from auditory cortex that control IC
neuron feature selectivity, plasticity, and certain forms of perceptual learning. Although
these corticofugal synapses primarily release the excitatory transmitter glutamate, many
physiology studies show that auditory cortical activity has a net inhibitory effect on IC
neuron spiking. Perplexingly, anatomy studies indicate that corticofugal axons primarily
target glutamatergic IC neurons while only sparsely innervating IC GABA neurons,
thereby suggesting that corticofugal inhibition of the IC occurs largely independently of
feedforward activation of local GABA neurons. We shed light on this paradox using *in*
vitro electrophysiology and optogenetics in acute IC slices from fluorescent reporter
mice. We find that corticofugal synaptic strength is indeed stronger onto IC glutamate
compared to GABA neurons. Nevertheless, two mechanisms enable reliable
corticofugal activation of local GABA neurons. 1) Many IC GABA neurons fire tonically
at rest, such that sparse and weak excitation suffices to significantly increase their spike
rates. 2) Corticofugal activity triggers barrages of large amplitude, polysynaptic EPSPs
in IC GABA neurons, owing to a dense intra-collicular connectivity from local axon
collaterals of IC glutamate neurons. Consequently, repetitive activity in corticofugal
fibers drives spikes in IC GABA neurons and generates substantial recurrent inhibition.
Thus, descending signals trigger intra-collicular inhibition despite apparent constraints
of monosynaptic connectivity between auditory cortex and IC GABA neurons.

Introduction
Feedback projections from the sensory neo-cortex to sub-cortical regions are ubiquitous
in the mammalian brain. These corticofugal pathways enable "high-level", cortical
computations to rapidly control the nature of ascending sensory signals, thereby
hypothetically supporting important "top-down" functions such as predictive coding, error
propagation, or stream segregation (Briggs and Usrey, 2011; Stebbings et al., 2014;

Usrey and Sherman, 2019; Asilador and Llano, 2020). Interestingly, corticofugal activity
often has a net inhibitory effect upon spontaneous and sensory-evoked activity in sub-
cortical circuits, consequently sharpening receptive fields and increasing the
sparseness of neuronal representations along the ascending sensory hierarchy (Syka
and Popelár, 1984; Zhang et al., 1997; Bledsoe et al., 2003; Boyd et al., 2012; Crandall
et al., 2015; Vila et al., 2019; Born et al., 2021; but see Kirchgessner et al., 2021).
Corticofugal projections originate almost exclusively from excitatory (glutamatergic)
neurons in cortical layers 5 and 6 (Hattox and Nelson, 2007; Schofield, 2009; Slater et
al., 2013; Asilador and Llano, 2020; Sherman and Usrey, 2021), the well-documented
inhibitory effects suggest that corticofugal activity primarily recruits sub-cortical,
feedforward inhibitory (GABAergic) interneurons rather than faithfully detonating spikes
in projection neurons. However, the cellular and circuit-level mechanisms that enable
corticofugal activity to reliably generate sub-cortical inhibition are poorly understood; this
knowledge gap persists owing to the difficulty of quantifying corticofugal transmission in
identified excitatory and inhibitory neurons.
56
In the central auditory system, the corticofugal projection from auditory cortex to the
inferior colliculus (IC) is of particular importance as the IC relays the majority of
ascending acoustic signals destined for forebrain circuits. Accordingly, corticofugal
activity powerfully shapes how IC neurons respond to diverse sound features, often via
inhibitory interactions: Stimulating the auditory cortex dampens or completely
suppresses IC acoustic responses (Syka and Popelár, 1984; Bledsoe et al., 2003; Vila
et al., 2019; Blackwell et al., 2020), a result further corroborated by intracellular data
showing that auditory cortex stimulation can drive synaptic inhibition in individual IC
neurons (Mitani et al., 1983; Qi et al., 2020). Conversely, silencing auditory cortex often
potentiates spontaneous and sound-evoked activity in the IC (Nwabueze-Ogbo et al.,
2002; Popelár et al., 2003; Popelář et al., 2016; but see Zhang and Suga, 1997),
indicating that ongoing auditory cortical activity can have a net inhibitory effect on the
moment-to-moment excitability of IC neurons. However, anatomical data show that IC
GABA neurons receive surprisingly few synapses from auditory cortex; the majority of
corticofugal axons instead targeting glutamatergic IC neurons (Nakamoto et al., 2013;
Chen et al., 2018). Thus, rather than recruiting *local* inhibitory neurons to sharpen IC
neuron tuning curves, auditory cortex may instead exert inhibitory control through long-
range inhibitory pathways (Beneyto et al., 1998; Budinger et al., 2000), via local effects
upstream of the IC (Kong et al., 2014), or through a newly discovered, monosynaptic
GABAergic corticofugal projection (Bertero et al., 2021). Alternatively, circuit
mechanisms beyond the orthodox feedforward inhibitory motif, whereby long-range
excitation preferentially drives GABAergic interneurons (Pouille and Scanziani, 2001;
Gabernet et al., 2005; Cruikshank et al., 2007; Boyd et al., 2012), may nevertheless
enable the auditory cortex to generate local inhibition in the IC.

We used patch-clamp electrophysiology and optogenetics in brain slices from
fluorescent GABA reporter mice to study corticofugal transmission onto identified IC
neurons. We focused specifically on neurons in the dorso-medial "shell" IC because this
sub-region receives the densest projection of corticofugal axons (Winer et al., 1998;
Song et al., 2018; Oberle et al., 2022). Our data confirm previous anatomical predictions
by showing that corticofugal transmission is on average stronger onto shell IC glutamate
compared to GABA neurons. However, we also find that shell IC GABA neurons are
densely contacted by potent, intra-collicular synapses from local glutamate neurons. As
such, repetitive corticofugal activity drives strong polysynaptic excitation in a subset of
shell IC GABA neurons and generates local inhibition. Our results identify a novel
mechanism for the neo-cortex to drive recurrent computations in a sub-cortical circuit.
More broadly, the data establish a biophysical context via which descending inhibitory
control operates in apparent absence of a dense, monosynaptic convergence onto
inhibitory networks.
95
96 **Results**
97
Cortical stimulation inhibits superficial IC neurons in vivo
Previous studies show that stimulating the auditory cortex profoundly reduces sound-
evoked spiking in IC neurons. As a first test to determine if we could similarly observe
descending inhibition of IC neurons *in vivo*, we electrically stimulated the auditory cortex
of urethane anesthetized mice during whole-cell patch-clamp recordings from neurons
in the cortico-recipient, superficial IC layers. Indeed, single shocks (Figure 1B) or train
stimulation (Figure 1C; 10 shocks delivered between 10-200 Hz) drove brief excitatory
postsynaptic potentials (EPSPs) followed by large inhibitory postsynaptic potentials
(IPSPs) in n = 4/7 cells tested. Similar results were obtained with optogenetic
stimulation of auditory cortex neurons expressing the light gated opsin Chronos: Single
flashes of 473 nm light delivered to auditory cortex neurons drove EPSP/IPSP
sequences in n = 6/21 cells tested (Figure 1E). Interestingly, inhibition was also seen in
5/17 IC neurons that did not receive any apparent excitation from auditory cortex. In n =
4 neurons, repetitive auditory cortical activity (80 stimuli at 20 Hz) caused a tonic or
phasic hyperpolarization of the membrane potential (Figure 1F), whereas in 1 neuron
tested, single light flashes generated a single, long-latency IPSP. Altogether these data
indicate that in addition to the monosynaptic depolarizations mediated by glutamatergic
corticofugal projections (Oberle et al., 2022), descending activity can also substantially
hyperpolarize IC neurons, presumably via polysynaptic inhibition.

Figure 1: *Auditory cortical stimulation generates IPSPs in IC neurons.* **A)** Cartoon of experiment. Whole cell recordings are obtained from superficial IC neurons in urethane anesthetized mice; A bipolar electrode is placed in auditory cortex to activate corticofugal fibers. **B)** Example average trace showing a brief EPSP followed by a large and long-lasting IPSP upon auditory cortex stimulation. **C)** Example average trace showing that synaptic inhibition summates upon repetitive stimulation of auditory cortex. Data in B and C are from the same neuron. Stimulation artifacts are blanked for clarity. **D)** Chronos is expressed in auditory cortex neurons via AAV injections and 2-4 weeks later, whole-cell patch-clamp recordings are obtained from superficial IC neurons. **E)** Example average traces showing an EPSP-IPSP sequence following single light flashes delivered to the auditory cortex. **F)** In a different experiment as that shown in E, repetitive stimulation of auditory cortex generates no apparent EPSPs, but rather causes a sustained hyperpolarization of the neuron's membrane potential.

Corticofugal synapses are stronger onto glutamate compared to GABA shell IC neurons
The hyperpolarization of IC neurons observed during auditory cortex stimulation is
intriguing, as anatomy data suggest that auditory cortico-collicular axons preferentially
synapse onto excitatory glutamatergic IC neurons; far fewer descending synapses
target GABA neurons in the IC (Nakamoto et al., 2013; Chen et al., 2018). We thus
wondered if the functional strength of auditory corticofugal synapses differed between
glutamatergic and GABAergic IC neurons in a manner consistent with anatomical, or
rather, physiological data (e.g., Figure 1). To this end, we expressed the red fluorescent
protein tdTomato specifically in GABAergic neurons by crossing crossed VGAT-ires-cre
and Ai14 fl/fl mouse lines. We then virally transduced the optogenetic activator Chronos
in auditory cortex neurons of these mice via stereotaxic injections, and 2-4 weeks later,
prepared acute brain slices for whole-cell patch-clamp electrophysiology (Figure 2A,B).

We targeted either tdTomato-positive or -negative somata in the dorso-medial shell IC
to record from GABAergic or presumptive glutamatergic neurons, respectively
(Supplemental Figure 1A,B). Single flashes of blue light delivered through the
microscope objective activated Chronos-expressing auditory cortico-collicular axons
and generated EPSPs that were >3-fold larger in glutamate compared to GABA neurons
(Figure 2B,C. Median amplitude: 1.86 vs. 0.58 mV, n=29 glutamate neurons from N = 9
mice and n=37 GABA neurons from N=10 mice, respectively. p=0.0013, Kruskal-Wallis
test). By contrast, median EPSP halfwidths were similar across the two neuron
populations (37.5 vs. 41.0 ms, p=0.7223, Kruskal-Wallis test), indicating that amplitude
differences were unlikely due to expression of postsynaptic glutamate receptors with
differential glutamate affinity. Furthermore, these results were unlikely to reflect a
preferential severing of GABA neuron dendrites during slice preparation, as

Figure 2: *Differential strength of corticofugal synapses onto shell IC GABA and glutamate neurons.* **A)** Diagram of experiment. The optogenetic activator Chronos is virally transduced into auditory cortex neurons of transgenic VGAT-cre x Ai14 mice. **B)** 2-4 weeks following surgery, corticofugal synaptic strength is assayed in presumptive GABA and glutamate neurons by recording optogenetically evoked EPSPs from visually targeted tdTomato-positive and negative neurons. Right, example average EPSPs evoked by single light flashes in a glutamate (black) and GABA (magenta) shell IC neuron. **C,D)** Summary data showing cumulative probability distributions for EPSP peak amplitudes (C) and half-widths (D) in glutamate and GABA neurons. Asterisks denote statistical significance.

instantaneous PSP rates were significantly higher in GABA compared to glutamate
neurons (Figure 3A-C; median rate 15.9 vs. 7.4 Hz, n = 43 GABA neurons from N = 10
mice and n = 26 glutamate neurons from N = 9 mice, respectively. p < 0.001, Kruskal-
Wallis test). Instead, our results imply a specific connectivity logic to the auditory
cortico-collicular pathway whereby glutamate, rather than GABA neurons are the major
recipients of monosynaptic corticofugal signals. By contrast, the differential background
PSP activity suggests that GABA neurons may receive a substantial number of
excitatory synapses from the intra-collicular circuit.

Figure 3: *IC glutamate and GABA neurons have different rates of spontaneous PSP activity.* **A,B)** Examples of spontaneous synaptic activity in single glutamate (A) or GABA (B) shell IC neurons. Scale bars apply to both panels. Of note is the higher rate and larger amplitude of spontaneous PSPs in the GABA neuron. **C)** Summary of instantaneous PSP rates for the two neuron classes.

A subset of IC GABA neurons tonically fire action potentials
As expected from previous studies (Ono et al., 2005; Naumov et al., 2019), IC GABA
and glutamate neurons displayed a range of overlapping cellular properties and could
not be obviously distinguished based solely on membrane responses to current
injections (Supplemental Figure 1C-H). However, we observed that GABA neurons
were significantly more likely to tonically fire spikes at rest (Figure 4A,B. 34/68 GABA
neurons fired tonically upon break-in compared to 1/25 glutamate neurons, χ^2=17.14,
p<0.001). Consequently, even sparse synaptic excitation might suffice to significantly
increase the mean firing rate of IC GABA neurons. Indeed, injecting small excitatory
postsynaptic current (EPSC)-like waveforms (10-20 pA peak amplitude) at the soma of
tonically firing GABA neurons significantly increased spike rates 1.82 ± 0.16 fold relative
to a pre-current injection baseline period Figure 4C,D; n = 8 cells from N = 2 mice,
p=0.0012, one-sample t-test). Thus, a dense convergence of descending axons is not
explicitly necessary to increase firing rates of shell IC GABA neurons.

Figure 4: *A subset of GABA neurons fire tonically, thus requiring only small currents to increase baseline firing rates* **A)** Example trace from a tonic firing IC GABA neuron recorded in a VGAT-cre x Ai14 mouse. **B)** Summary histogram of tonic firing rates in IC GABA neurons. Green bin denotes neurons that did not fire tonically. **C)** Top: Example overlaid traces from a tonically firing IC GABA neuron. Middle: An EPSC -like current waveform (20 pA peak amplitude) increases spike probability in the ~50 ms following current injection, as exemplified in the spike (lower panel). **D)** Summary data from n=8 cells as in panel C, showing that small EPSC waveforms (10-20 pA) nearly double the spike probability of tonically firing GABA neurons.

Shell IC GABA neurons receive powerful intra-collicular excitation
The high rate of spontaneous PSPs in IC GABA neurons (Figure 3) might reflect a
dense convergence of local, intra-collicular excitatory fibers. We tested this hypothesis
by crossing G42 mice (Chattopadhyaya et al., 2004), which express GFP in a subset of
IC GABA neurons (Supplemental Figure 2), with VGluT2-cre mice which express cre-
recombinase in nearly all IC glutamate neurons (Chen et al., 2018). We then injected
into the IC a cre-dependent virus packaging the red-shifted optogenetic activator
ChrimsonR (Figure 5A). This approach enabled targeting our patch-clamp recordings to
shell IC GABA neurons while selectively activating local glutamate neurons via an optic
fiber coupled to a 625 nm LED (Figure 5B). We estimated the strength of unitary
connections using a "minimal stimulation" paradigm designed to activate one or very
few local inputs: The LED power level was titrated such that optogenetic stimulation
fluctuated between synaptic successes and failures on a trial-by-trial basis (Figure 5C).

Under these conditions, successful EPSPs had a mean amplitude of 2.8 ± 0.3 mV
(Figure 5D, n = 11 cells from N = 6 mice). In addition, EPSP size grew monotonically
with increasing LED power levels: EPSP amplitudes at maximal LED power had a
median value 7.2 fold larger than EPSPs evoked with threshold stimulation (range: 2.2
to 37.7; Figure 5E,F. n=12 cells from N = 6 mice). These measurements likely under-
estimate the extent of synaptic connectivity due to the severing of connections during
the slicing procedure. Consequently, our values probably represent a lower bound for
the extent of intra-collicular convergence onto shell IC GABA neurons.
184
185   *Repetitive corticofugal activity drives polysynaptic excitation onto IC GABA neurons*
186   The dense convergence of intra-collicular excitation suggested by Figures 3 + 5 implies
187   that corticofugal activity might drive spiking in IC glutamate neurons, subsequently
188   causing polysynaptic EPSPs in IC GABA neurons. In this context, local recurrent

Figure 5: *IC GABA neurons pool strong unitary excitatory inputs from multiple local glutamate neurons.*
A,B) Cartoon of experiment. A flex-ChrimsonR virus is injected in the IC of G42 x VGluT2-cre mice. 2-4
weeks later, patch-clamp recordings are targeted to GFP+ neurons in the shell IC and glutamate neurons
are optically stimulated with 625 nm light via an optic fiber positioned above the slice (not shown). **C)**
Example recording from a shell IC GABA neuron showing transmission successes (black) and failures
(grey) following optical activation of ChrimsonR-expressing glutamate neurons. **D)** Summary data plotting
the average peak amplitude of putative unitary EPSPs during threshold stimulation. Black is individual
experiments, magenta is mean ± SEM. **E,F)** Increasing LED power beyond threshold generates larger
EPSPs, indicating that multiple presynaptic glutamate neurons converge onto a single postsynaptic
GABA neuron. Data are from a different experiment as the one in C.

excitation would amplify auditory corticofugal signals onto IC GABA neurons
independently of strong monosynaptic innervation. We tested this hypothesis by
repetitively stimulating Chronos-expressing corticofugal axons (25 flashes at 50 Hz)
while recording from GABA or glutamate neurons in VGAT-cre x Ai14 mice; these
stimulation parameters are within the physiological range of sound-evoked firing rates in
corticofugal neurons (Williamson and Polley, 2019).
195
In a subset of GABA neurons (n = 21 neurons from N = 16 mice), train stimuli triggered
asynchronous barrages of large amplitude EPSPs that could be evoked even when
single light flashes resulted in either weak or inapparent EPSPs (Figure 6A,B).
Strikingly, these EPSP barrages were sometimes powerful enough to drive spikes even
when the membrane potential was hyperpolarized with bias current to prevent tonic
firing (Supplemental Figure 3). Close inspection of the raw traces indicated that these
asynchronous EPSPs are generated by polysynaptic activity rather than monosynaptic
inputs from auditory cortex: EPSPs occurred with a variable delay and displayed
considerable temporal jitter in their onset following individual light flashes (Figure 6C).
Additional examples of polysynaptic EPSPs in IC GABA neurons are provided in
Supplemental Figure 4. Interestingly, EPSP barrages with long and variable latencies
were less common in glutamate neurons (n = 2 neurons from N = 2 mice). Instead,
single and train stimuli almost exclusively generated short latency EPSPs that were
temporally locked to individual light pulses (Figure 6D-F; n = 14 neurons from N = 8
mice), as expected from dense monosynaptic connectivity of corticofugal axons onto IC
glutamate neurons.
212
The difference in EPSP timing between GABA and glutamate neurons is further
apparent when the EPSP latencies after each pulse in the stimulus train are plotted as
latency histograms for the two example cells in panels A-F: Whereas EPSPs in the
glutamate neuron reliably occur ~3-4 ms following each pulse (Figure 6G), EPSP onsets
in the GABA neuron are temporally dispersed over 10 ms following light stimuli (Figure
6H). Consequently, double gaussian fits to the data reveal two clear peaks at ~3 and ~7
ms in the histogram from the GABA neuron, but not in the data from the example
glutamate neuron (see also Supplemental Figure 4C,F). A qualitatively similar result can
be observed when averaging the membrane potential waveform of the example
glutamate and GABA neurons following each light pulse in the 50 Hz train (Figure 6I):
The cycle averages from the glutamate and GABA neurons show one and two peaks,
respectively. We interpret the first short-latency peak as reflecting the monosynaptic
input from corticofugal synapses, while the later, more temporally-jittered EPSPs reflect
recurrent excitation from local IC glutamate neurons driven to spike by monosynaptic
corticofugal inputs. Altogether, these data show that repetitive corticofugal activity
triggers powerful recurrent excitation in shell IC GABA neurons.

Figure 6: *Repetitive corticofugal activity triggers polysynaptic excitation in the shell IC.*
A,B) Example EPSPs in a GABA neuron during single (A) or train (B) stimuli delivered to corticofugal axons. While single light flashes generate negligible synaptic depolarization in this example, train stimulation drives powerful EPSP barrages. Gray traces are individual trials, magenta is average of multiple trials. **C)** The second and third stimuli from panel B are shown at faster timebase. Blue dotted lines denote light flashes. Of note is the long latency and jittered onset of the large EPSPs, a hallmark of polysynaptic origin. **D,E)** Same experimental approach as in A-B, but for a glutamate neuron. Gray traces are individual trials, black is average. Scale bars in D and E also apply to panels A and B, respectively. **F)** The second and third stimuli in the train from panel E are shown at a faster time base. Of note is the short latency and low onset jitter. **G,H)** Histograms of the EPSP onset latency after each of the 25 stimuli in the trains for the GABA and glutamate neurons shown in A-C and D-F, respectively. **I)** The membrane voltage following each stimuli in the 50 Hz trains is averaged and peak normalized. Black and magenta traces are data from the example glutamate and GABA neuron, respectively. Of note is the dual component EPSP in the GABA neuron, suggesting a monosynaptic corticofugal input followed by a much larger recurrent EPSP from the local circuit.

Corticofugal activity preferentially triggers recurrent excitation in GABA neurons.
Our data thus far suggest that corticofugal transmission drives monosynaptic excitation
of IC glutamate neurons, which in turn generate polysynaptic EPSPs in IC GABA
neurons. This hypothesis predicts that the primarily polysynaptic EPSPs in IC GABA
neurons should be uniquely sensitive to experimental manipulations that dampen the
feedforward spiking of IC glutamate neurons during corticofugal stimulation. By contrast,
the mostly monosynaptic corticofugal EPSPs onto IC glutamate neurons should be
comparatively less reduced by the same manipulation. We tested this prediction by

quantifying how a sub-saturating concentration of the AMPA receptor antagonist NBQX
(50 nM) differentially impacts repetitive corticofugal activity (25 stimuli at 50 Hz) in IC
glutamate and GABA neurons. This concentration of NBQX blocks 20-50% of the AMPA
receptor charge transfer (Randle et al., 1992; Diamond and Jahr, 1997) and should
decrease the probability that glutamate neurons fire spikes during corticofugal
stimulation, thus non-linearly reducing polysynaptic EPSPs in GABA neurons.
Accordingly, 50 nM NBQX reduced the total voltage integral during stimulus trains by
22±7% in glutamate neurons (Figure 7A,C; n=10 cells from N = 8 mice). By contrast, 50
nM NBQX caused a significantly greater reduction in GABA neurons showing long-
latency and asynchronous EPSPs (61±3% reduction, Figure 7B,C; n = 11 cells from N =
8 mice; p<0.0001 compared to glutamate neurons, paired t-test). Altogether these data
argue that while auditory cortical activity drives spikes in GABA neurons (Supplemental
Figure 3), much of the underlying depolarization is driven by local excitatory inputs
rather than monosynaptic connections from corticofugal axons.

Figure 7: *Differential block of "corticofugal" excitation in glutamate and GABA neurons by 50 nM NBQX.* **A,B)** Example average train EPSPs from a shell IC glutamate (A) or GABA (B) neuron before and after bath application of 50 nM NBQX (black and magenta traces, respectively). Lower traces are the cumulative integral of the EPSP waveform. **C)** Summary data showing the fraction remaining in 50 nM NBQX for the for the cumulative integral of the EPSP waveform. Black and magenta symbols are individual experiments and mean ± SEM, respectively. Asterisks denote statistical significance.

Corticofugal inputs generates polysynaptic inhibition
Our data show that auditory cortical activity strongly depolarizes IC GABA neurons via a
combination of direct and polysynaptic excitation. If IC GABA neurons project their
axons locally, stimulating auditory cortico-collicular axons should thus suffice to
generate feed-forward inhibitory postsynaptic currents (IPSCs) in the shell IC. We
tested this hypothesis in Chronos-injected wild-type mice by voltage-clamping the
somata of shell IC neurons near the reversal potential for excitation (~ +5 to +10 mV)
and stimulating corticofugal axons with trains of blue light flashes (25x, 50 Hz).

Recordings under these conditions were characterized by a flurry of spontaneous,
outward inhibitory postsynaptic currents (IPSCs) that were preferentially sensitive to the
GABA$_A$ receptor antagonist SR95531 (5-10 µM; Supplemental Figure 5. Median IPSC
rate: 13.5 vs. 0.0557 Hz in control and SR95531, p=0.0156, sign-rank test, n = 7 cells
from N = 3 mice). Strikingly, stimulating auditory cortico-collicular axons caused a ~2-3
fold increase in IPSC rate relative to baseline (Figure 8A,B), as revealed by a main
effect of time in a two-way, repeated measures ANOVA (F(14,126)=11.7, p<0.0001, n =
10 cells from N = 5 mice). Importantly, the IPSC rate increase was abolished by bath
application of glutamate receptor antagonists NBQX (10 µM) and/or R-CPP (5 µM),
indicating that it was mediated by polysynaptic activation of IC GABA neurons
(Timepoint x drug interaction: F(14,126)=13.38, p<0.0001).
271
Functionally, this feedforward inhibition controlled the temporal integration of
corticofugal excitation: Blocking GABA$_A$ receptors significantly increased corticofugal
EPSP summation during repetitive stimulation in current clamp, measured as a
significant increase in the voltage integral following bath application of 5-10 µM
SR55931 (Figure 8C,D. Fractional change relative to control: 1.18 ± 0.06, n = 12 cells
from N = 11 mice. p=0.013, one sample t-test). Interestingly, we also noted a minor, but
statistically significant increase in membrane resistance following SR95531 (386 ± 61
vs. 397 ± 62 MΩ in control and SR95531, respectively. Fractional change: 1.04 ± 0.014,
p = 0.0204, one sample t-test). Although this observation argues that the high rates of
spontaneous IPSCs from the local circuit contribute a certain level of effectively tonic
inhibition (e.g., Supplemental Figure 5. See also Brickley et al., 1996, 2001), this small

Figure 8: *Corticofugal activity generates polysynaptic inhibition in the shell IC.* **A)** Example recording from a shell IC neuron voltage clamped near the reversal potential for synaptic excitation, before and after bath application of glutamate receptor blockers. Gray traces are overlays of individual trials, black is average. Of note is that the powerful increase in IPSCs during corticofugal stimulation (blue bar) in control conditions is abolished by blocking glutamate receptors. **B)** The normalized IPSC rate is binned every 100 ms for statistical comparisons. Asterisks denote p<0.001 following Sidak's post-hoc tests comparing appropriate control and drug timepoints. Of note is that significance is limited to timepoints during optical stimulation of corticofugal axons. **C)** Average traces showing temporal integration of corticofugal EPSPs in current-clamp before (black) and after (magenta) bath application of the GABA$_A$ receptor antagonist SR95531. Shading is ± SEM. **D)** Summary data for the effect of SR95531 on corticofugal transmission current-clamp. Values are the ratio of voltage integrals in drug and control conditions. Asterisk denotes statistical significance.

change in membrane resistance in SR95531 is unlikely to account for our observed
effect on EPSP summation: The relative increase in EPSP integral was more than 4-fold
larger than the change in membrane resistance, and there was no significant pairwise
correlation between the magnitude of EPSP enhancement and change in membrane
resistance after SR95531 application (Pearson's coefficient ± standard deviation from
100 bootstrapped iterations: -0.25 ± 0.36, p = 0.43). Thus, auditory cortico-collicular
transmission generates intra-collicular inhibition that regulates how IC neurons integrate
descending signals over time.
291
292 **Discussion**
293
Although decades of *in vivo* physiology show that auditory cortical activity inhibits IC
sound responses, the underlying synaptic mechanisms have been difficult to pin down.
Indeed, anatomy data suggest that auditory cortical axons preferentially target
glutamate, rather than GABA neurons in the IC (Nakamoto et al., 2013; Chen et al.,
2018). This anatomical motif is in striking contrast to the often studied pyramidal
neuron-based microcircuits of sensory cortices and hippocampus, where long-range
excitation powerfully drives feed-forward inhibition to sharpen temporal fidelity of
afferent signals: Glutamatergic synapses onto GABAergic interneurons in these circuits
are typically more numerous, have higher synaptic conductance, and/or higher release
probability than onto pyramidal projection neurons (Acsády et al., 1998; Lawrence et al.,
2004; Gabernet et al., 2005; Cruikshank et al., 2007). At first glance, the anatomical
constraints of the auditory cortico-collicular pathway suggest that descending inhibition
of the IC is unlikely to operate via a classical feedforward recruitment of locally
projecting IC GABA neurons. Instead, auditory cortex may effectively recruit long-range
inhibitory afferents to the IC (e.g., dorsal nucleus of the lateral lemniscus; Budinger et
al., 2000), or alternatively may provide monosynaptic inhibition via a newly discovered,
long-range projection from VIP GABAergic neurons (Bertero et al., 2021).
311
However, our experiments reveal cellular and circuit mechanisms that enable the
auditory cortex to trigger intra-collicular inhibition in absence of a dense innervation onto
IC GABA neurons. Consistent with the anatomical data (Nakamoto et al., 2013; Chen et
al., 2018), corticofugal synapses tended to generate smaller EPSPs in IC GABA
compared to glutamatergic neurons. Nevertheless, two cellular mechanisms enabled
corticofugal activity to effectively increase firing rates of IC GABA neurons. First,
approximately half of shell IC GABA neurons encountered in our study fired tonically,
similar to findings in the NPY-positive GABA neurons of the central IC (Silveira et al.,
2020). Thus, even apparently weak descending EPSPs sufficed to increase the basal
firing rates of GABA neurons. Second, repetitive firing of corticofugal axons drives

robust polysynaptic excitation in IC GABA neurons due to local excitatory circuitry,
thereby orchestrating a multisynaptic cascade leading to intra-collicular inhibition.
324
We did not observe direct monosynaptic GABAergic inputs from auditory cortex VIP
neurons to the shell IC (Bertero et al., 2021). *In vitro*, the IPSC barrages recruited
during auditory cortex stimulation were entirely blocked by glutamate receptor
antagonists, indicating the necessity for intermediary, excitatory synapses onto IC
GABA neurons with locally projecting axons. This result is perhaps surprising, as our
stimulation paradigm was quite strong (25 light flashes at 50 Hz) and likely sufficed to
activate descending synapses with low release probability. Furthermore, Bertero et al.
showed that VIP-GABA axons terminate in the dorsomedial IC shell where our
recordings were obtained. However, an absence of evidence is not conclusive evidence
of absence: This newly discovered pathway may instead operate via non-canonical
mechanisms such as extrasynaptic diffusion of GABA, or via release of the
neuropeptide VIP. These synaptic properties would be difficult to quantify with our
whole-cell patch-clamp recordings. In addition, selective viral tropism may have
hindered the expression of our optogenetic construct in auditory cortical VIP neurons;
extensive follow-up experiments are necessary to clarify these issues. Rather, our
results show that intra-collicular mechanisms suffice to explain at least some of the
auditory cortex' inhibitory actions on IC sound responses.
342
Functionally, the polysynaptic activity driven by descending transmission could enable a
myriad of "high-level" computations in the shell IC. For example, previous work in
piriform cortex shows that recurrent excitatory synapses from principal cells onto
interneurons trigger feedback inhibition and truncates the duration of principal cell
responses to incoming activity from the olfactory bulb. Although odorant signals from
the bulb's lateral olfactory tract are graded with respect to odorant concentration, the
output of piriform cortex principal cells is consequently rendered concentration invariant
by this feedback inhibition which powerfully limits spiking to odor onset (Bolding and
Franks, 2018). Our findings reveal a conceptually similar microcircuit architecture in the
shell IC. Corticofugal projections, which potentially transmit behaviorally relevant
information, could thus enhance IC glutamate neurons' ability to trigger feedback
inhibition and promote a level-invariant representation of behaviorally relevant sounds in
the IC. Such a computation may be particularly useful in creating invariant neural
responses necessary for sound localization in complex, reverberant environments
(Hartmann, 1983; Rakerd and Hartmann, 1986). Indeed, the onset spiking response of
IC neurons represents directional signals that are less corrupted by reverberant
reflections and inter-aural decorrelation than later, sustained responses (Devore et al.,
2009). It is not unreasonable to assume that onset spiking in glutamate neurons could
recruit feedback inhibition that minimizes sustained activity during the reverberant tail of

sounds, and that this mechanism is enhanced by "top-down" cortical signals during
periods of increased vigilance and attention. However, extensive future studies are
required to elucidate the behaviorally relevant conditions that trigger descending
inhibitory control, and to understand how corticofugal projections might flexibly switch
between amplifying and inhibiting tectal sound responses.
367
368 **Methods**
369
The experiments were approved by the University of Michigan's IACUC and performed
according to NIH's Guide for the care and use of laboratory animals. *In vitro*
experiments were performed on 5-8 week old male or female offspring of VGAT-ires-cre
x Ai14 fl/fl or VGluT2-cre x G42 breeder pairs from our colony (Figures 2-7) or 5-8 week
old C57/Bl6J mice ordered from Jackson labs (Figure 8). For *in vivo* experiments with
wild-type mice, we used 4-7 week old offspring of CBA x C57Bl6/J matings (Figure
1B,C) or 7-10 week old C57/BL6J mice ordered from Jackson labs (Figure 1E,F). The
raw data from some of these *in vivo* recordings were also included in a previous set of
distinct analyses (Oberle et al., 2022).
379
Surgery for intracranial virus injections: Our virus injection protocol is described in detail
in Oberle et al., 2022. Briefly, mice were anesthetized with 4-5% isoflurane in O_2,
mounted in a stereotaxic frame, and isoflurane was lowered to 1-2% while body
temperature was maintained near 37-38° C using a heating blanket. Mice were
administered 5 mg/kg carprofen as a pre-surgical analgesic, a small incision was made
in the scalp, and 2% lidocaine was then applied to the wound margins. A 200-400 μm
craniotomy was carefully drilled over the left auditory cortex (-2.75 mm from Bregma,
centered on the lateral ridge) or left IC (0.9 mm caudal and 1 mm lateral from lambda
suture) to inject 100-200 nL of pAAV-Syn-Chronos-GFP (Addgene #59170-AAV1) or
pAAV-syn-flex-rc[ChrimsonR-tdTomato] (Addgene #62723-AAV5) virus. At the end of
the surgery, the craniotomy was filled with bone wax and the skin was sutured.
Immediately following surgery, mice received an analgesic injection of buprenorphine
(0.03 mg/kg, s.c.) and recovered on a heating pad before returning to their home cage.
A post-operative dose of carprofen was administered ~24 hours later.
394
In vivo electrophysiology: Mice were deeply anesthetized with isoflurane, mounted in
the stereotaxic frame, and craniotomies were carefully opened over the left IC and
auditory cortex as described above and in Oberle et al. 2022. For optogenetic
stimulation, we left the dura intact and implanted a cranial window over the auditory
cortex. For electrical stimulation we made a small slit in the dura and the craniotomy
was sealed with silicone elastomer. The IC craniotomy was plugged with silicone
elastomer, a titanium headbar was implanted with dental cement, and the mouse was
removed from the stereotax before being re-anesthetized with urethane (1.5 g/kg, i.p.)

and head-fixed in a custom-made sound attenuation chamber. Body temperature during
the recording session was maintained at 37-38° C with a custom designed, feedback-
controlled heating blanket. Optogenetic stimulation was performed with a 0.5 NA, 400
μm core optic fiber (Thorlabs M45L02) coupled to a 470 nm LED (Thorlabs M470F3)
positioned <1 mm away from the auditory cortex cranial window. For electrical
stimulation experiments, a bipolar platinum-iridium electrode (FHC 30210) was carefully
inserted ~800 μm into auditory cortex at an angle roughly perpendicular to the cortical
layers and shocks were delivered via a custom stimulus isolator. Whole-cell current-
clamp recordings were obtained ~100-400 μm from the dura with pipettes containing (in
mM): 115 K-Gluconate, 4 KCl, 0.1 EGTA, 10 HEPES, 14 Tris-Phosphocreatine, 4 Mg-
ATP, 0.5 Tris-GTP, 4 NaCl, pH 7.2-7.3, 290 mOsm (open tip resistance: 5-10 MΩ).
414
In vitro electrophysiology: 2-4 weeks following viral injections, mice were deeply
anesthetized with isoflurane and 200-300 μm brain slices containing both IC
hemispheres were prepared in warm (~34° C) oxygenated ACSF containing (in mM):
119 NaCl, 25 $NaHCO_3$, 3 KCl, 1.25 NaH_2PO_4, 15 glucose, 1 $MgCl_2$, 1.3 $CaCl_2$, 1
ascorbate, 3 pyruvate. A small cut was typically made in the lateral portion of the right
cerebellum or right IC to visually identify the un-injected hemisphere. Slices were then
incubated at 34° C in a holding chamber filled ACSF for 25-30 min and then stored at
room temperature until use. During experiments, slices were mounted in a submersion
chamber continuously perfused with oxygenated ACSF heated to 32-34° C (2-4 mL/min;
chamber volume: ~ 1 mL). Neurons in the dorso-medial shell IC were visualized via DIC
or Dodt contrast optics using a 40x or 63x objective (Zeiss Axioskop 2 FS Plus or
Olympus BXW51 microscope). GABA neurons were identified from tdTomato or GFP
fluorescence. Presumptive glutamate neurons were identified based on lack of
fluorescence in VGAT-cre x Ai14 mice. Whole-cell current-clamp recordings were
obtained using pipettes filled with the same K^+ rich internal solution as employed for in
vivo recordings (open tip resistance: 3-6 MΩ). In some experiments, 30 μM Alexa 488
or 0.1% biocytin were added to the internal solution to visualize neuronal morphology
via online fluorescence or post-hoc histological reconstruction. Voltage clamp
recordings were obtained with a Cs^+ based solution containing (in mM): 110 Cesium
Methanesulfonate, 10 QX-314-Bromide, 0.1 EGTA, 10 HEPES, 0.5 Tris-GTP, 4.5
MgATP, 5 TEA-Cl, 10 Tris-phosphocreatine. Experiments were generally conducted
within 3-4 hours following slice preparation. Chronos-expressing axons were stimulated
via 2-5 ms wide-field flashes of blue light delivered through the microscope objective.
ChrimsonR-expressing IC neurons were stimulated via 0.5-1 ms flashes from a 400 μm
diameter optic fiber (0.5 NA) coupled to a 625 nm LED <1 mm above the slice.
440
Data Acquisition: Data were acquired using a Molecular Devices Multiclamp 700B or
AM Systems model 2400 patch-clamp amplifier, online filtered at 2-10 kHz, and digitized

at 50 kHz with a National Instruments PCI-6343 card + BNC2090A interface controlled
by Matlab based acquisition software (Wavesurfer). As shown in Figure 4, many GABA
neurons fired tonically *in vitro*. Consequently, negative bias current (-5 to -200 pA) was
often injected to hyperpolarize neurons and prevent spontaneous or optogenetically
triggered spiking when recording EPSPs in GABA neurons.
448
Data analysis: All experiments were analyzed using custom scripts written in Matlab.
Analyses in Figures 1, 2, 5, 7, and 8 were run on averages of multiple trials (typically
>10 trials per condition) after baseline subtraction and lowpass filtering at 1 kHz. EPSP
peak amplitudes were calculated by averaging data points ± 0.1 ms around the local
maximum following optogenetic stimulation. Halfwidths were defined as the full width at
half-maximum of the peak. Voltage integrals were calculated using the Matlab functions
trapz() or cumtrapz(). Event detection analyses in Figures 3, 4, 6, and 8 were conducted
by differentiating the membrane voltage or current traces, smoothing the data (1-5 ms
sliding window), and using threshold crossing or peak detection algorithms to identify
synaptic events. Synaptic current waveforms (Figure 4C,D) were generated in
Wavemetrics Igor Pro using the Neuromatic package (Rothman and Silver, 2018).
460
Statistics: In the text, n and N refer to the number of neurons and mice, respectively.
Sample sizes were not explicitly pre-determined prior to data collection. However, the
number of cells and mice in these experiments are in accordance with commonly
accepted standards in the field. Data were tested for normality using a Lilliefors test
prior to statistical comparisons. Parametric tests were used for normally distributed
data, while non-parametric tests were used if one or more of the datasets deviated from
normal. Alpha is corrected for multiple comparisons in post-hoc significance tests
following ANOVA. Statistics were run in Matlab or Graphpad Prism 9. Pearson's
coefficient and associated bootstraps were calculated in Matlab with the functions corr()
and bootstrp(), respectively.
471

472   **Author Contributions:** HMO + PFA conducted and analyzed experiments of Figures 1
473   + 8. ANF performed stereotaxic injections with contributions from HMO. HMO + ANF
474   conducted histology and confocal imaging. PFA performed and analyzed experiments of
475   Figures 2-7, interpreted the results, and wrote the paper.
476

Acknowledgments: Funding was provided by the Whitehall Foundation, Hearing
Health Foundation, and NIH/NIDCD R01DC019090 to PFA, as well as T32DC005356
and T32DC000011 to HMO. We thank Dr. Michael Roberts for critical comments on the
manuscript, Dr. Kevin Jones for providing the G42 mouse used for histology in
Supplemental Figure 2, Dr. Meike Rogalla for the illustrations in Figure 1, and Jordyn
Czarny for technical assistance with histology.

Supplemental Figure 1: *IC GABA and glutamate neurons have similar cellular properties.*
A) Tile scan of the IC in a VGAT-cre x Ai14 mouse. An area of interest is denoted by the dashed line and shown at higher magnification in panel B. Scale bar = 500 μm.
B) Magnification of dashed rectangle in panel A. The micrograph was contrast enhanced to highlight tdTomato-positive GABA neurons and tdTomato-negative, presumptive glutamate neurons visible as dark "shadows". Scale bar = 100 μm.
C-E) Examples of distinct types of hyperpolarizing and depolarizing responses in GABA neurons encountered in the shell IC.
F-H) Same as A-C, but for glutamate neurons. Of note is the qualitative similarity in firing patterns and membrane properties of both neuron classes.

Supplemental Figure 2: *G42 mouse line labels a sparse population of IC GABA neurons.*

A) Confocal micrograph from the IC of a G42-GFP mouse. Scale bar = 500 µm. Dashed rectangle denotes area of interest in the dorso-medial shell IC.

B) Area of interest from panel A shown at higher magnification. Scale bar = 100 µm.

C) Example current-clamp recording from a GFP+ neuron in the dorso-medial IC of a G42 mouse. The majority of GFP+ neurons recorded in G42 mice fired spontaneously when recorded in current-clamp with 0 pA bias current (8/11 neurons tested).

D) Example trace from the same neuron in panel C, but hyperpolarized with -30 pA bias current. A +40 pA current injection step reveals an adapting firing pattern. 18/21 neurons tested had similar adapting spike patterns when injected with current steps following membrane hyperpolarization to prevent spontaneous spiking. The remaining GFP+ neurons either fired a transient burst of spikes atop a T-type like hump (n=2 cells; See also Figure 3E from Oberle et al., 2022), whereas n = 1 cell showed minimal spike rate adaptation.

Supplemental Figure 3: *Polysynaptic excitation drives spikes in IC GABA neurons*

A) Example recording where repetitive corticofugal stimulation triggered spiking in an IC

GABA neuron. Records are multiple overlaid single trials.

B) EPSPs evoked by the first two light flashes in (A) are shown at a faster time base to

highlight the onset jitter and polysynaptic nature of the EPSPs.

Supplemental Figure 4: *Additional examples of polysynaptic EPSPs in GABA neurons.*

A) Example overlaid traces (10 consecutive trials) from an IC GABA neuron. Blue bar

denotes stimulation of corticofugal fibers.

B) Traces in A expanded to highlight EPSPs evoked during the first 10 light pulses. Blue

ticks indicate onset of individual light flashes.

C) Latency histogram for EPSPs of the neuron of A+B, showing two distinct peaks.

D-F) Same as panels A-C but for a different GABA neuron.

Supplemental Figure 5: *Spontaneous IPSCs in shell IC neurons are predominantly*
mediated by GABA$_A$ receptors.
A) Example voltage-clamp recording from a shell IC neuron held at +8 mV. Of note are
the spontaneous IPSCs often occurring in bursts, suggesting they are mediated by
action potentials in local GABA neurons. Asterisk denotes one such IPSC burst shown
at a faster time base in the inset.
B) In the same cell as shown in panel A, bath application of the GABA$_A$ receptor
antagonist SR95531 (10 µM) blocks the majority of spontaneous IPSCs. Although quite
rare, a few small amplitude events nevertheless remained in the presence of SR95531
(asterisk and inset; median IPSC rate in SR95531: one event per 17.95 seconds).
These SR95531 resistant outward currents may reflect long-range glycinergic inputs
from the ventral nucleus of the lateral lemniscus (Moore and Trussell, 2017).
C) Summary data for n = 7 cells showing that SR95531 profoundly reduces
spontaneous IPSC rates, indicating that inhibitory transmission in shell IC neurons is
mostly GABAergic.

References

Acsády L, Kamondi A, Sík A, Freund T, Buzsáki G (1998) GABAergic cells are the major postsynaptic targets of mossy fibers in the rat hippocampus. J Neurosci Off J Soc Neurosci 18:3386–3403.

Asilador A, Llano DA (2020) Top-Down Inference in the Auditory System: Potential Roles for Corticofugal Projections. Front Neural Circuits 14:615259.

Beneyto M, Winer JA, Larue DT, Prieto JJ (1998) Auditory connections and neurochemistry of the sagulum. J Comp Neurol 401:329–351.

Bertero A, Garcia C, Apicella AJ (2021) Corticofugal VIP Gabaergic Projection Neurons in the Mouse Auditory and Motor Cortex. Front Neural Circuits 15:714780.

Blackwell JM, Lesicko AM, Rao W, De Biasi M, Geffen MN (2020) Auditory cortex shapes sound responses in the inferior colliculus. eLife 9:e51890.

Bledsoe SC, Shore SE, Guitton MJ (2003) Spatial representation of corticofugal input in the inferior colliculus: a multicontact silicon probe approach. Exp Brain Res 153:530–542.

Bolding KA, Franks KM (2018) Recurrent cortical circuits implement concentration-invariant odor coding. Science 361:eaat6904.

Born G, Schneider-Soupiadis FA, Erisken S, Vaiceliunaite A, Lao CL, Mobarhan MH, Spacek MA, Einevoll GT, Busse L (2021) Corticothalamic feedback sculpts visual spatial integration in mouse thalamus. Nat Neurosci 24:1711–1720.

Boyd AM, Sturgill JF, Poo C, Isaacson JS (2012) Cortical feedback control of olfactory bulb circuits. Neuron 76:1161–1174.

Brickley SG, Cull-Candy SG, Farrant M (1996) Development of a tonic form of synaptic inhibition in rat cerebellar granule cells resulting from persistent activation of GABAA receptors. J Physiol 497 (Pt 3):753–759.

Brickley SG, Revilla V, Cull-Candy SG, Wisden W, Farrant M (2001) Adaptive regulation of neuronal excitability by a voltage-independent potassium conductance. Nature 409:88–92.

Briggs F, Usrey WM (2011) Corticogeniculate feedback and visual processing in the primate. J Physiol 589:33–40.

Budinger E, Heil P, Scheich H (2000) Functional organization of auditory cortex in the Mongolian gerbil (Meriones unguiculatus). IV. Connections with anatomically characterized subcortical structures. Eur J Neurosci 12:2452–2474.

Chattopadhyaya B, Di Cristo G, Higashiyama H, Knott GW, Kuhlman SJ, Welker E, Huang ZJ (2004) Experience and activity-dependent maturation of perisomatic GABAergic innervation in primary visual cortex during a postnatal critical period. J Neurosci Off J Soc Neurosci 24:9598–9611.

Chen C, Cheng M, Ito T, Song S (2018) Neuronal Organization in the Inferior Colliculus Revisited with Cell-Type-Dependent Monosynaptic Tracing. J Neurosci 38:3318–3332.

Crandall SR, Cruikshank SJ, Connors BW (2015) A corticothalamic switch: controlling the thalamus with dynamic synapses. Neuron 86:768–782.

Cruikshank SJ, Lewis TJ, Connors BW (2007) Synaptic basis for intense thalamocortical activation of feedforward inhibitory cells in neocortex. Nat Neurosci 10:462–468.

Devore S, Ihlefeld A, Hancock K, Shinn-Cunningham B, Delgutte B (2009) Accurate sound localization in reverberant environments is mediated by robust encoding of spatial cues in the auditory midbrain. Neuron 62:123–134.

Diamond JS, Jahr CE (1997) Transporters buffer synaptically released glutamate on a submillisecond time scale. J Neurosci Off J Soc Neurosci 17:4672–4687.

Gabernet L, Jadhav SP, Feldman DE, Carandini M, Scanziani M (2005) Somatosensory integration controlled by dynamic thalamocortical feed-forward inhibition. Neuron 48:315–327.

Hartmann WM (1983) Localization of sound in rooms. J Acoust Soc Am 74:1380–1391.

Hattox AM, Nelson SB (2007) Layer V neurons in mouse cortex projecting to different targets have distinct physiological properties. J Neurophysiol 98:3330–3340.

Kirchgessner MA, Franklin AD, Callaway EM (2021) Distinct "driving" versus "modulatory" influences of different visual corticothalamic pathways. Curr Biol CB:S0960-9822(21)01265-3.

Kong L, Xiong C, Li L, Yan J (2014) Frequency-specific corticofugal modulation of the dorsal cochlear nucleus in mice. Front Syst Neurosci 8

Lawrence JJ, Grinspan ZM, McBain CJ (2004) Quantal transmission at mossy fibre targets in the CA3 region of the rat hippocampus. J Physiol 554:175–193.

Mitani A, Shimokouchi M, Nomura S (1983) Effects of stimulation of the primary auditory cortex upon colliculogeniculate neurons in the inferior colliculus of the cat. Neurosci Lett 42:185–189.

Moore LA, Trussell LO (2017) Corelease of Inhibitory Neurotransmitters in the Mouse Auditory Midbrain. J Neurosci Off J Soc Neurosci 37:9453–9464.

Nakamoto KT, Mellott JG, Killius J, Storey-Workley ME, Sowick CS, Schofield BR (2013) Ultrastructural examination of the corticocollicular pathway in the guinea pig: a study using electron microscopy, neural tracers, and GABA immunocytochemistry. Front Neuroanat 7

Naumov V, Heyd J, de Arnal F, Koch U (2019) Analysis of excitatory and inhibitory neuron types in the inferior colliculus based on Ih properties. J Neurophysiol 121:2126–2139.

Nwabueze-Ogbo FC, Popelár J, Syka J (2002) Changes in the acoustically evoked activity in the inferior colliculus of the rat after functional ablation of the auditory cortex. Physiol Res 51 Suppl 1:S95–S104.

Oberle HM, Ford AN, Dileepkumar D, Czarny J, Apostolides PF (2022) Synaptic mechanisms of top-down control in the non-lemniscal inferior colliculus. eLife 10:e72730.

Ono M, Yanagawa Y, Koyano K (2005) GABAergic neurons in inferior colliculus of the GAD67-GFP knock-in mouse: electrophysiological and morphological properties. Neurosci Res 51:475–492.

Popelár J, Nwabueze-Ogbo FC, Syka J (2003) Changes in neuronal activity of the inferior colliculus in rat after temporal inactivation of the auditory cortex. Physiol Res 52:615–628.

Popelář J, Šuta D, Lindovský J, Bureš Z, Pysanenko K, Chumak T, Syka J (2016) Cooling of the auditory cortex modifies neuronal activity in the inferior colliculus in rats. Hear Res 332:7–16.

Pouille F, Scanziani M (2001) Enforcement of temporal fidelity in pyramidal cells by somatic feed-forward inhibition. Science 293:1159–1163.

Qi J, Zhang Z, He N, Liu X, Zhang C, Yan J (2020) Cortical Stimulation Induces Excitatory Postsynaptic Potentials of Inferior Colliculus Neurons in a Frequency-Specific Manner. Front Neural Circuits 14:591986.

Rakerd B, Hartmann WM (1986) Localization of sound in rooms, III: Onset and duration effects. J Acoust Soc Am 80:1695–1706.

Randle JC, Guet T, Cordi A, Lepagnol JM (1992) Competitive inhibition by NBQX of kainate/AMPA receptor currents and excitatory synaptic potentials: importance of 6-nitro substitution. Eur J Pharmacol 215:237–244.

Rothman JS, Silver RA (2018) NeuroMatic: An Integrated Open-Source Software Toolkit for Acquisition, Analysis and Simulation of Electrophysiological Data. Front Neuroinformatics 12:14.

640   Schofield BR (2009) Projections to the inferior colliculus from layer VI cells of auditory
641   cortex. Neuroscience 159:246–258.

642   Sherman SM, Usrey WM (2021) Cortical control of behavior and attention from an
643   evolutionary perspective. Neuron:S0896-6273(21)00462-1.

644   Silveira MA, Anair JD, Beebe NL, Mirjalili P, Schofield BR, Roberts MT (2020)
645   Neuropeptide Y Expression Defines a Novel Class of GABAergic Projection
646   Neuron in the Inferior Colliculus. J Neurosci 40:4685–4699.

647   Slater BJ, Willis AM, Llano DA (2013) Evidence for layer-specific differences in auditory
648   corticocollicular neurons. Neuroscience 229:144–154.

649   Song JH, Lucaci D, Calangiu I, Brown MTC, Park JS, Kim J, Brickley SG, Chadderton P
650   (2018) Combining mGRASP and Optogenetics Enables High-Resolution
651   Functional Mapping of Descending Cortical Projections. Cell Rep 24:1071–1080.

652   Stebbings KA, Lesicko AMH, Llano DA (2014) The auditory corticocollicular system:
653   Molecular and circuit-level considerations. Hear Res 314:51–59.

654   Syka J, Popelár J (1984) Inferior colliculus in the rat: neuronal responses to stimulation
655   of the auditory cortex. Neurosci Lett 51:235–240.

656   Usrey WM, Sherman SM (2019) Corticofugal circuits: Communication lines from the
657   cortex to the rest of the brain. J Comp Neurol 527:640–650.

658   Vila C-H, Williamson RS, Hancock KE, Polley DB (2019) Optimizing optogenetic
659   stimulation protocols in auditory corticofugal neurons based on closed-loop spike
660   feedback. J Neural Eng 16:066023.

661   Williamson RS, Polley DB (2019) Parallel pathways for sound processing and functional
662   connectivity among layer 5 and 6 auditory corticofugal neurons. eLife 8:e42974.

663   Winer JA, Larue DT, Diehl JJ, Hefti BJ (1998) Auditory cortical projections to the cat
664   inferior colliculus. J Comp Neurol 400:147–174.

665   Zhang Y, Suga N (1997) Corticofugal amplification of subcortical responses to single
666   tone stimuli in the mustached bat. J Neurophysiol 78:3489–3492.

667   Zhang Y, Suga N, Yan J (1997) Corticofugal modulation of frequency processing in bat
668   auditory system. Nature 387:900–903.

669