Correlated responses of energy metabolism, development and reproduction to

evolution for increasing heat tolerant in *Drosophila subobscura.*

Andrés Mesas[1,2] and Luis E. Castañeda[3]

[1]Millenium Institute of Oceanography (IMO), Universidad de Concepción, Concepción, Chile.

[2]Departemento de Sistemas Acuáticos, Facultad de Ciencias Ambientales, Universidad de

Concepción, Concepción, Chile.

[3]Programa de Genética Humana, Instituto Ciencias Biomédicas, Facultad de Medicina, Universidad

de Chile, Santiago, Chile.

Correspondence:

Andrés Mesas. Email: andresmesasp@gmail.com

Luis E. Castañeda. Email: luis.castaneda@uchile.cl

Running Title: Correlated evolution of metabolism and fitness

Keywords: Heat tolerance evolution, intensity of thermal stress, metabolic rate, G6P branchpoint,

early fecundity, egg-to-adult viability.

Funding information: This work was funded by Fondo Nacional de Investigación Científica y

Tecnológica grant number 1140066 (L.E.C.)

Abstract

1. Adaptation to increasing environmental temperatures implies an increase of the upper thermal

limit (CTmax), but also an enhanced capacity of the organisms to perform fundamental biological

functions at high temperatures.

2. Evolution of stress resistance is companied with a metabolic depression, which allows to

organisms have reduced maintaining costs and allocate the energy surplus to energy demanding

resistance mechanisms and reproduction.

3. Heat tolerance evolution depends on the heat intensity selection and it could have different

consequences on the evolution of correlated traits such as energy metabolism and fitness-related

traits. Thus, we evaluated the correlated responses of metabolic rate, metabolism-related enzymes

(G6P branchpoint enzymes), reproductive and preadult survival in control and selected lines for

increasing heat tolerance in *Drosophila subobscura*.

4. We did not find a correlated response of metabolic rate associated with heat tolerance evolution,

but we detected a reduced hexokinase activity in the slow-ramping selected lines. On the other hand,

we found an increase in fecundity in the fast- and slow-ramping selected lines, but an increase in

egg-to-adult viability was only found in the slow-ramping selected lines.

5. Our results show that the evolution of heat tolerance does not imply an increase in energy

demand neither a tradeoff with reproduction and preadult survival, which suggests that heat

tolerance evolution induces correlated response to increase the performance at high temperatures.

Therefore, spatial and temporal variability of thermal stress intensity should be taking into account

in future studies if we want to understand and predict the adaptive response to ongoing and future

climatic conditions.

Introduction

Environmental thermal stress involves strong selective pressures that can led to important

evolutionary changes depending on the presence of genetic variation in natural populations (Umina

et al, 2005; Crozier & Hutchings, 2014; Chick et al, 2020). The pervasive effects of the temperatures

on ectothermic organisms cause that fitness, abundance and distribution of their populations are

susceptible to the current and future increase of temperature (Deutsch et al, 2008; Angilletta, 2009;

Chen et al, 2011). Therefore, it is crucial to know and understand the capacity of ectothermic

organisms to withstand the global warming.

The ability to tolerate higher temperatures implies that together with an increase of the upper

thermal limit, organisms must exhibit an array of responses of at different levels of biological

organization, allowing them to maintain their biological performance at high temperatures (Huey &

Kingsolver, 1993; Pörtner et al, 2006; Mesas et al, 2021; McGaughran et al, 2021).

Among these responses, metabolic depression, defined as a reduction of the energy expenditure

associated to maintenance of organisms, is considered an important mechanism associated with

resistance to different types of environmental stress (Guppy & Withers, 1999; Storey & Storey,

2004). Plastic and evolutionary decreases of energy metabolism should have important ecological

and evolutionary implications in stressful conditions because it allows to organisms reduce their

minimum energetic requirements and redistribute the energy surplus to biological functions close to

fitness (Atkinson, 1977; Hochachka & Somero, 2002, Storey & Storey, 2004). For example, a

reduced standard metabolic rate (SMR) has evolved by natural selection in the garden snail (Artacho

& Nespolo, 2009), whereas evolutionary responses increasing tolerance to desiccation, starvation,

hypoxia, and high temperatures have been accompanied by a reduction in metabolic rate (Hoffmann

& Parson, 1989; Padfield et al, 2016; Regan et al, 2017; Brown et al, 2019).

In ectotherms, metabolism increases with environmental temperatures, inducing high metabolic

costs in individuals exposed to heat stress as expected by global warming (Lighton & Turner, 2004;

Huey & Kingsolver, 2019). Therefore, metabolic depression could be an important strategy to

prevent that metabolism exceeds the physiological limits and energy expenditure, which has

adversely effects on fitness under extreme heat conditions (Marshall & McQuaid, 2011; González-

Tokman et al, 2020; Pilakouta et al, 2020). This metabolic response to high temperature could

partially explain the counter gradient variation proposed in metabolic cold adaptation hypothesis

(Addo-Bediako, 2002; Gaston et al, 2009), which establishes that ectotherm species from temperate

habitats have lower metabolic rates than those living in colder environments (Addo-Bediako et al,

2002; Sylvestre et al, 2007; Schaefer & Walters 2010; Bruning et al, 2013; Sinnatamby et al, 2015;

Pilakouta et al, 2020). On other hand, experimental populations exposed to warm environmental

conditions during several generations showed an evolutionary increase of their thermal tolerance

accompanied with a reduced metabolic rate (Padfield et al, 2016, but see Mallard et al, 2018).

However, heat tolerance evolution and the correlated responses of associated traits depend on the

intensity of heat stress selection (Santos et al 2012; Mesas et al, 2021). For instance, Mesas et al

(2021) found evolutionary responses of heat tolerance regardless of intensity of heat selection, which

did have an effect on the evolutionary response of the thermal performance in *Drosophila subobscura*

(Mesas et al, 2021). The effects of thermal selection intensity on thermal-related traits were

previously predicted by Santos et al (2012) whom, using mathematical simulation of artificial

selection experiments, found a decrease of metabolic rate only when selection for increasing heat

tolerance was performed using slow-ramping protocols. This correlated response between heat

tolerance and metabolic rate can be explained because flies with low metabolic rate can withstand

longer exposure in slow-ramping assays, resulting in higher thermal tolerance (Rezende et al, 2011;

Santos et al, 2012).

Energy metabolism is emergent property of interactions between different cellular processes and

metabolic pathways (Berrigan & Hoang, 1999; Montooth et al, 2003; Brown et al, 2004; Matoo et al,

2019). Then, exposure to different heat stress intensities should results into thermal dependence of

metabolic rates and flux of metabolic pathways. For example, several genes associated with the

glucose-6-phosphate (G6P) branchpoint, which has an important role in the storage and use of

molecules as glucose, glycogen, triglycerides and ATP, have shown clinal variation in their allelic

frequencies along latitude (Oakeshott et al 1983; Duvernell & Eanes, 2000; Verrelli & Eanes, 2001;

Rank & Dahlhoff, 2002, Sezgin et al, 2004). These findings suggest that the evolution of functional

and structural properties of the enzymes associated with energy metabolism are sensitive to

environmental temperatures. Moreover, different haplotypes associated with G6P branchpoint can

differentially influence the energy flux in a particular metabolic pathway (Eanes & Hey, 1986; Zera,

1987; Verrelli & Eanes, 2001), but also affect the flux in parallel metabolic pathways, facilitating

correlated responses among fitness-related traits under stressful conditions (Zera & Zhao, 2003a).

For example, evolution of stress resistance was associated with changes in metabolic rate, fecundity,

and development in *D. melanogaster* (Hoffmann & Parson, 1989; Brown et al, 2019). On the other

hand, evolution under warm conditions resulted in a down-regulation of metabolic pathways (e.g.,

genes associated with Krebs cycle, oxidative phosphorylation, and glycolysis enzymes) and an

increase in fertility in *D. simulans* (Mallard et al, 2018). Notably, reproductive and developmental

traits have been negatively associated with heat stress in in ectotherms (Morrison & Hero, 2003; Qu

et al, 2011; Hamby et al, 2016; Porcelli et al, 2017), which should suggest that heat tolerance

evolution involves metabolic readjustments that impact on energy-demanding traits such as fitness-

related traits (Hochachka & Somero, 2002; Harshman & Zera, 2007; Tomanek & Zuzow, 2010).

In this work, we evaluated the evolutionary change on energy metabolism and life history traits in

response to artificial selection for increasing heat tolerance in *D. subobscura*, using two different

thermal selection protocols to measure heat tolerance: slow and fast ramping protocols, which

involve different heat intensity selection regimens. We hypothesized that increased heat tolerance

evolves associated with a reduction of energy metabolism, especially if thermal selection is

performed using slow-ramping protocols in comparison to fast-ramping protocols. Additionally, we

proposed that the metabolic reduction will decrease the maintenance energy requirements and

generate an energy surplus, which should positively affect to reproductive and developmental traits.

To evaluate these hypotheses, we measured the routine metabolic rate (RMR) and the activity of

enzymes associated with the G6P branchpoint as proxies of the energy metabolism in experimental

populations of *D. subobscura* previously selected for increasing heat tolerance (Mesas et al, 2021). We

measured the activity of these enzymes in flies exposed to two different thermal conditions (non-

stress: 21 ºC, and stress: 32 ºC) to evaluate if heat stress could impact on the enzymatic responses of

selected lines. Additionally, we evaluated the early fecundity and egg-adult viability the evolved

population to determine effect of heat tolerance evolution on life-history traits of *D. subobscura*.

Methodology

Heat knockdown temperature selection

A mass-bred population was established from the offspring of 100 isofemale lines derived of

inseminated females collected in Valdivia, Chile (39.8 ºS; 73.2 ºW). These flies were maintained as a

large outbred population, at 21 ºC (12:12 light:dark cycle) and feed with David's killed-yeast

Drosophila medium. After 6 generations, this population was split into 12 experimental lines, which

were assigned in triplicate to the different artificial selection protocols and their respective controls:

fast-ramping control, fast-ramping selected, slow-ramping control, and slow-ramping selected lines.

See details of the experimental evolution protocol in Mesas et al (2021).

For each replicated lines, 160 four-days old virgin females were randomly mated with two no-related

males from the same replicated line. After two days, males were discarded, and vials were checked

for positive oviposition. Then, 120 adult females were individually placed in a capped 5-mL glass

vial, which was attached to two racks, each one with capacity to attach 60-capped vials. Racks were

immersed in a water tank with an initial temperature of 28 ºC, which was controlled by a heating unit

(Model ED, Julabo Labortechnik, Seelbach, Germany). After an equilibration time of 10 min,

temperature was increased with a rate of 0.08 ºC min^{-1} for the flies belong to the slow-ramping

selection protocol, or 0.4 ºC min^{-1} for the flies belong to the fast-ramping selection protocol. Each

assay was photographed every 3 s with a high-resolution camera (D5100, Nikon, Tokyo, Japan). All

photos for each assay were visualized to score the knockdown temperature for each fly, which was

defined as the temperature at which each fly ceased to move. Flies were ranked according to their

knockdown temperatures and we selected vials containing offspring of the 40 most tolerant flies

(upper 30% of each assay) to found the next generation. For the fast-control and slow-control lines,

the knockdown temperature was measured as mentioned above but offspring to found the next

generation were randomly chosen.

Artificial selection for heat tolerance was performed for 16 generations, after which flies were

dumped in acrylic cages and maintained without selection at 21ºC (12:12 light-dark cycle) until the

measurement of metabolic and life-history traits. For logistic reasons, we used the fast -ramping

control experimental line as control line for statistical comparisons.

Metabolic rate

The routine metabolic rate (RMR) was measured as the amount of CO_2 produced by individuals

four days-old virgin females from the 26[th] generation. RMR was measured in 15 females from each

experimental lines, reaching a total of 135 flies (3 thermal selection protocols x 3 replicated lines x

15 flies). Before metabolic measurements, flies were ice-cold anesthetized and weighted using a

microbalance (MXA5, Radwag, Czech Republic). Then, flies were individually placed in a metabolic

chamber, which were made of 2 cm of a Bev-A-Line tube and ending connectors glued with a

cotton mesh to avoid that flies escape form the chamber. Metabolic chambers were connected to an

eight-channel multiplexer system (RM8, Sable Systems International, NV, USA), with seven

chamber containing individual flies and one empty chamber used for baseline. Metabolic chambers

were placed in dark conditions within a thermal cabinet (PC-1, Sable Systems International, NV,

USA) at a temperature of 21 °C, which was controlled by a temperature controller (PELT-5, Sable

Systems International, NV, USA). CO2-free air was pumped into chambers at a flow of 15 mL/min

(Sierra MFC-2, Sable Systems International, NV, USA) and after 30 min of equilibration, CO2

production was sequentially measured in each chamber for 15 min using an infrared CO_2 analyzer

with a resolution of 1 ppm (Li-6251, LI-COR Bioscience, NE, USA). Measurements of CO_2 were

recorded, transformed to metabolic rate (μl CO_2/h), and analyzed using the software Expedata

(Sable Systems International, NV, USA).

Enzyme activity

The catabolic activities of enzymes from the G6P branchpoint were measured in four days-old

females from the 27[th] generation through the appearance of NADPH in kinetic assays. we evaluated

the activity of hexokinase (HEX: E.C. 2.7.1.1) involved in glucose metabolism, phosphoglucomutase

(PGM; E.C. 5.4.2.2) involved in glycogen metabolism, phosphoglucose isomerase (PGI; E.C.

5.3.1.9) involved in ATP metabolism; and glucose-6-phosphate dehydrogenase (G6PD; E.C.

1.1.1.49) involved in lipids metabolism.

From each replicated line, 100 females were collected and pooled in groups of 10 flies each.

We measured the enzyme activity of the G6P branchpoint in flies exposed to two thermal

conditions: thermal-stressed flies (five replicates were exposed to 32 °C per 1 h and then, flies were

allowed to recover at 21 °C for 2 h); and non-stressed flies (five replicates were maintained at 21 °C

for 3 h). Then, flies were quickly frozen in liquid nitrogen, homogenized in 1 ml of homogenization

buffer (0.01 M KH_2PO_4, 179 1.0 mM EDTA, pH 7.4), and centrifuged at 2,000 rpm for 2 min at 4

°C. The supernatant was aliquoted and maintained at -80 °C until enzymatic assays. The buffers for

each assayed enzyme were as follow. **HEX:** 20 mM TrisHCl, 0.5 mM NADP, 0.2 mM $MgCl_2$, 0.36

mM ATP, 5 mM D-glucose, 0.23 units/mL G6PD, pH 8.0. **PGI:** 20 mM TrisHCl, 0.28 mM NADP,

3 mM fructose-6-phosphate, 1.37 units/mL G6PD, pH 8.0. **PGM:** 20 mM TrisHCl, 0.5 mM

NADP, 1 mM $MgCl_2$, 0.83 mM glucose-1-phosphate, 3.1 units/mL G6PD, pH 8.0. **G6PD:** 20 mM

TrisHCl, 0.2 mM NADP, 18.8 mM $MgCl_2$, 3.5 mM glucose-6-phosphate, pH 8.0. Enzymatic assays

were performed adding 25 μL extracts of flies with a protein concentration of 3500 μg/mL and 200

μL of respectively assay buffer in ultraviolet plates with flat bottom wells. Immediately, the optical

density (OD) to 340 nm was measured with multiple reads per well in a microplate reader (Infinite

200 Pro, Tecan) previously heated to 29 °C. The OD was measured each 25 s during 10 min to

HEX and PGI, every 10 s during 10 minutes to PGM, and every 45 s during 15 min to G6PD.

Enzymatic activities were estimated by triplicated and mean OD was analyzed. Blank wells

containing buffer assays and double-distilled water (instead fly extracts) were measured to establish

the basal optical density of each reaction. Finally, the enzymatic activity was calculated as increment

rate of optical density through assay (ΔOD/s).

Early fecundity and egg-to-adult viability

Early fecundity was measured in fly females from 26[th] generation. For each replicated line, 10 virgin

females were individually placed in vials with David's killed-yeast *Drosophila* medium together to two

unrelated males from the same replicated line. After 24 h, females were transferred to new vials and

the number of oviposited eggs in each vial were daily counted using a stereomicroscope. This

procedure was repeated every 24 h for eight days to obtain the accumulated fecundity for each

female. Additionally, we measured the egg-adult viability as follow. Vials containing eggs oviposited

by five days-old females (age of maximal activity of oviposition in *D. subobscura*) were retained to

collect eggs. For each vial, 20 eggs were randomly transferred to a new vial, which were maintained

to 21 °C until flies will hatch.

Statistical analysis

Normality and homoscedasticity were tested for all variables, and only the enzyme activity of HEX

was squared-root transformed to meet the parametric assumptions. Then, a mixed lineal model

(hereafter 'full model') was used to evaluate the effects of thermal selection (fixed effect) and

replicated lines nested within thermal selection (random effect) on body mass (eq. 1) and RMR

(including body mass as covariate) (eq. 2):

229                $Mb_{ijk} = \mu + Selection_i + Replicate(Selection)_{ij} + e_{ijk}$ (eq. 1)

230                $RMR_{ijk} = \mu + Selection_i * Mb_{ijk} + Replicate(Selection)_{ij} + e_{ijk}$ (eq. 2)

232   For the enzyme activities, the full model included the interaction between the thermal exposure and

233   thermal selection as fixed effects (eq. 3):

235                $Activity_{ijkl} = \mu + Selection_i * Exposure_j + Replicate(Selection)_{ik} + e_{ijkl}$ (eq. 3)

Early fecundity was analyzed using a generalized lineal model with a Poisson distribution with

thermal selection, oviposition days and their interaction were considered as fixed effects, while

replicated lines were nested within thermal selection were considered as a random effect (eq. 4). As

fecundity was measured for each female along 8 days, females' identity (ID) was included in the

model to accomplish for the repeated-measures design:

$\text{Fecundity}_{ijkl} = \mu + \text{Selection}_i * \text{Day}_j + \text{Replicate(Selection)}_{ik} + \text{ID}_l + e_{ijkl}$ (eq. 4)

Finally, egg-to-adult viability was estimated as the proportion of emerging flies per vial, considering

that 20 eggs were placed in a vial. This proportion was analyzed using a generalized lineal model

with a binomial distribution with thermal selection as fixed effect, and replicated lines were nested

within thermal selection were considered as a random effect (eq. 5):

$\text{Proportion}_{ijk} = \mu + \text{Selection}_i + \text{Replicate(Selection)}_{ij} + e_{ijk}$ (eq. 5)

All statistical analysis was performed in R (R Core Team, 2021) using the function *lmer* and *glmer* of

the *lme4* package for the generalized mixed model (Bates et al, 2014). Post-hoc analysis were

performed using a Tukey-adjusted method using the *emmeans* package (Lenth, 2021).

Results

We found that knockdown temperature was significantly higher in fast-ramping selected lines

compared to fast-ramping control lines ($\text{mean}_{\text{fast-ramping}} \pm \text{SD} = 37.71 \pm 0.68$ ºC and $\text{mean}_{\text{fast-control}} \pm$

$\text{SD} = 37.23 \pm 0.79$ ºC; $F_{1,4} = 32.0$, $P = 0.005$), and knockdown temperature in slow-ramping selected

lines was significantly higher than in slow-ramping control lines ($\text{mean}_{\text{slow-ramping}} \pm \text{SD} = 35.48 \pm 0.66$

ºC and $\text{mean}_{\text{slow-control}} \pm \text{SD} = 34.97 \pm 0.72$ ºC; $F_{1,4} = 41.7$, $P = 0.003$). The findings of the thermal

selection experiments are detailed in Mesas et al (2021).

RMR and body mass

Body mass was not significantly different between the control and selected lines ($\chi^2_2 = 0.46$, P =

0.79) and it did not differ between replicated lines ($\chi^2_1 = 0.03$, P = 0.86). On the other hand, as

expected, RMR and body mass were significantly correlated (r = 0.29, $t_{132} = 3.48$, P = 0.0007). Then,

body mass was included as covariate in the mixed linear model, which showed that RMR did not

differ between thermal selection regimens ($\chi^2_2 = 0.28$, P = 0.87; Fig. 1) and neither between the

replicated lines ($\chi^2_1 = 0$, P = 1).

Enzyme activity

For HEX activity (Fig. 2A), we found a significant effect of the thermal selection regimens ($\chi^2_2 =$

9.32, P = 0.009), a significant decrease of HEX activity after heat stress ($\chi^2_1 = 4.46$, P = 0.03), and a

non-significant interaction between thermal selection and thermal exposure ($\chi^2_2 = 2.28$, P = 0.32).

Looking for selection effects on each exposure treatment (Fig. 1A), we found that HEX basal

activity evolved to lower values in the slow-ramping selected lines (Tukey P-adjusted value = 0.02),

whereas fast-ramping selected lines showed similar HEX basal activity than control lines (Tukey P-

adjusted value = 0.35). On the other hand, HEX induced activity was not different between thermal

selection lines (Tukey P-adjusted value > 0.5). For PGI activity (Fig. 1B), we did not find significant

effects of thermal selection ($\chi^2_2 = 3.43$, P = 0.18), thermal exposure ($\chi^2_1 = 0.42$, P = 0.52), neither

significant interaction between these factors ($\chi^2_2 = 2.10$, P = 0.35). For PGM activity (Fig. 1C), we

did not find significant effect of the thermal selection regimens ($\chi^2_2 = 0.70$, P = 0.71), thermal

exposure ($\chi^2_1 = 1.42$, P = 0.23), neither significant interaction between these factors ($\chi^2_2 = 1.16$, P =

0.56). Finally, G6PD activity was significatively different between the thermal selection regimens (χ^2_2

= 6.7, P = 0.04; Fig. 1D). On the other hand, G6PD activity did not show significant differences

between thermal treatments (χ^2_1 = 1.35, P = 0.25) neither a significant interaction between thermal

selection and thermal exposure (χ^2_2 = 1.78, P = 0.41). Looking for selection effects on each

exposure treatment, probably because differences between thermal selection regimens were close to

significance, we did not find *a posteriori* significant differences between thermal selection regimens

within each thermal stress treatment (Tukey P-adjusted value ≥ 0.1).

Regarding to the variability present among replicated lines, we found that the replicated lines

showed a non-significant effect on the activity for all studied enzymes (χ^2_1 = 0, P = 1).

Early fecundity and egg-to-adult viability

Cumulative fecundity (Fig. 3) showed a significant effect of thermal selection (χ^2_2 = 17.83, P =

0.0001), oviposition day (χ^2_1 = 9248.33, P < 2.2 x 10^{-16}), and a significant interaction between both

factors (χ^2_2 =30.51, P = 2.4 x 10^{-7}). Specifically, the cumulative fecundity in the control lines was

lower than the slow-ramping selected lines (Tukey P-adjusted value = 0.02) and the fast-ramping

selected (Tukey P-adjusted value = 0.04). Whereas both selected lines showed similar cumulative

fecundity (Tukey P-adjusted value = 0.98). Looking for selection effects at each day (Fig. 3A), we

observed that females of the selected lines lay more eggs than females of the control lines since day

3 after mating. Additionally, cumulative fecundity was not different between replicated lines (χ^2_1 =

2.03, *P* = 0.15).

For egg-to-adult viability (Fig. A), we found significant effects of thermal selection (χ^2_2 = 16.24, *P* =

0.0003, Fig. 4). Specifically, slow-ramping selected lines showed a significantly higher egg-to-adult

viability than control lines (Tukey P-adjusted value = 0.0002), but not significantly different than

fast-ramping selected lines (Bonferroni p-adjusted value = 0.07). On the other hand, fast-ramping

selected lines showed a similar egg-to-adult viability compared to control lines (Bonferroni p-

adjusted value = 0.17). Finally, egg-to-adult viability were significantly different among replicated

lines (χ^2_4 = 3.18, P = 1.4 x 10^{-6}), but this among-replicates variability did not blur the effects of

thermal selection on the egg-to-adult viability.

Discussion

The mechanisms of tolerance to environmental stress are fundamental for the persistence of natural

populations and biological diversity. Temperature is a conspicuous environmental factor that has

influenced the evolution of metabolic and life-history traits in ectotherms, providing an interesting

system to study the evolution and constraints of stress tolerance (Gilloly et al, 2001 Addo-Bediako et

al, 2002; Brown et al, 2004; Padfield et al, 2016, Mallard et al, 2018). In this work, despite we did not

find evidence of changes of metabolic rate in response to heat tolerance evolution in *D. subobscura*,

we detected evolutionary correlated responses in a specific enzyme related to energy metabolism and

positive correlated responses of fitness-related traits. Interestingly, these correlated responses of

fitness-related traits to heat tolerance with no increase of the metabolic demand, which suggests that

low-cost phenotypes are positively selected during heat tolerance challenges. Previously, our research

team has been reported a correlated response of the locomotor performance to heat tolerance

evolution in *D. subobscura*, corroborating that this species can adapt to increasing environmental

temperatures (Mesas et al, 2021).

Despite the pervasive effects of temperature on metabolic traits, our findings show no evidence of a

correlated response of RMR to heat tolerance evolution in *D. subobscura*. Different studies along

geographical gradients have reported that organisms living in temperate environments have lower

metabolic rates than their counterparts inhabiting cold habitats (Addo-Bediako et al, 2002; Sylvestre

et al, 2007; Schaefer & Walters, 2010; Sinnatamby et al, 2015). In the same way, studies of selection

for stress resistance have found a reduction of metabolic rate in selected lines (Harshmann et al,

1999; Padfield et al, 2016). In addition, a study reported that populations of sticklebacks inhabiting

geothermally warmed lakes showed a lower RMR than populations from cold lakes, but that this

difference was only detected when both populations were acclimated to different temperatures

(Pilakouta et al, 2020). However, our results agree a study published by Messamah et al (2017), who

reported that the variation in the metabolic rate of 65 *Drosophila* species was not related to the annual

mean temperature of their habitats of origin. Similarly, some studies in *D. melanogaster* show no

differences in the metabolic rate between desiccation selected and control lines (Djawdan et al,

1997), and lines evolved in contrasting thermal condition (Alton et al, 2017). In this work, Alton et

al (2017) suggest that the evolution of metabolic rate could depend on the interaction between

temperature and resource availability, which would difficult the detection of metabolic response to

thermal variability under typical maintaining conditions. In this way, our control and evolved

populations were fed 'ad libitum' before measures, which could hide correlated responses of

metabolic rate to heat tolerance evolution. Additionally, metabolic rate was measured at non-

stressful conditions, which could have limited our capacity to detect differences between control and

evolved lines.

Another possible explanation for not finding a correlated response of RMR to heat tolerance

evolution is that metabolic rate is a complex trait that represents the total flux of energy at

organismic level and that its evolutionary response could be constrained by the evolution of

underlying traits such as metabolism-associated enzymes. For instance, Berrigan & Hoang (1999)

reported a positive correlation between SMR and the activity of enzymes associated with energy

metabolism across *Drosophila* genus. However, the moth *Amphion floridensis* showed that the oxygen

consumption was no affected by changes in activity of glycolytic and mitochondrial enzymes,

suggesting that changes in enzyme activities associated with energy metabolism were not enough to

modify the metabolic rate (Obrien & Suarez, 2001).

It is widely known that enzyme activity depends on the temperature: the lower the temperature, the

lower the enzyme activity (Sokolova & Pörtner, 2001). Indeed, a study in *D. simulans* reports a

reduction in the expression of genes related to glycolysis, Krebs cycle and oxidative phosphorylation

in populations evolving in thermally fluctuating environments (Mallard et al, 2018). Similarly, we

found that evolved lines showed a reduced activity in two out of four enzymes belong to the G6P

branchpoint, but this correlated response to heat tolerance evolution depended on how thermal

tolerance was selected: HEX evolved to lower activity in the slow-ramping selected lines, whereas

analysis of G6PD detected significant differences between thermal selection regimens, but we

cannot detect differences regarding to control lines. Interestingly, these correlated responses of

HEX were only observed when flies were exposed to non-stressful conditions. Additionally, these

same enzymes showed a reduced activity in flies exposed to 32 °C in comparison to those exposed

to non-stressful conditions (21 °C), reveling a plastic response of some G6P branchpoint enzymes to

temperature. These evolutionary and plastic responses of G6P branchpoint enzymes could be

associated with changes in transcription rate of the enzyme coding genes as well as changes of

thermostability and catalytic efficiency resulting from post-transcriptional modifications (Clark &

Wang, 1994). However, we did not performed qPCR or RNA-seq approaches, which would be very

useful to understand how heat tolerance evolution determine the specific or global patterns of gene

expression in ectotherms.

Several studies have investigated the stress resistance evolution and have found correlated responses

of metabolic enzymes of the G6P branchpoint and evolutionary responses in reproductive traits of

insects (Kageyama & Ohnishi, 1971; Clark & Fucito, 1998; Harshman et al, 1999; Mallard et al,

2018), suggesting a correlated response between energy metabolism and fitness. In our study, we

observed a reduced activity of HEX in the slow-ramping selected lines, suggesting that individuals

selected for high thermal tolerance also had lower metabolism (Santos et al, 2012). According to

pace-to-life syndrome, individuals with fast metabolism should grow faster than their counterparts

with slow metabolism, but at the cost to have shorter lifespan and lower fecundity (Stearns 1989;

Polverino et al, 2018). In agreement with this idea, our results show that the evolution of heat

tolerance is associated with a lower activity in the G6P branchpoint and higher fecundity. Our

findings also agree to those reported by Talbert et al (2015) who found that *D. melanogaster* flies that

express reduced HEX activity show longer life, whereas G6PD has been related to the reproduction

and development of insects and fishes (Downer et al, 1985; Zera & Zhao, 2003b; Tripathi & Verma,

2004). For example, seasonal changes in G6PD activity have been associated with the reproductive

cycle of catfish females, *Clarias batrachus* (Tripathi & Verma, 2004).

We also found a correlated response of the egg-to-adult viability to heat tolerance evolution, but

only in the slow-ramping selected lines. Likely, evolution of fitness-related traits is associated with

the pace-to-life syndrome as previously explained: the lower the activity of metabolic enzymes, the

higher the fitness. However, this explanation is not completely valid for the fast-ramping selected

lines, even though egg-to-adult viability is increased by 8% in these lines compared to control lines,

and with some vials reaching up to 100% viability. However, enzyme activities have been related to

development in *Drosophila* species: flies with short preadult development exhibit higher HEX and

G6PD activities than flies with long development times (Marinkovic et al, 1986). Similar results were

found in *D. simulans*, where populations evolved under different thermal regimens showed a low

expression of genes coding for enzymes related to energy metabolism and also high fecundity

(Mallard et al, 2018). Therefore, correlated responses of fitness-related traits suggest that

evolutionary responses heat tolerance to global warming should be adaptive. However, these

findings need to be evaluated under realistic conditions to clearly understand the ecological and

evolutionary implications of thermal evolution in natural populations.

In conclusion, we show that the evolution of tolerance to high temperatures does not involve a

metabolic depression neither an increase in energy demand. Furthermore, increased heat tolerance

evolution has positive consequences on fitness-related traits, including increased fecundity and

preadult survival. As we previously showed (see Mesas et al, 2021), heat tolerance evolution induces

correlated response on several traits, which finally result into an increased performance at high

temperatures. Despite some evidence shows that thermal limits have limited evolutionary potential

(Kellermann et al, 2012; Kelly et al, 2012), we have found through several works that upper thermal

limits have genetic variation (Castañeda et al, 2019), which allows to *D. subobscura* populations adapt

to local conditions (Castañeda et al, 2015) and respond to selection (Mesas et al, 2021). Thus, heat

tolerance can evolve under different thermal scenarios but with different outcomes on related traits

depending on the intensity of thermal stress, which change across spatial and temporal scales.

Therefore, spatial and temporal variability of thermal stress intensity should be taking into account

in future studies if we want to understand and predict the adaptive response to ongoing and future

climatic conditions.

References

Addo-Bediako, A., Chown, S. L., & Gaston, K. J. (2002). Metabolic cold adaptation in insects: a

large-scale perspective. Functional Ecology, 16(3), 332-338. https://doi.org/10.1046/j.1365-

2435.2002.00634.x

Alton, L. A., Condon, C., White, C. R., & Angilletta Jr, M. J. (2017). Colder environments did not
select for a faster metabolism during experimental evolution of *Drosophila melanogaster*. Evolution,
71(1), 145-152. https://doi.org/10.1111/evo.13094

Angilletta Jr, M. J., & Angilletta, M. J. (2009). Thermal adaptation: a theoretical and empirical
synthesis. Oxford, UK: Oxford University Press.

Artacho, P. & Nespolo, R.F. (2009). Natural selection reduces energy metabolism in the garden
snail, *Helix aspersa* (*Cornu Aspersum*). Evolution, 63: 1044-1050. https://doi.org/10.1111/j.1558-
5646.2008.00603.x

Atkinson D.E. (1977). Cellular Energy Metabolism and Its Regulation. Academic Press. New York.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using
lme4. arXiv preprint arXiv:1406.5823

Berrigan, D., & Hoang, A. (1999). Correlation between enzyme activities and routine metabolic rate
in *Drosophila*. Journal of Evolutionary Biology, 12(2), 258-262. https://doi.org/10.1046/j.1420-
9101.1999.00027.x

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a metabolic
theory of ecology. Ecology, 85(7), 1771-1789. https://doi.org/10.1890/03-9000

Brown, E. B., Slocumb, M. E., Szuperak, M., Kerbs, A., Gibbs, A. G., Kayser, M. S., & Keene, A. C.
(2019). Starvation resistance is associated with developmentally specified changes in sleep, feeding
and metabolic rate. Journal of Experimental Biology, 222(3), jeb191049.
https://doi.org/10.1242/jeb.191049

Bruning, A., González, A., Gaitán-Espitia, J. D., Bartheld, J. L., Toader-Williams, A., Mondaca, F., &

Nespolo, R. F. (2013). Energy metabolism, heart rate and physiological differentiation in the

pulmonate gastropod *Cornu aspersum*. Journal of Molluscan Studies, 79(3), 257-262.

https://doi.org/10.1093/mollus/eyt021

Castañeda, L. E., Rezende, E. L., & Santos, M. (2015). Heat tolerance in *Drosophila subobscura* along a

latitudinal gradient: Contrasting patterns between plastic and genetic responses. Evolution, 69, 2721-

2734. https://doi.org/10.1111/evo.12757

Castañeda, L.E., Romero-Soriano, V., Mesas, A., Roff, D. A., & Santos, M. (2019). Evolutionary

potential of thermal preference and heat tolerance in *Drosophila subobscura*. Journal of evolutionary

biology, 32(8), 818-824. https://doi.org/10.1111/jeb.13483

Chen, I. C., Hill, J. K., Ohlemüller, R., Roy, D. B., & Thomas, C. D. (2011). Rapid range shifts of

species associated with high levels of climate warming. Science, 333(6045), 1024-1026.

https://doi.org/doi/10.1126/science.1206432

Chick, L. D., Waters, J. S., & Diamond, S. E. (2021). Pedal to the metal: Cities power evolutionary

divergence by accelerating metabolic rate and locomotor performance. Evolutionary applications,

14(1), 36-52. https://doi.org/10.1111/eva.13083

Clark, A. G., & Fucito, C. D. (1998). Stress tolerance and metabolic response to stress in *Drosophila*

melanogaster. Heredity, 81(5), 514-527. https://doi.org/10.1046/j.1365-2540.1998.00414.x

Clark, A. G., & Wang, L. (1994). Comparative evolutionary analysis of metabolism in nine *Drosophila*

species. Evolution, 48(4), 1230-1243. https://doi.org/10.1111/j.1558-5646.1994.tb05308.x

Crozier, L. G., & Hutchings, J. A. (2014). Plastic and evolutionary responses to climate change in

fish. Evolutionary Applications, 7(1), 68-87. https://doi.org/10.1111/eva.12135

Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., &

Martin, P. R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude.

Proceedings of the National Academy of Sciences, 105(18), 6668-6672.

https://doi.org/10.1073/pnas.0709472105

Djawdan, M., Rose, M.R., & Bradley, T.J. (1997). Does selection for stress resistance lower

metabolic rate?. Ecology, 78, 828-837. https://doi.org/10.1890/0012-

9658(1997)078[0828:DSFSRL]2.0.CO;2

Downer, R.G.H. (1985). Lipid metabolism. En Kerkut, G.A. and L.I. Gilbert. (eds). Comprehensive

insect physiology, biochemistry and pharmacology. Pergamon, Oxford, U.K

Duvernell, D. D., & Eanes, W. F. (2000). Contrasting molecular population genetics of four

hexokinases in *Drosophila melanogaster, D. simulans* and *D. yakuba*. Genetics, 156(3), 1191-1201.

https://doi.org/10.1093/genetics/156.3.1191

Eanes, W. F., & Hey, J. (1986). In vivo function of rare G6pd variants from natural populations of

Drosophila melanogaster. Genetics, 113(3), 679-693. https://doi.org/10.1093/genetics/113.3.679

Gaston, K.J., Chown, S.L., Calosi, P., Bernardo, J., Bilton, D.T., Clarke, A., Clusella-Trullas, S.,

Ghalambor, C.K., Konarzewski, M., Peck, L.S., Porter, W.P., Pörtner, H.O., Rezende, E.L., Schulte,

P.M., Spicer, J.I., Stillman, J.H., Terblanche, J.S., & van Kleunen, M. (2009). Macrophysiology: a

conceptual reunification. American Naturalist,174(5), 595-612. https://doi.org/10.1086/605982

Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M., & Charnov, E. L. (2001). Effects of size and

temperature on metabolic rate. science, 293(5538), 2248-2251.

https://doi.org/10.1126/science.1061967

González-Tokman, D., Córdoba-Aguilar, A., Dáttilo, W., Lira-Noriega, A., Sánchez-Guillén, R. A.,

& Villalobos, F. (2020). Insect responses to heat: physiological mechanisms, evolution and ecological

implications in a warming world. Biological Reviews, 95(3), 802-821.

https://doi.org/10.1111/brv.12588

Guppy, M., & Withers, P. (1999). Metabolic depression in animals: physiological perspectives and

biochemical generalizations. Biological Reviews, 74(1), 1-40.

https://doi.org/10.1017/S0006323198005258

Hamby, K. A., Bellamy, D. E., Chiu, J. C., Lee, J. C., Walton, V. M., Wiman, N. G., ... & Biondi, A.

(2016). Biotic and abiotic factors impacting development, behavior, phenology, and reproductive

biology of *Drosophila suzukii*. Journal of pest science, 89(3), 605-619.

https://doi.org/10.1007/s10340-016-0756-5

Harshman, L. G., Hoffmann, A. A., & Clark, A. G. (1999). Selection for starvation resistance in

Drosophila melanogaster: physiological correlates, enzyme activities and multiple stress responses.

Journal of Evolutionary Biology, 12(2), 370-379. https://doi.org/10.1046/j.1420-9101.1999.00024.x

Harshman, L. G., & Zera, A. J. (2007). The cost of reproduction: the devil in the details. Trends in

ecology & evolution, 22(2), 80-86. https://doi.org/10.1016/j.tree.2006.10.008

Hochachka, P. W., & Somero, G. N. (2002). Biochemical adaptation: mechanism and process in

physiological evolution. Oxford university press.

Hoffmann, A. A., & Parsons, P. A. (1989). Selection for increased desiccation resistance in *Drosophila*

melanogaster: additive genetic control and correlated responses for other stresses. Genetics, 122(4),

837-845. https://doi.org/10.1093/genetics/122.4.837

Huey, R. B., & Kingsolver, J. G. (1993). Evolution of resistance to high temperature in ectotherms.

The American Naturalist,142, S21-S46. https://doi.org/10.1086/285521

Huey, R. B., & Kingsolver, J. G. (2019). Climate warming, resource availability, and the metabolic

meltdown of ectotherms. The American Naturalist, 194(6), E140-E150.

https://doi.org/10.1086/705679

Kageyama, T., & Ohnishi, E. (1971). Carbohydrate metabolism in the eggs of the silkworm, *Bombyx*

mori. I. Absence of phosphofructokinase in the diapausing egg. Development, Growth &

Differentiation, 13(2), 97-106. https://doi.org/10.1111/j.1440-169X.1971.00097.x

Kellermann, V., Loeschcke, V., Hoffmann, A.A., Kristensen, T.N., Fløjgaard, C., David, J.R.,

Svenning, J.C., & Overgaard, J. (2012). Phylogenetic constraints in key functional traits behind

species' climate niches: patterns of desiccation and cold resistance across 95 *Drosophila* species.

Evolution, 66, 3377-3389. https://doi.org/10.1111/j.1558-5646.2012.01685.x

Kelly, M. W., Sanford, E., & Grosberg, R.K. (2012). Limited potential for adaptation to climate

change in a broadly distributed marine crustacean. Proceedings of the Royal Society B: Biological

Sciences, 279(1727), 349-356. https://doi.org/10.1098/rspb.2011.0542

Lenth, R. (2021). Estimated Marginal Means, aka Least-Squares Means. R Package version 1.3.4.

https://cran.r-project.org/web/packages/emmeans/index.

Lighton, J. R., & Turner, R. J. (2004). Thermolimit respirometry: an objective assessment of critical

thermal maxima in two sympatric desert harvester ants, *Pogonomyrmex rugosus* and *P. californicus*.

Journal of Experimental Biology, 207(11), 1903-1913. https://doi.org/10.1242/jeb.00970

Mallard, F., Nolte, V., Tobler, R., Kapun, M., & Schlötterer, C. (2018). A simple genetic basis of

adaptation to a novel thermal environment result in complex metabolic rewiring in *Drosophila*.

Genome biology, 19(1), 1-15. https://doi.org/10.1186/s13059-018-1503-4

Marinkovié, D., Milošević, M., & Milanović, M. (1986). Enzyme activity and dynamics of *Drosophila*

development. Genetica, 70(1), 43-52. https://doi.org/10.1007/BF00123213

Marshall, D. J., & McQuaid, C. D. (2011). Warming reduces metabolic rate in marine snails:

adaptation to fluctuating high temperatures challenges the metabolic theory of ecology. Proceedings

of the Royal Society B: Biological Sciences, 278(1703), 281-288.

https://doi.org/10.1098/rspb.2010.1414

Matoo, O.B., Julick, C. R., & Montooth, K. L. (2019). Genetic Variation for Ontogenetic Shifts in

Metabolism Underlies Physiological Homeostasis in *Drosophila*. Genetics, 212(2), 537-552.

https://doi.org/10.1534/genetics.119.302052

McGaughran, A., Laver, R., & Fraser, C. (2021). Evolutionary responses to warming. Trends in

Ecology & Evolution. 36, 591-600. https://doi.org/10.1016/j.tree.2021.02.014

Mesas, A., Jaramillo, A., & Castañeda, L. E. (2021). Experimental evolution on heat tolerance and

thermal performance curves under contrasting thermal selection in *Drosophila subobscura*. Journal of

Evolutionary Biology, 34(5), 767-778. https://doi.org/10.1111/jeb.13777

Messamah, B., Kellermann, V., Malte, H., Loeschcke, V., & Overgaard, J. (2017). Metabolic cold

adaptation contributes little to the interspecific variation in metabolic rates of 65 species of

Drosophilidae. Journal of Insect Physiology, 98, 309-316.

https://doi.org/10.1016/j.jinsphys.2017.02.003

Montooth, K.L., Marden, J.H. & Clark, A.G. (2003). Mapping determinants of variation in energy

metabolism, respiration and flight in *Drosophila*. Genetics, 165(2), 623-635.

https://doi.org/10.1093/genetics/165.2.623

Morrison, C., & Hero, J. M. (2003). Geographic variation in life-history characteristics of

amphibians: a review. Journal of Animal Ecology, 72(2), 270-279. https://doi.org/10.1046/j.1365-

2656.2003.00696.x

Oakeshott, J. G., Chambers, G. K., Gibson, J. B., Eanes, W. F., & Willcocks, D. A. (1983).

Geographic variation in G-6-pd and Pgd allele frequencies in *Drosophila melanogaster*. Heredity, 50(1),

67-72.

O'Brien, D. M., & Suarez, R. K. (2001). Fuel use in hawkmoth (*Amphion floridensis*) flight muscle:

enzyme activities and flux rates. Journal of Experimental Zoology, 290(2), 108-114.

https://doi.org/10.1002/jez.1040

Padfield, D., Yvon-Durocher, G., Buckling, A., Jennings, S., & Yvon-Durocher, G. (2016). Rapid

evolution of metabolic traits explains thermal adaptation in phytoplankton. Ecology letters, 19(2),

133-142. https://doi.org/10.1111/ele.12545

Pilakouta, N., Killen, S. S., Kristjánsson, B. K., Skúlason, S., Lindström, J., Metcalfe, N. B., &

Parsons, K. J. (2020). Multigenerational exposure to elevated temperatures leads to a reduction in

standard metabolic rate in the wild. Functional ecology, 34(6), 1205-1214.

https://doi.org/10.1111/1365-2435.13538

Polverino, G., Santostefano, F., Díaz-Gil, C., & Mehner, T. (2018). Ecological conditions drive pace-

of-life syndromes by shaping relationships between life history, physiology and behaviour in two

populations of Eastern mosquitofish. Scientific reports, 8(1), 14673.

https://doi.org/10.1038/s41598-018-33047-0

Porcelli, D., Gaston, K. J., Butlin, R. K., & Snook, R. R. (2017). Local adaptation of reproductive

performance during thermal stress. Journal of Evolutionary Biology, 30(2), 422-429.

https://doi.org/10.1111/jeb.13018

Pörtner, H. O., Bennett, A. F., Bozinovic, F., Clarke, A., Lardies, M. A., Lucassen, M., Pelster, B.,

Schiemer, F., & Stillman, J. H. (2006). Trade-offs in thermal adaptation: the need for a molecular to

ecological integration. Physiological and Biochemical Zoology, 79(2), 295-313.

https://doi.org/10.1086/499986

Qu, Y. F., Li, H., Gao, J. F., & Ji, X. (2011). Geographical variation in reproductive traits and trade-

offs between size and number of eggs in the king ratsnake, *Elaphe carinata*. Biological Journal of the

Linnean Society, 104(3), 701-709. https://doi.org/10.1111/j.1095-8312.2011.01749.x

R Core Team. (2021). R: A language and environment for statistical computing (version 4.0.2). R

Foundation for Statistical Computing, Vienna, Austria. Retrieved from https://www.R-project.org

Rank, N. E., & Dahlhoff, E. P. (2002). Allele frequency shifts in response to climate change and

physiological consequences of allozyme variation in a montane insect. Evolution, 56(11), 2278-2289.

https://doi.org/10.1111/j.0014-3820.2002.tb00151.x

Regan, M. D., Gill, I. S., & Richards, J. G. (2017). Metabolic depression and the evolution of

hypoxia tolerance in threespine stickleback, *Gasterosteus aculeatus*. Biology letters, 13(11), 20170392.

https://doi.org/10.1098/rsbl.2017.0392

Rezende, E. L., Tejedo, M., & Santos, M. (2011). Estimating the adaptive potential of critical thermal

limits: methodological problems and evolutionary implications. Functional Ecology, 25(1), 111-121.

https://doi.org/10.1111/j.1365-2435.2010.01778.x

Santos, M., Castañeda, L. E., & Rezende, E. L. (2012). Keeping pace with climate change: what is

wrong with the evolutionary potential of upper thermal limits?. Ecology and evolution, 2(11), 2866-

2880. https://doi.org/10.1002/ece3.385

Schaefer, J., & Walters, A. (2010). Metabolic cold adaptation and developmental plasticity in

metabolic rates among species in the *Fundulus notatus* species complex. Functional Ecology, 24(5),

1087-1094. https://doi.org/10.1111/j.1365-2435.2010.01726.x

Sezgin, E., Duvernell, D. D., Matzkin, L. M., Duan, Y., Zhu, C. T., Verrelli, B. C., & Eanes, W. F.

(2004). Single-locus latitudinal clines and their relationship to temperate adaptation in metabolic

genes and derived alleles in *Drosophila melanogaster*. Genetics, 168(2), 923-931.

https://doi.org/10.1534/genetics.104.027649

Sinnatamby, R.N, Brian Dempson, J., Reist, J. D., & Power, M. (2015). Latitudinal variation in

growth and otolith-inferred field metabolic rates of Canadian young-of-the-year Arctic charr.

Ecology of Freshwater Fish, 24(3), 478-488. https://doi.org/10.1111/eff.12166

Sokolova, I., & Pörtner, H. O. (2001). Temperature effects on key metabolic enzymes in *Littorina*

saxatilis and *L. obtusata* from different latitudes and shore levels. Marine Biology, 139(1), 113-126.

https://doi.org/10.1007/s002270100557

Stearns, S.C. (1989). Trade-Offs in Life-History Evolution. Functional Ecology, 3(3), 259-268.

https://doi.org/10.2307/2389364

Storey, K. B., & Storey, J. M. (2004). Metabolic rate depression in animals: transcriptional and

translational controls. Biological Reviews, 79(1), 207-233. http:// doi.org/10.1007/s00360-007-

0143-x

Sylvestre, E. L., Lapointe, D., Dutil, J. D., & Guderley, H. (2007). Thermal sensitivity of metabolic

rates and swimming performance in two latitudinally separated populations of cod, *Gadus morhua* L.

Journal of Comparative Physiology B, 177(4), 447-460. https://doi.org/10.1007/s00360-007-0143-

x.pdf

Talbert, M. E., Barnett, B., Hoff, R., Amella, M., Kuczynski, K., Lavington, E., Koury, S., Brud, E.,

& Eanes, W. F. (2015). Genetic perturbation of key central metabolic genes extends lifespan in

Drosophila and affects response to dietary restriction. Proceedings. Biological sciences, 282(1815),

20151646. https://doi.org/10.1098/rspb.2015.1646

Tripathi, G., & Verma, P. (2004). Sex-specific metabolic changes in the annual reproductive cycle of

a freshwater catfish. Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular

Biology, 137(1), 101-106. https://doi.org/10.1016/j.cbpc.2003.10.005

Tomanek, L., & Zuzow, M. J. (2010). The proteomic response of the mussel congeners *Mytilus*

galloprovincialis and *M. trossulus* to acute heat stress: implications for thermal tolerance limits and

metabolic costs of thermal stress. Journal of Experimental Biology, 213(20), 3559-3574.

https://doi.org/10.1242/jeb.041228

Umina, P. A., Weeks, A. R., Kearney, M. R., McKechnie, S. W., & Hoffmann, A. A. (2005). A rapid

shift in a classic clinal pattern in *Drosophila* reflecting climate change. Science,308 (5722), 691-693.

https:/doi.org/ 10.1126/science.1109523

Verrelli, B. C., & Eanes, W. F. (2001). The functional impact of Pgm amino acid polymorphism on

glycogen content in *Drosophila melanogaster*. Genetics, 159(1), 201-210.

https://doi.org/10.1093/genetics/159.1.201

Zera, A. J. (1987). Temperature-dependent kinetic variation among phosphoglucose isomerase

allozymes from the wing-polymorphic water strider, *Limnoporus canaliculatus*. Molecular biology and

evolution, 4(3), 266-285. https://doi.org/10.1093/oxfordjournals.molbev.a040445

Zera, A. J., & Zhao, Z. (2003a). Morph-dependent fatty acid oxidation in a wing-polymorphic

cricket: implications for the trade-off between dispersal and reproduction. Journal of Insect

Physiology, 49(10), 933-943. https://doi.org/10.1016/S0022-1910(03)00152-5

Zera, A. J., & Zhao, Z. (2003b). Life-history evolution and the microevolution of intermediary

metabolism: Activities of lipid-metabolizing enzymes in life-history morphs of a wing-dimorphic

cricket. Evolution, 57(3), 586-596. https://doi.org/10.1111/j.0014-3820.2003.tb01550.x

 Figure legends

 Figure 1. Relationship between body mass and metabolic rate (central panel) for flies of control lines

 (black circles), slow-ramping selected lines (red circles), and fast-ramping selected lines (blue circles)

 for increasing heat tolerance in *Drosophila subobscura*. The grey line represents the linear regression

 between both traits. Distribution of body mass and metabolic rate are showed in the right and upper

 panels, respectively, including the mean of each thermal selection regimen (black line). Non-

 significant different between thermal selection regimens were found for both traits (see text).

 Figure 2. Activity of enzymes related to G6P branchpoint of flies of control lines (black circles),

 slow-ramping selected lines (red circles), and fast-ramping selected lines (blue circles) for increasing

 heat tolerance in *Drosophila subobscura* exposed to non-stressful (21 °C) and stressful conditions (32

 °C): (A) hexokinase (HEX); (B) phophoglucomutase (PGM); (C) phosphoglucoisomerase (PGI); and

 (D) glucose-6-phosphate dehydrogenase (G6PD). Horizontal lines show the mean for each group.

 P-values above whiskers show the results of *a posteriori* comparisons between control and selected

 lines using Tukey tests.

 Figure 3. Cumulative fecundity along first 8 days after mating for females of control lines (black

 circles and boxplots), slow-ramping selected lines (red circles and boxplots), and fast-ramping

 selected lines (blue circles and boxplots) for increasing heat tolerance in *Drosophila subobscura*.

 Boxplots show median and the interquartile range (IRQ) and vertical whiskers represent the

1.5*IQR. P-values above horizontal whiskers show the results of *a posteriori* comparisons between

control and selected lines using Tukey tests (ns: P > 0.05, *: P < 0.05, **: P < 0.01, ***: P < 0.001).

Figure 4. Egg-to-adult viability of flies of control lines (black circles), slow-ramping selected lines

(red circles), and fast-ramping selected lines (blue circles) for increasing heat tolerance in *Drosophila*

subobscura. Horizontal lines show the mean for each group. P-values above horizontal whiskers show

the results of *a posteriori* comparisons between control and selected lines using Tukey tests.