VE607 Stabilizes SARS-CoV-2 Spike In the "RBD-up" Conformation and Inhibits Viral

Entry

Shilei Ding[1], Shang Yu Gong[1,2], Jonathan Grover[3], Mohammadjavad Mohammadi[4], Yaozong

Chen[5], Dani Vézina[1,6], Guillaume Beaudoin-Bussières[1,6], Vijay Tailor Verma[7], Guillaume

Goyette[1], Jonathan Richard[1,6], Derek Yang[8], Amos B. Smith III[8], Marzena Pazgier[5],

Marceline Côté[9], Cameron Abrams[4], Walther Mothes[3], Andrés Finzi[1,2,6,10*] and Christian

Baron[7*]

[1]Centre de recherche du CHUM, Montréal, QC, Canada
[2]Department of Microbiology and Immunology, McGill University, Montreal, QC, Canada
[3]Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven, CT 06510,
USA
[4]Department of Biochemistry and Molecular Biology, Drexel University College of Medicine,
Philadelphia, PA 19104, USA.
[5]Infectious Disease Division, Department of Medicine, Uniformed Services University of the Health
Sciences, Bethesda, MD 20814-4712, USA
[6]Département de Microbiologie, Infectiologie et Immunologie, Université de Montréal, Montréal, QC,
Canada
[7]Department of Biochemistry and Molecular Medicine, Université de Montréal, Montréal, QC, Canada
[8]Department of Chemistry, School of Arts and Sciences, University of Pennsylvania, Philadelphia, PA,
USA
[9]Department of Biochemistry, Microbiology and Immunology, and Center for Infection, Immunity,
and Inflammation, University of Ottawa, Ottawa, ON K1H 8M5, Canada
25
26 [10]Lead Contact
27

* Correspondence: andres.finzi@umontreal.ca ; christian.baron@umontreal.ca

Running Title: Small molecule inhibitors of SARS-CoV-2 entry

Keywords: VE607, COVID-19, SARS-CoV-2, SARS-CoV-1, spike glycoprotein, small

molecule inhibitor, docking, single-molecule FRET

Summary

SARS-CoV-2 infection of host cells starts by binding of the Spike glycoprotein (S) to the

ACE2 receptor. The S-ACE2 interaction is a potential target for therapies against COVID-19

as demonstrated by the development of immunotherapies blocking this interaction. Here, we

present the commercially available VE607, comprised of three stereoisomers, that was

originally described as an inhibitor of SARS-CoV-1. We show that VE607 specifically

inhibits infection of SARS-CoV-1 and SARS-CoV-2 S-expressing pseudoviral particles as

well as authentic SARS-CoV-2. VE607 stabilizes the receptor binding domain (RBD) in its

"up" conformation. *In silico* docking and mutational analysis map the VE607 binding site at

the RBD-ACE2 interface. The IC_{50} values are in the low micromolar range for

pseudoparticles derived from SARS-CoV-2 Wuhan/D614G as well as from variants of

concern (Alpha, Beta, Gamma, Delta and Omicron), suggesting that VE607 has potential for

the development of drugs against SARS-CoV-2 infections.

Introduction

The COVID-19 pandemic continues to cause widespread morbidity and mortality (Wu et al., 2020; Zhu et al., 2020a). This is largely due to insufficient vaccination levels as vaccines offer good protection against infection and severe disease (Ball, 2021). The currently used vaccines exploit modified versions of the Spike (S) glycoprotein that is exposed on the surface of viral particles (Krammer, 2020) and infected cells (Ding et al., 2022). S is processed by cellular proteases furin and TMPRSS2 on host cells. After binding to ACE2 via its receptor binding domain (RBD), S undergoes significant conformational changes that ultimately lead to fusion of the viral membrane with human cells. Fusion allows translocation of the RNA genome and associated replicase proteins into mammalian cells, leading to viral replication (Harrison et al., 2020; Hoffmann et al., 2020a; Hoffmann et al., 2020b; Yang and Rao, 2021). S is a trimeric glycoprotein that is present in multiple conformations that have been resolved primarily by cryo-electron microscopy (Cai et al., 2020; Lan et al., 2020; Shang et al., 2020; Wrapp et al., 2020; Yan et al., 2020). Its conformational dynamics can be monitored by single molecule FRET (Li et al., 2021b; Lu et al., 2020; Ullah et al., 2021; Yang et al., 2021). Vaccine-elicited antibodies act in several ways including neutralizing viral particles, but also through Fc-mediated effector functions (Tauzin et al., 2022; Tauzin et al., 2021). The selective pressure during the pandemic has led to a growing list of variants carrying mutations in the S-glycoprotein (Gong et al., 2021a; Li et al., 2021a; Mannar et al., 2021; Nabel et al., 2021; Prevost and Finzi, 2021; Yang and Rao, 2021) resulting in different degrees of resistance to previous infection and vaccine-elicited antibody neutralization.

Despite the efficacy of currently used vaccines and ongoing work to generate broadly protective pan-coronavirus vaccines (Cohen, 2021; Nabel et al., 2021; Rappazzo et al., 2021),

there is an urgent need for efficient and specific treatments for infected patients. The viral

replication machinery offers different possible drug targets (Yang and Rao, 2021) and several

small molecule inhibitors targeting the SARS-CoV-2 protease (Dai et al., 2020; Zhang et al.,

2020) or replicase (Kokic et al., 2021; Yin et al., 2021) have been published with some

recently showing promise in clinical trials (Owen et al., 2021). In contrast, relatively little

attention has been given to the S-ACE2 interaction as a potential target for small molecule

inhibitors (Tong, 2009; Wang et al., 2021; Zhu et al., 2020b). Research on SARS-CoV-1 and

Middle East respiratory syndrome (MERS) has inspired work on potential drug targets, and

some previous studies explored the isolation of small molecule inhibitors against various

potential targets. Some of these molecules were described as potential inhibitors of the SARS-

CoV-1 RBD interaction with ACE2 (Adedeji et al., 2013; Kao et al., 2004), but the binding

was not demonstrated directly and there was no biological follow-up work to characterize

their mode of action.

Here we employed differential scanning fluorimetry (DSF) to identify the capacity of

the small molecule inhibitor VE607 (Kao et al., 2004), composed of three stereoisomers, to

bind the SARS-CoV-2 RBD. We found that this VE607 mixture of isomers (hereafter referred

to as "VE607") is capable of specific inhibition of infection of human cells with pseudoviral

particles that express the SARS-CoV-1 or SARS-CoV-2 S-glycoproteins. VE607 was also

able to inhibit the infection with authentic SARS-CoV-2 viruses. We found that VE607

inhibits the Spike by stabilizing the "up" conformation of the RBD. The mode of binding to

RBD was elucidated by *in silico* docking experiments followed by validation of critical

residues through mutagenesis and functional studies. Finally, VE607 remains potent against

current variants of concern (VOC) of SARS-CoV-2 suggesting that it may be an interesting

lead for the development of drugs for the prevention or treatment of COVID-19 infections.

Results

Differential scanning fluorimetry and docking suggest that VE607 may bind the RBD

We tested the ability of previously described SARS-CoV-1 inhibitors VE607 (Kao et al., 2004) and SSAA09E2 (Adedeji et al., 2013) to bind the SARS-CoV-2 RBD (Figure 1A). We used differential scanning fluorimetry (DSF) that, measures the effect of small molecules on the melting temperature of proteins (Mashalidis et al., 2013). Incubation with VE607 led to a significant decrease of the melting temperature (ΔT_m, -2.3°C) while SSAA09E2 had a smaller, yet measurable effect (ΔT_m, -0.7°C) (Figure 1B). Since this result suggested binding of VE607 to RBD, we next performed *in silico* docking against RBD using Glide (Schrödinger, 2020). We identified moderately favorable potential VE607 binding sites overlapping the ACE2 epitopes in both SARS-CoV-1 and SARS-CoV-2 RBDs (Figure 1C and D).

VE607 inhibits infection of pseudoviral particles and authentic SARS-CoV-2

To assess the effects of VE607 and SSAA09E2 on infection we expressed the SARS-CoV-1 and SARS-CoV-2 S glycoproteins on the surface of pseudoviral particles carrying a luciferase reporter gene. Pseudoviral particles carrying the VSV-G glycoprotein served as control. Infection was measured using ACE2-expressing 293T (293T-ACE2) cells (Prevost et al., 2020) in the presence of increasing concentrations of VE607 and SSAA09E2. VE607 specifically inhibited pseudoviral particles bearing the SARS-CoV-1 Spike (IC_{50} = 1.47 μM, Figure 2A), in agreement with previous findings (Kao et al., 2004). Interestingly, VE607 also

inhibited pseudoviral particles expressing the Spike from SARS-CoV-2 (IC_{50} = 3.06 μM,

Figure 2A), albeit slightly less efficiently than for SARS-CoV-1. No inhibition was observed

for pseudoviral particles bearing the VSV-G (IC_{50} > 100μM, Figure 2A). To ensure that the

inhibitory capacity of VE607 against SARS-CoV-2 was not limited to pseudoviral particles,

we evaluated whether the inhibitory capacity of VE607 was maintained against authentic

viruses. As shown in Figure 2B, VE607 inhibited authentic SARS-CoV-2 D614G with an

IC_{50} of 2.42 μM. No cell toxicity of VE607 or the three different enantiomers was observed

with concentrations up to 100 μM on 293T ACE2 cells or Vero-E6 cells (Figure 2C). In

contrast, SSAA09E2 at concentrations up to 100 μM did not inhibit infection of pseudoviral

particles (data not shown) and we did not further pursue work with this small molecule.

As stated above, commercially available VE607 is a mixture of three stereochemical isomers,

comprised of the (S,S)-VE607, (R,R)-VE607, and the meso (R,S)-VE607. We observed no

differences in the SARS-CoV-2 pseudoviral inhibition among these enantiomers obtained by

synthesis, and the commercially available mixture of all three isomers (Figures S1).

Initial *in silico* docking identified RBD residues Y505 and Q498 as potential specific

contact sites for VE607 (Figure 1D). We mutated these residues in the full-length SARS-

CoV-2 D614G Spike and prepared pseudoviral particles to test whether they affect VE607

inhibition. While the Q498V mutation had only a minor effect (IC_{50} = 1.80 μM), the Y505T

mutant was resistant to VE607 inhibition (IC_{50} > 40 μM, Figure 2D). These results are in

agreement with the *in silico* analysis, where a strong π-π interaction between Y505's aromatic

side-chain and the central aromatic ring of VE607 is predicted. Alignment of sACE2 on the

known ACE2 epitope of the VE607-bound model of RBD displayed significant steric clashes

between ACE2 and VE607, suggesting some direct competition for the ACE2 epitope.

VE607 stabilizes the "up" conformation of the S protein

We next assessed whether VE607 affects the RBD-ACE2 interaction. Briefly, we measured

by flow cytometry the capacity of VE607 to compete with soluble ACE2 (sACE2) for

interaction with the full SARS-CoV-2 Spike, expressed at the cell surface, as described

previously (Anand et al., 2020). We observed no competition between VE607 (100 μM) and

sACE2 (Figure 3A). Since the mode of action of some neutralizing antibodies such as CV3-1

involve S1 shedding (Li et al., 2022), we tested if VE607 acted in a similar manner. To this

effect, Spike expressing 293T cells were radioactively labeled followed by the

immunoprecipitation of cell lysates and supernatant, as described (Li et al., 2022). In contrast

to CV3-1, VE607 decreased shedding, resulting in a significantly more stable trimer (Figure

3B).

To evaluate whether the enhanced stability observed in presence of VE607 altered the

conformational landscape of the Spike, we performed single-molecule FRET (smFRET)

analysis using viral pseudoparticles carrying modified S glycoproteins labelled with FRET

donor and acceptor dyes enabling us to distinguish the "up" and "down" conformations (Lu et

al., 2020) (Li et al., 2022). In agreement with previous observations, the unliganded Spike

predominantly sampled the "down" conformation (Figure 3C). As expected, the addition of

sACE2 shifted the conformational landscape of the Spike to the "up" conformation reflecting

the receptor bound state (Figure 3D). Interestingly, we observed that VE607 stabilized the

"up" conformation mimicking sACE2 (Figure 3 E).

VE607 inhibits infection of some SARS-CoV-2 variants of concern

SARS-CoV-2 is in constant evolution as VOCs keep emerging. VE607 was identified as an

inhibitor of SARS-CoV-1, a related Beta-coronavirus, suggesting some inhibitory breadth.

Therefore, we tested whether it inhibits pseudoviral particles bearing the Spike glycoproteins

from the major VOCs (Alpha, Beta, Gamma, Delta and Omicron). In agreement with its broad

SARS coronavirus activity, VE607 inhibited all VOCs with similar potency with IC_{50} values

in the low micromolar range (Figure 4). These results demonstrate that the various amino acid

changes in the S-glycoprotein of these variants do not impact the inhibitory potential of

VE607 and show promise for the development a new generation of anti-SARS-CoV-2 small

molecule inhibitors blocking viral entry.

Discussion

Here we present data suggesting that small molecule inhibitors of SARS-CoV-2 entry, such as VE607, have potential for the development of drugs for the prevention and/or treatment of COVID-19. VE607 was originally described as an inhibitor of the S glycoprotein-ACE2 mediated SARS-CoV-1 entry (Kao et al., 2004) and we confirmed these results using a pseudovirus infection assay. The inhibition of SARS-CoV-1 carrying pseudoviruses was stronger than that of SARS-CoV-2 pseudoviruses, but the IC_{50} values remained in the low μM range in both cases. *In silico* docking identified a potential binding site at the RBD-ACE2 interface in both cases (Lan et al., 2020; Shang et al., 2020). The results of mutational analysis are consistent with the predicted binding site suggesting a critical role of Y505 for the activity of VE607. Therefore, VE607 may inhibit viral entry by blocking ACE2-mediated Spike conformational changes required for fusion.

Single molecule FRET data indicate that VE607 alone stabilized the Spike in the preferred "up" conformation for sACE2 binding. However, in contrast to RBD-targeting antibodies such as CV3-1 (Li et al., 2022), VE607 did not induce shedding of S1. These results suggest a unique mechanistic basis for the inhibition of infection by VE607. We hypothesize that VE607 stabilizes one ACE2-bound conformation of Spike but is an allosteric inhibitor of downstream ACE2-triggered conformational changes required for fusion. The binding site predicted by *in silico* docking was confirmed by mutational analysis suggesting that VE607 binds at the S-ACE2 interface where it may block the protein-protein interaction required to activate the Spike for fusion. The emergence of variants over the course of the pandemic is a continuing concern and many of them carry mutations in the S glycoprotein including the RBD domain that contribute to increased infectivity and immune escape. To validate the

binding site predicted by *in silico* docking we introduced the mutations Q498V and Y505T

and whereas the first change had only a modest effect, the change Y505T led to an S

glycoprotein that is resistant to VE607.

These results are consistent with the fact that pseudoviruses carrying S from the variants

Alpha (B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (B.167.2) and Omicron (B.1.1.529) are

inhibited by VE607 at low μM IC_{50} values, suggesting that common mutations in the vicinity

of the binding site such as N501Y do not impact the inhibitory potential of VE607. Changes

of Q498 are rarely observed in the Beta variant (less than 0.1% of all sequences), but the

mutation Q498R is very frequent in Omicron (90% of all sequences) which is rapidly

spreading and leading to the next phase of the pandemic (https://outbreak.info/compare-

lineages). Interestingly, the Y505H mutation is also frequent in Omicron (90% of all

sequences), but it is rarely present in other variants such as Alpha (less than 0.1% of all

sequences). Nevertheless, the inhibitory effect of VE607 on pseudoviral particles carrying

Spike from Omicron was comparable to other VOCs. It is presently unclear why the presence

of Y505H does not affect the inhibitory activity of VE607 against Omicron when the Y505T

change renders pseudoviral particles resistant to the molecule. It is possible that the role of

Y505 in VE607 binding depends on the overall conformation of the Spike, which might differ

in Omicron since it accumulated more than 30 mutations in the Spike. Altogether, our results

constitute a proof of concept showing that small molecules targeting the SARS-CoV-2 Spike

have potential for the development of drugs that may contribute to the fight against COVID-

19.

Acknowledgments

The authors thank the CRCHUM BSL3 and Flow Cytometry Platforms for technical assistance. The authors thank Hughes Charest and the LSPQ for the authentic SARS-CoV-2 virus. This work was supported by "Ministère de l'Économie et de l'Innovation du Québec, Programme de soutien aux organismes de recherche et d'innovation", by the Fondation du CHUM, by a Canadian Institutes of Health Research (CIHR) foundation grant #352417 and by an Exceptional Fund COVID-19 from the Canada Foundation for Innovation (CFI) #41027 to A.F, , and by R01 AI163395 from NIH NIAID to W.M. Work on the presented variants was also supported by the Sentinelle COVID Quebec network led by the LSPQ in collaboration with Fonds de Recherche du Québec Santé (FRQS) to A.F. The work was also supported by the UdeM Faculty of Medicine fund "Combattre la COVID-19: de la prévention au contrôle" to C.B. and A.F.; A.F. is the recipient of a Canada Research Chair on Retroviral Entry # RCHS0235 950-232424. G.B.B. is the recipient of a Fonds de Recherche Québec—Santé (FRQS) PhD fellowship. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The graphical abstract has been generated with BioRender website.

Author Contributions

A.F. and C.B. designed the studies. S.D., S.Y.G., J.G., M.M., D.V., G.B.B., V.T.V., G.G., J.R., D.Y., A.B.S., M.P., M.C., C.A., W.M., A.F., and C.B., performed the experiments and interpreted the results. Y.C., A.B.S., M.P., and M.C. contributed unique reagents. A.F. S.D., and C.B. wrote the manuscript with inputs from others. Every author has read, edited and approved the final manuscript.

Competing interests

The authors declare no-competing interests

269 **DISCLAIMER**

270 The views expressed in this manuscript are those of the authors and do not reflect the official

271 policy or position of the Uniformed Services University, US Army, the Department of

272 Defense, or the US Government.

273
274

Figure Legends

Fig. 1. Potential interactions of SARS-CoV-1 inhibitors with the RBD. (A) Chemical

structures of VE607 and SSAA09E2; (B) Differential scanning fluorimetry of the SARS-

CoV-2 RBD in the presence of SARS-CoV-1 inhibitors, results from two experiments (eight

replicates total) are shown; (C) Virtual docking of VE607 to SARS-CoV-1 and (D) SARS-

CoV-2 RBD. Left panels, the electrostatic potential is displayed over molecular surface of the

RBD and colored red and blue for negative and positive potential, respectively. Right panels,

scheme showing a docking model of VE607 to the RBD. The presumable RBD contact

residues are shown as spheres.

Fig. 2. VE607 inhibits infection of SARS-CoV-1 and SARS-CoV-2 pseudoviral particles

and of authentic SARS-CoV-2. (A) VE607 inhibition of SARS-CoV-1, SARS-CoV-2 or

VSV-G (specificity control) pseudovirus; (B) VE607 inhibition of authentic live SARS-CoV-

2 virus; (C) VE607 and the three different enantiomers are not toxic on 293T-ACE2 (left) or

VERO-E6 (right) cells, as measured by CellTiter-Glo One Solution Assay for the

quantitation of ATP presented in live cells. (D) Pseudovirus neutralization of SARS-CoV-2 S

mutants predicted by our *in silico* analysis to modulate the inhibition by VE607. Data

represents the average of at least four independent experiments ± SEM.

Fig. 3. VE607 stabilizes SARS-CoV-2 S in the "up" conformation. (A) VE607 does not

compete for sACE2 interaction, as measured by flow cytometry. The values represent the

median fluorescence intensities (MFI) normalized to binding signals obtained with the

conformationally independent CV3-25 Ab. Five experiments are represented as mean ± SEM

and statistical significance was tested using unpaired t test. (B) SARS-CoV-2 Spike stability

was measured by radioactive labelling of 293T Spike expressing cells followed by

immunoprecipitation of cell lysates and supernatants. At least four experiments are

represented as mean ± SEM and statistical significance was tested using unpaired t test, * $p <$

0.05. (C - E) Single molecule FRET analysis of SARS-CoV-2 S unliganded (C), in presence

of sACE2 (D) or VE607 (E).

Fig. 4. VE607 inhibits infection of SARS-CoV-2 variant Alpha, Beta, Gamma, Delta and

Omicron pseudovirus particles. VE607 inhibits SARS-CoV-2 pseudoviral particles

infection of 293T-hACE2 cells. IC_{50} values are shown next to the different VOCs Spikes.

Data represents the average of at least four independent experiments ± SEM.

STAR Methods

Lead Contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact andres.finzi@umontreal.ca.

Materials Availability

All unique reagents generated during this study are available from the Lead contact

andres.finzi@umontreal.ca with a completed Materials Transfer Agreement.

Data and Code Availability

This study did not generate new code.

Plasmids

The plasmids expressing the human coronavirus Spike of SARS-CoV-2 was kindly provided

by Stefan Pöhlmann and was previously reported (Hoffmann et al., 2020b). The pNL4.3 R-E-

Luc was obtained from NIH AIDS Reagent Program. The codon-optimized RBD sequence

(encoding residues 319-541) fused to a C-terminal hexahistidine tag was cloned into the

pcDNA3.1(+) expression vector and was reported elsewhere (Beaudoin-Bussieres et al.,

2020). The plasmids encoding the SARS-CoV-2 variants Spikes D614G, Alpha (B.1.1.7),

Beta (B.1.351), Gamma (P.1) were codon-optimized and synthesized by Genscript. Plasmid

encoding the Delta (B.1.617.2) and Omicron (B.1.1.529) Spikes were generated by

overlapping PCR using a codon-optimized wild-type SARS-CoV-2 Spike gene (GeneArt,

ThermoFisher) that was synthesized (Biobasic) and cloned in pCAGGS as a template

(Chatterjee et al., 2021; Gong et al., 2021b; Tauzin et al., 2022). The vesicular stomatitis

virus G (VSV-G)-encoding plasmid (pSVCMV-IN-VSV-G) was previously described (Emi et

al., 1991). Plasmids used to generate SARS-CoV-2 pseudoviral particles for smFRET analysis were described previously (Lu et. al., 2020).

Cell lines

293T human embryonic kidney cells (obtained from ATCC) and Vero E6 cells (ATCC CRL-1586™) were maintained at 37°C under 5% CO2 in Dulbecco's modified Eagle's medium (DMEM) (Wisent) containing 5% fetal bovine serum (VWR), 100 UI/ml of penicillin and 100μg/ml of streptomycin (Wisent). The 293T-ACE2 cell line was previously reported (Prevost et al., 2020).

Virus

Authentic SARS-CoV-2 was isolated, sequenced, and amplified from clinical samples obtained from patients infected with SARS-CoV-2 D614G by the Laboratoire de Santé Publique du Québec (LSPQ) and was previously described (Prevost et al., 2021). The virus was sequenced by MinION technology (Oxford Nanopore technologies, Oxford, UK). All work with the infectious SARS-CoV-2 authentic virus was performed in Biosafety Level 3 (BSL3) facilities at CRCHUM using appropriate positive-pressure air respirators and personal protective equipment.

Methods detail

Purification of SARS-CoV-2 RBD

FreeStyle 293 F cells (Invitrogen) were grown in FreeStyle 293F medium (Invitrogen) to a density of 1×10^6 cells/mL at 37°C with 8% CO_2 with regular agitation (150 rpm). Cells were transfected with a plasmid coding for SARS-CoV-2 S RBD using ExpiFectamine 293

transfection reagent, as directed by the manufacturer (Invitrogen). One week later, cells were

pelleted and discarded. Supernatants were filtered using a 0.22 μm filter (Thermo Fisher

Scientific). The recombinant RBD proteins were purified by nickel affinity columns, as

directed by the manufacturer (Invitrogen). The RBD preparations were dialyzed against

phosphate-buffered saline (PBS) and stored in aliquots at -80°C until further use. To assess

purity, recombinant proteins were loaded on SDS-PAGE gels and stained with Coomassie

Blue.

Differential scanning fluorimetry

DSF experiments were essentially performed as described previously (Sharifahmadian et al.,

2017). DSF was conducted using 5 μM of purified RBD, 10x concentration of SYPRO

Orange (from 5000x stock solution, ThermoFisher) in 50 mM HEPES, 100 mM NaCl, pH 7.5

and 5% final concentration of DMSO. The small molecules were added to final

concentrations of 5 mM. SYPRO Orange fluorescence was monitored over 20–95 °C with a

LightCycler® 480 instrument (Roche, USA). The LightCycler® 480 Software (Roche) was

used to calculate the first derivate of the resulting melting curve, with the steepest point of the

slope being the Tm.

Molecular modeling

System preparation, modeling, and docking calculation were performed using the Schrödinger

Suite package (Schrödinger, 2020), using default parameters unless otherwise noted. The

target structures were taken from SARS-CoV-1 RBD (PDB ID: 6waq) and SARS-CoV-2

RBD (PDB ID: 6w41) prepared using the Protein Preparation Wizard (Sastry et al., 2013). To

prepare the structures, force field atom types and bond orders were assigned, missing atoms

and side-chains were added, protonation states of ionizable amino acid side-chains were

determined using PROPKA (Olsson et al., 2011), water orientations were sampled, and

hydrogen bond networks were subsequently optimized by flipping Asn/Gln/His residues and

sampling hydroxyl/thiol hydrogen. Constrained energy minimization was then performed

using the imperf module from impact (Schrödinger, 2020) to generate the structure to be used

in the subsequent modeling calculations. Potential binding sites were explored and

characterized using the SiteMap tool (Halgren, 2007; Halgren, 2009). VE607 compound was

structurally preprocessed using LigPrep (Schrödinger, 2020) to generate multiple states for

stereoisomers, tautomers, ring conformations, and protonation states at a selected pH range.

Then, energy minimization was performed with the OPLS3e force field (Roos et al., 2019).

The prepared molecular structures were docked into the putative binding sites using Glide

(Friesner et al., 2004; Halgren et al., 2004) with the standard precision (SP) scoring function

to evaluate enrichment of the calculated docked models.

Neutralization assay using pseudoviral particles

Target cells were infected with single-round luciferase-expressing lentiviral particles as

described previously (Prevost et al., 2020). Briefly, 293T cells were transfected by the

calcium phosphate method with the lentiviral vector pNL4.3 R-E- Luc (NIH AIDS Reagent

Program) and a plasmid encoding for SARS-CoV-2 Spike at a ratio of 5:4. Two days post-

transfection, cell supernatants were harvested and stored at –80°C until use. 293T-ACE2

target cells were seeded at a density of 1×10^4 cells/well in 96-well luminometer-compatible

tissue culture plates (Perkin Elmer) 24h before infection. Recombinant viruses in a final

volume of 100µl were incubated with the indicated concentrations of small molecules (VE607

or SSAA009E2) up to concentrations of 100 µM for 1h at 37°C and were then added to the

target cells followed by incubation for 48h at 37°C; cells were lysed by the addition of 30µl of

passive lysis buffer (Promega) followed by one freeze-thaw cycle. An LB941 TriStar

luminometer (Berthold Technologies) was used to measure the luciferase activity of each well

after the addition of 100µl of luciferin buffer (15mM $MgSO_4$, 15mM KPO_4 [pH 7.8], 1mM

ATP, and 1mM dithiothreitol) and 50µl of 1mM d-luciferin potassium salt (Prolume). The

neutralization half-maximal inhibitory dilution (ID_{50}) represents the sera dilution to inhibit

50% of the infection of 293T-ACE2 cells by recombinant viruses.

Cell surface staining and flow cytometry analysis

Using the standard calcium phosphate method, 10 µg of Spike expressor and 2.5 µg of a green

fluorescent protein (GFP) expressor (pIRES-GFP) were transfected into 2×10^6 293T cells.

48h post-transfection Spike-expressing cells were incubated with 100 µM of VE607 or

equivalent volume of vehicle (DMSO) and incubated for 30 min at room temperature. CV3-

25 (5 µg/ml) or sACE2 (100 µg/ml) was added to the cells and incubated for 45min at 37ºC

and sACE2 binding was detected using a polyclonal Goat anti-human ACE2 (RND Systems)

at 1/100 dilution at room temperature for 30min. AlexaFluor-647-conjugated goat anti-human

IgG (H+L) Ab (Invitrogen) and AlexaFluor-conjugated donkey anti-goat IgG (H+L) Ab

(Invitrogen) was used as secondary antibodies. The percentage of transfected cells (GFP+

cells) was determined by gating the living cell population based on viability dye staining

(Aqua Vivid, Invitrogen). sACE2 binding levels were normalized to signals obtained with the

conformationally independent anti-S2 CV3-25 mAb (Li et al., 2022; Prevost et al., 2021;

Tauzin et al., 2022).   Samples were acquired on a LSRII cytometer (BD Biosciences,

Mississauga, ON, Canada) and data analysis was performed using FlowJo vX.0.7 (Tree Star,

Ashland, OR, USA).

Radioactive labeling and immunoprecipitation

For pulse-labeling experiments, 5×10^5 293T cells were transfected by the calcium phosphate method with SARS-CoV-2 D614G Spike expressor. One day after transfection, cells were metabolically labeled for 16-24 h with 100 µCi/ml [^{35}S]methionine-cysteine ([^{35}S] protein labeling mix; Perkin-Elmer) in Dulbecco's modified Eagle's medium lacking methionine and cysteine and supplemented with 10% of dialyzed fetal bovine serum and 1% GlutaMAX™. Simultaneously, cells were treated with or without 100 µM VE607. Cells were subsequently lysed in radio immunoprecipitation assay (RIPA) buffer (140 mM NaCl, 8 mM Na$_2$HPO4, 2 mM NaH$_2$PO$_4$, 1% IGEPAL® CA-630, 0.05% sodium dodecyl sulfate [SDS], 1.2 mM sodium deoxycholate [DOC]). Precipitation of radiolabeled SARS-CoV-2 D614G Spike glycoprotein from cell lysates or supernatant was performed with CV3-25 and polyclonal rabbit antiserum raised against SARS-CoV-2 RBD protein, for 1 h at 4°C in the presence of 45 µL of 10% protein A-Sepharose beads (GE Healthcare). Samples were washed twice with RIPA buffer and then boiled 5 min in Laemmli buffer with β-mercaptoethanol before being separated by SDS-PAGE. After migration, gels were dried with a Model 583 gel dryer (Bio-Rad) and exposed to a storage phosphor screen. Densitometry data were acquired with a Typhoon Trio Variable Mode Imager (Amersham Biosciences) in storage phosphor acquisition mode and analyzed using ImageQuant 5.2 (Molecular Dynamics). Association index was determined by precipitation of radiolabeled cell lysates and supernatants with CV3-25 and polyclonal rabbit antiserum raised against SARS-CoV-2 RBD protein. The association index is a measure of the ability of the VE607 treated S1 subunit to remain associated with the trimeric spike (S) protein on the expressing cell relative to that of the mock-treated S1 and was calculated with the following formula: association index = ([cell S1]$_{treated}$/[supernatant S1]$_{treated}$)/([cell S1]$_{mock-treated}$/[supernatant S1]$_{mock-treated}$).

Microneutralization with authentic virus

One day prior to infection, 2×10^4 Vero E6 cells were seeded per well in the 96-well flat bottom plate and incubated overnight to permit Vero E6 cell adherence. Compounds dilutions ranged from 0, 0.316, 1, 3.16, 10, 31.6 and 100 µM were performed in a separate 96 well culture plate using DMEM supplemented with penicillin (100 U/mL), streptomycin (100 µg/mL), HEPES, 0.12% sodium bicarbonate, 2% FBS and 0.24% BSA. 10^4 $TCID_{50}$/mL of SARS-CoV-2 virus was prepared in DMEM + 2% FBS and combined with an equivalent volume of diluted compounds for one hour. After this incubation, all media was removed from the 96 well plate seeded with Vero E6 cells and virus: compounds mixture was added to each respective well at a volume corresponding to 600 $TCID_{50}$ per well and incubated for one hour further at 37°C. Both virus only and media only (MEM + 2% FBS) conditions were included in this assay. All virus-compounds supernatant was removed from wells without disrupting the Vero E6 monolayer. Each diluted compound (100 µL) was added to its respective Vero E6-seeded well in addition to an equivalent volume of MEM + 2% FBS and was then incubated for 48 hours. Media was then discarded and replaced with 10% formaldehyde for 24 hours to cross-link Vero E6 monolayer. Formaldehyde was removed from wells and subsequently washed with PBS. Cell monolayers were permeabilized for 15 minutes at room temperature with PBS + 0.1% Triton X-100, washed with PBS and then incubated for one hour at room temperature with PBS + 3% non-fat milk. An anti-mouse SARS-CoV-2 nucleocapsid protein (Clone 1C7, Bioss Antibodies) primary antibody solution was prepared at 1 µg/mL in PBS + 1% non-fat milk and added to all wells for one hour at room temperature. Following extensive washing (3×) with PBS, an anti-mouse IgG HRP secondary antibody solution was formulated in PBS + 1% non-fat milk. One hour post-room temperature incubation, wells were washed with 3× PBS, substrate (ECL) was added and an

LB941 TriStar luminometer (Berthold Technologies) was used to measure the signal each

well.

Cell viability test

To measure the cytotoxicity of VE607 and its stereoisomers on 293T-ACE2 or Vero-E6 cells,

a cell viability assay using CellTiter-Glo® One Solution Assay (Promega) was performed.

Briefly, 293T-ACE2 or Vero-E6 cells were seeded at a density of 1×10^4 cells/well in 96-well

luminometer-compatible tissue culture plates (Perkin Elmer); After 24h, indicated

concentrations of VE607, (S,S)-VE607, (R,R)-VE607 or (R,S)-VE607 up to concentrations of

100 μM were added to the cells followed by incubation for 48h at 37°C, same volume of its

vehicle, DMSO, was added as control. Then a volume of CellTiter-Glo® One Solution equal

to the volume of cell culture medium present in each well was added, followed by 2 min

mixing on shaker and 10 min incubation at room temperature. An LB941 TriStar luminometer

(Berthold Technologies) was used to measure the luciferase activity of each well.

Chemical synthesis of the three enantiomers of VE607

All reactions were conducted in oven-dried glassware under an inert atmosphere of nitrogen,

unless otherwise stated. All solvents were reagent or high-performance liquid chromatography

(HPLC) grade. Anhydrous THF was obtained from the Pure SolveTM PS-400 system under

an argon atmosphere. All reagents were purchased from commercially available sources and

used as received. Reactions were magnetically stirred under a nitrogen atmosphere, unless

otherwise noted and were monitored by thin layer chromatography (TLC) was performed on

pre-coated silica gel 60 F-254 plates (40-55 micron, 230-400 mesh) and visualized by UV

light or staining with KMnO4 and heating. Yields refer to chromatographically and

spectroscopically pure compounds. Optical rotations were measured on a JASCO P-200

polarimeter. Proton (^{1}H) and carbon (13C) NMR spectra were recorded on a Bruker Avance

III 500-MHz spectrometer. Chemical shifts (δ) are reported in parts per million (ppm) relative

to chloroform (δ 7.26) or methanol (δ 3.31) for ^{1}H NMR, and chloroform (δ 77.2) or methanol

(δ 49.0). High resolution mass spectra (HRMS) were recorded at the University of

Pennsylvania Mass Spectroscopy Service Center on either a VG Micromass 70/70H or VG

ZAB-E spectrometer. Lyophilization was performed in a Labconco FreeZone 12 Plus

lyophilizer (0.148 mbar). The purity of new compounds were judged by NMR and LCMS

(>95%).

smFRET analysis

Pseudoviral particles bearing labeled CoV2$_{WH01}$ S protein were prepared, imaged, and

analyzed as described previously (Li et al., 2021c; Lu et al., 2020). Samples were pre-

incubated with sACE2 (200 μg/ml) or VE607 (100 μM) for 90 minutes at room temperature

prior to imaging.

Quantification and statistical analysis

Statistics were analyzed using GraphPad Prism version 8.0.2 (GraphPad, San Diego, CA,

(USA). Every data set was tested for statistical normality and this information was used to

apply the appropriate (parametric or nonparametric) statistical test. P values <0.05 were

considered significant; significance values are indicated as * $p<0.05$; ** $p<0.01$; *** $p<0.001$;

**** $p<0.0001$.

References

Adedeji, A.O., Severson, W., Jonsson, C., Singh, K., Weiss, S.R., and Sarafianos, S.G. (2013). Novel inhibitors of severe acute respiratory syndrome coronavirus entry that act by three distinct mechanisms. J Virol *87*, 8017-8028.

Anand, S.P., Chen, Y., Prevost, J., Gasser, R., Beaudoin-Bussieres, G., Abrams, C.F., Pazgier, M., and Finzi, A. (2020). Interaction of Human ACE2 to Membrane-Bound SARS-CoV-1 and SARS-CoV-2 S Glycoproteins. Viruses *12*.

Ball, P. (2021). The lightning-fast quest for COVID vaccines - and what it means for other diseases. Nature *589*, 16-18.

Beaudoin-Bussieres, G., Laumaea, A., Anand, S.P., Prevost, J., Gasser, R., Goyette, G., Medjahed, H., Perreault, J., Tremblay, T., Lewin, A.*, et al.* (2020). Decline of Humoral Responses against SARS-CoV-2 Spike in Convalescent Individuals. mBio *11*.

Cai, Y., Zhang, J., Xiao, T., Peng, H., Sterling, S.M., Walsh, R.M., Jr., Rawson, S., Rits-Volloch, S., and Chen, B. (2020). Distinct conformational states of SARS-CoV-2 spike protein. Science *369*, 1586-1592.

Chatterjee, D., Tauzin, A., Marchitto, L., Gong, S.Y., Boutin, M., Bourassa, C., Beaudoin-Bussières, G., Bo, Y., Ding, S., Laumaea, A.*, et al.* (2021). SARS-CoV-2 Omicron Spike recognition by plasma from individuals receiving BNT162b2 mRNA vaccination with a 16-weeks interval between doses. 2021.2012.2021.473679.

Cohen, J. (2021). The dream vaccine. Science *372*, 227-231.

Dai, W., Zhang, B., Su, H., Li, J., Zhao, Y., Xie, X., Jin, Z., Liu, F., Li, C., Li, Y.*, et al.* (2020). Structure-based design of antiviral drug candidates targeting the SARS-CoV-2 main protease. Science.

Ding, S., Adam, D., Beaudoin-Bussières, G., Tauzin, A., Gong, S.Y., Gasser, R., Laumaea, A., Anand, S.P., Privé, A., Bourassa, C.*, et al.* (2022). SARS-CoV-2 Spike Expression at the Surface of Infected Primary Human Airway Epithelial Cells. Viruses *14*, 5.

Emi, N., Friedmann, T., and Yee, J.K. (1991). Pseudotype formation of murine leukemia virus with the G protein of vesicular stomatitis virus. J Virol *65*, 1202-1207.

Friesner, R.A., Banks, J.L., Murphy, R.B., Halgren, T.A., Klicic, J.J., Mainz, D.T., Repasky, M.P., Knoll, E.H., Shelley, M., Perry, J.K.*, et al.* (2004). Glide: a new approach for rapid, accurate docking and scoring. 1. Method and assessment of docking accuracy. J Med Chem *47*, 1739-1749.

Gong, S.Y., Chatterjee, D., Richard, J., Prevost, J., Tauzin, A., Gasser, R., Bo, Y., Vezina, D., Goyette, G., Gendron-Lepage, G.*, et al.* (2021a). Contribution of single mutations to selected SARS-CoV-2 emerging variants spike antigenicity. Virology *563*, 134-145.

Gong, S.Y., Chatterjee, D., Richard, J., Prévost, J., Tauzin, A., Gasser, R., Bo, Y., Vézina, D., Goyette, G., Gendron-Lepage, G.*, et al.* (2021b). Contribution of single mutations to selected SARS-CoV-2 emerging variants Spike antigenicity. 2021.2008.2004.455140.

Halgren, T. (2007). New method for fast and accurate binding-site identification and analysis. Chem Biol Drug Des *69*, 146-148.

Halgren, T.A. (2009). Identifying and characterizing binding sites and assessing druggability. J Chem Inf Model *49*, 377-389.

Halgren, T.A., Murphy, R.B., Friesner, R.A., Beard, H.S., Frye, L.L., Pollard, W.T., and Banks, J.L. (2004). Glide: a new approach for rapid, accurate docking and scoring. 2. Enrichment factors in database screening. J Med Chem *47*, 1750-1759.

Harrison, A.G., Lin, T., and Wang, P. (2020). Mechanisms of SARS-CoV-2 Transmission and Pathogenesis. Trends Immunol *41*, 1100-1115.

Hoffmann, M., Kleine-Weber, H., and Pohlmann, S. (2020a). A Multibasic Cleavage Site in
the Spike Protein of SARS-CoV-2 Is Essential for Infection of Human Lung Cells. Molecular
cell.
Hoffmann, M., Kleine-Weber, H., Schroeder, S., Kruger, N., Herrler, T., Erichsen, S.,
Schiergens, T.S., Herrler, G., Wu, N.H., Nitsche, A., *et al.* (2020b). SARS-CoV-2 Cell Entry
Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor.
Cell *181*, 271-280 e278.
Kao, R.Y., Tsui, W.H., Lee, T.S., Tanner, J.A., Watt, R.M., Huang, J.D., Hu, L., Chen, G.,
Chen, Z., Zhang, L., *et al.* (2004). Identification of novel small-molecule inhibitors of severe
acute respiratory syndrome-associated coronavirus by chemical genetics. Chem Biol *11*,
1293-1299.
Kokic, G., Hillen, H.S., Tegunov, D., Dienemann, C., Seitz, F., Schmitzova, J., Farnung, L.,
Siewert, A., Hobartner, C., and Cramer, P. (2021). Mechanism of SARS-CoV-2 polymerase
stalling by remdesivir. Nat Commun *12*, 279.
Krammer, F. (2020). SARS-CoV-2 vaccines in development. Nature *586*, 516-527.
Lan, J., Ge, J., Yu, J., Shan, S., Zhou, H., Fan, S., Zhang, Q., Shi, X., Wang, Q., Zhang, L., *et*
al. (2020). Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2
receptor. Nature *581*, 215-220.
Li, J., Lai, S., Gao, G.F., and Shi, W. (2021a). The emergence, genomic diversity and global
spread of SARS-CoV-2. Nature *600*, 408-418.
Li, W., Chen, Y., Prevost, J., Ullah, I., Lu, M., Gong, S.Y., Tauzin, A., Gasser, R., Vezina,
D., Anand, S.P., *et al.* (2021b). Structural Basis and Mode of Action for Two Broadly
Neutralizing Antibodies Against SARS-CoV-2 Emerging Variants of Concern. bioRxiv.
Li, W., Chen, Y., Prevost, J., Ullah, I., Lu, M., Gong, S.Y., Tauzin, A., Gasser, R., Vezina,
D., Anand, S.P., *et al.* (2021c). Structural Basis and Mode of Action for Two Broadly
Neutralizing Antibodies Against SARS-CoV-2 Emerging Variants of Concern. Cell reports
S2211-1247(21)01714-9.
Li, W., Chen, Y., Prevost, J., Ullah, I., Lu, M., Gong, S.Y., Tauzin, A., Gasser, R., Vezina,
D., Anand, S.P., *et al.* (2022). Structural basis and mode of action for two broadly neutralizing
antibodies against SARS-CoV-2 emerging variants of concern. Cell reports *38*, 110210.
Lu, M., Uchil, P.D., Li, W., Zheng, D., Terry, D.S., Gorman, J., Shi, W., Zhang, B., Zhou, T.,
Ding, S., *et al.* (2020). Real-Time Conformational Dynamics of SARS-CoV-2 Spikes on
Virus Particles. Cell Host Microbe *28*, 880-891 e888.
Mannar, D., Saville, J.W., Zhu, X., Srivastava, S.S., Berezuk, A.M., Zhou, S., Tuttle, K.S.,
Kim, A., Li, W., Dimitrov, D.S., *et al.* (2021). Structural analysis of receptor binding domain
mutations in SARS-CoV-2 variants of concern that modulate ACE2 and antibody binding.
Cell Rep *37*, 110156.
Mashalidis, E.H., Sledz, P., Lang, S., and Abell, C. (2013). A three-stage biophysical
screening cascade for fragment-based drug discovery. Nature protocols *8*, 2309-2324.
Nabel, K.G., Clark, S.A., Shankar, S., Pan, J., Clark, L.E., Yang, P., Coscia, A., McKay,
L.G.A., Varnum, H.H., Brusic, V., *et al.* (2021). Structural basis for continued antibody
evasion by the SARS-CoV-2 receptor binding domain. Science, eabl6251.
Olsson, M.H., Sondergaard, C.R., Rostkowski, M., and Jensen, J.H. (2011). PROPKA3:
Consistent Treatment of Internal and Surface Residues in Empirical pKa Predictions. J Chem
Theory Comput *7*, 525-537.
Owen, D.R., Allerton, C.M.N., Anderson, A.S., Aschenbrenner, L., Avery, M., Berritt, S.,
Boras, B., Cardin, R.D., Carlo, A., Coffman, K.J., *et al.* (2021). An oral SARS-CoV-2 M(pro)
inhibitor clinical candidate for the treatment of COVID-19. Science *374*, 1586-1593.
Prevost, J., and Finzi, A. (2021). The great escape? SARS-CoV-2 variants evading
neutralizing responses. Cell Host Microbe *29*, 322-324.

Prevost, J., Gasser, R., Beaudoin-Bussieres, G., Richard, J., Duerr, R., Laumaea, A., Anand,
S.P., Goyette, G., Benlarbi, M., Ding, S., *et al.* (2020). Cross-Sectional Evaluation of
Humoral Responses against SARS-CoV-2 Spike. Cell Rep Med *1*, 100126.
Prevost, J., Richard, J., Gasser, R., Ding, S., Fage, C., Anand, S.P., Adam, D., Gupta Vergara,
N., Tauzin, A., Benlarbi, M., *et al.* (2021). Impact of temperature on the affinity of SARS-
CoV-2 Spike glycoprotein for host ACE2. J Biol Chem *297*, 101151.
Rappazzo, C.G., Tse, L.V., Kaku, C.I., Wrapp, D., Sakharkar, M., Huang, D., Deveau, L.M.,
Yockachonis, T.J., Herbert, A.S., Battles, M.B., *et al.* (2021). Broad and potent activity
against SARS-like viruses by an engineered human monoclonal antibody. Science *371*, 823-
829.
Roos, K., Wu, C., Damm, W., Reboul, M., Stevenson, J.M., Lu, C., Dahlgren, M.K., Mondal,
S., Chen, W., Wang, L., *et al.* (2019). OPLS3e: Extending Force Field Coverage for Drug-
Like Small Molecules. J Chem Theory Comput *15*, 1863-1874.
Sastry, G.M., Adzhigirey, M., Day, T., Annabhimoju, R., and Sherman, W. (2013). Protein
and ligand preparation: parameters, protocols, and influence on virtual screening enrichments.
J Comput Aided Mol Des *27*, 221-234.
Schrödinger (2020). Release 2020-4. Induced Fit Docking protocol. Glide, Schrödinger, LLC,
New York, NY.
Shang, J., Ye, G., Shi, K., Wan, Y., Luo, C., Aihara, H., Geng, Q., Auerbach, A., and Li, F.
(2020). Structural basis of receptor recognition by SARS-CoV-2. Nature.
Sharifahmadian, M., Arya, T., Bessette, B., Lecoq, L., Ruediger, E., Omichinski, J.G., and
Baron, C. (2017). Monomer-to-dimer transition of Brucella suis type IV secretion system
component VirB8 induces conformational changes. The FEBS journal *284*, 1218-1232.
Tauzin, A., Gong, S.Y., Beaudoin-Bussieres, G., Vezina, D., Gasser, R., Nault, L., Marchitto,
L., Benlarbi, M., Chatterjee, D., Nayrac, M., *et al.* (2022). Strong humoral immune responses
against SARS-CoV-2 Spike after BNT162b2 mRNA vaccination with a 16-week interval
between doses. Cell Host Microbe *30*, 97-109 e105.
Tauzin, A., Nayrac, M., Benlarbi, M., Gong, S.Y., Gasser, R., Beaudoin-Bussieres, G.,
Brassard, N., Laumaea, A., Vezina, D., Prevost, J., *et al.* (2021). A single dose of the SARS-
CoV-2 vaccine BNT162b2 elicits Fc-mediated antibody effector functions and T cell
responses. Cell Host Microbe *29*, 1137-1150 e1136.
Tong, T. (2009). Therapies for coronaviruses. Part I of II – viral entry inhibitors. Expert
Opinion on Therapeutic Patents *19*.
Ullah, I., Prevost, J., Ladinsky, M.S., Stone, H., Lu, M., Anand, S.P., Beaudoin-Bussieres, G.,
Symmes, K., Benlarbi, M., Ding, S., *et al.* (2021). Live imaging of SARS-CoV-2 infection in
mice reveals that neutralizing antibodies require Fc function for optimal efficacy. Immunity
54, 2143-2158 e2115.
Wang, G., Yang, M.L., Duan, Z.L., Liu, F.L., Jin, L., Long, C.B., Zhang, M., Tang, X.P., Xu,
L., Li, Y.C., *et al.* (2021). Dalbavancin binds ACE2 to block its interaction with SARS-CoV-
2 spike protein and is effective in inhibiting SARS-CoV-2 infection in animal models. Cell
Res *31*, 17-24.
Wrapp, D., Wang, N., Corbett, K.S., Goldsmith, J.A., Hsieh, C.L., Abiona, O., Graham, B.S.,
and McLellan, J.S. (2020). Cryo-EM structure of the 2019-nCoV spike in the prefusion
conformation. Science *367*, 1260-1263.
Wu, F., Zhao, S., Yu, B., Chen, Y.M., Wang, W., Song, Z.G., Hu, Y., Tao, Z.W., Tian, J.H.,
Pei, Y.Y., *et al.* (2020). A new coronavirus associated with human respiratory disease in
China. Nature *579*, 265-269.
Yan, R., Zhang, Y., Li, Y., Xia, L., Guo, Y., and Zhou, Q. (2020). Structural basis for the
recognition of SARS-CoV-2 by full-length human ACE2. Science *367*, 1444-1448.

Yang, H., and Rao, Z. (2021). Structural biology of SARS-CoV-2 and implications for
therapeutic development. Nature reviews Microbiology *19*, 685-700.
Yang, Z., Han, Y., Ding, S., Finzi, A., Mothes, W., and Lu, M. (2021). SARS-CoV-2 variants
exhibit increased kinetic stability of open spike conformations as an evolutionary strategy.
2021.2010.2011.463956.
Yin, W., Luan, X., Li, Z., Zhou, Z., Wang, Q., Gao, M., Wang, X., Zhou, F., Shi, J., You, E.*,
et al.* (2021). Structural basis for inhibition of the SARS-CoV-2 RNA polymerase by suramin.
Nat Struct Mol Biol *28*, 319-325.
Zhang, L., Lin, D., Sun, X., Curth, U., Drosten, C., Sauerhering, L., Becker, S., Rox, K., and
Hilgenfeld, R. (2020). Crystal structure of SARS-CoV-2 main protease provides a basis for
design of improved alpha-ketoamide inhibitors. Science *368*, 409-412.
Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu,
R.*, et al.* (2020a). A Novel Coronavirus from Patients with Pneumonia in China, 2019. N
Engl J Med *382*, 727-733.
Zhu, Z.L., Qiu, X.D., Wu, S., Liu, Y.T., Zhao, T., Sun, Z.H., Li, Z.R., and Shan, G.Z.
(2020b). Blocking Effect of Demethylzeylasteral on the Interaction between Human ACE2
Protein and SARS-CoV-2 RBD Protein Discovered Using SPR Technology. Molecules *26*.
697

Figure 1

Figure 2

Figure 3

Figure 4

Supplemental Figure 1. Stereochemical isomers of VE607 have the same capacity as commercially available VE607 to neutralize SARS-CoV-2 pseudovirus particles. All three stereochemical isomers of VE607, (S,S)-VE607, (R,R)-VE607 and (R,S)-VE607 were tested for their inhibition of VSV-G pseudovirus or SARS-CoV-2 D614G pseudovirus. IC50 values are shown next to the different VOC Spikes. Data represents the average of at least four independent experiments ± SEM.