Mast Cell Derived Histamine Negatively Regulates Hematopoiesis

Bailey R. Klein[1], Julianne N.P. Smith[1], Ramachandra Katabathula[1], Rahul Chaudhary[1],

Zhenxiang Gao[1], Brittany A. Cordova[1], Frederick Petroze, Riya Tiwari[1], Stanton L. Gerson[1,2],

Rong Xu[1], Sanford D. Markowitz[1,2], Amar B. Desai[1]

[1]Department of Medicine, and Case Comprehensive Cancer Center Case Western Reserve

University, Cleveland OH 44106 USA

[2]University Hospitals Seidman Cancer Center, Cleveland OH 44106 US

Author Contribution: BRK conceived and designed the study, collected and interpreted the data,

and wrote the manuscript. JNPS conceived and designed the study and collected and interpreted

the data. RK designed studies, interpreted data, and contributed to manuscript writing. RC

collected the data. ZG collected and interpreted the data. BAC collected the data. FP collected

the data. RT collected the data. SLG designed studies and interpreted data. RX designed studies,

interpreted data, and contributed to manuscript writing. SDM conceived and designed studies

and interpreted data. ABD conceived and designed the study, interpreted the data, wrote the

manuscript, and provided final approval of the manuscript.

*Correspondence:

Amar B. Desai

Department of Medicine

Case Western Reserve University

Abd10@case.edu

Running title: Mast Cells Negatively Regulate HSCs

Total word count: 3310

Abstract word count: 236

Figures: 5

Tables: 1

References: 46

ABSTRACT

Hematopoietic stem cells (HSCs) are essential for generating all blood cell types and maintaining

immune function and oxygen transport. This requires tight regulation of self-renewal,

differentiation, and quiescence, driven by intrinsic and extrinsic signals. While the influence of

many HSC progeny on HSC decisions are recognized, the role of mast cells (MCs) remain

understudied. MCs are known for their immunomodulatory functions through the secretion of

factors such as histamine and could offer new insights into HSC regulation. In this study, we

describe a novel role for MC-derived histamine in modulating HSC behavior. We observed that

genetically MC-deficient "SASH" mice exhibit increased hematopoietic output and bone marrow

(BM) HSCs, characterized by an enhanced quiescent signature that increases resistance to

myeloablative chemotherapy. The SASH microenvironment also contained increased frequencies

of HSC-supportive cell types and expression of genes conducive to HSC maintenance, which

together accelerated HSC engraftment when wild-type BM was transplanted into SASH

recipients. Moreover, we found lower serum histamine levels in SASH mice, and that the

enhanced hematopoietic phenotype observed in these mice could be reversed by administering

exogenous histamine. Subsequent experiments with FDA-approved antihistamines in wild-type

mice revealed that cetirizine, an H1R inverse agonist, notably increased HSC frequency in the

BM. Overall, our findings implicate MCs are negative regulators of HSC function. This lays the

groundwork for future studies to elucidate the underlying mechanisms and explore the

therapeutic potential of modulating histamine signaling to promote hematopoiesis.

INTRODUCTION

Hematopoietic stem cells (HSCs) are primitive multipotent cells that reside in the bone marrow

(BM) and differentiate into all the cellular components of blood, including myeloid,

megakaryocytic, lymphoid, and erythroid lineage cells[1]. This process of differentiation, known as

hematopoiesis, occurs life-long from a small pool of self-renewing HSCs, which employ

protective mechanisms to ensure lifelong functionality. These include residing in a hypoxic niche

to minimize exposure to reactive oxygen species, maintaining a quiescent state to protect

themselves from genetic insults, and the undergoing apoptosis in response to excessive cellular

damage[2-4]. The BM stroma consists of mesenchymal cells, endothelial cells, and some HSC

progeny, such as megakaryocytes and macrophages. It provides a supportive microenvironment

for HSC functions by producing essential factors including stem cell factor (SCF), CXCL12, and

thrombopoietin (TPO), which are necessary to maintain the BM niche[5-9]. Among these cellular

interactions and signaling mechanisms, recent discoveries have highlighted a distinct role for

histamine derived from bone marrow myeloid cells. Specifically, histamine has been found to

induce quiescence and self-renewal by activating the H2 receptor on myeloid-biased HSCs,

pointing to a nuanced interplay of local and systemic factors that collectively safeguard HSC

integrity[10]. Currently, there are no studies linking mast cells (MCs), which are one of the largest

producers of histamine, to HSC function. This may be due to a lack of proximity as MC

precursors originate in the BM, but then migrate to peripheral tissues for further development,

resulting in few to no mature MCs within the BM[11-14].

MCs are tissue-resident immune cells of the myeloid lineage located in mucosal and epithelial

tissues throughout the body[15]. These cells are canonically known to be first responders to

infection and mediators of allergy through activation of FcεRI and pattern recognition

receptors[16, 17]. Activation of MCs causes a release of mediators such as proteases, leukotrienes,

and histamine that elicit physiological responses including vasodilation, inflammatory cell

recruitment, and angiogenesis[18, 19]. Degranulation releases copious amounts of histamine, but

MCs also constitutively secrete a basal level of histamine systemically where it can bind and

activate one of its four distinct cognate G-coupled protein receptors: H1R, H2R, H3R, and

H4R[20-23]. Antihistamines, a well-tolerated class of drugs, bind to these receptors to inhibit their

activation and physiological effects. H1R and H2R antihistamines, approved by the FDA and

used by millions of people daily are used to treat allergy symptoms and gastric ulcers

respectively[24, 25].

In this study, we showcase a unique phenotype wherein MCs exhibit a negative regulatory

influence on hematopoiesis and hematopoietic output. Additionally, we present evidence that this

phenotype can be replicated pharmacologically by targeting H1R with the FDA approved

antihistamine cetirizine. These findings suggest a potential strategy for enhancing hematopoietic

output in multiple scenarios including chemotherapy induced myelosuppression, transplantation,

and bone marrow failure.

RESULTS

MC deficient Kit^(W-sh/W-sh) "SASH" mice display leukocytosis and BM HSPC expansion.

We characterized the steady state hematopoietic compartments of the commercially available

MC deficient mouse model (SASH c-Kit^(w-sh), Jackson Labs Stock #030764). The Kit^(W-sh/W-sh)

(hereinafter referred to as SASH) mice harbor an inversion mutation impacting the

transcriptional regulatory elements of the *c-Kit* gene leading to a total ablation of their MC

population shortly after birth[26, 27]. Complete blood count analysis on peripheral blood

demonstrated SASH mice have a 1.2-fold increase in total white blood cells (WBCs), 1.8-fold

increase in neutrophils (NE), 1.6-fold increase in monocytes (MO), and 1.4-fold increase in

platelets (PLT) (Figure 1A). Despite the increases in peripheral blood WBCs and specific

myeloid populations, we found no impact of MC deficiency on peripheral blood lymphocytes

(LY) or RBCs (Supplemental Figure 1). In the BM, we observed a 1.4-fold increase in total

cellularity and through immunophenotypic analysis (gating strategy outlined in Supplemental

Figure 7) we identified significant 1.7-fold increase in lineage$^-$Sca-1$^+$c-Kit$^+$ (LSK) cells and 2.1-

fold increase in CD150$^+$CD48$^-$ long-term HSCs (LT-HSC) (Figure 1B). Additionally, we

investigated the myeloid and lymphoid frequencies within the BM (gating strategy outlined in

Supplemental Figure 8) and observed a significant increase in total myeloid frequency and a

significant decrease in B cells and CD8$^+$T cells (Supplemental Figure 1).

We additionally characterized spleens of SASH mice and found these mice displayed splenic

extramedullary hematopoiesis as indicated by significant increases in splenic HSPCs and CFU

potential (Supplemental Figure 2). This is consistent with previously reported findings of

extramedullary hematopoiesis in the spleen of SASH mice[28]. Taken together, these data

demonstrate significantly enhanced hematopoietic output following loss of MCs. To next test the

functional consequence of the observed increase in hematopoietic output we assessed the

capacity of SASH mice to recover following repeated 5-Fluorouracil (5-FU) administration. 5-

FU, a pyrimidine analog, disrupts DNA synthesis and repair, preferentially targeting and

eliminating actively cycling cells within the hematopoietic system. Therefore, 5-FU can be

utilized in hematopoietic recovery studies to induce genotoxic stress, creating a selective

pressure against cycling HSPCs. Consequently, the ability to recover from 5-FU relies on the

reserve of quiescent HSCs to replenish the depleted hematopoietic cell populations. This makes

5-FU an ideal agent for evaluating the depth of the HSC pool, as resistance to its cytotoxic

effects suggests the presence of a substantial reservoir of quiescent HSCs[29, 30]. Indeed, we

observed that genetic loss of MCs resulted in a significant survival extension compared to WT

mice following weekly 150 mg/kg 5-FU administrations (Figure 1C), indicating that the

increased phenotypic HSCs and progenitors are functional and confer protection against

hematopoietic injury.

SASH lineage depleted BM exhibit transcriptional signatures indicative of HSPC

quiescence.

Next, to elucidate the observed hematopoietic phenotype in SASH mice we performed RNA

sequencing on both bulk BM cells and lineage depleted BM cells (a population enriched for

hematopoietic stem and progenitor cells) from both WT and SASH mice (Figure 2A). Principal

component analysis plots indicate no obvious stratification between bulk bone marrow

populations of SASH and WT mice (teal vs. red). However, lineage depleted populations

demonstrated a clear divergence in expression profiles (green vs. purple), suggesting a unique

HSPCs signature in SASH mice (Figure 2B). Gene set enrichment analyses of lineage depleted

cells (Supplemental Figure 3) in SASH mice revealed a downregulation in pathways related to

cellular proliferation and division, such as "DNA replication", "cell cycle checkpoint signaling",

and "nuclear chromosome segregation" coupled with an increase in "negative regulation of cell

activation" within the M5 ontology gene set (BP) defined by the Gene Ontology consortium

(Figure 2C). This signature is consistent with our 5-FU data, demonstrating a quiescent

transcriptional signature within the lineage depleted portion of the BM. Moreover, the observed

upregulation in "leukocyte migration" and "myeloid leukocyte activation" pathways indicates

that, alongside a quiescent transcriptional profile, these cells are primed for WBC production

(Figure 2D). This suggests a balance between maintaining quiescence and the potential for

activation.

The SASH BM microenvironment is primed for hematopoietic engraftment.

We next investigated the frequency of HSC supportive cell types through immunophenotypic

characterization within the CD45$^-$/Ter119$^-$ BM population (gating strategy outlined in

Supplemental Figure 9). We found that leptin receptor positive cells (LepR$^+$), had a 1.4-fold

increase and adipocytes had a 1.9-fold increase in frequency within SASH BM (Figure 3A).

LepR+ cells, which comprise skeletal stem cells, along with osteogenic and adipogenic

progenitor cells are pivotal for their dual role in the niche; they not only contribute to the

physical scaffolding that determines the structural integrity of the niche but also secrete a range

of soluble factors that are vital for HSC maintenance, such as cytokines and growth factors.

Similarly, the increase in adipocytes, which are known to secrete adipokines and other regulatory

molecules, suggests an enriched environment for HSC support[31].

We next investigated a panel of different niche genes within the CD45⁻ population of the BM

(Supplemental Figure 4) and observed significant increases in angiopoietin-1 (*Angpt1*)

(important for maintaining a quiescent state within HSCs[32], and notably secreted by LepR+

cells[33]), *Vegfa* (important for HSC survival[34]), and *Il6* (needed for granulopoiesis[35]) (Figure 3B).

To determine the functional effects of the observed gene expression data, we performed

transplantation studies in which WT donor BM was transplanted either into WT or SASH

recipient mice. We sacrificed mice on day 20 following transplantation and found 1.2-fold higher

BM cellularity in SASH recipients compared to WT controls, 1.6-fold greater LSKs and trends

towards increased total LT-HSC (1.76-fold) (Figure 3C). This data suggests that even after

myeloablation, the SASH microenvironment is more favorable to hematopoietic engraftment and

thus peripheral reconstitution following transplantation compared to WT counterparts.

SASH mice contain lower histamine levels and histamine administration impairs

hematopoietic output.

To determine the role of histamine in modulating hematopoietic output in SASH mice we

compared the steady-state levels of serum histamine between SASH and WT mice and observed

that WT mice contained 1.6-fold higher histamine than SASH (Figure 4A). To further investigate

this connection, we assessed the expression of histidine decarboxylase (*Hdc*), the sole enzyme

that converts L-histidine to histamine, thus serving as a specific marker for the biosynthesis of

histamine[36]. We observed no significant differences in *Hdc* expression between SASH and WT

mice in both bulk BM and CD45⁻ populations, suggesting that the decrease in histamine levels

was due specifically to the loss of MCs (Figure 4B). Moreover, we investigated the expression

levels of *Hrh1*, *Hrh2*, and *Cysltr1* in both bulk and CD45⁻ populations in SASH and WT mice

(Supplemental Figure 5). Interestingly, we observed a decrease in the expression of *Hrh1* within

the CD45⁻ population of SASH BM, indicating reduced H1-mediated signaling in response to

histamine in these cells.

To explore the functional consequences of reduced histamine in SASH mice, we administered

exogenous histamine to these animals and evaluated its effects on peripheral blood counts and

BM HSPC frequencies. Treatment of SASH mice at 100 mg/kg histamine BID for 7 days

resulted in a 2.9, 2.6, and 3.1-fold reduction in WBC, NE, and LY respectively, demonstrating

that histamine supplementation reduces peripheral blood output in SASH mice (Figure 4C).

Notably, RBC and PLT counts were not altered by the 7-day histamine treatment (Supplemental

Figure 6). We tested a lower dose of 10 mg/kg and a 4-day blood analysis time point on both

doses but saw no alterations (Supplemental Figure 6). We also examined BM HSPCs and

observed a trend towards decreased cell populations only at the higher histamine dose (Figure 4D

and supplemental Figure 6). Together this data suggests histamine negatively regulates

hematopoietic function and the loss of MC derived histamine in SASH mice contributes to the

observed increase in hematopoietic output.

***In vivo* antihistamine treatment results in steady-state BM progenitor cell expansion.**

Having identified a potential causative role for histamine in driving the hematopoietic phenotype

in SASH mice, we sought to pharmacologically mimic this effect using antihistamines. Mice

were treated twice daily for 7 days with a selection of FDA-approved antihistamines.

Specifically, we administered 20 mg/kg of cetirizine (H1R inverse agonist[37]), 10 mg/kg of

famotidine (H2R antagonist[38]), 5 mg/kg of montelukast (leukotriene receptor antagonist[39]), and

10 mg/kg of ketotifen (H1R inverse agonist and MC stabilizer which prevents the release of

additional inflammatory mediators from MCs[40]). While additional studies will be required to

identify optimal dosing regimens, our initial findings revealed a statistically significant 1.7-fold

increase in both LSK and LT-HSC populations specifically following cetirizine treatment (Figure

5B). In addition, we performed a myeloid and lymphoid population analysis within the BM and

found that cetirizine induced significant changes including an increase in total myeloid cells, B

cells, and CD8$^+$ T cells (Figure 5C).

EHR data identifies association between antihistamine use and white blood cell counts.

To determine the clinical relevance of our murine data, we analyzed electronic health record

(EHR) data from TrinetX in patients diagnosed with allergy and prescribed antihistamines vs

other anti-allergy medications. The 'exposure risk' represents the proportion of patients taking

antihistamines who had elevated white blood cell counts. Conversely, the 'control risk' refers to

the proportion of patients taking other anti-allergy medications who had elevated WBC counts.

Antihistamines were associated with a significantly increased risk for elevated white blood cell

counts compared with other anti-allergy medications during a 1-year follow-up in patients with

allergy and no history of elevated white blood cell counts (HR: 2.09, 95% CI: 1.57-2.78),

suggesting a strong association in a relative short time follow-up (**Table 1**). While this data does

not establish causality, it intriguingly parallels our animal model results and prompts further

investigation into whether this association is indicative of a conserved biological mechanism. It

also provides a compelling link between antihistamine use and hematopoietic output in a clinical

setting, thereby reinforcing the translational relevance of our animal model findings.

Table 1. Outcome: elevated white blood cell count (follow up: 1 year)			
Cohort setting	Exposure risk	Control Risk	Hazard Ratio (95% CI)

Antihistamines group vs Decongestants and antiallergics group	1.15%	0.55%	2.09 (1.23,3.56)
Antihistamines group vs Corticosteroids group	0.58%	0.27%	2.09 (1.57,2.78)

DISCUSSION

This work elucidates a previously unexplored regulatory mechanism within the hematopoietic

system demonstrating how MCs and histamine signaling via H1R influence both HSC function

and overall hematopoietic output. We found that MC deficient SASH mice display leukocytosis,

expanded HSPC populations in the BM, and increased splenic extramedullary hematopoiesis.

Our transcriptomic analysis indicated that the lineage depleted BM is predisposed to increased

WBC output without compromising its quiescent state. Increased resistance to repeated 5-FU

treatment validated this predisposition. Further analyses of cell frequencies and gene signatures

within the BM niche suggested a more supportive microenvironment, which correlated with

improved engraftment following transplantation into SASH recipient animals.

Our work additionally posits that the reduced histamine levels observed in SASH mice

contribute to their hematopoietic phenotype. We demonstrate that administration of exogenous

histamine reverses the expanded peripheral blood output and BM HSPCs. Collectively these

findings indicate that histamine acts a negative regulator of hematopoietic function.

In addition to elucidating a novel mechanism in SASH mice by which MC derived histamine

regulates hematopoiesis, we also found that inhibition of histamine signaling with cetirizine

treatment, an H1R inverse agonist, resulted in significant expansion of HSPC populations in the

BM, alongside increased myeloid populations and lymphoid cell subsets. Importantly we also

identified a consistent result from retrospective analysis of electronic health record data, in which

patients taking antihistamines had significantly higher white blood cell counts than untreated

cohorts.

Our data suggests cetirizine could be utilized as a novel approach to promote hematopoietic

function. Repurposing FDA-approved therapies for alternative disease indications offers

advantages such as shorter development time (5-7 years), a higher rate of approval for the

alternative indication, a substantial reduction in development costs (ranging from 40-70%), and

access to a wealth of existing data from the extensive FDA approval process these compounds

have undergone[41]. Based on our findings, this work may have implications for several clinical

needs including:

-   Poor graft function (PGF)- a complication of allogeneic transplantation characterized by

unexplained persistent cytopenia despite full donor chimerism which occurs in 5 to 27

percent of patients and is associated with increased morbidity and mortality[42]. The

management of PGF is limited and often ineffective, typically involving supportive

treatments such as hematopoietic growth factors (like G-CSF), eltrombopag, and stem cell

reinfusion or boosts. By bolstering the HSC pool, cetirizine could effectively augment

hematopoietic regeneration, potentially mitigating the persistent cytopenia characteristic

of PGF and improving overall patient outcomes post-transplant.

-   Myelofibrosis- a chronic myeloproliferative disorder marked by progressive scarring of the

bone marrow, leading to severe anemia, weakness, fatigue, and often, splenomeglia[43]. MF

patients also display clonal hematopoiesis, which is characterized by HSC production of a

small population of dominant clones rather than a healthy polyclonal population, and which

is associated with a significantly higher risk of transformation[44]. While targeting MCs in

fibrotic disease has recently been proposed as a novel therapeutic strategy[45, 46], there is a

possibility that antihistamines may display dual efficacy in MF by acting to not only limit

clonal dominance but also reduce fibrotic onset.

Overall, this work uncovers a previously unrecognized role of MCs in regulating hematopoiesis
and provides insights into the underlying molecular mechanisms. Further investigations into the
precise mechanisms mediating MC-HSPC interactions are warranted, along with exploring the
therapeutic potential of MC-targeted interventions in additional settings of insufficient
hematopoiesis including chemotherapy induced neutropenia, various anemias, and transplantation
settings. This research not only contributes to a deeper understanding of the mechanisms governing
normal and regenerative hematopoiesis, but also demonstrates the potential of repurposing MC
targeting therapies as a cost effective method to promote hematopoietic function in the clinical
setting.

Acknowledgments

This work was supported by NIH grants R00 HL135740, RM1GM42002, R21AG075573, and
by the Radiation Resources Core Facility, the Translational Therapeutics Core Facility, the
Hematopoietic Biorepository and Cellular Therapy Core Facility, and the Cytometry &
Microscopy Core Facility of the Case Comprehensive Cancer Center (P30CA043703). This work
was also supported by a generous award from the Ohio Cancer Research Foundation. This
project was also supported by the Clinical and Translational Science Collaborative of Northern
Ohio, which is funded by the National Institutes of Health, National Center for Advancing
Translational Sciences, Clinical and Translational Science Award grant, UM1TR004528.

METHODS

Animals: Animals were housed in the AAALAC-accredited facilities of the CWRU School of Medicine. The Case Western Reserve University Institutional Animal Care and Use Committee (IACUC) approved the husbandry and experimental procedures in accordance with approved IACUC protocols 2019-0065. C-Kit[W-sh] mice were purchased from The Jackson Laboratory. Littermate wild-type control animals were used as comparator animals for all studies, and a combination of male and female mice, aged 8-12 weeks old, were used for all studies. Mice were housed in standard microisolator cages and maintained on a defined, irradiated diet and autoclaved water. All animals were observed daily for signs of illness. Animals that were used for tissue analysis were sacrificed humanely with CO_2 and then by cervical dislocation. Bone marrow was obtained by removing the tibia and femur and flushing the cells into FACS buffer. Splenocytes were obtained by removing the spleen and mincing them into a single-cell suspension.

Complete Blood Count Analysis: Peripheral blood was collected by submandibular bleeding into Microtainer EDTA tubes (Becton-Dickson cat # 365974). Blood counts were analyzed by a Hemavet 950 FS hematology analyzer.

Flow Cytometry: Cells were resuspended and stained with 200 µL of FACS with fluorescently labeled antibodies for 15 minutes and then fixed in 1% PFA for 15 minutes. Samples were then washed with FACS buffer and finally resuspended in 200 µL of FACS buffer. Data was acquired on an LSRII flow cytometer (BD Biosciences). Analysis was performed on FlowJo software (TreeStar).

5-FU Administration: C-Kit[W-sh] and WT mice were treated with 150 mg/kg of 5-FU (Sigma cat # 7F6627) in a 10% DMSO solution in PBS every seven days via IP injection. Mice were

monitored daily and sacrificed when they exhibited >25% weight loss or became moribund.

Animal survival was monitored throughout this study.

Histamine ELISAs: Peripheral blood was collected from mice and allowed to clot before

centrifugation to collect serum. Samples were frozen until day of assay. ELISA assays performed

using Abcam kits (cat # AB213975) according to manufacturer's protocol.

Histamine and Antihistamine Administration: Histamine dihydrochloride was solubilized

directly into PBS. Cetirizine dihydrochloride (Sigma cat # C3618), ketotifen fumarate salt (Sigma

cat # K2628), famotidine (Sigma cat # F6889), and montelukast sodium (Sigma cat # PHR1603)

were dissolved in DMSO at 60, 30, 30 and 15 mg/mL respectively. Each compound was then

diluted in water to a final concentration of 2, 1, 1, and 0.5 mg/mL respectively, diluting the DMSO

to 3.3%. Naïve SASH C57BL/6 were administered histamine and antihistamines respectively via

IP injection on a BID schedule (A.M. and P.M. administration) for 7 days and were sacrificed 1

hour after the 15^{th} dose.

Bone Marrow Transplantation: Mice were exposed to a single dose of 10 Gy total body

irradiation from a cesium source. 16-24 hrs later, mice received 1e6 whole bone marrow cells by

retroorbital injection. Recipients were monitored daily for signs of distress or illness and sacrificed

on day 20 to assess bone marrow and spleen cellularity in addition to cytometric assessment of

cell populations.

REFERENCES

1. Lee JY, Hong SH. Hematopoietic Stem Cells and Their Roles in Tissue Regeneration. Int J Stem Cells. 2020;13(1):1-12.

2. Nakamura-Ishizu A, Takizawa H, Suda T. The analysis, roles and regulation of quiescence in hematopoietic stem cells. Development. 2014;141(24):4656-4666.

3. Alenzi FQ, Alenazi BQ, Ahmad SY, Salem ML, Al-Jabri AA, Wyse RK. The haemopoietic stem cell: between apoptosis and self renewal. Yale J Biol Med. 2009;82(1):7-18.

4. Zhang CC, Sadek HA. Hypoxia and metabolic properties of hematopoietic stem cells. Antioxid Redox Signal. 2014;20(12):1891-1901.

5. Chow A, Lucas D, Hidalgo A, et al. Bone marrow CD169+ macrophages promote the retention of hematopoietic stem and progenitor cells in the mesenchymal stem cell niche. J Exp Med. 2011;208(2):261-271.

6. Mitroulis I, Kalafati L, Bornhauser M, Hajishengallis G, Chavakis T. Regulation of the Bone Marrow Niche by Inflammation. Front Immunol. 2020;11(1540.

7. Park D, Spencer JA, Koh BI, et al. Endogenous bone marrow MSCs are dynamic, fate-restricted participants in bone maintenance and regeneration. Cell Stem Cell. 2012;10(3):259-272.

8. Poulos MG, Guo P, Kofler NM, et al. Endothelial Jagged-1 is necessary for homeostatic and regenerative hematopoiesis. Cell Rep. 2013;4(5):1022-1034.

9. Mendelson A, Frenette PS. Hematopoietic stem cell niche maintenance during homeostasis and regeneration. Nat Med. 2014;20(8):833-846.

10. Chen X, Deng H, Churchill MJ, et al. Bone Marrow Myeloid Cells Regulate Myeloid-Biased Hematopoietic Stem Cells via a Histamine-Dependent Feedback Loop. Cell Stem Cell. 2017;21(6):747-760 e747.

11. Kitamura Y, Oboki K, Ito A. Development of mast cells. Proc Jpn Acad Ser B Phys Biol Sci. 2007;83(6):164-174.

12. Orfao A, Escribano L, Villarrubia J, et al. Flow cytometric analysis of mast cells from normal and pathological human bone marrow samples: identification and enumeration. Am J Pathol. 1996;149(5):1493-1499.

13. Schnedl WJ, Enko D. Histamine Intolerance Originates in the Gut. Nutrients. 2021;13(4):

14. Shulpekova YO, Nechaev VM, Popova IR, et al. Food Intolerance: The Role of Histamine. Nutrients. 2021;13(9):

15. Krystel-Whittemore M, Dileepan KN, Wood JG. Mast Cell: A Multi-Functional Master Cell. Front Immunol. 2015;6(620.

16. Gilfillan AM, Rivera J. The tyrosine kinase network regulating mast cell activation. Immunol Rev. 2009;228(1):149-169.

17. Moon TC, Befus AD, Kulka M. Mast cell mediators: their differential release and the secretory pathways involved. Front Immunol. 2014;5(569.

18. Kunder CA, St John AL, Abraham SN. Mast cell modulation of the vascular and lymphatic endothelium. Blood. 2011;118(20):5383-5393.

19. Varricchi G, Rossi FW, Galdiero MR, et al. Physiological Roles of Mast Cells: Collegium Internationale Allergologicum Update 2019. Int Arch Allergy Immunol. 2019;179(4):247-261.

20. Brooks AC, Whelan CJ, Purcell WM. Reactive oxygen species generation and histamine release by activated mast cells: modulation by nitric oxide synthase inhibition. Br J Pharmacol. 1999;128(3):585-590.

21. Nakamura Y, Ishimaru K, Shibata S, Nakao A. Regulation of plasma histamine levels by the mast cell clock and its modulation by stress. Sci Rep. 2017;7(39934.

22. Neumann J, Grobe JM, Weisgut J, et al. Histamine can be Formed and Degraded in the Human and Mouse Heart. Front Pharmacol. 2021;12(582916.

23.   Borriello F, Iannone R, Marone G. Histamine Release from Mast Cells and Basophils. Handb Exp
Pharmacol. 2017;241(121-139.
24.   Leurs R, Church MK, Taglialatela M. H1-antihistamines: inverse agonism, anti-inflammatory actions
and cardiac effects. Clin Exp Allergy. 2002;32(4):489-498.
25.   Simons FE, Simons KJ. H1 antihistamines: current status and future directions. World Allergy Organ
J. 2008;1(9):145-155.
26.   Grimbaldeston MA, Chen CC, Piliponsky AM, Tsai M, Tam SY, Galli SJ. Mast cell-deficient W-sash c-
kit mutant Kit W-sh/W-sh mice as a model for investigating mast cell biology in vivo. Am J Pathol.
2005;167(3):835-848.
27.   Lotinun S, Krishnamra N. Disruption of c-Kit Signaling in Kit(W-sh/W-sh) Growing Mice Increases
Bone Turnover. Sci Rep. 2016;6(31515.
28.   Michel A, Schuler A, Friedrich P, et al. Mast cell-deficient Kit(W-sh) "Sash" mutant mice display
aberrant myelopoiesis leading to the accumulation of splenocytes that act as myeloid-derived suppressor
cells. J Immunol. 2013;190(11):5534-5544.
29.   Carnevalli LS, Scognamiglio R, Cabezas-Wallscheid N, et al. Improved HSC reconstitution and
protection from inflammatory stress and chemotherapy in mice lacking granzyme B. J Exp Med.
2014;211(5):769-779.
30.   Schepers K, Hsiao EC, Garg T, Scott MJ, Passegue E. Activated Gs signaling in osteoblastic cells alters
the hematopoietic stem cell niche in mice. Blood. 2012;120(17):3425-3435.
31.   Pinho S, Frenette PS. Haematopoietic stem cell activity and interactions with the niche. Nat Rev
Mol Cell Biol. 2019;20(5):303-320.
32.   Arai F, Hirao A, Ohmura M, et al. Tie2/angiopoietin-1 signaling regulates hematopoietic stem cell
quiescence in the bone marrow niche. Cell. 2004;118(2):149-161.
33.   Zhou BO, Ding L, Morrison SJ. Hematopoietic stem and progenitor cells regulate the regeneration
of their niche by secreting Angiopoietin-1. Elife. 2015;4(e05521.
34.   Gerber HP, Malik AK, Solar GP, et al. VEGF regulates haematopoietic stem cell survival by an
internal autocrine loop mechanism. Nature. 2002;417(6892):954-958.
35.   Sasaki Y, Guo YM, Goto T, et al. IL-6 Generated from Human Hematopoietic Stem and Progenitor
Cells through TLR4 Signaling Promotes Emergency Granulopoiesis by Regulating Transcription Factor
Expression. J Immunol. 2021;207(4):1078-1086.
36.   Haak-Frendscho M, Darvas Z, Hegyesi H, et al. Histidine decarboxylase expression in human
melanoma. J Invest Dermatol. 2000;115(3):345-352.
37.   Shimizu T, Nishihira J, Watanabe H, Abe R, Ishibashi T, Shimizu H. Cetirizine, an H1-receptor
antagonist, suppresses the expression of macrophage migration inhibitory factor: its potential anti-
inflammatory action. Clin Exp Allergy. 2004;34(1):103-109.
38.   Dizaj KA, Monfared AS, Mozdarani H, et al. Combined effect of oral famotidine and cimetidine on
the survival of lethally irradiated mice: An in vivo study. J Cancer Res Ther. 2021;17(4):865-869.
39.   Roa J, Morikawa H, Crawford L, Baatjes A, Duong M, Denburg JA. The effects of montelukast on
tissue inflammatory and bone marrow responses in murine experimental allergic rhinitis: interaction with
interleukin-5 deficiency. Immunology. 2007;122(3):438-444.
40.   Hu Y, Jin Y, Han D, et al. Mast cell-induced lung injury in mice infected with H5N1 influenza virus. J
Virol. 2012;86(6):3347-3356.
41.   Pushpakom S, Iorio F, Eyers PA, et al. Drug repurposing: progress, challenges and
recommendations. Nat Rev Drug Discov. 2019;18(1):41-58.
42.   Lv WR, Zhou Y, Xu J, et al. Haploidentical donor transplant is associated with secondary poor graft
function after allogeneic stem cell transplantation: A single-center retrospective study. Cancer Med.
2021;10(23):8497-8506.

43.   Jacquelin S, Kramer F, Mullally A, Lane SW. Murine Models of Myelofibrosis. Cancers (Basel).
2020;12(9):

44.   Jaiswal S, Ebert BL. Clonal hematopoiesis in human aging and disease. Science. 2019;366(6465):

45.   Bradding P, Pejler G. The controversial role of mast cells in fibrosis. Immunol Rev. 2018;282(1):198-
231.

46.   Overed-Sayer C, Miranda E, Dunmore R, et al. Inhibition of mast cells: a novel mechanism by which
nintedanib may elicit anti-fibrotic effects. Thorax. 2020;75(9):754-763.

Figure 1: MC deficient Kit$^{W\text{-}sh/W\text{-}sh}$ "SASH" mice display leukocytosis and BM HSPC expansion

Figure 1: A. Complete blood count analysis of circulating WBC, MO, NE, and PLT at steady-state in WT and SASH mice. N=24 mice/group, except monocytes was N=10 mice/group. **B.** Immunophenotypic analysis of HSPCs and LSKs per hindlimb. N=9 mice/arm. Total nucleated bone marrow cells per hindlimb were obtained. N=9 mice/group. **C.** Survival curve of mice subjected to weekly 150 mg/kg injections of 5-FU. N=15 mice/group. **A, B.** Error bars represent SEM. Students two-tailed T test was used for statistical analysis with p>0.05=*, p>0.01=**, p>0.001=***,p>0.0001=****

Figure 2: SASH lineage depleted BM exhibit transcriptional signatures indicative of HSPC quiescence

Figure 2: A. Schematic depicting the experimental design for RNA sequencing study. **B.** Principal component analysis (PCA) plot of SASH and WT bulk and lineage depleted populations. **C,D.** Heatmap of top differentially regulated genes between SASH and WT mice showing **C.** pathways related to cellular activation and replication **D.** pathways related to WBC activation and migration. N=5 mice/group.

Figure 3: The SASH BM microenvironment is primed for hematopoietic engraftment

Figure 3: A. Flow cytometric analysis of BM stromal populations. N=10 mice/arm. **B.** RT-PCR analysis of niche factor genes within BM stromal cells. N=10 mice/arm. **C.** BM cellularity and quantification of LSK and LT-HSC cells per mouse (2 hindlimbs) of recipients 20 days post-transplant. N=18-22 mice/arm. **A-C.** Error bars represent SEM. Students two-tailed T test was used for statistical analysis with p>0.05=*, p>0.01=**, p>0.001=***, p>0.0001=****

Figure 4: SASH mice contain lower histamine levels and histamine administration impairs hematopoietic output

Figure 4: A. ELISA analysis of histamine concertation in the serum of WT and SASH mice. N=8-9 mice/arm. **B.** RT-PCR analysis of HDC and H1R within bulk and stromal BM cells. N=4-5 mice/arm. **C.** Complete blood count analysis of circulating WBC, NE, LY, and MO after 7 days of BID 100 mpk histamine administration. N=4-5 mice/group. **D.** Immunophenotypic analysis of HSPCs and LSKs per hindlimb after 7 days of BID 100 mpk histamine administration. N=4-5 mice/arm. **A-D.** Error bars represent SEM. Students two-tailed T test was used for statistical analysis with p>0.05=*, p>0.01=**, p>0.001=***,p>0.0001=****

Figure 5: *In vivo* antihistamine treatment results in BM progenitor cell expansion at steady-state

Figure 5: A. Schematic depicting the mechanism of action for FDA-approved drugs that target histamine and leukotriene receptors and for drugs that stabilize MCs. **B.** Immunophenotypic analysis of HSPCs and LSKs per hindlimb of WT mice treated BID with 2, 1, 0.5, and 1 mg/kg of cetirizine, famotidine, montelukast, and ketotifen respectively. N=30 mice/arm. **C.** Immunophenotypic analysis of myeloid, B and T cells per hindlimb of the same mice in B. N=30 mice/arm (15/sex). **B, C.** Error bars represent SEM. Students two-tailed T test was used for statistical analysis with p>0.05=*, p>0.01=**, p>0.001=***,p>0.0001=****