Title: Priority effects determine how dispersal affects biodiversity in seasonal metacommunities

Authors: Heng-Xing Zou[1], Volker H. W. Rudolf[2]

ORCIDs: [1] 0000-0002-1965-6910, [2] 0000-0002-9214-2000

Address: [1,2] Program in Ecology and Evolutionary Biology, BioSciences, Rice University, 6100 Main St, Houston, TX 77005

Corresponding Author: Zou, H.-X. (hengxingzou@rice.edu)

Running title: Priority Effects in Metacommunities

Keywords: priority effects, metacommunity, dispersal-diversity relationships, phenology, seasonal variation

Manuscript type: Major article

Number of words in the abstract: 189

Number of words in the main text: 5338

Number of references: 62

Number of figures: 5

Number of tables: 1

Abstract

The arrival order of species frequently determines the outcome of their interactions. This

phenomenon, called the priority effect, is ubiquitous in nature and determines local community

structure, but we know surprisingly little about how it influences biodiversity across different

spatial scales. Here, we use a seasonal metacommunity model to show that biodiversity

patterns and the homogenizing effect of high dispersal depend on the specific mechanisms

underlying priority effects. When priority effects are only driven by positive frequency

dependence, dispersal-diversity relationships are sensitive to initial conditions but generally

show a hump-shaped relationship: biodiversity declines when dispersal rates become high and

allow the dominant competitor to exclude other species across patches. When spatiotemporal

variation in phenological differences alters species' interaction strengths (trait-dependent

priority effects), local, regional, and temporal diversity are surprisingly insensitive to variation

in dispersal, regardless of the initial numeric advantage. Thus, trait-dependent priority effects

can strongly reduce the effect of dispersal on biodiversity, preventing the homogenization of

metacommunities. Our results suggest an alternative mechanism that maintains local and

regional diversity without environmental heterogeneity, highlighting that accounting for the

mechanisms underlying priority effects is fundamental to understanding patterns of biodiversity.

Introduction

Dispersal promotes the exchange of individuals between communities and thus links local population and community dynamics to regional patterns of species distributions. As a consequence, dispersal can play a key role in shaping biodiversity patterns at both local and regional scales (Kerr et al. 2002; Cadotte 2006; Grainger and Gilbert 2016). However, the relationship between dispersal and biodiversity patterns is also shaped by local conditions and processes, including biotic interactions (e.g. Kneitel and Miller 2003; Shurin et al. 2004; Altermatt et al. 2011; Carrara et al. 2012; Vanschoenwinkel et al. 2013; Catano et al. 2017). Although recent studies identify local competitive interactions as a major contributor to dispersal-diversity relationships (Pu and Jiang 2015; Lu 2021; Miller and Allesina 2021), one ubiquitous biotic process in metacommunities, the temporal sequence of community assembly, has rarely received attention.

Communities are not assembled all at once; instead, species arrive at different times and differences in the sequence of arrival times can alter the outcome of species interactions and thereby also community composition. Such priority effects (or historical contingencies) are widespread in nature and occur in systems ranging from microbes to plants and vertebrates (Alford and Wilbur 1985; Drake 1991; Kardol et al. 2013; Fukami 2015). In a metacommunity context, priority effects become important because they can alter the fate of immigrating individuals into a patch. For instance, with an early arriver advantage, individuals who arrive at a patch may either successfully establish themselves if they arrive before competitors or perform poorly and even be excluded if they arrive after competitors. By altering the outcome of local competitive interactions, priority effects have the potential to shape the relationship between dispersal and diversity patterns at local and regional scales. Although several theories

predict that priority effects could affect how regional dispersal shapes biodiversity (e.g. Shurin

et al. 2004; Fukami 2015; Grainger and Gilbert 2016; Miller and Allesina 2021), few have

incorporated the diverse mechanisms that can generate priority effects in natural systems. Yet

accounting for these mechanisms may be key to explaining why dispersal-diversity

relationships are often contrary to theoretical expectations (e.g., Pu and Jiang 2015; Vannette

and Fukami 2017; Toju et al. 2018).

Priority effects can be generated by at least two fundamentally different mechanisms,

broadly categorized as "frequency-dependent" and "trait-dependent". Traditionally, frequency-

dependent priority effects have received the most attention and dominate the theoretical

literature. They arise when fitness differences between species are small and destabilizing

differences are large (i.e., positive frequency-dependent growth rates; Ke and Letten 2018). If

conditions are met, the system exhibits alternative stable states, where neither species can

invade when rare. Arrival time itself does not affect any per-capita rates and only matters if

differences in arrival time allow for shifts in the relative frequency of species when interactions

start. In contrast, trait-dependent priority effects arise when per-capita interactions between

species change with their relative arrival times (Rudolf 2019). In nature, this occurs when

differences in arrival time are correlated with changes in traits that affect species interactions,

such as size and behavior, or modified environment from the difference in arrival times (Poulos

and McCormick 2014; Rasmussen et al. 2014; Rudolf 2018; Blackford et al. 2020). In contrast

to frequency-dependent priority effects, trait-dependent priority effects shift what outcomes are

possible (e.g., from coexistence to competitive exclusion) by altering the parameter space (e.g.,

changes in interspecific competition) itself instead of favoring one of two possible states

(Rudolf 2019; Zou and Rudolf 2020; Fragata et al. 2022).

These two mechanisms or priority effects are not mutually exclusive but operate at

different temporal scales. Frequency-dependent priority effects require that arrival times are

separated by several generations to allow populations to reproduce and increase before

competitors arrive (Figure 1A; Fukami 2004; Grainger et al. 2019). Trait-dependent priority

effects typically occur when arrival times are separated by less than one generation, e.g., when

phenological differences between competitors lead to differences in traits such as body size

(Figure 1B; Shorrocks and Bingley 1994; Rudolf 2018; Blackford et al. 2020). This distinction

is especially important in a seasonal context where both can occur. Consider a community

where species complete one generation per year (Figure 1C). Here, frequency-dependent

priority effects can arise when one species colonizes a patch several years before others (a

"first colonizer"; Figure 1C), but a later colonizing species may have earlier phenology (e.g.,

earlier germination, emergence from dormancy, or reproduction) and thus become "active" first

within a year, allowing for trait-dependent priority effects in a year (an "early emerger"; Figure

1D). Therefore, both the different timing between and within years can determine the local

outcomes of competition.

These differences in priority effects in local patches have the potential to alter dispersal-

diversity relationships. With frequency-dependent priority effects, positive frequency

dependence could enable the more abundant first colonizer to spread over the landscape

through dispersal and competitively exclude late arrivers. High dispersal will therefore

decrease both local and regional diversity because one or a few regionally dominant species

will eventually establish and maintain a numeric advantage in all patches, similar to the classic

prediction that dispersal homogenizes (Tilman 1994; Mouquet and Loreau 2003; Leibold et al.

2004; Vanoverbeke et al. 2016; Fodelianakis et al. 2019). Following this premise, alpha (local)

diversity is likely highest at intermediate dispersal while beta (dissimilarity between patches)

and gamma (regional) diversity likely decreases with dispersal (Mouquet and Loreau 2002,

2003; Cadotte 2006). However, regional diversity may be retained even under high dispersal if

priority effects are driven by trait-dependent mechanisms, such as when early species alter the

environment or rapidly adapts to the habitat (Shurin et al. 2004; Fukami 2015; Grainger and

Gilbert 2016; Leibold et al. 2019; Miller and Allesina 2021). In these cases, outcomes of

competition are not only driven by relative abundances but also by within-year arrival times, or

phenology. Early emergers therefore will not be excluded even if other species are more

abundant. Consequences of this process should be particularly important in seasonal systems,

as the frequent, periodic assembly of local communities creates many opportunities for spatial

and temporal variations in arrival orders (Sheriff et al. 2011; Diez et al. 2012; Theobald et al.

2017; Carter et al. 2018; Rudolf 2019). In these seasonal systems, trait-dependent priority

effects and natural variations in phenology could lead to high temporal turnover of local and

regional community composition despite the homogenizing effect of dispersal.

Here, we use a spatially implicit, multispecies metacommunity model to reveal how

different mechanisms of priority effects interact with dispersal to shape biodiversity at local

and regional scales in competitive metacommunities. We modeled a seasonal system with

spatial and temporal variations in annual emergence times (representing species phenology)

and compared the observed dispersal-diversity relationships with and without trait-dependent

priority effects across different initial distributions of the population. We asked the following

questions: (1) How will different mechanisms of priority effects change dispersal-diversity

relationships? (2) When will local and regional diversity be maintained under high dispersal?

Methods

Model Setup

We modeled natural habitats with seasonal disturbance, such as temperate grassland or

ephemeral ponds, in which species survive periodic, unfavorable environments (e.g., winter,

droughts) and reemerge at the beginning of the next growing season. We define a growing

season as the growth period between a previous periodic disturbance to the next disturbance. At

the beginning of the growing season (hereafter season for short), each species emerges from

dormancy in a patch at an assigned time (phenology). Patches are homogeneous throughout the

landscape and are not spatially structured.

Within each season, local population dynamics of species i in a given patch are modeled

with a modified Beverton-Holt model (Rudolf 2019):

$$N_i'(t) = \frac{\lambda_i N_i(t)}{1+\sum_{j=1}^{n} \alpha_{ij}\lambda_i N_j(t)} \text{ (Eqn. 1)},$$

where the population at the end of time step t (before dispersal and end-of-season disturbance),

$N_i'(t)$, is determined by population growth of species i, $\lambda_i N_i(t)$, where λ_i is the intrinsic

growth rate and $N_i(t)$ is the population of species i at the beginning of the season (after

dispersal and disturbance of the previous season). Population growth is dependent on densities

of all species with respective pairwise competition coefficients α_{ij}. Note that this process

assumes one reproduction event per season, which applies to systems such as annual plants or

amphibians.

If the community is driven by trait-dependent priority effects, interspecific competition is a

function of the relative emergence time:

$$\alpha_{ij}(\Delta p_{it}) = \frac{B}{1+exp(\Delta p_{it}/c)} \text{(Eqn. 2)}$$

where Δp_{it} is the difference between actual emergence times between species i and j at the start

of season t, calculated as $p_{jt} - p_{it}$; $\Delta p_{it} > 0$ when species i emerges first, and vice versa. B

and c are constants that determine the asymptotic maximum and shape of the sigmoidal

function respectively (Figure 2). When two species arrive simultaneously ($\Delta p_{it} = 0$), the

interspecific competition coefficient is $B/2$. This nonlinear competition-phenology function

has been validated by several empirical studies in animal and plant systems (Shorrocks and

Bingley 1994; Rudolf 2018; Blackford et al. 2020). With trait-dependent priority effects, α_{ij} is

calculated based on the respective Δp_{it} of a given patch at season t. In the absence of trait-

dependent priority effects, interspecific competition does not depend on the relative emergence

time. We therefore calculate α_{ij} as if $\Delta p_{it} = 0$: $\alpha_{ij}(0) = B/2$. Using Eqn. 2 in this way ensures

the consistency between the presence and absence of trait-dependent priority effects. We

created a competitive hierarchy across all five species by differentiating intraspecific

competition (intensity of self-limitation). For each simulation, we drew intraspecific

competition coefficients from a normal distribution around a mean (Table 1).

At the end of the season, species disperse in the metacommunity, and all patches are

subject to an end-of-season disturbance. We modeled the dispersal process as follows: first,

emigrating populations of each species in each patch are determined by the number of

successes of a binomial draw with $N_i'(t)$ (rounded to the nearest integer) trials and probability

r (Thompson et al. 2020), and r represents the dispersal rate. Emigrants are then pooled and

randomly redistributed to all patches. This spatially implicit method assumes that dispersal is

not affected by distance or patch choice. Spatially implicit models have been widely used in

theoretical studies (e.g. Mouquet and Loreau 2002; 2003) and represent a group of empirical

studies that manually transfer individuals between patches (Pu and Jiang 2015; Grainger and

Gilbert 2016). We modeled the end of the season by a reduction in survival (s_i), as it usually

represents a period when individuals face harsh environmental conditions (drought, cold). We

allowed s_i to vary stochastically across seasons and patches by drawing it from a normal

distribution with a mean of 0.5 and a standard deviation of 0.05. Each species then emerges at

its assigned times in the next season.

Generating Emergence Times

We randomly selected the emergence time (phenology) of species i from a normal

distribution with mean $\overline{p}_i$ and standard deviation v. We separated $\overline{p}_i$ of each species by a

constant interval along a gradient, with species 1 emerging on average the earliest and species

5 the latest (Figure 2). The average phenology $\overline{p}_i$ of each species is fixed, meaning that on

average, the order of emergence within a season is fixed. However, the actual phenology is

drawn from a normal distribution with average $\overline{p}_i$ and standard deviation v, allowing for shifts

in the order of emergence. We avoided negative phenology values by normalizing the earliest

phenology to 0. We repeated this process each season and for each patch and species (Figure

S1). If $v > 0$, this process creates variations in the phenology of co-occurring species and

mimics the environmental stochasticity across space and time in natural systems (Sheriff et al.

2011; Diez et al. 2012; Theobald et al. 2017; Carter et al. 2018).

Based on field and experimental data, the phenological variation (v) can be up to 5-10

times the pairwise phenological differences between interacting species (Menzel et al. 2006;

Kharouba et al. 2018; Rudolf 2018). Because emergence times are in a sequence with fixed

intervals in our model, the pairwise differences depend on the specific species combination. To

encompass possible scenarios of natural variation, we explored a wide range of v and picked

three representative values, 0.5 (smallest pairwise difference in $\overline{p}_i$), 2 (largest pairwise

difference in $\overline{p}_i$), and 8 (four times the largest pairwise difference in $\overline{p}_i$), corresponding small,

medium, and large variations (Table 1). We focused on results with $v = 2$.

Initial Scenarios

To identify the different and combined effects of each type of priority effects (frequency-

vs. trait-dependent) we examined the model under two different starting scenarios. The first

scenario starts each patch with all five species at the same initial population ("Equal Initials").

The second scenario assumes that the first arriving species colonizes and establishes in a patch

for several seasons before others arrive ("First Colonizer"). We created the latter scenario by

running simulations with only one species per patch and no dispersal for 10 seasons (burn-in),

with each species occupying equal numbers of patches. This ensures that the first colonizer or

the "resident" species reached quasi-equilibrium abundance in the patch and only varied over

time due to stochastic end-of-season survival rates before we allowed dispersal (Table 1).

We intentionally selected ranges of intra- vs. interspecific competition coefficients to allow

frequency-dependent priority effects to occur in all scenarios (Table 1). However, the "First

Colonizer" scenario emphasizes this frequency-dependent priority effect by allowing the initial

colonizer to gain a numeric advantage over rare immigrants, while the "Equal Initial" scenario

establishes a baseline that only captures the effect of seasonal variations in abundance, which is

present in all cases but is not the focus of our model. We then compared the resulting

population dynamics and biodiversity measures with and without trait-dependent priority

effects under each scenario, resulting in a total of four different starting conditions.

Measuring Biodiversity

We examined dispersal-diversity relationships by calculating alpha, beta, and gamma

diversity of the metacommunity at the end of each simulation. We calculated alpha and gamma

diversity using the Simpson's index on local patches and the whole metacommunity

respectively. Alpha diversity was averaged among all patches of a metacommunity. We

quantified beta diversity (dissimilarity in community composition based on numeric

abundances of species) by averaging all pairwise Bray-Curtis distances between patches. We

calculate these metrics for each of the 50 simulations of a given parameter combination using

respective functions in the R package *vegan* (Oksanen et al. 2013). We also calculated temporal

beta diversity to quantify changes in metacommunity composition over time using the R

package *adespatial* (Dray et al. 2018). Specifically, we calculated the dissimilarity (*D* metric in

Legendre 2019) in species composition of a metacommunity between consecutive time steps

separated by 10 seasons, excluding the first 10 seasons as burn-in. A high dissimilarity index

indicates that relative abundances of species vary greatly over time.

Model Simulations

Although our model can be extended to any number of species and patches, we simulated

metacommunities with 5 and 10 competing species and 50 patches. We focus on results with 5

species here to highlight the individual populations within each patch. Each simulation was run

for 110 seasons, including a 10-season burn-in period; patterns such as extinction and

dominance qualitatively stabilized within 25-50 seasons. We ran 50 simulations under the same

set of parameters.

All simulations were performed in R version 4.1.1 (R Core Team 2021). Code and data

were uploaded to Dryad (Zou and Rudolf 2023).

Results

Overall, we observed remarkable differences in population dynamics between scenarios

with and without trait-dependent priority effects. Without trait-dependent priority effects,

dispersal-diversity relationships are consistent with classic predictions but are highly dependent

on initial scenarios (Equal Initials or First Colonizer). With trait-dependent priority effects,

high dispersal does not lower alpha, beta, and gamma diversity in metacommunities regardless

of the initial scenarios.

Scenario 1: Equal Initials

The local population dynamics immediately highlight the difference with and without trait-

dependent priority effects. Without trait-dependent priority effects, all species have an

approximately equal chance of dominating a patch because the competitive hierarchy is

randomly determined for each simulation. This pattern is not affected by dispersal rates, which

only increases the fluctuation of populations (Figure 3). We observed that the species with the

largest regional population has also the highest abundance in almost all patches, indicating both

local and regional dominance that increases with dispersal rates (Figure S1). We also found a

sharp decline in the temporal turnover of local communities, indicating fast competitive

exclusion (Figure S2). These observations clearly show that higher dispersal promotes

competitive exclusion and therefore homogenizes the metacommunity, which is consistent with

the previous theory.

In contrast, with trait-dependent priority effects, immigrants can successfully establish and

potentially become the most abundant species in patches even at the lowest level of dispersal

rates (Figure 3). This establishment is possible because the relative timing of emergence in a

season determines per-capita effects and thus competitive dominance. Consequently, the

species with the largest regional population does not always have the highest abundance in all

patches because the rank of overall population size does not correspond to the strengths of self-

limitation (Figure S1). Rather, the species that emerges the earliest also generally has the

largest population (Figure S3). This also means that any species that is competitively excluded

from a patch can always recolonize given sufficient seasonal variations in phenology. This

recolonization is reflected in a consistently high temporal turnover in community composition

within each patch (Figure S2).

Without trait-dependent priority effects, we found a weak hump-shaped pattern between

alpha, beta, and gamma diversity and dispersal rates in metacommunities (Figure 5). Beta

diversity is low at low dispersal rates because all local communities started with the same

composition. The low alpha and gamma diversity at low dispersal rates reflects the eventual

local dominance of the species with the least self-limitation. In contrast, alpha diversity

increases with dispersal rates in metacommunities with trait-dependent priority effects, beta

diversity remains largely constant and is not influenced by dispersal rates, and gamma diversity

slightly increases with higher dispersal under trait-dependent priority effects (Figure 5). These

striking differences correspond to observations of local population dynamics.

Scenario 2: First Colonizer

Like Scenario 1, population dynamics with and without trait-dependent priority effects are

remarkably different. Without trait-dependent priority effects, the first colonizer remains at

high abundance in a patch at lower dispersal rates while the population of other immigrating

species remains small (Figure 4; Figure S1). This pattern only changes when dispersal rates are

so high that immigrating species are abundant enough to overcome the numeric advantage of

the first colonizer; once this condition is met, immigrants can invade and exclude the resident

(Figure 4; Figure S1). These observations are captured by temporal turnover: low dispersal

rates lead to little temporal turnover of local communities that reflects the stochastic "noise" of

random dispersal events. With higher dispersal rates, temporal turnover is initially high but

quickly decreases to low levels because the species with the least self-limitation spreads within

the metacommunity, driving others to local extinction (Figure S2). With trait-dependent priority

effects, observed patterns are very similar to that of Scenario 1: the local community

composition fluctuates over time (Figure S2), but the species with the earliest emergence time

on average (species 1) still has the largest total population (Figure 4; Figure S4). These

observations indicate that with trait-dependent priority effects, the community dynamics

depend less on relative abundances but more on variations of emergence times.

These differences in within-patch dynamics strongly affect the local and regional dispersal-

diversity relationships with and without trait-dependent priority effects (Figure 5). Without

trait-dependent priority effects, alpha diversity shows a clear hump-shaped relationship with

dispersal rates. Beta diversity starts highest due to both the numeric advantage and low

dispersal but decreases with higher dispersal rates, indicating that dispersal homogenizes the

metacommunity and leads to regional dominance of a single species. Gamma diversity

decreases rapidly at higher dispersal rates as species are driven to extinction by competitive

exclusion. However, with trait-dependent priority effects, all dispersal-diversity relationships

are strikingly similar to those observed under Scenario 1 (compare left and right panels of

Figure 5): alpha diversity increases, beta diversity slightly decreases, and gamma diversity

remains the same with dispersal. Note that compared to metacommunities without trait-

dependent priority effects, beta diversity is lower at low dispersal with trait-dependent priority

effects because immigrants can establish in patches occupied by the first colonizer, increasing

the similarity of local community composition. Overall, this shows that systems with trait-

dependent priority effects are not sensitive to initial conditions, which is the opposite of

systems with only frequency-dependent priority effects.

Sensitivity Analyses

We simulated the main model with 10 species and found qualitatively the same dispersal-

diversity patterns, indicating the results reflect the dynamics of more diverse systems (Figure

S5). Additionally, we simulated the main model with smaller and larger variations around

average phenology (i.e., the magnitude of v). We found that the exact dispersal-diversity

relationships are strongly dependent on the magnitude of phenological variations: alpha, beta,

and gamma diversity are positively correlated with phenological variation (v) (Figure S6, S7).

This positive correlation arises because high variation increases the probability that realized

phenological differences vary over time and space, thereby reducing the competitive

dominance of the species which emerges 1^{st} on average. However, we still observed a

consistent difference between biodiversity patterns with and without trait-dependent priority

effects regardless of specific levels of phenological variations.

We then simulated the main model with five species but with two higher ranges of

intraspecific competition that favor coexistence rather than frequency-dependent priority

effects. We observed qualitative changes in both population dynamics and dispersal-diversity

relationships without trait-dependent priority effects, but results with trait-dependent priority

effects are robust (Supplementary Material, Section II). Finally, to further determine the

robustness of our results we relaxed each of the following assumptions of our model: (1) each

season consists of one generation, (2) dispersal happens once at the end of a season, (3) all

species have equal dispersal rates (r), and (4) no additional spatial disturbance. Overall, we did

not find a qualitative change, as we still observed remarkable differences in population

dynamics and dispersal-diversity relationships between mechanisms of priority effects (results

not shown).

Discussion

Priority effects are ubiquitous in natural systems, yet their role in shaping biodiversity

patterns is still poorly understood (Fukami 2015). Here we show that priority effects can play a

key role in shaping the local and regional biodiversity of seasonal communities and how they

are influenced by dispersal. However, the shape of this dispersal-diversity relationship depends

on what type of priority effects are present. Metacommunities with only frequency-dependent

priority effects are highly sensitive to the initial conditions and exhibit high spatial diversity at

low dispersal rates but become homogenized as dispersal increases. In contrast, trait-dependent

priority effects promote local and regional species coexistence even under very high levels of

dispersal and the patterns are highly robust to variation in initial conditions. Together, these

results provide novel insights into how priority effects shape the dispersal-diversity

relationships in nature and highlight the importance of considering the seasonal nature of local

community assembly to predict regional patterns.

Positioning Priority Effects in Current Dispersal-Diversity Framework

Classic models predict that the dispersal-diversity relationship should be maximized at

intermediate dispersal, while between-patch and regional diversity should constantly decrease

with dispersal (Mouquet and Loreau 2002; 2003). Diversity peaks at intermediate dispersal

because species are present in both source and sink habitats, and the population in the latter can

be rescued by immigration from the former (Leibold et al. 2004). Traditionally, source-sink

dynamics are expected to be driven by spatial heterogeneity in habitat conditions (Pulliam

1988). However, we still observed source-sink dynamics even in the absence of any

environmental heterogeneity. In our model, the species with the highest population in a patch

poses a barrier for rare immigrants to establish themselves. These patches serve as sources for

the resident and sinks for all other (immigrating) species. At low to moderate dispersal rates,

frequency-dependent priority effects thus maintain high beta diversity within metacommunities.

At high dispersal rates, species with the least self-limitation can reach patches in large numbers

and overcome frequency-dependent priority effects. The source-sink dynamics dissolve, and

the homogenization effect of dispersal decreases local and regional diversity (Mouquet and

Loreau 2003; Grainger and Gilbert 2016; Catano et al. 2017). Our results agree with several

modeling studies involving patch preemption, which prevents immigrants from colonizing

patches occupied by previous or current residents. Here, patch preemption can be considered a

source of environmental heterogeneity that can arise from trait-dependent priority effects via

habitat modification. Past models show that this mechanism promotes regional coexistence (Yu

and Wilson 2001; Calcagno et al. 2006; Miller and Allesina 2021).

Furthermore, we found that the shape of dispersal-diversity relationship is highly sensitive

to initial conditions (i.e., the presence or absence of initial numeric advantage) but only in

metacommunities without trait-dependent priority effects. Notably, our scenario where all

species start at the same densities (Scenario 1) coincides with a design often used in lab

experiments (Grainger and Gilbert 2016), but the resulting population dynamics and dispersal-

diversity relationships are different from those obtained under a scenario where a species was

allowed to establish in a patch before others could colonize (Scenario 2). This result is not

previously emphasized in theory (but see Lu 2021) and could at least partly help explain why

the diversity-stability relationship differs among studies that use different initial settings

(Grainger and Gilbert 2016). We did not observe this sensitivity in models with trait-dependent

priority effects because the high spatiotemporal variation in interspecific competition

determined population dynamics, masking any effect of initial conditions.

On the contrary, diversity patterns in metacommunities with trait-dependent priority effects

are much less affected by dispersal or initial conditions. In these metacommunities, immigrants

may successfully colonize a patch regardless of their numeric disadvantage as their

colonization success is more dependent on the order of emergence. If an immigrating species

emerges early enough within a season, this advantage (relative increase in interspecific

competitive ability) can overcome its numeric disadvantage toward the resident species,

allowing it to establish in the patch. Given the temporal and spatial variation in within-season

emergence times (phenology), trait-dependent priority effects can promote local and regional

coexistence of species by preventing competitive exclusion and homogenization at a high

dispersal rate. Indeed, smaller phenological variations lead to a more definitive competitive

hierarchy, lowering local and regional biodiversity. Our results suggest that seasonality and

phenological variations are important factors promoting coexistence in spatial communities, in

addition to other classic mechanisms such as spatial heterogeneity (Leibold et al. 2004;

Schreiber and Killingback 2013).

The trait-dependent priority effect in our model appears conceptually similar to processes

implemented by some previous models (e.g. biotic heterogeneity in Shurin et al. 2004;

monopolization effects by local adaptation in Vanoverbeke et al. 2016). Yet, they found that

positive feedback from early arrivers (in our model, first colonizers) enables their regional

dominance. This result represents what we found in metacommunities with high dispersal and

initial numeric advantage, but not in those with trait-dependent priority effects. This

discrepancy arises because we incorporated seasonality and realistic spatial and temporal

phenological variations in our model. These variations lead to a constant reshuffling of

competitive hierarchies across time and space that reduces or prevents local and regional

extinction, as predicted by previous theoretical results in non-spatial models (Rudolf 2019; Zou

and Rudolf 2020). In our model, phenological variations changed interspecific competition

coefficients and therefore the relative strengths between them and each species' self-limitation.

When aggregated on a spatial or temporal scale, this reshuffling of competitive hierarchies can

also lead to intransitive competition, which is predicted to stabilize multispecies systems

(Allesina and Levine 2011; Levine et al. 2017).

Our results provide a theoretical basis for interpreting many empirical results, especially

those explicitly considering the role of priority effects. For instance, experiments in freshwater

protists found that community composition in homogeneous patches was strongly correlated to

the temporal sequence of assembly despite dispersal (Pu and Jiang 2015). Our results suggest

that this counterintuitive pattern can be easily explained by trait-dependent priority effects.

Consistent with this prediction, Pu and Jiang (2015) reported changes in interspecific

competition based on arrival orders of species, indicating the presence of trait-dependent

priority effects. Our observations also corroborate the result that priority effects in local nectar

microbial communities help maintain beta diversity (Vannette and Fukami 2017), although we

do not observe a consistent increase of beta diversity with higher dispersal, as in the

experiment. Toju et al. (2018) also found a persistent numeric advantage of initially abundant

species after prolonged dispersal in nectar microbial communities. This may indicate that

dispersal in this empirical study was not high enough to homogenize all local patches or

priority effects initiated by numeric advantage were maintained in other trait-dependent

mechanisms, such as habitat modification.

Future Directions

Phenology and priority effects are key factors that structure population dynamics and

coexistence patterns in natural communities. Although the importance of priority effects is

increasingly recognized (Fukami 2015; Rudolf 2019), they are rarely studied in an explicitly

spatial context. Our model indicates the need to consider specific mechanisms of priority

effects when studying spatial dynamics of metacommunities and vice versa. For instance,

systems that display robust diversity measurements regardless of dispersal might be more

influenced by trait-dependent processes. In these systems, considering specific trait-dependent

mechanisms of priority effects may help to understand otherwise unexpected patterns (e.g. Pu

and Jiang 2015; Vannette and Fukami 2017; Toju et al. 2018). Conversely, metacommunities

with species known to display trait-dependent priority effects (e.g. amphibians, Rudolf 2018;

dragonflies, Rasmussen et al. 2014; plants, Kardol et al. 2013; Blackford et al. 2020) may

show different dynamics at a regional scale compared to metacommunities with frequency-

dependent priority effects. Yet, few studies have considered this aspect when studying

dispersal-diversity relationships in these systems, although many metacommunities are driven

by stochastic processes that promote priority effects (e.g. Chase 2007).

Our model predicts that seasonal metacommunities under spatial and temporal variations

in phenology may behave differently based on the mechanism of priority effects. This is

relevant to many systems in nature, including ephemeral ponds, nectar microbiome, pitcher

plant inquilines, rocky intertidal pools, and any other periodic communities where such

variations could occur (Berlow and Navarrete 1997; Kneitel and Miller 2003; Chase 2007;

Sheriff et al. 2011; Theobald et al. 2017; Vannette and Fukami 2017). Although recent studies

of these systems indicate that phenology can vary considerably between communities (Sheriff

et al. 2011; Diez et al. 2012; Theobald et al. 2017; Carter et al. 2018), few have addressed the

local and regional level consequences of these variations. As climate change could exacerbate

phenological shifts and variations (Parmesan 2006; Diez et al. 2012; Wolkovich et al. 2014;

Pearse et al. 2017; but see Stemkovski et al. 2022), the arrival order of species may differ

greatly over space and time, leading to increasing opportunities for priority effects.

Understanding how metacommunities respond to this increasing reshuffling of arrival order

(and likely competitive hierarchy) in the long term could help evaluate the responses of

biodiversity to climate change.

We have extensively investigated different forms of dispersal and incorporated variations

in phenology, but the influence of another aspect of spatial communities, environmental

heterogeneity, remains unexplored. Heterogeneous patches could serve as a spatial refuge for

some species, promoting regional coexistence (Shurin et al. 2004; Schreiber and Killingback

2013). Environmental heterogeneity may also induce trait-dependent priority effects, leading to

monopolization effects (Urban and De Meester 2009; De Meester et al. 2016) or legacy effects

(Miller and Allesina 2021) in local patches. Furthermore, environmental heterogeneity could

covary with other factors that are coupled with another process, such as spatial variation of

arrival time (e.g., species may arrive early in warmer patches), leading to more complex

dynamics. Finally, we focused on systems with one generation per growing season (e.g., annual

plants, amphibians), but the life history could also affect the importance of trait-dependent

priority effects: with multiple generations per season (e.g., aphids, protists), frequency-

dependent priority effects could arise from within-season population dynamics, leading to

alternative patterns of biodiversity (Zou and Rudolf 2020). Collectively, our results emphasize

that the temporal sequence of community assembly is a critical process shaping biodiversity in

spatial communities. A mechanistic understanding of this process is fundamental to our

knowledge of how biodiversity responds to critical environmental processes in a changing

world.

Acknowledgement

We thank members of Miller and Rudolf labs for their feedback on the project and Lydia Beaudrot, Joshua Fowler, Tom E. X. Miller, and Chuliang Song for their comments on the manuscript. Feedback from Erol Akçay, Karen Abbott and four anonymous reviewers greatly improved the manuscript. Funding was provided by NSF DEB-1655626.

Statement of Authorship

H.-X.Z. conceived the idea, designed, and performed the research with assistance from V.H.W.R. H.-X.Z. wrote the first draft, and all authors contributed to revisions.

Data and Code Accessibility

Data and code are deposited at Dryad (Zou and Rudolf 2023).

478   **Table 1. Major parameters and values used in simulations.**

	Definition	Symbol	Value(s) Used
Population dynamics	Intrinsic growth rate	λ	100
	Intraspecific competition	α_{ii}	Randomly drawn from normal distribution with means 0.07 (shown in main text), 0.10 or 0.15 (shown in Figure S8-S13) and standard deviation 0.01
	Constants for calculating interspecific competition	B, c	0.225, -0.5
	End-of-season survival	s	Randomly drawn from normal distribution with mean 0.5 and standard deviation 0.05
Metacommunity	Number of patches	P	50
	Number of species	n	5 (shown in main text), 10 (shown in Figure S5)
	Number of generations in a season	T	1
	Initial population	N_0	10
	Dispersal rate	r	Range: 0.001-0.1; representative values: 0.01 (low) and 0.1 (high)
Emergence time	Average emergence time	$\overline{p}_i$	0, 0.5, 1, 1.5, 2 for species 1-5, respectively
	Seasonal variation	v	2 (shown in main text), 0.5, and 8 (shown in Figure S6-S7)

Figure Captions

Figure 1. Illustration of frequency-dependent and trait-dependent priority effects and how they

operate at different temporal scales. A species may arrive several seasons before its competitor,

allowing it to reproduce and thus reach a higher population than the immigrating population of

the late arriver (A, C). However, species can also differ in their phenology, i.e., when they

become "active" (or emerge) within a season (B, D). This within-season timing can lead to

changes in species interaction via changes in traits (e.g., size; B) and therefore trait-dependent

priority effects. Note that a species can arrive several seasons earlier at a patch, but it can still

have a later phenology within a season (C, D): the species with a frequency-dependent priority

effect does not necessarily imply a trait-dependent priority effect.

Figure 2. Basic processes of the metacommunity model. Colors represent species. In the

hypothetical metacommunity each patch is occupied by one species and the season starts when

species emerge in patches (step 1). Emergence time can vary across seasons and patches within

a given species-specific range. After emergence, species compete locally (step 2) according to

competition coefficients that either: (A) do not depend on the relative emergence time (without

trait-dependent priority effects), or (B) depend on the pairwise difference in relative emergence

time (Δp) (with trait-dependent priority effects). These local patch dynamics together

determine regional patterns (step 3) at the end of the season. Finally, the season ends with

dispersal (step 4) and end-of-season disturbance.

Figure 3. Example of population dynamics of one randomly selected patch under Scenario 1

(Equal Initials) and different dispersal rates, with and without trait-dependent priority effects.

Line colors represent species. The left column shows dynamics without trait-dependent priority

effects, and the right column shows dynamics with trait-dependent priority effects. Rows show

results for low (0.01) and high (0.1) dispersal rates (r). Each line of the same color in a panel

represents the population dynamics of that species from one out of 50 simulations. All

population densities are transformed by natural log. See Table 1 for other parameters used.

Figure 4. Example of population dynamics of one randomly selected patch under Scenario 1

(First Colonizer) and different dispersal rates, with and without trait-dependent priority effects.

Line colors represent species. The left column shows dynamics without trait-dependent priority

effects, and the right column shows dynamics with trait-dependent priority effects. Rows show

results for low (0.01) and high (0.1) dispersal rates (r). Each line of the same color in a panel

represents the population dynamics of that species from one out of 50 simulations. All

population densities are transformed by natural log. See Table 1 for other parameters used.

Figure 5. Relationship of alpha, beta, and gamma diversity and dispersal rates under the two

initial scenarios, with or without trait-dependent priority effects. Left and right columns show

dispersal-diversity relationships under Scenario 1 (Equal Initials) and 2 (First Colonizer),

respectively. Each line is rendered from data of all 50 simulations. See Table 1 for other

parameters used.

Reference

Alford, R. A., and H. M. Wilbur. 1985. Priority effects in experimental pond communities: competition between Bufo and Rana. Ecology.

Allesina, S., and J. M. Levine. 2011. A competitive network theory of species diversity. Proceedings of the National Academy of Sciences of the United States of America 108:5638–5642.

Altermatt, F., S. Schreiber, and M. Holyoak. 2011. Interactive effects of disturbance and dispersal directionality on species richness and composition in metacommunities. Ecology.

Berlow, E. L., and S. A. Navarrete. 1997. Spatial and temporal variation in rocky intertidal community organization: Lessons from repeating field experiments. Journal of Experimental Marine Biology and Ecology 214:195–229.

Blackford, C., R. M. Germain, and B. Gilbert. 2020. Species differences in phenology shape coexistence. The American Naturalist 195.

Cadotte, M. W. 2006. Dispersal and species diversity: A meta-analysis. American Naturalist.

Calcagno, V., N. Mouquet, P. Jarne, and P. David. 2006. Coexistence in a metacommunity: the competition–colonization trade-off is not dead. Ecology Letters 9:897–907.

Carrara, F., F. Altermatt, I. Rodriguez-Iturbe, and A. Rinaldo. 2012. Dendritic connectivity controls biodiversity patterns in experimental metacommunities. Proceedings of the National Academy of Sciences.

Carter, S. K., D. Saenz, and V. H. W. Rudolf. 2018. Shifts in phenological distributions reshape interaction potential in natural communities. Ecology Letters 21:1143–1151.

Catano, C. P., T. L. Dickson, and J. A. Myers. 2017. Dispersal and neutral sampling mediate contingent effects of disturbance on plant beta-diversity: a meta-analysis. Ecology Letters 20:347–356.

Chase, J. M. 2007. Drought mediates the importance of stochastic community assembly. Proceedings of the National Academy of Sciences of the United States of America.

De Meester, L., J. Vanoverbeke, L. J. Kilsdonk, and M. C. Urban. 2016. Evolving Perspectives on Monopolization and Priority Effects. Trends in Ecology and Evolution 31:136–146.

Diez, J. M., I. Ibáñez, A. J. Miller-Rushing, S. J. Mazer, T. M. Crimmins, M. A. Crimmins, C. D. Bertelsen, et al. 2012. Forecasting phenology: From species variability to community patterns. Ecology Letters.

Drake, J. A. 1991. Community-assembly mechanics and the structure of an experimental species ensemble. American Naturalist 137:1–26.

Dray, S., G. Blanchet, D. Borcard, G. Guenard, T. Jombart, G. Larocque, P. Legendre, et al. 2018. Package 'adespatial.' R package version 3–8.

Fodelianakis, S., A. Lorz, A. Valenzuela-Cuevas, A. Barozzi, J. M. Booth, and D. Daffonchio. 2019. Dispersal homogenizes communities via immigration even at low rates in a simplified synthetic bacterial metacommunity. Nature Communications 10:1314.

Fragata, I., R. Costa-Pereira, M. Kozak, A. Majer, O. Godoy, and S. Magalhães. 2022. Specific sequence of arrival promotes coexistence via spatial niche pre-emption by the weak competitor. Ecology Letters n/a.

Fukami, T. 2004. Assembly history interacts with ecosystem size to influence species diversity. Ecology 85:3234–3242.

———. 2015. Historical contingency in community assembly: integrating niches, species pools, and priority effects. Annual Review of Ecology, Evolution, and Systematics.

Grainger, T. N., and B. Gilbert. 2016. Dispersal and diversity in experimental metacommunities: linking theory and practice. Oikos.

Grainger, T. N., A. D. Letten, B. Gilbert, and T. Fukami. 2019. Applying modern coexistence theory to priority effects. Proceedings of the National Academy of Sciences of the United States of America 116:6205–6210.

Kardol, P., L. Souza, and A. T. Classen. 2013. Resource availability mediates the importance of priority effects in plant community assembly and ecosystem function. Oikos.

Ke, P. J., and A. D. Letten. 2018. Coexistence theory and the frequency-dependence of priority effects. Nature Ecology and Evolution 2:1691–1695.

Kerr, B., M. A. Riley, M. W. Feldman, and B. J. M. Bohannan. 2002. Local dispersal promotes biodiversity in a real-life game of rock-paper-scissors. Nature.

Kharouba, H. M., J. Ehrlén, A. Gelman, K. Bolmgren, J. M. Allen, S. E. Travers, and E. M. Wolkovich. 2018. Global shifts in the phenological synchrony of species interactions over recent decades. Proceedings of the National Academy of Sciences of the United States of America 115:5211–5216.

Kneitel, J. M., and T. E. Miller. 2003. Dispersal Rates Affect Species Composition in Metacommunities of Sarracenia purpurea Inquilines. American Naturalist 162:165–171.

Legendre, P. 2019. A temporal beta‐diversity index to identify sites that have changed in exceptional ways in space–time surveys. Ecology and Evolution 9:3500–3514.

Leibold, M. A., M. Holyoak, N. Mouquet, P. Amarasekare, J. M. Chase, M. F. Hoopes, R. D. Holt, et al. 2004. The metacommunity concept: A framework for multi-scale community ecology. Ecology Letters.

Leibold, M. A., M. C. Urban, L. De Meester, C. A. Klausmeier, and J. Vanoverbeke. 2019. Regional neutrality evolves through local adaptive niche evolution. Proceedings of the National Academy of Sciences of the United States of America 116:2612–2617.

Levine, J. M., J. Bascompte, P. B. Adler, and S. Allesina. 2017. Beyond pairwise mechanisms of species coexistence in complex communities. Nature 546:56–64.

Lu, M. 2021. Complex relationships between beta diversity and dispersal in meta-community models. Ecography n/a.

Menzel, A., T. H. Sparks, N. Estrella, E. Koch, A. Aasa, R. Ahas, K. Alm-Kübler, et al. 2006. European phenological response to climate change matches the warming pattern. Global Change Biology 12:1969–1976.

Miller, Z. R., and S. Allesina. 2021. Metapopulations with habitat modification. Proceedings of the National Academy of Sciences 118.

Mouquet, N., and M. Loreau. 2002. Coexistence in metacommunities: The regional similarity hypothesis. American Naturalist 159:420–426.

———. 2003. Community Patterns in Source-Sink Metacommunities. American Naturalist.

Oksanen, J., F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. O'hara, G. L. Simpson, et al. 2013. Package 'vegan.' Community ecology package, version 2:1–295.

Parmesan, C. 2006. Ecological and Evolutionary Responses to Recent Climate Change. Annual Review of Ecology, Evolution, and Systematics.

Pearse, W. D., C. C. Davis, D. W. Inouye, R. B. Primack, and T. J. Davies. 2017. A statistical estimator for determining the limits of contemporary and historic phenology. Nature Ecology and Evolution 1:1876–1882.

Poulos, D. E., and M. I. McCormick. 2014. Who wins in the battle for space? The importance of priority, behavioural history and size. Animal Behaviour 90:305–314.

Pu, Z., and L. Jiang. 2015. Dispersal among local communities does not reduce historical contingencies during metacommunity assembly. Oikos 124:1327–1336.

Pulliam, H. R. 1988. Sources, Sinks, and Population Regulation. The American Naturalist 132:652–661.

R Core Team. 2021. R: A language and environment for statistical computing.

Rasmussen, N. L., B. G. Van Allen, and V. H. W. Rudolf. 2014. Linking phenological shifts to species interactions through size-mediated priority effects. Journal of Animal Ecology.

Rudolf, V. H. W. 2018. Nonlinear effects of phenological shifts link interannual variation to species interactions. Journal of Animal Ecology.

———. 2019. The role of seasonal timing and phenological shifts for species coexistence. Ecology Letters.

Schreiber, S. J., and T. P. Killingback. 2013. Spatial heterogeneity promotes coexistence of rock–paper–scissors metacommunities. Theoretical Population Biology 86:1–11.

Sheriff, M. J., G. J. Kenagy, M. Richter, T. Lee, Ø. Tøien, F. Kohl, C. L. Buck, et al. 2011. Phenological variation in annual timing of hibernation and breeding in nearby populations of Arctic ground squirrels. Proceedings of the Royal Society B: Biological Sciences 278:2369–2375.

Shorrocks, B., and M. Bingley. 1994. Priority effects and species coexistence: Experiments with fungal-breeding Drosophila. Journal of Animal Ecology 63:799–806.

Shurin, J. B., P. Amarasekare, J. M. Chase, R. D. Holt, M. F. Hoopes, and M. A. Leibold. 2004. Alternative stable states and regional community structure. Journal of Theoretical Biology 227:359–368.

Stemkovski, M., J. R. Bell, E. R. Ellwood, B. D. Inouye, H. Kobori, S. D. Lee, T. Lloyd-Evans, et al. 2022. Disorder or a new order: How climate change affects phenological variability. Ecology n/a:e3846.

Theobald, E. J., I. Breckheimer, and J. HilleRisLambers. 2017. Climate drives phenological reassembly of a mountain wildflower meadow community. Ecology 98:2799–2812.

Thompson, P. L., L. M. Guzman, L. De Meester, Z. Horváth, R. Ptacnik, B. Vanschoenwinkel, D. S. Viana, et al. 2020. A process-based metacommunity framework linking local and regional scale community ecology. Ecology Letters 23:1314–1329.

Tilman, D. 1994. Competition and Biodiversity in Spatially Structured Habitats. Ecology 75:2–16.

Toju, H., R. L. Vannette, M. P. L. Gauthier, M. K. Dhami, and T. Fukami. 2018. Priority effects can persist across floral generations in nectar microbial metacommunities. Oikos 127:345–352.

Urban, M. C., and L. De Meester. 2009. Community monopolization: local adaptation enhances priority effects in an evolving metacommunity. Proceedings of the Royal Society B-Biological Sciences.

Vannette, R. L., and T. Fukami. 2017. Dispersal enhances beta diversity in nectar microbes. Ecology Letters 20:901–910.

Vanoverbeke, J., M. C. Urban, and L. De Meester. 2016. Community assembly is a race between immigration and adaptation: eco-evolutionary interactions across spatial scales. Ecography 39:858–870.

Vanschoenwinkel, B., F. Buschke, and L. Brendonck. 2013. Disturbance regime alters the impact of dispersal on alpha and beta diversity in a natural metacommunity. Ecology 94:2547–2557.

Wolkovich, E. M., B. I. Cook, K. K. McLauchlan, and T. J. Davies. 2014. Temporal ecology in

the Anthropocene. Ecology Letters.

Yu, D. W., and H. B. Wilson. 2001. The Competition☐Colonization Trade☐off Is Dead; Long

Live the Competition☐Colonization Trade☐off. The American Naturalist 158:49–63.

Zou, H.-X., and V. H. W. Rudolf. 2020. Priority effects and season length shape long-term

competition dynamics. bioRxiv 2020.08.14.251926.

Zou, H.-X., and V. H. W. Rudolf. 2023. Data from: Priority effects determine how dispersal

affects biodiversity in seasonal metacommunities. American Naturalist, Dryad Digital

Repository, https://doi.org/10.5061/dryad.sbcc2frb4.

A. Density-dependent priority effects

reproduction

Time

B. Trait-dependent priority effects

growth

Time

C. Arrivals across years (sequence of colonization)

colonizes   colonizes

reproduction reproduction reproduction reproduction

Years

D. Arrivals within a season (sequence of phenology)

emerges   emerges   Time within a year

4. End-of-Season Disturbance, then Next Season

Species 1
Species 2
Species 3
Species 4
Species 5

Δp

Actual emergence Time
Average emergence Time
Range of possible emergence time

Time

Actual emergence order: 1-3-2-5-4
Average emergence order: 1-2-3-4-5

3. Dispersal   **1. Emergence**

2. Local Dynamics

Competition Coefficients   Population Dynamics

I. No Trait-dependent Priority Effects

α α_{12} α_{21}

Δp

N N_2 N_1

Time within a season

II. Trait-dependent Priority Effects

α α_{21} α_{12}

Δp

N N_1 N_2

Time within a season