**Title:** Beta-cell Cre expression and reduced *Ins1* gene dosage protect mice from type 1 diabetes

**Authors:** Søs Skovsø[1]*, Peter Overby[1]*, Jasmine Memar-Zadeh[1], Jason T.C. Lee[1], Jenny C.C. Yang[1],

Iryna Shanina[1], Vaibhav Sidarala[2], Elena Levi-D'Ancona[2], Jie Zhu[2], Scott A. Soleimanpour[2], Marc S.

Horwitz[1], James D. Johnson*[1]

* These authors contributed equally

Correspondence should be addressed to James D. Johnson: james.d.johnson@ubc.ca, 2350 Health

Sciences Mall, 5358 Life Sciences Building, Vancouver, BC, V6T 1Z3

**Affiliations:** [1] Life Sciences Institute, University of British Columbia, Vancouver, BC, Canada; [2]

Department of Molecular and Integrative Physiology, Division of Metabolism, Endocrinology, and

Diabetes of the Department of Internal Medicine, University of Michigan, Ann Arbor, MI, USA

**Correspondence:** J.D.J.

**Keywords:** type 1 diabetes animal models, Cre recombinase, knock-in mice, pancreatic β-cell

**Funding:** This project was primarily supported by an operating grant to J.D.J. from the Canadian

Institutes for Health Research (PJT-152999) and the JDRF Centre of Excellence at UBC (3-COE-2022-

1103-M-B). S.A.S. was supported by the JDRF (CDA-2016-189, COE-2019-861), the NIH (R01

DK108921, U01 DK127747), and the Department of Veterans Affairs (I01 BX004444).

**Acknowledgements:** We thank colleagues in the β-cell biology field for helpful discussions, including

on Twitter.

**Disclosure statement:** The authors have nothing to disclose

## Author Contributions

S. S. co-conceived experiments, designed and executed in vivo experiments, analyzed data, co-wrote the manuscript

P. O. executed in vivo experiments, analyzed data, co-wrote the manuscript

J. M-Z. analyzed data and co-wrote the manuscript

J. L. designed and analyzed flow cytometry experiments

J. C.C Y. executed in vivo experiments

I. S. designed and executed flow cytometry experiments

V. S. conducted *in vivo* validation experiments

E. L-D. conducted *in vivo* validation experiments

J. Z. conducted *in vivo* validation experiments

S. A. S. oversaw and interpreted *in vivo* validation experiments

M. H. oversaw and interpreted *in vivo* experiments

J. D. J. designed the model, co-conceived experiments, analyzed data, co-wrote the manuscript, and is the ultimate guarantor of this work.

**Abstract word count:** 222

**Body word count:** 4023

**Reference count:** 38

**Figure count:** 6

## Abstract

A central goal of physiological research is the understanding of cell-specific roles of disease-associated genes. Cre-mediated recombineering is the tool of choice for cell type-specific analysis of gene function in pre-clinical models. In the type 1 diabetes research field, multiple lines of NOD mice have been engineered to express Cre recombinase in pancreatic β-cells using insulin promoter fragments, but tissue promiscuity remains a concern. Constitutive $Ins1^{tm1.1(cre)Thor}$ ($Ins1^{Cre}$) mice on the C57/bl6-J background has high β-cell specificity and with no reported off-target effects. We explored if NOD:$Ins1^{Cre}$ mice could be used to investigate β-cell gene deletion in type 1 diabetes disease modeling. We studied wildtype ($Ins1^{WT/WT}$), $Ins1$ heterozygous ($Ins1^{Cre/WT}$ or $Ins1^{Neo/WT}$), and $Ins1$ null ($Ins1^{Cre/Neo}$) littermates on a NOD background. Female $Ins1^{Neo/WT}$ mice exhibited significant protection from diabetes, with further near-complete protection in $Ins1^{Cre/WT}$ mice. The effects of combined neomycin and Cre knock-in in $Ins1^{Neo/Cre}$ mice were not additive to the Cre knock-in alone. In $Ins1^{Neo/Cre}$ mice, protection from diabetes was associated with reduced insulitis at 12 weeks of age. Collectively, these data confirm previous reports that loss of $Ins1$ alleles protects NOD mice from diabetes development and demonstrates, for the first time, that Cre itself may have additional protective effects. This has significant implications for the experimental design and interpretation of pre-clinical type 1 diabetes studies using β-cell-specific Cre in NOD mice.

## Introduction

Type 1 diabetes (T1D) is a chronic disorder precipitated by immune-mediated pancreatic β-cell destruction and associated with the presence of autoantibodies against β-cell proteins (1,2). Due to the progressive loss of β-cells and consequent insulin deficiency, individuals living with type 1 diabetes have lifelong dependency on exogenous insulin (3). Higher levels of endogenous insulin secretion are associated with better short- and long-term outcomes in type 1 diabetes. Preservation of residual β-cells and islet function at the time of diagnosis is therefore imperative (4).

Exogenous insulin administration is not a cure of type 1 diabetes. Research efforts using pre-clinical

animal models continue to produce new therapeutic possibilities. Initially developed as a model for

spontaneous onset of cataracts, the female non-obese diabetic (NOD/ShiLt) mouse (commonly known

as NOD) is the most well-established and extensively used model of type 1 diabetes (5). The polygenic

NOD mouse model recapitulates multiple pathophysiological features of human type 1 diabetes,

including the development of autoantibodies in prediabetic NOD mice (6), circulating autoreactive T cells

(7), and subsequent onset of hyperglycemia as progressive β-cell loss materializes (8). While the

emergence of hyperglycemia most often occurs between 12-15 weeks of age, islet immune infiltration

is established earlier at 8-12 weeks with insulitis present across the pancreas(9). The cleanliness of

housing facilities and the animal breeding approach are just two factors which affect the age at which

these features present.

Mice express two nonallelic insulin genes. The insulin 2 gene (*Ins2*) is the murine homologue of the

human insulin gene and is located on chromosome 7 (10). The insulin 1 gene (*Ins1*) is the result of an

RNA-mediated gene duplication event. *Ins1* has a simpler gene structure lacking the second intron

present in *Ins2,* and is found on the murine chromosome 19 (10). *Ins1* is expressed specifically in

pancreatic β-cells, whereas *Ins2* is expressed predominantly in β-cells, with trace expression in other

tissues including thymus and brain (11). Previous studies have examined the effects of insulin gene

knockout in NOD mice. Complete knockout of *Ins2* on the NOD background accelerated diabetes onset

(2), a phenomenon attributed to a failure in central tolerization to insulin. Thymus-specific deletion of

*Ins2* was reported to be sufficient to cause spontaneous diabetes, even outside the NOD background

(12). Although we have not observed autoimmune diabetes in globally deficient $Ins2^{-/-}$ mice (13,14) it

has previously reported that male, but not female, NOD:$Ins1^{Neo/WT}$;$Ins2^{-/-}$ mice, with a single remaining

*Ins1* allele have been shown to succumb to insulin insufficiency (15), a finding we confirmed on other

backgrounds (16). These observations were found to be dependent on housing conditions (11). In

contrast, NOD:$Ins1^{Neo/Neo}$ mice have previously been shown to be largely protected from insulitis

diabetes (10), which was proposed to be due to the loss of autoantigenic *Ins1*-derived peptides despite

presence of insulin autoantibodies (IAA). Insulin is a primary auto-antigen in both murine and human

type 1 diabetes pathogenesis (1). Replacing *Ins1* with non-antigenic human insulin also protects NOD

mice from onset of diabetes (17). Together, these previous observations indicate that insulin 1 gene

dosage is a key driver of diabetes in NOD mice.

To advance the understanding of the underlying mechanisms of T1D pathogenesis, powerful genetic

engineering tools are employed by researchers across the world. *Ins1*[Cre] mice take advantage of the

tissue specificity of *Ins1* expression to excise *lox*P site-flanked (floxed) DNA segments. As the Cre DNA

recombinase allele is inserted into exon 2 of *Ins1* in these mice, floxed target genes are removed

specifically in beta-cells at the cost of the loss of 1 *Ins1* allele. Both *Ins1*[Cre] and *Ins1*[Neo] mice have

reduced *Ins1* gene dosages with 50% or 100% in their heterozygous and homozygous states,

respectively. For *Ins1*[Neo] mice, the neomycin cassette obliterates the entire *Ins1* promotor and gene

sequence in addition to deletion of 7-9kb upstream and downstream of *Ins1* (see Figure 1A).

Since the early days of Cre/*lox*P system usage, concerns of side effects such as Cre toxicity have

been raised. More than two decades ago, Loonstra *et al* observed increased sister chromatid exchange

frequency, leading to a cellular halt in the $G_2$/M phase and a reduction in proliferation of mouse

embryonic fibroblasts (MEF) transduced with a bicitronic retroviral vector encoding Cre (18). Silver and

Livingston demonstrated similar results including ceased proliferation in 293xLac cells, NIH 3T3 cells

and MEFs transfected with Cre-expressing retroviral vectors (19). The same study revealed

chromosomal abnormalities in Cre-expressing MEFs. Lentiviral Cre administration and expression have

likewise been shown to reduce proliferation through cellular accumulation in the $G_2$M phase of Cre-

expressing CV-1 and COS cells, two kidney cell lines derived from monkeys (20). Cre toxicity has been

shown to be dose dependent, even when expressed transiently, when using an adenovirus vector (21).

Past studies have confirmed Cre toxicity and apoptosis in mouse cardiac tissue (22,23) and p53[-/-] thymic

lymphoma cells (24). Collectively, these findings emphasize the necessity of using Cre-only (no flox)

controls into *in vivo* research study designs since Cre recombinase expression may affect cells beyond

the intended target gene deletion.

For this study, we generated and characterized a new NOD *Ins1*[Cre] knock-in mouse line. Our

intension was to establish an *in vivo* tool facilitating the unravelling of type 1 diabetes pathogenesis. The

NOD $Ins1^{Cre}$ model would ideally enable studying of β-cell specific deletion of target genes in mice on an NOD background. As a predecessor, the goal of this study was to investigate the impact of 1) deleting one $Ins1$ allele and 2) introducing beta cell specific Cre expression, on the spontaneous diabetes development NOD mice are well known for. As previous studies by the Eisenbarth group provided evidence that $Ins1^{Neo/WT}$ mice, with only one intact $Ins1$ allele, exhibit decreased and delayed diabetes incidence, we were expecting similar results for $Ins1^{Cre/WT}$ mice. We incorporated both NOD:$Ins1^{Neo/Wt}$ NOD;$Ins1^{WtTWT}$ littermates in our study design, to control for both the loss of one $Ins1$ allele and introduction of Cre expression in NOD $Ins1^{Cre/WT}$ mice. We compared diabetes incidence, insulitis severity, and immune activation between groups within each sex. Our data demonstrate that $Ins1$-driven Cre expression has further protective effects beyond the loss of a single $Ins1$ allele deleted via $Ins1^{Neo}$. Carefully chosen proper controls and caution are therefore required when interpreting experiments including β-cell specific Cre expression in mice on an NOD background.

**Methods**

*Mice*

All animal procedures and ethical standards were in accordance with the Canadian Council for Animal Care guidelines. All animal studies and protocols were approved by the University of British Columbia Animal Care Committee and Institutional Care and Use Committee (IACUC) at the University of Michigan. At UBC, mice were housed in the Centre for Disease Modelling Specific Pathogen-Free (SFP) facility on a standard 12-h light/12-h dark cycle with ad libitum access to chow diet (PicoLab, Mouse Diet 20–5058). To generate NOD:$Ins1^{Cre}$ and NOD:$Ins1^{Neo}$ mice, we contracted The Jackson Laboratories to backcross (>12 times) B6(Cg)-$Ins1^{tm1.1(cre)Thor}$/J (The Jackson Laboratory, US, #026801) ($Ins1^{Cre}$) and NOD.129S2(B6)-$Ins1^{tm1Jja}$/GseJ (The Jackson Laboratory, US, #0005035) $Ins1^{Neo}$) mice onto a NOD/ShiLtJ (The Jackson Laboratory, US, #001976) (NOD) background. The $Ins1^{Cre}$ and $Ins1^{Neo}$ mice were originally on a mixed, largely C57Bl/6J, background. Subsequently, we designed a strict breeding strategy for our study. An $Ins1^{Cre}$ maternal parent colony as well as a $Ins1^{Neo}$ paternal parent colony was established (Fig. 1B). Each parent colony was backcrossed every five

generations. Female *Ins1*^Cre/WT and male *Ins1*^Neo/WT mice were set up as breeders, at 7-8 weeks of age,

to generate experimental mice. Mice from the parental colonies were only included once as breeders to

eliminate risk of onset of hyperglycemia during pregnancy and weaning at later ages. This breeding

strategy generated littermates of four genotypes:  wildtype NOD:*Ins1*^WT/WT mice with both insulin 1

alleles,  heterozygous NOD:*Ins1*^Cre/WT mice with an *Ins1* replaced with Cre-recombinase, heterozygous

NOD:*Ins1*^Neo/WT mice with one *Ins1* allele replaced with a neomycin cassette, and full *Ins1* null mice with

both *Ins1* alleles replaced NOD:*Ins1*^Neo/Cre. Specific cohorts were monitored twice per week for the sole

purpose of determining hyperglycemia onset incidence and body mass changes. Any mice that

developed diabetes, defined as two consecutive blood glucose measurements $\geq$16 mmol/L or one

measurement $\geq$22 mmol/L, were euthanized. Specific cohorts were generated for tissue analysis

terminated at 12 weeks of age in a pre-diabetic phase (for the Vancouver housing facility) or at 1 year

of age, all animals were monitored for diabetes prior to euthanasia.

An additional colony of NOD:*Ins1*^Cre mice were generated independently through in-house

backcrossing at a second site (University of Michigan) via the speed congenic approach in consultation

with Charles River Laboratories (Wilmington, MA). Following each backcross, *Ins1*^Cre/WT offspring with

allelic profiles most closely matching the NOD strain (determined by MAX-BAX mouse 384 SNP panel

screening), were selected as breeders for the subsequent backcross. Following 8 generations of

backcrossing, animals with an allelic profile percent match >99.9% were utilized to generate *Ins1*^WT/WT

and *Ins1*^Cre/WT experimental mice. Animals were housed in an SPF facility on a standard 12-h light/12-h

dark cycle with access to ad libitum chow diet (LabDiets, Rodent Diet 5L0D) and water. Drinking water

was provided at a pH level of 2.5-3 upon advice from Jackson laboratories that acidified water supports

the diabetes frequency in many NOD colonies (25)Blood glucose measurements were taken once-twice

per week. Incidence of diabetes, defined as blood glucose levels > 16 mM for 5 consecutive

measurements, was recorded. Diabetic mice were euthanized and excluded from future analysis.

*Tissue processing and histology*

Immediately following euthanasia, pancreata were collected according to a pre-established protocol

with the exception that extracted pancreases were not further treated to obtain isolated islets and were

instead processed as a whole(26). The dissected pancreata were fixed in 4% PFA for 24 hours prior to

storage in 70% ethanol at 4 °C. Paraffin-embedded sections were prepared, stained with hematoxylin

and eosin (H&E), and imaged by WaxIT Histology Services Inc. (Vancouver, BC).

*Islet infiltration scoring*

Images from H&E-stained pancreatic sections were analyzed with QuPath software (27). Islets in

pancreatic sections were scored blindly for pancreatic islet infiltration of mononuclear immune cells

according to the previously established 4-point scale(28). In brief, 0- no insulitis, 1- peri-insulitis marked

by less than 25% peripheral immune-islet infiltration, 2- insulitis marked by 25-75% immune cell

infiltration, 3- severe insulitis marked by greater than 75% immune-islet infiltration. Random samples

were scored by a second person to ensure consistency. 25-30 islets per mouse.

*Insulin auto-antibodies and plasma insulin*

Blood for plasma insulin and autoantibodies measurements was collected at study endpoint by cardiac

puncture from anesthetized Animals (2.5% isoflurane) and put directly on ice. Insulin plasma where

measured using Mouse Insulin ELISA kit (#80-INSMS-E10, RRID: AB_2923075; ALPCO). Samples

were shipped to the Insulin Antibody Core Lab at the University of Colorado, Barbara Davis Diabetes

Centre for auto-antibody analysis.  The IAA data are expressed as an index against a standard

positive and negative controls as procedure at the Insulin Antibody Core lab

*Flow cytometry*

Immediately following euthanasia lymph nodes and spleen were isolated and splenic single-cell

suspensions were counted and stained with fluorescently conjugated monoclonal antibodies (mAbs)

for cell-surface markers (see Table 1 for list of antibodies used). Fixable Viability Dye eFluor™ 506

was used as viability dye (#65-0866-14; ThermoFischer). Following staining, cells were analyzed by

flow cytometry and Flow Jo software (Tree Star Inc., Ashland, Oregon).

*Statistics*

Statistical significance was assessed using 2-way ANOVA analysis with Tukey's multiple

comparisons test, at a threshold of $p < 0.05$. Plasma insulin, Insulitis and insulin auto-antibodies

significance were quantified by 1-way ANNOVA analyses. We used the Mantel-Cox-log rank test,

corrected with the Benjamini-Hochberg procedure, to analyse the Kaplain Meier survival plots. Repeated

measured Mixed effect models were applied for RBG and Body mass analyses.

Prism 9 (GraphPad Software Inc., USA) was used for statistical analyses and generation of most

figure panels.  Data are expressed as mean ± SEM unless otherwise specified.

**Results**

*Type 1 diabetes incidence*

A greater proportion of female, in comparison to male, mice develop diabetes and the timing of the

development of the disease is more consistent. We observed the highest diabetes incidence in female

NOD:*Ins1*$^{WT/WT}$ mice, with 65% diabetes incidence at 1 year (Fig. 1C). The majority of these mice (7/9)

were diabetic before 20 weeks of age. The diabetes incidence and disease time-course of this study

was comparable with previous NOD:*Ins1*$^{WT/WT}$ cohorts in our facilities (28), although slightly delayed

relative to some other facilities (see below). Female NOD:*Ins1*$^{Cre/WT}$ exhibited both a delay in diabetes

onset and a diabetes incidence by the end of the study of only 27%, which was significantly different

when compared to NOD:*Ins1*$^{WT/WT}$ mice at one year of age ($p=0.033$; $p_{adjust}=0.078$; Fig. 1C). We also

observed at delay in the timing of the onset between female NOD:*Ins1*$^{Neo/WT}$ mice (Fig. 1C), in agreement

with previous studies of NOD:*Ins1*$^{Neo/WT}$ mice (10), but there was no difference in the final diabetes

incidence between female littermate NOD:*Ins1*$^{Neo/WT}$ mice (64%) and  NOD:*Ins1*$^{WT/WT}$ mice (64%)

($p=0.40$; $p_{adjust}=0.48$; Fig. 1C). Double mutant female NOD:*Ins1*$^{Neo/Cre}$ mice (lacking both wildtype alleles

of *Ins1*) were protected from diabetes (29% diabetes incidence) when compared to NOD:*Ins1*$^{WT/WT}$ mice

(64%), ($p=0.019$; $p_{adjust}=0.078$), to a similarly extent as  NOD:*Ins1*$^{Cre/WT}$ (27%) mice ($p=0.98$; $p_{adjust}=0.98$;

Fig. 1C). NOD:*Ins1*$^{Neo/WT}$ mice had a significantly lower diabetes incidence than  NOD:*Ins1*$^{Cre/Neo}$ mice

(p=0.038; $p_{adjust}$=0.078; Fig. 1C) further suggesting Cre expression rather than loss of one *Ins1* allele

protects against diabetes onset in NOD mice. No differences were observed in random blood glucose

(Fig. 1D,E) and body mass (Fig. 1F,G) prior to diabetes onset in female mice. Together, these data

confirm previous findings that reduced *Ins1* gene dosage protects NOD mice from diabetes, but also

reveal a further, additive protective effect of β-cell Cre expression.

Male NOD:*Ins1*^WT/WT mice demonstrated a low diabetes incidence as expected (20% incidence),  We

observed no cases of hyperglycemia in any of the male mice with reduced *Ins1* gene dosage (Fig. 2A).

There were no differences in random blood glucose (Fig. 2 B,C) or body mass (Fig. 2D,E) between any

of the groups. These results confirm the strong sex bias in this NOD model and suggest that the

protection conferred by reducing *Ins1* gene dosage may not be sex-specific.

*Insulitis and Insulin auto-antibodies*

Next, we examined the effects of reduced *Ins1* gene dosage and Cre expression on insulitis in female

mice, the pathological evidence of islet directed autoimmunity. We found no significant difference in

plasma insulin or autoantibodies between any of the genotypes (Fig. 3A,B). H&E-stained pancreas

sections (Fig. 3C) were blindly scored for immune islet infiltration in a pre-diabetic cohort of littermates

euthanized at 12 weeks of age, and also in the mice that survived to 1 year. Insulitis scores from 12-

week-old mice with reduced *Ins1* gene dosages were not significantly different from NOD:*Ins1*^WT/WT

littermates. We noticed the least amount of immune islet infiltration in double mutant NOD:*Ins1*^Neo/Cre

mice  (Fig. 3D), consistent with their more complete protection from diabetes. The outcomes of the

samples gathered at 1 year are unfortunately inconclusive due to low sample size (NOD:*Ins1*^WT/WT, n=3)

(Fig 3E). The lower survival rate may therefore mask any potential significant changes for both insulitis

scoring (Fig. 3E)  and insulin auto-antibodies (Fig. 3B). Unfortunately, we did not collect blood for this

analysis at the 12-week timepoint for all cohorts due to challenges brought about by the COVID19-

pandemic.

*Immune cell characterization*

To assess immune cell populations in the pancreatic lymph nodes and spleen at 50 weeks of age,

we used a panel of validated antibodies for flow cytometry. While we were able to confidently identify

many key immune cell populations (Fig. 4), there were no significant differences between groups (Fig.

5). These observations demonstrate that β-cell specific insulin gene manipulations alter type 1 diabetes

incidence without robust effects on the lymphocytes found in the pancreatic lymph nodes and spleen.

*Independent Validation Cohort*

To ensure the protective effects of β-cell Cre expression were not solely limited to a single animal

housing facility, we additionally studied female NOD:*Ins1*[Cre/WT] mice that were independently generated

at a separate site (Fig. 6) in parallel to the cohorts studied in Figure 1. The overall incidence of diabetes

development in female NOD mice in this second animal colony was as expected (approximately 65-

80% by 25 weeks of age, Fig. 6A). Similar to our data shown in Figure 1, we observed that female

NOD:*Ins1*[Cre/WT] animals were protected from diabetes incidence (25% by 1 year) and had significantly

improved mean blood glucose (Fig. 6C) when compared to NOD:*Ins1*[WT/WT] littermates (75% incidence

by 1 year). Again, these studies confirm the protective effects of reduced *Ins1* gene dosage and β-cell

Cre expression in NOD mice and suggest that these findings are not due to a consequence of

environment or housing.

**Discussion**

In this study, we used a rigorous littermate control study design to examine the effects, of replacing

one *Ins1* allele with Cre-recombinase, on the onset of diabetes in mice of an NOD background. To

investigate, whether potential effects were due to the loss of *Ins1* or the introduction of Cre we included

*Ins1*[Neo/WT] mice, in our study. Both *Ins1*[Cre] and *Ins1*[Neo] mice have reduced *Ins1* gene dosages with 50%

or 100% in their heterozygous and homozygeous states, respectively. We found a similar reduction in

the diabetes incidence in female *Ins1*[Cre/WT] mice and *Ins1*[Neo/Cre] mice when compared to littermate control

*Ins1*[WT/WT] mice. This work has implications for our understanding of the pathogenesis of type 1 diabetes,

as well as for the use of Cre-recombinase as a tool for *in vivo* genome engineering in mouse models of

the disease.

Our findings concur with previous work that showed a similar reduction in diabetes incidence in

female NOD mice lacking 1 or 2 alleles of *Ins1* (10,15), although we did not observe additional protection

with the double *Ins1* knockout, beyond what was seen with the Cre replacement allele. Similarly, our

data are in alignment with a previous study showing replacing the murine *Ins1* gene with the human *INS*

gene was found to protect female NOD mice from diabetes in both heterozygous and homozygous

states (17). We were unable to detect differences in diabetes incidence in male $Ins1^{Cre/WT}$, $Ins1^{Neo/WT}$,

and $Ins1^{Neo/Cre}$ NOD mice, perhaps due to an insufficient study period. A previous study found that

removal of a single *Ins1* allele is sufficient to abolish spontaneous diabetes in 50-week-old male NOD

mice (10). Together with the work of others, our experiments support the contention that proinsulin 1 is

a key player in the generation of autoimmunity in mice. While there are several autoantigens targeted

by autoreactive T cells in type 1 diabetes (2), insulin and proinsulin are particularly common

autoantibody targets in prediabetic humans (29,30). Our experiments were underpowered to detect

subtle differences in the levels of insulin autoantibodies, as we were limited by only examining a single,

late time point. However, we did examine insulitis at two time points, and insulin antibodies are often

correlated with insulitis (31). In our hands, there was a qualitative difference in the number of islets that

did not exhibit insulitis at 50 weeks in mice with at least 1 *Ins1* allele replaced. At 12 weeks, there was

a slight trend towards more insulitis-free islets in the $Ins1^{Neo/Cre}$ mice, consistent with the greater

protection from type 1 diabetes incidence. These observations are consistent with previous studies

showing that *Ins1* knockout in NOD mice is protective against the development and severity of insulitis

(10). Thus, our results show that reducing the *Ins1* gene dosage lowers the threshold required for

diabetes onset, likely by removing the source of primary autoantigens and suppressing insulitis.

An important observation of our study is that Cre expression itself has protective effects in NOD mice,

beyond the protection afforded by the loss of 1 *Ins1* allele. To examine the specific consequences of

Cre expression, we compared NOD;$Ins1^{Cre/WT}$ to a different knock-in (neo) $Ins1^{Neo/WT}$ mouse line and

found roughly twice as much final diabetes protection in Ins$1^{Cre/WT}$ mice versus $Ins1^{Neo/WT}$ mice. Though

it is possible diabetes could have been delayed further than 1 year, these findings suggest that Cre itself

may affect diabetes rates in female NOD mice, with the caveat the Neo and Cre insertions are different,

altering the local gene structure. Mechanistically, we are able to attribute the observed effects to

differences in insulitis, suggesting the possibility of a β-cell autonomous effect of Cre expression.

Previous studies highlighted the potential of Cre recombinase to result in toxicity due to DNA damage

(32,33) Mammalian genomes contain pseudo loxP sites, and even though these sequences can deviate

considerably from the consensus loxP site, they can still serve as functional recognition sites for Cre

(32). It is predicted that the frequency of pseudo-loxP sites could be as many as 250 and 300 in mouse

and human genomes, respectively (33). The sustained presence of high levels of Cre in fibroblasts can

cause growth arrest and chromosomal abnormalities (18-20,22,23). Cre-dependent DNA damage and

accumulation of cytoplasmic DNA have been shown to initiate a STING-dependent immune response

(34). STING is an intracellular adaptor molecule, associated with the endoplasmic reticulum membrane

(35), that can play a critical role in detecting pathogen-derived DNA in the cytoplasm (36). There is

precedence for diabetes protection in NOD mice with early exposure to pathogen in the coxsackievirus

mode (37). A recent paper reported that STING is required for normal β-cell function in mice (38),

although, ironically, the study did not employ Cre-only controls. Theoretically, Cre expressed in β-cells

could delay the onset of diabetes in a STING-dependent manner. Future studies, beyond the scope of

this work, will be required to delineate the molecular mechanisms by which Cre expression induces

further protection that *Ins1* loss in the NOD mouse model. To unequivocally demonstrate an effect of

Cre activity on progression to T1D/insulitis, mice would have to be generated with enzyme-dead Cre

knocked into the same locus, but such studies are also beyond the scope of this work.

As with all studies, this work has a number of limitations. For example, the broadly observed

phenomenon that diabetes incidences differ between NOD mouse colonies means that we cannot

directly compare the results between sites. We do not know the reasons for the apparent difference in

diabetes incidence in wildtype NOD mice between our colonies, but it could be related to many

environmental factors, including native microbiome, water, food, or bedding, as well as subtle

differences in genotype. Another caveat of our study is that we could not simultaneously address *Ins1*

gene dosage and the effects of the Cre transgene, because generating both homozygous and

heterozygous littermates with each of the knock-in alleles is impossible. In order to reduce the potential

impacts of environment and genotype, mentioned above, it was imperative that we prioritized using

littermates. Another limitation is the natural uncertainty if $Ins1^{Cre/WT}$ mice could have proceeded to

develop diabetes beyond the end of this study. While our study has limitations, we believe it is important

to report these results that will help guide those in the field who use type 1 diabetes mouse models or

Cre in any context.

In summary, our observations suggest caution when interpreting experiments that involve Cre

recombinase in NOD mice. Our data showed that Cre expression itself has protective effects in NOD

mice, beyond the protection afforded by the loss of 1 $Ins1$ allele. Cre-loxP systems are vital tools for

research, however, there are multiple caveats that should be considered related to off-target effects and

the determination of correct controls. At the bare minimum, Cre-only controls are essential. Additional

tools for $in\ vivo$ genome engineering are required to advance the field. Many studies will need to be re-

interpreted.

**Acknowledgements**

We thank the BC Diabetes Research community and the Johnson laboratory members for valuable

feedback and discussions at local meetings. We thank Dr. Liping Yu and his team at the Insulin Antibody

Core Lab at the University of Colorado, Barbara Davis Diabetes Centre for insulin auto-antibody

analysis.  We thank Dr. Cara Ellis, at University of Alberta for helpful statistics discussions and analysis

of survival/ diabetes incidence data. We thank the amazing animal care services staff for their daily

attention and care of our NOD mice housed at UBC, especially during the challenging COVID-19

pandemic.

**Data Availability**

Data generated during and/or analyzed during the current study are not publicly available but are

available from the corresponding author on reasonable request.

**References**

1.  Daniel D, Gill RG, Schloot N, Wegmann D. Epitope specificity, cytokine production profile and diabetogenic activity of insulin-specific T cell clones isolated from NOD mice. *Eur J Immunol*. 1995;25(4):1056-1062.
2.  Thebault-Baumont K, Dubois-Laforgue D, Krief P, Briand JP, Halbout P, Vallon-Geoffroy K, Morin J, Laloux V, Lehuen A, Carel JC, Jami J, Muller S, Boitard C. Acceleration of type 1 diabetes mellitus in proinsulin 2-deficient NOD mice. *J Clin Invest*. 2003;111(6):851-857.
3.  Atkinson MA, Eisenbarth GS, Michels AW. Type 1 diabetes. *Lancet*. 2014;383(9911):69-82.
4.  Ehlers MR. Immune interventions to preserve beta cell function in type 1 diabetes. *J Investig Med*. 2016;64(1):7-13.
5.  Makino S, Kunimoto K, Muraoka Y, Mizushima Y, Katagiri K, Tochino Y. Breeding of a non-obese, diabetic strain of mice. *Jikken Dobutsu*. 1980;29(1):1-13.
6.  Melanitou E, Devendra D, Liu E, Miao D, Eisenbarth GS. Early and quantal (by litter) expression of insulin autoantibodies in the nonobese diabetic mice predict early diabetes onset. *J Immunol*. 2004;173(11):6603-6610.
7.  Gregori S, Giarratana N, Smiroldo S, Adorini L. Dynamics of pathogenic and suppressor T cells in autoimmune diabetes development. *J Immunol*. 2003;171(8):4040-4047.
8.  Ize-Ludlow D, Lightfoot YL, Parker M, Xue S, Wasserfall C, Haller MJ, Schatz D, Becker DJ, Atkinson MA, Mathews CE. Progressive erosion of beta-cell function precedes the onset of hyperglycemia in the NOD mouse model of type 1 diabetes. *Diabetes*. 2011;60(8):2086-2091.
9.  Mathews CE, Xue S, Posgai A, Lightfoot YL, Li X, Lin A, Wasserfall C, Haller MJ, Schatz D, Atkinson MA. Acute Versus Progressive Onset of Diabetes in NOD Mice: Potential Implications for Therapeutic Interventions in Type 1 Diabetes. *Diabetes*. 2015;64(11):3885-3890.
10. Moriyama H, Abiru N, Paronen J, Sikora K, Liu E, Miao D, Devendra D, Beilke J, Gianani R, Gill RG, Eisenbarth GS. Evidence for a primary islet autoantigen (preproinsulin 1) for insulitis and diabetes in the nonobese diabetic mouse. *Proc Natl Acad Sci U S A*. 2003;100(18):10376-10381.
11. Mehran AE, Templeman NM, Brigidi GS, Lim GE, Chu KY, Hu X, Botezelli JD, Asadi A, Hoffman BG, Kieffer TJ, Bamji SX, Clee SM, Johnson JD. Hyperinsulinemia drives diet-induced obesity independently of brain insulin production. *Cell Metab*. 2012;16(6):723-737.
12. Fan Y, Rudert WA, Grupillo M, He J, Sisino G, Trucco M. Thymus-specific deletion of insulin induces autoimmune diabetes. *Embo J*. 2009;28(18):2812-2824.
13. Martin-Pagola A, Pileggi A, Zahr E, Vendrame F, Damaris Molano R, Snowhite I, Ricordi C, Eisenbarth GS, Nakayama M, Pugliese A. Insulin2 gene (Ins2) transcription by NOD bone marrow-derived cells does not influence autoimmune diabetes development in NOD-Ins2 knockout mice. *Scand J Immunol*. 2009;70(5):439-446.
14. Botezelli JD, Overby P, Lindo L, Wang S, Haida O, Lim GE, Templeman NM, Pauli JR, Johnson JD. Adipose depot-specific upregulation of Ucp1 or mitochondrial oxidative complex proteins are early consequences of genetic insulin reduction in mice. *Am J Physiol Endocrinol Metab*. 2020;319(3):E529-E539.
15. Babaya N, Nakayama M, Moriyama H, Gianani R, Still T, Miao D, Yu L, Hutton JC, Eisenbarth GS. A new model of insulin-deficient diabetes: male NOD mice with a single copy of Ins1 and no Ins2. *Diabetologia*. 2006;49(6):1222-1228.
16. Zhang AMY, Magrill J, de Winter TJJ, Hu X, Skovso S, Schaeffer DF, Kopp JL, Johnson JD. Endogenous Hyperinsulinemia Contributes to Pancreatic Cancer Development. *Cell Metab*. 2019;30(3):403-404.
17. Elso CM, Scott NA, Mariana L, Masterman EI, Sutherland APR, Thomas HE, Mannering SI. Replacing murine insulin 1 with human insulin protects NOD mice from diabetes. *PLoS One*. 2019;14(12):e0225021.

18. Loonstra A, Vooijs M, Beverloo HB, Allak BA, van Drunen E, Kanaar R, Berns A, Jonkers J.
Growth inhibition and DNA damage induced by Cre recombinase in mammalian cells. *Proc*
*Natl Acad Sci U S A*. 2001;98(16):9209-9214.
19. Silver DP, Livingston DM. Self-excising retroviral vectors encoding the Cre recombinase
overcome Cre-mediated cellular toxicity. *Mol Cell*. 2001;8(1):233-243.
20. Pfeifer A, Brandon EP, Kootstra N, Gage FH, Verma IM. Delivery of the Cre recombinase by a
self-deleting lentiviral vector: efficient gene targeting in vivo. *Proc Natl Acad Sci U S A*.
2001;98(20):11450-11455.
21. Baba Y, Nakano M, Yamada Y, Saito I, Kanegae Y. Practical range of effective dose for Cre
recombinase-expressing recombinant adenovirus without cell toxicity in mammalian cells.
*Microbiol Immunol*. 2005;49(6):559-570.
22. Lexow J, Poggioli T, Sarathchandra P, Santini MP, Rosenthal N. Cardiac fibrosis in mice
expressing an inducible myocardial-specific Cre driver. *Dis Model Mech*. 2013;6(6):1470-1476.
23. Bersell K, Choudhury S, Mollova M, Polizzotti BD, Ganapathy B, Walsh S, Wadugu B, Arab S,
Kuhn B. Moderate and high amounts of tamoxifen in alphaMHC-MerCreMer mice induce a
DNA damage response, leading to heart failure and death. *Dis Model Mech*. 2013;6(6):1459-
1469.
24. Li Y, Choi PS, Casey SC, Felsher DW. Activation of Cre recombinase alone can induce
complete tumor regression. *PLoS One*. 2014;9(9):e107589.
25. Sofi MH, Gudi R, Karumuthil-Melethil S, Perez N, Johnson BM, Vasu C. pH of drinking water
influences the composition of gut microbiome and type 1 diabetes incidence. *Diabetes*.
2014;63(2):632-644.
26. Villarreal D, Pradhan G, Wu CS, Allred CD, Guo S, Sun Y. A Simple High Efficiency Protocol
for Pancreatic Islet Isolation from Mice. *J Vis Exp*. 2019(150).
27. Bankhead P, Loughrey MB, Fernandez JA, Dombrowski Y, McArt DG, Dunne PD, McQuaid S,
Gray RT, Murray LJ, Coleman HG, James JA, Salto-Tellez M, Hamilton PW. QuPath: Open
source software for digital pathology image analysis. *Sci Rep*. 2017;7(1):16878.
28. Lee JTC, Shanina I, Chu YN, Horwitz MS, Johnson JD. Carbamazepine, a beta-cell protecting
drug, reduces type 1 diabetes incidence in NOD mice. *Sci Rep*. 2018;8(1):4588.
29. Kuglin B, Gries FA, Kolb H. Evidence of IgG autoantibodies against human proinsulin in
patients with IDDM before insulin treatment. *Diabetes*. 1988;37(1):130-132.
30. Yu L, Robles DT, Abiru N, Kaur P, Rewers M, Kelemen K, Eisenbarth GS. Early expression of
antiinsulin autoantibodies of humans and the NOD mouse: evidence for early determination of
subsequent diabetes. *Proc Natl Acad Sci U S A*. 2000;97(4):1701-1706.
31. Robles DT, Eisenbarth GS, Dailey NJ, Peterson LB, Wicker LS. Insulin autoantibodies are
associated with islet inflammation but not always related to diabetes progression in NOD
congenic mice. *Diabetes*. 2003;52(3):882-886.
32. Thyagarajan B, Guimaraes MJ, Groth AC, Calos MP. Mammalian genomes contain active
recombinase recognition sites. *Gene*. 2000;244(1-2):47-54.
33. Karimova M, Abi-Ghanem J, Berger N, Surendranath V, Pisabarro MT, Buchholz F. Vika/vox, a
novel efficient and specific Cre/loxP-like site-specific recombination system. *Nucleic Acids Res*.
2013;41(2):e37.
34. Pepin G, Ferrand J, Honing K, Jayasekara WS, Cain JE, Behlke MA, Gough DJ, BR GW,
Hornung V, Gantier MP. Cre-dependent DNA recombination activates a STING-dependent
innate immune response. *Nucleic Acids Res*. 2016;44(11):5356-5364.
35. Barber GN. STING: infection, inflammation and cancer. *Nat Rev Immunol*. 2015;15(12):760-
770.
36. Ishikawa H, Barber GN. STING is an endoplasmic reticulum adaptor that facilitates innate
immune signalling. *Nature*. 2008;455(7213):674-678.
37. Serreze DV, Wasserfall C, Ottendorfer EW, Stalvey M, Pierce MA, Gauntt C, O'Donnell B,
Flanagan JB, Campbell-Thompson M, Ellis TM, Atkinson MA. Diabetes acceleration or
prevention by a coxsackievirus B4 infection: critical requirements for both interleukin-4 and
gamma interferon. *J Virol*. 2005;79(2):1045-1052.

38. Qiao J, Zhang Z, Ji S, Liu T, Zhang X, Huang Y, Feng W, Wang K, Wang J, Wang S, Meng
ZX, Liu M. A distinct role of STING in regulating glucose homeostasis through insulin sensitivity
and insulin secretion. *Proc Natl Acad Sci U S A*. 2022;119(7).

**Figure Legends**

**Figure 1. *Ins1* replacement with Cre and Neo protects female NOD mice from type 1 diabetes.**

**(A)** Structure of the wildtype (WT) *Ins1* locus, recombinant alleles result of the neomycin (neo) targeting

vector, or the *Ins1* Cre locus. Created with Biorender.com. **(B)** Overview of breeding strategy. Created

with Biorender.com. **(C)** Kaplan-Meier plot denoting diabetes incidence in NOD mice by *Ins1* genotype,

including $Ins1^{WT/WT}$, $Ins1^{Cre/WT}$, $Ins1^{Neo/WT}$ and $Ins1^{Neo/Cre}$. Survival analysis was performed using Log-

rank (Mantel-Cox) test (*p* value shown). **(D,E)** Individual and mean random blood glucose of female

mice. The mean blood glucose of the $Ins1^{Neo/Cre}$ was significantly lower than that of the $Ins1^{WT/WT}$

colonies, with an adjusted p-value of 0.0023. Mean blood glucose of the $Ins1^{Neo/WT}$ colony was higher

than those of the $Ins1^{WT/WT}$ and $Ins1^{Cre/WT}$ colonies, with adjusted p-values of 0.0002 and 0.0058

respectively. **(F,G)** Individual and mean body mass traces female mice. The mean body mass of the

$Ins1^{Neo/Cre}$ mice was significantly lower compared to the $Ins1^{Cre/WT}$ colony (adjusted p-value<0.0001). The

mean blood glucose of the $Ins1^{Cre/WT}$ was also higher than the $Ins1^{WT/WT}$ and the $Ins1^{Cre/WT}$ colonies, both

with an adjusted p-value<0.0001. Error bars represent SEM

**Figure 2. Effects of *Ins1* replacement with Cre and Neo in male NOD mice.**

**(A)** Kaplan-Meier plot denoting diabetes incidence in NOD mice by *Ins1* genotype. **(B,C)** Individual and

mean random blood glucose in male mice. **(D,E)** Individual and mean body mass traces in male mice .

Error bars represent SEM.

**Figure 3. Insulitis scoring in female NOD mice with *Ins1* replacement.**

**(A)** Plasma insulin concentration Nanomolar (nM) collected at 14 weeks by cardiac puncture. **(B)** Mouse

insulin autoantibodies collected at 14 weeks by cardiac puncture, result of IAA is expressed as an index,

against internal standard positive and negative controls. **(C)** Representative images of H&E stained

pancreata used for insulitis scoring. Scale bars are 100 μm. **(D)** Mean percent insulitis scores at 12-

week-old of each genotype, categorized by score, 0 - No Insulitis, 1 - Peri-Insulitis (<25%), 2 - Insulitis

(25-75%) and 3 - Severe Insulitis (>75%) respectively **(E)** 1-year-old time points. Error bars represent

SEM.

**Figure 4. Gating strategy for flow cytometry of pancreatic lymph node and spleen cells.**

**(A)** Singlets were obtained with use of FSC-A x FSC-H parameters and viable cells were identified by

selecting viability dye negative cells for subsequent analysis. The populations were subsequently split

into three groups of interest with dendritic cells identified by CD11c, B cells identified by CD19, and T

cells identified by CD3. T cells were further categorized into cytotoxic and helper phenotypes with use

of CD4 and CD8 markers and their respective single marker populations were assessed for activation

and priming status (naïve, effector, memory) with the use of CD69, CD44, and CD62L. Regulatory T

cell populations were further selected for with the use of a Foxp3 marker.

**Figure 5. Immune profiling in female NOD mice with *Ins1* replacement**

Flow cytometric analysis of cell populations within the pancreatic lymph node and spleen at 50 weeks

of age. **(A)** Percentage of CD3+ positive T cells and CD19+ B cells from the pancreatic lymph node. **(B)**

Percentage of CD11+ Dendritic cells from the pancreatic lymph node. **(C)** Percentage of CD8+CD4-

Cytotoxic T cells, CD8+CD4- T helper cells and CD4+Foxp3+ Treg cells from the pancreatic lymph

node. **(D)** Percentage of CD8+CD4+ immature cells and CD8-CD4- DN T cells from the pancreatic

lymph node. **(E)** Percentage of CD69+ Activated cytotoxic T cells from the pancreatic lymph node. **(F)**

CD44loCD62L + Naive cytotoxic T cells from the pancreatic lymph node. **(G)** Percentage of

CD44hiCD62 – Memory cytotoxic T cells from the pancreatic lymph node. **(H)** Percentage of

CD44midCD62 – Effector cytotoxic T cells from the pancreatic lymph node. **(I)** Percentage of CD69+

Activated T helper cells from the pancreatic lymph node. **(J)** CD44loCD62L + Naive T helper cells from the pancreatic lymph node. **(K)** Percentage of CD44hiCD62 + Memory T helper cells from the pancreatic lymph node. **(L)** Percentage of CD44midCD62 – Effector T helper cells from the pancreatic lymph node. **(M)** Percentage of CD3+ positive T cells and CD19+ B cells from the Spleen. **(N)** Percentage of CD11+ Dendritic cells from the spleen. (**O**) Percentage of CD8+CD4- Cytotoxic T cells, CD8+CD4- T helper cells and CD4+Foxp3+ Treg cells from the spleen. **(P)** Percentage of CD8+CD4+ immature cells and CD8-CD4- DN T cells from the spleen. **(Q)** Percentage of CD69+ Activated cytotoxic T cells from the spleen. **(R)** CD44loCD62L + Naive cytotoxic T cells from the spleen. **(S)** Percentage of CD44hiCD62 – Memory cytotoxic T cells from the spleen. **(T)** Percentage of CD44midCD62 – Effector cytotoxic T cells from the spleen. **(U)** Percentage of CD69+ Activated T helper cells from the spleen. **(V)** CD44loCD62L + Naive T helper cells from the spleen. **(W)** Percentage of CD44hiCD62 + Memory T helper cells from the spleen. **(X)** Percentage of CD44midCD62 – Effector T helper cells from the spleen. Error bars represent SEM.

**Figure 6. *Ins1* replacement with Cre protects female NOD mice from type 1 diabetes in an independent facility.**

**(A)** Kaplan-Meier plot denoting diabetes incidence. **(B,C)** Individual and mean random blood glucose of female NOD colonies from a second, independent site. The mean blood glucose of the *Ins1*[Neo/Cre] (green) was significantly lower than that of the *Ins1*[WT/WT] (blue) littermates, with an adjusted p-value < 0.05. Female NOD non-littermate controls used to track overall diabetes incidence in the colony are listed in red. Error bars represent SEM.

**Table 1. Summary of Antibodies used for Flow cytometry**

| Antibody | Manufacturer | Colour | Laser (nM) | Filter | Catalogue# | RRID |
|---|---|---|---|---|---|---|
| CD3 | ThermoFisher | e-Flour 540 | 405 | 450/45 | 48-003382 | AB_2016704 |
| CD4 | ThermoFisher | BV650 | 405 | 660/10 | 64-0042-82 | AB_2662401 |
| CD8 | BioLegend | PE-TR | 561 | 610/20 | 100762 | AB_2564027 |

| CD11c | ThermoFisher | PE | 561 | 585/42 | 12-0114-81 | AB_465551 |
|-------|--------------|-----|-----|--------|------------|-----------|
| CD19 | ThermoFisher | SB780 | 405 | 763/43 | 78-0193-82 | AB_2722936 |
| CD44 | ThermoFisher | APC | 633 | 660/10 | 17-0441-82 | AB_469390 |
| CD62L | ThermoFisher | PerCP-Cy5.5 | 488 | 690/50 | 45-0621-82 | AB_996667 |
| CD69 | ThermoFisher | FITC | 488 | 525/40 | 11-0692-82 | AB_11069282 |
| Foxp3 | ThermoFisher | AF-700 | 633 | 712/25 | 565773-82 | AB_1210557 |

**A**

Ins1^WT — Exon 1 — Exon 2

Ins1^neo — neo

Ins1^cre — Exon 1 — cre

**B**

♂ Ins1^Neo/WT  X  Ins1^Cre/WT ♀

Ins1^WT/WT; 100% Ins1 dosage
Ins1^Cre/WT; 50% Ins1 dosage
Ins1^Neo/WT; 50% Ins1 dosage
Ins1^Neo/Cre; 0% Ins1 dosage

**C**

| Comparison | p-value |
|---|---|
| Ins1^wt/wt vs. Ins1^Cre/wt | **0.0325** |
| Ins1^wt/wt vs. Ins1^Neo/wt | 0.4030 |
| Ins1^wt/wt vs. Ins1^Cre/Neo | **0.0189** |
| Ins1^Cre/wt vs. Ins1^Neo/wt | 0.0801 |
| Ins1^Cre/wt vs. Ins1^Cre/Neo | 0.983 |
| Ins1^Neo/wt vs. Ins1^Cre/Neo | 0.0388 |

Ins1^wt/wt (n=14)
Ins1^cre/wt (n=11)
Ins1^neo/wt (n=14)
Ins1^neo/cre (n=14)

A

*Michigan cohort*

♂ *Ins1*<sup>wt/wt</sup> x *Ins1*<sup>cre/wt</sup> ♀

*Ins1*<sup>cre/wt</sup>                                    (n=4)
*Ins1*<sup>wt/wt</sup> Littermates              (n=7)
*Ins1*<sup>wt/wt</sup> Non-Littermates    (n=11)