1 TITLE
2
3 # A heterogeneous drug tolerant persister state in BRAF-mutant
4 melanoma is characterized by ion channel dysregulation and
5 susceptibility to ferroptosis
6
7 AUTHORS
8

9 Corey E. Hayford[1], Philip E. Stauffer[1], Blake Baleami[1], B. Bishal Paudel[1,2], Darren R. Tyson[3], Aziz
10 Al'Khafaji[4,5], Kaitlyn E. Johnson[5], Leonard A. Harris[6,7,8], Amy Brock[4,5], and Vito Quaranta[3,9]
11
[1]Chemical and Physical Biology Graduate Program, Vanderbilt University School of Medicine, Nashville, TN USA
[2]Department of Biomedical Engineering, University of Virginia, Charlottesville, VA USA
[3]Department of Biochemistry, Vanderbilt University School of Medicine, Nashville, TN USA
[4]Institute for Cellular and Molecular Biology, University of Texas at Austin, TX USA
[5]Department of Biomedical Engineering, University of Texas at Austin, TX USA
[6]Department of Biomedical Engineering, University of Arkansas, Fayetteville, AR USA
[7]Interdisciplinary Graduate Program in Cell and Molecular Biology, University of Arkansas, Fayetteville, AR USA
[8]Cancer Biology Program, Winthrop P. Rockefeller Cancer Institute, University of Arkansas for Medical Sciences, Little
Rock, AR USA
[9]Department of Pharmacology, Vanderbilt University School of Medicine, Nashville, TN USA

22
23 ABSTRACT
24

There is increasing interest in cancer cell subpopulations that can withstanding treatment via
non-genetic mechanisms, such as tumor cell plasticity and adaptation. These cell populations
may be comprised of cells with diverse phenotypes, e.g., quiescent or slow cycling. Such
populations have been broadly termed "drug-tolerant persisters" (DTPs) and may be responsible
for minimal residual disease following anticancer treatment and acquired resistance.
Understanding molecular mechanisms that drive emergence of DTPs could lead to new strategies
to improve therapeutic outcomes. Recently, we reported that BRAF-mutant melanoma cells
under prolonged BRAF inhibition enter a DTP state with balanced cell death and division, which
we termed "idling." Here, we apply single cell barcoding to show that idling DTP populations
emerge via cell state transitions, rather than selection of a few pre-existing drug-tolerant clones.
Within the time frame of our experiments, DTPs exhibit varying proportions of fast- and slow-
cycling cells within each lineage, suggesting that entry into the DTP state is a stochastic process.
Furthermore, single-cell transcriptomics and bulk epigenomics reveal common gene expression
and ontology signatures in DTP lineages that are consistent with rebalancing of ion channels.
Calcium flux experiments uncover a reduction of divalent cation reserves in intracellular
organelles, likely leading to endoplasmic reticulum stress. Accordingly, idling DTPs are more
prone to ferroptotic cell death, as indicated by increased sensitivity to inhibition of glutathione
peroxidase 4 (GPX4), which prevents removal of toxic lipid peroxides. In summary, we propose
that ion channel homeostasis is a central process underlying idling DTP emergence in BRAF-
mutated melanoma. Future studies will investigate translational aspects of this insight.

INTRODUCTION

Cancer is a complex and dynamic disease characterized by intratumoral cell-to-cell variability that has been implicated in treatment evasion and acquired resistance to therapy.[1,2] Tumor cells exhibit multiple forms of variability, due to a complex interplay between genetics, epigenetics, and other non-genetic sources of variation.[3] Historically, genetic variability has received the most attention, with pre-existing or acquired genetic resistance mutations implicated as the primary cause of treatment resistance and tumor relapse. However, recently, epigenetic factors have become increasingly recognized as major sources of tumor heterogeneity and consequent treatment failure.[4–6] Epigenetic heterogeneity, defined as co-existence of multiple distinct phenotypes within a population of genetically identical cells,[6] may result in long-term drug tolerance, from which new genetic resistance mutations can ultimately arise.[7,8] As such, it has been suggested that a window of opportunity might exist between the onset of drug tolerance and the acquisition of genetic resistance mutations during which a targeted secondary treatment can be deployed to further reduce, or even eliminate, the residual tumor mass.[7,8] Accomplishing this will require a detailed understanding of the molecular drivers underlying drug tolerance so that potential vulnerabilities can be identified and exploited clinically.

Multiple investigators have reported cancer cell subpopulations capable of withstanding drug treatments via non-genetic mechanisms. These populations have been broadly termed "drug-tolerant persisters" (DTPs) and described, variously, as either quiescent[9] or slow cycling.[10–12] Recently, we described a similar drug-tolerant state, termed "idling," that arises in BRAF-mutant melanoma cell populations under prolonged BRAF inhibition.[7,13] Importantly, in contrast to previously reported DTPs, the idling phenotype refers to the state of a cell population, rather than of individual cells. More precisely, we found that the growth dynamics of drug-treated BRAF-mutant melanoma cell populations were best described by a model in which cells are distributed across multiple phenotypic states, each with a distinct proliferation rate.[7] In the idling state, cells are proportionally distributed across these states, such that the proliferation rate of the population as a whole is approximately zero. Given the significant interest in drug tolerance as a precursor to treatment failure and acquired resistance in tumors,[14] recognizing that DTP populations may, in fact, be heterogenous, composed of multiple phenotypic states that act collectively to evade and survive drug treatment, is crucial for developing effective treatment strategies against them.

Here, we demonstrate experimentally in BRAF-mutant melanoma that the idling phenotype is a heterogeneous collection of phenotypic states, which is less heterogeneous than the collection of states in untreated populations. This reduction in heterogeneity under drug pressure suggests idling cells may potentially be vulnerable to a targeted secondary treatment. Single-cell RNA sequencing demonstrates that idling populations comprise multiple distinct transcriptomic states that are more similar to each other than the states observed in untreated populations. Furthermore, we use DNA barcoding to show that the idling state is populated by cells from most lineages present in the untreated population, suggesting that phenotypic state transitions, rather

than clonal selection, underlies the establishment of drug tolerance. A gene ontology (GO)
analysis, combining data from both transcriptomics and epigenomics, indicates that ion channel
expression is significantly altered in idling cells relative to untreated cells, pointing to a role for
mitochondrial metabolism in the idling phenotype. This is supported by calcium flux assays that
show store-operated calcium entry (SOCE) is significantly altered in idling cells. Finally, idling cell
populations are shown to have increased susceptibility to death by ferroptosis, supporting the
notion that DTPs are vulnerable to targeted secondary treatments.
93
94 RESULTS
95
96 **Drug-tolerant melanoma populations include cells from almost all untreated lineages**
97
We have previously shown that BRAF-mutant melanoma cell lines exhibit non-linear drug
response dynamics to BRAF inhibitor (BRAFi). The initial phase of drug response is characterized
by variable degrees of proliferation rate reduction. This is followed by entry of the treated cell
population into a state of near-zero proliferation rate, termed idling.[7] To determine whether the
idling state populations are comprised of a collection of DTP clones, we barcoded the BRAF-
mutant SKMEL5 melanoma cell line with a gRNA barcoding library.[15] This approach allowed for
high-depth coverage of dynamics for multiple lineages in response to selective pressures (i.e.,
drug treatment) and the ability to distinguish between clonal selection and phenotypic state
transitions (FIG. 1A). Upon treatment with BRAFi PLX 4720 (8 μM for 8 days), the barcode library
complexity was reduced by less than 10% (FIG. 1B). These surviving barcodes were shared in a
large majority of replicates within each condition (untreated and idling; FIG. 1C), as well as
between conditions (FIG. 1D). These results indicate that most clonal lineages survive treatment
and persist at a similar relative proportion after BRAFi treatment. Simply, the idling state consists
of cells from an overwhelming majority of the untreated population, providing direct evidence
against clonal selection in response to BRAFi.
113
Some fluctuations in relative barcode abundance do exist after treatment with BRAFi (FIG. 1E,
see next section). The underlying fold change distribution for the entire barcoded cell population
from untreated to idling is a normal distribution centered at zero, with no obvious exceptions;
the top 25 most populous barcodes reflect this distribution and are used for downstream
analyses (FIG. 1F). Notably, nearly all the lineages that do not survive treatment come from clones
that have an exceedingly small representation in the overall distribution, suggesting that the loss
of those lineages is due to random chance. These results indicate that phenotypic state
transitions likely dominate the response to BRAFi, leading a diverse set of clonal lineages to adopt
a phenotypic state(s) akin to DTPs.
123
124 **Melanoma populations become less transcriptomically heterogeneous following prolonged**
125 **BRAF inhibition**
126
We performed single-cell transcriptomics on the barcoded cell line in untreated and in idling
conditions (see Methods). Using Uniform Manifold Approximation and Projection[16] (UMAP) to
project the transcriptomic profiles for all cells into two dimensions, we see that untreated and

idling cells clearly fall in different regions of the UMAP space, with minimal overlap (FIG. 2A).
Additionally, the idling cells reside in a more constrained region of the space, while in the
untreated cells two distinct clusters are evident.
133
To quantify the variability within and between populations, pairwise distances were calculated
between cells in each condition (FIG. 2B). Nearly all idling cells fall closer to one another than
untreated cells fall to each other in the principal component analysis (PCA) space using 10
principal components. The Earth Mover's Distance (EMD) was calculated between the two
pairwise distance distributions (EMD = 4.38) to measure the separation between the untreated
and idling populations. In summary, the untreated population is more heterogeneous than the
idling one in transcriptomic space.
141
Next, we aimed to determine the biological factors that differentiate the axes that separate
population clusters. UMAP_2 (e.g., separation between untreated clusters) largely separates
based on metabolic processes, consistent with our previous report of metabolic heterogeneity in
untreated melanoma cells[7] (Supplementary FIG. S1A). Specifically, the large untreated (UT_L)
cluster is enriched in a glycolysis gene signature compared to the small untreated (UT_S) cluster
(see Methods, Supplementary FIG. S1D). Conversely, the idling population is enriched in an
oxidative phosphorylation gene signature compared to both untreated clusters (Supplementary
FIG. S1D). This is consistent with multiple reports in the literature, where BRAF-mutant cells rely
primarily on glycolysis for growth and energy production, and BRAFi interferes with glycolytic
processes, presumably favoring a switch to oxidative phosphorylation (see Discussion).
152
Alternatively, UMAP_1 primarily separates based on cell cycle stage, even after cell cycle gene
signature regression, within the context of each cluster (Supplementary FIG. S1B). We used a
previously established cell cycle gene signature (see Methods) to classify individual cells by cell
cycle stage (G1, G2M, S). We then projected cell cycle stage on cells in the single-cell
transcriptomics UMAP space (FIG. 2C). For visualization clarity, cell cycle stages were simplified
into non-cycling (G1) and cycling (G2/M/S) states. The number and proportion of cells in cycling
vs. non-cycling states were calculated (FIG. 2D), and the number of cycling cells is dramatically
decreased in the treated population (67% vs 15%).
161
162   **Changes in barcode abundance between untreated and drug-tolerant populations can be**
163   **explained by simple random effects, not clonal selection**
164
The barcoding system we adopted integrates droplet cell barcodes with transduced gRNA lineage
barcodes, establishing a direct connection between the abundance of clonal lineages and their
location in the single-cell transcriptomics state space. The barcoding library was designed with a
relative low complexity (~65k potential barcodes, ~425 identified barcoded lineages), which
allowed for multiple instances of the same lineage barcode in gene expression space. We
projected the lineage distribution of SKMEL5 barcoded cells onto scRNA-seq UMAP space
(examples in FIG. 3A). Barcodes exhibit various UMAP transcriptomic state occupancies and fall
in both clusters (large and small) and cycling states across treatment conditions. In the untreated
clusters, barcodes tend to have an approximately similar breakdown between small and large

clusters as the average of the population. However, in the idling cell cluster, the breakdown
between non-cycling and cycling cell states tends to be predictive of clonal lineage drug response
(FIG. 1E; FIG. 3B). From these examples, three key behaviors emerge. The first is exemplified by
barcodes 5 and 9, where only a few cells (~5%) from the barcoded cell lineages fall in the cycling
idling state region of the cluster. This low proportion of cycling cells potentially leads to a
reduction in the relative proportion of these barcodes post-treatment (FIG. 1E-F). The second
behavior, as seen in barcodes 2 and 13, is the opposite, where an increased proportion of cells in
each barcode fall in the idling cycling state (~23%). This increase in cells in the cycling state
potentially leads to an increase in the relative proportion of both lineages after treatment (FIG.
1E-F). The third is characteristic of most lineages and falls somewhere in between, which causes
a middling effect on treatment response (Supplementary FIG. S2). Thus, a strong correlation
(Pearson, R=0.55) exists between the proportion of cells in the cycling idling state and the relative
fold change of barcode abundance after treatment (FIG. 3C), indicating that this is a shared
feature of all barcoded cell lineages.
188
189 **Multi-omics analysis points to ion channel dysregulation as a central factor in drug tolerance**
190
Due to the time scale of the appearance of DTPs, it is generally accepted that DTPs may arise by
epigenetic regulation rather than the accumulation of mutations.[9–12] To determine whether this
is the case for transitions to the idling state, we performed bulk ATAC-seq on SKMEL5 cells before
and after BRAFi treatment (see Methods). In each condition, we identified a fragment size
distribution characteristic of a successful run (Supplementary FIG. S3B; library complexity in
Supplementary FIG. S3A). Regions of open chromatin were enriched in cells from both conditions,
resulting in unique and shared peaks between conditions (see Methods; FIG. 4A). These peaks
were normalized to the transcription start site (TSS), and the distribution of binding loci
(Supplementary FIG. S3C) was used to quantify the peak feature set (FIG. 4B). Idling peaks have
much fewer proximal features (e.g., promoter regions) and more distal elements (e.g., distal
intergenic and intronic regions) compared to the untreated condition. Distal elements have been
known to be involved in short-term epigenetic regulation. Unique peaks were also assigned to a
corresponding gene, and genes associated with unique peaks were input into a GO over-
representation analysis, which identified ion transport and activity as differentiators of idling
(FIG. 4C).
206
To confirm this finding independently, we subjected clonal sublines[7] of SKMEL5 (SC01, SC07, and
SC10) to bulk RNA sequencing (RNA-seq) over the course of BRAFi treatment (0, 3, and 8 days
post-BRAFi). Even though these sublines have marked short-term differences in terms of BRAFi
response,[7] they were chosen because they all converge to a long-term near-zero proliferation
rate characteristic of idling (3-8 days). To understand the time evolution of clonal subline
response to BRAFi, normalized counts of each subline were projected into reduced
dimensionality space by PCA (FIG. 4D).
214
On day 0, prior to BRAFi treatment, sublines are distinct on the first PCA axis (PC1), with SC07
and SC10 showing the most similarity. In short-term treatment (day 0 → 3), sublines
predominately change on PC2 and maintain the overall variance between sublines observed at

day 0 (FIG. 4D). In the long-term response (day 3 → 8), the sublines begin to converge on both
PC1 and PC2 (FIG. 4D). Considering that these three sublines are clonal derivatives of the SKMEL5
population, this result is consistent with the *en masse* transition of single cells to a common
UMAP space after treatment (FIG. 3A).
222
A differential expression analysis was performed between untreated (day 0) and idling (day 8)
across all three clones to identify transcriptomic signatures characteristic of idling. Differentially
expressed genes were input into a GO over-enrichment analysis to reveal processes upregulated
in the idling state. GO terms associated with ion transport and homeostasis were also found to
be upregulated in idling cells (MF – FIG. 4E; BP – Supplementary FIG. S3D; CC – Supplementary
FIG. S3E). In fact, statistically enriched GO terms shared a strong correlation between data
modalities, with the most significant terms pointing to the ion channel activity shown in previous
analyses (FIG. 4C and 4E). In summary, both epigenomics and transcriptomics data point toward
ion channel activity as a major molecular determinant of the idling state.
232
233   **Intracellular calcium ion flux is significantly reduced in drug-tolerant melanoma populations**
234
Next, we experimentally tested calcium channel activity in idling versus untreated cells, using a
calcium flux assay that measures the amount of ER resident calcium and propensity of SOCE
activity. In this assay, cyclopiazonic acid (CPA) is used to inhibit the activity of Sarcoendoplasmic
Reticulum Calcium ATPases (SERCAs), leading to a release of free calcium from the ER to the
cytoplasm where it is detected by the calcium dye, Fluo-8-AM. (first peak, FIG. 5A). To replenish
ER calcium upon depletion, cells undergo store operated calcium entry (SOCE) which brings in
extracellular calcium that can then be pumped into the ER by SERCAs. In order to isolate ER
calcium release from SOCE activity, CPA is added in a calcium free buffer, followed by addition of
calcium to the assay (second peak, FIG. 5A). ER calcium release stimulates the opening of plasma
membrane resident calcium channels (e.g., ORAI), but calcium cannot flow in until it is added to
the assay buffer, allowing ER calcium content/release and propensity of SOCE activity to be
studied separately. Idling cells exhibited decreased SOCE activity compared to untreated cells,
demonstrated by the second peak (FIG. 5A). These results verify differences in ion channel
activity in idling cells, and further suggest ER stress may be a vulnerability of idling cells.
249
To see if these differences in calcium flux may be a result of gene expression changes driven by
BRAFi treatment, a differential gene expression analysis was conducted on bulk RNAseq data
obtained from three SKMEL5 clonal sublines (FIG. 5B). Many of the queried calcium handling
genes are differentially expressed in sublines after 3 days and 8 days of BRAF inhibition.
Interestingly, trends in gene expression patterns seem relatively consistent across the sublines
(FIG. 5B). These results show that calcium handling genes are differentially expressed in cell
populations treated with BRAFi compared to untreated cell populations.
257
258   **Drug-tolerant melanoma populations exhibit increased susceptibility to death via ferroptosis**
259

Previous reports have indicated a connection between ion channels and ferroptosis, a type of
regulated cell death, by way of increased ER stress.[17] Thus, induction of ferroptosis may provide
a potential way to eradicate idling cancer cells. Indeed, expression of several genes in the
ferroptosis signature is altered in melanoma cells upon BRAFi and entry into the idling state (FIG.
6A; Supplementary FIG. S4). Notably, glutathione metabolism gene expression is increased and
genes for polyunsaturated fatty acid (PUFA) enzymes are decreased after BRAFi treatment, along
with differences in various other parts of ferroptosis-related signaling. Interestingly, the
expression pattern results in increased supply to the Fenton reaction, which produces the free
radical precursors to reactive oxygen species (ROS), the step directly before commitment to
ferroptosis (Supplementary FIG. S1E and S4).

To directly test susceptibility to ferroptosis, we subjected both drug-naïve and idling cells to
treatment (see Methods) with RSL3; RSL3 is a compound that induces ferroptosis by targeting
GPX4, a key regulator of glutathione oxidation and PUFA reduction. Interestingly, this treatment
would reduce glutathione metabolism while preventing reduction of PUFA intermediates, leading
to more precursors of the Fenton reaction and potentially to increased ROS that would drive cell
death by ferroptosis. Interestingly, idling cells were ~3-fold more sensitive to RSL3 by potency
(FIG. 6B; difference in IC50s), suggesting idling cells are susceptible to ferroptosis. Furthermore,
addition of ferrostatin-1 (Fer-1), a drug that inhibits the production of lipid peroxides by the
Fenton reaction (i.e., ROS), rescued the differential drug-response behavior of both idling and
untreated cells in the presence of RSL3 (FIG. 6C). These results indicate that idling cells are
vulnerable to ferroptosis, presumably due to the increased ROS that results from sequential GPX4
inhibition, providing a potential route to DTP elimination.

DISCUSSION

In this study, we show that BRAF inhibition (BRAFi) causes BRAF-mutant melanoma cells to
transition *en masse* into idling populations of plastic drug-tolerant persisters (DTPs), which
exhibit ion channel dysregulation and susceptibility to ferroptosis. Using cellular barcoding, we
showed that idling cells result from an overwhelming majority of untreated clones, rather than
clonal selection of a special idling clone (FIG. 1B-D). Distinct transcriptomic signatures were
identified that differentiate untreated and idling cells, with the idling cells represented in a more
restrained transcriptomic space (FIG. 2A-B). The idling cell population consisted of cells in cycling
and non-cycling states, as defined by their phase in the cell cycle (FIG. 2C); though, the idling
population had a lower proportion of cycling cells in its transcriptomic space than untreated cell
populations (FIG. 2D). Barcoded clonal lineages were distributed across both transcriptomic
states in the untreated condition (FIG. 3A). However, relative barcode abundances for lineages
that have a larger proportion of cells in the cycling versus non-cycling idling state result in a larger
proportion of that barcode in the idling population after treatment (FIG. 3A-C; c.f. FIG. 1E). Bulk
epigenomics and time-series transcriptomics of the SKMEL5 cell line and clonal lineages identify
a convergent idling molecular signature shared across modalities, which points towards ion
dysregulation as a characteristic of the idling state (FIG. 4). The calcium ion channel dysregulation
was validated in a calcium flux assay (FIG. 5A) and a differential gene expression analysis of ion
channels (FIG. 5B). This evidence of calcium ion channel dysregulation further established a

potential connection to ferroptosis, a type of regulated cell death. A ferroptosis gene signature
(FIG. 6A) suggested potential susceptibility to GPX4 inhibition through increased ROS; this
susceptibility was validated to have a clear effect on idling cells (FIG. 6B). Rescue of the drugged
phenotype by a lipid ROS scavenger verified the connection between idling drug tolerance and
susceptibility to ferroptosis (FIG. 6C).
309
Many studies have remarked on the epigenetic plasticity as a way to understand decreased drug
sensitivity,[18] and others have postulated using the epigenetic landscape as a mechanism to
understand how cancer cells transition between states.[19] However, little experimental evidence
exists showing how epigenetic state transitions lead to drug tolerance and eventual resistance.
Together, these data provide evidence for a view of tumor cell plasticity where cells fall into
basins across an epigenetic landscape. This study puts forth data that suggests a timeline for
epigenetic plasticity of cancer cells prior to and after treatment. Tumor initiation from a single
clone creates a population with the same genetic background. The genetic clone emanates an
epigenetic landscape, comprised of several basins of attraction over which cells populate to
create multiple cell types, each with different molecular phenotypes. Over time, cells in the
landscape reach a dynamic equilibrium, i.e., cells can still transition between basins but the
population is in a state of balance, a process known as "bet hedging".[20,21] The introduction of a
perturbation, such as an anticancer drug treatment, upends the equilibrated landscape and drops
cells into a new landscape. Cells re-equilibrate to the new landscape and adopt cell fates
corresponding to the state in which they now reside. In the case of a drug with good efficacy,
most cells will fall into a state of the drug-treated landscape that results in death. However, if the
new landscape includes a state where cells have a positive proliferation rate in drug, the
population will invariably rebound. In our case, most of the cells matriculate into the large non-
cycling state upon treatment with BRAFi, but some end up in a smaller cycling state (FIG. 2C). This
feature is consistent with several different types of DTPs, from quiescent to slow-cycling.[9–12]
However, some barcoded clonal lineages disproportionally fall into one of the two states (FIG.
3A), leading to a differential short-term drug fitness in response to BRAFi. Previous studies in the
lab show that even sublines eventually adopt a near-zero proliferation rate in prolonged BRAFi,[7]
suggesting that lineages fully equilibrate to the new landscape over time. This occurrence is
largely consistent with other oncogene-addicted cancers treated with targeted therapies, which
exhibit a noisy short-term drug tolerance that often leads to long-term genetic resistance.[22,23]
336
An actionable result of a drug-modified landscape is new biochemical network properties that
can lead to new treatment sensitivities. This phenomenon, in which resistance (or tolerance) to
one drug treatment confers sensitivity to a different drug (or drug class), is commonly known as
"collateral sensitivity." Interesting screens have been performed to identify drugs that have
increased sensitivity for cells treated with an initial drug, including in cancer.[24] However, these
methods ignore the heterogeneity present in treated populations and often lead to traditional
up-front combination therapies. These combination therapies create an entirely new epigenetic
landscape that has new molecular properties with unknown vulnerabilities,[6,25] making the new
tumor difficult to treat. The heterogeneity identified in our idling cells exemplifies this problem
and, therefore, requires a different approach. Sequential therapy is an alternative approach in
which cells are able to equilibrate to the new drug-treated landscape with the hope that it is

more sensitive as a whole.[26] Molecular analyses in our study show that the drug-treated
landscape is indeed more homogeneous (FIG. 2A) and has a common thread in ion channel
activity across transcriptomics and epigenomics data (FIG. 4). By targeting the idling state with a
sequential therapy that leverages the increased ion channel activity to promote targeted
ferroptotic cell death, we seemingly create another epigenetic landscape where all of the basins
have a negative proliferation rate in drug treatment and result in tumor eradication. Therefore,
potential future studies could be aimed at identifying vulnerabilities of epigenetic landscapes
that result after primary drug treatment and finding patient-specific secondary drugs used for
sequential drug treatment regimens that eradicate the entire tumor. Together, these results
indicate that BRAF-mutant melanoma cells converge to a confined idling population after BRAFi
treatment, yet still exhibits cycling and non-cycling transcriptomic states. This breakdown
appears to encapsulate previously described DTPs that exhibit various cycling behaviors,
including quiescence, active division, and slow cycling.

361
362 **Figure captions**

363
Figure 1: *Most clonal lineages survive treatment with BRAFi into idling.* (A) Schematic of
example lineage tracing experiments using cellular barcoding. (B) Number of unique barcodes in
each treatment condition. Lines correspond to the means of three experimental replicates
(points). A minimum cutoff of 100 counts per million (CPM) was used. (C) Proportional sharing of
barcodes among experimental replicates (i.e., R1 = replicate 1) for each treatment condition. (D)
Heatmap of relative barcode abundances ($\log_{10}$ CPM) for each experimental replicates across all
captured barcodes. Heatmap is organized by decreasing barcode abundance in untreated
condition. (E) Relative fraction of the top 25 ranked (in untreated) barcoded cell lineages in
untreated and idling conditions. Bar height corresponds to the average of three experimental
replicates (line is standard deviation). (F) Distribution of ($\log_2$) fold change for barcoded clonal
lineages from untreated to idling. Means of fold changes were compiled into a distribution for all
captured lineages (grey), as well as the top 25 most abundant lineages noted in *E*.

376
Figure 2: *Idling cells represent a convergent, yet still heterogeneous, transcriptomic state with
less idling cells in the cycling state than untreated cells. (A) UMAP projection of untreated and
idling single-cell transcriptomes. 6410 cells are shown, with an approximately equal split between
conditions. (B) CDF of pairwise cell distances (random sampling of 15,000) with 10 principal
components on the PCA space of the untreated and idling single-cell transcriptomic data. An
Earth Mover's Distance (EMD) was calculated between the distributions to quantify their
separation from each other. (C) Overlay of cell cycle state (see Methods) on UMAP projection of
single cell transcriptomes (colored contours represent information in *A*)**.** (D) Relative proportion
of cells in cell cycle state for major untreated and idling clusters. The total number of cells in each
cluster (n) is noted above each bar.

387
Figure 3: *Lineage distribution across cell cycle states is reflective of clonal dynamics.* (A)
Projections of lineage transcriptomic distributions on UMAP projection in *FIG. 2A*. Lineages
correspond to colored dots, while contours reflect treatment condition. (B) Proportion of cells in
the idling cycling transcriptomic state for the top 25 most abundant barcodes. Dashed line

represents average of all barcodes. (C) A Pearson correlation analysis was performed between
the percentage of cycling cells in the idling state and the $\log_2$ fold change in barcode abundance
following BRAFi treatment for the top 25 most abundant barcodes.

Figure 4: *Unique elements of the idling transcriptome and epigenome have differential features*
for ion activity. (A) Venn diagram of ATAC-seq identified peaks of open chromatin. (B) Alignment
of peaks to the TSS allows for prediction of epigenomic features. Predicted feature distribution
in untreated, idling, or shared peaks. Peaks were assigned to features based on proximity to TSS
(see Methods). (C) GO over-enrichment analysis of genes associated to unique idling peaks. The
top 10 terms with the largest gene ratio are shown. (D) PCA projection of subclones (SC01, SC07,
and SC10) at multiple times (0, 3, and 8 days) in BRAFi. Each condition (e.g. SC01 at day 0) was
completed in triplicate. Lines are drawn between the centroids of triplicates across the time
series. (E) GO over-representation analysis on differentially expressed genes with increased
expression between untreated and idling. GO terms with the top 10 largest gene ratios (see
Methods) are shown.

Figure 5: *Idling cells have decreased calcium flux and altered expression of calcium handling*
genes. (A) Calcium flux assay of untreated and idling cells using cyclopiazonic acid (CPA) to
deplete ER calcium without extracellular Ca^{2+}, addition of extracellular Ca^{2+} to test store operated
calcium entry (SOCE) activity, and ionomycin to control for number of cells in the assay. (B)
Differential gene expression analysis of calcium handling genes in sublines transcriptomics data
from *4D*.

Figure 6: *Idling cell populations are susceptible to ferroptosis.* (A) Gene expression values (z-
scores; see Methods) for multiple clonal sublines (SC01, SC07, SC10) across the BRAFi treatment
time course (0, 3, and 8 days post treatment). Genes are further grouped by their associated
processes related to ferroptosis. (B) DIP rate dose response curves for untreated and idling cells
treated with ferroptosis inducer RSL3. (C) Same as *B,* except with rescue experiments on the post-
treated cells using ferroptosis inhibitor Fer-1.

Figure S1: *Analysis of the UMAP single cell transcriptomic space.* (A) GO analysis of differentially
expressed between the untreated clusters in the UMAP single cell transcriptomics space shown
in *FIG. 2A.* (B) Same as *A* except the analysis was applied to the idling cluster and a subpopulation
of cells that were predominantly in the cycling state. (C) A representation of UMAP single cell
transcriptomic space of untreated and idling cells used for differential gene expression and GO
analyses in *D* and *E*. (D) Distributions of the signature scores for cells in each cluster in the
differential gene expression analysis from the GO analysis related to glycolysis and oxidative
phosphorylation. Colors of each distribution correspond to the colors of the clusters in *C*, where
pink is cluster 1 (i.e., UT_S), yellow is cluster 2 (i.e., UT_L), and blue is cluster 3 (i.e., idling). (E) Same
as *D* except with other processes.

Figure S2: *Lineage distributions for barcodes with a middle effect in clonal dynamics to*
treatment response. Same as *FIG. 3A* for the remaining top 25 barcodes in the single cell
transcriptomics analysis.

Figure S3: *Quality control of bulk epigenomics ATA-seq data and correlation analysis of differentially expressed genes in idling transcriptome and epigenome.* (A) Plot of unique reads versus sequenced reads for the bulk ATAC-seq data used in epigenome analyses. (B) Insert size distribution of aligned reads from ATAC-seq data on untreated and idling cells. Both conditions follow traditional nucleosome patterning. (C) Peak binding site distribution for untreated, idling, and shared peaks. X-axis represents kb distances from the TSS. (D) Correlation analysis of the differentially expressed genes from a Biological Process (BP)-type of GO analysis for the epigenome and transcriptome of the idling cell population. (E) Same as *D* for a Cellular Component (CC)-type of GO analysis.

Figure S4: *Ferroptosis pathway.* A schematic diagram of the ferroptosis pathway with boxes that correspond to the colors of the groups for the genes in *FIG. 6A*.

MATERIALS AND METHODS

Cell culture and reagents

We chose BRAF-mutant melanoma cell line SKMEL5 as a preferred model, since it exhibits median BRAFi sensitivity compared to other BRAF-mutant melanoma cell lines.[27] SKMEL5 cell line was purchased from ATCC© and labeled with either a fluorescent histone H2B conjugated to the monomeric red fluorescent protein (H2BmRFP) and a cellular barcoding library (see "Cellular barcoding" below) or H2B conjugated to the green fluorescent protein (H2B-GFP). Single cell-derived subclones of SKMEL5 were selected and derived by limiting dilution, as described previously. Cells were cultured in a mixed media of DMEM and Ham F-12 media (DMEM:F12 1:1; catalog no. 11330-032), supplemented with 10% fetal bovine serum (FBS). Cells were incubated at 37°C, 5% CO_2, and passaged twice a week using TrpLE (Gibco). Cell lines and sublines were tested for mycoplasma contamination using the MycoAlert[TM] mycoplasma detection kit (Lonza), according to manufacturer's instructions, and confirmed to be mycoplasma-free. BRAF inhibitor PLX 4720 (analog to Vemurafenib), ferroptosis inducer RSL3 and ferroptosis inhibitor Fer-1 were obtained from MedChem Express (Monmouth Junction, NJ) and solubilized in dimethyl sulfoxide (DMSO) at a stock concentration of 10mM and stored at -20°C. Cell lines were originally stored at -80°C, then moved into liquid nitrogen.

Cellular barcoding

Setup: Cellular barcoding library was constructed by cloning a guide RNA (gRNA) library of barcodes into a CROP-seq-BFP-TSO vector as previously described. The vector was engineered such that barcodes can be isolated by isolation and amplification (barcode sampling) or mRNA capture in a single-cell RNA sequencing (scRNA-seq) experiment. gRNAs were built as a 20-nucleotide sequence of four nucleotides identical among all barcodes, followed by a 16 strong-weak (SW) paired nucleotides (i.e., XXXXSWSWSWSWSWSWSWSW). The SW pairing of the barcode sequence was designed to prevent polymerase chain reaction (PCR) amplification bias.

The maximum complexity of this library is 2^{16} (~65,000 unique barcodes). The barcode library
vector was used to produce lentiviral libraries using a lipofectamine transfection of HEK293T
cells. Media containing lentiviral particles were collected at 48- and 72-hours post-transfection,
pooled, filtered through a 0.45 um Nalgene syringe filter and concentrated using a 50 mL size-
exclusion column by centrifugation at 2200 RCF at 4°C for two hours. Concentrated virus was
stored in -80°C. SKMEL5 cells were seeded in a 6-well plate at ~$1x10^6$ cells per well in 2.5 mL
culture media. Cells were transduced with the barcoded CROP-seq-BFP-TSO-Barcode_sgRNA
lentivirus using 0.8 μg/mL in each well and a multiplicity of infection (MOI) of 0.05. Twenty-four
hours after incubation, transduction media (containing polybrene) was exchanged for fresh
culture media. Forty-eight hours after incubation, barcoded cells were isolated by fluorescence-
activated cell sorting (FACS) and subsequently cultured until confluence in a T-150 dish and then
cryopreserved. Cryopreserved cells were thawed in a T-25 dish and scaled up for ~2 weeks in two
separate sets. The first set of thawed cells were treated with 8uM PLX4720 (and an untreated
control) for eight days and subjected to barcode sampling (see "Barcode Sampling Analysis"
subsection below). The second set was plated in three T-75 flasks (parallel replicates) and
independently treated with 8μM PLX4720 (or untreated control) for eight days and subjected to
scRNA-seq by the 10X genomics Chromium platform (version 2 chemistry; see "RNA single-cell
transcriptome sequencing" section below for more details). In both cases, treated cells had media
and drug replaced every three days. Untreated cells were expanded completely over the course
of the time course (i.e., no cell splitting).
499
Barcode Sampling: After PLX4720 treatment for eight days (or no treatment expansion), cells in
the first set were pelleted for genomic DNA (gDNA) extraction using the DNeasy Blood and Tissue
Kit (Qiagen) per manufacturers' instructions. Barcode sequences were amplified for each
replicate by polymerase chain reaction (PCR; 98°C for 30 seconds, followed by 22 cycles of
denaturation - 98°C for 10 seconds, annealing - 63°C for 30 seconds, extension - 72°C for 10
seconds, and a final extension of 72°C for 5 minutes) using primers containing flanking regions
and Illumina adapter index sequences. 2ug gDNA was used in each PCR reaction, and a
combination of 5 distinct pooled forward primers were utilized to minimize sequencing error.
Reactions were purified using a 1.8x AMPure XP bead (Beckman Coulter) cleanup. Reaction
products were confirmed using agarose gel confirmation (band at ~215bp). The resulting libraries
were quantified using a Qubit fluorometer (ThermoFisher), Bioanalyzer 2100 (Agilent) for library
profile assessment, and qPCR (Kapa Biosciences Cat: KK4622) to validate ligated material,
according to the manufacturer's instructions. The libraries were sequenced using the NovaSeq
6000 with 150 bp paired end reads as sequencing spike-ins (targeting ~200k reads). RTA (version
2.4.11; Illumina) was used for base calling and MultiQC (version 1.7) for quality control.
515
Barcode Sampling Analysis: Barcodes were identified from amplified sequence reads by trimming
flanking adapter sequences. Barcodes abundances were totaled and normalized to library read
depth, resulting in reads per million (RPM). Barcodes less than 100 CPM were removed from the
analysis. Numbers of unique barcodes were calculated based on this threshold. Overlaps
between experimental replicates were calculated to determine the proportion of barcodes
shared across different runs. Total barcode abundance (including low abundance barcodes) was

calculated using the $\log_{10}$ of barcode RPM for each replicate. Relative barcode fraction was
calculated for each sample across three replicates. $\log_2$ fold change of the idling to untreated
mean barcode fractions was calculated for all barcodes above the CPM threshold.
525
526 **RNA single-cell transcriptome sequencing**
527
Data Collection: After PLX4720 treatment for eight days (or no treatment expansion), cells in the
second set were prepared targeting ~3000 cells per sample, washed, and resuspended in 0.04%
bovine serum albumin (BSA) in phosphate-buffered saline (PBS). Cell suspensions were subjected
to 10X Genomics single-cell gene expression protocol (version 2, 3' counting) in two separate
wells, according to manufacturer's guidelines. Single-cell mRNA expression libraries were
prepared according to manufacturer instructions. Due to the nature of gRNA barcoding library
construction, mRNAs resulting from gRNA barcodes were captured along with other mRNAs.
Libraries were cleaned using SPRI beads (Beckman Coulter) and quantified using a Bioanalyzer
2100 (Agilent). The libraries were sequenced using the NovaSeq 6000 with 150 bp paired-end
reads targeting 50M reads per sample for the mRNA library (including barcode library). RTA
(version 2.4.11; Illumina) was used for base calling and MultiQC (version 1.7) for quality control.
Gene counting, including alignment, filtering, barcode counting, and unique molecular identifier
(UMI) counting was performed on each library using the *count* function in the 10X Genomics
software *Cell Ranger* (version 3.0.2) with the GRCh38 (hg38) reference transcriptome.
542
Transcriptome Analysis: Cell Ranger output two single-cell gene expression matrices, for
untreated and idling cells. scUniFrac was performed to quantify the degree of overlap between
conditions. Finding minimal overlaps, and since cells were prepared and processed in parallel, no
computational batch correction was performed. Seurat was used to perform gene expression
analysis. The *SCTransform* function was used to regress out mitochondrial gene expression
(percent.mt), number of features (genes; nFeature_RNA), number of RNA molecules in the cell
(nCount_RNA), and cell cycle variables (S.Score and G2M.Score). Feature selection was
performed according to Seurat guidelines using a variance stabilizing transformation of the top
2000 most variable features. Data was normalized and scaled according to Seurat guidelines.
Data between conditions were combined and visualized using the Uniform Manifold
Approximation and Projection (UMAP) dimensionality reduction algorithm as implemented in
Seurat. In addition to UMAP, t-distributed Stochastic Neighbor Embedding (t-SNE) and Principal
Component Analysis (PCA) were also performed, using the Seurat implementation. Clustering
was performed in the joint UMAP space using the default Seurat implementation, a shared
nearest neighbor (SNN) modularity optimization-based method. Pairwise Euclidean distances
were calculated between cells in each condition with 10 principal components in the PCA space
and plotted as a cumulative density function (CDF). An Earth Mover's Distance (EMD) was
calculated between 15000 randomly sampled pairwise distances using the *wasserstein1d*
function in the *transport* R package. Differential expression analysis was performed between
conditions (untreated, idling), between combined clusters (large, small) within each condition,
and between clusters across conditions (e.g., untreated$_{large}$ vs. idling$_{large}$). Differential expression
was performed using the Seurat *FindMarkers* function and DEGs (adjusted-p < 0.05) were input
into a GO over-enrichment analysis. The GO analysis identified cell cycle as a major factor

separating idling clusters and was not present for untreated clusters. Therefore, using cell cycle
scores, a cell cycle phase (G1, G2M, S) was assigned to each cell, which was further simplified
into cycling (S, G2M) and non-cycling (G1), which we call cell cycle state. Cluster proportion was
calculated by cell cycle state to quantify the differences between clusters. The number of cells in
each cell cycle state were tallied and calculated as a percentage across each cluster.

Barcode Analysis: After calculation of scRNA-seq gene expression matrices, barcode abundances
were incorporated to the matrices. To do this, gRNA lineage barcodes had to be mapped to their
associated 10X cell barcodes. First, unmapped scRNA-seq BAM files were cleaned to only include
the mRNA transcript ID, scRNA-seq cell barcode, and scRNA-seq unique molecular identifier
(UMI). Mapped scRNA-seq BAM files (3' heavy) were cleaned to only include the mRNA transcript
ID and lineage barcode (from gRNA library). Unmapped and mapped subsets were merged on
the mRNA transcript ID to assign a lineage barcode to each cell barcode and UMI. The resulting
merged dataset was paired down to a single cell barcode – lineage barcode pair, which was
appended to each cell in the gene expression matrix as a metadata tag. Barcode abundances
were totaled across all cells in the experiment that captured a barcode, and strongly reflected
barcode sampling relative abundances and fold changes upon treatment (FIG. 1E - scRNA-seq
barcode fractions). Barcodes were overlaid on UMAP projections of scRNA-seq data, and further
categorized into the cycling and non-cycling transcriptomic states (see Transcriptome Analysis
subsection). Total number of cells from each barcode were tallied across each transcriptomic cell
cycle state, and a percentage (relative to each barcode) in each state was calculated. A
correlation analysis (Pearson) was performed on the percentage of cycling cells in idling vs. the
log$_2$ barcode fold change after treatment.

Functional interpretation analysis: The single-cell transcriptome count matrix (see "RNA single-
cell transcriptome sequencing:Data Collection" above) was scaled by multiplying counts by the
median RNA molecules across all cells and dividing that number by the number of RNA molecules
in each cell (as recommended). Gene signature files were obtained from the molecular signatures
database (MSigDB). *Hallmark* gene sets (50 in total) were downloaded from MSigDB (gsea-
msigdb.org/gsea/msigdb/genesets.jsp?collection=H). Both the scaled counts matrix and each of
the hallmark gene sets were input into VISION[28] to identify gene signature scores for each cell-
signature pair. Four hallmark gene sets (KRAS_SIGNALING_UP, KRAS_SIGNALING_DOWN,
UV_RESPONSE_UP, UV_RESPONSE_DOWN) were condensed into two (KRAS_SIGNALING,
UV_RESPONSE) by VISION to leave 48 total gene signatures. Scores were compiled into a
distribution and plotted by cluster (k=3) for each gene set.

Bulk RNA transcriptome sequencing

Data acquisition: Total RNA was isolated from untreated SKMEL5 single cell-derived subclones,
each in triplicate, using Trizol isolation method (Invitrogen), according to the manufacturer's
instructions. RNA samples were submitted to Vanderbilt VANTAGE Core services for quality
check, where mRNA enrichment and cDNA library preparation were done with Illumina Tru-Seq
stranded mRNA sample prep kit. Sequencing was done at Paired-End 75 bp on the Illumina HiSeq

3000. Reads were aligned to the GRCh38 human reference genome using 'HISAT2' and gene
counts were obtained using 'featureCounts'.

Data analysis: RNA-seq data was analyzed using the DESeq2[29] R package. Cells with less than 18
reads per condition were removed, according to DESeq2 vignettes. Counts were transformed
using the regularized logarithm (rlog) normalization algorithm. PCA was performed using the
prcomp function in R. The path between time series data points was visualized as a line between
subline-time point replicate means in the PCA space. EMDs were calculated between pairwise
distance distributions for each subline across the treatment time series and for each time point
across the sublines. Hierarchical clustering was performed using the *hclust* R function with a
Ward's minimum variance method. Differential expression analysis was performed using a model
design to quantify both changing variables and their interaction (~ subline + treatment time +
subline:treatment time). DEGs across sublines between untreated (pre-treatment, day 0) and
idling (day 8 post-treatment) were identified (adjusted-$p < 0.05$, log2 fold change > 2). DEGs were
input into a GO enrichment analysis, which identified GO terms associated with "Biological
Process" (BP), "Molecular Function" (MF), and "Cellular Component" (CC) GO types.

Ferroptosis gene signature analysis: A ferroptosis gene signature was obtained from the Kyoto
Encyclopedia of Genes and Genomes (KEGG). For each subline, genes were normalized to the 0-
day time point and a $\log_2$ fold change was calculated compared to the 0-day baseline. Groupings
of genes were annotated by separate signaling processes (Supplementary FIG. S5).

Bulk ATAC epigenome sequencing

Data acquisition: Data was collected using the omni-ATAC protocol for bulk ATAC sequencing
(ATAC-seq). After PLX4720 treatment for eight days (or no treatment expansion), cells in the
second set (in parallel to barcode sampling data collection) were pelleted at 50k cells and
resuspended in a cold ATAC-seq resuspension and lysis buffer containing NP40 (0.1%), Tween20
(0.1%), and Digitonin (0.01%) and incubated on ice. A resuspension buffer was added (0.1%
Tween20, no NP40 or Digitonin) to wash out the lysis reaction. Cells were pelleted and
resuspended in a transposition mix (5x Tris-DMF, PBS, 1% digitonin, 10% Tween20, nuclease-free
H_2O), including transposase Tn5, followed by a 30-minute incubation at 37°C, with shaking to
enhance tagmentation. After 30 minutes, the reaction was stopped by adding a DNA binding
buffer (Zymo) and purified using a DNA Clean and Concentrate kit (D4004, Zymo). The final
product was eluted in nuclease-free H_2O. PCR amplification was performed on the eluate with an
NEBNext 2X High Fidelity PCR Mix (NEB, M0541S) N7, and N5 index sequencing primers
(extension at 72°C for 5 minutes; denaturation at 90°C for 30 seconds; 12 cycles: denaturation at
98°C for 10 seconds, annealing at 62°C for 30 seconds, extension at 72°C for 30 seconds; final
extension at 72°C for 5 minutes). The PCR product was purified with the Zymo DNA Clean and
Concentrate kit and eluted in 22μL nuclease-free H_2O. ATAC-seq PCR libraries were visualized by
agarose gel electrophoresis for a quick check for the nucleosome ladder pattern (bands over ~150
bp). Libraries were also quantified using a Qubit fluorometer (ThermoFisher), Bioanalyzer 2100
(Agilent) for library profile assessment, and qPCR (Kapa Biosciences Cat: KK4622) to validate

ligated material, according to the manufacturer's instructions. The libraries were sequenced
using the NovaSeq 6000 with 150 bp paired-end reads as spike-ins on the sequencing chip
(untreated: ~160m, idling: ~130m reads). RTA (version 2.4.11; Illumina) was used for base calling
and MultiQC (version 1.7) for quality control.
656
Data Analysis: Reads were trimmed using *cutadapt* (paired-end) to remove primer sequences
and aligned to the hg38 reference genome using the *bwa mem* function in Burrows-Wheeler
Aligner (BWA, version 0.7.17). Aligned reads were sorted and duplicates were marked using
Picard (version 2.17.10). Untreated reads had more detected duplicates (~78% compared to
~32% in idling). Reads were deduplicated, but to address the discrepancy in deduplicated library
complexity, idling reads were subsampled (25% of original library, Supplementary FIG. S4A-B) to
achieve a similar complexity to the untreated. Reads were further cleaned according to sequence
quality guidelines. Insert sizes were plotted from the output of *InsertSizeMetrics* after
deduplication in Picard. Peaks of open chromatin were called using the *MACS2 callpeak* function
according to recommended guidelines for ATAC-seq data (BAM paired-end method, q-threshold:
0.05, no MACS2 model, shift: -100, extension size: 200). Peaks were subjected to a further round
of quality control and cleaning using *ChIPQC* (peak mapping, peak duplication, blacklist peak
detection), and blacklisted peaks were removed. Peaks were converted to consensus counts
using the *runConsensusCounts* function in *soGGi*. Intersections of and unique cleaned peaks were
determined and visualized as a Venn diagram using the *vennDiagram* function in the *limma*
package. Unique and intersection peaks were annotated with the nearest neighbor genes using
the *annotatePeak* function and hg38 transcriptome in the *ChIPseeker* package. These peaks were
also re-aligned to the transcription start site (TSS) for each gene. Average profiles of read subsets
across all genes (nucleosome-free, mono-nucleosome, and di-nucleosome; normalized to the
TSS). Peaks were classified based on closeness to the TSS and assigned to predicted feature
(promoter, UTR, exon, intron, downstream, distal intergenic). Genes associated with unique and
intersections of peaks were input into a GO enrichment analysis for BP, MF, and CC GO types
(same as "Bulk Transcriptome Analysis" above). Transcription factor (TF) footprinting in the
region around TSSs was performed on untreated and idling unique peaks for key TFs.
681
682   **Gene ontology analysis**
683
Setup: Genes associated with unique ATAC-seq peaks (see "Bulk ATAC Epigenome Sequencing:
Data Analysis" above) were identified for each condition (i.e., untreated or idling). Additionally,
DEGs from the bulk RNA-seq statistical analysis were determined (8-day vs. 0-day time points
across all clonal sublines; see "Bulk RNA Transcriptome Sequencing:Data Analysis" above). The
two gene lists were independently subjected to a GO enrichment analysis using '*clusterProfiler*'.
Genes were compared to BP, MF, CC GO types. GO terms significantly enriched in the unique
ATAC-seq peaks ($p < 0.05$) and in DEGs ($p < 0.05$) were identified and stored independently as
separate GO term lists for untreated and idling datasets.
692
Correlation Analysis: The $-\log_{10}$(p-value) was calculated for terms shared between the lists
associated with the separate modalities, ranking terms based on statistical significance.

Spearman correlation was calculated between the significant GO terms using the *ggpubr* package
(version 0.4.0) for each GO type.

Calcium flux assays

Data acquisition: SKMEL5 barcoded cells were plated onto a 384 well, tissue culture treated plate
24 hours before imaging at a density of 10,000 cells/well. For the treated condition, cells were
treated with 8 μM PLX4720, a vemurafenib derivative, for 8 days with fresh media/drug swapped
out every 3 days. Untreated control cells were taken from the same cell line culture which was
maintained separately while the treatment condition was exposed to PLX4720, being split down
when necessary to maintain healthy growth conditions. On the day of experimentation, the 384
well plate with all treated and untreated cells were incubated with 4 μM Fluo-8-AM in fresh
culture media (10% FBS) for 1 hour at room temperature, as recommended by the manufacturer
(AAT Bioquest). Dye containing media was removed and Hanks Buffered Saline Solution (HBSS,
10 mM HEPES, no Ca^{2+}) was used to wash the wells of excess dye, followed by removal and
addition of 20 μL of fresh HBSS (no Ca^{2+}) for use as the assay buffer. The plate was loaded into a
kinetic imaging instrument (Panoptic, by Waveguide Biosciences), which records the fluorescent
intensity emitted by each well of the 384 well plate. A three-addition protocol was used to add
the various drugs and assay conditions to the plate during the SOCE assay. Drug Addition plates
were loaded with assay buffer (HBSS, with or without Ca^{2+}) and thoroughly mixed immediately
before imaging. Add conditions were split into three parts: 1) Cyclopiazonic Acid (CPA; final
concentration of 50 μM) to inhibit activity of Sarcoendoplasmic Reticulum Calcium ATPase
(SERCA), leading ER Ca^{2+} release; 2) Addition of Ca^{2+} to the assay condition to activate SOCE
activity; and 3) Addition of the Ca^{2+} ionophore, Ionomycin (final concentration of 5 μM), was used
to generate maximal signal intensity to control for variations in cell count in individual wells (this
was particularly important since drug treated idling cells experienced increased washout due to
the stressful nature of sustained BRAF inhibition). Fluo-8-AM was excited with [480 nm] and
imaged at [538 nm], with a frequency of 1 Hz. The CPA treatment condition was imaged for 260
seconds before addition of Ca^{2+}, followed by 270 seconds of imaging before addition of 5 μM
Ionomycin. Treatment conditions were replicated in sets of 8 and average values traced with 95%
confidence intervals.

Data analysis: Fluorescence data were normalized by dividing data from each well by the first
fluorescence value at the start of the experiment. The ionomycin peak was used as a control for
the number of cells in each treatment group, as the idling population tends to have less cell than
the untreated population. Therefore, the mean ionomycin treatment peak was calculated for
each treatment group (i.e., idling and untreated), and data from each treatment group were
divided by the corresponding mean ionomycin peak value to account for the change in the
number of cells in the assay. For comparison of calcium flux between the treatment groups, the
lowest values were the zero starting point (on the y-axis) in the plot for each of the curves. A
mean and 95% confidence interval were calculated for each time point and plotted using *ggplot*.

Ferroptosis-induction experiments

Plates of H2B-GFP-labeled SKMEL5 cells were treated with either vehicle (DMSO) or BRAFi ($8\mu M$ PLX4720) for eight days incubated at 37°C and 5% CO_2, changing media (with vehicle/drug) every three days. BRAFi-treated cells were plated at ~2500 cells per well in a black, clear-bottom 96-well plate (Falcon). After cell seeding, RSL3, with or without Fer-1, was added the following morning, with media changes every three days (six replicates per condition). Plates were imaged using automated fluorescence microscopy (Cellavista Instrument, Synentec). Twenty-five non-overlapping fluorescent images (20X objective, 5x5 montage) were taken twice daily for a total of 150 hours or until confluency. Cellavista image segmentation software (Synentec) was utilized to calculate nuclear count (i.e., cell count) per well at each time point (Source = FITC, Dichro = FITC, Filter = FITC, Emission Time = $800\mu s$, Gain = 20x, Quality = High, Binning = 2x2). Cell nuclei counts across wells were normalized to time of drug treatment and used to calculate a DIP rate (stable linear growth rate). A dose-response curve was calculated across replicates using the *drm* R package with a 4-parameter log-logistic function, with DIP rate as the drug effect. Replicates were used to calculate means and 95% confidence intervals for the dose-response curves. IC_{50} values were calculated for each condition and plotted as vertical dashed lines. Data was visualized using the ggplot2R package (version 3.2.0).

Model and Experimental Analysis Code Availability. The codes used to generate model simulations and analyze experimental data are publicly available via GitHub, or from the corresponding author upon request.

Data Availability. The sequencing datasets generated in this study can be found in the gene expression omnibus and sequence read archive. Additional experimental data will be available from the corresponding author upon request.

ACKNOWLEDGEMENTS

We thank Emily Hodges and Kelly Barnett for ATAC-seq reagents and experimental advice. We also thank Jing Hao for reagent acquisition and Tony Capra, Christian Meyer, Sarah Maddox Groves, Carlos Lopez, Alissa Weaver, John McLean, Bruce Damon, and Ken Lau for useful discussions.

Sequencing studies were performed by the Vanderbilt Technologies for Advanced Genomics (VANTAGE) core facility at Vanderbilt University Medical Center with help from Angela Jones, Karen Beeri, Jamie Roberson, Latha Raju, and Matthew Scholz. Cellular barcode library construction and transduction were performed in the Brock laboratory at The University of Texas at Austin. Data processing and model simulations were performed using resources at the Advanced Computing Center for Research and Education (ACCRE) at Vanderbilt University.

This work was supported by the following funding sources: C.E.H., National Institutes of Health (NIH) Ruth L. Kirschstein National Research Service Award (NRSA, F31-CA221147) and Chemical-Biology Interface Training Grant (T32-GM0650); L.A.H., National Cancer Institute (NCI) Transition

781 Career Development Award to Promote Diversity (K22-CA237857-01A1); D.R.T., NIH Research
782 Specialist Award (1R50CA243783); V.Q., NCI Emerging Questions in Cancer Systems Biology (U01
783 CA215845), NCI Research Centers for Cancer Systems Biology (U54 CA217450), NCI Predictive
784 Multiscale Models for Biomedical, Biological, Behavioral, Environmental and Clinical Research
785 (U01 CA174706), and Quantitative Systems Biology Center at Vanderbilt University.

786
787 COMPETING INTERESTS
788
789 The authors declare no competing interests.
790
791 AUTHOR CONTRIBUTIONS
792
Conceptualization, C.E.H., L.A.H., and V.Q.; Investigation and experiments, A.A., B.B.P., C.E.H.,
D.R.T., K.E.J., P.E.S.; Bioinformatic Analysis, B.B, C.E.H.; Modeling, C.E.H. and L.A.H.; Writing,
B.B., C.E.H. and V.Q.; Review and Editing, A.B., B.B., C.E.H., L.A.H., and V.Q.

REFERENCES

1   Liu J, Dang H, Wang XW. The significance of intertumor and intratumor heterogeneity in
liver cancer. *Exp Mol Med* 2018; **50**: e416.

2   Inde Z, Dixon SJ. The impact of non-genetic heterogeneity on cancer cell death. *Crit Rev
Biochem Mol Biol* 2018; **53**: 99–114.

3   Hayford CE, Tyson DR, Robbins CJ, Frick PL, Quaranta V, Harris LA. An in vitro model of
tumor heterogeneity resolves genetic, epigenetic, and stochastic sources of cell state
variability. *PLOS Biol* 2021; **19**: e3000797.

4   Huang S. Genetic and non-genetic instability in tumor progression: link between the
fitness landscape and the epigenetic landscape of cancer cells. *Cancer Metastasis Rev*
2013; **32**: 423–448.

5   Brock A, Chang H, Huang S. Non-genetic heterogeneity--a mutation-independent driving
force for the somatic evolution of tumours. *Nat Rev Genet* 2009; **10**: 336–42.

6   Pisco AO, Huang S. Non-genetic cancer cell plasticity and therapy-induced stemness in
tumour relapse: 'What does not kill me strengthens me'. *Br J Cancer* 2015; **112**: 1725–
1732.

7   Paudel BB, Harris LA, Hardeman KN, Abugable AA, Hayford CE, Tyson DR *et al.* A
nonquiescent 'idling' population state in drug-treated, BRAF-mutated melanoma.
Biophys J 2018; **114**: 1499–1511.

8   Aranda-Anzaldo A, Dent MAR. Landscaping the epigenetic landscape of cancer. *Prog
Biophys Mol Biol* 2018; **140**: 155–174.

9   Sharma S V, Lee DY, Li B, Quinlan MP, Takahashi F, Maheswaran S *et al.* A chromatin-
mediated reversible drug-tolerant state in cancer cell subpopulations. *Cell* 2010; **141**: 69–
80.

10   Hata AN, Niederst MJ, Archibald HL, Gomez-Caraballo M, Siddiqui FM, Mulvey HE *et al.*
Tumor cells can follow distinct evolutionary paths to become resistant to epidermal
growth factor receptor inhibition. *Nat Med* 2016; **22**: 262–269.

11   Ramirez M, Rajaram S, Steininger RJ, Osipchuk D, Roth MA, Morinishi LS *et al.* Diverse
drug-resistance mechanisms can emerge from drug-tolerant cancer persister cells. *Nat
Commun* 2016; **7**: 10690.

12   Oren Y, Tsabar M, Cuoco MS, Amir-Zilberstein L, Cabanos HF, Hütter J-C *et al.* Cycling
cancer persister cells arise from lineages with distinct programs. *Nat 2021 5967873* 2021;
596: 576–582.

13   Jia D, Paudel BB, Hayford CE, Hardeman KN, Levine H, Onuchic JN *et al.* Drug-tolerant
idling melanoma cells exhibit theory-predicted metabolic low-low phenotype. *Front
Oncol* 2020; **10**: 1426.

14   Workman P, Travers J. Cancer: drug-tolerant insurgents. *Nature* 2010; **464**: 844–5.

15   Gutierrez C, Al'Khafaji AM, Brenner E, Johnson KE, Gohil SH, Lin Z *et al.* Multifunctional
barcoding with ClonMapper enables high-resolution study of clonal dynamics during
tumor evolution and treatment. *Nat Cancer 2021 27* 2021; **2**: 758–772.

16   Becht E, McInnes L, Healy J, Dutertre CA, Kwok IWH, Ng LG *et al.* Dimensionality
reduction for visualizing single-cell data using UMAP. *Nat Biotechnol* 2019; **37**: 38–47.

17   Xie Y, Hou W, Song X, Yu Y, Huang J, Sun X *et al.* Ferroptosis: process and function. *Cell
Death Differ* 2016; **23**: 369–379.

18 Boumahdi S, de Sauvage FJ. The great escape: tumour cell plasticity in resistance to
targeted therapy. *Nat Rev Drug Discov 2019 191* 2019; **19**: 39–56.
19 Gupta PB, Pastushenko I, Skibinski A, Blanpain C, Kuperwasser C. Phenotypic plasticity:
driver of cancer initiation, progression, and therapy resistance. *Cell Stem Cell* 2019; **24**:
65–78.
20 Rees M, Jessica C, Metcalf E, Childs DZ. Bet-hedging as an evolutionary game: the trade-
off between egg size and number. *Proc Biol Sci* 2010; **277**: 1149–51.
21 Jolly MK, Kulkarni P, Weninger K, Orban J, Levine H. Phenotypic plasticity, bet-hedging,
and androgen independence in prostate cancer: Role of non-genetic heterogeneity.
Front. Oncol. 2018; **8**: 50.
22 Shaffer SM, Dunagin MC, Torborg SR, Torre EA, Emert B, Krepler C *et al.* Rare cell
variability and drug-induced reprogramming as a mode of cancer drug resistance. *Nature*
2017; **546**: 431–435.
23 Emert BL, Cote CJ, Torre EA, Dardani IP, Jiang CL, Jain N *et al.* Variability within rare cell
states enables multiple paths toward drug resistance. *Nat Biotechnol 2021 397* 2021; **39**:
865–876.
24 Gray JW, Mills GB. Large-Scale Drug Screens Support Precision Medicine. *Cancer Discov*
2015; **5**: 1130–2.
25 Huang S, Kauffman S. How to escape the cancer attractor: Rationale and limitations of
multi-target drugs. *Semin Cancer Biol* 2013; **23**: 270–278.
26 Lee MJ, Ye AS, Gardino AK, Heijink AM, Sorger PK, MacBeath G *et al.* Sequential
application of anticancer drugs enhances cell death by rewiring apoptotic signaling
networks. *Cell* 2012; **149**: 780–94.
27 Hardeman KN, Peng C, Paudel BB, Meyer CT, Luong T, Tyson DR *et al.* Dependence on
glycolysis sensitizes BRAF-mutated melanomas for increased response to targeted BRAF
inhibition. *Sci Rep* 2017; **7**: 42604.
28 DeTomaso D, Jones MG, Subramaniam M, Ashuach T, Ye CJ, Yosef N. Functional
interpretation of single cell similarity maps. *Nat Commun* 2019; **10**: 4376.
29 Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* 2014; **15**: 1–21.
870

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6