TITLE

Heterogeneous, population-level drug-tolerant persisters exhibit ion-channel remodeling and ferroptosis susceptibility

AUTHORS

Corey E. Hayford[1†], Blake Baleami[2†], Philip E. Stauffer[1†‡], B. Bishal Paudel[1#], Aziz Al'Khafaji[3,4§], Amy Brock[3,4], Vito Quaranta[5,6¶], Darren R. Tyson[6‖*], and Leonard A. Harris[7,8,9*]

[1]Chemical and Physical Biology Graduate Program, Vanderbilt University School of Medicine, Nashville, TN, USA
[2]Human Genetics Graduate Program, Vanderbilt University School of Medicine, Nashville, TN, USA
[3]Institute for Cellular and Molecular Biology, University of Texas at Austin, TX, USA
[4]Department of Biomedical Engineering, University of Texas at Austin, TX, USA
[5]Department of Biochemistry, Vanderbilt University School of Medicine, Nashville, TN, USA
[6]Department of Pharmacology, Vanderbilt University School of Medicine, Nashville, TN, USA
[7]Department of Biomedical Engineering, University of Arkansas, Fayetteville, AR, USA
[8]Interdisciplinary Graduate Program in Cell and Molecular Biology, University of Arkansas, Fayetteville, AR, USA
[9]Cancer Biology Program, Winthrop P. Rockefeller Cancer Institute, University of Arkansas for Medical Sciences, Little Rock, AR, USA

†These authors contributed equally to this work.
*Corresponding author. Email: darren.tyson@duke.edu (D.R.T.); harrisl@uark.edu (L.A.H.)

‡Current address: Center of Experimental Rheumatology, Department of Rheumatology, University Hospital of Zürich, Zürich, Switzerland
#Current address: UVA Comprehensive Cancer Center, University of Virginia School of Medicine, Charlottesville, VA, USA
§Current address: Broad Institute of MIT and Harvard, Cambridge, MA, USA
¶Current address: Department of Pharmaceutical Sciences, Bouvé College of Health Sciences, Northeastern University, Boston, MA, USA
‖Current address: Department of Pharmacology and Cancer Biology, Duke University School of Medicine, Durham, NC, USA

ABSTRACT

Drug-tolerant persisters (DTPs) represent a major obstacle to durable responses in targeted cancer therapy. DTPs are commonly described as distinct single-cell states that survive drug treatment via reversible, non-genetic mechanisms and drive tumor recurrence. Recent work demonstrates that multiple DTPs can coexist, reflecting diversity in lineage, signaling programs, or stress responses. However, each DTP is still generally viewed as a uniform cellular phenotype. Building on our prior work describing a

population-level DTP termed "idling" [Paudel et al., *Biophys. J.* (2018) 114, 1499–1511], here we present evidence supporting a fundamentally different view: that DTPs are not single-cell states, but rather heterogeneous populations composed of multiple sub-states with distinct division and death rates that balance to produce near-zero net population growth. Using single-cell transcriptomics and lineage barcoding, we identify multiple phenotypic states within idling DTP populations, with reduced heterogeneity compared to untreated populations, and find that idling DTP cells emerge from nearly all lineages. Transcriptomic and functional analyses further reveal altered ion-channel activity in idling DTPs, which we confirm experimentally. Moreover, drug-response assays reveal increased susceptibility of idling DTPs to ferroptosis, a non-apoptotic form of regulated cell death, indicating the emergence of vulnerabilities associated with drug tolerance. Altogether, our results support a population-level view of tumor drug tolerance in which DTPs comprise stable collections of phenotypic states, shaped by treatment-defined phenotypic landscapes, which are potentially vulnerable to subsequent interventions. This perspective implies that eradicating DTPs will require a fundamental shift away from cell-type-centric strategies toward sequential treatments that progressively reduce phenotypic heterogeneity by modulating the molecular and cellular processes that establish the DTP landscape, an approach previously termed "targeted landscaping."

INTRODUCTION

Cancer is a complex, dynamic disease characterized by intratumoral heterogeneity, which has been implicated in treatment evasion and acquired drug resistance (1–3). Tumor heterogeneity is traditionally viewed in terms of clonal cell populations with distinct genetic profiles that pre-exist or emerge during the course of treatment. However, non-genetic sources of tumor heterogeneity are receiving increased attention (4–10). Multiple investigators have reported cancer-cell subpopulations, broadly termed drug-tolerant persisters (DTPs), that withstand drug treatments via non-genetic mechanisms (11–16). DTPs have been variously described as quiescent or slow cycling and as reservoirs from which genetic resistance mutations may emerge. As such, eliminating DTPs before acquired resistance could delay or even prevent tumor recurrence, significantly improving anticancer treatment outcomes. Doing so, however, will require identifying vulnerabilities within the molecular and cellular processes underlying drug tolerance (6, 17–19), which remain poorly understood.

Recently, we described a heterogeneous DTP state, termed "idling," in BRAF-mutant melanoma cell populations under prolonged BRAF inhibition (17, 20). The idling state is not quiescent but rather characterized by balanced, non-zero rates of division and death, such that the population size remains nearly constant over time. It is also reversible, with the population returning to baseline proliferation after drug removal and reentering the DTP state upon drug reapplication. Mathematical modeling of several drug-tolerant BRAF-mutant melanoma cell populations suggested the idling DTP state is composed of multiple phenotypic sub-states, each with individual rates of division and death, across which cells can transition (17). This characterization of a DTP as a heterogeneous

population of single-cell states that contribute collectively to drug tolerance distinguishes idling DTPs from the current consensus view of DTPs (21). Specifically, while prior studies have reported phenotypic heterogeneity within drug-tolerant populations, it has been from the perspective of multiple co-existing DTPs, each considered to be a uniform, single-cell state that either pre-exists or emerges during treatment (11–16). In contrast, while our framework can support the co-existence of multiple DTPs within a population, we add an additional layer of complexity: phenotypic heterogeneity *within* the DTP population itself. This distinction is crucial because cell-type-centric interventions targeting DTPs are likely to fail if DTPs are heterogeneous collections of phenotypic states across which cells can transition during drug treatment. Eliminating any one phenotypic state, or presumed "persister" subtype, will fail to eradicate the drug-tolerant population because cells can reorganize to occupy remaining or newly emergent viable phenotypes. This may explain why effective treatments targeting DTPs have yet to be developed (21), despite numerous ongoing efforts (22–28).

In this work, we use single-cell and bulk RNA sequencing of BRAF-mutant melanoma cell populations to demonstrate that idling DTPs, while less heterogeneous than untreated cell populations, are still composed of multiple phenotypic states. We also use single-cell DNA barcoding to show that idling DTPs emerge from nearly all untreated cell lineages, rather than by competitive selection and expansion of pre-existing drug-tolerant clones. Gene-ontology (GO) analyses and calcium-flux assays indicate that ion-channel activity, including store-operated calcium entry (SOCE), is significantly altered in idling cells. These observations suggest idling DTPs can be induced into ferroptotic cell death (11, 29, 30), which we confirm via drug-response assays demonstrating increased sensitivity to inhibition of glutathione peroxidase 4 (GPX4), a key regulator of ferroptosis. Taken together, we provide experimental support for a population-level view of drug tolerance in which tumor cell populations remodel essential intracellular programs and retain a degree of phenotypic heterogeneity that confounds cell-type-focused targeting. However, vulnerabilities also emerge within DTP populations that may be actionable via secondary or sequential interventions. In what follows, we present the results of our experimental studies probing the transcriptomic, clonal lineage, molecular, and drug-response characteristics of idling DTPs. We conclude with a discussion of the broader implications of the view that DTPs are heterogeneous collections of interchanging phenotypes for future efforts aimed at targeting the biological processes underlying the phenotypic landscape in which these cells reside.

RESULTS

Prolonged BRAF inhibition reduces transcriptomic heterogeneity in BRAF-mutant melanoma cell populations

We previously showed that BRAF-mutant melanoma cell populations exhibit complex drug-response dynamics to targeted BRAF inhibition (BRAFi) (17). Across cell lines and single cell-derived subclones, the initial phase of drug response can vary, with

proliferation rates ranging from negative to positive. In almost all cases, however, this is followed by entry into a DTP state of near-zero net growth, termed "idling," in which cell division and death remain balanced. To assess the level of phenotypic heterogeneity in the idling DTP state relative to drug-naïve cells, we performed single-cell RNA sequencing on untreated and idling BRAF-mutant melanoma cell populations (Materials and Methods). When the high-dimensional transcriptomic data are visualized using a two-dimensional Uniform Manifold Approximation and Projection (UMAP) (31), we see that the untreated and idling DTP cells reside in distinct regions of the projected space, with minimal overlap (FIG. 1A). We also identify at least two distinct clusters in each case, which we refer to as "large" and "small" untreated and idling DTP clusters (FIGS. 1A and S1A). Importantly, the large and small untreated clusters are much farther apart in the UMAP space than the two idling clusters, indicating the untreated clusters are more dissimilar from each other than are the idling clusters (31). Taken together, these results suggest that phenotypic heterogeneity is reduced in idling DTPs relative to untreated melanoma cell populations. These observations are consistent with our prior work (17), in which we proposed that a "drug-naïve epigenetic landscape" composed of multiple phenotypic states is modified upon exposure to BRAFi, and that the resulting "drug-modified epigenetic landscape" also comprises multiple phenotypic states but is less heterogeneous overall.

As additional evidence for the reduced phenotypic heterogeneity of the idling DTP state, we performed bulk RNA sequencing on three single-cell-derived subclones of the same BRAF-mutant melanoma cell line (Materials and Methods). These subclones were previously shown to have disparate short-term BRAFi responses, but all transition into an idling DTP state within 3−8 days of treatment (17). Using a principal component analysis (PCA), we see that prior to BRAFi treatment (day 0) there is a significant difference along the first PCA axis (PC1) between one subclone and the other two (FIG. 1B). After BRAFi treatment, all subclones move along the second PCA axis (PC2) and then begin to converge along PC1 by the end of the experiment (day 8), as the populations reach the idling DTP state. Since these subclones are single-cell-derived, it is exceedingly unlikely that these changes are genetic in origin. Therefore, these results provide further evidence that the idling DTP state is less heterogeneous than the untreated population, but still retains a degree of phenotypic heterogeneity.

Untreated and idling DTP transcriptomic states are associated with different metabolic processes and cell cycle stages

To determine the biological factors that distinguish the single-cell transcriptomic clusters identified in the untreated and idling DTP populations (FIGS. 1A and S1A), we performed a functional interpretation analysis using hallmark gene sets from the Molecular Signatures Database (MSigDB) (32–34) (Materials and Methods). For the untreated population, we considered the large and small clusters separately, across which the population is split approximately 80/20. We see that the large untreated cluster is enriched in a glycolysis gene signature compared to the small untreated cluster (FIG. 1C), which

is consistent with our previous report of metabolic heterogeneity within untreated BRAF-
mutant melanoma cell populations (20). Furthermore, the large untreated cluster has a
higher DNA repair gene signature score, while the small cluster is enriched in a p53
pathway gene signature (FIG. 1C). Comparing the idling population (considered as a
single cluster in this analysis) to both untreated clusters, we see enrichment in gene
signatures for DNA repair, fatty-acid metabolism, reactive oxygen species (ROS)
pathways, and oxidative phosphorylation (FIG. 1C). The latter is consistent with multiple
reports in the literature that BRAF-mutant melanoma cells rely primarily on glycolysis for
growth and energy production (35, 36) and treatment with BRAFi interferes with glycolytic
processes, presumably favoring a switch to oxidative phosphorylation (27, 37).
189
We also performed differential expression analyses between the two untreated single-cell
transcriptomic clusters (FIG. S1B) and between the two idling DTP clusters (the large and
small idling clusters are considered separately in this analysis; FIG. S1C). After
regressing out cell-cycle-associated genes (along with various other genes; see Materials
and Methods), a GO over-enrichment analysis (38) returned numerous cell-cycle-related
GO terms (e.g., "nuclear division", "DNA replication"; FIG. S1C). Thus, even with cell cycle
genes removed from the analysis, cell cycle-related processes emerge as major factors
distinguishing cells in the large and small idling DTP clusters. To further explore this
result, we used a previously established cell cycle gene signature (39) to classify
individual cells (both untreated and idling) by cell cycle stage (G1, G2/M, S). For visual
clarity, we condensed the cell cycle stages into "slow dividing" (G1) and "fast dividing" (S-
G2-M) categories and colored all cells in the single-cell transcriptomics UMAP space by
category (FIG. 1D). Within the large and small untreated clusters, we see both slow- and
fast-dividing cells, although they are relatively well separated into different regions within
the clusters. This may indicate the presence of phenotypic sub-states within the two
untreated clusters, which would be consistent with our recent report on transcriptomic
heterogeneity within untreated EGFR-mutant non-small cell lung cancer cell populations
(6). Conversely, the large idling DTP cluster is highly populated by slow-dividing cells,
while the small cluster is composed almost entirely of fast-dividing cells (FIG. 1D,E).
Furthermore, the proportion of fast-dividing cells is much lower in the idling DTP
population (~15%) than in the untreated population (~67%), which is consistent with
BRAFi prolonging time spent in G1 (40–42) and the reduced (near-zero) proliferation rate
of idling DTPs. Altogether, these results and those from the functional interpretation
analysis above (using MSigDB), further support the view that the idling DTP state
comprises a heterogeneous collection of phenotypic states.
215

Idling DTP cell populations comprise cells from most untreated lineages

217
218   To determine whether cells in the idling DTP population are clonally selected from a pre-
219   existing subpopulation or emerge from all untreated lineages after drug treatment, we
220   barcoded the same BRAF-mutant melanoma cell line as above with a guide RNA (gRNA)
221   barcoding library (43, 44). Our barcoding system establishes a direct connection between
222   the abundance of a clonal lineage and the positions of cells within the single-cell

transcriptomics UMAP space (Materials and Methods). Upon treatment with BRAFi for eight days, we see only modest changes in the relative abundances of individual barcodes between the untreated and idling DTP populations (FIGS. 2A and S2A). We also see that the barcode library complexity is reduced by less than 10% (FIG. S2B), barcodes are largely shared across replicates for both the untreated and idling DTP populations (FIG. S2C), and fold-changes in barcode abundances are approximately normally distributed with a mean of zero (FIG. 2B). Notably, nearly all the lineages that do not survive BRAFi treatment come from clones with exceedingly small abundances (data not shown), suggesting that the loss of those lineages is due to random chance (6, 45–47).

Additionally, we mapped barcoded cells for the 24 most abundant barcodes onto the single-cell transcriptomics UMAP space (FIGS. 2C and S2D). In all cases, we see barcoded cells residing in all untreated and idling DTP clusters. Moreover, the changes in the relative abundances of barcodes (FIGS. 2A and S2A) correspond to the proportion of cells in a lineage that reside in the fast-dividing (small) idling DTP cluster, which we assume is populated randomly as the cell population re-equilibrates following drug treatment (17). Lineages in which only a few cells (<5%) reside in the fast-dividing cluster (e.g., barcodes 5 and 9; see FIG. 2C,D) show a reduction in their relative proportions after BRAFi treatment (FIG. 2A, *stars*). Conversely, lineages in which many cells (>20%) reside in the fast-dividing cluster (e.g., barcodes 2 and 13; see FIG. 2C,D) significantly increase their relative proportions following treatment (FIG. 2A, *circles*). Indeed, a strong positive correlation exists (Pearson, R=0.55) between the proportion of cells in a lineage residing in the fast-dividing idling DTP state and the relative fold change of barcode abundance after treatment (FIG. 2E). In all, these results indicate that most clonal lineages survive BRAFi treatment and persist in the idling DTP state in similar proportions to the untreated population. Thus, rather than clonal selection, the experimental evidence points to all BRAF-mutant melanoma cells having the capacity to enter the idling DTP state, which is consistent with numerous recent reports on the behaviors and characteristics of DTPs (11–16).

Multi-omics and calcium-flux analyses indicate changes in ion channel homeostasis and activity in drug-tolerant cells

We further analyzed the bulk RNA sequencing data from above for the three single-cell-derived BRAF-mutant melanoma subclones (FIG. 1B) by performing a differential expression analysis between the untreated (day 0) and idling DTP (day 8) populations. Inputting differentially expressed genes (DEGs) for all three subclones into a GO over-enrichment analysis returned numerous terms associated with ion transport and homeostasis (FIG. 3A). This led us to look at expression levels for numerous genes related to calcium handling (Materials and Methods). We see that many are upregulated or downregulated across the subclones after 3 and 8 days of BRAFi (FIG. 3B). These include genes encoding for sarcoendoplasmic reticulum Ca^{2+}-ATPases (SERCAs; e.g., ATP2A1), inositol 1,4,5-trisphosphate receptors (ITPRs), stromal interaction molecule 1

(STIM1), and solute carrier (SLC) genes that encode for various membrane transport
proteins. Additionally, we probed the epigenomic status of the idling DTP state by
performing bulk ATAC (assay for transposase-accessible chromatin) sequencing (48) on
samples of the same BRAF-mutant melanoma cell line, before and after 8 days of BRAFi
treatment (FIG. S3A,B; see Materials and Methods). Regions of open chromatin on the
DNA, referred to as "peaks," were identified for both conditions (FIG. S3C). The
distribution of binding loci (FIG. S3D) and corresponding peak feature set (FIG. S3E)
show that idling peaks have fewer proximal features (e.g., promoter regions) and more
distal elements (e.g., distal intergenic and intronic regions) compared to untreated cells.
Distal elements have been known to be involved in short-term epigenetic regulation (49).
Peaks unique to idling DTPs were then assigned to corresponding genes and these
genes input into a GO over-enrichment analysis. As with the bulk RNA-sequencing data,
numerous GO terms were returned associated with ion transport and activity (FIG. 3C).
Indeed, GO terms from bulk RNA and ATAC sequencing data strongly correlated with
each other, with the most significant terms being associated with ion-channel activity
(FIGS. 3D and S3F,G). Thus, two independent data modalities (transcriptomics and
epigenomics) provide strong evidence that alterations in ion-channel homeostasis and
activity are characteristic of the idling DTP state.
285
To assess the functional consequences of these findings, we quantified calcium-channel
activity in untreated and idling DTP cells using a calcium-flux assay that measures the
amount of endoplasmic reticulum (ER)-resident calcium and the propensity for store-
operated calcium entry (SOCE). In this assay, free ER calcium is measured first by using
cyclopiazonic acid (CPA) in a calcium-free buffer to inhibit the activity of SERCAs, leading
to the release of free calcium from the ER to the cytoplasm, where it is detected by a
calcium-binding dye (FIG. 3E, *first peak*). Under normal conditions, cells undergo SOCE
to replenish ER calcium upon depletion, i.e., ER calcium release stimulates the opening
of plasma membrane-resident calcium channels (e.g., ORAI by STIM1), allowing
extracellular calcium to flow into the cytoplasm before being pumped into the ER by
SERCAs. Thus, to assess SOCE activity, calcium is added to the assay buffer after CPA
addition and the subsequent increase in cytoplasmic calcium is quantified (FIG. 3E,
second peak). Our results show that while untreated and idling DTP cells have essentially
equal ER calcium content, SOCE activity in idling DTP cells is significantly reduced
compared to untreated cells (FIG. 3E). These results verify that ion-channel activity is
significantly altered in the idling DTP state.
302
303 **Idling DTPs exhibit increased susceptibility to ferroptotic cell death**
304
Previous reports indicate a complex interplay between ion channels, ER stress, and
ferroptosis, a form of regulated cell death characterized by the iron-dependent
accumulation of lipid peroxides, leading to oxidative damage to cell membranes (50–53).
ER stress occurs when there is an imbalance between the protein-folding load in the ER
and the ER's folding capacity, which can be caused by various factors, including
disturbances in calcium homeostasis (54). ER stress may, in turn, lead to increased

cellular iron levels, which can promote the generation of ROS and lipid peroxidation,
contributing to ferroptosis (29, 30). In our bulk RNA-sequencing data, we see altered
expression of several genes related to ferroptosis in idling DTP cells (FIGS. 4A and S4).
Notably, glutathione-metabolism gene expression is increased and genes for
polyunsaturated fatty-acid (PUFA) enzymes are decreased in idling DTP cells.
Interestingly, many DEGs feed into the Fenton reaction, which produces the free-radical
precursors to ROS, the step directly before commitment to ferroptosis (FIG. S4).

To directly test susceptibility to ferroptosis of BRAF-mutant melanoma cell populations,
we subjected both untreated and idling DTP cells to treatment with RSL3 (Materials and
Methods), a compound that induces ferroptosis by targeting GPX4, a key regulator of
glutathione oxidation and PUFA reduction (11). Interestingly, this treatment reduces
glutathione metabolism while preventing the reduction of PUFA intermediates, leading to
more precursors of the Fenton reaction and increased ROS, driving cell death by
ferroptosis. We see that idling DTPs are ~3-fold more sensitive to RSL3 by potency (IC50)
than untreated cells (FIG. 4B). Furthermore, the addition of ferrostatin-1, a drug that
inhibits the production of ROS by the Fenton reaction, rescues the enhanced sensitivity
of the idling DTP population to RSL3 (FIG. 4C). These results indicate, therefore, that
idling DTP cells are susceptible to death by ferroptosis induction, presumably because of
increased ROS levels resulting from BRAFi-induced alterations to ion channel
homeostasis and activity (FIG. 3). This is a potential vulnerability of idling DTPs that may
be able to be exploited therapeutically.

DISCUSSION

Drug tolerance has been observed and described in many cancer types (14, 16, 27, 37,
55–57), yet the processes that give rise to DTPs have been interpreted in various ways
(56). Some studies report treatment-induced protective programs in which drug exposure
rapidly triggers new transcriptional or epigenetic responses that were not evident before
treatment (12, 14, 16, 58). Others emphasize rare, pre-existing phenotypic lineages that
differ from the bulk population and become enriched under therapy (13, 15, 59). A third
group of studies highlights stochastic, reversible phenotypic states that cells already visit
before treatment but become stabilized once drug is applied (6, 14, 16, 59, 60). These
varying perspectives emphasize different origins for drug tolerance but acknowledge that
treatment exposure changes which phenotypic states persist within a cancer cell
population and that multiple drug-tolerant states may co-exist (27, 37, 55, 57, 61–63).
However, these prior reports also describe DTPs, at least implicitly, as uniform states in
which each cell within the population is essentially identical.

In this work, we build on these prior studies but provide evidence for a fundamentally
different view of the nature of DTPs, i.e., not as uniform phenotypes but rather as
heterogeneous collections of phenotypic sub-states that, together, achieve near-zero net
population growth under treatment conditions. Specifically, across multiple assays,
untreated and idling DTP populations in BRAF-mutant melanoma exhibited significant

population-level differences, consistent with prior reports (17, 20). In bulk and single-cell transcriptomic data, the idling DTP population showed reduced but still appreciable heterogeneity relative to the untreated population (FIG. 1). This treatment-exposed population occupied a narrower range of phenotypic states, some of which showed distinct functional programs and some that partially overlapped with those observed in untreated conditions (FIG. 1C). Lineage tracing indicated that drug tolerance does not arise from a small set of pre-existing lineages but rather emerges from cells in virtually all lineages (FIG. 2). Single-cell RNA-seq, bulk RNA-seq, and bulk ATAC-seq, together with GO analyses, identified enrichment of ion-channel and calcium-handling pathways, consistent with treatment-associated changes observed in calcium-flux assays (FIG. 3) and with similar treatment-associated shifts in calcium regulation reported in previous melanoma studies (18, 35, 36, 54, 64). Consistent with these transcriptional and functional changes, the idling DTP population exhibited increased sensitivity to ferroptosis induction relative to the untreated population (FIG. 4). Similar vulnerabilities have been reported in other cancer systems, including ferroptosis sensitivity of DTPs and therapy-modulated redox and lipid metabolism (11, 29, 30, 51–53).

It is important to note that a limitation of the present study is that our data represent population snapshots rather than time-lapse trajectories. Thus, we do not directly observe how individual cells move through the landscape over time. Additionally, our ferroptosis experiments establish a population-level vulnerability in this melanoma model but do not identify the specific molecular drivers of that susceptibility, and we cannot assume all DTP populations will share the same vulnerability (11, 30, 51–53). We also did not isolate and analyze drug-tolerant populations following ferroptosis induction, which are also expected to be composed of multiple phenotypic sub-states. Nonetheless, the observations reported here build on prior work (17) and provide substantial experimental support for the view that drug treatment modifies the epigenetic landscape on which cancer cells reside. The population re-equilibrates across this drug-modified landscape that, while less heterogeneous than the untreated landscape, still comprises multiple phenotypic states (FIG. 5A). Moreover, cells from all parts of the untreated landscape transition into the DTP sub-states, which are characterized by distinct molecular processes (20), such as increased oxidative phosphorylation and decreased calcium flux (FIG. 5A). This perspective helps unify diverse observations in the DTP literature (56, 65–68). For example, treatment-induced programs, persistence of pre-existing lineages, and stabilization of fluctuating phenotypes (13–16, 59) can all be viewed as different ways in which a population adjusts to treatment-driven phenotypic changes, rather than fundamentally distinct mechanisms of drug tolerance emergence. The co-existence of multiple DTPs within a drug-tolerant tumor can also be accommodated within this conceptual framework. Not only can multiple DTP states co-exist, across which cells can transition, but each is itself a collection of traversable sub-states that either expand or deplete the total cell population (FIG. 5B).

This view adds an additional layer of complexity to the issue of DTP heterogeneity, with consequences for future therapies aimed at eradicating drug-tolerant populations. The

current philosophy for targeting DTPs focuses on the isolation and molecular
characterization of DTP states to identify actionable vulnerabilities, under the assumption
that the DTP population is phenotypically homogeneous (21–28). Within the framework
proposed here, this approach will be unsuccessful because the phenotypic landscape
that cancer cells occupy is heterogeneous and adjusts in response to perturbations. This
provides a means of escape, whereby cells transition into remaining or emergent viable
sub-states. A game of "whack-a-mole" provides a useful metaphor for this scenario,
where a player constantly chases the mole but is unable to strike it because it moves to
a new location with each whack. Rather than engaging in this futile pursuit, a more fruitful
strategy would be to change the rules of the game and reshape the board itself, e.g., by
filling holes or altering its slope, thereby blocking avenues for the mole's escape.
Analogously, a more durable approach to treating DTPs would be to alter the phenotypic
landscape on which the cells reside. This "targeted landscaping" strategy would entail
multiple sequential interventions to progressively modify the topography of the landscape
to favor regressing over expanding sub-states, until the entire DTP population is
eliminated (FIG. 5C). By shifting the focus to how treatment reshapes the phenotypic
landscape and the nature of the sub-states within it, the framework proposed in this work
offers a generalizable path toward controlling, and potentially eliminating, DTPs before
acquired treatment resistance arises.
418
419　**MATERIALS AND METHODS**
420
421　**Cell culture and reagents**
422
The SKMEL5 cell line, chosen for this study because it exhibits median BRAFi sensitivity
compared to other BRAF-mutant melanoma cell lines (36), was purchased from ATCC©
and labeled with a fluorescent histone H2B conjugated to either the monomeric red
fluorescent protein (H2BmRFP) or the green fluorescent protein (H2B-GFP), as
previously described (6, 17). Cells were initially stored at -80°C and then moved into liquid
nitrogen. Single cell-derived subclones of SKMEL5 were selected and derived by limiting
dilution, as described previously (17). Cells were cultured in a mixed media (1:1) of
Dulbecco's Modified Eagle Medium (DMEM) and Ham's F-12 medium (both from Gibco,
Grand Island, NY, USA), supplemented with 10% fetal bovine serum (FBS) (Thermo
Fisher Scientific, Waltham, MA, USA), incubated at 37°C in 5% CO_2, and passaged twice
a week using TrypLE (Gibco, Grand Island, NY, USA). Cells were also tested for
mycoplasma contamination using the MycoAlert™ mycoplasma detection kit (Lonza,
Basel, Switzerland), according to the manufacturer's instructions, and confirmed to be
mycoplasma-free. SKMEL5-H2BmRFP cells were further transduced with a cellular
barcoding library (see "Cellular barcoding") using HEK293T cells (Thermo Fisher
Scientific, Waltham, MA, USA) to produce lentiviral vectors.
439
440　The BRAFi PLX4720 (research analog to Vemurafenib), ferroptosis inducer RSL3, and
441　ferroptosis inhibitor ferrostatin-1 (Fer-1) were obtained from MedChem Express
442　(Monmouth Junction, NJ, USA) and solubilized in dimethyl sulfoxide (DMSO) at a stock

concentration of 10mM and stored at -20°C. Bovine serum albumin (BSA; Research Products International, Mt. Prospect, IL, USA) and phosphate-buffered saline (PBS; Corning Inc., Corning, NY, USA) were used to prepare cells for single-cell RNA sequencing. Detergents NP-40, Tween20, and Digitonin (MilliporeSigma, Burlington, MA, USA) were used to lyse cells and permeabilize nuclei in preparation for bulk ATAC sequencing. For calcium flux assays, we used calcium chloride ($CaCl_2$; Thermo Fisher Scientific, Waltham, MA, USA, Cat #: J63122.AE) as the source of Ca^{2+}, the Ca^{2+} ionophore ionomycin (Cayman Chemical Company, Ann Arbor, MI, USA, Cat #: 11932), green-fluorescent calcium binding dye Fluo-8-AM (AAT Bioquest, Pleasanton, CA, USA), and Hanks' Balanced Salt Solution (HBSS; no Ca^{2+}; Gibco, Grand Island, NY, USA, Cat #: 14175-095) supplemented with 10 mM HEPES buffer (Corning Inc., Corning, NY, USA, Cat #: 25-060-CI).

Cellular barcoding

Setup: The cellular barcoding library was constructed by cloning a guide RNA (gRNA) library of barcodes into a CROP-seq-BFP-TSO vector, as previously described (43, 44). The vector was engineered such that barcodes can be found by isolation and amplification (barcode sampling) or mRNA capture in a single-cell RNA sequencing (scRNA-seq) experiment. gRNAs were built as a 20-nucleotide sequence of four nucleotides identical among all barcodes (XXXX), followed by 16 strong-weak (SW) paired nucleotides (i.e., XXXXSWSWSWSWSWSWSWSW). The SW pairing of the barcode sequence was designed to prevent polymerase chain reaction (PCR) amplification bias. The maximum complexity of this library is 2^{16} (~65,000 unique barcodes). The barcode library vector was used to produce lentiviral libraries using a lipofectamine transfection of HEK293T cells. Media containing lentiviral particles were collected at 48- and 72-hours post-transfection, pooled, filtered through a 0.45 μm Nalgene syringe filter (Thermo Fisher Scientific, Waltham, MA, USA), and concentrated using a 50 mL size-exclusion column (MilliporeSigma, Burlington, MA, USA) by centrifugation at 2200 relative centrifugal field (RCF) at 4°C for two hours. Concentrated virus was stored in -80°C. SKMEL5 cells were seeded in a 6-well plate at ~1x10^6 cells per well in 2.5 mL culture media. Cells were transduced with the barcoded CROP-seq-BFP-TSO-Barcode_sgRNA lentivirus using 0.8 mg/mL in each well and a multiplicity of infection (MOI) of 0.05, as previously described (43, 44). Twenty-four hours after incubation, transduction media (containing polybrene) was exchanged for fresh culture media. Forty-eight hours after incubation, barcoded cells were isolated by fluorescence-activated cell sorting (FACS) and subsequently cultured until confluence in a T-150 dish and cryopreserved. Cells were then thawed in a T-25 dish and scaled up for ~2 weeks in two separate sets. The first set of thawed cells was treated with 8 μM PLX4720 for eight days and subjected to barcode sampling (see "Cellular Barcoding: *Barcode Sampling Analysis*"), along with an untreated control sample. The second set of thawed cells was plated in three T-75 flasks (parallel replicates) and independently treated with 8 M PLX4720 for eight days and subjected to scRNA-seq (along with an untreated control sample) by the 10X genomics Chromium platform (version 2 chemistry; see "Single-cell

RNA transcriptomic sequencing"). For all drug-treated samples, media and drug were
replaced every three days. Untreated cells were expanded completely over the entire time
course (no cell splitting).

Barcode Sampling: For the first set of thawed cells (see "Cellular Barcoding: *Setup*"),
untreated and 8-day PLX4720-treated cells were pelleted for gDNA extraction using the
DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany), per manufacturers'
instructions. Barcode sequences were amplified for each replicate by PCR (98°C for 30
seconds, followed by 22 cycles of denaturation at 98°C for 10 seconds, annealing at 63°C
for 30 seconds, extension at 72°C for 10 seconds, and a final extension at 72°C for 5
minutes) using primers containing flanking regions and Illumina adapter index sequences
(Illumina, San Diego, CA, USA). For each PCR reaction, 2 μg gDNA was used and a
combination of five distinct pooled forward primers was utilized to minimize sequencing
error. Reactions were purified using a 1.8x AMPure XP bead (Beckman Coulter, Brea,
CA, USA) cleanup. Reaction products were confirmed using agarose gel confirmation
(band at ~215 bp). The resulting libraries were quantified using a Qubit fluorometer
(Thermo Fisher Scientific, Waltham, MA, USA), Bioanalyzer 2100 (Agilent Technologies,
Santa Clara, CA, USA) for library profile assessment, and quantitative PCR (qPCR; Kapa
Biosystems, Wilmington, MA, USA, Cat #: KK4622) to validate ligated material, according
to the manufacturer's instructions. The libraries were sequenced using the NovaSeq 6000
with 150 bp paired-end reads as sequencing spike-ins (targeting ~200,000 reads). 'Real-
Time Analysis' ('RTA') software (version 2.4.11; Illumina, San Diego, CA, USA) was used
for base calling and 'MultiQC' (version 1.7) (69) was used for quality control. Barcodes
were identified from amplified sequence reads by trimming flanking adapter sequences
(5': XXXX; 3': XXXX). Barcode abundances were totaled and normalized to the library
read depth, resulting in reads per million (RPM). Barcodes less than 100 counts per
million (CPM) were removed from the analysis. Numbers of unique barcodes were
calculated based on this threshold.

Single-cell RNA transcriptomic sequencing

Data Collection: For the second set of thawed cells (see "Cellular Barcoding: *Setup*"),
untreated and 8-day PLX4720-treated cells were prepared targeting ~3000 cells per
sample, washed, and resuspended in 0.04% BSA in PBS. Cell suspensions were
subjected to a 10X Genomics single-cell gene expression protocol (version 2, 3'
counting; 10xgenomics.com/support/single-cell-gene-expression/documentation/steps/
library-prep/assay-scheme-and-configuration-of-chromium-single-cell-3-v-2-libraries) in
two separate wells, according to the manufacturer's guidelines. Single-cell mRNA
expression libraries were prepared according to the manufacturer's instructions. Due to
the nature of the gRNA barcoding library construction (43, 44), mRNAs resulting from
gRNA barcodes were captured along with other mRNAs. Libraries were cleaned using
solid-phase reversible immobilization (SPRI) beads (Beckman Coulter, Brea, CA, USA)
and quantified using a Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA, USA).
The libraries were sequenced using the NovaSeq 6000 (Illumina, San Diego, CA, USA)

with 150 bp paired-end reads targeting 50 million reads per sample for the mRNA library
(including the barcode library). 'RTA' software (version 2.4.11; Illumina, San Diego, CA,
USA) was used for base calling and 'MultiQC' (version 1.7) (69) for quality control. Gene
counting, including alignment, filtering, barcode counting, and unique molecular identifier
(UMI) counting was performed on each library using the '*count*' function in the 10X
Genomics software 'Cell Ranger' (version 3.0.2) (70) with the GRCh38 (hg38) reference
transcriptome (71).
538
Transcriptome Analysis: We used 'sc-UniFrac' (60) to quantify the degree of overlap
between the two scRNA-seq count matrices (untreated and idling DTP) obtained from
'Cell Ranger' (see "Single-cell RNA transcriptomic sequencing: *Data Collection*"). Since
overlaps were minimal and cells were prepared and processed in parallel, no
computational batch correction was performed. We used 'Seurat' (version 3) (72) to
perform gene expression analysis. The '*SCTransform*' function was used to regress out
mitochondrial gene expression ('percent.mt'), number of features (i.e., genes;
'nFeature_RNA'), number of RNA molecules in the cell ('nCount_RNA'), and cell cycle
variables ('S.Score' and 'G2M.Score'). Feature selection was performed according to
'Seurat' guidelines using a variance stabilizing transformation of the top 2000 most
variable features. Data was normalized and scaled according to 'Seurat' guidelines. Data
between conditions were combined and visualized using the UMAP dimensionality
reduction algorithm (31), as implemented in 'Seurat'. Clustering was performed in the joint
UMAP space using the default 'Seurat' implementation, a shared nearest neighbor (SNN)
modularity optimization-based method (73). Differential expression analysis was
performed using the '*FindMarkers*' function in 'Seurat'. Differentially expressed genes
(DEGs; adjusted-p < 0.05) were input into a GO over-enrichment analysis (38).
556
Barcode Analysis: After calculating the scRNA-seq gene expression matrices with 'Cell
Ranger' (see above), barcode abundances were incorporated into the matrices. To do
this, gRNA lineage barcodes were mapped to their associated 10X cell barcodes. First,
unmapped scRNA-seq Binary Alignment Map (BAM) files were cleaned to only include
the mRNA transcript ID, scRNA-seq cell barcode, and scRNA-seq UMI. Mapped scRNA-
seq BAM files (3' heavy) were cleaned to only include the mRNA transcript ID and lineage
barcode (from the gRNA library). Unmapped and mapped subsets were then merged on
the mRNA transcript ID to assign a lineage barcode to each cell barcode and UMI. The
resulting merged dataset was paired down to a cell barcode/lineage barcode pair, which
was appended to each cell in the gene expression matrix as a metadata tag. Barcode
abundances were totaled across all cells in the experiment that captured a barcode.
Barcodes overlaid on UMAP projections of scRNA-seq data were further categorized into
"fast-dividing" (G1) and "slow-dividing" (S-G2-M) states based on cell cycle score (39)
(see "Single-cell RNA transcriptomic sequencing: *Transcriptome Analysis*").
571
Functional Interpretation Analysis: The single-cell transcriptome count matrices obtained
from 'Cell Ranger' (see "Single-cell RNA transcriptomic sequencing: *Data Collection*")
were scaled by multiplying counts by the median RNA molecules across all cells and

dividing that number by the number of RNA molecules in each cell. Hallmark gene sets
(50 in total) were downloaded from the molecular signatures database (MSigDB) (32–
34). The scaled count matrix and each hallmark gene set were input into 'VISION' (74) to
identify gene signature scores for each cell-signature pair. Four hallmark gene sets
(KRAS_SIGNALING_UP, KRAS_SIGNALING_DOWN, UV_RESPONSE_UP,
UV_RESPONSE_DOWN) were condensed into two (KRAS_SIGNALING,
UV_RESPONSE) by 'VISION' to leave 48 total gene signatures. Scores were compiled
into a distribution and plotted by cluster (k=3) for each gene set.
583
Bulk RNA transcriptomic sequencing
585
Data Acquisition: Total RNA was isolated from untreated and BRAFi-treated SKMEL5
single cell-derived subclones, each in triplicate, using the TRIzol isolation method
(Invitrogen, Carlsbad, CA, USA), according to the manufacturer's instructions. RNA
samples were submitted to Vanderbilt VANTAGE Core services for quality check, where
mRNA enrichment and cDNA library preparation were done with Illumina TruSeq
stranded mRNA sample prep kit (Illumina, San Diego, CA, USA). Sequencing was done
at Paired-End 75 bp on the Illumina HiSeq 3000. Reads were aligned to the hg38 human
reference genome (71) using 'HISAT2' (75) and gene counts were obtained using
'featureCounts' (76).
595
Data Analysis: RNA-seq data was analyzed using the 'DESeq2' R package (77). Cells
with less than 18 reads per condition were removed, according to 'DESeq2' vignettes.
Counts were transformed using the regularized logarithm ('*rlog*') normalization algorithm.
PCA was performed using the '*prcomp*' function in R (rdrr.io/r/stats/prcomp.html).
Differential expression analysis was performed using a 'DESeq2' model design to quantify
both changing variables and their interaction ($\sim$ subline + treatment time +
subclone:treatment time). DEGs across subclones between untreated (pre-treatment,
day 0) and idling DTP (post-treatment, day 8) were identified (adjusted-$p < 0.05$, log2 fold-
change > 2) and input into a GO enrichment analysis (38), which identified GO terms
associated with "Biological Process" (BP), "Molecular Function" (MF), and "Cellular
Component" (CC) GO categories.
607
Calcium Handling Gene Signature Analysis: Using the 'AmiGO' web application (78)
genes associated with calcium handling were identified by the GO term "intracellular
calcium ion homeostasis" (Accession #: GO:0006874; Release date: 2023-11-17; Version
#: 10.5281/zenodo.10162580). Inositol 1,4,5 trisphosphate receptors (ITPRs) and other
transient receptor potential melastatin (TRPM) receptors were also added manually to the
gene set.
614
615 *Ferroptosis Gene Signature Analysis:* A ferroptosis gene signature was obtained from the
616 WikiPathways database (wikipathways.org/pathways/WP4313.html) (79, 80). For each
617 single cell-derived subclone, mRNA levels were normalized to the 0-day time point and a
618 log_2 fold-change was calculated compared to the 0-day baseline.

Bulk ATAC epigenomic sequencing

Data Acquisition: Data was collected using the Omni-ATAC protocol (81) for bulk assay for transposase-accessible chromatin with sequencing (ATAC-seq). For the second set of thawed cells (see "Cellular Barcoding: *Setup*"), untreated and 8-day PLX4720-treated cells were pelleted at ~50,000 cells and resuspended in a cold ATAC-seq resuspension and lysis buffer containing NP-40 (0.1%), Tween20 (0.1%), and Digitonin (0.01%) and incubated on ice. A resuspension buffer was added (0.1% Tween20, no NP-40 or Digitonin) to wash out the lysis reaction. Cells were pelleted and resuspended in a transposition mix (5x Tris-DMF, phosphate-buffered saline (PBS), 1% Digitonin, 10% Tween20, nuclease-free H_2O), including transposase Tn5, followed by a 30-minute incubation at 37°C, with shaking to enhance tagmentation. After 30 minutes, the reaction was stopped by adding a DNA binding buffer (Zymo Research, Irvine, CA, USA) and purified using a DNA Clean and Concentrate kit (Zymo Research, Irvine, CA, USA, Cat #: D4004). The final product was eluted in nuclease-free H_2O. PCR amplification was performed on the eluate with an NEBNext High-Fidelity 2X PCR Master Mix (New England Biolabs, Ipswich, MA, USA, Cat #: M0541S) and custom primer indexes (extension at 72°C for 5 minutes; denaturation at 90°C for 30 seconds; 12 cycles: denaturation at 98°C for 10 seconds, annealing at 62°C for 30 seconds, extension at 72°C for 30 seconds; final extension at 72°C for 5 minutes). The PCR product was purified with the Zymo DNA Clean and Concentrate kit and eluted in 22 mL nuclease-free H_2O. ATAC-seq PCR libraries were visualized by agarose gel electrophoresis for a quick check for the nucleosome ladder pattern (bands over ~150 bp). Libraries were also quantified using a Qubit fluorometer (Thermo Fisher Scientific, Waltham, MA, USA), Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA, USA) for library profile assessment, and qPCR (Kapa Biosystems, Wilmington, MA, USA, Cat #: KK4622) to validate ligated material, according to the manufacturer's instructions. The libraries were sequenced using the NovaSeq 6000 with 150 bp paired-end reads as spike-ins on the sequencing chip (untreated: ~160 million reads, idling DTP: ~130 million reads). 'RTA' software (version 2.4.11; Illumina, San Diego, CA, USA) was used for base calling and 'MultiQC' (version 1.7) (69) for quality control.

Data Analysis: Reads were trimmed using 'cutadapt' (82) (paired-end) to remove primer sequences and aligned to the hg38 reference genome (71) using the '*bwa-mem*' function in 'Burrows-Wheeler Aligner' (BWA; version 0.7.17) (83). Aligned reads were sorted and duplicates were marked using the 'Picard' software tool (version 2.17.10; broadinstitute.github.io/picard). Untreated reads had more detected duplicates (~78% compared to ~32% in idling DTPs). Reads were deduplicated, but to address the discrepancy in deduplicated library complexity, idling DTP reads were subsampled (25% of the original library) to achieve a similar complexity to the untreated sample (FIG. S3A,B). Insert sizes were plotted from the output of '*InsertSizeMetrics*' in 'Picard' after deduplication. Peaks of open chromatin were called using the *'callpeak'* function in 'MACS2' (84), according to recommended guidelines for ATAC-seq data (BAM paired-

end method, q-threshold: 0.05, no 'MACS2' model, shift: -100, extension size: 200).
Peaks were subjected to a further round of quality control and cleaning using 'ChIPQC'
(85) (peak mapping, peak duplication, blacklist peak detection) and blacklisted peaks
were removed. Peaks were converted to consensus counts using the
'*runConsensusCounts*' function in 'soGGi' (rdrr.io/bioc/soGGi/). Intersections and unique
cleaned peaks were determined and visualized as a Venn diagram (FIG. S3C) using the
'*vennDiagram*' function in the 'limma' software package (86). Unique and intersection
peaks were annotated with nearest neighbor genes using the '*annotatePeak*' function and
the hg38 transcriptome (71) in the 'ChIPseeker' software package (87). These peaks
were also re-aligned to the transcription start site (TSS) for each gene (FIG. S3D). Peaks
were classified based on closeness to the TSS and assigned to a predicted feature
(promoter, UTR, exon, intron, downstream, distal intergenic; FIG. S3E). Genes
associated with unique peaks and intersections of peaks were also input into a GO
enrichment analysis for BP, MF, and CC GO categories (same as "Bulk RNA
Transcriptomic Sequencing: *Data Analysis*" above).
678
679 **Gene ontology analysis**
680
Genes associated with unique ATAC-seq peaks (see "Bulk ATAC Epigenomic
Sequencing: *Data Analysis*") and DEGs from bulk RNA-seq (for all single cell-derived
subclones; see "Bulk RNA Transcriptomic Sequencing: *Data Analysis*") were identified
for untreated and idling DTP cell populations. The two gene lists were independently
subjected to a GO enrichment analysis using 'clusterProfiler' (88, 89) and genes were
compared to BP, MF, and CC GO categories. For each category, GO terms significantly
enriched in the unique ATAC-seq peaks ($p < 0.05$) and in DEGs ($p < 0.05$) were identified
and stored independently as separate GO term lists. For all GO terms shared between
the two lists, we calculated $-\log_{10}$(p-value) and ranked terms based on statistical
significance. Spearman correlation (90) was calculated between the significant GO terms
using 'ggpubr' (version 0.4.0; rpkgs.datanovia.com/ggpubr).
692
693 **Calcium flux assays**
694
Data Acquisition: Barcoded SKMEL5 cells were plated onto a 384-well tissue-culture-
treated plate 24 hours before imaging at a density of 10,000 cells/well. For the treated
condition, cells were treated with 8 μM PLX4720 for eight days, with fresh media/drug
swapped out every three days. Cells were trypsinized, counted, and seeded to imaging
plates, all while in the continuous presence of 8 μM PLX4720. Untreated control cells
were taken from the same cell line culture, which was maintained separately. On the day
of experimentation, the 384-well plate with all treated and untreated cells was incubated
with 4 μM Fluo-8-AM in fresh culture media (10% FBS) for 1 hour at room temperature,
as recommended by the manufacturer. Dye-containing medium was removed and HBSS
(10 mM HEPES, no Ca^{2+}) was used to wash the wells of excess dye, followed by removal
and addition of 20 μL of fresh HBSS for use as the assay buffer. The plate was loaded
into a Panoptic kinetic imaging plate reader (WaveFront Biosciences, Franklin, TN, USA),

which combines liquid handling capabilities and the ability to record fluorescent intensity
from each well of the 384-well plate. A custom three-addition protocol was developed to
add the various drugs and assay conditions to the plate during the SOCE assay. Drug-
addition reservoir plates were loaded with assay buffer (HBSS, with or without Ca^{2+}) and
thoroughly mixed immediately before imaging. Add conditions were divided into three
parts: 1) cyclopiazonic acid (CPA; final concentration of 50 μM) to inhibit the activity of
Sarcoendoplasmic Reticulum Calcium ATPase (SERCA), leading to ER Ca^{2+} release; 2)
addition of Ca^{2+} to the assay condition to activate SOCE activity; and 3) addition of
ionomycin, the Ca^{2+} ionophore (final concentration of 5 μM), to generate maximal signal
intensity to control for variations in cell count in individual wells (this was particularly
important since drug-treated idling DTP cells experienced increased washout due to the
stressful nature of sustained BRAFi treatment). Fluo-8-AM was excited at 480 nm and
imaged at 538 nm, with a frequency of 1 Hz. The CPA treatment condition was imaged
for 260 seconds before the addition of Ca^{2+}, followed by 270 seconds of imaging before
the addition of 5 μM ionomycin. Treatment conditions were replicated in sets of eight and
average values traced with 95% confidence intervals.
723
Data Analysis: For each condition (untreated and idling), fluorescence data were
normalized by subtracting from each well the lowest average intensity value near the start
of the experiment (t < 10 s) and then dividing by the highest average intensity value during
the ionomycin treatment phase (t > 530 s) to control for the number of cells in each
treatment group (the idling DTP population tended to have fewer cells than the untreated
population). For comparing calcium flux between the treatment groups, means and
standard errors ($\sigma/\sqrt{N}$) were calculated for each time point and plotted using 'ggplot2'
(version 3.2.0) (91). For visualization purposes, the earliest time points (t < 10 s) and data
from the ionomycin treatment phase (t > 530 s) were excluded from the plots.
733
734 **Ferroptosis-induction experiments**
735
Plates of H2B-GFP-labeled SKMEL5 cells (see "Cell culture and reagents") were treated
with either vehicle (DMSO) or BRAFi (8 μM PLX4720) for eight days, incubated at 37°C
and 5% CO_2, changing media (with vehicle/drug) every three days. BRAFi-treated cells
were plated at ~2500 cells per well in a black, clear-bottom 96-well plate (Corning Inc.,
Corning, NY, USA). After cell seeding, RSL3, with or without Fer-1, was added the
following morning, with media changes every three days (six replicates per condition).
Plates were imaged using automated fluorescence microscopy (Cellavista Instrument,
Synentec, Elmshorn, Germany). Twenty-five non-overlapping fluorescent images (20X
objective, 5x5 montage) were taken twice daily for a total of 150 hours or until confluency.
Cellavista image segmentation software (Synentec, Elmshorn, Germany) was utilized to
calculate nuclear count (i.e., cell count) per well at each time point (Source = FITC, Dichro
= FITC, Filter = FITC, Emission Time = 800 ms, Gain = 20x, Quality = High, Binning =
2x2). Cell nuclei counts across wells were normalized to the time of drug treatment and
used to calculate drug-induced proliferation (DIP) rates, obtained as the slope of a linear
fit to the log2-scaled population growth curve (92, 93). A dose-response curve was

calculated across replicates using the 'drm' R package (94) with a 4-parameter log-logistic
function, with DIP rate as the drug effect, as previously described (93). Replicates were
used to calculate means and 95% confidence intervals for the dose-response curves.
Data was visualized using the 'ggplot2' R package (version 3.2.0) (91).
755

756 **Model and experimental analysis code availability.** The codes used to generate model
757 simulations and analyze experimental data are publicly available at GitHub
758 (github.com/SysBioCollab-UArk/Hayford_Melanoma_DTP_2026).
759

760 **Data availability.** Sequencing data generated in this study have been deposited in the
761 Gene Expression Omnibus (GEO) under accession GSE324655. Raw sequencing reads
762 are available through the Sequence Read Archive (SRA) under BioProject accession
763 PRJNA1431156. Processed data and analysis codes are available at GitHub
764 (github.com/SysBioCollab-UArk/Hayford_Melanoma_DTP_2026).
765

766 **ACKNOWLEDGEMENTS**
767

We thank Emily Hodges and Kelly Barnett for ATAC-seq reagents and experimental
advice. We thank Jing Hao for reagent acquisition; Tony Capra, Christian Meyer, Sarah
Maddox Groves, Carlos Lopez, Alissa Weaver, John McLean, Bruce Damon, and Ken
Lau for useful discussions; and Rachana Nitin for providing feedback on an earlier version
of this manuscript. B.B. thanks Nancy Cox and Christopher Wright for continued
mentorship and support, and Jennifer (Piper) Below, Lea Davis, and Eric Gamazon for
valued feedback and guidance. Sequencing studies were performed at the Vanderbilt
Technologies for Advanced Genomics (VANTAGE) core facility at Vanderbilt University
Medical Center with help from Angela Jones, Karen Beeri, Jamie Roberson, Latha Raju,
and Matthew Scholz. Data processing and bioinformatics analyses were performed using
computational resources at the Advanced Computing Center for Research and Education
(ACCRE) at Vanderbilt University. We also acknowledge the Vanderbilt High Throughput
Screening (HTS) core for contributions to this work.
781

This work was supported by the following funding sources: C.E.H., National Institutes of
Health (NIH) Ruth L. Kirschstein National Research Service Award (NRSA, F31-
CA221147) and Chemical-Biology Interface Training Grant (T32-GM065086); B.B.,
Integrated Biological Systems Training in Oncology Training Grant (T32-CA119925);
P.E.S., Molecular Endocrinology Training Grant (T32-MH064913); B.B.P., Vanderbilt
Institute for Clinical and Translational Research (VICTR) Grants (VR16721 and
VR16721.1.); V.Q., NIH Clinical and Translational Science Award (U54-CA113007);
D.R.T., NIH Research Specialist Award (R50-CA243783); L.A.H., Vanderbilt Biomedical
Informatics Training Program (NLM 5T15-LM007450-14), Quantitative Systems Biology
Center at Vanderbilt, and National Cancer Institute (NCI) Transition Career Development
Award (K22-CA237857). In addition, this material is based upon work supported by the
National Science Foundation Graduate Research Fellowship Program under Grant No.
1937963 (to B.B). Any opinions, findings, and conclusions or recommendations

expressed in this material are those of the authors and do not necessarily reflect the views
of the National Science Foundation.
797
798 **COMPETING INTERESTS**
799
800 V.Q. is an academic cofounder and equity holder of Duet BioSystems and shareholder of
801 Incendia Therapeutics. The other authors declare no competing interests.
802
803 **AUTHOR CONTRIBUTIONS**
804
805 Conceptualization: C.E.H., B.B., P.E.S., V.Q., D.R.T., L.A.H.; Experimental Data
806 Collection: C.E.H., P.E.S., B.B.P., A.A., D.R.T.; Analysis and Interpretation of Results:
807 C.E.H., B.B., P.E.S., V.Q., D.R.T., L.A.H.; Bioinformatics Analysis: C.E.H., B.B., P.E.S.;
808 Writing: C.E.H., B.B., V.Q., L.A.H.; Review and Editing: C.E.H., B.B., P.E.S., A.B., V.Q.,
809 D.R.T., L.A.H.
810
811 **GLOSSARY**
812 *This glossary defines key terms as used in this study, which may differ from standard*
813 *usage in the literature.*
814
815 *Phenotype*: The observable characteristics of a cell, such as morphology, migratory
816 ability, and metastatic potential.
817
818 *Phenotypic state*: A discrete category of closely related phenotypes that share common
819 molecular or functional features. Individual cells are assigned to one phenotypic state at
820 any given time. Although cells within a phenotypic state may vary at the molecular level,
821 they are considered sufficiently similar to be classified together. Phenotypic states are
822 also commonly referred to as "cell states" or "single-cell states."
823
824 *Population state*: A collection of cells distributed across a discrete set of phenotypic
825 states. The population state refers to the equilibrium distribution of cells across these
826 states under a defined set of conditions. At equilibrium, individual phenotypic states may
827 be expanding or regressing, but their combined contributions determine the overall
828 population behavior.
829
DTP population state: A drug-induced population state in which the equilibrium
distribution of cells across discrete phenotypic states results in a near-zero net
proliferation rate for the population as a whole. Cells may transition among individual
phenotypic states with differing proliferation rates under drug treatment, but their
combined contributions balance so that the overall population size remains effectively
constant.
836
837 **REFERENCES**
838

1. Dagogo-Jack, I., and A.T. Shaw. 2018. Tumour heterogeneity and resistance to
cancer therapies. *Nat. Rev. Clin. Oncol.* 15:81–94, doi:
10.1038/nrclinonc.2017.166.
2. Marusyk, A., and K. Polyak. 2010. Tumor heterogeneity: Causes and
consequences. *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer*
1805:105–117, doi: 10.1016/J.BBCAN.2009.11.002.
3. Marusyk, A., V. Almendro, and K. Polyak. 2012. Intra-tumour heterogeneity: a
looking glass for cancer? *Nat. Rev. Cancer* 12:323–334, doi: 10.1038/nrc3261.
4. Brock, A., H. Chang, and S. Huang. 2009. Non-genetic heterogeneity - a mutation-
independent driving force for the somatic evolution of tumours. *Nat. Rev. Genet.*
10:336–42, doi: 10.1038/nrg2556.
5. Burdziak, C., D. Alonso-Curbelo, T. Walle, J. Reyes, F.M. Barriga, D. Haviv, Y. Xie,
Z. Zhao, C.J. Zhao, H.A. Chen, O. Chaudhary, I. Masilionis, Z.N. Choo, V. Gao, W.
Luan, A. Wuest, Y.J. Ho, Y. Wei, D.F. Quail, R. Koche, L. Mazutis, R. Chaligné, T.
Nawy, S.W. Lowe, and D. Pe'er. 2023. Epigenetic plasticity cooperates with cell-
cell interactions to direct pancreatic tumorigenesis. *Science (1979).* 380:eadd5327,
doi: 10.1126/science.add5327.
6. Hayford, C.E., D.R. Tyson, C.J. Robbins, P.L. Frick, V. Quaranta, and L.A. Harris.
2021. An in vitro model of tumor heterogeneity resolves genetic, epigenetic, and
stochastic sources of cell state variability. *PLoS Biol.* 19:e3000797, doi:
10.1371/journal.pbio.3000797.
7. Huang, S. 2013. Genetic and non-genetic instability in tumor progression: link
between the fitness landscape and the epigenetic landscape of cancer cells.
Cancer and Metastasis Reviews 32:423–448, doi: 10.1007/s10555-013-9435-7.
8. Raghavan, S., P.S. Winter, A.W. Navia, H.L. Williams, A. DenAdel, K.E. Lowder, J.
Galvez-Reyes, R.L. Kalekar, N. Mulugeta, K.S. Kapner, M.S. Raghavan, A.A.
Borah, N. Liu, S.A. Väyrynen, A.D. Costa, R.W.S. Ng, J. Wang, E.K. Hill, D.Y.
Ragon, L.K. Brais, A.M. Jaeger, L.F. Spurr, Y.Y. Li, A.D. Cherniack, M.A. Booker,
E.F. Cohen, M.Y. Tolstorukov, I. Wakiro, A. Rotem, B.E. Johnson, J.M. McFarland,
E.T. Sicinska, T.E. Jacks, R.J. Sullivan, G.I. Shapiro, T.E. Clancy, K. Perez, D.A.
Rubinson, K. Ng, J.M. Cleary, L. Crawford, S.R. Manalis, J.A. Nowak, B.M. Wolpin,
W.C. Hahn, A.J. Aguirre, and A.K. Shalek. 2021. Microenvironment drives cell
state, plasticity, and drug response in pancreatic cancer. *Cell* 184:6119-6137.e26,
doi: 10.1016/J.CELL.2021.11.017.
9. Su, Y., M.E. Ko, H. Cheng, R. Zhu, M. Xue, J. Wang, J.W. Lee, L. Frankiw, A. Xu,
S. Wong, L. Robert, K. Takata, D. Yuan, Y. Lu, S. Huang, A. Ribas, R. Levine, G.P.
Nolan, W. Wei, S.K. Plevritis, G. Li, D. Baltimore, and J.R. Heath. 2020. Multi-omic
single-cell snapshots reveal multiple independent trajectories to drug tolerance in
a melanoma cell line. *Nat. Commun.* 11:2345, doi: 10.1038/s41467-020-15956-9.
10. Tavernari, D., E. Battistello, E. Dheilly, A.S. Petruzzella, M. Mina, J. Sordet-
Dessimoz, S. Peters, T. Krueger, D. Gfeller, N. Riggi, E. Oricchio, I. Letovanec, and
G. Ciriello. 2021. Nongenetic evolution drives lung adenocarcinoma spatial
heterogeneity and progression. *Cancer Discov.* 11:1490–1507, doi: 10.1158/2159-
8290.CD-20-1274.

11.   Hangauer, M.J., V.S. Viswanathan, M.J. Ryan, D. Bole, J.K. Eaton, A. Matov, J.
Galeas, H.D. Dhruv, M.E. Berens, S.L. Schreiber, F. McCormick, and M.T.
McManus. 2017. Drug-tolerant persister cancer cells are vulnerable to GPX4
inhibition. *Nature* 551:247–250, doi: 10.1038/nature24297.

12.   Hata, A.N., M.J. Niederst, H.L. Archibald, M. Gomez-Caraballo, F.M. Siddiqui, H.E.
Mulvey, Y.E. Maruvka, F. Ji, H.-E.C. Bhang, V. Krishnamurthy Radhakrishna, G.
Siravegna, H. Hu, S. Raoof, E. Lockerman, A. Kalsy, D. Lee, C.L. Keating, D.A.
Ruddy, L.J. Damon, A.S. Crystal, C. Costa, Z. Piotrowska, A. Bardelli, A.J. Iafrate,
R.I. Sadreyev, F. Stegmeier, G. Getz, L. V Sequist, A.C. Faber, and J.A. Engelman.
2016. Tumor cells can follow distinct evolutionary paths to become resistant to
epidermal growth factor receptor inhibition. *Nat. Med.* 22:262–269, doi:
10.1038/nm.4040.

13.   Oren, Y., M. Tsabar, M.S. Cuoco, L. Amir-Zilberstein, H.F. Cabanos, J.-C. Hütter,
B. Hu, P.I. Thakore, M. Tabaka, C.P. Fulco, W. Colgan, B.M. Cuevas, S.A. Hurvitz,
D.J. Slamon, A. Deik, K.A. Pierce, C. Clish, A.N. Hata, E. Zaganjor, G. Lahav, K.
Politi, J.S. Brugge, and A. Regev. 2021. Cycling cancer persister cells arise from
lineages with distinct programs. *Nature* 596:576–582, doi: 10.1038/s41586-021-
03796-6.

14.   Ramirez, M., S. Rajaram, R.J. Steininger, D. Osipchuk, M.A. Roth, L.S. Morinishi,
L. Evans, W. Ji, C.-H. Hsu, K. Thurley, S. Wei, A. Zhou, P.R. Koduru, B.A. Posner,
L.F. Wu, and S.J. Altschuler. 2016. Diverse drug-resistance mechanisms can
emerge from drug-tolerant cancer persister cells. *Nat. Commun.* 7:10690, doi:
10.1038/ncomms10690.

15.   Rehman, S.K., J. Haynes, E. Collignon, K.R. Brown, Y. Wang, A.M.L. Nixon, J.P.
Bruce, J.A. Wintersinger, A. Singh Mer, E.B.L. Lo, C. Leung, E. Lima-Fernandes,
N.M. Pedley, F. Soares, S. McGibbon, H.H. He, A. Pollet, T.J. Pugh, B. Haibe-
Kains, Q. Morris, M. Ramalho-Santos, S. Goyal, J. Moffat, and C.A. O'Brien. 2021.
Colorectal cancer cells enter a diapause-like DTP state to survive chemotherapy.
Cell 184:226–242, doi: 10.1016/j.cell.2020.11.018.

16.   Sharma, S. V, D.Y. Lee, B. Li, M.P. Quinlan, F. Takahashi, S. Maheswaran, U.
McDermott, N. Azizian, L. Zou, M.A. Fischbach, K.-K. Wong, K. Brandstetter, B.
Wittner, S. Ramaswamy, M. Classon, and J. Settleman. 2010. A chromatin-
mediated reversible drug-tolerant state in cancer cell subpopulations. *Cell* 141:69–
80, doi: 10.1016/j.cell.2010.02.027.

17.   Paudel, B.B., L.A. Harris, K.N. Hardeman, A.A. Abugable, C.E. Hayford, D.R.
Tyson, and V. Quaranta. 2018. A nonquiescent "idling" population state in drug-
treated, BRAF-mutated melanoma. *Biophys. J.* 114:1499–1511, doi:
10.1016/j.bpj.2018.01.016.

18.   Stauffer, P.E., J. Brinkley, D.A. Jacobson, V. Quaranta, and D.R. Tyson. 2024.
Purinergic Ca2+ signaling as a novel mechanism of drug tolerance in BRAF-mutant
melanoma. *Cancers (Basel).* 16:2426, doi: 10.3390/CANCERS16132426/S1.

19.   Aranda-Anzaldo, A., and M.A.R. Dent. 2018. Landscaping the epigenetic
landscape of cancer. *Prog. Biophys. Mol. Biol.* 140:155–174, doi:
10.1016/j.pbiomolbio.2018.06.005.

20. Jia, D., B.B. Paudel, C.E. Hayford, K.N. Hardeman, H. Levine, J.N. Onuchic, and
V. Quaranta. 2020. Drug-tolerant idling melanoma cells exhibit theory-predicted
metabolic low-low phenotype. *Front. Oncol.* 10:1426, doi:
10.3389/FONC.2020.01426.
21. Russo, M., M. Chen, E. Mariella, H. Peng, S.K. Rehman, E. Sancho, A. Sogari, T.S.
Toh, N.Q. Balaban, E. Batlle, R. Bernards, M.J. Garnett, M. Hangauer, E. Leucci,
J.-C. Marine, C.A. O'Brien, Y. Oren, E.E. Patton, C. Robert, S.M. Rosenberg, S.
Shen, and A. Bardelli. 2024. Cancer drug-tolerant persister cells: from biological
questions to clinical opportunities. *Nat. Rev. Cancer* 24:694–717, doi:
10.1038/s41568-024-00737-z.
22. Cabanos, H.F., and A.N. Hata. 2021. Emerging insights into targeted therapy-
tolerant persister cells in cancer. *Cancers (Basel).* 13:2666, doi:
10.3390/CANCERS13112666.
23. Cai, S.F., C.W. Chen, and S.A. Armstrong. 2015. Drugging chromatin in cancer:
recent advances and novel approaches. *Mol. Cell* 60:561–570, doi:
10.1016/J.MOLCEL.2015.10.042.
24. Mikubo, M., Y. Inoue, G. Liu, and M.S. Tsao. 2021. Mechanism of drug tolerant
persister cancer cells: the landscape and clinical implication for therapy. *Journal of
Thoracic Oncology* 16:1798–1809, doi: 10.1016/j.jtho.2021.07.017.
25. Pu, Y., L. Li, H. Peng, L. Liu, D. Heymann, C. Robert, F. Vallette, and S. Shen.
2023. Drug-tolerant persister cells in cancer: the cutting edges and future
directions. *Nat. Rev. Clin. Oncol.* 20:799–813, doi: 10.1038/s41571-023-00815-5.
26. Settleman, J., J.M. Fernandes Neto, and R. Bernards. 2021. Thinking differently
about cancer treatment regimens. *Cancer Discov.* 11:1016–1023, doi:
10.1158/2159-8290.CD-20-1187.
27. Shen, S., S. Vagner, and C. Robert. 2020. Persistent cancer cells: the deadly
survivors. *Cell* 183:860–874, doi: 10.1016/J.CELL.2020.10.027.
28. Zhang, Z., Y. Tan, C. Huang, and X. Wei. 2023. Redox signaling in drug-tolerant
persister cells as an emerging therapeutic target. *EBioMedicine* 89:104483, doi:
10.1016/j.ebiom.2023.104483.
29. Pope, L.E., and S.J. Dixon. 2023. Regulation of ferroptosis by lipid metabolism.
Trends Cell Biol. 33:1077–1087, doi: 10.1016/J.TCB.2023.05.003.
30. Zheng, X., X. Jin, F. Ye, X. Liu, B. Yu, Z. Li, T. Zhao, W. Chen, X. Liu, C. Di, and
Q. Li. 2023. Ferroptosis: a novel regulated cell death participating in cellular stress
response, radiotherapy, and immunotherapy. *Exp. Hematol. Oncol.* 12:65, doi:
10.1186/S40164-023-00427-W.
31. Becht, E., L. McInnes, J. Healy, C.A. Dutertre, I.W.H. Kwok, L.G. Ng, F. Ginhoux,
and E.W. Newell. 2019. Dimensionality reduction for visualizing single-cell data
using UMAP. *Nat. Biotechnol.* 37:38–47, doi: 10.1038/nbt.4314.
32. Subramanian, A., P. Tamayo, V.K. Mootha, S. Mukherjee, B.L. Ebert, M.A. Gillette,
A. Paulovich, S.L. Pomeroy, T.R. Golub, E.S. Lander, and J.P. Mesirov. 2005.
Gene set enrichment analysis: A knowledge-based approach for interpreting
genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* 102:15545–15550,
doi: 10.1073/pnas.0506580102.

33. Liberzon, A., A. Subramanian, R. Pinchback, H. Thorvaldsdóttir, P. Tamayo, and
J.P. Mesirov. 2011. Molecular signatures database (MSigDB) 3.0. *Bioinformatics*
27:1739–1740, doi: 10.1093/BIOINFORMATICS/BTR260.

34. Liberzon, A., C. Birger, H. Thorvaldsdóttir, M. Ghandi, J.P. Mesirov, and P. Tamayo.
2015. The molecular signatures database hallmark gene set collection. *Cell Syst.*
1:417–425, doi: 10.1016/J.CELS.2015.12.004.

35. Parmenter, T.J., M. Kleinschmidt, K.M. Kinross, S.T. Bond, J. Li, M.R. Kaadige, A.
Rao, K.E. Sheppard, W. Hugo, G.M. Pupo, R.B. Pearson, S.L. McGee, G. V. Long,
R.A. Scolyer, H. Rizos, R.S. Lo, C. Cullinane, D.E. Ayer, A. Ribas, R.W. Johnstone,
R.J. Hicks, and G.A. McArthur. 2014. Response of BRAF-mutant melanoma to
BRAF inhibition is mediated by a network of transcriptional regulators of glycolysis.
Cancer Discov. 4:423–433, doi: 10.1158/2159-8290.CD-13-0440.

36. Hardeman, K.N., C. Peng, B.B. Paudel, C.T. Meyer, T. Luong, D.R. Tyson, J.D.
Young, V. Quaranta, and J.P. Fessel. 2017. Dependence on glycolysis sensitizes
BRAF-mutated melanomas for increased response to targeted BRAF inhibition.
Sci. Rep. 7:42604, doi: 10.1038/srep42604.

37. Shen, S., S. Faouzi, S. Souquere, S. Roy, E. Routier, C. Libenciuc, F. André, G.
Pierron, J.Y. Scoazec, and C. Robert. 2020. Melanoma persister cells are tolerant
to BRAF/MEK inhibitors via ACOX1-mediated fatty acid oxidation. *Cell Rep.*
33:108421, doi: 10.1016/J.CELREP.2020.108421.

38. Tomczak, A., J.M. Mortensen, R. Winnenburg, C. Liu, D.T. Alessi, V. Swamy, F.
Vallania, S. Lofgren, W. Haynes, N.H. Shah, M.A. Musen, and P. Khatri. 2018.
Interpretation of biological experiments changes with evolution of the Gene
Ontology and its annotations. *Sci. Rep.* 8:5115, doi: 10.1038/s41598-018-23395-2.

39. Tirosh, I., B. Izar, S.M. Prakadan, M.H. Wadsworth, D. Treacy, J.J. Trombetta, A.
Rotem, C. Rodman, C. Lian, G. Murphy, M. Fallahi-Sichani, K. Dutton-Regester, J.-
R. Lin, O. Cohen, P. Shah, D. Lu, A.S. Genshaft, T.K. Hughes, C.G.K. Ziegler, S.W.
Kazer, A. Gaillard, K.E. Kolb, A.-C. Villani, C.M. Johannessen, A.Y. Andreev, E.M.
Van Allen, M. Bertagnolli, P.K. Sorger, R.J. Sullivan, K.T. Flaherty, D.T. Frederick,
J. Jané-Valbuena, C.H. Yoon, O. Rozenblatt-Rosen, A.K. Shalek, A. Regev, and
L.A. Garraway. 2016. Dissecting the multicellular ecosystem of metastatic
melanoma by single-cell RNA-seq. *Science (1979).* 352:189–196, doi:
10.1126/science.aad0501.

40. Kim, A., and M.S. Cohen. 2016. The discovery of vemurafenib for the treatment of
BRAF-mutated metastatic melanoma. *Expert Opin. Drug Discov.* 11:907–916, doi:
10.1080/17460441.2016.1201057.

41. Lee, J.T., L. Li, P.A. Brafford, M. Van Den Eijnden, M.B. Halloran, K. Sproesser,
N.K. Haass, K.S.M. Smalley, J. Tsai, G. Bollag, and M. Herlyn. 2010. PLX4032, a
potent inhibitor of the B-Raf V600E oncogene, selectively inhibits V600E-positive
melanomas. *Pigment Cell Melanoma Res.* 23:820–827, doi: 10.1111/J.1755-
148X.2010.00763.X.

42. Joseph, E.W., C.A. Pratilas, P.I. Poulikakos, M. Tadi, W. Wang, B.S. Taylor, E.
Halilovic, Y. Persaud, F. Xing, A. Viale, J. Tsai, P.B. Chapman, G. Bollag, D.B. Solit,
and N. Rosen. 2010. The RAF inhibitor PLX4032 inhibits ERK signaling and tumor

cell proliferation in a V600E BRAF-selective manner. *Proc. Natl. Acad. Sci. U. S. A.*
107:14903–14908, doi: 10.1073/PNAS.1008990107.

43.   Al'Khafaji, A.M., D. Deatherage, and A. Brock. 2018. Control of lineage-specific
gene expression by functionalized gRNA barcodes. *ACS Synth. Biol.* 7:2468–2474,
doi: 10.1021/acssynbio.8b00105.

44.   Gardner, A., D. Morgan, A. Al'Khafaji, and A. Brock. 2022. Functionalized lineage
tracing for the study and manipulation of heterogeneous cell populations. In
Methods in Molecular Biology. Rasooly A, H Baker, MR Ossandon, editors.
Humana Press Inc.: New York, NY, pp. 109–131.

45.   Korolev, K.S., J.B. Xavier, and J. Gore. 2014. Turning ecology and evolution
against cancer. *Nat. Rev. Cancer* 14:371–80, doi: 10.1038/nrc3712.

46.   Coates, J., B.R. Park, D. Le, E. Şimşek, W. Chaudhry, and M. Kim. 2018. Antibiotic-
induced population fluctuations and stochastic clearance of bacteria. *Elife*
7:e32976, doi: 10.7554/eLife.32976.

47.   Wolf, D.M., V. V Vazirani, and A.P. Arkin. 2005. Diversity in times of adversity:
probabilistic strategies in microbial survival games. *J. Theor. Biol.* 234:227–53, doi:
10.1016/j.jtbi.2004.11.020.

48.   Buenrostro, J.D., B. Wu, H.Y. Chang, and W.J. Greenleaf. 2015. ATAC-seq: A
method for assaying chromatin accessibility genome-wide. *Curr. Protoc. Mol. Biol.*
109:21.29.1-21.29.9, doi: 10.1002/0471142727.mb2129s109.

49.   Sakabe, N.J., D. Savic, and M.A. Nobrega. 2012. Transcriptional enhancers in
development and disease. *Genome Biol.* 13:238, doi: 10.1186/gb-2012-13-1-238.

50.   Dixon, S.J., and D.A. Pratt. 2023. Ferroptosis: A flexible constellation of related
biochemical mechanisms. *Mol. Cell* 83:1030–1042, doi:
10.1016/J.MOLCEL.2023.03.005.

51.   Dixon, S.J., and B.R. Stockwell. 2019. The hallmarks of ferroptosis. *Annu. Rev.*
Cancer Biol. 3:35–54, doi: 10.1146/ANNUREV-CANCERBIO-030518-055844.

52.   Dos Santos, A.F., G. Fazeli, T.N. Xavier da Silva, and J.P. Friedmann Angeli. 2023.
Ferroptosis: mechanisms and implications for cancer development and therapy
response. *Trends Cell Biol.* 33:1062–1076, doi: 10.1016/j.tcb.2023.04.005.

53.   Xie, Y., W. Hou, X. Song, Y. Yu, J. Huang, X. Sun, R. Kang, and D. Tang. 2016.
Ferroptosis: process and function. *Cell Death Differ.* 23:369–379, doi:
10.1038/cdd.2015.158.

54.   Groenendyk, J., L.B. Agellon, and M. Michalak. 2021. Calcium signaling and
endoplasmic reticulum stress. *Int. Rev. Cell Mol. Biol.* 363:1–20, doi:
10.1016/BS.IRCMB.2021.03.003.

55.   Pu, Y., L. Li, H. Peng, L. Liu, D. Heymann, C. Robert, F. Vallette, and S. Shen.
2023. Drug-tolerant persister cells in cancer: the cutting edges and future
directions. *Nat. Rev. Clin. Oncol.* 20:799–813, doi: 10.1038/s41571-023-00815-5.

56.   Russo, M., M. Chen, E. Mariella, H. Peng, S.K. Rehman, E. Sancho, A. Sogari, T.S.
Toh, N.Q. Balaban, E. Batlle, R. Bernards, M.J. Garnett, M. Hangauer, E. Leucci,
J.-C. Marine, C.A. O'Brien, Y. Oren, E.E. Patton, C. Robert, S.M. Rosenberg, S.
Shen, and A. Bardelli. 2024. Cancer drug-tolerant persister cells: from biological

questions to clinical opportunities. *Nat. Rev. Cancer* 24:694–717, doi: 10.1038/s41568-024-00737-z.

57. Liu, S., A. Jiang, F. Tang, M. Duan, and B. Li. 2025. Drug-induced tolerant persisters in tumor: mechanism, vulnerability and perspective implication for clinical treatment. *Mol. Cancer* 24:150, doi: 10.1186/s12943-025-02323-9.

58. Hsu, H.-E., M.J. Martin, S.-H. Weng, R.B. Kitata, S. Nagelli, C.-Y. Chang, S. Hess, and Y.-J. Chen. 2025. Phosphoproteomics of osimertinib-tolerant persister cells reveals targetable kinase-substrate signatures. *Mol. Syst. Biol.* 21:1547–1562, doi: 10.1038/s44320-025-00141-1.

59. Su, Y., M.E. Ko, H. Cheng, R. Zhu, M. Xue, J. Wang, J.W. Lee, L. Frankiw, A. Xu, S. Wong, L. Robert, K. Takata, D. Yuan, Y. Lu, S. Huang, A. Ribas, R. Levine, G.P. Nolan, W. Wei, S.K. Plevritis, G. Li, D. Baltimore, and J.R. Heath. 2020. Multi-omic single-cell snapshots reveal multiple independent trajectories to drug tolerance in a melanoma cell line. *Nat. Commun.* 11:2345, doi: 10.1038/s41467-020-15956-9.

60. Liu, Q., C.A. Herring, Q. Sheng, J. Ping, A.J. Simmons, B. Chen, A. Banerjee, W. Li, G. Gu, R.J. Coffey, Y. Shyr, and K.S. Lau. 2018. Quantitative assessment of cell population diversity in single-cell landscapes. *PLoS Biol.* 16:e2006687, doi: 10.1371/JOURNAL.PBIO.2006687.

61. Marusyk, A., and K. Polyak. 2010. Tumor heterogeneity: Causes and consequences. *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer* 1805:105–117, doi: 10.1016/j.bbcan.2009.11.002.

62. Marusyk, A., V. Almendro, and K. Polyak. 2012. Intra-tumour heterogeneity: a looking glass for cancer? *Nat. Rev. Cancer* 12:323–334, doi: 10.1038/nrc3261.

63. Hinohara, K., and K. Polyak. 2019. Intratumoral heterogeneity: more than just mutations. *Trends Cell Biol.* 29:569–579, doi: 10.1016/j.tcb.2019.03.003.

64. Yang, C., C. Tian, T.E. Hoffman, N.K. Jacobsen, and S.L. Spencer. 2021. Melanoma subpopulations that rapidly escape MAPK pathway inhibition incur DNA damage and rely on stress signalling. *Nat. Commun.* 12:1747, doi: 10.1038/s41467-021-21549-x.

65. Huang, S. 2009. Non-genetic heterogeneity of cells in development: more than just noise. *Development* 136:3853–3862, doi: 10.1242/dev.035139.

66. Huang, S., and S. Kauffman. 2013. How to escape the cancer attractor: Rationale and limitations of multi-target drugs. *Semin. Cancer Biol.* 23:270–278, doi: 10.1016/j.semcancer.2013.06.003.

67. Aranda-Anzaldo, A., and M.A.R. Dent. 2018. Landscaping the epigenetic landscape of cancer. *Prog. Biophys. Mol. Biol.* 140:155–174, doi: 10.1016/j.pbiomolbio.2018.06.005.

68. Harris, L.A., S. Beik, P.M.M. Ozawa, L. Jimenez, and A.M. Weaver. 2019. Modeling heterogeneous tumor growth dynamics and cell–cell interactions at single-cell and cell-population resolution. *Curr. Opin. Syst. Biol.* 17:24–34, doi: 10.1016/j.coisb.2019.09.005.

69. Ewels, P., M. Magnusson, S. Lundin, and M. Käller. 2016. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* 32:3047–3048, doi: 10.1093/BIOINFORMATICS/BTW354.

1102   70.   Zheng, G.X.Y., J.M. Terry, P. Belgrader, P. Ryvkin, Z.W. Bent, R. Wilson, S.B.
1103   Ziraldo, T.D. Wheeler, G.P. McDermott, J. Zhu, M.T. Gregory, J. Shuga, L.
1104   Montesclaros, J.G. Underwood, D.A. Masquelier, S.Y. Nishimura, M. Schnall-Levin,
1105   P.W. Wyatt, C.M. Hindson, R. Bharadwaj, A. Wong, K.D. Ness, L.W. Beppu, H.J.
1106   Deeg, C. McFarland, K.R. Loeb, W.J. Valente, N.G. Ericson, E.A. Stevens, J.P.
1107   Radich, T.S. Mikkelsen, B.J. Hindson, and J.H. Bielas. 2017. Massively parallel
1108   digital transcriptional profiling of single cells. *Nat. Commun.* 8:14049, doi:
1109   10.1038/ncomms14049.
1110   71.   Schneider, V.A., T. Graves-Lindsay, K. Howe, N. Bouk, H.C. Chen, P.A. Kitts, T.D.
1111   Murphy, K.D. Pruitt, F. Thibaud-Nissen, D. Albracht, R.S. Fulton, M. Kremitzki, V.
1112   Magrini, C. Markovic, S. McGrath, K.M. Steinberg, K. Auger, W. Chow, J. Collins,
1113   G. Harden, T. Hubbard, S. Pelan, J.T. Simpson, G. Threadgold, J. Torrance, J.M.
1114   Wood, L. Clarke, S. Koren, M. Boitano, P. Peluso, H. Li, C.S. Chin, A.M. Phillippy,
1115   R. Durbin, R.K. Wilson, P. Flicek, E.E. Eichler, and D.M. Church. 2017. Evaluation
1116   of GRCh38 and de novo haploid genome assemblies demonstrates the enduring
1117   quality of the reference assembly. *Genome Res.* 27:849–864, doi:
1118   10.1101/GR.213611.116.
1119   72.   Stuart, T., A. Butler, P. Hoffman, C. Hafemeister, E. Papalexi, W.M. Mauck, Y. Hao,
1120   M. Stoeckius, P. Smibert, and R. Satija. 2019. Comprehensive integration of single-
1121   cell data. *Cell* 177:1888-1902.e21, doi: 10.1016/J.CELL.2019.05.031.
1122   73.   Waltman, L., and N.J. Van Eck. 2013. A smart local moving algorithm for large-
1123   scale modularity-based community detection. *Eur. Phys. J. B* 86:471, doi:
1124   10.1140/EPJB/E2013-40829-0.
1125   74.   DeTomaso, D., M.G. Jones, M. Subramaniam, T. Ashuach, C.J. Ye, and N. Yosef.
1126   2019. Functional interpretation of single cell similarity maps. *Nat. Commun.*
1127   10:4376, doi: 10.1038/s41467-019-12235-0.
1128   75.   Kim, D., J.M. Paggi, C. Park, C. Bennett, and S.L. Salzberg. 2019. Graph-based
1129   genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat.*
1130   *Biotechnol.* 37:907–915, doi: 10.1038/s41587-019-0201-4.
1131   76.   Liao, Y., G.K. Smyth, and W. Shi. 2014. FeatureCounts: An efficient general
1132   purpose program for assigning sequence reads to genomic features. *Bioinformatics*
1133   30:923–930, doi: 10.1093/bioinformatics/btt656.
1134   77.   Love, M.I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change
1135   and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15:550, doi:
1136   10.1186/s13059-014-0550-8.
1137   78.   Carbon, S., A. Ireland, C.J. Mungall, S. Shu, B. Marshall, S. Lewis, J. Lomax, C.
1138   Mungall, B. Hitz, R. Balakrishnan, M. Dolan, V. Wood, E. Hong, and P. Gaudet.
1139   2009. AmiGO: online access to ontology and annotation data. *Bioinformatics*
1140   25:288–289, doi: 10.1093/BIOINFORMATICS/BTN615.
1141   79.   Pico, A.R., T. Kelder, M.P. Van Iersel, K. Hanspers, B.R. Conklin, and C. Evelo.
1142   2008. WikiPathways: pathway editing for the people. *PLoS Biol.* 6:e184, doi:
1143   10.1371/JOURNAL.PBIO.0060184.
1144   80.   Agrawal, A., H. Balcı, K. Hanspers, S.L. Coort, M. Martens, D.N. Slenter, F. Ehrhart,
1145   D. Digles, A. Waagmeester, I. Wassink, T. Abbassi-Daloii, E.N. Lopes, A. Iyer, J.

Millán Acosta, L.G. Willighagen, K. Nishida, A. Riutta, H. Basaric, C.T. Evelo, E.L.
Willighagen, M. Kutmon, and A.R. Pico. 2024. WikiPathways 2024: next generation
pathway database. *Nucleic Acids Res.* 52:D679–D689, doi:
10.1093/NAR/GKAD960.

81. Corces, M.R., A.E. Trevino, E.G. Hamilton, P.G. Greenside, N.A. Sinnott-
Armstrong, S. Vesuna, A.T. Satpathy, A.J. Rubin, K.S. Montine, B. Wu, A. Kathiria,
S.W. Cho, M.R. Mumbach, A.C. Carter, M. Kasowski, L.A. Orloff, V.I. Risca, A.
Kundaje, P.A. Khavari, T.J. Montine, W.J. Greenleaf, and H.Y. Chang. 2017. An
improved ATAC-seq protocol reduces background and enables interrogation of
frozen tissues. *Nat. Methods* 14:959–962, doi: 10.1038/nmeth.4396.

82. Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput
sequencing reads. *EMBnet. J.* 17:10–12, doi: 10.14806/EJ.17.1.200.

83. Li, H. 2013. Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. *arXiv:1303.3997v2*, doi: 10.48550/arXiv.1303.3997.

84. Zhang, Y., T. Liu, C.A. Meyer, J. Eeckhoute, D.S. Johnson, B.E. Bernstein, C.
Nussbaum, R.M. Myers, M. Brown, W. Li, and X.S. Shirley. 2008. Model-based
analysis of ChIP-Seq (MACS). *Genome Biol.* 9:R137, doi: 10.1186/gb-2008-9-9-
r137.

85. Carroll, T.S., Z. Liang, R. Salama, R. Stark, and I. de Santiago. 2014. Impact of
artifact removal on ChIP quality metrics in ChIP-seq and ChIP-exo data. *Front.*
Genet. 5:75, doi: 10.3389/fgene.2014.00075.

86. Ritchie, M.E., B. Phipson, D. Wu, Y. Hu, C.W. Law, W. Shi, and G.K. Smyth. 2015.
limma powers differential expression analyses for RNA-sequencing and microarray
studies. *Nucleic Acids Res.* 43:e47, doi: 10.1093/NAR/GKV007.

87. Wang, Q., M. Li, T. Wu, L. Zhan, L. Li, M. Chen, W. Xie, Z. Xie, E. Hu, S. Xu, and
G. Yu. 2022. Exploring epigenomic datasets by ChIPseeker. *Curr. Protoc.* 2:e585,
doi: 10.1002/CPZ1.585.

88. Yu, G., L.G. Wang, Y. Han, and Q.Y. He. 2012. clusterProfiler: an R package for
comparing biological themes among gene clusters. *OMICS* 16:284–287, doi:
10.1089/OMI.2011.0118.

89. Wu, T., E. Hu, S. Xu, M. Chen, P. Guo, Z. Dai, T. Feng, L. Zhou, W. Tang, L. Zhan,
X. Fu, S. Liu, X. Bo, and G. Yu. 2021. clusterProfiler 4.0: A universal enrichment
tool for interpreting omics data. *The Innovation* 2:100141, doi:
10.1016/j.xinn.2021.100141.

90. Zar, J.H. 2005. Spearman Rank Correlation. In Encyclopedia of Biostatistics. John
Wiley & Sons, Ltd.

91. Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag:
New York.

92. Frick, P.L., B.B. Paudel, D.R. Tyson, and V. Quaranta. 2015. Quantifying
heterogeneity and dynamics of clonal fitness in response to perturbation. *J. Cell.*
Physiol. 230:1403–1412, doi: 10.1002/jcp.24888.

93. Harris, L.A., P.L. Frick, S.P. Garbett, K.N. Hardeman, B.B. Paudel, C.F. Lopez, V.
Quaranta, and D.R. Tyson. 2016. An unbiased metric of antiproliferative drug effect
in vitro. *Nat. Methods* 13:497–500, doi: 10.1038/nmeth.3852.

94. Ritz, C., F. Baty, J.C. Streibig, and D. Gerhard. 2015. Dose-response analysis using
R. *PLoS One* 10:e0146021, doi: 10.1371/journal.pone.0146021.

1192

FIGURE LEGENDS

1194

Figure 1: *Idling DTPs are composed of a heterogeneous collection of phenotypic*
***states.* (A)** Two-dimensional UMAP space of untreated and idling DTP single-cell
transcriptomes. A total of 6410 cells are shown, split ~50/50 across conditions. UTS:
untreated small, UTL: untreated large, IS: idling DTP small, IL: idling DPT large. **(B)**
Principal component analysis (PCA) based on bulk RNA sequencing of three drug-treated
single-cell-derived subclones (SC01, SC07, SC10) at multiple time points (0, 3, 8 days).
All experiments were performed in triplicate. Lines are drawn between the centroids of
the triplicates across the time series. **(C)** Distributions of gene signature scores from the
functional interpretation analysis for cells in the UTS, UTL, and idling DTP clusters
(considered as one cluster in this analysis) in *A*. **(D)** Overlay of cell-cycle state ("fast" =
S-G2-M, "slow" = G1) onto all single cells in the single-cell transcriptomic UMAP space.
Colored contours (blue and red) behind the cells are cell densities calculated from *A*. **(E)**
Relative proportions of cells in each cell-cycle state for the two untreated and two idling
DTP clusters in *D*. Numbers of cells in each cluster (n) are shown above each bar.

1209

Figure 2: *Most clonal lineages survive BRAFi treatment and enter the idling DTP*
***state.* (A)** Relative fractions of the top 24 most abundant barcoded cell lineages in the
untreated and idling DTP populations. Bar heights correspond to the average of three
experimental replicates; lines are the standard deviations. Bars are ordered by
abundance in the untreated condition. Stars (circles) signify example barcodes (plotted in
C) with reduced (increased) abundance after treatment. **(B)** Distribution of (log2) fold-
change from untreated to idling DTP state for barcoded clonal lineages. Means of fold-
changes were compiled into a distribution for all captured lineages (grey) and the top 24
most abundant lineages from *A*. **(C)** Overlay of single cells from four barcoded lineages
(stars and circles from *A*) on the transcriptomic UMAP space from FIG. 1A. Colored
contours (red and blue) reflect cell density for both treatment conditions. **(D)** Proportion
of cells in the fast-dividing (small) idling DTP cluster (see FIG. 1D) for the top 24 most
abundant barcodes. The dashed line indicates the average for all barcodes. **(E)** Pearson
correlation analysis between the percentage of cells in the fast-dividing idling cluster and
the $\log_2$ fold-change in barcode abundance following BRAFi treatment for the top 24 most
abundant barcodes.

1226

Figure 3: *Idling DTPs are characterized by altered ion channel homeostasis and*
***reduced calcium flux.* (A)** Gene ontology (GO) over-representation analysis of
differentially expressed genes from bulk RNA sequencing of untreated and idling DTP
populations. Shown are GO terms with the top 10 largest gene ratios. **(B)** Differential gene
expression of calcium handling genes 3 and 8 days after BRAFi treatment in three single-
cell-derived BRAF-mutant melanoma subclones. **(C)** GO over-enrichment analysis of
genes associated with unique epigenomic peaks in idling DTP populations from ATAC

sequencing. Shown are the top 10 terms with the largest gene ratio. **(D)** Correlation
analysis between Molecular Function (MF)-type GO terms from transcriptomic data (panel
A) and epigenomic data (panel *C*). **(E)** Calcium flux assay of untreated and idling DTP
cells to assess endoplasmic reticulum calcium content/release and store-operated
calcium entry (SOCE) activity. CPA: cyclopiazonic acid. Note that calcium is added to the
culture medium in the presence of CPA and ionomycin is used to control for the number
of cells in the assay (see Materials and Methods).
1241

Figure 4: *Idling DTP populations are susceptible to death by ferroptosis induction.*
(A) Gene expression values (log$_2$ fold-change) at 3 and 8 days post-BRAFi treatment for
three single-cell-derived subclones (SC01, SC07, SC10). Genes are grouped by
ferroptosis-associated process (see FIG. S4). **(B)** Dose-response curves for untreated
and idling DTP populations treated with the ferroptosis inducer RSL3 (*top*) and RSL3
together with ferroptosis inhibitor ferrostatin-1 (Fer-1; *bottom*). Dashed lines correspond
to the half-maximal inhibitory concentration (IC$_{50}$) for each condition, included in units of
*μ*M. DIP: drug-induced proliferation (see Materials and Methods).
1250

Figure 5. *Conceptual framework describing drug-induced phenotypic landscape*
reshaping, DTP heterogeneity, and targeted treatment strategies. **(A)** Illustrative
interpretation of the drug-induced transition of a BRAF-mutant melanoma population to
the idling DTP state. The clusters in the UMAP space correspond to those observed
experimentally (FIG. 1A). Within the framework proposed in this work, the untreated
phenotypic landscape (*red*) comprises multiple, disparate sub-states across which cells
can transition (*solid arrows*). BRAFi treatment modifies the landscape, reducing but not
eliminating phenotypic heterogeneity (*blue*). Cells from any untreated sub-state can
transition into the idling DTP population state (*dashed arrows*) and occupy any of its sub-
states, across which cells can transition (*solid arrows*). Consistent with GO analyses
(FIGS. 3A,C and S1B,C), different phenotypic sub-states within the untreated and idling
DTP populations exhibit distinct functional programs, including metabolic pathways, ion
channel activity, and stress response processes. **(B)** Illustration of multiple levels of DTP
heterogeneity. (*Left*) Consistent with prior reports, multiple distinct DTP states can coexist
within a tumor ("DTP 1" and "DTP 2"), among which cells can transition. (*Right*) Novel to
the framework presented here, each DTP state is a collection of phenotypic sub-states,
with proliferation rates ranging from positive ("Expanding") to negative ("Regressing").
Cells dynamically transition across these sub-states until a steady-state is reached, in
which the population as a whole exhibits a near-zero net proliferation rate. **(C)**
Hypothetical "targeted landscaping" treatment strategy. Based on the framework
presented in this work, sequential therapies are applied to reshape the phenotypic
landscape and reduce the available phenotypic sub-states within the DTP population
state. Ultimately, the goal is to produce a phenotypic landscape composed entirely of
regressing states, resulting in the elimination of all drug-tolerant cells. All panels were
created with BioRender.com.
1276

Figure S1: *Gene ontology analysis of single-cell transcriptomic clusters.* **(A)** Ten single-cell clusters identified by the 'Seurat' software package used in this work (see Materials and Methods). In this UMAP projection, the "untreated small" (UTS) cluster corresponds to Seurat cluster 8, the "untreated large" (UTL) cluster to Seurat clusters 1, 2, 4, and 9, the "idling small" (IS) cluster to Seurat cluster 6, and the "idling large" (IL) cluster to Seurat clusters 0, 3, 5, and 7. **(B)** GO analysis of differentially expressed genes between the UTS and UTL clusters in the UMAP single-cell transcriptomics space. **(C)** Same as *B* but applied to the IS and IL clusters.

Figure S2: *DNA barcode abundances and distributions across untreated and idling DTP states.* **(A)** Heatmap of relative barcode abundances for all experimental replicates of the untreated and idling DTP populations. The heatmap is organized by decreasing barcode abundance in replicate 1 (R1) of the untreated condition. RPM: reads per million. **(B)** Number of unique barcodes in each treatment condition and number of shared barcodes between conditions. Lines correspond to the means of three experimental replicates. A minimum cutoff of 100 counts per million (CPM) was used. Note that the number of unique barcodes is higher in the drug-treated (idling) condition than in the untreated cells. We attribute this to biological noise, i.e., more lineages went extinct, by chance, in the untreated sample than in the idling sample. Overall, we interpret these data as evidence that lineages are not being clonally selected for upon drug treatment. **(C)** Proportional sharing of barcodes among experimental replicates for untreated and idling DTP populations. **(D)** Overlay of single cells from 20 barcoded lineages on the transcriptomic UMAP space from FIG. 1A. Colored contours (red and blue) reflect cell density for both treatment conditions.

Figure S3: *Quality control of bulk ATAC-seq data and correlation analyses of transcriptome- and epigenome-based GO terms.* **(A)** Unique reads versus sequenced reads for the bulk ATAC-seq data used in the epigenome analyses. **(B)** Insert size distribution of aligned reads from ATAC-seq data for untreated and idling DTP cells. Both conditions follow traditional nucleosome patterning. **(C)** Venn diagram of identified ATAC-seq peaks of open chromatin in the untreated and idling DTP populations. **(D)** Peak binding site distribution for untreated, idling DTP, and shared peaks. The x-axis represents the percentage of peaks; colors correspond to distances (kb) from the transcription start site (TSS). **(E)** Alignment of peaks to the TSS allows for the prediction of epigenomic features in untreated and idling DTP populations. **(F)** Correlation analysis between Biological Process (BP)-type GO terms from transcriptomic data (FIG. 3A) and epigenomic data (FIG. 3C). **(G)** Same as *F*, but for Cellular Component (CC)-type GO terms.

Figure S4: *Ferroptosis pathway.* Schematic diagram of the ferroptosis pathway from the WikiPathways database (wikipathways.org/pathways/WP4313.html) (79, 80). Colored boxes group genes associated with the GO terms listed at the bottom. Genes that are shaded are the ones included in the heatmap in FIG. 4A.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5