3 **Title**

4 Monotreme-specific conserved proteins derived from retroviral reverse transcriptase
5

Authors

Koichi Kitao[1], Takayuki Miyazawa[1*], So Nakagawa[2*]

Affiliations

[1]Laboratory of Virus-Host Coevolution, Institute for Frontier Life and Medical Sciences,

Kyoto University, Sakyo-ku, Kyoto 606-8507, Japan

[2]Department of Molecular Life Science, Tokai University School of Medicine, Isehara,

Kanagawa 259-1193, Japan.

*Correspondence to: takavet@infront.kyoto-u.ac.jp (TM) and so@tokai.ac.jp (SN)

Abstract

Endogenous retroviruses (ERVs) have played an essential role in the evolution of

mammals. Many ERV-derived genes are reported in the therians that are involved in the

placental development. However, the contribution of the ERV-derived genes in

monotremes, which are oviparous mammals, remains to be uncovered. Here, we

conducted a comprehensive search for possible ERV-derived genes in platypus and

echidna genomes and identified three reverse transcriptase-like genes, named "*RTOM1*,

2, and *3*." They were found to be clustered in the *GRIP2* intron. Phylogenetic analysis

revealed that *RTOM1*, *2*, and *3* are strongly conserved between these species, and they

were generated by tandem duplications before the divergence of platypus and echidna.

The *RTOM* transcripts were specifically expressed in the testis, suggesting the

physiological importance of *RTOM* genes. This is the first study reporting monotreme-

specific *de novo* gene candidates derived from ERVs, which provides new insights into

the unique evolution of monotremes.

Endogenous retroviruses (ERVs) are remnants of retroviral genomes found in the host

genomes. ERVs are retroviruses that infect the host germline cells and are integrated into

the host genome (Johnson 2019). Young ERVs retain their viral open reading frames

(ORFs), but gradually lose their intact ORFs due to the accumulation of mutations.

However, proteins expressed from ERVs sometimes evolve as functional genes in the host

(Ueda et al. 2020). A typical example is the syncytin genes, ERV-derived fusogenic genes,

which are expressed in the human placenta (Mi et al. 2000; Blond et al. 2000; Blaise et

al. 2003) and are required for mouse placenta formation (Dupressoir et al. 2009;

Dupressoir et al. 2011). Syncytin genes have been independently acquired from different

ERVs in different mammalian lineages, which is a representative example of the

convergent evolution (Imakawa et al. 2015). In addition, other ERV-derived genes have

also been found to be expressed in the placenta. For example, *HEMO* encoding a secreted

envelope protein (Heidmann et al. 2017) as well as *gagV1* and *pre-gagV1* genes (Boso et

al. 2021) are highly expressed in the human placenta. However, it is unknown whether

ERV-derived genes are co-opted in monotremes that are egg-laying mammals.

Comparative studies for the detection of ERV-derived genes have been conducted in

mammalian genomes, including the platypus (Nakagawa and Takahashi 2016; Wang and

Han 2020). For monotremes, however, only the genome sequence of one species, the

platypus, was available (OANA5). and the quality was limited (Warren et al. 2008).

Recently, high-quality monotreme genomes of platypus (mOrnAna1.p.v1) and echidna

(mTacAcu1.pri) were sequenced using long-read sequencing technology (Zhou et al.

2021), making it possible to search for conserved ERV genes in monotremes. Here, we

performed a comparative analysis of the genomes of these two monotremes and found

three ERV-derived genes that are specific to the monotreme lineage.

To comprehensively search for ERV genes in monotremes, we extracted ORFs from the

genomes of platypus and echidna. The amino acid sequences obtained by the virtual

translation of these ORFs were used as queries for the sequence search. We used the

hidden Markov model (HMM) of the retroviral genes in the Gypsy Database 2.0 (GyDB)

(Llorens et al. 2011) as the subject of the sequence search (supplementary table S1). We

identified ORFs similar to *gag*, *pro*, *pol*, and *env* genes (fig. 1A). These ORFs are

presumed to be a mixture of (1) ORFs that are evolutionarily conserved and (2) ORFs of

young transposons that retain their ORFs. To exclude young ERV ORFs, we performed

the clustering analysis based on the amino acid sequence identity. Since young ERVs are

thought to be included in large clusters due to their mutual similarity to each other, we

removed sequences that belonged to large clusters consisting of more than 10 sequences.

Next, using the platypus ORFs as queries, and the echidna ORFs as the subjects, we

conducted a sequence similarity search using BLASTp. We obtained nine ORF pairs with

high amino acid similarity and the same synteny between platypus and echidna

(supplementary table S2). For six pairs among these, we found respective homologs in

the human genome, indicating that they were either ERV genes acquired in the common

mammalian ancestor or host genes with high similarity to ERVs. Indeed, one of the six

genes is *ASPRV1* that is a known ERV-derived protease gene acquired in the common

ancestor of mammals and is responsible for skin maintenance (Matsui et al. 2011). The

remaining three genes were not found in the human genome. They were located tandemly

in the intron of the *GRIP2* gene in the opposite direction (fig. 1B). All three ORFs showed

high similarity to the reverse transcriptase (RT) of spumaretrovirus in GyDB

(supplementary table S3). Therefore, we designated these genes as *RTOM* [RT-like ORF

in Monotreme], and three genes were named as *RTOM1*, *RTOM2*, and *RTOM3* in order

of their location from the 5′ direction (fig. 1B). The *RTOM* coding sequences were

searched in the genomes of 6 mammals, 2 birds, 8 reptilians, and 2 amphibians

(supplementary table S4); however, significant hits were not obtained other than in

platypus and echidna (BLASTn: E-value < 1E-5). Therefore, the *RTOM* genes could be

monotreme-specific and by originated more than 55 million years ago, the divergence

time of platypus and echidna (Zhou et al. 2021).

We found that the gene structures of *RTOM* genes in the platypus genome were annotated

in the RefSeq database (fig. 2A). *RTOM1*, *2*, and *3* genes of platypus contained two

introns in the 5′ UTR, and the entire *RTOM* ORFs are expressed as mRNA excluding a

second splicing variant of *RTOM3* that partially lost its ORF (fig. 2A). In echidna,

RTOM2 and *RTOM3* gene structures were annotated in the RefSeq transcripts; however,

RTOM1 was not annotated. By conducting transcriptome assemblies of RNA-seq data of

echidna tissues (supplementary table S5), we reconstructed the *RTOM1* transcript (fig.

2B; supplementary data S1). As a result, all echidna *RTOM* transcripts have two introns

in the 5′ UTR, which was similar to observations for platypus. According to the alignment

of the six amino acid sequences of platypus and echidna *RTOM* genes, RTOM2 lacks a

region shared by RTOM1 and RTOM3, but the C-terminal region was conserved among

the RTOM proteins without insertion or deletion (fig. 2C). To investigate the tissue-

specific expression of *RTOM* genes, we analyzed the RNA-seq data of platypus and

echidna (supplementary table S5). In platypus, *RTOM1*, *2*, and *3* were all highly expressed

in the testis (fig. 2D). *GRIP2* was expressed not only in the testis but also in the brain,

and its expression level was lower than that of the *RTOM* genes. This suggests that the

RTOM expression was not a result of the *GRIP2* expression. We further investigated the

mapped reads using Interactive Genome Viewer (Thorvaldsdóttir et al. 2013) and found

that *RTOM3* showed a splicing variant with an intron in the coding region, as shown in

the RefSeq transcript (supplementary fig S1). In echidna, we found that all *RTOM*

transcripts were specifically expressed in the testis, similar to platypus. Expression of

GRIP2 in echidna testis was also relatively low, strengthening the idea that the *RTOM*

expression is independent of *GRIP2* expression (fig. 2E). Given the higher expression

level of *RTOM2* in both platypus and echidna, this gene may play a central role of the

RTOM proteins. It is still possible that the relative expression levels of three genes may

change according to tissues and developmental stages that were not examined in this study.

To obtain insights into the viral origin of the *RTOM* genes, we performed a BLASTp

search of the amino acid sequence of platypus RTOM1 against the NCBI virus database.

We found that retrovirus Pol proteins from various distinct lineages, namely

gammaretrovirus, deltaretrovirus, epsilonretrovirus, and spumaretrovirus, are similar to

the RTOM1 proteins (BLASTp: E-value < 1E-20). In all hits, the retroviral Pol proteins

showed high similarity to the latter half of RTOM1 (about 370-607aa). Domain search

against the Pfam database (Mistry et al. 2021) in the HMMER web service (Finn et al.

2011) revealed that the latter half of RTOM1 and RTOM3 contain RT domains (fig. 3A;

supplementary fig. S2). A phylogenetic tree was constructed from the RT regions of the

RTOM proteins and the retroviral Pol proteins (fig. 3B). The RTOM proteins appear to

be more related to class III retroviruses, including spumaviruses or spumavirus-related

MuERV-L (Llorens et al. 2009). The tree topology suggested that RTOM1 emerged at

first, and RTOM2 and RTOM3 were then generated by tandem gene duplications before

the divergence of platypus and echidna (fig. 3C). In the non-RT region of RTOM1

(approximately 1-369aa), no significant hits for retroviruses were obtained (fig. 3A). We

performed a BLASTp search for all non-redundant proteins in the GenBank database for

the non-RT region of RTOM1; however, no similar proteins were found except for

RTOM2 and 3 (E-value < 0.05). Therefore, the non-RT region of the *RTOM* genes dons

not seem to be derived from ERV genes or conserved host genes. Considering the

structural divergence of the non-RT region, such as deletion of RTOM2 and splicing

variant of platypus RTOM3 (fig. 2C), the RT region is a core domain of the RTOM

proteins, and the non-RT region may provide functional modifications specific to each

RTOM protein.

During the 187-million-years history after diverging from monotremes (Zhou et al. 2021),

therians have acquired many ERV genes and evolved their unique features, especially the

placenta (Imakawa and Nakagawa 2017). Our work revealed that monotremes also

domesticated ERV genes. The functional inference of the RTOM proteins is difficult as

co-opted RT genes, such as RTOMs, have not been reported in other vertebrates to the

best of our knowledge. One possibility is that RTOM proteins may function as restrictive

factors against ERVs and retrotransposons. For example, gag-derived *Fv1* (Best et al.

1996) and env-derived *Fv4* (Ikeda and Sugimura 1989) inhibit retroviral infection in mice.

It is possible that the RT domains in the *RTOM* genes compete with retrotransposition as

antagonists. Another possibility is that RTOM proteins are involved in physiological

functions unique to monotremes. In future studies, it would be important to clarify which

cells, viz. germ cells or somatic cells, in testis express the *RTOM* genes. Further studies

pertaining to *RTOM1*, *2*, and *3* in platypus and echidna will expand our understanding of

ERV co-option during the evolution of mammals.

Materials and Methods

Identification of conserved ERV genes

The platypus genome (mOrnAna1.p.v1, GCF_004115215.1) and the echidna genome

(mTacAcu1.pri, GCF_015852505.1) were used for the ERV gene screening (please see

fig. 1). The 240-nt ORF flanked by stop codons were retrieved using the getorf program

in the European Molecular Biology Open Software Suite (Rice et al. 2000). For HMM-

based sequence search, hmmscan was used (Expected threshold: 1E-5) in HMMER3

v3.2.2 (Eddy 2011). ORFs were clustered using CD-HIT v4.8.1 (Li and Godzik 2006)

with 50% amino acid identity. The sequence search for platypus ORFs against echidna

ORFs was conducted using BLASTp v2.10.0+ with an e-value < 1E-50 (Camacho et al.

2009). Hits with a bitscore > 400 were retrieved and checked the synteny was checked

using the UCSC genome browser (https://genome.ucsc.edu/index.html).

Expression analysis

RNA-seq data of platypus (20 samples from 6 tissues) (Marin et al. 2017) and echidna

(11 samples from 7 tissues) (Zhou et al. 2021) were used (supplementary file S5). Low-

quality reads were trimmed and filtered using fastp v0.19.5 with default options (Chen et

al. 2018). The filtered reads were mapped to the each reference genome using HISAT2

v2.1.0 (Pertea et al. 2016). Based on the 11 RNA-seq sequencing data mapped on the

echidna genome, we obtained the echidna *RTOM1* transcript by conducting transcriptome

assembly using Stringtie2 v2.1.6 with "--merge" option (Kovaka et al. 2019). We added

the coordinates of the echidna *RTOM1* transcript (supplementary data S1) to the RefSeq

gene coordinates. We then calculated the expression levels for 20 platypus and 11 echidna

RNA-seq samples using the Stringtie2 program with default options (Kovaka et al. 2019).

Phylogenetic analysis

Representative retroviral Pol amino acid sequences were retrieved from the GyDB

collection

(https://gydb.org/index.php/Alignment?alignment=POL_retroviridae_Biology_Direct_4

_41_2009&format=txt) (Llorens et al. 2009). A multiple alignment was generated using

MAFFT v7.487 (Katoh and Standley 2013), and poorly aligned regions were removed

using trimAl v1.4.rev15 (Capella-Gutiérrez et al. 2009). A phylogenetic tree was

constructed using IQ-TREE2 v2.0.8 (Minh et al. 2020) with 1000 replicates of ultrafast-

bootstrap (Hoang et al. 2018). The tree was visualized using FigTree v1.4.4

(http://tree.bio.ed.ac.uk/software/figtree/).

Acknowledgments

We would like to thank Editage for English language editing. This work was supported

by Grant-in-Aid for JSPS fellows 20J22607 to K.K. and JSPS KAKENHI 20K06775 and

20H03150 to T.M. and S.N. The super-computing resource was partially supported by the

NIG supercomputer at ROIS National Institute of Genetics.

References

Best S, Le Tissier P, Towers G, Stoye JP. 1996. Positional cloning of the mouse
retrovirus restriction gene Fv1. *Nature*. 382:826–829.

Blaise S, de Parseval N, Bénit L, Heidmann T. 2003. Genomewide screening for
fusogenic human endogenous retrovirus envelopes identifies syncytin 2, a gene
conserved on primate evolution. *Proc Natl Acad Sci U S A*. 100:13013–13018.

Blond JL, Lavillette D, Cheynet V, Bouton O, Oriol G, Chapel-Fernandes S, Mandrand
B, Mallet F, Cosset FL. 2000. An envelope glycoprotein of the human
endogenous retrovirus HERV-W is expressed in the human placenta and fuses
cells expressing the type D mammalian retrovirus receptor. *J Virol*. 74:3321–
3329.

Boso G, Fleck K, Carley S, Liu Q, Buckler-White A, Kozak CA. 2021. The oldest co-
opted gag gene of a human endogenous retrovirus shows placenta-specific
expression and is upregulated in diffuse large B-cell lymphomas. *Mol Biol Evol*.
38:5453–5471.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.
2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: A tool for
automated alignment trimming in large-scale phylogenetic analyses.
Bioinformatics 25:1972–1973.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics* 34:i884–i890.

Dupressoir A, Vernochet C, Bawa O, Harper F, Pierron G, Opolon P, Heidmann T.
2009. Syncytin-A knockout mice demonstrate the critical role in placentation of a
fusogenic, endogenous retrovirus-derived, envelope gene. *Proc Natl Acad Sci U
S A.* 106:12127–12132.

Dupressoir A, Vernochet C, Harper F, Guégan J, Dessen P, Pierron G, Heidmann T.
2011. A pair of co-opted retroviral envelope syncytin genes is required for
formation of the two-layered murine placental syncytiotrophoblast. *Proc Natl
Acad Sci U S A* 108:E1164–E1173.

Eddy SR. 2011. Accelerated profile HMM Searches. *PLoS Comput Biol.* 7:e1002195.

Finn RD, Clements J, Eddy SR. 2011. HMMER web server: interactive sequence
similarity searching. *Nucleic Acids Res.* 39:W29–W37.

Heidmann O, Béguin A, Paternina J, Berthier R, Deloger M, Bawa O, Heidmann T.
2017. HEMO, an ancestral endogenous retroviral envelope protein shed in the
blood of pregnant women and expressed in pluripotent stem cells and tumors.
Proc Natl Acad Sci U S A. 114:E6642–E6651.

Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2:
Improving the ultrafast bootstrap approximation. *Mol Biol Evol.* 35:518–522.

Ikeda H, Sugimura H. 1989. Fv-4 resistance gene: a truncated endogenous murine
leukemia virus with ecotropic interference properties. *J. Virol.* 63:5405–5412.

Imakawa K, Nakagawa S. 2017. The phylogeny of placental evolution through dynamic
integrations of retrotransposons. *Prog Mol Biol Transl Sci.* 145:89–109.

Imakawa K, Nakagawa S, Miyazawa T. 2015. Baton pass hypothesis: successive
incorporation of unconserved endogenous retroviral genes for placentation during
mammalian evolution. *Genes Cells.* 20:771–788.

Johnson WE. 2019. Origins and evolutionary consequences of ancient endogenous
retroviruses. *Nat Rev Microbiol.* 17:355–370.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol.* 30:772–780.

Kovaka S, Zimin A V., Pertea GM, Razaghi R, Salzberg SL, Pertea M. 2019.

248   Transcriptome assembly from long-read RNA-seq alignments with StringTie2.
249    *Genome Biol.* 20:278.

250 Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of
251    protein or nucleotide sequences. *Bioinformatics.* 22:1658–1659.

252 Llorens C, Muñoz-Pomer A, Bernad L, Botella H, Moya A. 2009. Network dynamics of
253    eukaryotic LTR retroelements beyond phylogenetic trees. *Biol Direct* 4:41.

254 Llorens C, Futami R, Covelli L, Domínguez-Escribá L, Viu JM, Tamarit D, Aguilar-
255    Rodríguez J, Vicente-Ripolles M, Fuster G, Bernet GP, et al. 2011. The Gypsy
256    Database (GyDB) of mobile genetic elements: release 2.0. *Nucleic Acids Res.*
257    39:D70–D74.

258 Marin R, Cortez D, Lamanna F, Pradeepa MM, Leushkin E, Julien P, Liechti A, Halbert
259    J, Brüning T, Mössinger K, et al. 2017. Convergent origination of a Drosophila-
260    like dosage compensation mechanism in a reptile lineage. *Genome Res.* 27:1974–
261    1987.

262 Matsui T, Miyamoto K, Kubo A, Kawasaki H, Ebihara T, Hata K, Tanahashi S,
263    Ichinose S, Imoto I, Inazawa J, et al. 2011. SASPase regulates stratum corneum
264    hydration through profilaggrin-to-filaggrin processing. *EMBO Mol Med.* 3:320–
265    333.

266 Mi S, Lee X, Li X, Veldman GM, Finnerty H, Racie L, LaVallie E, Tang XY, Edouard
267    P, Howes S, et al. 2000. Syncytin is a captive retroviral envelope protein
268    involved in human placental morphogenesis. *Nature* 403:785–789.

269 Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A,
270    Lanfear R. 2020. IQ-TREE 2: New models and efficient methods for
271    phylogenetic inference in the genomic era. *Mol Biol Evol.* 37:1530–1534.

272 Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL,
273    Tosatto SCE, Paladin L, Raj S, Richardson LJ, et al. 2021. Pfam: The protein
274    families database in 2021. *Nucleic Acids Res.* 49:D412–D419.

275 Nakagawa S, Takahashi MU. 2016. gEVE: a genome-based endogenous viral element
276    database provides comprehensive viral protein-coding sequences in mammalian
277    genomes. *Database.* 2016:1–8.

Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. 2016. Transcript-level expression
analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nat*
Protoc. 11:1650–1667.

Rice P, Longden I, Bleasby A. 2000. EMBOSS: The European Molecular Biology Open
Software Suite. *Trends Genet.* 16:276–277.

Thorvaldsdóttir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer
(IGV): high-performance genomics data visualization and exploration. *Brief*
Bioinform. 14:178–192.

Ueda MT, Kryukov K, Mitsuhashi S, Mitsuhashi H, Imanishi T, Nakagawa S. 2020.
Comprehensive genomic analysis reveals dynamic evolution of endogenous
retroviruses that code for retroviral-like protein domains. *Mob DNA* 11:29.

Wang J, Han GZ. 2020. Frequent retroviral gene co-option during the evolution of
vertebrates. *Mol Biol Evol.* 37:3232–3242.

Warren WC, Hillier LW, Marshall Graves JA, Birney E, Ponting CP, Grützner F, Belov
K, Miller W, Clarke L, Chinwalla AT, et al. 2008. Genome analysis of the
platypus reveals unique signatures of evolution. *Nature* 453:175–183.

Zhou Y, Shearwin-Whyatt L, Li J, Song Z, Hayakawa T, Stevens D, Fenelon JC, Peel
E, Cheng Y, Pajpach F, et al. 2021. Platypus and echidna genomes reveal
mammalian biology and evolution. *Nature* 592:756–762.

Figure legends

FIG. 1. Identification of RTOM1, 2, and 3. (A) Schematic representation of the in silico screening for conserved ERV-derived genes in platypus and echidna. (B) Genomic context of RTOM1, 2, and 3.

FIG. 2. Expression of *RTOM1*, *2*, and *3*. (A) Schematic representation of the RefSeq

transcripts of the *RTOM* genes in platypus. (B) Schematic representation of the

reconstructed *RTOM1* transcript and RefSeq transcripts of the *RTOM2* and *3* genes in

echidna. (C) Multiple alignment of the amino acid sequences of RTOM proteins. The

amino acid sequence of echidna RTOM1 was obtained from the genomic ORF.

"RTOM3A_plasypus" and "RTOM3B_platypus" are protein isoforms derived from

309   "XM_039910408.1" and "XM_029051290.2," respectively. The regions showing

310   similarity to the HMM of spumaretrovirus RT domain in GyDB are indicated in red. (D

311   and E) Tissue-specific expression of *RTOM* genes and *GRIP2* in (D) platypus and (E)

312   echidna. Normalized expression levels are presented as transcript per million (TPM).

FIG. 3. Evolution of RTOM1, 2, and 3. (A) Comparison between platypus RTOM1 and

retroviral Pol protein. Walleye epidermal hyperplasia virus 1 is represented as an example.

A region showing similarity to the Pol protein by BLASTp was designated as "RT region."

A region that did not show similarity to any retroviral genes was designated as "non-RT

region." (B) A phylogenetic tree constructed from the amino acid sequences of RT regions

of the six RTOM proteins and the retroviral Pol proteins in GyDB. The multiple alignment

is available in supplementary data S2. Ultrafast-bootstrap values obtained from 1000

times replication are shown in major branches. (C) Detailed representation of the clade

of the RTOM genes in (B).

Supplementary Materials

Supplementary fig. S1. Screenshots of Interactive Genomic Viewer of RNA-seq reads on

the RTOM genes.

Supplementary fig. S2. Protein domains in RTOM1, 2, and 3.

Supplementary table S1. The HMM profiles in GyDB used in this study.

Supplementary table S2. ERV-like ORFs shared between platypus and echidna

Supplementary table S3. The GyDB HMMs hit to the RTOM genes

Supplementary table S4. Species and genomes used for genes similar to the RTOM genes.

Supplementary table S5. RNA-seq data used in this study.

Supplementary data S1. Nucleotide sequence of the echidna *RTOM1* transcript.

Supplementary data S2. Alignment of representative retroviral *pol* genes and the *RTOM*

genes.

Supplementally FIG. S1. Screenshots of Interactive Genomic Viewer of RNA-seq reads

on the *RTOM* genes. The transcript tracks in blue lines display the coordinates from the

RefSeq GTF files. Thick blue lines indicate the coding sequences. Since there is no

corresponding RefSeq transcript for echidna *RTOM1*, its gene coordinate was manually

added from assembled transcripts in this study (Materials and Methods).

Supplementally FIG. S2. Protein domains in RTOM1, 2, and 3. The domain search was

conducted using hmmscan in HMMER web server with default options

(https://www.ebi.ac.uk/Tools/hmmer/search/hmmscan).

349