Title: In vitro effect of a non-immunosuppressive FKBP ligand, FK1706, on SARS-CoV-2

replication in combination with antivirals

Running Title: FK1706-remdesivir in vitro synergy against SARS-CoV-2

William E. Fitzsimmons[a,b] #, Tracy L. Hartman[c], Michelle Mendenhall[d] , Catherine Z. Chen[e]

a.   University of Illinois at Chicago, Chicago, IL
b.   Tutela Pharmaceuticals Inc, Vernon Hills, IL
c.   ImQuest Biosciences, Frederick, MD;
d.   Institute for Antiviral Research, Utah State University, Logan, UT;
e.   National Center for Advancing Translational Sciences, NIH, Rockville, MD;

#Address correspondence to: William E. Fitzsimmons
wfitzsim@uic.edu
*Present Address: University of Illinois at Chicago, College of Pharmacy
833 S. Wood Street
Chicago, IL 60612

Abstract

FKBP, a naturally occurring ubiquitous intracellular protein, has been proposed as a potential

target for coronavirus replication. A non-immunosuppressive FKBP ligand, FK1706, was studied

in vitro in a Vero cell model to assess potential activity alone and in combination with antivirals

against SARS-CoV-2 replication. When combined with remdesivir, synergistic activity was seen

(summary synergy score 24.7$\pm$9.56). FK1706 warrants in vivo testing as a potential new

combination therapeutic for the treatment of COVID-19 infections.

FKBP is one of the naturally occurring ubiquitous intracellular proteins called immunophilins that

has enzymatic activity as a peptidyl prolyl cis-trans isomerase and is also essential to the

pharmacologic activity of immunosuppressants. The binding of tacrolimus, everolimus, and

sirolimus, to FKBP is necessary but not sufficient to produce immunosuppression (1,2).

Replication of human coronaviruses is dependent on active immunophilin binding and inhibition

of cyclophilins, an intracellular immunophilin, by cyclosporine blocks the replication of CoVs of

all genera tested, including SARS-CoV, human CoV-229E and -NL-63, feline CoV, as well as

avian infectious bronchitis virus (3-6). More recently, the immunophilin FKBP has been

described as one of the potential targets for SARS-CoV-2 (7,8).

Two ligands to FKBP that are not immunosuppressive, FK1706 (9,10) and ElteN378 (11,12)

were studied. These compounds are structurally distinct; both bind to the core structure for

FKBP but do not have intact calcineurin or mTOR binding domains that produce

immunosuppression. Because these drugs target host cells and may work by a unique

mechanism to inhibit coronavirus replication, the additive or synergistic effect with known virus-

targeting antivirals with mechanisms of RNA polymerase inhibition (e.g., remdesivir), viral error

catastrophe or viral lethal mutagenesis (e.g., molnupiravir), or protease inhibition (e.g.,

M128533) were evaluated.

Vero E6 cells were infected with the live SARS-CoV-2 virus (USA-WA1/2020; World Reference

Center for Emerging Viruses and Arboviruses (WRCEVA)) at low MOI (multiplicity of infection)

and multiple rounds of viral replication occurred over the course of the assay. Percent CPE in

compound-treated virus-infected cells were normalized to infected untreated cells as 0% and

uninfected cells as 100% CPE protection. Based on these data, a concentration-response curve

was created. Toxicity was assessed and compared in untreated, uninfected cells compared to

treated cells.

In vitro testing was conducted at two independent laboratories in sequence. The details of the

protocol followed by each laboratory are included in the appendix materials.

FK1706 (Shanghai SIMR Biotechnology Co. LQY20200910), ElteN378 (Glixx Laboratories Inc.

GLXC -20448), remdesivir, molnupiravir, and M128533 were solubilized in DMSO and were

diluted in culture test media to prepare compound concentrations.

Synergy was calculated using SynergyFinder 2.0 software (13). A summary synergy score

greater than 10 was considered synergistic.

The initial results of FK1706 alone and in combination with remdesivir, molnupiravir, and

M128533 are summarized in Table 1.

When combined, FK1706 (11-90 µM) and remdesivir (3 µM) were effective in inhibiting SARS

CoV-2 viral CPE (93-100%, see Appendix Fig A1). FK1706 (2.85-90 µM) and molnupiravir (0.3

µM) inhibited SARS CoV-2 CPE (up to 70% reduction in viral CPE at 90 µM FK1706 with 0.3

µM molnupiravir; see Appendix Fig A2). FK1706 (11-90 µM) and M128533 (1 µg/mL) reduced

SARS CoV-2 CPE (64-100%, see Appendix Fig A3).

Although FK1706 alone did not exhibit inhibitory activity against SARS-CoV-2, when combined

with suboptimal concentrations (less than the EC_{50}) of all three antivirals, increased inhibition

was observed. Additive effects of ElteN378 with either remdesivir or M128533 were also

demonstrated (see Appendix Table A1).

In follow-up confirmatory combination studies, FK1706 at multiple concentrations was tested in

combination with multiple concentrations of remdesivir. When combined, remdesivir and

FK1706 exhibited synergistic activity inhibiting SARS-CoV-2 and shifting the EC50 value of both

compounds when in combination with the other (Figures 1A,B). The summary synergy scores

were 24.7+9.56 by the ZIP (Supplementary Fig A4,A5), 24.8+9.56 by the Bliss and 24.9+9.56

by the HSA models. Scores >10 in all 3 models indicate synergy.

Molnupiravir alone, nor in combination with FK1706, did not demonstrate activity in the follow-up

confirmatory study. There was no evidence of cytotoxicity with FK1706 or remdesivir alone or in

combination (Appendix Figs A6,A7).

The synergistic effects of FK1706 in combination remdesivir were demonstrated in a live SARS-

CoV-2 virus assay measuring the ability of compounds to inhibit viral-induced CPE in Vero E6

host cells in vitro. The CPE reduction assay is a popular and widely used assay format to

screen for antiviral agents because of its ease of use in quantitative high-throughput screening.

The CPE reduction assay indirectly monitors the ability of compounds to inhibit viral replication

and infection through various mechanisms, including direct inhibition of viral entry or enzymatic

processes as well as acting on host pathways that modulate viral replication. This assay was

previously used to screen 8,810 approved and investigational drugs from the National Center for

Advancing Translational Sciences (NCATS) small molecule collections (14). A cytotoxicity

counter-screen was conducted in parallel in host cells without addition of virus and

demonstrated no substantial cytotoxicity of any of the test agents alone or in combination.

Since two chemically distinct FKBP ligands, FK1706 and ElteN378, both demonstrated activity,

it is likely that FKBP is the key target. This target is in the host cells and complements the virus-

targeted antivirals. The combination activity of these FKBP ligands was not limited to a single

virus-targeted mechanism as the three antivirals have distinctly different mechanisms.

Remdesivir (Veklury), currently the only FDA-approved antiviral for COVID-19 infections, is

administered intravenously to patients (15). Molnupiravir has received Emergency Use

Authorization as oral therapy for outpatient COVID-19 infections (16). Although both of these

antivirals have demonstrated clinical efficacy, there is a need for higher response rates and

FK1706 may have utility in both settings. Additionally, these combinations should be active

against variants with mutations in spike protein.

Both live virus assays use Vero E6 as host cells. Vero E6 cells have been shown to have high

drug efflux transporter P-glycoprotein (P-gp) activity, which can reduce cellular concentrations of

test articles, and remdesivir is a known P-gp substrate (17). Therefore, synergy observed in

Vero E6 cells could be due to P-gp inhibition, which enhances the exposure of remdesivir *in*

vitro, and warrants repeating in other cell-based models.

FK1706 has completed all nonclinical safety pharmacology, ADME, and GLP toxicity studies to

support clinical development. Phase 1 healthy volunteer and Phase 2 studies in patients with

neuropathy have been completed (9). This clinical experience would expedite the introduction of

FK1706 into clinical studies of patients infected with SARS-CoV-2.

In conclusion, these data demonstrate that FKBP is a valid target for coronavirus infections in

combination with virus-targeted antivirals such as remdesivir and molnupiravir. FK1706 warrants

testing in an in vivo animal model of SARS-CoV-2 and if promising, rapid introduction into

COVID-19 infection clinical trials.

Acknowledgements:

The authors thank Diane M. Coniglio, Pharm.D., President, Opus Medical Communications for

editorial assistance with the manuscript.

Funding:

This research was funded by Tutela Pharmaceuticals Inc., a 501(c)(3) not-for-profit

pharmaceutical company. William E. Fitzsimmons is the Founder and Chair of Tutela

Pharmaceuticals Inc. This work was supported by the Intramural Research Program of National

Center for Advancing Translational Sciences, Sciences, National Institutes of Health.

Figure Captions:

Figure 1A. Concentration response of FK1706 when combined with remdesivir (RDM).

Figure 1B. Concentration response of remdesivir when combined with FK1706.

References:

1. Fitzsimmons WE. 2012. Tacrolimus. In: Kaplan B., Burckart G.J., Lakkis F.G., eds.

Immunotherapy in Transplantation Principles and Practice. Wiley-Blackwell. Chichester, West

Sussex, UK, pp. 224-240.

2. Tamura K, Fujimura T, Iwasaki K, Sakuma S, Fujitsu T, Nakamura K, Shimomura K, Kuno T,

Tanaka C, Kobayashi M. 1994. Interaction of tacrolimus (FK506) and its metabolites with FKBP

and calcineurin. Biochem Biophys Res Commun. 202:437-43.

3. Carbajo-Lozoya J, Müller MA, Kallies S, Thiel V, Drosten C, von Brunn A. 2012. Replication

of human coronaviruses SARS-CoV HCoV-NL63 and HCoV-229E is inhibited by the drug

FK506. Virus Res. 165: 112-117.

4. de Wilde AH, Falzarano D, Zevenhoven-Dobbe JC, Beugeling C, Fett C, Martellaro C,

Posthuma CC, Feldmann H, Perlman S, Snijder EJ. 2017. Alisporivir inhibits MERS-and SARS-

coronavirus replication in cell culture, but not SARS-coronavirus infection in a mouse model.

Virus Res. 228: 7-13.

5. Pfefferle S, Schöpf J, Kögl M, Friedel CC, Müller MA, Carbajo-Lozoya J, Stellberger T, von

Dall'Armi E, Herzog P, Kallies S, Niemeyer D, Ditt V, Kuri T, Züst R, Pumpor K, Hilgenfeld R,

Schwarz F, Zimmer R, Steffen I, Weber F, Thiel V, Herrler G, Thiel HJ, Schwegmann-Wessels

C, Pöhlmann S, Haas J, Drosten C, von Brunn A. 2011. The SARS-coronavirus-host

interactome:identification of cyclophilins as target for pan-coronavirus inhibitors. PLoS Pathog.

7(10):e1002331.

6. Tanaka Y., Sato Y., Sasaki T. 2013. Suppression of coronavirus replication by cyclophilin

inhibitors. Viruses. 5:1250-60.

7. Gordon DE, Jang GM, Bouhaddou M, Xu J, Obernier K, White KM, O'Meara MJ, Rezelj VV,

Guo JZ, Swaney DL, Tummino TA, Hüttenhain R, Kaake RM, Richards AL, Tutuncuoglu B,

Foussard H, Batra J, Haas K, Modak M, Kim M, Haas P, Polacco BJ, Braberg H, Fabius JM,
Eckhardt M, Soucheray M, Bennett MJ, Cakir M, McGregor MJ, Li Q, Meyer B, Roesch F, Vallet
T, Mac Kain A, Miorin L, Moreno E, Naing ZZC, Zhou Y, Peng S, Shi Y, Zhang Z, Shen W, Kirby
IT, Melnyk JE, Chorba JS, Lou K, Dai SA, Barrio-Hernandez I, Memon D, Hernandez-Armenta
C, Lyu J, Mathy CJP, Perica T, Pilla KB, Ganesan SJ, Saltzberg DJ, Rakesh R, Liu X,
Rosenthal SB, Calviello L, Venkataramanan S, Liboy-Lugo J, Lin Y, Huang XP, Liu Y,
Wankowicz SA, Bohn M, Safari M, Ugur FS, Koh C, Savar NS, Tran QD, Shengjuler D, Fletcher
SJ, O'Neal MC, Cai Y, Chang JCJ, Broadhurst DJ, Klippsten S, Sharp PP, Wenzell NA,
Kuzuoglu-Ozturk D, Wang HY, Trenker R, Young JM, Cavero DA, Hiatt J, Roth TL, Rathore U,
Subramanian A, Noack J, Hubert M, Stroud RM, Frankel AD, Rosenberg OS, Verba KA, Agard
DA, Ott M, Emerman M, Jura N, von Zastrow M, Verdin E, Ashworth A, Schwartz O, d'Enfert C,
Mukherjee S, Jacobson M, Malik HS, Fujimori DG, Ideker T, Craik CS, Floor SN, Fraser JS,
Gross JD, Sali A, Roth BL, Ruggero D, Taunton J, Kortemme T, Beltrao P, Vignuzzi M, García-
Sastre A, Shokat KM, Shoichet BK, Krogan NJ. 2020. A SARS-CoV-2 protein interaction map
reveals targets for drug repurposing. Nature. 583(7816):459-468.

8. Shigdel UK, Lee SJ, Sowa ME, Bowman BR, Robison K, Zhou M, Pua KH, Stiles DT,
Blodgett JAV, Udwary DW, Rajczewski AT, Mann AS, Mostafavi S, Hardy T, Arya S, Weng Z,
Stewart M, Kenyon K, Morgenstern JP, Pan E, Gray DC, Pollock RM, Fry AM, Klausner RD,
Townson SA, Verdine GL. 2020. Genomic discovery of an evolutionarily programmed modality
for small-molecule targeting of an intractable protein surface. Proc Natl Acad Sci USA.
117:17195-17203.

9. Minematsu T, Lee J, Zha J, Moy S, Kowalski D, Hori K, Ishibashi K, Usui T, Kamimura
H.2010. Time-dependent inhibitory effects of
(1R,9S,12S,13R,14S,17R,18E,21S,23S,24R,25S,27R)-1,14-dihydroxy-12-(E)-2-[(1R,3R,4R)-4-
hydroxy-3-methoxycyclohexyl]-1-methylvinyl-23,25-dimethoxy-13,19,21,27-tetramethyl-17-(2-
oxopropyl)-11,28-dioxa-4-azatricyclo[22.3.1.0(4.9)]octacos-18-ene-2,3,10,16-tetrone (FK1706),
a novel nonimmunosuppressive immunophilin ligand, on CYP3A4/5 activity in humans in vivo
and in vitro. Drug Metab Dispos. 38:249-59.

10. Price RD, Yamaji T, Yamamoto H, Higashi Y, Hanaoka K, Yamazaki S, Ishiye M, Aramori I,

Matsuoka N, Mutoh S, Yanagihara T, Gold BG. 2005. FK1706, a novel non-immunosuppressive

immunophilin: neurotrophic activity and mechanism of action. Eur J Pharmacol. 509:11-9.

11. Caminati G, Martina MR, Menichetti S, Procacci P. 2019. Blocking the FKBP12 induced

dendrimeric burst in aberrant aggregation of a-synuclein by using the ElteN378 synthetic

inhibitor. J Enz Inhib Med Chem. 34: 1711-1715.

12. Martina MR, Tenori E, Bizzarri M, Menichetti S, Caminati G, Procacci P. 2013. The precise

chemical-physical nature of the pharmacore in FK506 binding protein inhibition: ElteX, a New

class of nanomolar FKBP12 ligands. J Med Chem. 56:1041-51.

13. Ianevski A., Giri A.K., Aittokallio T. 2020. SynergyFinder 2.0: visual analytics of multi-drug

combination synergies.Nucleic Acids Research.48:W488–W493.

14. Chen CZ, Shinn P, Itkin Z, Eastman RT, Bostwick R, Rasmussen L, Huang R, Shen M, Hu

X, Wilson KM, Brooks BM, Guo H, Zhao T, Klump-Thomas C, Simeonov A, Michael SG, Lo DC,

Hall MD, Zheng W. 2021. Drug repurposing screen for compounds inhibiting the cytopathic

effect of SARS-CoV-2. Front Pharmacol. 11:592737.

15. Rubin D, Chan-Tack K, Farley J, Sherwat A. 2020. FDA approval of remdesivir- a step in

the right direction. N. Engl. J. Med. 383:2598-2600.

16. Jayk Bernal A, Gomes da Silva MM, Musungaie DB, Kovalchuk E, Gonzalez A, Delos

Reyes V, Martín-Quirós A, Caraco Y, Williams-Diaz A, Brown ML, Du J, Pedley A, Assaid C,

Strizki J, Grobler JA, Shamsuddin HH, Tipping R, Wan H, Paschke A, Butterton JR, Johnson

MG, De Anda C; MOVe-OUT Study Group. 2021. Molnupiravir for oral treatment of Covid-19 in

nonhospitalized patients. N Engl J Med. 2021 Dec 16:NEJMoa2116044.

17. Kumar D., Trivedi, N. 2021. Disease-drug and drug-drug interaction in COVID-19: Risk and
assessment. Biomed Pharmacother. 2021.139: 111642.

	90uM FK1706	45uM FK1706	22.5uM FK1706	11.3uM FK1706
EC50	1.789	3.468	18.55	20.76

	6.7uM RDM	3.3uM RDM	1.7uM RDM
EC50	9.926	25.96	137.9

Table 1. Anti-SARS-CoV-2 Cytoprotection Assay Results for FK1706 and antivirals against SARS-CoV-2 (USA-WA1/2020).

Compound or combination	EC$_{50}$ (µM)	TC$_{50}$ (µM)	TI
FK1706	>90	>90	--
FK1706 + Remdesivir (3 µM)	<11.3	>90	>7.96
Remdesivir (3 µM)	>3	>3	--
Remdesivir	3.63	>100	>27.5
FK1706 + M128533 (1 µg/mL)	<11.3	>90	>7.96
M128533 (1 µM)	>1	>1	--
M128533 (µg/mL)	1.53	86.8	56.7
FK1706 + Molnupiravir (0.3 µM)	28.7	>90	>3.14
Molnupiravir (single conc.)	>0.3	>0.3	--