Interference with lipoprotein maturation sensitizes methicillin-resistant *Staphylococcus aureus* to human

group IIA secreted phospholipase A₂ and daptomycin.

Marieke M. Kuijk[1], Yongzheng Wu[2]*, Vincent P. van Hensbergen[3]*, Gizem Shanlitourk[3], Christine Payré[4], Gérard

Lambeau,[4] Jennifer Herrmann[5], Rolf Müller[5], Jos A.G. van Strijp[6], Yvonne Pannekoek[1], Lhousseine Touqui[7, 8], Nina

M. van Sorge[1,9,#]

[1] Medical Microbiology and Infection Prevention, Amsterdam University Medical Centers, location AMC, University

of Amsterdam, Amsterdam, The Netherlands

[2] Unité de Biologie Cellulaire de l'Infection Microbionne, CNRS UMR3691, Institut Pasteur, Paris, France

[3] Medical Microbiology, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands

[4] Université Côte d'Azur, CNRS, Institut de Pharmacologie Moléculaire et Cellulaire, Valbonne Sophia Antipolis,

France

[5] Helmholtz-Institute for Pharmaceutical Research Saarland (HIPS), Helmholtz Center for Infection Research (HZI),

Department of Pharmacy at Saarland University, Saarbrücken, Germany

[6] Medical Microbiology, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands

[7] Mucoviscidose et Bronchopathies Chroniques, Institut Pasteur, Université de Paris, Paris, France

[8] Sorbonne Université, INSERM UMR S 938, Centre de Recherche Saint-Antoine (CRSA), Paris, France

[9] Netherlands Reference Laboratory for Bacterial Meningitis, Amsterdam University Medical Centers, location AMC,

Amsterdam, The Netherlands

* These authors contributed equally

Corresponding author: Prof. dr. N.M. van Sorge, Amsterdam UMC, location AMC, Department of Medical

Microbiology and Infection Prevention, Meibergdreef 9, IWO building IA3.159, 1105 AZ Amsterdam. Email:

n.m.vansorge@amsterdamumc.nl; Phone: +31 (0)20 566 4862.

Running title

29 *Staphylococcus aureus* sensitization to innate and antibiotic killing

Keywords

31 *Staphylococcus aureus*, host defense, human group IIA secreted phospholipase A₂, daptomycin,
32 lipoprotein

Abstract

Methicillin-resistant *Staphylococcus aureus* (MRSA) has been classified as a high priority pathogen by the World Health Organization underlining the high demand for new therapeutics to treat infections. Human group IIA secreted phospholipase A_2 (hGIIA) is among the most potent bactericidal proteins against Gram-positive bacteria, including *S. aureus*. To determine hGIIA-resistance mechanisms of MRSA we screened the Nebraska Transposon Mutant Library using a sublethal concentration of recombinant hGIIA. We identified and confirmed the role of *lspA,* encoding the lipoprotein signal peptidase LspA, as a new hGIIA resistance gene in both *in vitro* assays and an infection model in hGIIA-transgenic mice. Increased susceptibility of the *lspA* mutant was associated with faster and increased cell wall penetration of hGIIA. Moreover, *lspA* deletion also increased susceptibility to daptomycin, a last-resort antibiotic to treat MRSA infections. Exposure of MRSA wild-type to the LspA-specific inhibitors globomycin and myxovirescin A1 induced a *lspA* mutant phenotype with regard to hGIIA and daptomycin killing. Analysis of >26,000 *S. aureus* genomes showed that LspA is highly sequence-conserved, suggesting that LspA inhibition could be applied universally. The role of LspA in hGIIA resistance was not restricted to MRSA since *Streptococcus mutans* and *Enterococcus faecalis* were also more hGIIA-susceptible after *lspA* deletion or LspA inhibition, respectively. Overall, our data suggest that pharmacological blocking of LspA may disarm Gram-positive pathogens, including MRSA, to enhance clearance by innate host defense molecules and clinically-applied antibiotics.

Introduction

Infectious diseases are a significant cause of morbidity and mortality worldwide and are estimated to increase tremendously in the coming decades due to the rise of antimicrobial resistance [1]. The rapid development of antibiotic resistance does not just limit the success of treatment but also of prophylaxis of infections. Methicillin-resistant *Staphylococcus aureus* (MRSA) is a prominent example of a bacterium that has developed rapid antibiotic resistance over the past decades [2, 3]. Indeed, MRSA is ranked as one of the high priority pathogens by the World Health Organization with regard to the need for new therapeutic strategies [4]. While this bacterium is a common member of the human microbiota and asymptomatically colonizes the skin, gut, and nasal cavity, it can cause a wide spectrum of clinical diseases both in the hospital and in the community once *S. aureus* breaches host barriers.

The discovery of new antibiotics is slower than the emergence of new resistance mechanisms of pathogens [5-7]. Antibiotics are classified as substances that are able to kill bacteria (bactericidal) or inhibit their growth (bacteriostatic) [5]. Consequently, antibiotics target molecules or processes in the cell that are either essential or at least critical for the growth of bacteria. An alternative strategy to target bacterial pathogens could include anti-virulence or sensitizing drugs. These drugs may not affect bacterial viability or growth under laboratory conditions, but would affect bacterial fitness or even allow killing of bacteria in the context of specific host immune components, thereby clearing the infection. Indeed, *S. aureus* expresses a wide array of virulence molecules allowing for persistence in different host compartments through interference with a range of immune defense mechanisms and molecules [8].

The human group IIA secreted phospholipase A2 (hGIIA, also known as sPLA$_2$-IIA) is a bactericidal enzyme that represents an important innate host defense molecule [9, 10]. hGIIA is highly cationic and effectively kills Gram-positive bacteria through hydrolysis of bacterial membrane phospholipids [11]. The enzyme is constitutively present at low levels (<5 ng/mL) in the blood circulation and its concentration increases rapidly to levels as high as 1000 ng/mL upon bacterial infection associated with sepsis [12, 13]. hGIIA requires anionic structures in the bacterial cell wall for binding to and penetration of the Gram-positive cell wall [14, 15]. Once at the membrane, hGIIA hydrolyzes membrane phospholipids resulting in bacterial lysis. hGIIA has been implicated in host defense against *S. aureus.* First, blocking hGIIA in acute phase serum results in loss of bactericidal effects against *S. aureus*, whereas addition of hGIIA to normal serum conferred anti-staphylococcal activity [16]. A bactericidal role of hGIIA has also been observed at barrier sites for example in human tears [17]. Second, hGIIA-transgenic (Tg) mice show higher survival rates compared to control littermates that are naturally sPLA$_2$-IIA-deficient, after an experimental lethal dose of *S. aureus* [18, 19]. As a result, *S. aureus* has evolved resistance strategies against hGIIA-mediated killing, which are geared towards changing the overall charge of the

membrane or cell wall. For example, *S. aureus* increases its surface charge by adding D-alanine residues

to teichoic acids through the DltABCD machinery and L-lysine residues to membrane phospholipids

through the activities of the enzyme MprF [14, 20]. The two-component regulatory system GraRS

controls the expression of both *mrpF* and *dltABCD*, thereby controlling *S. aureus* resistance to cationic

antimicrobial peptides and proteins such as hGIIA [21, 22]. Interestingly, the same bacterial genes are

involved in *S. aureus* resistance to daptomycin, the antibiotic of last-resort to treat MRSA infections.

Indeed, increased expression or gain-of-function mutations in *mprF* and *dltABCD* confer daptomycin

non-susceptibility to *S. aureus* [23, 24]. Therefore, insight into hGIIA resistance mechanisms could

provide new clues for the resistance against clinically-important antibiotics.

S. aureus is predicted to express between 50 to 70 lipoproteins, many of unknown function [25,

26]. Some lipoproteins are involved in antibiotic resistance, for example the beta-lactamase BlaZ and

Dsp1 [27-29]. Before lipoproteins are considered mature, they need to be sequentially processed by the

prolipoprotein diacylglyceryl transferase (Lgt) and lipoprotein signal peptidase II (LspA) enzymes. Lgt

anchors prolipoproteins into the cell membrane through diacylglycerol and LspA subsequently

generates the mature lipoprotein by removal of the signal peptide [30]. Both enzymes are conserved in

all bacteria and marked as essential in Gram-negative but not Gram-positive bacteria [30]. Nonetheless,

incorrect processing of lipoproteins changes the immune interaction of *S. aureus*; the deletion of *lgt*

results in hypervirulence, whereas mutation of *lspA* attenuates virulence in a murine systemic infection

model [31]. In addition, two screens, one designed to identify virulence genes and the other to identify

MRSA resistance mechanisms to polymyxin B-mediated killing, identified *lspA* as a resistance

determinant [32, 33].

The mechanisms by which *S. aureus* or MRSA resist hGIIA-mediated killing have never been

studied in a comprehensive unbiased manner. Here, we screened the Nebraska Transposon Mutant

Library (NTML) to identify hGIIA-susceptible mutants [34]. In addition to previously implicated genes

[14, 35], we identified and confirmed that deletion of *lspA*, which we show to be extremely sequence-

conserved, sensitizes *S. aureus* to hGIIA-mediated killing both *in vitro* and *in vivo*. Moreover, LspA

confers resistance to the last-resort antibiotic daptomycin. Both hGIIA- and daptomycin susceptibility

could be induced by treatment of MRSA with the LspA inhibitors globomycin and myxovirescin A1. The

contribution of LspA to hGIIA resistance was not *S. aureus*-specific but was also observed in

Streptococcus mutans (*S. mutans*) and in *Enterococcus faecalis* (*E. faecalis*). In conclusion, we identify

LspA as a possible new therapeutic target to break resistance of *S. aureus* and possibly other Gram-

positive pathogens to both endogenous antimicrobials and antibiotics routinely used in clinic.

Materials and Methods

Materials

Recombinant hGIIA was produced as described previously [36]. HEPES and CaCl$_2$ were purchased from Sigma Aldrich and Merck, respectively. Albumin Bovine Fraction V, pH 7.0 (BSA) was purchased from Serva. SYTOX Nucleic acid stain was purchased from ThermoFisher and DiOC$_2$(3) was obtained at Promokine / Bio-Connect B.V.. All antibiotics (chloramphenicol, erythromycin, daptomycin, gentamicin, and globomycin) were purchased from Sigma Aldrich.

Bacterial culture

The NTML [34] was grown in Tryptic Soy Broth (TSB, Oxoid) supplemented with 5 µg/mL erythromycin. All other *S. aureus* strains and *Enterococcus* species (*E. faecalis* and *E. faecium*) used in this study (Table 1) were grown in Todd-Hewitt Broth (THB, Oxoid) with continuous shaking at 37°C. After overnight culture, strains were sub-cultured to an optical density at 600 nm (OD$_{600}$) of 0.4 (early logarithmic phase; ≈1x10^8 colony-forming units (CFU)/mL). The plasmid complemented strains were grown in THB supplemented with 20 µg/mL chloramphenicol. *S. mutans* was grown statically in Brain Heart Infusion (BHI) at 37 °C with 5% CO$_2$. The following day, sub-cultures were grown to OD$_{600}$ of 0.2 (early logarithmic phase). Plasmid complemented *S. mutans* strains were grown in the presence of 3 µg/mL chloramphenicol. *Escherichia coli* (*E. coli*) strains were grown in Lysogeny broth (LB) medium supplemented with appropriate antibiotics with continuous shaking.

Table 1. Overview of strains and plasmids used in this study.		
Strains/plasmids	Description	Reference
E. coli		
DC10b	Host strain for cloning vectors *S. aureus*	[37]
MC1061	Host strain for cloning vectors *S. mutans*	
S. aureus		
NRS384	Wild-type, USA300-0114, CA-MRSA	NARSA strain collection
NRS384 Δ*lspA*	NRS384 background with a deletion of *lspA*	This study
NRS384 Δ*lspA*+p*lspA*	NRS384 Δ*lspA* background complemented with *lspA*	This study
S. mutans		
UA159	Wild-type, ATCC 700610, serotype c	[38]
UA159 Δ*lspA*	UA159 background with a deletion of *lspA*	This study
UA159 Δ*lspA*+p*lspA*	UA159 Δ*lspA* background complemented with *lspA*	This study
E. faecalis		
V583	Clinical isolate, ATCC 700802	[39]

E. faecium		
U0317	Clinical isolate	[40]
Plasmids		
pKOR1-MCS	Temperature-sensitive shuttle vector for allelic exchange in *S. aureus*	[41]
pDC123	Complementation vector for gene *lspA*	[42]

Screening the NTML for MRSA hGIIA resistance genes

All 1,920 mutants of the NTML were grown overnight in 96-well round bottom plates. After overnight culture, all transposon-mutant cultures were diluted 20 times in TSB supplemented with 5 µg/mL erythromycin and grown to early exponential phase. Cultures were subsequently diluted 20-fold in HEPES solution (20 mM HEPES, 2 mM CaCl$_2$, pH=7.4) and exposed to 1.25 µg/mL recombinant hGIIA. After incubation for 1 hour at 37 °C, 5 µL droplets were plated on TS agar plates. Mutants with visibly reduced number of CFU were identified as putative hGIIA sensitive mutants.

Construction of *lspA* deletion and *lspA* complemented strains

The markerless *lspA* (SAUSA300_1089) deletion mutant (MRSA Δ*lspA*) was generated in *S. aureus* strain USA300 NRS384. The temperature-sensitive and modified pKOR1 plasmid was used as described earlier [41, 43]. A fusion PCR of the upstream region of 1,008 base pairs (bp) and downstream region of 986 bp flanking the *lspA* gene was generated using NRS384 genomic DNA as template. The fusion PCR product was ligated into the pKOR1-MCS plasmid and amplified in *E. coli* DC10b before electroporation into *S. aureus* NRS384. Allelic exchange was performed through temperature shifts and counter selection [43].

To generate a *lspA* (SMU_853) deletion mutant in *S. mutans* strain UA159, the flanking regions (upstream fragment of 635 bp, downstream fragment of 574 bp) were fused with an erythromycin cassette into a single PCR product. For transformation, *S. mutans* was grown in BHI supplemented with heat-inactivated horse serum and the PCR fusion construct was added at 0.5 µg/mL.

Complementation of both *S. aureus* (MRSA Δ*lspA*::p*lspA*) and *S. mutans* strains was performed with pDC123 containing the full length *lspA* (SAUSA300_1089 for *S. aureus* or SMU_853 for *S. mutans*, respectively). Successful transformation was checked with chloramphenicol resistance and colony PCR. An overview of all strains, plasmids and primers used in this study are shown in Tables 1 and 2. All transformants were plated on selective plates containing appropriate antibiotics and successful transformation was checked with PCR and sequencing.

Table 2. Overview of primers used in this study.			
Primers	Orientation	RE site	Sequence
S. aureus NRS384			
lspA up	Forward	KpnI	GCG**GGTACC**GAATGGCTATTATCAACATTTGGC
lspA up	Reverse		<u>GGAAGTATCCTTTAATAAGGC</u>GCATTTTCGTTCCTCCAATCAATC
lspA down	Forward		GATTGATTGGAGGAACGAAAATGC<u>GCCTTATTAAAGGATACTTCC</u>
lspA down	Reverse	EcoRI	GCG**GAATTC**CGTAATTATAGCACGACACAATTATGCATC
Complementation *lspA*	Forward	EcoRI	GCG**GAATTC**CATGGACGATTGATTGGAG
Complementation *lspA*	Reverse	BglII	GCG**AGATCT**CATTACTTAACCTCCTTCTCC
S. mutans UA159			
lspA up	Forward		GCCAGTCAGCACTATGATTTCTTACCGCC
lspA up	Reverse		<u>GTTTTGAGAATATTTTATATTTTTGTTCAT</u>AAGATCTCCTAAGGCTTATAAGTTTC
lspA down	Forward		*AGTTATCTATTATTTAACGGGAGGAAATAA*GTGTTGGTGTAGCACTTC
lspA down	Reverse		GGTCATTTGGCAAGTTGCCGTGTACAAGGG
Erythromycin cassette	Forward		<u>ATGAACAAAAATATAAAATATTCTCAAAACTTTTTAACG</u>
Erythromycin cassette	Reverse		*TTATTTCCTCCCGTTAAATAATAGATAACT*
Complementation *lspA*	Forward	XbaI	GC**TCTAGA**GCCTTAGGAGATCTTATGCG
Complementation *lspA*	Reverse	BamHI	CGC**GGATCC**GCCTTATCCAGACGCACTCCTGC
Underlined and italic bases indicate overlapping sequences to generate fusion construct. Bases in bold indicate either restriction enzyme (RE) sites.			

CFU killing assay

Survival after hGIIA, daptomycin, or gentamicin exposure was determined by quantifying CFU on TH

agar. Early log-phase bacteria (OD$_{600}$ of 0.2 for S*. mutans* or 0.4 for *S. aureus* and *Enterococcus* spp.)

were washed and resuspended in HEPES solution supplemented with 1% BSA (HEPES 1% BSA) and cell

density was adjusted to the original OD$_{600}$. Bacterial suspensions (containing 10^3 CFU of *S. aureus*, 2×10^3

CFU of *S. mutans* or 10^5 CFU of *Enterococcus* spp.) were mixed 1:1 with increasing concentrations of

recombinant hGIIA, daptomycin, or gentamicin in HEPES 1% BSA and incubated for 1 hour at 37°C.

Samples were then serially diluted in phosphate buffered saline (PBS, pH 7) and plated on TH agar plates.

After overnight incubation at 37°C, CFU were counted and bacterial survival was calculated compared

to untreated bacteria. To investigate the effect of the LspA inhibitor globomycin or myxovirescin A1 on

hGIIA- or daptomycin-mediated killing, the compounds were added to wild-type (WT) bacteria during

sub-culturing to early exponential phase at a concentration of 100 µg/mL for globomycin and 10 µg/mL

for myxovirescin A1, which were produced and purified as previously described [44] and dissolved in

DMSO. The maximum concentration of DMSO was 1%, which was also added to other bacterial cultures

as a control.

Scanning Electron Microscopy (SEM)

MRSA WT, MRSA Δ$lspA$, and MRSA Δ$lspA$::p$lspA$ at stationary phase and early exponential phase (OD$_{600}$ 0.4) were washed, fixed, and dehydrated as described previously [45]. Samples were mounted on 12.5 mm specimen stubs (Agar scientific, Stansted, Essex, UK) and coated with 1 nm gold using the Quorum Q150R S sputter coater at 20 mA. Microscopy was performed with a Phenom PRO desktop SEM (Phenom-World BV) operating at an acceleration voltage of 10 kV.

Growth curve

MRSA WT, MRSA Δ$lspA$, and MRSA Δ$lspA$::p$lspA$ were grown overnight and sub-cultured the following day to an OD$_{600}$ of 0.4 in THB supplemented with antibiotics when appropriate. The early exponential phase bacteria were diluted to OD$_{600}$ 0.025 in THB. OD$_{600}$ was measured every 5 minutes over 20 hours (shaking) in a Biotek Synergy H1.

MRSA infection experiment in hGIIA-Tg mice

Tg mice overexpressing hGIIA were from Taconic (Denmark). They were generated by inserting the 6.2 kb full-length of human gene (PLA_2G2A) into the mouse genome and were bred to a sPLA$_2$-IIA naturally-deficient C57BL/6 female mouse that lacks the functional mouse homologue (Pla2g2a) [19, 46]. The animals were housed at Institut Pasteur animal facility accredited by the French Ministry of Agriculture for performing experiments on live rodents. The study on animals was performed in compliance with the French and European regulations on care and protection of laboratory animals (EU Directive 2010/63, French Law 2013-118, February 6th, 2013). The experimental protocol was approved by the Institut Pasteur Ethics Committee and registered under the reference 2014–0014 with the infection protocol 21.185 (AC 0419).

Mice, both males and females (Supplementary Table 1) of 7–9 weeks old, were bred at Institut Pasteur animal facility and infected intra-peritoneally with MRSA WT or the isogenic Δ$lspA$ mutant ($1x10^7$ or $5x10^7$ CFU) suspended in 100 μL PBS. Mortality and weight loss of mice were monitored twice daily up to 5 days after infection.

Surface charge

Bacterial surface charge was determined as previously described [47]. Briefly, early-exponential phase bacteria (OD$_{600}$ = 0.4) were washed twice in 20 mM MOPS buffer (pH 7.0, Sigma-Aldrich) and adjusted to OD$_{600}$ 0.7. Bacteria were concentrated 10 times, of which 200 μL aliquots were added to 0.5 mg/mL cytochrome c (from Saccharomyces cerevisiae, Sigma-Aldrich) in a sterile 96 well round-bottom plate. Suspensions were incubated for 10 minutes at room temperature and subsequently centrifuged at 3,500 rpm for 8 minutes. Supernatant was transferred to a sterile 96 well flat-bottom plate and

217 absorbance was recorded at 530 nm. The percentage of residual cytochrome c was calculated using
218 samples containing MOPS buffer only (100% binding) and samples containing MOPS buffer and
219 cytochrome c (0% binding).

Membrane potential and permeability assays

Changes in hGIIA-dependent membrane potential were determined using the membrane potential
probe $DiOC_2(3)$ (PromoKine) [15, 48]. Bacterial suspensions (OD_{600} of 0.4) were diluted 100 times
(~$1x10^6$ CFU/mL) and incubated with serial dilutions of hGIIA. After incubation at 37˚C, 3 mM $DiOC_2(3)$
was added and incubated at room temperature for 5 minutes in the dark. Changes in green and red
fluorescence emissions were analyzed by flow cytometry. Bacterial staining with the DNA stain SYTOX
Green (Invitrogen) is a measurement for membrane permeabilization and an indication of bacterial cell
death [49]. Serial dilutions of hGIIA in HEPES solutions were added to wells of a sterile flat-bottom 96
well plate. Bacteria were resuspended in HEPES solution containing 1 µM SYTOX green (OD_{600} of 0.4)
and added to hGIIA dilutions in a final volume of 100 µL. Fluorescence over time was recorded using
Optima Fluostar (green fluorescence 520 nm emission and excitation 485 nm) at 37˚C.

PubMLST database analysis of *S. aureus lspA*

The PubMLST database, assessed at https://pubmlst.org/organisms/staphylococcus-aureus [50] was
used to analyze the presence and sequence conservation of *lspA* (SAUR1197) across the *S. aureus*
population. Alignments were made using the locus explorer of the PubMLST database and
nucleotide and amino acid identity was calculated using the NCBI BLAST tool
(https://blast.ncbi.nlm.nih.gov/blast.cgi). *LspA* gene sequences of 26,036 *S. aureus* strains were
downloaded from the database in February 2021. We excluded whole genome sequences for data
analysis that were unlikely to be *S. aureus*, contained > 300 contigs or an N50 contig length shorter than
20,000 bp, contained an internal stop codon rendering a truncated LspA or when *lspA* was located at
the end of a contig.

Statistical analysis

Statistical analysis was performed using GraphPad Prism 9. We used the Student's *t* test and one- and
two-way ANOVA's with Bonferroni statistical hypothesis testing to correct for multiple comparisons. All
values are reported as mean with standard error of the mean of three biological replicates unless
indicated otherwise. A *p* value of < 0.05 was considered statistically significant.

Results

Identification of hGIIA resistance genes in MRSA.

To unravel new hGIIA resistance mechanisms of MRSA, we screened 1,920 individual MRSA mutants of

the Nebraska Transposon Mutant Library (NTML). Exponentially-grown transposon mutants were

exposed to recombinant hGIIA for one hour and subsequently spotted on agar plates for semi-

quantitative assessment of survival (Supplementary Figure 1A). In total, 39 mutants were identified with

potential increased susceptibility to hGIIA-mediated killing (Supplementary Table 2). These hits included

the transposon mutant NE1360 (*mrpF*), which displays an increased positive charge of membrane

phospholipids and was previously linked to hGIIA resistance [14]. Additionally, transposon insertion in

genes encoding the two-component system GraRS and its ABC-transporter VraFG also rendered MRSA

more susceptible to hGIIA. These genes are important for the regulation of the before mentioned *mprF*

and *dltABCD* operon [22], which has a known role in hGIIA resistance [14]. Transposon mutants in

individual genes of the *dltABCD* operon were not identified since these mutants are absent in the NTML

[34].

264   To confirm the phenotype of individual transposon mutants identified in our screen, we

assessed their susceptibility in a quantitative killing assay across a hGIIA concentration range. As

expected, disruption of previously-identified genes *graR*, *graS*, and *mprF* rendered MRSA more

susceptible to hGIIA-mediated killing (Supplementary Figure 1B). In contrast, mutants with transposons

inserted in the genes *esaC*, *srtB*, *ltaA*, and *asp1* were not differently affected by hGIIA (Supplementary

Figure 1B). Interestingly, the *lspA* transposon mutant (NE1757), showed increased susceptibility to hGIIA

(Supplementary Figure 1B). *LspA* is conserved among Gram-positive and Gram-negative bacteria and

encodes the lipoprotein signal peptidase A, an enzyme involved in the lipoprotein maturation pathway

[30, 51].

Deletion of *lspA* attenuates MRSA resistance to hGIIA *in vitro* and virulence in a hGIIA-Tg mouse model.

To verify the contribution of LspA to hGIIA resistance, we constructed a markerless *lspA* deletion mutant

in the MRSA strain NRS384 (MRSA Δ*lspA*) and a plasmid complemented mutant strain (MRSA

Δ*lspA*::p*lspA*). In accordance to results from our NTML screen, MRSA Δ*lspA* was 5 to 10-fold more

susceptible to hGIIA-mediated killing and the phenotype was rescued by complementation with the full

length *lspA* gene (Fig. 1A). Deletion of *lspA* in the MRSA background did not result in morphological

differences as assessed by scanning electron microscopy (Fig. 1B). Moreover, in accordance with

previous literature of other Gram-positive bacteria [52-55], growth of MRSA in bacterial broth was not

affected in the *lspA* deletion mutant (Fig. 1C).

It was previously shown that mutation of *lspA* resulted in attenuated virulence of *S. aureus* but

had no effect on median lethal dose (LD_{50}) values in a mouse infection model [31]. Interestingly, the

mouse strain used in this study was C57BL/6, which lacks a functional mouse $sPLA_2$-IIA homologue due

to a natural frameshift mutation [19]. Although hGIIA-Tg mice, generated in this naturally-deficient

strain background, showed enhanced survival compared to control littermates after infection with WT

S. aureus [18], it has not yet been determined how *lspA* mutation affects *S. aureus* virulence in a mouse

strain with a functional GIIA gene. Therefore, we infected hGIIA-Tg C57BL/6 mice with MRSA WT or its

isogenic mutant Δ*lspA* at 2 different doses ($1x10^7$ or $5x10^7$ CFU/mouse). All mice survived the challenge.

However, as judged by weight loss, mice infected with either $1x10^7$ or $5x10^7$ MRSA WT bacteria showed

significantly more weight loss compared to mice infected with Δ*lspA* bacteria (Fig. 1D). This suggests

that in the presence of a functional hGIIA enzyme, LspA contributes to MRSA virulence in this infection

model.

hGIIA shows faster cell wall penetration and membrane permeabilization in the absence of LspA.

To gain further insights into the underlying mechanisms of hGIIA susceptibility in the absence of LspA,

we assessed the effects of *lspA* deletion on hGIIA binding and cell wall penetration. Since charge-

dependent binding is an important first step in hGIIA's mechanism of action, we determined the surface

charge of the three strains using the cationic compound cytochrome c [47]. Equal binding levels of

cytochrome c was observed for all three strains (Fig. 2A), suggesting that *lspA* does not affect surface

charge. However, we did observe that MRSA Δ*lspA* was not only more sensitive to hGIIA, but that killing

kinetics were also faster for the mutant compared to WT (Fig. 2B). To assess whether hGIIA trafficking

across the cell wall was different, we compared how hGIIA affected membrane depolarization (early

effect of hGIIA activity) and membrane permeabilization (late effect of hGIIA activity). Membrane

depolarization was measured with the fluorescent voltage-sensitive dye $DiOC_2(3)$ that exhibits green

fluorescence (FITC) in all bacterial cells dependent on cell size and red fluorescence (PerCP) dependent

on membrane potential. Deletion of *lspA* resulted in a faster and more extensive membrane

depolarization (Fig. 2C, Supplementary Figure 2). Loss of LspA also caused increased SYTOX intensity, an

indication of membrane permeabilization [15, 47], compared to MRSA WT and complemented strain

starting from 9 min (Fig. 2D).

Interruption of lipoprotein maturation sensitizes MRSA towards daptomycin.

314   The antibiotic daptomycin is clinically important to treat MRSA infections. Interestingly, the mechanism

315   of action of daptomycin displays similarities with hGIIA, since it is dependent on its positive charge and

316   targets the cell membrane [9, 56]. Correspondingly, the identified *S. aureus* resistance genes, i.e.

317   *dltABCD*, *graRS*, and *mprF* overlap for daptomycin and hGIIA [14, 22-24, 57]. We therefore investigated

whether *lspA* deletion affected daptomycin resistance. Indeed, MRSA Δ*lspA* was about 5-fold more susceptible to daptomycin killing, whereas the *lspA* plasmid complemented strain became even more resistant compared to WT (Fig. 3A). As comparison, we assessed whether an intracellular acting antibiotic, gentamicin, was differentially effective in the presence and absence of LspA. Only at one concentration did we observe that loss of *lspA* rendered MRSA more susceptible to gentamicin killing (Fig. 3B), indicating that LpsA has minimal impact on gentamicin-mediated killing.

LspA inhibitors sensitize MRSA towards hGIIA and daptomycin.

The antibiotics globomycin and myxovirescin A1 are directly bactericidal towards Gram-negative bacteria with minimum inhibitory concentration values of 12.5 and 1 µg/mL for *E. coli*, respectively [58, 59]. Interestingly, both compounds are LspA inhibitors [60, 61] and do not kill *S. aureus* growth even at concentrations of 30 µg/mL myxovirescin A1 and >100 µg/mL globomycin [58, 59]. The co-crystal structures of *S. aureus* LspA with both of these inhibitors were recently published [44]. We assessed whether MRSA could also be sensitized to hGIIA and daptomycin through pharmacological inhibition of LspA. To this end, we pre-incubated MRSA WT with either of these compounds during growth to exponential phase and subsequently exposed the bacterial culture to hGIIA or daptomycin. Indeed, pharmacological interference with LspA by either compound rendered MRSA more susceptible to killing by hGIIA and daptomycin compared to untreated bacteria (Fig. 4A, B). This suggests that these compounds may be interesting sensitizing agent in the context of *S. aureus* infections.

LspA is highly sequence-conserved within the *S. aureus* population.

In considering LspA as a drug target, it is important to assess the sequence conservation over bacterial species. LspA contains five conserved domains, including the catalytic residues, across several bacteria [62]. Moreover, LspA amino acid sequence identity is in-between 35% and 95% across 485 different bacterial species [63].

To investigate the presence and sequence conservation of *lspA*, the genomes of 25,243 *S. aureus* isolates were surveyed using PubMLST [50]. These isolates originated from different continents and from a wide variety of hosts as well as human patients and carriers. A *lspA* gene was present in all isolates examined. Only 5 isolates contained a gene with an internal stop codon rendering a truncated LspA. In total, 141 *lspA* alleles were observed.

The majority of the isolates contained *lspA*4 (14,000 isolates, 54%), *lspA*5 (6,000 isolates, 23%) or *lspA*1 (4,000 isolates, 16%). All other *lspA* alleles were at frequencies < 2.5% (Table 3). Interestingly, specific clonal complexes were associated with a single dominant allele (Table 3). Among 141 *lspA* alleles 110 polymorphic positions out of a total gene length of 492 nucleotides were found. These 110 polymorphic sites represented 124 single nucleotide polymorphisms (SNPs). None of the SNPs found in

critical residues were synonymous, emphasizing the high degree of conservation. The most frequently observed SNPs were found at nucleotide positions 331 and 402 (Fig. 5A). Only one of these, at nucleotide position 331, and present in *lspA*1, *lspA*3, *lspA*7 and *lspA*26, results in an amino acid substitution (Ile111Val). A second non-synonymous SNP at nucleotide position 230 is found in *lspA*3, but this allele is present in only 1% of the isolates (Table 3 and Fig. 5B). All other SNPs, as found in the most frequently observed *lspA* alleles among the *S. aureus* population studied are synonymous (Fig. 5B).

Thus, only two amino acid differences are found when comparing the protein sequences encoded by the six most frequently found alleles among a total population of 25,243 isolates analyzed.

Table 3. Distribution of *lspA* among 25,243 *S. aureus* isolates.			
lspA allele	# isolates	Percentage [a]	Dominant in cc [b]
4	14,000	53.7%	1, 8, 15, 22, 97
5	6,000	22.6%	5
1	4,000	15.7%	30
7	500	2.2%	45
26	400	1.7%	93
3	300	1.1%	-
Other	700	2.9%	

[a]Alleles with at least 1% occurrence in isolates are shown

[b]The allele was considered most dominant in the clonal complex (cc) with an occurrence percentage of 92% or higher.

Numbers are rounded off to thousands and to tenths for number of isolates and percentage, respectively.

LspA contribution to hGIIA resistance is not restricted to *S. aureus*.

Streptococcus mutans is a Gram-positive bacterium that resides in the human oral cavity and is the major cause of dental caries [64]. To assess whether LspA-mediated resistance to hGIIA is restricted to *S. aureus* or more widespread, we created a *lspA* deletion mutant in *S. mutans* strain UA159 by replacing the *lspA* gene with an erythromycin cassette. Complementation of this deletion mutant was accomplished by introducing the plasmid pDC123 containing the full *lspA* gene of *S. mutans*. Results from the killing assay revealed that *lspA* deletion renders *S. mutans* more susceptible to hGIIA and complementation fully restored this phenotype (Fig. 6A).

In addition, we tested two clinical isolates of the enterococcal strains *E. faecalis* V583 and *E. faecium* U0317. These species are part of a group that consists of clinically-relevant and antibiotic-resistant pathogens, collectively called ESKAPE pathogens (*Enterococcus* spp*., Staphylococcus aureus, Klebsiella pneumoniae, Acinetobacter baumannii, Pseudomonas aeruginosa* and *Enterobacter* spp.) [65, 66]. Of the Gram-positive *Enterococci*, the species *E. faecalis* and *E. faecium* are most abundant and are

376 responsible for 75% of all enterococcal infections [67]. We observed that *E. faecalis* was 5-fold more

377 sensitive to hGIIA compared to *E. faecium* (Supplementary Figure 3). Pretreating the clinical

378 enterococcal isolates with 10 µg/mL myxovirescin A1 sensitized *E. faecalis*, but not *E. faecium,* to hGIIA

379 killing compared to the untreated bacteria (Fig. 6B, C). Also, higher concentrations of myxovirescin A1

380 (i.e. 50 µg/mL) did not increase hGIIA killing of *E. faecium*.

Discussion

New treatment strategies against MRSA are in high demand due to the rise of antibiotic resistance even against the last-resort antibiotic daptomycin. The current antibiotic arsenal as well as many therapeutic agents under development aim to be directly bactericidal or stop bacterial growth [6]. The drawback of these compounds is the high selective pressure leading to antimicrobial resistance. Non-traditional antibacterial agents, such as anti-virulence drugs, can offer new therapies in the race against antimicrobial resistance by interfering with bacterial strategies that normally allow survival in the context of immune defenses [68]. Such strategies are expected to be less affected by resistance development as there is no direct pressure on survival [69]. Although sensitizing agents still have to prove their clinical use, the concept is appealing. Many of these strategies against *S. aureus* are under active investigation and some are already in preclinical development [70]. For example, inhibition of staphyloxanthin production increased susceptibility to killing in human blood and decreased the virulence of *S. aureus* in mouse infection models [71, 72]. The present work shows that interfering with lipoprotein maturation by inhibition of LspA enhances innate immune killing of MRSA through the modulation of the bactericidal effects of hGIIA. LspA inhibition also enhances daptomycin-mediated killing, which may provide an add-on strategy in antibiotic treatment.

To identify resistance genes against hGIIA in MRSA, we screened the NTML and confirmed increased susceptibility for hits in *graR*, *graS*, and *mprF*. These three genes have previously been linked to cationic antimicrobial resistance [22], and also specifically to hGIIA resistance [14]. We also identified *vraF* and *vraG*, which is also in line with expectations, since these genes encode the ABC-transporter linked to the GraRS two-component system [73]. This confirms that the screen, although semi-quantitative, does allow the identification of hGIIA-susceptible mutants. However, the screen likely lacks sensitivity to provide a comprehensive list of hGIIA-susceptible mutants. This is illustrated by the fact that we did not identify *graX*, the gene encoding GraX, which was shown to be involved in cationic antimicrobial peptide resistance and interacts with the GraRS system [73, 74]. Therefore, additional hGIIA sensitive mutants are likely to be identified using another set-up of the screening assay.

In our unbiased genetic screen, we identified the transposon mutant NE1757 (*lspA*) to be more susceptible to hGIIA-mediated killing. To exclude the possibility that the *lspA* transposon mutant was identified as a result of growth defects or polar effects of the transposon insertion, we constructed a *lspA* deletion strain in the MRSA background NRS384 that was exposed to a hGIIA concentration range and quantified for bacterial survival. With this quantitative killing assay as well as an infection model in hGIIA-Tg mice, we confirmed *lspA* as a novel hGIIA resistance determinant. Additionally, MRSA Δ*lspA* was also more effectively killed by daptomycin compared to WT. This makes LspA an interesting

therapeutic target as its inhibition would simultaneously increase susceptibility to endogenous and
specific clinically-used antibiotics.

Indeed, we provided proof-of-principle that inhibition of LspA by two known pharmacological
inhibitors, globomycin and myxovirescin, renders MRSA more susceptible to hGIIA and daptomycin
killing. A previous study has shown that the *S. aureus* LspA enzyme is inhibited by these compounds, but
has no direct bactericidal effects [44]. This is in line with the observation that deletion of *lspA* does not
affect growth and morphological appearance of MRSA. Hence, selective pressure of this anti-virulence
strategy is likely to be minimal. LspA inhibition as a therapeutic strategy may have other advantages.
For example, the extracellular location of LspA makes it accessible to drug while no LspA analogs are
found in eukaryotic cells, thereby reducing the risk of off-target effects [44, 62, 63]. In addition, we
showed that LspA is highly conserved among *S. aureus* strains with only 1 amino acid substitution in
>96% of the *S. aureus* collection in the PubMLST database (>26,000 isolates at the time of this analysis).
Conserved proteins are less likely to mutate, making them ideal targets as the inhibitor compounds are
longer lasting and more effective [75]. The natural antibiotics globomycin and myxovirescin A1
specifically inhibit LspA and have similar binding sites on LspA, docking to the catalytic dyad and
clustering around 14 conserved residues [44, 60, 61, 63]. Although they have a distinct chemical
structure and biosynthesis, there is a remarkable similarity in their mode of action. This might point
towards a co-evolution that advanced to prevent resistance [44].

LspA processes prolipoproteins that are anchored into the cell membrane by the enzyme Lgt
[30]. The mechanism by which LspA mediates hGIIA and daptomycin resistance is currently not clear.
We explored the possibility that LspA deletion altered surface charge, thereby facilitating hGIIA binding.
However, no difference in binding of the cationic protein cytochrome c was observed, suggesting no
large effects on the net charge. Since hGIIA binding to bacteria is based on electrostatic interactions
[76], we expect that hGIIA binds similar to WT and *lspA* knock-out strains. On the other hand, we did
observe that loss of LspA affected both kinetics and concentration-dependent effects on membrane
depolarization and membrane permeabilization, with Δ*lspA* mutants showing faster disruption after
exposure to hGIIA. Since LspA is a transmembrane protein [62], lack of LspA may change membrane
properties such as membrane fluidity. However, since we observed the same effects in MRSA WT after
pretreatment with globomycin or myxovirescin, which inhibit LspA enzymatic activity, this explanation
is unlikely. Nonetheless, the presence of multiple immature lipoproteins that still carry the signal
peptide may affect membrane characteristics as these prolipoproteins likely accumulate in the
membrane. In some Gram-positive bacteria other putative signal peptidases are present that could take
over the role of LspA [26], but it is not known if this is the case in *S. aureus*. Another explanation could
be that the function of a single lipoprotein is abolished by deletion of *lspA*, resulting in the observed
phenotypes. However, our screen did not identify mutants in individual lipoprotein-encoding genes. In

addition, lipoproteins may retain their function even without proper processing by LspA [77]. Based on

these considerations and observations, we currently favor the hypothesis that differences in membrane

composition due to the presence of the signal peptide are responsible for the observed phenotypes.

We observed that *lspA* deletion affected antibiotic susceptibility, most pronounced for

daptomycin and marginally for gentamicin. In addition, daptomycin susceptibility could also be

conferred by pharmacological inhibition of LspA. These findings suggest that LspA is involved in

daptomycin resistance. However, the role of LspA in daptomycin-resistance is not necessarily

straightforward, since *lspA* was not identified in two previous screens aimed at identifying daptomycin

resistance determinants [78, 79]. The study using the same NTML as we did here [78], only identified a

single daptomycin-susceptible mutant (SAUSA300_1003). This may indicate that the assay set up was

unable to identify all susceptible mutants, since even *mprF*, a well-known daptomycin resistance

determinant [24], was not identified. The second study used methicillin-sensitive *S. aureus* instead of

MRSA to screen for antibiotic susceptibility, including daptomycin [79]. It may well be that strain

background affects the contribution of *lspA* to daptomycin susceptibility. This is illustrated by a recent

comparative transposon sequencing (Tn-seq) screen where only one of five *S. aureus* strains showed

significant changes in *lspA* insertions after daptomycin exposure [80]. This observation may suggest that

despite high protein sequence conservation, therapeutic efficacy of LspA inhibition may be strain-

specific. This should be addressed in future studies when considering anti-virulence strategies.

Earlier *in vivo* experiments performed with a *S. aureus lspA* deletion strain showed that the

mutant was less virulent [31, 32]. Interestingly, these experiments were performed in inbred C57BL/6

mice or outbred CD-1 mice, which carry a natural homozygous or heterozygous inactivating mutation in

the mouse sPLA$_2$-IIA-encoding gene, respectively [19]. Thus, to assess the contribution of *lspA* mutation

to *S. aureus* virulence in an animal with a functional sPLA$_2$-IIA enzyme, we performed a mouse infection

experiment using hGIIA-Tg C57BL/6 mice [46]. These hGIIA-Tg mice have increased resistance to lethal

S. aureus infection compared to control littermates [18]. In this hGIIA-Tg background, mice infected with

MRSA Δ*lspA* did not display weight loss whereas mice infected with MRSA WT showed on average 5 to

10% weight loss depending on the infectious dose. Altogether, we conclude that LspA-dependent

virulence occurs in a hGIIA-dependent and -independent manner as the effects are observed in

naturally-deficient C57BL/6 mice and hGIIA-Tg mice.

The hGIIA susceptibility phenotype was not only observed in *S. aureus*, but also in *S. mutans*

after *lspA* deletion or *E. faecalis* upon LspA inhibition. LspA inhibitors can bind LspA from multiple Gram-

positive bacteria [44, 63], which may broaden the scope of therapeutic application. However, LspA

inhibition does not universally sensitize Gram-positive bacteria to hGIIA killing, since hGIIA killing of *E.*

faecium was not affected by myxovirescin A1 pretreatment. It is possible that myxovirescin could not

reach LspA in sufficient amounts due differences in cell wall architecture between species and strains.

Alternatively, LspA has no role in hGIIA resistance in the *E. faecium* strain, therefore inhibition had no
effect on susceptibility. Similar differences have been observed with regard to daptomycin resistance
mechanisms, where mutations in the LiaFSR system caused a rearrangement of anionic membrane
phospholipids in *E. faecalis* and daptomycin resistance but this was not observed for *E. faecium* [81].
More research is needed to clarify the potential application of LspA inhibitors as therapeutic add on for
different Gram-positive pathogens.

hGIIA is considered as an acute phase protein [82]. It is strongly expressed by innate immune
cells upon infection [10] and rises high levels in blood and organs that could be exploited for the
development of new treatment strategies for MRSA infections. Deletion of *lspA* or its pharmacological
inhibition renders MRSA more susceptible to hGIIA-mediated killing possibly due to altered membrane
properties. Moreover, hGIIA resistance mechanisms overlap partially with daptomycin resistance
mechanisms and indeed interference with LspA enhanced MRSA susceptibility to daptomycin. We only
focused on hGIIA and clinically-relevant antibiotics, but it is possible that LspA inhibition has broader
effects on virulence. We provided proof-of-concept for this potential add-on therapy by demonstrating
that the antibiotics globomycin and myxovirescin A1 sensitizes MRSA for hGIIA-mediated killing,
although strain-specific effects should be investigated. In addition to MRSA, *S. mutans* and *E. faecalis*
were sensitized by pharmacological inhibition of LspA, increasing the impact of LspA as an sensitizing
target. Therefore, interference with lipoprotein maturation through LspA inhibition is a strategy that
warrants further exploration.

Statements

Conflict of interest

The authors have no conflicts of interest to declare.

Funding sources

This work was supported by grants to G.L. from the Centre National de la Recherche Scientifique (CNRS),
the Fondation Jean Valade/Fondation de France (Award FJV_FDF-00112090), the National Research
Agency (grants MNaims (ANR-17-CE17-0012-01), AirMN (ANR-20-CE14-0024-01) and "Investments for
the Future" Laboratory of Excellence SIGNALIFE, a network for innovation on signal transduction
pathways in life sciences (ANR-11-LABX-0028-01 and ANR-15-IDEX-01), and the Fondation de la
Recherche Médicale (DEQ20180339193L). Part of this work was supported by "Fondation Air Liquide"
(Grant: S-CM19006) granted to L.T.. This study was supported by project 91713303 of the Vidi research

Author contributions

M.M.K., Y.W., V.P.v.H., G.S., C.P., and J.H. carried out the experiments. Y.W., C.P., G.L., J.H., R.M., and

L.T. provided essential reagents. M.K. and V.H. took the lead in writing the manuscript. Y.W., G.L., J.H.,

R.M., Y.P., and L.T. revised the manuscript. N.S conceptualized the study and acquired funding. N.M.v.S.,

Y.P., and J.A.G.v.S. supervised the project.

Data availability statement

Data and resources are available upon request from the corresponding author.

References

1. O'Neill, J., *Tackling drug-resistant infections globally: final report and recommendations*. 2016.
2. Enright, M.C., et al., *The evolutionary history of methicillin-resistant Staphylococcus aureus (MRSA).* Proc Natl Acad Sci U S A, 2002. **99**(**11**): p. 7687-92.
3. Lee, A.S., et al., *Methicillin-resistant Staphylococcus aureus.* Nat Rev Dis Primers, 2018. **4**: p. 18033.
4. Tacconelli, E., et al., *Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis.* Lancet Infect Dis, 2018. **18**(3): p. 318-327.
5. Ferri, M., et al., *Antimicrobial resistance: A global emerging threat to public health systems.* Crit Rev Food Sci Nutr, 2017. **57**(13): p. 2857-2876.
6. Theuretzbacher, U., et al., *Critical analysis of antibacterial agents in clinical development.* Nat Rev Microbiol, 2020. **18**(5): p. 286-298.
7. Miethke, M., et al., *Towards the sustainable discovery and development of new antibiotics.* Nature Reviews Chemistry, 2021. **5**(10): p. 726-749.
8. Koymans, K.J., et al., *Staphylococcal Immune Evasion Proteins: Structure, Function, and Host Adaptation.* Curr Top Microbiol Immunol, 2017. **409**: p. 441-489.
9. van Hensbergen, V.P., et al., *Type IIA Secreted Phospholipase A2 in Host Defense against Bacterial Infections.* Trends Immunol, 2020. **41**(4): p. 313-326.
10. Dore, E. and E. Boilard, *Roles of secreted phospholipase A2 group IIA in inflammation and host defense.* Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids, 2019. **1864**(6): p. 789-802.
11. Lambeau, G. and M.H. Gelb, *Biochemistry and physiology of mammalian secreted phospholipases A2.* Annu Rev Biochem, 2008. **77**: p. 495-520.
12. Rintala, E.M. and T.J. Nevalainen, *Group II phospholipase A2 in sera of febrile patients with microbiologically or clinically documented infections.* Clin Infect Dis, 1993. **17**(5): p. 864-70.
13. Nevalainen, T.J., et al., *Time-resolved fluoroimmunoassays of the complete set of secreted phospholipases A2 in human serum.* Biochim Biophys Acta, 2005. **1733**(2-3): p. 210-23.
14. Koprivnjak, T., et al., *Role of charge properties of bacterial envelope in bactericidal action of human group IIA phospholipase A2 against Staphylococcus aureus.* J Biol Chem, 2002. **277**(49): p. 47636-44.
15. van Hensbergen, V.P., et al., *Streptococcal Lancefield polysaccharides are critical cell wall determinants for human Group IIA secreted phospholipase A2 to exert its bactericidal effects.* PLoS Pathog, 2018. **14**(10): p. e1007348.
16. Weinrauch, Y., et al., *Mobilization of potent plasma bactericidal activity during systemic bacterial challenge. Role of group IIA phospholipase A2.* J Clin Invest, 1998. **102**(3): p. 633-8.
17. Qu, X.D. and R.I. Lehrer, *Secretory phospholipase A2 is the principal bactericide for staphylococci and other gram-positive bacteria in human tears.* Infect Immun, 1998. **66**(6): p. 2791-7.
18. Laine, V.J., D.S. Grass, and T.J. Nevalainen, *Protection by Group II Phospholipase A2 against Staphylococcus aureus.* J Immunol, 1999. **162**(12): p. 7402-8.
19. Kennedy, B.P., et al., *A natural disruption of the secretory group II phospholipase A2 gene in inbred mouse strains.* J Biol Chem, 1995. **270**(38): p. 22378-85.
20. Slavetinsky, C.J., et al., *Sensitizing Staphylococcus aureus to antibacterial agents by decoding and blocking the lipid flippase MprF.* Elife, 2022. **11**.
21. Kraus, D., et al., *The GraRS regulatory system controls Staphylococcus aureus susceptibility to antimicrobial host defenses.* BMC Microbiol, 2008. **8**: p. 85.
22. Yang, S.J., et al., *The Staphylococcus aureus two-component regulatory system, GraRS, senses and confers resistance to selected cationic antimicrobial peptides.* Infect Immun, 2012. **80**(1): p. 74-81.
23. Bayer, A.S., et al., *Dysregulation of mprF and dltABCD expression among daptomycin-non-susceptible MRSA clinical isolates.* J Antimicrob Chemother, 2016. **71**(8): p. 2100-4.

24.   Ernst, C.M. and A. Peschel, *MprF-mediated daptomycin resistance.* Int J Med Microbiol, 2019.
309(5): p. 359-363.

25.   Babu, M.M., et al., *A database of bacterial lipoproteins (DOLOP) with functional assignments to*
predicted lipoproteins. J Bacteriol, 2006. **188**(8): p. 2761-73.

26.   Kovacs-Simon, A., R.W. Titball, and S.L. Michell, *Lipoproteins of bacterial pathogens.* Infect
Immun, 2011. **79**(2): p. 548-61.

27.   Zhang, H.Z., et al., *A proteolytic transmembrane signaling pathway and resistance to beta-*
lactams in staphylococci. Science, 2001. **291**(5510): p. 1962-5.

28.   Foster, T.J., *Antibiotic resistance in Staphylococcus aureus. Current status and future prospects.*
FEMS Microbiol Rev, 2017. **41**(3): p. 430-449.

29.   Jousselin, A., et al., *The posttranslocational chaperone lipoprotein PrsA is involved in both*
glycopeptide and oxacillin resistance in Staphylococcus aureus. Antimicrob Agents Chemother,
2012. **56**(7): p. 3629-40.

30.   Nguyen, M.T. and F. Gotz, *Lipoproteins of Gram-Positive Bacteria: Key Players in the Immune*
Response and Virulence. Microbiol Mol Biol Rev, 2016. **80**(3): p. 891-903.

31.   Bubeck Wardenburg, J., W.A. Williams, and D. Missiakas, *Host defenses against Staphylococcus*
aureus infection require recognition of bacterial lipoproteins. Proc Natl Acad Sci U S A, 2006.
103(37): p. 13831-6.

32.   Mei, J.M., et al., *Identification of Staphylococcus aureus virulence genes in a murine model of*
bacteraemia using signature-tagged mutagenesis. Mol Microbiol, 1997. **26**(2): p. 399-407.

33.   Vestergaard, M., et al., *Inhibition of the ATP Synthase Eliminates the Intrinsic Resistance of*
Staphylococcus aureus towards Polymyxins. mBio, 2017. **8**(5).

34.   Fey, P.D., et al., *A genetic resource for rapid and comprehensive phenotype screening of*
nonessential Staphylococcus aureus genes. mBio, 2013. **4**(1): p. e00537-12.

35.   Koprivnjak, T., et al., *Wall teichoic acid deficiency in Staphylococcus aureus confers selective*
resistance to mammalian group IIA phospholipase A(2) and human beta-defensin 3. Infect
Immun, 2008. **76**(5): p. 2169-76.

36.   Ghomashchi, F., et al., *Preparation of the Full Set of Recombinant Mouse- and Human-Secreted*
Phospholipases A2. Methods Enzymol, 2017. **583**: p. 35-69.

37.   Monk, I.R., et al., *Transforming the untransformable: application of direct transformation to*
manipulate genetically Staphylococcus aureus and Staphylococcus epidermidis. mBio, 2012.
3(2).

38.   Ajdic, D., et al., *Genome sequence of Streptococcus mutans UA159, a cariogenic dental*
pathogen. Proc Natl Acad Sci U S A, 2002. **99**(22): p. 14434-9.

39.   Paulsen, I.T., et al., *Role of mobile DNA in the evolution of vancomycin-resistant Enterococcus*
faecalis. Science, 2003. **299**(5615): p. 2071-4.

40.   van Schaik, W., et al., *Pyrosequencing-based comparative genome analysis of the nosocomial*
pathogen Enterococcus faecium and identification of a large transferable pathogenicity island.
BMC Genomics, 2010. **11**(1): p. 239.

41.   Stapels, D.A., et al., *Staphylococcus aureus secretes a unique class of neutrophil serine protease*
inhibitors. Proc Natl Acad Sci U S A, 2014. **111**(36): p. 13187-92.

42.   Chaffin, D.O. and C.E. Rubens, *Blue/white screening of recombinant plasmids in Gram-positive*
bacteria by interruption of alkaline phosphatase gene (phoZ) expression. Gene, 1998. **219**(1-2):
p. 91-9.

43.   Bae, T. and O. Schneewind, *Allelic replacement in Staphylococcus aureus with inducible counter-*
selection. Plasmid, 2006. **55**(1): p. 58-63.

44.   Olatunji, S., et al., *Structures of lipoprotein signal peptidase II from Staphylococcus aureus*
complexed with antibiotics globomycin and myxovirescin. Nat Commun, 2020. **11**(1): p. 140.

45.   van der Beek, S.L., et al., *GacA is essential for Group A Streptococcus and defines a new class of*
monomeric dTDP-4-dehydrorhamnose reductases (RmlD). Mol Microbiol, 2015. **98**(5): p. 946-
62.

46. Grass, D.S., et al., *Expression of human group II PLA2 in transgenic mice results in epidermal*
hyperplasia in the absence of inflammatory infiltrate. J Clin Invest, 1996. **97**(10): p. 2233-41.
47. Carvalho, F., et al., *L-Rhamnosylation of Listeria monocytogenes Wall Teichoic Acids Promotes*
Resistance to Antimicrobial Peptides by Delaying Interaction with the Membrane. PLoS Pathog,
2015. **11**(5): p. e1004919.
48. Shapiro, H.M., *Membrane potential estimation by flow cytometry.* Methods, 2000. **21**(3): p. 271-
9.
49. Saar-Dover, R., et al., *D-alanylation of lipoteichoic acids confers resistance to cationic peptides*
in group B streptococcus by increasing the cell wall density. PLoS Pathog, 2012. **8**(9): p.
e1002891.
50. Jolley, K.A., J.E. Bray, and M.C.J. Maiden, *Open-access bacterial population genomics: BIGSdb*
software, the PubMLST.org website and their applications. Wellcome Open Res, 2018. **3**: p. 124.
51. Braun, V. and K. Hantke, *Lipoproteins: Structure, Function, Biosynthesis.* Subcell Biochem, 2019.
92: p. 39-77.
52. Khandavilli, S., et al., *Maturation of Streptococcus pneumoniae lipoproteins by a type II signal*
peptidase is required for ABC transporter function and full virulence. Mol Microbiol, 2008. **67**(3):
p. 541-57.
53. Weston, B.F., A. Brenot, and M.G. Caparon, *The metal homeostasis protein, Lsp, of*
Streptococcus pyogenes is necessary for acquisition of zinc and virulence. Infect Immun, 2009.
77(7): p. 2840-8.
54. Reglier-Poupet, H., et al., *Maturation of lipoproteins by type II signal peptidase is required for*
phagosomal escape of Listeria monocytogenes. J Biol Chem, 2003. **278**(49): p. 49469-77.
55. de Greeff, A., et al., *Lipoprotein signal peptidase of Streptococcus suis serotype 2.* Microbiology
(Reading), 2003. **149**(Pt 6): p. 1399-1407.
56. Taylor, S.D. and M. Palmer, *The action mechanism of daptomycin.* Bioorg Med Chem, 2016.
24(24): p. 6253-6268.
57. Slavetinsky, C.J., et al., *Sensitizing Staphylococcus aureus to antibacterial host defense by*
decoding and blocking the lipid flippase MprF. bioRxiv, 2021: p. 2020.11.12.379776.
58. Gerth, K., et al., *The myxovirescins, a family of antibiotics from Myxococcus virescens*
(Myxobacterales). J Antibiot (Tokyo), 1982. **35**(11): p. 1454-9.
59. Kiho, T., et al., *Structure-activity relationships of globomycin analogues as antibiotics.* Bioorg
Med Chem, 2004. **12**(2): p. 337-61.
60. Inukai, M., et al., *Mechanism of action of globomycin.* J Antibiot (Tokyo), 1978. **31**(11): p. 1203-
5.
61. Xiao, Y., et al., *Myxobacterium-produced antibiotic TA (myxovirescin) inhibits type II signal*
peptidase. Antimicrob Agents Chemother, 2012. **56**(4): p. 2014-21.
62. Paetzel, M., *Bacterial Signal Peptidases.* Subcell Biochem, 2019. **92**: p. 187-219.
63. Vogeley, L., et al., *Structural basis of lipoprotein signal peptidase II action and inhibition by the*
antibiotic globomycin. Science, 2016. **351**(6275): p. 876-80.
64. Nakano, K. and T. Ooshima, *Serotype classification of Streptococcus mutans and its detection*
outside the oral cavity. Future Microbiol, 2009. **4**(7): p. 891-902.
65. Rice, L.B., *Federal funding for the study of antimicrobial resistance in nosocomial pathogens: no*
ESKAPE. J Infect Dis, 2008. **197**(8): p. 1079-81.
66. Pendleton, J.N., S.P. Gorman, and B.F. Gilmore, *Clinical relevance of the ESKAPE pathogens.*
Expert Rev Anti Infect Ther, 2013. **11**(3): p. 297-308.
67. Weiner-Lastinger, L.M., et al., *Antimicrobial-resistant pathogens associated with adult*
healthcare-associated infections: Summary of data reported to the National Healthcare Safety
Network, 2015-2017. Infect Control Hosp Epidemiol, 2020. **41**(1): p. 1-18.
68. Theuretzbacher, U. and L.J.V. Piddock, *Non-traditional Antibacterial Therapeutic Options and*
Challenges. Cell Host Microbe, 2019. **26**(1): p. 61-72.
69. Dickey, S.W., G.Y.C. Cheung, and M. Otto, *Different drugs for bad bugs: antivirulence strategies*
in the age of antibiotic resistance. Nat Rev Drug Discov, 2017. **16**(7): p. 457-471.

70. Ford, C.A., I.M. Hurford, and J.E. Cassat, *Antivirulence Strategies for the Treatment of*
Staphylococcus aureus Infections: A Mini Review. Front Microbiol, 2020. **11**: p. 632706.
71. Chen, F., et al., *Small-molecule targeting of a diapophytoene desaturase inhibits S. aureus*
virulence. Nat Chem Biol, 2016. **12**(3): p. 174-9.
72. Liu, C.I., et al., *A cholesterol biosynthesis inhibitor blocks Staphylococcus aureus virulence.*
Science, 2008. **319**(5868): p. 1391-4.
73. Falord, M., et al., *GraXSR proteins interact with the VraFG ABC transporter to form a five-*
component system required for cationic antimicrobial peptide sensing and resistance in
Staphylococcus aureus. Antimicrob Agents Chemother, 2012. **56**(2): p. 1047-58.
74. Herbert, S., et al., *Molecular basis of resistance to muramidase and cationic antimicrobial*
peptide activity of lysozyme in staphylococci. PLoS Pathog, 2007. **3**(7): p. e102.
75. Rao, C.V.S., et al., *Antibiotic targeting of the bacterial secretory pathway.* Biochim Biophys Acta,
2014. **1843**(8): p. 1762-83.
76. Weiss, J.P., *Molecular determinants of bacterial sensitivity and resistance to mammalian Group*
IIA phospholipase A2. Biochim Biophys Acta, 2015. **1848**(11 Pt B): p. 3072-7.
77. Shahmirzadi, S.V., M.T. Nguyen, and F. Gotz, *Evaluation of Staphylococcus aureus Lipoproteins:*
Role in Nutritional Acquisition and Pathogenicity. Front Microbiol, 2016. **7**: p. 1404.
78. Vestergaard, M., et al., *Genome-Wide Identification of Antimicrobial Intrinsic Resistance*
Determinants in Staphylococcus aureus. Front Microbiol, 2016. **7**: p. 2018.
79. Rajagopal, M., et al., *Multidrug Intrinsic Resistance Factors in Staphylococcus aureus Identified*
by Profiling Fitness within High-Diversity Transposon Libraries. mBio, 2016. **7**(4).
80. Coe, K.A., et al., *Multi-strain Tn-Seq reveals common daptomycin resistance determinants in*
Staphylococcus aureus. PLoS Pathog, 2019. **15**(11): p. e1007862.
81. Miller, W.R., A.S. Bayer, and C.A. Arias, *Mechanism of Action and Resistance to Daptomycin in*
Staphylococcus aureus and Enterococci. Cold Spring Harb Perspect Med, 2016. **6**(11).
82. Crowl, R.M., et al., *Induction of phospholipase A2 gene expression in human hepatoma cells by*
mediators of the acute phase response. J Biol Chem, 1991. **266**(4): p. 2647-51.
709

Figures

Figure 1. LspA contributes to hGIIA resistance *in vitro* as well as virulence in a hGIIA-Tg mouse model. (A)
Survival of MRSA WT, MRSA Δ*lspA*, and MRSA Δ*lspA*::p*lspA* after exposure to a concentration range of
recombinant hGIIA. (B) Representative scanning electron microscopy (SEM) images of MRSA WT and
MRSA Δ*lspA* in early exponential phase. (C) Growth curves of MRSA WT, MRSA Δ*lspA*, and MRSA
Δ*lspA*::p*lspA*. (D) Relative weight of male and female hGIIA-Tg C57BL/6 mice injected i.p. with either
MRSA WT or MRSA Δ*lspA* (1×10^7 or 5×10^7 CFU). Statistical significance was determined using one- or
two-way ANOVA + Bonferroni's Multiple Comparison Test. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p <$
0.0001. A, D: Data represent mean with standard error of the mean of three biological replicates.

Figure 2. Deletion of *lspA* results in faster hGIIA cell wall penetration and membrane permeabilization. (A) Surface charge of MRSA WT, MRSA Δ*lspA*, and MRSA Δ*lspA*::p*lspA* as determined in a cytochrome c binding assay. (B) Survival of MRSA WT, MRSA Δ*lspA*, and MRSA Δ*lspA*::p*lspA* over time after incubation with 500 ng/mL recombinant hGIIA. (C) Flow cytometric analysis of PerCP-positive cells of MRSA WT, MRSA Δ*lspA*, and MRSA Δ*lspA*::p*lspA* stained with $DiOC_2(3)$ after exposure to a concentration range of recombinant hGIIA. (D) Kinetic analysis of SYTOX intensity for MRSA WT, MRSA Δ*lspA*, and MRSA Δ*lspA*::p*lspA* in the presence of 250 ng/mL recombinant hGIIA. Statistical significance was determined using a one- or two-way ANOVA + Bonferroni's Multiple Comparison Test. ns = not significant, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. Data represent mean with standard error of the mean of three biological replicates.

Figure 3. Impact of LspA on killing by clinically-relevant antibiotics. Survival of MRSA WT, MRSA Δ*lspA*, and MRSA Δ*lspA*::p*lspA* after exposure to (A) daptomycin or (B) gentamicin. Statistical significance was determined between MRSA WT and MRSA Δ*lspA* using a two-way ANOVA + Bonferroni's Multiple Comparison Test. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. Data represent mean with standard error of the mean of three biological replicates.

Figure 4. Globomycin and myxovirescin A1 increase MRSA killing by hGIIA and daptomycin. Survival of MRSA WT, MRSA Δ*lspA*, MRSA WT + 100 µg/mL globomycin, and MRSA WT + 10 µg/mL myxovirescin A1 after subsequent exposure to (A) recombinant hGIIA (250 ng/mL) or (B) daptomycin (1 µg/mL). Statistical significance was determined using a one-way ANOVA + Bonferroni's Multiple Comparison Test. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$. Data represent mean with standard error of the mean of three biological replicates.

Figure 5. *lspA* and the encoded LspA protein are highly sequence conserved across the *S. aureus* population. (A) Polymorphic site frequencies of 141 alleles of *lspA* among 25,243 *S. aureus* genomes. Consensus sequence is depicted with color coding for the occurrence in percentages. (B) Alignment and consensus sequence at amino acids level encoded by the 6 most common *lspA* alleles. The five conserved domains across all bacterial species are depicted below with roman numerals [62]. *LspA*4 is the reference allele.

Figure 6. *S. mutans* and *E. faecalis*, but not *E. faecium*, are sensitized to hGIIA via *lspA* deletion or LspA inhibition. (A) Survival of *S. mutans* UA159 WT, *S. mutans* Δ*lspA*, and *S. mutans* Δ*lspA*::p*lspA* after exposure to concentration range of recombinant hGIIA. (B) Survival of *E. faecalis* and *E. faecalis* + 10 µg/mL myxovirescin A1 after subsequent exposure to recombinant hGIIA (0.5 ng/mL). (C) Survival of *E. faecium* and *E. faecium* + 10 µg/mL myxovirescin A1 after subsequent exposure to recombinant hGIIA (0.5 ng/mL). Statistical significance was determined using a one- or two-way ANOVA + Bonferroni's Multiple Comparison Test or an unpaired two-tailed Student's *t* test. ns = not significant, *$p < 0.05$, **$p < 0.01$, ****$p < 0.0001$. Data represent mean with standard error of the mean of three biological replicates.

757 **Supplemental Figures**

759 **Supplementary Table 1.** The four different groups including number and sex of mice used in the MRSA

760 infection experiment using hGIIA-Tg C57BL/6 mice.

Bacteria	Infectious dose	No. of animals (male)
MRSA WT	1×10^7 CFU	8 (3)
MRSA $\Delta lspA$	1×10^7 CFU	9 (4)
MRSA WT	5×10^7 CFU	7 (2)
MRSA $\Delta lspA$	5×10^7 CFU	9 (4)

762   **Supplementary Table 2.** The 39 MRSA transposon mutants that showed decreased survival on plate

763   after hGIIA exposure in a non-biased genetic screen.

Strain name	Gene	Function	Accession number
NE43	*ureF*	urease accessory protein	SAUSA300_2242
NE45	*asp1*	accessory secretory protein	SAUSA300_2587
NE70	*vraG*	ABC transporter of GraRS	SAUSA300_0648
NE77	-	conserved hypothetical protein	SAUSA300_0465
NE150	-	conserved hypothetical protein	SAUSA300_0097
NE229	*fakA*	conserved hypothetical protein	SAUSA300_1119
NE235	-	conserved hypothetical protein	SAUSA300_0847
NE256	-	putative pyridoxal phosphate-dependent acyltransferase	SAUSA300_0535
NE257	-	conserved hypothetical protein	SAUSA300_1623
NE259	*pnpA*	polyribopolyribonucleotide nucleotidyltransferase	SAUSA300_1167
NE264	*lipA*	lipoic acid synthetase	SAUSA300_0829
NE340	*ilvA*	threonine dehydratase	SAUSA300_2014
NE352	*rsgA*	ribosome small subunit-dependent GTPase A	SAUSA300_1114
NE428	*moaE*	molybdopterin converting factor	SAUSA300_2222
NE462	*ltaA*	proton-coupled antiporter flippase	SAUSA300_0917
NE481	*graR*	part of two component system GraRS	SAUSA300_0645
NE592	*atpA*	ATP synthase F1, alpha subunit	SAUSA300_2060
NE603	*moeA*	molybdopterin biosynthesis protein A	SAUSA300_2224
NE604	*esaC*	protein within ESAT-6 gene cluster	SAUSA300_0284
NE605	*brnQ2*	branched-chain amino acid transport system II carrier protein	SAUSA300_0306
NE630	-	acetyltransferase, GNAT family	SAUSA300_0665
NE631	-	phiPVL ORF39-like protein	SAUSA300_1962
NE645	*vraF*	ABC transporter of GraRS	SAUSA300_0647
NE788	*trkA*	potassium uptake protein	SAUSA300_0988
NE883	*xerC*	tyrosine recombinase xerC	SAUSA300_1145
NE885	-	pyruvate ferredoxin oxidoreductase, alpha subunit	SAUSA300_1182
NE891	-	sodium transport family protein	SAUSA300_0924
NE1193	*sarA*	accessory regulator A	SAUSA300_0605
NE1334	-	hypothetical protein	SAUSA300_1494
NE1360	*mprF*	phosphatidylglycerol lysyltransferase	SAUSA300_1255
NE1363	*srtB*	sortase B	SAUSA300_1034
NE1371	-	conserved hypothetical phage protein	SAUSA300_1967
NE1504	-	Na+/H+ antiporter	SAUSA300_0617
NE1531	*pdxT*	glutamine amidotransferase subunit	SAUSA300_0505
NE1536	*gcvH*	glycine cleavage system protein H	SAUSA300_0791
NE1756	*graS*	part of two component system GraRS	SAUSA300_0646
NE1757	*lspA*	lipoprotein signal peptidase	SAUSA300_1089
NE1828	*pdxS*	pyridoxal biosynthesis lyase	SAUSA300_0504
NE1894	-	hypothetical membrane protein	SAUSA300_1908

Supplementary Figure 1. Identification of MRSA transposon mutants with increased susceptibility to

hGIIA-mediated killing. (A) Representative image of an agar plate with spotted MRSA transposon

mutants after exposure to 1.25 μg/mL recombinant hGIIA. Two transposon mutants on this plate

showed decreased viability; at position C4 is NE0646 (*graS*) and at position C5 is the transposon mutant

NE1757 (*lspA*). (B) Survival of potentially hGIIA susceptible mutants from the NTML using quantitative

concentration-dependent killing assay. Data represent mean with standard error of the mean from

three technical replicates.

Supplementary Figure 2. Faster membrane depolarization in *lspA* deletion mutant compared to WT and

complemented strain after hGIIA exposure. Flow cytometric analysis of PerCP-positive cells of MRSA WT,

MRSA Δ*lspA*, and MRSA Δ*lspA*::p*lspA* stained with DiOC$_2$(3) at different time point after exposure to 500

ng/mL hGIIA. Data represent mean ± standard error of the mean of three independent experiments.

Statistical significance was determined using a two-way ANOVA + Bonferroni's Multiple Comparison

Test. *p < 0.05. Data represent mean with standard error of the mean of three biological replicates.

Supplementary Figure 3. *E. faecalis* is more sensitive to hGIIA killing compared to *E. faecium*. Survival of

E. faecalis V583 and *E. faecium* U0317 after exposure to concentration range of recombinant hGIIA.

Data represent mean with standard error of the mean of three biological replicates.