Inflammatory monocytes promote pyogranuloma formation to counteract

Yersinia **blockade of host defense**

Daniel Sorobetea[1,2], Rina Matsuda[1,2], Stefan T. Peterson[1], James P. Grayczyk[1],

Indira Rao[1], Elise Krespan[1], Matthew Lanza[1], Charles-Antoine Assenmacher[1],

Daniel Beiting[1], Enrico Radaelli[1] & Igor Brodsky[1]*

*e-mail: ibrodsky@vet.upenn.edu

Affiliations

1. Department of Pathobiology, School of Veterinary Medicine, University of

Pennsylvania, Philadelphia, USA

2. These authors contributed equally

Abstract

Granulomas are organized immune cell aggregates that form in response to chronic

infection or antigen persistence. *Yersinia pseudotuberculosis* (*Yp*) blocks innate

inflammatory signaling and phagocytosis, inducing formation of neutrophil-rich

pyogranulomas within lymphoid tissues. Here, we uncover that *Yp* triggers

pyogranuloma formation within the murine intestinal mucosa, a site not known to

contain such structures. Mice lacking circulating monocytes fail to form defined

pyogranulomas, have defects in neutrophil activation, and succumb to *Yp* infection.

Yersinia lacking the virulence factors that block phagocytosis did not induce

pyogranulomas, indicating that intestinal pyogranulomas form in response to *Yp*

disruption of phagocytosis. Notably, mutation of a single anti-phagocytic virulence

factor, YopH, restored pyogranuloma formation and control of *Yp* infection in

monocyte-deficient mice, demonstrating that monocytes override YopH-dependent

blockade of innate immune defense. This work reveals an unappreciated site of

Yersinia intestinal invasion, and defines host and pathogen drivers of intestinal

granuloma formation.

Introduction

Microbial pathogens utilize diverse mechanisms to subvert host immunity in order to replicate and spread to new hosts. While acute infections can be cleared rapidly by the immune system, some pathogens evade immune defenses to cause chronic disease. Inability to clear an infection often results in an alternative strategy of containment within structures termed granulomas to limit pathogen dissemination and tissue damage[1]. Granulomas are characterized by the presence of activated phagocytes, notably monocytes and macrophages, and form in response to a wide variety of infectious organisms[2]. Monocytes are rapidly recruited to infected tissues where they produce inflammatory cytokines and anti-microbial effector molecules, contributing to clearance of multiple pathogens[3–8]. Some pathogens, however, exploit monocytes as a means of dissemination, including *Salmonella enterica*, *Yersinia pestis,* and *Mycobacterium* species[9–12]. The pathogen-specific signals that induce granuloma formation remain poorly defined, raising questions about the precise bacterial and host factors that mediate granuloma formation.

The enteropathogenic *Yersinia, Y. pseudotuberculosis* (*Yp*) and *Y. enterocolitica* (*Ye*), cause self-limiting gastroenteritis and mesenteric lymphadenopathy following enteric infection[13,14]. In immune-compromised patients, bacteria can disseminate systemically to cause a fatal plague-like disease, indicating that the intestinal immune system is critical for control of acute infection. Indeed, studies in mice demonstrate that the intestinal compartment constitutes a major bottleneck against *Yersinia* dissemination, and that bacteria in systemic organs originate predominantly from the intestine rather than gut-associated lymphoid tissues such as the mesenteric lymph nodes (MLN)[15]. A hallmark of *Yersinia* infections is the presence of chronic inflammatory lesions in lymphoid tissue, termed

pyogranulomas (PG)[16,18,22]. Like classical granulomas, pyogranulomas are

characterized by nodular infiltrates of activated phagocytes, including monocytes and

macrophages. Pyogranulomas in particular also contain a core of activated

neutrophils. How these structures form and which cell types are required for this

process are not known. Notably, the contribution of monocytes to pyogranuloma

formation and their role in *Yersinia* restriction remain to be fully understood[17,18].

Interestingly, pyogranulomas have been observed in the human intestine in

response to both *Yp* and *Ye* infection[14,16,19–21] but have not been observed in murine

infection models.

Here we identify the formation of pyogranulomas in the murine intestinal

mucosa during acute enteric *Yersinia* infection. These lesions are enriched in

neutrophils and inflammatory monocytes, and contain comparable numbers of live

bacteria to Peyer's patches (PP). Notably, CCR2-deficient mice, which are deficient

in circulating inflammatory monocytes[37,38], fail to form defined PG, are unable to

contain bacteria within the lesions, and succumb rapidly to infection. Moreover, we

find that mice lacking CCR2 exhibit reduced levels of IL-1 cytokines and blunted

neutrophil activation within intestinal pyogranulomas. Critically, *Yp* lacking either the

virulence plasmid itself, which encodes the type III secreted *Yersinia* Outer Proteins

(Yops) present in pathogenic *Yersinia*, or lacking the Yops that block phagocytosis,

do not induce detectable pyogranulomas, indicating that pyogranulomas are

triggered by *Yersinia* blockade of phagocytosis. Moreover, bacterial restriction and

formation of defined granulomatous lesions in CCR2-deficient mice is restored in

infections with a YopH mutant unable to block phagocytosis, indicating that

inflammatory monocytes enable the host immune system to overcome *Yersinia*

blockade of phagocytosis. Altogether, our study identifies an unappreciated site of

Yersinia colonization within the murine intestinal mucosa, and reveals an essential

function for inflammatory monocytes in establishing pyogranuloma architecture that

enables *Yersinia* clearance.

Results

Intestinal pyogranulomas form acutely following oral Yersinia infection

      Yersinia pseudotuberculosis colonizes gut-associated lymphoid tissues, resulting in pyogranuloma formation within five days following oral infection[22]. Interactions between *Yersinia* and immune cells within systemic tissues have been extensively documented[23–25]. In our efforts to investigate the interaction between *Yp* and the intestinal immune system, we observed numerous nodular lesions in the gastrointestinal tract five days post-infection, which appeared as punctate areas of increased opacity along the length of the intestine (Fig. 1a). The total number of lesions in any individual mouse ranged from two to over forty, with a mean of twenty. The lesions were non-uniformly distributed along the intestinal tract and were over-represented in the jejunum and ileum (Extended Data Fig. 1a). Histologic analysis of *Yp*-infected intestines revealed focal regions of inflammation characterized by crypt hyperplasia, edema, and submucosal to transmural cellular infiltration, which contrasted with non-lesional areas of the infected intestine that appeared largely unaffected, similar to uninfected controls (Fig. 1b). Furthermore, the lesions contained infiltrates of macrophages and neutrophils surrounding colonies of coccobacilli, similar to structures we and others have observed in lymphoid tissues[22,26], and were therefore termed pyogranulomas (Fig. 1c). Consistently, flow-cytometric analysis of intestinal punch biopsies containing pyogranulomas (PG+), adjacent non-granulomatous tissue (PG-), and uninfected control tissue (uninf) revealed that neutrophils were the most enriched cell type specifically within PG+ tissue, followed by macrophages and inflammatory monocytes (Fig. 1d, Extended Data Fig. 1b). Only minor changes were observed in the frequency or number of

eosinophils, dendritic cells, B cells, and T cells in PG+ compared to PG- or

uninfected tissue (Extended Data Fig. 1c).

Yp pyogranulomas in lymphoid tissues consist of a central bacterial colony,

surrounded proximally by neutrophils, bordered in turn by monocytes and

macrophages[18,26]. Similarly, confocal microscopy of whole mount PG+ samples

demonstrated that intestinal pyogranulomas also contained a central *Yp* colony

surrounded by a dense population of Ly-6G$^+$ neutrophils (Fig. 1e). Interestingly, in

contrast to pyogranuloma architecture in lymphoid tissues, CCR2$^+$ monocytes and

macrophages formed a dense mesh-like network of cells that both overlapped with

and bordered the neutrophils (Fig. 1e). Consistent with the presence of bacterial

colonies detected by histology and fluorescence microscopy, PG+ tissue harbored

high numbers of viable bacteria, in contrast to neighboring PG- tissue (Fig. 1f).

Moreover, the total bacterial burden in PG+ tissue was equivalent to that found in the

PP (Fig. 1g). Since Peyer's patches are viewed as the main entry point and

replicative niche for enteropathogenic *Yersinia* following oral innoculation[27,28], our

data demonstrate that intestinal pyogranulomas represent a previously

unappreciated location of *Yersinia* invasion within the intestinal mucosa, and a likely

site of bacterial restriction or potential dissemination[29,30]. Intestinal lesions also

formed in response to *Yersinia enterocolitica (Ye)* infection (Fig. 1h), indicating that

these lesions are a conserved response to enteric *Yersinia* infection. Together, these

findings uncover the first description to our knowledge of intestinal pyogranulomas

that contain high numbers of viable bacteria within the murine intestinal mucosa

following oral infection and may serve as an intestinal niche for enteric *Yersinia*.

The intestinal innate inflammatory response to Yersinia *is localized to*

pyogranulomas

Intestinal pyogranulomas represent an unrecognized site for enteric *Yersinia* to interact with the murine mucosal immune system. To define the PG-specific inflammatory response, we performed RNA sequencing of punch biopsies containing pyogranulomas (PG+), adjacent non-pyogranuloma tissue (PG-), and uninfected control tissue. Principal-component analysis showed distinct clustering by sample type (Fig. 2a), and comparison of PG+ and PG- samples revealed 355 upregulated and 363 downregulated genes (Fig. 2b). Top upregulated genes included granulocyte- and monocyte-recruiting chemokines (*Cxcl1, Cxcl2, Cxcl3, Cxcl5*), pro-inflammatory cytokines (*Il1b, Il22*), antimicrobial metal-sequestration proteins (*S100a8, S100a9*), and matrix metalloproteases (*Mmp3*) (Fig. 2c). Gene ontology analysis revealed that chemotaxis of myeloid cells and defense against bacterial pathogens predominated the top upregulated responses within PG+ biopsies (Fig. 2d, Extended Data Table 1). These features of the transcriptional response were strikingly similar to those previously reported in *Yp*-infected PP[31]. Indeed, gene set-enrichment analysis indicated that the reported top 50 upregulated genes in *Yp*-infected PP were also enriched in PG+ samples (Fig. 2e). Consistently, protein levels of pro-inflammatory cytokines TNF, CCL2, IL-1β, IL-1α, and IL-6 were elevated within PG+ biopsies compared to PG- and uninfected biopsies (Fig. 2f). Consistent with the histology and microscopy analyses, the inflammatory transcriptional response was localized to PG+ tissue, as PG- samples did not exhibit enrichment of genes or ontology terms related to myeloid cell migration or innate immune-cell activation (Extended Data Fig. 2, Extended Data Table 2). Likewise, production of pro-inflammatory cytokines was not detected above uninfected levels in PG- tissue

(Fig. 2e). Altogether, our transcriptional and cytokine analyses indicate that the pro-

inflammatory response to *Yp* infection in the gut mucosa is driven by innate immune

cells and spatially restricted to pyogranulomas.

Inflammatory monocytes are required for organized pyogranuloma formation and

restriction of Yersinia within intestinal tissue

Inflammatory monocytes promote host defense by differentiating into

phagocytes and antigen presenting cells[3,5,7,32,33], producing pro-inflammatory

cytokines and antimicrobial molecules[4], and modulating the function of other immune

cells to promote pathogen clearance or prevent immunopathology[6,34]. However,

monocyte-derived cells can also promote pathogen replication or pathogen

dissemination to new infection sites[9,29,30,35,36]. Mice lacking the chemokine receptor

CCR2 have a ten-fold reduction in circulating monocytes due to defective bone

marrow-egress[37,38]. Monocytes are rapidly recruited to intestinal pyogranulomas (Fig.

1) and systemic *Yp* infection sites[18,22,26,39,40]. CCR2-deficiency was reported to be

associated with more rapid bacterial clearance from the MLN following enteric

infection[18], but increased susceptibility to intravenous *Yp* infection[17]. We therefore

sought to dissect the contribution of inflammatory monocytes to anti-*Yersinia* innate

immune defense and formation of pyogranulomas. Although infection of CCR2-

deficient mice[41,42] resulted in similar numbers of macroscopic intestinal lesions to WT

mice (Fig. 3a, Extended Data Fig. 3a), blinded analysis by a board-certified

pathologist revealed that intestinal lesions in *Ccr2*^{gfp/gfp} mice (in which insertion of

EGFP into the translation initiation site of *Ccr2* disrupts its expression) had a

disorganized appearance, and many showed a central region of caseation containing

substantial tissue necrosis that was not observed in intestinal PG of WT mice (Fig.

3b). Whereas WT pyogranulomas exhibited robust inflammatory infiltrates and a

defined cellular organization encapsulating central bacterial colonies, *Ccr2[gfp/gfp]*

intestinal lesions contained extensively distributed bacterial colonies with limited

inflammatory immune cell recruitment (Fig. 3c). Two independent lines of CCR2-

deficient mice had significantly higher bacterial burdens in both PG+ and PG-

intestinal tissue biopsies than corresponding tissues of WT mice, suggesting that

monocytes restrict enteric *Yp* infection within pyogranulomas and limit dissemination

to surrounding intestinal tissue (Fig. 3d, Extended Data Fig. 3b). Consistent with their

reduced cellularity in histologic analyses, flow cytometry of *Ccr2[gfp/gfp]* lesions

revealed an overall decrease in viable CD45+ hematopoietic cells compared to WT

pyogranulomas (Fig. 3e). Consistent with the known role of CCR2 in egress of

monocytes out of bone marrow and into circulation[37,38], *Ccr2[gfp/gfp]* intestinal lesions

contained significantly lower numbers of monocytes and macrophages compared

with pyogranulomas from WT mice. Surprisingly, despite similar frequencies of

neutrophils in the MLN and spleen upon infection (Extended Data Fig. 3c), *Ccr2[gfp/gfp]*

intestinal lesions showed reduced neutrophil numbers (Fig. 3e), suggesting that

monocytes or monocyte-derived cells promote recruitment, retention, or survival of

neutrophils within intestinal pyogranulomas during *Yp* infection. Notably,

immunofluorescence microscopy of the lesions revealed that the bacteria expanded

outside range of the neutrophil marker Ly-6G, suggesting that neutrophils were

unable to effectively contain *Yp* in the absence of monocytes. In contrast, *Yp* was

fully encapsulated by neutrophils and CCR2+ cells in CCR2-sufficient PGs (Fig. 3f).

Intriguingly, neutrophil surface expression of the integrin CD11b, an established

marker of neutrophil activation[43–46], was also significantly reduced in both *Ccr2[gfp/gfp]*

PGs and MLN compared to WT mice (Fig. 3g, Extended Data Fig. 3d), indicating a

defect in neutrophil activation in these tissues in the absence of monocytes.

Moreover, IL-1α and IL-1β levels were significantly decreased in *Ccr2$^{gfp/gfp}$* PGs (Fig.

3h), suggesting that monocytes or monocyte-derived cells were required to either

directly release IL-1 or promote IL-1 production by other cells within intestinal PGs.

CCR2-deficiency specifically impacted IL-1 levels within PGs, as levels of other pro-

inflammatory cytokines was not altered in CCR2-deficient mice (Extended Data Fig.

3e). Importantly, we found no effect of CCR2 deficiency on CD4$^+$ or CD8$^+$ T cell

numbers[47] in *Ccr2$^{gfp/gfp}$* intestinal PGs, suggesting that the defect in enteric control of

Yp in CCR2-defficient mice was not due an effect on T cell numbers (Extended Data

Fig. 3f). Altogether, these results demonstrate that monocyte-derived cells are

required to establish robust, functional granulomas that limit intestinal bacterial

replication and dissemination.

Inflammatory monocytes are required for systemic control of Yersinia and host

survival

Following dissemination from the intestine, *Yp* colonizes and induces PG

formation in lymphoid tissues[18,22,26]. Critically, CCR2-deficient mice had significantly

higher *Yp* burdens in the MLN and systemic organs (Fig. 4a, Extended Data Fig. 3g).

Histopathological analysis revealed that similar to intestinal tissues, infected

Ccr2$^{gfp/gfp}$ spleens lacked defined pyogranulomas and exhibited widespread tissue

necrosis, free bacterial colonies, and sparse immune cell recruitment, which

contrasted with WT spleens that contained neutrophils and monocytes that

effectively surrounded *Yp* microcolonies (Fig. 4b-c). Notably, mice lacking CCR2

succumbed rapidly to acute infection (Fig. 4d, Extended Data Fig. 3h). Importantly,

infection of co-housed littermate progeny from heterozygous mating demonstrated

that $Ccr2^{+/+}$ and $Ccr2^{gfp/+}$ mice were equally resistant to *Yp* infection while $Ccr2^{gfp/gfp}$

littermates succumbed (Fig. 4e), indicating that differences in intestinal microbiota

between $Ccr2^{gfp/gfp}$ and WT mice were unlikely to account for differences in

susceptibility to *Yp* infection. Collectively, our findings demonstrate that inflammatory

monocytes are required for formation of organized pyogranulomas in host tissues,

thereby limiting tissue necrosis and systemic bacterial dissemination, ultimately

enabling bacterial control and host survival following oral *Yp* infection.

Intestinal pyogranulomas form in response to Yersinia virulence

Mammalian pathogenic *Yersinia* inject their target cells with effector proteins

(Yops) that are encoded on a virulence plasmid (pYV)[48]. Yops block antimicrobial

defenses, including phagocytosis, production of reactive oxygen species (ROS), and

degranulation[24,49–52]. *Yp* lacking the virulence plasmid (pYV-) are still able to colonize

gut-associated lymphoid tissues, including the MLN, during the acute phase of

infection without causing disease[53]. As granuloma formation occurs in response to

pathogens that thwart immune defenses, we hypothesized that formation of intestinal

pyogranulomas is triggered by the activity of *Yp* effector proteins. Indeed, even at a

ten-fold higher infectious dose, we were unable to detect any macroscopic intestinal

lesions at day 5 post-infection with pYV- bacteria (Fig. 5a), despite significant, albeit

lower, bacterial burdens in the intestinal mucosa (Fig. 5b). We likewise found lower

bacterial burdens within gut-associated and systemic lymphoid tissues of mice

infected with pYV- compared to WT bacteria (Fig. 5c). Intriguingly, monocytes were

still recruited to both the intestinal mucosa and MLN in response to of pYV- infection,

indicating that absence of detectable pyogranulomas was not due to an absolute

lack of inflammatory cell infiltration into the tissue of pYV- infected mice (Fig. 5d,e).

Notably, however, neutrophil recruitment to the intestine or MLN was absent in

response to pYV- infection, demonstrating that neutrophil recruitment or retention in

the intestinal tissue is a response to *Yp* virulence activities (Fig. 5d,e).

To test whether specific effectors were responsible for triggering

pyogranuloma formation, we infected mice with individual or combined mutants of

different Yops. Intestinal lesions formed at similar numbers in mice infected with *Yp*

lacking either YopM (*ΔyopM*) or deficient in YopJ enzymatic activity (YopJ[C172A]),

which block pyrin inflammasome assembly and MAPK signaling[54–60], respectively,

suggesting that neither YopM nor YopJ are solely required to trigger pyogranuloma

formation (Extended Data Fig. 4a). A critical virulence function contained on the pYV

plasmid is disruption of the actin cytoskeleton, leading to blockade of phagocytosis

and inhibition of the reactive oxygen burst[24,61–63]. These Yops (E, H, and T), have

partially overlapping functions and can compensate for one another in certain

settings[62–69]. Notably, *Yp* containing point mutations in each of the key catalytic

residues of these Yops (YopE[R144A], YopT[C139A], and YopH[R409A]) did not induce

intestinal lesions and had burdens similar to pYV- infection (Fig. 5a-c). In contrast,

bacteria lacking only YopE (*ΔyopE*) or both YopE and YopT (*yopET*) induced

formation of similar numbers of intestinal lesions (Fig. 5a) and had wild-type levels of

intestinal colonization (Fig. 5b,c). These data indicate that YopH is sufficient, in the

absence of YopE and YopT, to induce pyogranuloma formation. Interestingly,

bacteria lacking only YopH enzymatic activity or YopE alone had similar numbers of

macroscopic lesions to wild-type-infected mice (Fig. 5a), suggesting that YopH is

sufficient, but not absolutely required to trigger pyogranuloma formation. Notably,

while bacteria lacking either YopE or YopH alone induced similar numbers of

intestinal lesions to wild-type *Yersinia*, bacteria lacking both YopE and YopH

(*yopEH*) induced no detectable lesions, and showed similar levels of colonization to

yopETH bacteria (Fig. 5a-c). Altogether, these data indicate that intestinal

pyogranuloma formation occurs in response to *Yersinia* disruption of host actin

cytoskeletal dynamics and is primarily driven by YopE and YopH.

Yp restriction by CCR2-deficient mice is restored in absence of YopH activity

Our results demonstrate that YopH is sufficient, but not strictly necessary to

induce formation of intestinal pyogranulomas. YopH potently blocks phagocytosis by

innate immune cells, including macrophages and neutrophils[24,49,50,62,69,70]. YopH-

deficient *Yp* therefore have enhanced susceptibility to neutrophil killing *in vitro* and

are attenuated *in vivo*[71–73]. While YopH potently blocks neutrophil functions,

neutrophil depletion increases susceptibility to *Yp* and restores virulence to *yopH-*

mutant bacteria[50,70], indicating that neutrophil defenses nevertheless counteract

YopH activity *in vivo*. How neutrophils counteract the activity of YopH is not

understood. Monocyte and neutrophil crosstalk is essential for optimal immune

defense in many infectious contexts[74]. However, whether monocytes are required to

combat YopH-dependent blockade of neutrophil function is unknown. To test this

possibility, we infected CCR2-deficient mice with WT or YopH-mutant *Yp*. Strikingly,

CCR2-deficient mice infected with YopH[R409A] bacteria exhibited robust neutrophil

recruitment and containment of bacterial colonies similar to WT mice, in contrast to

CCR2-deficient mice infected with WT *Yp* (Fig. 6a). Moreover, both *Ccr2*[gfp/gfp] and

Ccr2[-/-] mice restricted YopH[R409A] bacteria within the intestine (Fig. 6b, Extended Data

Fig. 4a) and systemically (Fig. 6c, Extended Data Fig. 4b). While CCR2-deficient

mice rapidly succumbed to WT *Yp*, they survived infection with YopH[R409A] bacteria

similarly to WT mice (Extended Data Fig. 4c). Altogether, our results uncover a key

role for monocytes in establishment and function of intestinal pyogranulomas,

revealing monocytes as essential in overcoming *Yersinia* suppression of neutrophil

function *in vivo* and preventing acute mortality.

Discussion

308
Granulomas are a conserved immunological response to persistent or long-

term infectious and non-infectious stimuli[1,2]. The natural rodent and human pathogen

Yersinia pseudotuberculosis induces formation of neutrophil-rich lesions termed

pyogranulomas in infected lymphoid tissues[14,16]. Here, we report that pyogranulomas

form throughout the intestine during murine enteropathogenic *Yersinia* infection.

Peyer's patches are considered the primary site of intestinal infection by *Yersinia*.

However, intestinal pyogranulomas harbored a similar total bacterial burden as the

PPs, suggesting that PGs are a previously unrecognized niche for enteropathogenic

Yersinia intestinal colonization. Notably, wild-type mice almost entirely restricted

intestinal *Yp* and inflammation to these granulomatous foci. The pro-inflammatory

transcriptional profile of PGs was similar to *Yp*-infected Peyer's patches[31],

suggesting a shared response to intestinal *Yp* infection driven by the recruitment and

activation of innate immune cells. These structures may therefore serve to limit

bacterial spread and tissue damage within the gut mucosa.

We found that monocyte-derived cells were critical for the architectural

integrity of PGs in the intestine and deeper tissues during *Yersinia* infection. While

other cell types express CCR2, notably activated T cells, we did not observe altered

numbers of these cells within intestinal PGs in CCR2-deficient mice. Absence of

circulating monocytes was associated with widespread tissue necrosis and elevated

bacterial burdens in intestinal and systemic tissues, indicating that the monocyte

lineage is critical for containment of *Yersinia* infection. Furthermore, CCR2-

deficiency was associated with defective pro-inflammatory IL-1 production and

neutrophil activation within intestinal PGs.

A previous study reported that monocytes were dispensable for acute control

of *Yp* following oral inoculation[18]. These studies employed a distinct *Ccr2*[-/-] mouse

line[32,38], originally generated on the 129 mouse background and backcrossed to

C57BL/6[37], raising the possibility that distinct polymorphisms in this mouse line might

account for this difference. *Ccr2*[gfp/gfp] mice were generated on the C57BL/6J

background[41], making it unlikely that our findings are due to immunologically

impactful polymorphisms that co-segregate with the *Ccr2* locus. Infection of co-

housed littermates also demonstrated that increased susceptibility of CCR2-deficient

mice is not due to differences in maternally-transmitted or environmentally-acquired

intestinal microbiota.

Formation of intestinal PGs also required the anti-phagocytic Yops. While

combined deficiency in two effector proteins that block phagocytosis, Yops E and H,

abrogated PG formation, either effector alone was sufficient to induce PG formation.

This suggests that PG formation occurs in response to pathogens that resist immune

clearance, and that the anti-phagocytic activities of YopH and YopE independently

mediate PG formation. Precisely how blockade of phagocytosis by YopsH and E

lead to recruitment of phagocytic cells and assembly of pyogranulomas remains to

be determined in future studies.

Monocytes and neutrophils comprise a large proportion of Yop-injected

cells *in vivo*[75]. YopH blocks neutrophil degranulation, ROS production, phagocytosis,

and release of neutrophil extracellular traps[24,50,70,73,76]. Notably, while monocytes

were required for host survival in response to WT bacteria, they were dispensable for

control of *Yp* lacking YopH activity. Our findings imply that monocytes are critical for

host defense against pathogen blockade of neutrophil function, either by directly

restricting bacterial growth or enhancing neutrophil-mediated bacterial control.

Indeed, intestinal PGs in CCR2-deficient mice exhibited decreased neutrophil

numbers and activation, suggesting that monocytes promote neutrophil function

during intestinal *Yp* infection. Altogether, our study reveals an unappreciated site of

Yersinia-host interaction within the intestine, and provides insight into mechanisms of

Yersinia granuloma formation and function.

Methods

Animals

C57BL/6 wild-type and *Ccr2^gfp/gfp* mice[41] (in which insertion of EGFP into the translation initiation site of *Ccr2* disrupts its expression) were acquired from the Jackson Laboratory and bred at the University of Pennsylvania. *Ccr2^-/-* mice[42] were provided by Dr. Sunny Shin (University of Pennsylvania). Unless specifically noted, all animals were bred by homozygous mating and housed separately by genotype. All animal studies were performed in strict accordance with University of Pennsylvania Institutional Animal Care and Use Committee-approved protocols.

Bacteria

Wild-type *Yp* (strain 32777, serogroup O1)[77] and isogenic mutants[58,77–79] were provided by Dr. James Bliska (Dartmouth College). *Ye* (strain 8081, serogroup O8)[80] was provided by Dr. Stanley Falkow (Stanford University). Additional mutants lacking YopE (*ΔyopE*), enzymatic activity of YopH (YopH^{R409A}) or YopE/YopT/YopH (YopE^{R144A}T^{C139A}H^{R409A}, denoted *yopETH*) were generated by two-step allelic recombination as previously described[79] with plasmids provided by Dr. James Bliska. Fluorescent *Yp* (mCherry$^+$) was generated from plasmids provided by Dr. Kimberly Davis (Johns Hopkins University).

Infections

Yp and *Ye* were cultured to stationary phase at 28°C and 250 rpm shaking for 16 hours in 2xYT broth supplemented with 2 μg/ml triclosan (Millipore Sigma). Mice of either sex between 8-12 weeks of age were fasted for 16 hours and subsequently inoculated by oral gavage with 100-200 μl phosphate-buffered saline (PBS). All

bacterial strains were administered at 2×10^8 colony-forming units (CFU) per mouse

with the exception of pYV- which was administered at 20×10^8 CFU per mouse.

Bacterial CFU quantifications

In order to determine bacterial burdens, tissue biopsies were collected in

sterile PBS, weighed, homogenized for 40 seconds with 6.35 mm ceramic spheres

(MP Biomedical) using a FastPrep-24 bead beater (MP Biomedical) and serially

diluted tenfold in PBS before being plated on LB agar supplemented with 2 µg/ml

triclosan and incubated for two days at room temperature. Dilutions of each sample

were plated in triplicate and expressed as the mean CFU per gram of tissue or per

biopsy.

Protein quantifications

Cytokines were measured in supernatants from homogenized tissue using

Cytometric Bead Array (BD Biosciences) according to manufacturer's instructions

with the following modification: the amount of capture beads, detection reagents, and

sample volumes was scaled down tenfold. Data were collected on an LSRFortessa

flow cytometer (BD Biosciences) and analyzed with FlowJo v10 (BD Biosciences).

Tissue preparation and cell isolation

Lymph nodes were homogenized in R10 buffer consisting of RPMI 1640

(Millipore Sigma) supplemented with 10 mM HEPES (Millipore Sigma), 10% fetal

bovine serum (Omega Scientific), 1 mM sodium pyruvate (Thermo-Fisher Scientific)

and 100 U/ml penicillin + 100 µg/ml streptomycin (Thermo Fisher Scientific), then

passed through 70 µm cell strainers (Fisher Scientific). Intestines were excised,

flushed luminally with sterile PBS to remove the feces, opened longitudinally along

the mesenteric side and placed luminal side down on cutting boards (Epicurean).

Small-intestinal tissue containing pyogranulomas (PG+), adjacent non-

granulomatous areas (PG-) and uninfected control tissue (uninf) were excised using

a 2 mm-ø dermal punch-biopsy tool (Keyes). Biopsies within each mouse were

pooled groupwise, suspended in epithelial-dissociation buffer consisting of calcium

and magnesium-free HBSS (Thermo Fisher Scientific) supplemented with 15 mM

HEPES, 10 mg/ml bovine serum albumin (Millipore Sigma), 5 mM EDTA and 100

U/ml penicillin + 100 μg/ml streptomycin, then incubated for 30 minutes at 37°C

under continuous agitation. To isolate immune cells from the lamina propria, the

tissue was enzymatically digested in R10 buffer, along with 0.5 Wünsch units/ml

liberase TM (Roche), 30 μg/ml DNase I (Roche), and 5 mM $CaCl_2$ for 20 min at 37°C

under continuous agitation. The resulting cell suspensions were filtered through 100

μm cell strainers (Fisher Scientific) and subjected to density-gradient centrifugation

using Percoll (GE Healthcare). Briefly, cells were suspended in 40% Percoll and

centrifuged over a 70% Percoll layer for 20 min at 600 × g with lowest brake at room

temperature. Cells collected between the layers were washed with R10 for

downstream analysis.

Flow cytometry

Unspecific Fc binding was blocked for 15 minutes on ice with unconjugated

anti-CD16/CD32 (clone 93; Thermo-Fisher Scientific). Cells were subsequently

fluorescently labeled for 30 minutes on ice with the following antibodies and

reagents: PE-conjugated rat anti-mouse Siglec-F (E50-2440; BD Biosciences), PE-

TxR-conjugated rat anti-mouse CD11b (M1/70.15; Thermo Fisher Scientific), PE-

Cy5-conjugated mouse anti-mouse NK1.1 (PK136; BioLegend), PE-Cy5.5-

conjugated rat anti-mouse CD4 (RM4-5; Thermo Fisher Scientific), PE-Cy7-

conjugated rat anti-mouse CD3 (17A2; BioLegend), FITC-conjugated Armenian

hamster anti-mouse CD11c (N418; BioLegend), PerCP-Cy5.5-conjugated rat anti-

mouse Ly-6C (HK1.4; Thermo Fisher Scientific), PB-conjugated rat anti-mouse

CD90.2 (53-2.1; BioLegend), BV510-conjugated rat anti-mouse CD19 (1D3; BD

Biosciences), BV605-conjugated Armenian hamster anti-mouse TCRβ (H57-597; BD

Biosciences), BV650-conjugated rat anti-mouse I-A/I-E (M5/114.15.2; BD

Biosciences), BV711-conjugated rat anti-mouse CD8α (53-6.7; BD Biosciences),

BV785-conjugated rat anti-mouse Ly-6G (1A8; Thermo Fisher Scientific), AF647-

conjugated mouse anti-mouse CD64 (X54-5/7.1; BD Biosciences) and AF700-

conjugated mouse anti-mouse CD45.2 (104; BioLegend) along with eF780 viability

dye (BioLegend) diluted in PBS according to manufacturer's instructions. Cells were

fixed for 20 minutes on ice with Cytofix/Cytoperm Fixation/Permeabilization solution

(BD Biosciences). Cells were acquired on an LSRFortessa flow cytometer and data

was analyzed with FlowJo v10. Dead and clustered cells were removed from all

analyses.

Histology

Tissues were fixed in 10% neutral-buffered formalin (Fisher Scientific) and

stored at 4°C until further processed. Tissue pieces were embedded in paraffin,

sectioned by standard histological techniques and stained with hematoxylin and

eosin for subsequent histopathological disease scoring by blinded board-certified

pathologists. Tissue sections were given scores between 0-4 (healthy-severe) for

multiple parameters, including degree of inflammatory cell infiltration, necrosis, and

free bacterial colonies along with tissue-specific parameters such as villus blunting

and crypt hyperplasia. Healthy mice were characterized by and subsequently scored

as having none or low levels of the parameters described, whereas severely afflicted

mice presented with high amounts of the respective parameters.

Fluorescence microscopy

Small-intestinal tissue was dissected and luminal contents were flushed out

with PBS. Intestines were opened longitudinally and ~0.5 cm tissue pieces

containing macroscopically visible lesions were excised. Tissues were fixed in 1%

paraformaldehyde overnight then blocked for two hours at room temperature in

blocking solution containing 10% BSA, 1 µg/ml anti-CD16/32, and 0.5% normal rat

IgG in PBS. AF647-conjugated anti-Ly-6G antibody (1A8; Biolegend) was added at

0.01 mg/ml in 100 µl PBS per sample, then whole tissue was stained for 24 hours at

4°C. Samples were washed three times with PBS, mounted whole onto slides in

Prolong Glass Antifade Mountant (Thermo Fisher Scientific) and cured for two days

at room temperature. Images were acquired on a DMI 6000 laser-scanning confocal

microscope (Leica) with a 20x NA 0.75 oil-immersion objective. The center of the

sample was determined in the Z direction, then imaged. Images were analyzed using

ImageJ v2.1.

RNA sequencing

Intestinal punch biopsies were collected as described above. Five biopsies

per sample type were pooled for each mouse. RNA was extracted using the RNeasy

Plus Mini Kit (Qiagen). Sequence-ready libraries were prepared using the Illumina

TruSeq Stranded Total RNA kit with Ribo-Zero Gold rRNA depletion (Illumina).

Quality assessment and quantification of RNA preparations and libraries were

carried out using an Agilent 4200 TapeStation and Qubit 3, respectively. Samples

were sequenced on an Illumina NextSeq 500 to produce 150–base pair single end

reads with a mean sequencing depth of 9 million reads per sample. Raw reads from

this study were mapped to the mouse reference transcriptome (Ensembl; Mus

musculus GRCm38) using Kallisto v0.46.2[81]. Raw sequence data are available on

the Gene Expression Omnibus (GEO; accession no. GSE194334).

All subsequent analyses were carried out using the statistical computing

environment R v4.0.3 in RStudio v1.2.5042 and Bioconductor[82]. Briefly, transcript

quantification data were summarized to genes using the tximport package[83] and

normalized using the trimmed mean of M values (TMM) method in edgeR[84]. Genes

with <1 CPM in 3 samples were filtered out. Normalized filtered data were variance-

stabilized using the voom function in limma[85], and differentially expressed genes

were identified with linear modeling using limma (FDR ≤ 0.05; absolute $\log_2$FC ≥ 1)

after correcting for multiple testing using Benjamini-Hochberg.

Statistics

Statistical analyses were performed using Prism v9.0 (GraphPad Software).

Independent groups were compared by Mann-Whitney U test. Survival curves were

compared by Mantel-Cox test. Statistical significance is denoted * ($p<0.05$), **

($p<0.01$), *** ($p<0.001$), **** ($p<0.0001$), NS (not significant).

References

1. Pagán, A. J. & Ramakrishnan, L. The Formation and Function of Granulomas. *Annu. Rev. Immunol.* **36**, 639–665 (2018).

2. Petersen, H. J. & Smith, A. M. The Role of the Innate Immune System in Granulomatous Disorders. *Front. Immunol.* **4**, 1–11 (2013).

3. Serbina, N. V, Salazar-Mather, T. P., Biron, C. A., Kuziel, W. A. & Pamer, E. G. TNF/iNOS-producing dendritic cells mediate innate immune defense against bacterial infection. *Immunity* **19**, 59–70 (2003).

4. Casson, C. N. *et al.* Neutrophils and Ly6Chi monocytes collaborate in generating an optimal cytokine response that protects against pulmonary Legionella pneumophila infection. *PLoS Pathog.* **13**, e1006309 (2017).

5. Dunay, I. R. *et al.* Gr1+ Inflammatory Monocytes Are Required for Mucosal Resistance to the Pathogen Toxoplasma gondii. *Immunity* **29**, 306–317 (2008).

6. Espinosa, V. *et al.* Inflammatory Monocytes Orchestrate Innate Antifungal Immunity in the Lung. *PLoS Pathog.* **10**, e1003940 (2014).

7. Seo, S.-U. *et al.* Intestinal macrophages arising from CCR2+ monocytes control pathogen infection by activating innate lymphoid cells. *Nat. Commun.* **6**, 8010 (2015).

8. Petermann, M. *et al.* CCR2 Deficiency Impairs Ly6Clo and Ly6Chi Monocyte Responses in Orientia tsutsugamushi Infection. *Front. Immunol.* **12**, 670219 (2021).

9. Antonelli, L. R. *et al.* Intranasal Poly-IC treatment exacerbates tuberculosis in mice through the pulmonary recruitment of a pathogen-permissive monocyte/macrophage population. *J. Clin. Invest.* **120**, 1674–1682 (2010).

10. St. John, A. L. *et al.* S1P-Dependent Trafficking of Intracellular Yersinia pestis

through Lymph nodes establishes buboes and systemic infection. *Immunity* **41**,

440–450 (2014).

11.   Cambier, C. *et al.* Mycobacteria manipulate macrophage recruitment through

coordinated use of membrane lipids. *Nature* **505**, 218–222 (2014).

12.   Eisele, N. A. *et al.* Salmonella require the fatty acid regulator PPARδ for the

establishment of a metabolic environment essential for long term persistence.

14, 171–182 (2013).

13.   Paff, J. R., Triplett, D. A. & Saari, T. N. Clinical and Laboratory Aspects of

Yersinia pseudotuberculosis Infections, with a Report of Two Cases. *Am. J.*

Clin. Pathol. **66**, 101–110 (1976).

14.   El-Maraghi, N. R. H. & Mair, N. S. The Histopathology of Enteric Infection with

Yersinia pseudotuberculosis. *Am. J. Clin. Pathol.* **71**, 631–639 (1979).

15.   Barnes, P. D., Bergman, M. A., Mecsas, J. C. & Isberg, R. R. Yersinia

pseudotuberculosis disseminates directly from a replicating bacterial pool in

the intestine. *J. Exp. Med.* **203**, 1591–601 (2006).

16.   Lamps, L. W. *et al.* The Role of Yersinia enterocolitica and Yersinia

pseudotuberculosis in Granulomatous Appendicitis: A Histologic and Molecular

Study. *Am. J. Surg. Pathol.* **25**, 508–515 (2001).

17.   Zhang, Y., Tam, J. W., Mena, P., van der Velden, A. W. M. & Bliska, J. B.

CCR2+ Inflammatory Dendritic Cells and Translocation of Antigen by Type III

Secretion Are Required for the Exceptionally Large CD8+ T Cell Response to

the Protective YopE69-77 Epitope during Yersinia Infection. *PLoS Pathog.* **11**,

e1005167 (2015).

18.   Zhang, Y., Khairallah, C., Sheridan, B. S., van der Velden, A. W. M. & Bliska,

J. B. CCR2+ Inflammatory Monocytes Are Recruited to Yersinia

558  pseudotuberculosis Pyogranulomas and Dictate Adaptive Responses at the

559  Expense of Innate Immunity during Oral Infection. *Infect. Immun.* **86**, e00782-

560  17 (2018).

561 19. Kojima, M. *et al.* Immunohistological findings of suppurative granulomas of

562  Yersinia enterocolitica appendicitis: A report of two cases. *Pathol. Res. Pract.*

563  **203**, 115–119 (2007).

564 20. Rohena, F. J., Almira-Suárez, M. I. & González-Keelan, C. Granulomatous

565  Enterocolitis Secondary to Yersinia in an 11-year-old Boy from Puerto Rico,

566  Confirmed by PCR: A Case Report. *P. R. Health Sci. J.* **33**, 27–30 (2014).

567 21. Richardson, T., Jones, M., Akhtar, Y. & Pollard, J. Suspicious Yersinia

568  granulomatous enterocolitis mimicking appendicitis. *BMJ Case Rep.* bcr-2018-

569  224177 (2018). doi:10.1136/bcr-2018-224177

570 22. Peterson, L. W. *et al.* RIPK1-dependent apoptosis bypasses pathogen

571  blockade of innate signaling to promote immune defense. *J. Exp. Med.* **214**,

572  3171–3182 (2017).

573 23. Tsutomu, U. Studies on the Pathogenicity of Yersinia enterocolitica III.

574  Comparative Studies between Y. enterocolitica and Y. pseudotuberculosis.

575  *Microbiol. Immunol.* **21**, 505–516 (1977).

576 24. Bliska, J. B., Brodsky, I. E. & Mecsas, J. Role of the Yersinia

577  pseudotuberculosis Virulence Plasmid in Pathogen-Phagocyte Interactions in

578  Mesenteric Lymph Nodes. *EcoSal Plus* **9**, eESP-0014-2021 (2021).

579 25. Davis, K. M. All Yersinia Are Not Created Equal: Phenotypic Adaptation to

580  Distinct Niches Within Mammalian Tissues. *Front. Cell. Infect. Microbiol.* **8**, 1–8

581  (2018).

582 26. Davis, K. M., Mohammadi, S. & Isberg, R. R. Community Behavior and Spatial

Regulation within a Bacterial Microcolony in Deep Tissue Sites Serves to

Protect against Host Attack. *Cell Host Microbe* **17**, 21–31 (2015).

27. Clark, M. A., Hirst, B. H. & Jepson, M. A. M-cell Surface β1 Integrin Expression

and Invasin-Mediated Targeting of Yersinia pseudotuberculosis to Mouse

Peyer's Patch M Cells. *Infect. Immun.* **66**, 1237–1243 (1998).

28. Fasciano, A. C. *et al.* Yersinia pseudotuberculosis YopE prevents uptake by M

cells and instigates M cell extrusion in human ileal enteroid-derived

monolayers. *Gut Microbes* **13**, 1988390 (2021).

29. Clay, H. *et al.* Dichotomous Role of the Macrophage in Early Mycobacterium

marinum Infection of the Zebrafish. *Cell Host Microbe* **2**, 29–39 (2007).

30. Davis, J. M. & Ramakrishnan, L. The Role of the Granuloma in Expansion and

Dissemination of Early Tuberculous Infection. *Cell* **136**, 37–49 (2009).

31. Nuss, A. M. *et al.* Tissue dual RNA-seq allows fast discovery of infection-

specific functions and riboregulators shaping host-pathogen transcriptomes.

Proc. Natl. Acad. Sci. U. S. A. **114**, E791–E800 (2017).

32. Zigmond, E. *et al.* Ly6C hi Monocytes in the Inflamed Colon Give Rise to

Proinflammatory Effector Cells and Migratory Antigen-Presenting Cells.

Immunity **37**, 1076–1090 (2012).

33. Hohl, T. M. *et al.* Inflammatory Monocytes Facilitate Adaptive CD4 T Cell

Responses during Respiratory Fungal Infection. *Cell Host Microbe* **6**, 470–481

(2009).

34. Grainger, J. R. *et al.* Inflammatory monocytes regulate pathologic responses to

commensals during acute gastrointestinal infection. *Nat. Med.* **19**, 713–721

(2013).

35. Repasy, T. *et al.* Intracellular bacillary burden reflects a burst size for

Mycobacterium tuberculosis in vivo. *PLoS Pathog.* **9**, e1003190 (2013).

36.   Cronan, M. R. *et al.* Macrophage Epithelial Reprogramming Underlies

Mycobacterial Granuloma Formation and Promotes Infection. *Immunity* **45**,

861–876 (2016).

37.   Kuziel, W. A. *et al.* Severe reduction in leukocyte adhesion and monocyte

extravasation in mice deficient in CC chemokine receptor 2. *Proc. Natl. Acad.*

Sci. U. S. A. **94**, 12053–12058 (1997).

38.   Serbina, N. V & Pamer, E. G. Monocyte emigration from bone marrow during

bacterial infection requires signals mediated by chemokine receptor CCR2.

Nat. Immunol. **7**, 311–317 (2006).

39.   Philip, N. H. *et al.* Caspase-8 mediates caspase-1 processing and innate

immune defense in response to bacterial blockade of NF-κB and MAPK

signaling. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 7385–7390 (2014).

40.   Philip, N. H. *et al.* Activity of Uncleaved Caspase-8 Controls Anti-bacterial

Immune Defense and TLR-Induced Cytokine Production Independent of Cell

Death. *PLoS Pathog.* **12**, e1005910 (2016).

41.   Satpathy, A. T. *et al.* Notch2-dependent classical dendritic cells orchestrate

intestinal immunity to attaching-and-effacing bacterial pathogens. *Nat.*

Immunol. **14**, 937–948 (2013).

42.   Boring, L. *et al.* Impaired Monocyte Migration and Reduced Type 1 (Th1)

Cytokine Responses in C-C Chemokine Receptor 2 Knockout Mice. *J. Clin.*

Invest. **100**, 2552–2561 (1997).

43.   Borjesson, D. L., Simon, S. I., Hodzic, E., Ballantyne, C. M. & Barthold, S. W.

Kinetics of CD11b/CD18 Up-Regulation During Infection with the Agent of

Human Granulocytic Ehrlichiosis in Mice. *Lab. Investig.* **82**, 303–311 (2002).

44.   Mann, B. S. & Chung, K. F. Blood neutrophil activation markers in severe

asthma: lack of inhibition by prednisolone therapy. *Respir. Res.* **7**, 1–10

(2006).

45.   Yipp, B. G. *et al.* The lung is a host defense niche for immediate neutrophil-

mediated vascular protection. *Sci. Immunol.* **2**, eaam8929 (2017).

46.   Yoon, J. W., Pahl, M. V & Vaziri, N. D. Spontaneous leukocyte activation and

oxygen-free radical generation in end-stage renal disease. *Kidney Int.* **71**,

167–172 (2007).

47.   Bakos, E. *et al.* CCR2 Regulates the Immune Response by Modulating the

Interconversion and Function of Effector and Regulatory T Cells. *J. Immunol.*

198, 4659–4671 (2017).

48.   Portnoy, D. A., Wolf-Watz, H., Bolin, I., Beeder, A. B. & Falkow, S.

Characterization of Common Virulence Plasmids in Yersinia Species and Their

Role in the Expression of Outer Membrane Proteins. *Infect. Immun.* **43**, 108–

114 (1984).

49.   Zhang, L. *et al.* The Functions of Effector Proteins in Yersinia Virulence. *Polish*

J. Microbiol. **65**, 5–12 (2016).

50.   Shaban, L. *et al.* Yersinia pseudotuberculosis YopH targets SKAP2-dependent

and independent signaling pathways to block neutrophil antimicrobial

mechanisms during infection. *PLOS Pathog.* **16**, e1008576 (2020).

51.   Green, E. R. *et al.* Fis Is Essential for Yersinia pseudotuberculosis Virulence

and Protects against Reactive Oxygen Species Produced by Phagocytic Cells

during Infection. *PLoS Pathog.* **12**, e1005898 (2016).

52.   Songsungthong, W., Higgins, M. C., Rolán, H. G., Murphy, J. L. & Mecsas, J.

ROS-inhibitory activity of YopE is required for full virulence of Yersinia in mice.

Cell. Microbiol. **12**, 988–1001 (2010).

53.   Balada-Llasat, J. M. & Mecsas, J. C. Yersinia Has a Tropism for B and T Cell

Zones of Lymph Nodes That Is Independent of the Type III Secretion System.

PLoS Pathog. **2**, e86 (2006).

54.   Yoon, S., Liu, Z., Eyobo, Y. & Orth, K. Yersinia Effector YopJ Inhibits Yeast

MAPK Signaling Pathways by an Evolutionarily Conserved Mechanism. *J. Biol.*

Chem. **278**, 2131–2135 (2003).

55.   Mukherjee, S. *et al.* Yersinia YopJ Acetylates and Inhibits Kinase Activation by

Blocking Phosphorylation. *Science* **312**, 1211–1214 (2006).

56.   Palmer, L. E., Hobbie, S., Galán, J. E. & Bliska, J. B. YopJ of Yersinia

pseudotuberculosis is required for the inhibition of macrophage TNF-α

production and downregulation of the MAP kinases p38 and JNK. *Mol.*

Microbiol. **27**, 953–965 (1998).

57.   Brodsky, I. E. & Medzhitov, R. Reduced Secretion of YopJ by Yersinia Limits

In Vivo Cell Death but Enhances Bacterial Virulence. *PLoS Pathog.* **4**,

e1000067 (2008).

58.   Chung, L. K. *et al.* The Yersinia Virulence Factor YopM Hijacks Host Kinases

to Inhibit Type III Effector-Triggered Activation of the Pyrin Inflammasome. *Cell*

Host Microbe **20**, 296–306 (2016).

59.   Larock, C. N. & Cookson, B. T. The Yersinia Virulence Effector YopM Binds

Caspase-1 to Arrest Inflammasome Assembly and Processing. *Cell Host*

Microbe **12**, 799–805 (2012).

60.   Ratner, D. *et al.* The Yersinia pestis Effector YopM Inhibits Pyrin

Inflammasome Activation. *PLoS Pathog.* **12**, 1–17 (2016).

61.   Grosdent, N., Maridonneau-Parini, I., Sory, M.-P. & Cornelis, G. R. Role of

Yops and Adhesins in Resistance of Yersinia enterocolitica to Phagocytosis.

Infect. Immun. **70**, 4165–4176 (2002).

62. Green, S. P., Hartland, E. L., M., Robins-Browne, R. M. & Phillips, W. A. Role

of YopH in the suppression of tyrosine phosphorylation and respiratory burst

activity in murine macrophages infected with Yersinia enterocolitica. *J. Leukoc.*

Biol. **57**, 972–977 (1995).

63. Bliska, J. B. & Black, D. S. Inhibition of the Fc Receptor-Mediated Oxidative

Burst in Macrophages by the Yersinia pseudotuberculosis Tyrosine

Phosphatase. *Infect. Immun.* **63**, 681–685 (1995).

64. Bliska, J. B., Guan, K., Dixon, J. E. & Falkow, S. Tyrosine phosphate

hydrolysis of host proteins by an essential Yersinia virulence determinant.

Proc. Natl. Acad. Sci. U. S. A. **88**, 1187–1191 (1991).

65. Rosqvist, R., Bölin, I. & Wolf-Watz, H. Inhibition of Phagocytosis in Yersinia

pseudotuberculosis: a Virulence Plasmid-Encoded Ability Involving the Yop2b

Protein. *Infect. Immun.* **56**, 2139–2143 (1988).

66. Rosqvist, R., Forsberg, Å. & Wolf-Watz, H. Intracellular Targeting of the

Yersinia YopE Cytotoxin in Mammalian Cells Induces Actin Microfilament

Disruption. *Infect. Immun.* **59**, 4562–4569 (1991).

67. Galyov, E. E., Håkansson, S., Forsberg, Å. & Wolf-Watz, H. A secreted protein

kinase of Yersinia pseudotuberculosis is an indispensable virulence

determinant. *Nature* **361**, 33–50 (1993).

68. Black, D. S. & Bliska, J. B. The RhoGAP activity of the Yersinia

pseudotuberculosis cytotoxin YopE is required for antiphagocytic function and

virulence. *Mol. Microbiol.* **37**, 515–527 (2000).

69. Mecsas, J., Raupach, B. & Falkow, S. The Yersinia Yops inhibit invasion of

Listeria, Shigella and Edwardsiella but not Salmonella into epithelial cells. *Mol.*

Microbiol. **28**, 1269–1281 (1998).

70. Rolán, H. G., Durand, E. A. & Mecsas, J. Identifying Yersinia YopH-Targeted

Signal Transduction Pathways that Impair Neutrophil Responses during In

Vivo Murine Infection. *Cell Host Microbe* **14**, 306–317 (2013).

71. Logsdon, L. K. & Mecsas, J. C. Requirement of the Yersinia

pseudotuberculosis Effectors YopH and YopE in Colonization and Persistence

in Intestinal and Lymph Tissues. *Infect. Immun.* **71**, 4595–4607 (2003).

72. Fisher, M. L., Castillo, C. & Mecsas, J. Intranasal Inoculation of Mice with

Yersinia pseudotuberculosis Causes a Lethal Lung Infection That Is

Dependent on Yersinia Outer Proteins and PhoP. *Infect. Immun.* **75**, 429–442

(2007).

73. Westermark, L., Fahlgren, A. & Fällman, M. Yersinia pseudotuberculosis

Efficiently Escapes Polymorphonuclear Neutrophils during Early Infection.

Infect. Immun. **82**, 1181–1191 (2014).

74. Silva, M. T. When two is better than one: macrophages and neutrophils work in

concert in innate immunity as complementary and cooperative partners of a

myeloid phagocyte system. *J. Leukoc. Biol.* **87**, 93–106 (2010).

75. Durand, E. A., Maldonado-Arocho, F. J., Castillo, C., Walsh, R. L. & Mecsas, J.

C. The presence of professional phagocytes dictates the number of host cells

targeted for Yop translocation during infection. *Cell. Microbiol.* **12**, 1064–1082

(2010).

76. Taheri, N., Fahlgren, A. & Fällman, M. Yersinia pseudotuberculosis Blocks

Neutrophil Degranulation. *Infect. Immun.* **84**, 3369–3378 (2016).

77. Simonet, M. & Falkow, S. Invasin expression in Yersinia pseudotuberculosis.

Infect. Immun. **60**, 4414–4417 (1992).

78.   McPhee, J. B., Mena, P. & Bliska, J. B. Delineation of Regions of the Yersinia

YopM Protein Required for Interaction with the RSK1 and PRK2 Host Kinases

and Their Requirement for Interleukin-10 Production and Virulence. *Infect.*

Immun. **78**, 3529–3539 (2010).

79.   Zhang, Y., Murtha, J., Roberts, M. A., Siegel, R. M. & Bliska, J. B. Type III

secretion decreases bacterial and host survival following phagocytosis of

Yersinia pseudotuberculosis by macrophages. *Infect. Immun.* **76**, 4299–4310

(2008).

80.   Portnoy, D. A., Moseley, S. L. & Falkow, S. Characterization of Plasmids and

Plasmid-Associated Determinants of Yersinia enterocolitica Pathogenesis.

Infect. Immun. **31**, 775–782 (1981).

81.   Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic

RNA-seq quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

82.   Huber, W. *et al.* Orchestrating high-throughput genomic analysis with

Bioconductor. *Nat. Methods* **12**, 115–121 (2015).

83.   Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq:

transcript-level estimates improve gene-level inferences. *F1000Research* **4**,

1521 (2016).

84.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor

package for differential expression analysis of digital gene expression data.

Bioinformatics **26**, 139–140 (2010).

85.   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-

sequencing and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).

Acknowledgements

We thank Dr. James Bliska for generously providing plasmids for Yop mutant

Yp strains, as well as Dr. Kimberly Davis for generously providing the mCherry+ *Yp*

plasmid. We thank the staff at the PennVet Comparative Pathology Core for their

help in preparing the histological samples. We thank Dr. David Christian and Dr.

Andrea Stout for key advice on confocal microscopy methods, and Dr. Sunny Shin

for constructive editorial comments and scientific discussion. This work was

supported by NIH Awards R01AI128530, R01139102A1 (IEB) and a BWF

Investigator in the Pathogenesis of Infectious Disease Award (IEB); the Foundation

Blanceflor Postdoctoral Scholarship (DS), the Swedish Society for Medical Research

postdoctoral fellowship (DS) and the Sweden-America Foundation J. Sigfrid Edström

award (DS); NIH NRSA F31AI160741-01 (RM); Mark Foundation Grant 19-011MIA,

and F32 AI164655 (JPG); NSF GRFP Award (SP). We thank members of the

Brodsky lab for scientific discussion and Dr. Daniel Grubaugh for comments on the

manuscript.

Author contributions

DS established the initial findings of intestinal granulomas. DS, RM, and IEB conceptualized the study and devised experiments. DS and RM devised the methodology and performed experiments. SP, JPG, and IR performed experiments. CAA, ER, and ML performed the histology and histopathological scoring. EK prepared the RNA-sequencing libraries. RM and DB analyzed the RNA-sequencing data. IEB acquired the funding and supervised the study. DS, RM, and IEB wrote the original draft. All authors reviewed and edited the manuscript.

781 **Competing interests**

782 The author declare no competing interests.

Figure legends

Figure 1. Intestinal pyogranulomas form acutely following oral *Yersinia* infection

(a) Longitudinal small-intestinal segments from uninf (top) and *Yp*-infected (bottom) mice with arrows depicting lesions. Scale bar = 3 mm. Representative image of >3 independent experiments.

(b) H&E-stained paraffin-embedded longitudinal small-intestinal sections from uninf (left) and *Yp*-infected (middle and right) mice. Scale bars = 200 µm. Representative images of three independent experiments.

(c) H&E-stained paraffin-embedded longitudinal small-intestinal PG+ section from a *Yp*-infected mouse depicting an encircled bacterial colony. Scale bar = 10 µm. Representative image of three independent experiments.

(d) Flow-cytometry plots (left side) identifying CD11b$^+$ Ly-6G$^+$ neutrophils (top), CD11b$^+$ CD64$^+$ Ly-6C$^+$ monocytes, and CD11b$^+$ CD64$^+$ Ly-6C$^-$ MHC-II$^+$ macrophages (bottom) in small-intestinal uninf (left), PG- (middle) and PG+ (right) tissue with quantification (right side) of frequency (top) and total number (bottom) of neutrophils (left), monocytes (middle) and macrophages (right) in small-intestinal uninf, PG- and PG+ tissue. Each circle represents one mouse. Lines represent median. Pooled data from four independent experiments. Representative images of four independent experiments.

(e) Fluorescently labeled whole-mount small-intestinal PG+ tissue from an *Yp*-infected *Ccr2$^{gfp/+}$* mouse. White (*Yp*-mCherry), magenta (Ly-6G-AF647), green (CCR2-GFP). Scale bars = 100 µm. Representative images of two independent experiments.

(f) Bacterial burdens in small-intestinal uninf, PG-, and PG+ tissue. Each circle

represents one mouse. Lines represent geometric mean. Pooled data from three

independent experiments.

(g) Cumulative bacterial burdens per intestine in PG+ tissue and PP. Each circle

represents one mouse. Lines represent geometric mean. Pooled data from five

independent experiments.

(h) Quantification of total number of intestinal lesions upon *Yp* or *Ye* infection. Each

circle represents one mouse. Lines represent median. Pooled data from three

independent experiments.

Mann-Whitney U test was performed for all statistical analyses. Abbreviations: *Yp*

(*Yersinia pseudotuberculosis*), PP (Peyer's patch), uninf (uninfected control), PG+

(pyogranuloma), PG- (non-granulomatous tissue), H&E (hematoxylin and eosin),

CFU (colony-forming units), ND (not detected), NS (not significant, $p > 0.05$).

Figure 2. The intestinal innate inflammatory response to *Yersinia* is localized

to pyogranulomas

(**a**) Principal component analysis of PG+ (pink), PG- (green), and uninf (gray)

samples at day 5 post-infection. Five biopsies were pooled per mouse.

(**b**) Heatmap of all differentially expressed genes in PG+ compared to PG- samples.

False discovery rate < 0.05 using Benjamini-Hochberg procedure.

(**c**) Heatmap of top 30 significantly upregulated genes in PG+ compared to PG-

samples in descending order by fold change. False discovery rate < 0.05 using

Benjamini-Hochberg procedure.

(**d**) Gene ontology analysis of top 30 upregulated genes by fold change only in PG+

compared to PG- samples.

(**e**) Gene set enrichment analysis (GSEA) of top 50 upregulated genes in *Yp*-infected

Peyer's patches. NES = normalized enrichment score; FDR = false discovery rate.

(**f**) Cytokine levels in homogenates of tissue punch biopsies were measured by

cytometric bead array. Lines represent group mean. Mann-Whitney U test was

performed for statistical analysis. NS = not significant. Data from one or two pooled

independent experiments.

Figure 3. Inflammatory monocytes are required for organized pyogranuloma

formation and restriction of Yersinia within intestinal tissue

(**a**) Quantification of total number of intestinal lesions at day 5 post infection. Each

circle represents one mouse. Lines represent median. Pooled data from four

independent experiments.

(**b**) H&E-stained paraffin-embedded longitudinal small intestinal sections from *Yp*-

infected mice. Scale bars = 250 µm (top) and 50 µm (bottom). Representative

images of two independent experiments.

(**c**) Histopathological scores of small intestinal tissue from uninfected and *Yp*-

infected mice. Each mouse was given a score between 0-4 (healthy-severe) for

signs of pathology. Each circle represents one mouse. Lines represent median.

Pooled data from two independent experiments.

(**d**) Bacterial burdens in small-intestinal PG- and PG+ tissue. Each circle represents

the mean *Yp*-CFU of 3-5 pooled punch biopsies from one mouse. Lines represent

geometric mean. Pooled data from six independent experiments.

(**e**) Total number of CD45$^+$ cells, monocytes, macrophages, and neutrophils in small

intestinal uninf, PG-, and PG+ tissue. Each circle represents the mean of 3-10

pooled punch biopsies from one mouse. Lines represent median. Pooled data from

five independent experiments.

(**f**) Fluorescently labeled whole-mount small intestinal PG+ tissue from *Yp*-infected

Ccr2^{gfp/+} (top) and *Ccr2*^{gfp/gfp} (bottom) mice. Scale bars = 100 μm. Representative

images of two independent experiments.

(**g**) PG+ neutrophil CD11b expression was measured by flow cytometry. Lines

represent median. Representative of four independent experiments.

(**h**) Cytokine levels in homogenates of tissue punch biopsies were measured by

cytometric bead array. Lines represent median. Pooled data from three independent

experiments.

Mann-Whitney U test was performed for all statistical analyses. NS = not significant.

Figure 4. Inflammatory monocytes are required for systemic control of *Yersinia*

and host survival

(**a**) Bacterial burdens in indicated organs at day 5 post infection. Each circle

represents one mouse. Lines represent geometric mean. Pooled data from four

independent experiments.

(**b**) H&E-stained paraffin-embedded longitudinal spleen sections from WT and

Ccr2^{gfp/gfp} mice. Scale bars = 500 μm (top) and 50 μm (top). Representative images

of two independent experiments.

(**c**) Histopathological scores of spleens from uninfected and *Yp*-infected mice. Each

mouse was given a score between 0-4 (healthy-severe) for signs of pathology. Each

circle represents one mouse. Lines represent median. Pooled data from two

independent experiments.

(**d**) Survival of infected WT (n=16) and *Ccr2^{gfp/gfp}* (n=13) mice. Pooled data from two

independent experiments.

(**e**) Survival of infected littermate wild-type *Ccr2^{+/+}* (n=20), heterozygous *Ccr2^{gfp/+}*

(n=19)*,* and homozygous *Ccr2^{gfp/gfp}* (n=15) mice. Pooled data from three independent

experiments.

Statistical analyses by (a, c) Mann-Whitney U test and (d-e) Mantel-Cox test. NS =

not significant.

Figure 5. Intestinal pyogranulomas form in response to *Yersinia* virulence

(**a**) Quantification of total number of intestinal lesions upon infection. Each symbol

represents one mouse. Lines represent median. Pooled data from 2-3 independent

experiments.

(**b**) Bacterial burdens in small-intestinal PG- (left) and PG+ (right) tissue. Each

symbol represents one mouse. Lines represent geometric mean. Pooled data from

2-3 independent experiments.

(**c**) Bacterial burdens in indicated organs. Each symbol represents one mouse. Lines

represent geometric mean. Pooled data from 2-3 independent experiments.

(**d**) Frequency of monocytes (left) and neutrophils (right) in small-intestinal uninf, PG-

and PG+ tissue. Each symbol represents one mouse. Lines represent median.

Pooled data from two independent experiments.

(**e**) Frequency of monocytes (left) and neutrophils (right) in MLN. Each symbol

represents one mouse. Lines represent median. Representative data from two

independent experiments.

Mann-Whitney U test was performed for all statistical analyses.

(**d**) Survival of infected WT (n=16) and *Ccr2*$^{gfp/gfp}$ (n=13) mice. Pooled data from two

independent experiments.

(**e**) Survival of infected littermate wild-type *Ccr2*$^{+/+}$ (n=20), heterozygous *Ccr2*$^{gfp/+}$

(n=19)*,* and homozygous *Ccr2*$^{gfp/gfp}$ (n=15) mice. Pooled data from three independent

experiments.

Statistical analyses by (a, c) Mann-Whitney U test and (d-e) Mantel-Cox test. NS =

not significant.

Figure 5. Intestinal pyogranulomas form in response to *Yersinia* virulence

(**a**) Quantification of total number of intestinal lesions upon infection. Each symbol

represents one mouse. Lines represent median. Pooled data from 2-3 independent

experiments.

(**b**) Bacterial burdens in small-intestinal PG- (left) and PG+ (right) tissue. Each

symbol represents one mouse. Lines represent geometric mean. Pooled data from

2-3 independent experiments.

(**c**) Bacterial burdens in indicated organs. Each symbol represents one mouse. Lines

represent geometric mean. Pooled data from 2-3 independent experiments.

(**d**) Frequency of monocytes (left) and neutrophils (right) in small-intestinal uninf, PG-

and PG+ tissue. Each symbol represents one mouse. Lines represent median.

Pooled data from two independent experiments.

(**e**) Frequency of monocytes (left) and neutrophils (right) in MLN. Each symbol

represents one mouse. Lines represent median. Representative data from two

independent experiments.

Mann-Whitney U test was performed for all statistical analyses.

Figure 6. *Yp* restriction by CCR2-deficient mice is restored in absence of YopH

activity

(**a**) H&E-stained paraffin-embedded transverse small intestinal sections from WT

(top) and *Ccr2^{gfp/gfp}* (bottom) mice infected with WT (left) or YopH^{R409A} (right) *Yp* at

day 5 post infection. Scale bars = 100 μm. Representative images of two

independent experiments.

(**b**) Bacterial burdens in small-intestinal PG- and PG+ tissue at day 5 post infection.

Each symbol represents one mouse. Lines represent geometric mean. Pooled data

from two independent experiments.

(**c**) Bacterial burdens in indicated organs at day 5 post infection. Each symbol

represents one mouse. Lines represent geometric mean. Pooled data from two

independent experiments.

Mann-Whitney U test was performed for all statistical analyses. NS = not significant.

Extended Data Figure 1. Intestinal pyogranulomas from acutely following oral

***Yersinia* infection**

(a) Frequency distribution (left) of lesions along the intestinal tract, with graphical key

of anatomical segments (right). Each colored bar represents the mean frequency of

lesions in a given segment (n = 20 mice). Only mice with >9 total pyogranulomas

were included. Pooled data from three independent experiments.

(b) Flow cytometry plots displaying the gating strategy employed to identify

eosinophils, dendritic cells, B cells and T cells in small-intestinal tissue.

Representative images of four independent experiments.

(c) Frequency (top) and total number (bottom) of eosinophils, dendritic cells, B cells,

CD4+ T cells and CD8+ T cells (left to right) in small-intestinal uninf, PG- and PG+

tissue. Each circle represents the mean of 3-10 pooled punch biopsies (ø = 2 mm)

from one mouse. Lines represent median. Pooled data from four independent

experiments.

Mann-Whitney U test was performed for all statistical analyses. NS = not significant.

Extended Data Figure 2. Non-pyogranuloma tissue does not undergo a pro-

inflammatory response during intestinal *Yp* infection

(**a**) Heatmap of top 30 significantly upregulated genes in PG- compared to uninfected

samples in descending order by fold change. False discovery rate < 0.05 using

Benjamini-Hochberg procedure.

(**b**) Gene ontology analysis of top 30 upregulated genes by fold change only in PG-

compared to uninfected samples. Dotted line denotes p = 0.05.

Extended Data Figure 3. *Ccr2*[-/-] mice have increased susceptibility to oral

Yersinia infection*.*

(**a**) Quantification of total number of intestinal lesions at day 3 post infection. Each

circle represents one mouse. Line represents median. Pooled data from two

independent experiments.

(**b**) Bacterial burdens in small intestinal PG- and PG+ tissue at day 3 post infection.

Each circle represents the mean *Yp*-CFU of 3-5 pooled punch biopsies from one

mouse. Lines represent geometric mean. Pooled data from four independent

experiments.

(**c**) Frequency of neutrophils in MLN and spleen at day 5 post infection. Each circle

represents the mean of 3-10 pooled punch biopsies from one mouse. Lines

represent median. Pooled data from two independent experiments.

(**d**) MLN neutrophil CD11b expression was measured by flow cytometry. Lines

represent median. Representative of two independent experiments.

(**e**) Cytokine levels in homogenates of tissue punch biopsies at day 5 post infection

were measured by cytometric bead array. Lines represent median. Pooled data from

three independent experiments.

(**f**) Total number of CD4$^+$ and CD8$^+$ T cells in small intestinal PG+ tissue. Each circle

represents the mean of 3-10 pooled punch biopsies from one mouse. Lines

represent median. Pooled data from 4 independent experiments.

(**g**) Bacterial burdens in indicated organs at day 3 post infection. Each circle

represents one mouse. Lines represent geometric mean. Pooled data from four

independent experiments.

(**h**) Survival of infected mice. Pooled data from two independent experiments.

Statistical analyses by (a-g) Mann-Whitney U test or (h) Mantel-Cox test. NS = not

significant.

Extended Data Figure 4. Monocytes are dispensable for restriction of YopH

mutant *Yp*

(**a**) Quantification of total number of intestinal lesions upon infection. Each symbol

represents one mouse. Lines represent median. Pooled data from 2 independent

experiments.

(**b**) Bacterial burdens in small-intestinal PG- and PG+ tissue at day 3 post infection.

Each symbol represents one mouse. Lines represent geometric mean. Pooled data

from two independent experiments.

(**c**) Bacterial burdens in indicated organs at day 3 post infection. Each symbol

represents one mouse. Lines represent geometric mean. Pooled data from two

independent experiments.

(**d**) Survival of *Yp*-infected mice. Pooled data from two independent experiments.

Statistical analyses by (a-c) Mann-Whitney U test and (d) Mantel-Cox test.

Extended Data Table 1. Gene ontology analysis of PG+ vs PG- samples

Extended Data Table 2. Gene ontology analysis of PG- vs uninfected samples

Supplementary code file

Figure 1. Intestinal pyogranulomas form acutely following oral *Yersinia* infection

(**a**) Longitudinal small-intestinal segments from uninf (top) and *Yp*-infected (bottom) mice with arrows depicting lesions. Scale bar = 3 mm. Representative image of >3 independent experiments.

(**b**) H&E-stained paraffin-embedded longitudinal small-intestinal sections from uninf (left) and *Yp*-infected (middle and right) mice. Scale bars = 200 μm. Representative images of three independent experiments.

(**c**) H&E-stained paraffin-embedded longitudinal small-intestinal PG+ section from a *Yp*-infected mouse depicting an encircled bacterial colony. Scale bar = 10 μm. Representative image of three independent experiments.

(**d**) Flow-cytometry plots (left side) identifying CD11b⁺ Ly-6G⁺ neutrophils (top), CD11b⁺ CD64⁺ Ly-6C⁺ monocytes, and CD11b⁺ CD64⁺ Ly-6C⁻ MHC-II⁺ macrophages (bottom) in small-intestinal uninf (left), PG- (middle) and PG+ (right) tissue with quantification (right side) of frequency (top) and total number (bottom) of neutrophils (left), monocytes (middle) and macrophages (right) in small-intestinal uninf, PG- and PG+ tissue. Each circle represents one mouse. Lines represent median. Pooled data from four independent experiments. Representative images of four independent experiments.

(**e**) Fluorescently labeled whole-mount small-intestinal PG+ tissue from an *Yp*-infected *Ccr2^gfp/+* mouse. White (*Yp*-mCherry), magenta (Ly-6G-AF647), green (CCR2-GFP). Scale bars = 100 μm. Representative images of two independent experiments.

(**f**) Bacterial burdens in small-intestinal uninf, PG-, and PG+ tissue. Each circle represents one mouse. Lines represent geometric mean. Pooled data from three independent experiments.

(**g**) Cumulative bacterial burdens per intestine in PG+ tissue and PP. Each circle represents one mouse. Lines represent geometric mean. Pooled data from five independent experiments.

(**h**) Quantification of total number of intestinal lesions upon *Yp* or *Ye* infection. Each circle represents one mouse. Lines represent median. Pooled data from three independent experiments.

Mann-Whitney U test was performed for all statistical analyses. Abbreviations: *Yp* (*Yersinia pseudotuberculosis*), PP (Peyer's patch), uninf (uninfected control), PG+ (pyogranuloma), PG- (non-granulomatous tissue), H&E (hematoxylin and eosin), CFU (colony-forming units), ND (not detected), NS (not significant, p > 0.05).

Figure 2. The intestinal innate inflammatory response to *Yersinia* is localized to pyogranulomas

(**a**) Principal component analysis of PG+ (pink), PG- (green), and uninf (gray) samples at day 5 post-infection. Five biopsies were pooled per mouse.

(**b**) Heatmap of all differentially expressed genes in PG+ compared to PG- samples. False discovery rate < 0.05 using Benjamini-Hochberg procedure.

(**c**) Heatmap of top 30 significantly upregulated genes in PG+ compared to PG- samples in descending order by fold change. False discovery rate < 0.05 using Benjamini-Hochberg procedure.

(**d**) Gene ontology analysis of top 30 upregulated genes by fold change only in PG+ compared to PG- samples.

(**e**) Gene set enrichment analysis (GSEA) of top 50 upregulated genes in *Yp*-infected Peyer's patches. NES = normalized enrichment score; FDR = false discovery rate.

(**f**) Cytokine levels in homogenates of tissue punch biopsies were measured by cytometric bead array. Lines represent group mean. Mann-Whitney U test was performed for statistical analysis. NS = not significant. Data from one or two pooled independent experiments.

Figure 3. Inflammatory monocytes are required for organized pyogranuloma formation and restriction of Yersinia within intestinal tissue

(**a**) Quantification of total number of intestinal lesions at day 5 post infection. Each circle represents one mouse. Lines represent median. Pooled data from four independent experiments.

(**b**) H&E-stained paraffin-embedded longitudinal small intestinal sections from *Yp*-infected mice. Scale bars = 250 μm (top) and 50 μm (bottom). Representative images of two independent experiments.

(**c**) Histopathological scores of small intestinal tissue from uninfected and *Yp*-infected mice. Each mouse was given a score between 0-4 (healthy-severe) for signs of pathology. Each circle represents one mouse. Lines represent median. Pooled data from two independent experiments.

(**d**) Bacterial burdens in small-intestinal PG- and PG+ tissue. Each circle represents the mean *Yp*-CFU of 3-5 pooled punch biopsies from one mouse. Lines represent geometric mean. Pooled data from six independent experiments.

(**e**) Total number of CD45+ cells, monocytes, macrophages, and neutrophils in small intestinal uninf, PG-, and PG+ tissue. Each circle represents the mean of 3-10 pooled punch biopsies from one mouse. Lines represent median. Pooled data from five independent experiments.

(**f**) Fluorescently labeled whole-mount small intestinal PG+ tissue from *Yp*-infected *Ccr2gfp/+* (top) and *Ccr2gfp/gfp* (bottom) mice. Scale bars = 100 μm. Representative images of two independent experiments.

(**g**) PG+ neutrophil CD11b expression was measured by flow cytometry. Lines represent median. Representative of four independent experiments.

(**h**) Cytokine levels in homogenates of tissue punch biopsies were measured by cytometric bead array. Lines represent median. Pooled data from three independent experiments.

Mann-Whitney U test was performed for all statistical analyses. NS = not significant.

Figure 4. Inflammatory monocytes are required for systemic control of *Yersinia* and host survival

(**a**) Bacterial burdens in indicated organs at day 5 post infection. Each circle represents one mouse. Lines represent geometric mean. Pooled data from four independent experiments.

(**b**) H&E-stained paraffin-embedded longitudinal spleen sections from WT and *Ccr2$^{gfp/gfp}$* mice. Scale bars = 500 µm (top) and 50 µm (top). Representative images of two independent experiments.

(**c**) Histopathological scores of spleens from uninfected and *Yp*-infected mice. Each mouse was given a score between 0-4 (healthy-severe) for signs of pathology. Each circle represents one mouse. Lines represent median. Pooled data from two independent experiments.

(**d**) Survival of infected WT (n=16) and *Ccr2$^{gfp/gfp}$* (n=13) mice. Pooled data from two independent experiments.

(**e**) Survival of infected littermate wild-type *Ccr2$^{+/+}$* (n=20), heterozygous *Ccr2$^{gfp/+}$* (n=19)*,* and homozygous *Ccr2$^{gfp/gfp}$* (n=15) mice. Pooled data from three independent experiments.

Statistical analyses by (a, c) Mann-Whitney U test and (d-e) Mantel-Cox test. NS = not significant.

Figure 5. Intestinal pyogranulomas form in response to *Yersinia* virulence

(**a**) Quantification of total number of intestinal lesions upon infection. Each symbol represents one mouse. Lines represent median. Pooled data from 2-3 independent experiments.

(**b**) Bacterial burdens in small-intestinal PG- (left) and PG+ (right) tissue. Each symbol represents one mouse. Lines represent geometric mean. Pooled data from 2-3 independent experiments.

(**c**) Bacterial burdens in indicated organs. Each symbol represents one mouse. Lines represent geometric mean. Pooled data from 2-3 independent experiments.

(**d**) Frequency of monocytes (left) and neutrophils (right) in small-intestinal uninf, PG- and PG+ tissue. Each symbol represents one mouse. Lines represent median. Pooled data from two independent experiments.

(**e**) Frequency of monocytes (left) and neutrophils (right) in MLN. Each symbol represents one mouse. Lines represent median. Representative data from two independent experiments.

Mann-Whitney U test was performed for all statistical analyses.

Figure 6. *Yp* restriction by CCR2-deficient mice is restored in absence of YopH activity

(**a**) H&E-stained paraffin-embedded transverse small intestinal sections from WT (top) and *Ccr2$^{gfp/gfp}$* (bottom) mice infected with WT (left) or YopH^{R409A} (right) *Yp* at day 5 post infection. Scale bars = 100 μm. Representative images of two independent experiments.

(**b**) Bacterial burdens in small-intestinal PG- and PG+ tissue at day 5 post infection. Each symbol represents one mouse. Lines represent geometric mean. Pooled data from two independent experiments.

(**c**) Bacterial burdens in indicated organs at day 5 post infection. Each symbol represents one mouse. Lines represent geometric mean. Pooled data from two independent experiments.

Mann-Whitney U test was performed for all statistical analyses. NS = not significant.

Extended Data Figure 1. Intestinal pyogranulomas from acutely following oral *Yersinia* infection

(**a**) Frequency distribution (left) of lesions along the intestinal tract, with graphical key of anatomical segments (right). Each colored bar represents the mean frequency of lesions in a given segment (n = 20 mice). Only mice with >9 total pyogranulomas were included. Pooled data from three independent experiments.

(**b**) Flow cytometry plots displaying the gating strategy employed to identify eosinophils, dendritic cells, B cells and T cells in small-intestinal tissue. Representative images of four independent experiments.

(**c**) Frequency (top) and total number (bottom) of eosinophils, dendritic cells, B cells, CD4+ T cells and CD8+ T cells (left to right) in small-intestinal uninf, PG- and PG+ tissue. Each circle represents the mean of 3-10 pooled punch biopsies (ø = 2 mm) from one mouse. Lines represent median. Pooled data from four independent experiments.

Mann-Whitney U test was performed for all statistical analyses. NS = not significant.

Extended Data Figure 2. Non-pyogranuloma tissue does not undergo a pro-inflammatory response during intestinal *Yp* infection
(**a**) Heatmap of top 30 significantly upregulated genes in PG- compared to uninfected samples in descending order by fold change. False discovery rate < 0.05 using Benjamini-Hochberg procedure.
(**b**) Gene ontology analysis of top 30 upregulated genes by fold change only in PG- compared to uninfected samples. Dotted line denotes p = 0.05.

Extended Data Figure 3. *Ccr2*[-/-] mice have increased susceptibility to oral *Yersinia* infection.
(**a**) Quantification of total number of intestinal lesions at day 3 post infection. Each circle represents one mouse. Line represents median. Pooled data from two independent experiments.
(**b**) Bacterial burdens in small intestinal PG- and PG+ tissue at day 3 post infection. Each circle represents the mean *Yp*-CFU of 3-5 pooled punch biopsies from one mouse. Lines represent geometric mean. Pooled data from four independent experiments.
(**c**) Frequency of neutrophils in MLN and spleen at day 5 post infection. Each circle represents the mean of 3-10 pooled punch biopsies from one mouse. Lines represent median. Pooled data from two independent experiments.
(**d**) MLN neutrophil CD11b expression was measured by flow cytometry. Lines represent median. Representative of two independent experiments.
(**e**) Cytokine levels in homogenates of tissue punch biopsies at day 5 post infection were measured by cytometric bead array. Lines represent median. Pooled data from three independent experiments.
(**f**) Total number of CD4+ and CD8+ T cells in small intestinal PG+ tissue. Each circle represents the mean of 3-10 pooled punch biopsies from one mouse. Lines represent median. Pooled data from 4 independent experiments.
(**g**) Bacterial burdens in indicated organs at day 3 post infection. Each circle represents one mouse. Lines represent geometric mean. Pooled data from four independent experiments.
(**h**) Survival of infected mice. Pooled data from two independent experiments.
Statistical analyses by (a-g) Mann-Whitney U test or (h) Mantel-Cox test. NS = not significant.

Extended Data Figure 4. Monocytes are dispensable for restriction of YopH mutant *Yp*
(**a**) Quantification of total number of intestinal lesions upon infection. Each symbol represents one mouse. Lines represent median. Pooled data from 2 independent experiments.
(**b**) Bacterial burdens in small-intestinal PG- and PG+ tissue at day 3 post infection. Each symbol represents one mouse. Lines represent geometric mean. Pooled data from two independent experiments.
(**c**) Bacterial burdens in indicated organs at day 3 post infection. Each symbol represents one mouse. Lines represent geometric mean. Pooled data from two independent experiments.
(**d**) Survival of *Yp*-infected mice. Pooled data from two independent experiments.
Statistical analyses by (a, b) Mann-Whitney U test and (c) Mantel-Cox test.