Complex genotype-phenotype relationships shape the response to treatment of Down Syndrome Childhood Acute Lymphoblastic Leukaemia

Christoph Lutz[1,7,8*], Virginia A. Turati[1*], Ruth Clifford[7], Petter S. Woll[2,3,10], Thomas Stiehl[4], Anders Castor[5], Sally A. Clark[2,3], Helen Ferry[2,3], Veronica Buckle[2] Andreas Trumpp[6], Anthony Ho[7], Anna Marciniak-Czochra[4], Javier Herrero[1], Anna Schuh[9], Sten-Eirik W. Jacobsen[2,3,10] and Tariq Enver[1,11]

[1]University College London, Cancer Institute, London WC1E 6BT, UK

[2]MRC Molecular Haematology Unit, MRC Weatherall Institute of Molecular Medicine,

[3]University of Oxford, Oxford, OX3 9DS, UK

[3]Haematopoietic Stem Cell Biology, MRC Weatherall Institute of Molecular Medicine, University of Oxford, Oxford, OX3 9DS, UK

[4]Institute of Applied Mathematics Mathematikon, Heidelberg University, Germany

[5] Department of Paediatric Haematology/Oncology, Skane University Hospital, Sweden

[6] Deutsches Krebsforschungszentrum Heidelebrg (DKFZ), Germany

[7] Department of Internal Medicine V, Heidelberg University, Germany

[8]Praxis für Onkologie, Neversstrasse 5, 56077 Koblenz, Germany

[9]Department of Haematology, Oxford Radcliffe Hospitals NHS Trust, Oxford OX3 9DU, UK

[10]Center for Hematology and Regenerative Medicine, Karolinska Institutet and Karolinska University Hospital Huddinge, Stockholm, Sweden

[11] Division of Molecular Medicine and Gene Therapy, Lund University, A12, 221 84 Lund Lund, Sweden

* These authors contributed equally to this work

Correspondence to:

Tariq Enver PhD

UCL Cancer Institute

Stem Cell Laboratory

Paul O'Gorman Building

72 Huntley Street

London WC1E 6BT

email: t.enver@cancer.ucl.co.uk

phone: +44 (0)20 7679

Fax: +44 (0)20 7679 6914

Abstract

Extensive genetic and epigenetic variegation has been demonstrated in many malignancies. Importantly, their interplay has the potential to contribute to disease progression and treatment resistance. To shed light on the complex relationships between these different sources of intra-tumour heterogeneity, we explored their relative contributions to the evolutionary dynamics of Acute Lymphoblastic Leukemia (ALL) in children with Down syndrome, which has particularly poor prognosis. We quantified the tumour propagating potential of genetically distinct sub-clones using serial transplantation assays and SNP-arrays. While most leukaemias were characterized by a single dominant subclone, others were highly heterogeneous. Importantly, we provide clear and direct evidence that genotypes and phenotypes with functional relevance to leukemic progression and treatment resistance can co-segregate within the disease. Hence, individual genetic lesions can be restricted to well-defined cell immunophenotypes, corresponding to different stages of the leukemic differentiation hierarchy and varied proliferation potentials. As a result of this difference in fitness, which can be accurately quantified via competitive transplantation assays, matching diagnostic, post-treatment, and relapse leukemias can be dominated by different genotypes, including pre-leukemic clones persisting throughout the disease progression and treatment. Intriguingly, plasticity also appears to be a temporally defined property that can segregate with genotype. These results suggest that Down Syndrome ALL should be viewed as a complex matrix of cells exhibiting genetic and epigenetic heterogeneity that foster extensive clonal evolution and competition. Therapeutic intervention reshapes this 'eco-system' and may provide the right conditions for the preferential expansion of selected compartments and subsequently relapse.

Introduction:

The existence of intra-tumour genetic heterogeneity, both genetic and epigenetic, has been documented in both solid and haematological malignancies[1–3]. In studies of childhood acute lymphoblastic leukaemia, we and others previously used multicolour-FISH and whole genome sequencing to show that these tumours are clonally variegated, and to demonstrate through serial transplantation assays, that leukaemia tumour propagating cells (TPC) are also genetically heterogeneous[4,5]. This finding was supported by our later genetic analysis of immunophenotypically distinct ALL diagnostic subpopulations which demonstrated similar distribution of subclones within a disease-specific primitive B-cell compartment (StemB, $34^+38^-19^+$) and the disease bulk in most, but, crucially, not all leukaemias[6]. Such

experiments relied on high-depth bulk whole-genome sequencing followed by single-cell qPCR and, together with our earlier work, suggested that genotype and phenotypes related to differentiation stage and cell-cycle state might not necessarily co-segregate within the disease. Importantly, as chemotherapy-induced selection acts on these same phenotypic traits[6–8], this lack of segregation also underpinned the survival of most genotypes to the early stages of childhood ALL treatment (namely, induction chemotherapy – where maximum cytotoxicity is achieved); as we demonstrated using a xenograft model of transplantation and treatment of primary leukaemias[6].

Large-scale comparisons of matching diagnostic and relapse samples from paediatric ALL patients, have nonetheless previously reported that in some patients a subset of diagnostic subclones, or of their later evolutionary descendants, can be selectively enriched at the time of disease recurrence[9,10]. Assuming that these clones did not become dominant as a result of stochastic selection, this finding opens to the possibility that fitness-based genotypic selection might potentially occur on a longer timescale than that of induction chemotherapy. It is not unreasonable to speculate that in a treatment protocol that lasts up to four years even subtle differences in fitness might eventually lead to clonal dominance. Another possibility is that in certain settings, genotype, and treatment-associated phenotypes do, in fact, segregate (either already at diagnosis or later because of further genomic evolution). In support of this latter hypothesis, our analysis of $34^{+}38^{-}19^{+}$ (StemB) ALL cells indicated that in a small proportion of leukaemias (1 out of 5 analysed), specific genotypes might indeed be preferentially enriched within developmentally more primitive and quiescent disease compartments[6]. Furthermore, recent evidence suggests thar mutations induced by chemotherapy treatment itself (particularly those targeting p53) might be responsible for - post-diagnosis – generating new, fitter subclones, and inducing drug-resistance; therefore, leading to relapse[11].

To shed light on the complex genotype-phenotype relationship that shape ALL, we here conduct a comprehensive investigation into the relative contribution of different sources of intratumor heterogeneity, namely genetic makeup and epigenetically determined properties - particularly cell cycle state, TPC activity and differentiation stage - to the disease initiation, sustenance, and treatment resistance. In doing so we focused our attention Down's syndrome associated ALL as this rarer disease subtype has so far remained poorly characterised while being associated with considerably worse prognosis than ALL in children without Down syndrome[12,13].

Altogether our results indicate that ALL should be viewed as a complex matrix of cells
exhibiting genetic and epigenetic heterogeneity. Our work provides evidence for, in some
patients, diversification in functional properties amongst immunophenotypically-distinct
leukemic subpopulations. In these setting, those rare leukemic cells which remain after
induction chemotherapy, and primarily account for minimal residual disease (MRD), are
highly enriched for the stem/B phenotype; suggesting that the more primitive and less
proliferative leukemic compartments likely constitute the pool of cells from which relapse
arises. Crucially, in this setting where genotype and resistance-associated phenotypes co-
segregate, bottleneck selection at phenotype level is accompanied by evidence of a clonal
sweep.
118

Methods:
Patient samples
Diagnostic and follow-up ALL bone marrow (BM) samples were obtained after informed
consent and the approval of the relevant research ethics committee from patients at the
Paediatrics Haematology Unit, Lund University Hospital, Sweden. Samples from 4 children
with Down's ALL were analyzed. For further patient characteristics see Table 1.
125

Cell separation, phenotyping, and sorting
Total mononuclear cells (MNCs) were isolated by ficoll gradient centrifugation and directly
cryopreserved in DMSO for later use. In some cases, $CD34^+$ cells were enriched by
magnetic bead separation (StemCell Technologies or Miltenyi). After thawing, dead cells
were evaluated and excluded by FACS after staining with DAPI. CD34-enriched cells or
MNCs from BM were stained with anti-CD19 PE (BD-Pharmingen), CD34 FITC (BD-
Pharmingen) and CD38 APC (BD-Pharmingen). Cells were analyzed in staining buffer
containing DAPI at 0·1µg/ml, for live cell analysis. I.) HSC ($34^+38^-19^-$); II.) Stem/B (34^+38^-
19^+) III.) Progenitor ($34^+38^+19^-$) IV.) ProB ($34^+38^+19^+$) and V.) PreB/Mature B-cells (34^-19^+)
were purified by flow cytometry (FACS Aria, BD-Pharmingen). Data acquisition and analysis
were done with CellQuest (Beckon Dickinson) or FlowJo (Tree star) software.
137

Cell cycle analysis of leukaemia subpopulations
Cells were stained with CD19-PECy5 (BioLegend), then fixed and permeabilized with 1.6%
paraformaldehyde and 90% ice-cold methanol. This was followed by staining with CD34-
APC (BD), CD38-PETxR (Invitrogen) and Ki67-FITC (Becton-Dickinson). For DNA content
analysis cells were incubated with 0·5µg/mL DAPI. Stained cells were analyzed by excitation

of DAPI with a violet laser on a FACS LSRII SORP (Becton-Dickinson) at the University of
Oxford (UK).

FISH analysis
Cells were fixed on slides in methanol/acetic acid fixative (3:1 vol/vol) and then hybridized
with BCR-ABL1 FISH probes using the LSI BCR/ABL1 Dual Color Dual Fusion Translocation
probes (Abbot). The slides were analyzed with an Olympus BX51 microscope equipped with
epifluorescence and a triple band pass filter. Images were captured by using a Sensys
charge-coupled device camera (Photometrics, Tucson, AZ) and MacProbe software (Applied
Imaging, Newcastle upon Tyne, U.K.). When possible, and unless otherwise specified at
least 100-150 nuclei were analyzed in each sample

SNP-Array
SNP-Arrays we performed as previously describe in Schuh et al., 2012[14]

NSG-mouse transplantation assay
Primary childhood ALL cells were transplanted into 8-12 weeks old NOD/SCID IL2Rγ[hull]
(NSG) sub-lethally irradiated mice (either females or males) via intramedullary injection. To
minimize possible adverse effects of sublethal irradiation, mice were administered acid water
for a week prior to the procedure, and Baytril (resuspended at 25.5 mg/kg in the drinking
water) for the 2 weeks following it. Sub-lethal irradiation was achieved with a single dose of
375 cGy. Unless otherwise stated, each mouse received 2×10^5 primary leukemia cells
resuspended in 40μl PBS 0.5% FBS. In the case of secondary limiting dilution assays, a
specified equal dose of treated and control leukemic cells harvested from the BM of primary
recipients was injected. Twelve-week post-injection mice were sampled by bone marrow
aspiration and the percentage of human engraftment was evaluated by flow cytometry
(hCD45/(hCD45$^+$ + mCD45$^+$)). At the same time, human cells were also FACS sorted for
downstream applications. Mice displaying at least 70% human engraftment were then
randomly assigned to either control or treatment groups (see below for details on the
treatment protocol). At the end of the treatment window all mice were culled, and their tibias,
femurs, pelvises, spleen, and brain harvested, and cells were then stained for FACS sorting.

Modeling of clonal evolution
See supplementary materials.

Small cell numbers RNA sequencing

Equivalent cell numbers (400 cells per sample) were flow sorted directly into 800µl Trizol
reagent (Invitrogen) and snap frozen in dry ice (long term storage at -80C). At the time of
extraction, the samples were thawed at RT and 160ul of chloroform was added to each.
Following a centrifugation step the RNA was isolated from the aqueous phase and
precipitated through the addition of equal volumes of isopropanol supplemented with 20µg
linear polyacrylamide. Samples were washed twice in 80% ethanol (first wash over night at
4°C, second wash 5 minutes at RT). RNA pellets were resuspended in 3-15µl of
diethylpyrocarbonate treated water (DEPC). RNA was then quantified by loading of 0.5-1ul
on an Agilent Bionalyser RNA 6,000 pico chip. Where possible equivalent amounts of total
RNA (100pg) from all samples were used for first strand synthesis with the SmartERv3 kit
(Takara Clontech) followed by 15-18 cycles of amplification (according to manufacturers'
instruction). cDNA was purified on Agencourt AMPureXP magnetic beads, washed twice
with fresh 80% ethanol and eluted in 17µl elution buffer. 1µl cDNA was quantified with Qubit
dsDNA HS (Molecular Probes) and checked on an Agilent Bioanalyser high sensitivity DNA
chip. Sequencing libraries were produced from 150pg input cDNA using Illumina Nextera XT
library preparation kit. A 1:4 miniaturized version of the protocol was adopted (see "Fluidigm
Single-Cell cDNA Libraries for mRNA sequencing", PN_100-7168_L1). Tagmentation time
was 5mins, followed by 12 cycles of amplification using Illumina XT 24 or 96 index primer kit.
Libraries were then pooled (1-2ul per sample depending on the total number of samples)
and purified with equal volumes (1:1) of Agencourt AMPureXP magnetic beads. Final elution
was in 66-144ul of resuspension buffer (depending on the total number of pooled samples).
Libraries were checked on an Agilent Bioanalyser high sensitivity DNA chip (size range 150-
2000bp) and quantified by Qubit dsDNA HS (Molecular Probes). Libraries were sequenced
on Illumina® NextSeq 500 using 150bp paired end kits as per manufacturer's instructions.
203

Bioinformatics

Sequencing data was assessed to detect sequencing failures using FASTQC and lower
quality reads were filtered or trimmed using *TrimGalore*
(https://github.com/FelixKrueger/TrimGalore). Outlier samples containing low sequencing
coverage or high duplication rates were discarded.

Processing of bulk RNAseq

Bulk RNAseq samples were mapped to the human reference GRCh38 using *tophat2*
(https://ccb.jhu.edu/software/tophat/index.shtml). Analyses were performed within the R
statistical computing framework, version 3.5 using packages from *BioConductor version 3.7*
(https://Bioconductor.org). Data was combined into a per-gene count matrix using
featureCounts from the subread package. The *DEseq2* BioConductor package was used for
outlier detection, normalization and differential gene expression analyses. All downstream
analyses used Rlog transformed data.

Data availability

SNP-array and RNAseq data are available at EGA under accession number
EGAS00001005968.

Results

We here explored the role of intra-tumour heterogeneity in the progression and response to
treatment of Down's associated ALL. We initially focused on genetic variegation and
described the phylogenetic hierarchies of diagnostic material from 4 patients. Serial
transplantation of bulk ALL samples into immune-deficient NSG-mice was concomitantly
utilized to evaluate the tumour propagating cell activity (TPC) and dynamic behaviours of the
different subclones identified. SNP-arrays were performed on diagnostic specimens as well
as cells isolated from the BM of primary and secondary transplanted mice (see schemata in
Figure 1A). In all patients, but one, a single dominant clone was detected both at diagnosis
and in all primary and secondary transplanted mice (Figure 1B). In patient 2, instead, a more
complex picture was observed, with several subclones coexisting within the tumour and
contributing to a different extent to engraftment across recipients. In this patient we detected
two deletions, on chromosomes 9 and 14 respectively, as well as an amplification on
chromosome X (Figure 1C). Considering the mutational mosaicism, we observed in some of
the engrafted mice, these observations allowed re-construction of a simple clonal
phylogenetic architecture (Figure 1D, and Supp Material for Mosaicism data). However, a
higher-depth genetic analysis of chromosome 9 suggested a far more complicated clonal
landscape. A heterozygous chromosome 9 deletion was detected across engrafted animals
with different degrees of homozygous involvement: to the extreme of copy neutral loss of
heterozygosity (cnLOH) (Figure 2A). A simple explanation to this observation is that pre-
existing sub-clones present at diagnosis at sub-threshold levels, were expanded through
secondary transplantation until reaching detection levels. Alternatively, continuous
broadening of the chromosome 9 deletion could have occurred over the short time frame of
the serial transplantation assays. This latter hypothesis would be indicative of active ongoing

clonal evolution. Based on the mosaicism data, and assuming no new mutations developed during transplantation, we were able to reconstruct a more comprehensive phylogenetic tree. To do so, we developed a custom-made mathematical model which we used to calculate all the theoretically possible diagnostic clone-combinations supported by our SNPs data analysis (Supplementary Table 1). In total 15-29 diagnostic clones were necessary to explain the mutational read-out from the xenografts. Of note, since SNP-arrays are limited in both genomic resolution (i.e., deletion size) and sensitivity to low abundance copy number variants (CNV) our model would if anything underestimate the number of clones (Figure 2B and supplementary methods). Nonetheless, this finding clearly illustrates the high clonal complexity of ALL and is line with our earlier single-cell WGS based characterization of genetic heterogeneity in non-Down Syndrome ALL, which also identified branching clonal architectures with up to 19 descendant subclones[6].

We subsequently sought to shed light on the phenotype-genotype relationship that defines the properties of individual leukemic subclones, by performing SNP analysis on bulk tumour cells and flow-sorted immunophenotypically-distinct ALL subpopulations from three Down's syndrome ALL patients both before and after xenotransplantation (see schemata in Figure 3A). Analysis of samples from patients 2 and 3 (supplementary Figure 1) revealed identical clonal architectures in diagnostic bulk, stem/B and proB-like cells for both patients (Figure 3BC). All samples analysed from patient 3 confirmed the existence of a single dominant leukemic clone (Figure 3C and supplementary table 1). In addition to the founding chromosome 21 amplification, a third patient (pt.4) we examined harboured a BCR-ABL translocation. FISH analysis showed that in this leukemia the fusion gene marked proB- and preB-like sub-compartments, but not cells with the more primitive and disease-specific stem/B ($34^+38^-19^+$) phenotype[15] (Figure 4A). Notably, in serial transplantation experiments flow-sorted stem/B cells only gave rise to grafts with stem/B like cells but not more differentiated (proB-preB) B-cell phenotypes. These stem/B like grafts remained BCR-ABL-negative. Similarly, proB cells derived grafts only contained proB and preB cells, all of which retained the BCR-ABL-fusion. Collectively these data suggest lack of phenotypic plasticity amongst distinct diagnostic cells from this patient.

We next compared the relative fitness of the two subpopulations (stem/B and proB) in a competitive transplantation setting facilitated by (i) the presence of the BCR-ABL translocation as a clonotypic marker and (ii) the non-interchangeability of their immunophenotypes. We transplanted NSG-mice with different ratios of stem/B and proB-like ALL cells and read out engraftment. In all cases the BCR-ABL$^+$ ProB-like cells displayed

increased engraftment fitness, most clearly exemplified in the setting where proB cells

represented only 10% of the initial inoculum and yet constituted 95% of the resultant

leukaemia (Figure 4B). Reasoning that enhanced fitness may stem from a proliferative

advantage we next analysed the cell cycle status of engrafted cells. In all engrafted

recipients proB cells displayed a more active cycling status compared to their stem/B

counterparts, which had a significantly larger proportion of dormant (G_0) cells (Figure 4C and

Supplementary Figure 2).

We subsequently speculated that the different cycling properties of these compartments

could be expected to affect their relative sensitivity to therapy. To experimentally determine

how these different ALL sub-compartments behaved during therapy, we analysed leukemic

subpopulations from serially collected diagnosis, remission, and relapse specimens. ProB-

like ALL cells were indeed preferentially eradicated during chemotherapy. Indeed, while both

populations were shrank by treatment to similar frequency (0.01%), ProB cells initially

represented approximately 70% of leukaemic cells detected at diagnosis while stem/B-like

ALL cells were only 8% (Figure 5A). After induction chemotherapy extremely few proB-like

cells remained, none of which carried the BCR-ABL translocation. While the behaviour of

these subpopulations during treatment contrasts starkly with their fitness as assessed by

xenotransplantation, it is entirely consistent with the known relative vulnerability of cycling

versus dormant cells to chemotherapy. Similarly, to what has previously been observed in

the context of bacteria and antibiotics resistance, these data also support the idea of the

existence of a trade-off on phenotypic traits between environments with and without drug[16].

The reduced competitive ability of the less proliferative species (StemB cells) in the absence

of treatment hence becomes an advantage in the presence of chemotherapy, while the

highly proliferative ProB cells pay a cost to their initial fitness.

Interestingly, the relapse sample was populated by both stemB and proB-like cells (Figure

5A). However, in contrast to the situation at diagnosis leukemic proB cells present at relapse

were BCR-ABL-negative and both relapse stemB and proB cells showed phenotypic

plasticity upon serial transplantation (Figure 5B). This data is somewhat counterintuitive in

that it argues against the simplest origins of relapse from the predominant stemB or proB

clones present at diagnosis. The stem/B compartment at presentation lacked phenotypic

plasticity in that it was unable to regenerate proB- or preB-like cells. Since the proB cells at

relapse were BCR-ABL-negative, relapse could also not have been seeded by the

predominant diagnostic proB-like compartment that was positive for BCR-ABL.

Taking advantage of the xenograft system we further investigated the phenotypic properties
of diagnostic and relapse cells at the molecular level by performing small cell number bulk
RNA sequencing (RNAseq) on cells harvested from either primary and or secondary
xenografts. We hence characterised six distinct cell populations i) diagnostic stemB cells
from primary xenografts, ii) diagnostic proB cells from primary xenografts iii) relapse stemB
cells harvested from secondary xenografts injected with stemB cells isolated from primary
xenografts iv) relapse proB cells harvested from secondary xenografts injected with stemB
cells isolated from primary xenografts, v) relapse proB cells harvested from secondary
xenografts injected with proB cells isolated from primary xenografts, and vi) relapse stemB
cells harvested from secondary xenografts injected with proB cells isolated from primary
xenografts. Principal Component Analysis (PCA) showed clustering of samples based on
disease stage. Interestingly, diagnostic cells also clustered separately based on
differentiation state, while all relapse populations clustered closely together (Figure 5C). This
finding resonates well with our observation that in this patient leukaemia, relapse but not
diagnostic cells displayed high cell plasticity. While the PC1 component of the PC captured
most of the sample variance (47.15), PC2 clearly separated diagnostic proB cells and all
relapse subpopulations from diagnostic stemB cells. Pathways analysis on the genes driving
this PC component revealed enrichment for TNF-alpha via NF-KB signalling cascade, which
has previously been implicated in regulation of cell survival, proliferation and stemness[17].
Gene set enrichment analysis (GSEA) using publicly available gene expression
signatures[18,19] also validated our FACS analysis results in that compared to diagnostic proB
cells and relapse cells, diagnostic stemB cells downregulated cell proliferation associated
genes and expressed higher levels of markers typical of more immature stages of the
haematopoietic and B-cell differentiation hierarchy (such as HSC and MLP cells) (Figure 5C-
D and Supplementary Figure 3A-B). Most B-lineage cells from ALL patients are thought to
be arrested at some point of the transition from the proB to preB stage, therefore broadly
allowing the classification of ALL leukaemias into two distinct subtypes based on preBCR
function. Those arrested at the preBI stage rely upon IL7R/STAT5 signaling (preBCR$^-$), while
those arrested at preBII stage signal by the preBCR (preBCR$^+$)[20]. The frequency of pre-
BCR$^+$ or pre-BCR$^-$ cases has also been suggested to associate with different cytogenetic
subtypes of childhood ALL. For example, the majority of BCR-ABL1$^+$ leukaemias tend to
display constitutive active cytokine signalling via activation of STAT5 and repression of
BCL6. Herein, however, we found that these two genetically and phenotypically distinct
disease subsets can coexist within the same leukaemia. Interestingly, contrary to
expectations, diagnostic proB cells, which carried the BCR-ABL translocation and expressed
markers associated with later stages of the B-cell differentiation (Figure 5C), displayed

stronger expression of the preBCR receptor components and signalling cascade, while the

BCR-ABL⁻ diagnostic stemB cells preferentially upregulated various genes involved in the

STAT5 signalling pathway (Supplementary Figure 3D).

We subsequently used SNP arrays to genetically characterise different phenotypic

compartments at presentation, relapse and after serial transplantation and track the

phylogenetic origins of the relapse propagating clones (Figure 6). SNP analysis of the

relapse samples indicated that it lacked several of the distinctive genetic abnormalities which

characterised either the dominant stemB compartment at diagnosis or the dominant proB.

However, the relapse hierarchy shared with the diagnostic compartments deletions on both

chromosomes 12 and 14. Of note, diagnosis and relapse clones also shared a common VDJ

rearrangement (data not shown).

Altogether, this data implies the survival over treatment of a common evolutionary ancestor

not detectable at diagnosis which gave rise to the relapse hierarchy through further

diversification. Accordingly, analysis of a bulk remission sample identified mosaic deletion of

chromosome 14 in the absence of del12 and any other obvious genetic abnormality.

Therefore, suggesting that this mosaic deletion is not a germline event and, likely, the

earliest genomic alteration (also preceding del12). To understand the basis of the mosaicism

observed in the remission sample, we separated CD3+ and CD3- cells and performed SNP

analysis. The CD3- compartment contained no evidence of the 14del. In contrast the CD3+

cells showed a complex pattern of chromosome 14 deletions indicative of a progressive

multi-step evolutionary process (Supplementary table 2). We next traced these 14del

variants back to the diagnostic and relapse stemB and proB compartments, as well as the

CD3+ compartment isolated at diagnosis. This spectrum of 14 deletions provides a branched

framework for subsequent evolution of stemB and proB clones seen at diagnosis and

relapse. Thus, using chromosomal deletions as markers of lineal ancestry we have here

reconstructed the evolutionary history of a Down's ALL from the predisposing event (Trisomy

21), through the acquisition of initiating lesions giving rise to a preleukemic hierarchy, into

further transformation marked by the onset of frank leukaemia. We also monitored the

response of this complex leukemic hierarchy to chemotherapy through the analysis of

matching remission and relapse specimens. Beyond the genetic considerations our work

emphasizes the role of cellular state primarily at the level of dormancy and cell cycle

indicating vulnerability to chemotherapy.

Discussion

Many studies have explored the extent and impact of different, individually assessed,
sources of intratumor heterogeneity on the evolution of tumours (particularly under
treatment). Our work, however, argues that cancer should be viewed as a complex matrix of
genetic and epigenetic heterogeneity such as cellular phenotype and cell cycle properties
which, altogether rather than in isolation, provide a substrate for tumour evolution during
disease progression in response to the selective impetus of therapeutic intervention.

The idea that epigenetically determined cell states, contribute to functional heterogeneity
amongst tumour cells by controlling fundamental cellular properties such as, for example,
cell identity, phenotype and cell-cycle stage is paradigmatically exemplified by the cancer
stem cell (CSC) hypothesis; which postulates that cells within a tumour are organized in a
hierarchical fashion reflecting lineage relationships and tumorigenic potential, and that the
maintenance of cancer clones is uniquely dependent on the most primitive CSCs with self-
renewal capacity[21–23]. This notion, which was pioneered in the context of blood malignancies,
remains however extensively debated. While phenotypic heterogeneity - primarily at the level
of immune-phenotype – has been widely observed in ALL, disparate data exist concerning
the existence of LSCs within this disease. Earlier work by Rehe et al., showed that many
cellular fractions contain LSC-potential at a similar frequency and that the majority of - if not
all - cells may be able to propagate the disease[24]. Importantly, the same study also found
evidence for phenotypic plasticity within different subpopulations upon serial transplantation
in immune-deficient mice. Distinct leukaemia stem cells (LSCs) have nonetheless been
identified in acute myeloid leukaemia (AML), where functional evidence for their existence
has been provided[25–28]. However, even in this context, their exact phenotype remains
elusive, with more recent studies uncovering LSC activity across different cellular subsets[29].
We here provide evidence that although different Down's ALL disease compartments can all
contribute to leukaemia propagation *in vivo,* these same compartments display largely
distinct fitness; both in the presence and in the absence of chemotherapy. In this scenario,
more primitive stem-like compartments are outcompeted by highly proliferative early-B cell
progenitors in the absence of treatment. However, this selective advantage is reversed in the
presence of chemotherapy, which selects for quiescent cells. While the genetically
heterogenous nature of childhood ALL has been demonstrated by multiple earlier studies,
our current work provides the first direct evidence for genetic diversity of cancer propagating
cells in patients with Down's syndrome ALL. Our SNP array analysis of serially transplanted
ALLs shows that leukemic cells undergo a process of gradual branching clonal evolution
demonstrated by the coexistence within a leukaemia of multiple subclones bearing
alterations of the same genomic locus characterised by distinct breakpoints.

Our earlier work showed that in many ALL leukemias genotype and phenotypes with
relevance to chemoresistance do not segregate. As a result, induction chemotherapy does
not select at the level of genotype. We here identified and studied the evolutionary history of
an unusual case of BCR-ABL+ Down's ALL in which genotype and phenotype do instead
clearly segregate. In this patient's leukaemia the stem/B cells were fusion-gene-negative
while proB/pre-B cells were fusion-gene-positive. This case allowed us to reconstruct in
detail the complex genotype-phenotype relationship characterizing different leukemic cell
fractions labelled by both distinct genetic makeup and fitness. We hence observed that
highly quiescent and developmentally more primitive cells preferentially escaped treatment.
As these cells carried only a subset of diagnostic genetic alterations, the relapse disease
was genetically divergent from the diagnostic specimen. Reconstruction of the leukaemia
phylogenetic tree demonstrated that the dominant population at relapse originated from a
rare pre-leukemic clone. This was likely present at diagnosis below detection level and, in
line with its likely low proliferative activity, did not extensively expand upon transplantation.

We first demonstrated the existence of pre-leukemic cells in ALL in studies of mono-
chorionic twins of whom one was diagnosed with TEL-AML1-positive ALL[15,30]. Blood spot
studies from Guthrie-cards further showed that these pre-leukemic cells can precede overt
leukaemia by up to 12 years and are also detectable in patients that never go on to overt
disease[31,32]. A case report of a 7-year old boy diagnosed with ALL with a mixed lympho-
myeloid phenotype previously suggested that BCR-ABL+ non-Down leukaemic cells could
reactivate to drive disease recurrence more than two decades from initial diagnosis[33]. Our
new data add to our understanding of this elusive population by showing that Down
Syndrome ALL chr21+ cells, which have not yet acquired the BCR-ABL translocation
labelling the more proliferative proB compartment disease, can preferentially survive
chemotherapy and, therefore, represent a reservoir for relapse initiation. Whether these cells
are quiescent principally as a function of their unique developmental origin[34] or rather of their
genotype[35] or a combination of both remains to be investigated and would be important to
understand if new therapeutic interventions targeting these cells are to be designed. This
notwithstanding, pre-leukemic clones, even when rare, should clearly be considered part of
the fully transformed leukaemia landscape and may upon selection by therapy provide an
additional cellular source for subsequent relapse.

Acknowledgments

We are grateful for the support of Leukaemia and Lymphoma Research (TE; Specialist Programme Grant), Cancerfonden (TE; Project grants CAN 2013/712 and 20 1334 PjF) , MRC Molecular Haematology Unit core award (SEJ; MC_UU_12009/5), Blood Cancer UK (SEJ), Swedish Childhood Cancer Fund (SEJ), the National Institute for Health Research Biomedical Research Centre Programme, the CBRC, UCL, Haematolinne, the Leukemia and Lymphoma Society, USA (Career Development Program fellowship to PSW), the Deutsche Forschungsgemeinschaft (fellowship to CL; LU1474/1-1 and programme grant as part of the Sonderforschungsbereich; SFB873, Teilprojekt A13).

Authorship Contributions

CL and VT performed experiments; RC analysed the SNP array; JH analysed the RNAseq data with input from VT; AS interpreted and supervised SNP-array data analyses; TS developed the mathematical model; CL and PSW performed the cell cycle analysis, VB carried out FISH analyses; AC provided patient data and samples; SAC, HF and PSW performed cell sorting. CL, VT, SEWJ and TE designed the study and interpreted data. CL, VT and TE wrote the paper with critical input from other authors.

Conflict of Interest Disclosures

The authors declare no conflict of interest.

References

1. Rosenthal, R., Mcgranahan, N., Herrero, J. & Swanton, C. Deciphering Genetic Intratumor Heterogeneity and Its Impact on Cancer Evolution. *Annu. Rev. Cancer Biol. 2017* **1**, 223–240 (2016).

2. Marusyk, A., Almendro, V. & Polyak, K. Intra-tumour heterogeneity: A looking glass for cancer? *Nature Reviews Cancer* vol. 12 323–334 (2012).

3. Landau, D. A., Carter, S. L., Getz, G. & Wu, C. J. Clonal evolution in hematological malignancies and therapeutic implications. *Leukemia* vol. 28 34–43 (2014).

4. Anderson, K. *et al.* Genetic variegation of clonal architecture and propagating cells in leukaemia. *Nature* **469**, 356–361 (2011).

5. Notta, F. *et al.* Evolution of human BCR-ABL1 lymphoblastic leukaemia-initiating cells. *Nature* **469**, 362–7 (2011).

6. Turati, V. A. *et al.* Chemotherapy induces canalization of cell state in childhood B-cell precursor acute lymphoblastic leukemia. *Nat. Cancer* doi:10.1038/s43018-021-00219-3.

7. Ebinger, S. *et al.* Characterization of Rare, Dormant, and Therapy-Resistant Cells in

Acute Lymphoblastic Leukemia. *Cancer Cell* **30**, 849–862 (2016).

8.   Lutz, C. *et al.* Quiescent leukaemic cells account for minimal residual disease in

childhood lymphoblastic leukaemia. *Leukemia* (2013) doi:10.1038/leu.2012.306.

9.   Waanders, E. *et al.* Mutational Landscape and Patterns of Clonal Evolution in

Relapsed Pediatric Acute Lymphoblastic Leukemia. *Blood Cancer Discov.* **1**, 96–111

(2020).

10.   Dobson, S. M. *et al.* Relapse-Fated Latent Diagnosis Subclones in Acute B Lineage

Leukemia Are Drug Tolerant and Possess Distinct Metabolic Programs. *Cancer*

Discov. (2020) doi:10.1158/2159-8290.cd-19-1059.

11.   Kirschner-Schwabe, R. *et al.* Chemotherapy and mismatch repair deficiency

cooperate to fuel TP53 mutagenesis and ALL relapse. *Nat. Cancer*

doi:10.1038/s43018-021-00230-8.

12.   Buitenkamp, T. D. *et al.* Acute lymphoblastic leukemia in children with Down

syndrome: A retrospective analysis from the Ponte di Legno study group. *Blood* **123**,

70–77 (2014).

13.   Xavier, A. C. & Taub, J. W. Acute leukemia in children with down syndrome.

Haematologica vol. 95 1043–1045 (2010).

14.   Schuh, A. *et al.* Monitoring chronic lymphocytic leukemia progression by whole

genome sequencing reveals heterogeneous clonal evolution patterns. *Blood* **120**,

4191–4196 (2012).

15.   Hong, D. *et al.* Initiating and cancer-propogating cells in TEL-AML1-associated

childhood leukemia. *Science (80-.).* **319**, 336–339 (2008).

16.   Zur Wiesch, P. S., Engelstädter, J. & Bonhoeffer, S. Compensation of fitness costs

and reversibility of antibiotic resistance mutations. *Antimicrob. Agents Chemother.* **54**,

2085–2095 (2010).

17.   Vazquez-Santillan, K., Melendez-Zajgla, & J., Jimenez-Hernandez, & L., Martínez-

Ruiz, G. & Maldonado, & V. NF-κB signaling in cancer stem cells: a promising

therapeutic target? doi:10.1007/s13402-015-0236-6.

18.   Sergushichev, A. A. An algorithm for fast preranked gene set enrichment analysis

using cumulative statistic calculation. *bioRxiv* 060012 (2016).

19.   Laurenti, E. *et al.* The transcriptional architecture of early human hematopoiesis

identifies multilevel control of lymphoid commitment. *Nat. Immunol.* **14**, 756–763

(2013).

20.   Geng, H. *et al.* Self-Enforcing Feedback Activation between BCL6 and Pre-B Cell

Receptor Signaling Defines a Distinct Subtype of Acute Lymphoblastic Leukemia.

Cancer Cell **27**, 409–425 (2015).

21. Dick, J. E. Stem cell concepts renew cancer research. *Blood* **112**, 4793–4807 (2008).

22. Kreso, A. & Dick, J. E. Evolution of the cancer stem cell model. *Cell Stem Cell* **14**,
275–291 (2014).

23. Ailles, L. E. & Weissman, I. L. Cancer stem cells in solid tumors. *Curr. Opin.*
Biotechnol. **18**, 460–466 (2007).

24. Rehe, K. *et al.* Acute B lymphoblastic leukaemia-propagating cells are present at high
frequency in diverse lymphoblast populations. *EMBO Mol. Med.* **5**, 38–51 (2013).

25. Shlush, L. I. *et al.* Tracing the origins of relapse in acute myeloid leukaemia to stem
cells. *Nature* **547**, 104–108 (2017).

26. Velten, L. *et al.* Identification of leukemic and pre-leukemic stem cells by clonal
tracking from single-cell transcriptomics. *Nat. Commun.* **12**, 1–13 (2021).

27. Lapidot, T. *et al.* A cell initiating human acute myeloid leukaemia after transplantation
into SCID mice. *MacKinnon, R. Yallen. G. Sci.* **362**, 31–39 (1993).

28. Dick, J. E. & Bonnet, D. Human Acute Myeloid Leukaemia is organised as a heirarchy
that originates from a primitive haematopoetic cell. *Nat. Med.* **3**, 730–737 (1997).

29. Taussig, D. C. *et al.* Leukemia-initiating cells from some acute myeloid leukemia
patients with mutated nucleophosmin reside in the CD34- fraction. *Blood* **115**, 1976–
1984 (2010).

30. Ford, A. M. *et al.* Fetal origins of the TEL-AML1 fusion gene in identical twins with
leukemia. *Med. Sci.* **95**, 4584–4588 (1998).

31. Greaves, M. F., Maia, A. T., Wiemels, J. L. & Ford, A. M. Leukemia in twins: Lessons
in natural history. *Blood* **102**, 2321–2333 (2003).

32. Gale, K. B. *et al.* Backtracking leukemia to birth: Identification of clonotypic gene
fusion sequences in neonatal blood spots. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 13950–
13954 (1997).

33. Ford, A. M. *et al.* Protracted dormancy of pre-leukemic stem cells. *Leukemia* **29**,
2202–2207 (2015).

34. Böiers, C. *et al.* A Human IPS Model Implicates Embryonic B-Myeloid Fate Restriction
as Developmental Susceptibility to B Acute Lymphoblastic Leukemia-Associated
ETV6-RUNX1. *Dev. Cell* **44**, 362-377.e7 (2018).

35. Wray, J. P. et al. Cell cycle corruption in a pre-leukemic ETV6-RUNX1 model exposes
RUNX1 addiction as a therapeutic target in acute lymphoblastic leukemia. *bioRxiv*
Prepr. (2021).

36. Tirosh, I. *et al.* Single-cell RNA-seq supports a developmental hierarchy in human
oligodendroglioma. *Nat. Publ. Gr.* **539**, (2016).

Figure legends

Figure 1: Study of clonal dynamics through serial transplantation assays. A.

Experimental setup. BM samples before and after serial transplantation in NSG-mice were

analyzed by SNP-arrays. **B.** SNP array results of patient 1. Deletions on Chromosomes 1,

12 and a single amplification on Chromosome 17 were identified and shown to be stable

during serial transplantation. Relationships between transplants recipients are indicated with

arrows. **C.** SNP array results of patient 2. Deletions on Chromosomes 9, 14 and a single

amplification on Chromosome X were identified. During serial transplantation copy number

abnormalities (CNA) disappeared and re-appeared suggesting the co-existence of different

sub-clones. Relationships between transplants recipients are indicated with arrows. **D.** Initial

reconstruction of the clonal hierarchy of patient 2 leukemia considering Chromosomes 9, 14

and X. * Indicate secondary transplants from a primary transplant for which SNP-array data

are unavailable due to insufficient numbers of cells for analysis.

Figure 2: Detailed analysis on chromosome 9 reveals higher clonal complexity

(Patient 2). A. In depth analysis of CNA on chromosome 9 during serial transplantation

suggests a more complex clonal hierarchy with multiple deletions of different size targeting

the locus and occurrence of cnLOH. Each box indicates an individual mouse. **B.**

Phylogenetic tree of leukemia cells based on SNP-array data obtained at diagnosis and

during serial transplantation. Mosaicim of CNA suggested the existence of sub-clones within

analyzed samples. The presented tree represents the minimum number of diagnostic clones

required to explain the mutational landscape identified in the engrafted leukaemias. Minimal

clone numbers were calculated by mathematical modelling (see supplementary materials).

Figure 3: Correlation of phenotype and genotype. A. Experimental setup. BM bulk,

StemB (CD34$^+$CD38$^{-/low}$CD19$^+$) and ProB (CD34$^+$CD38$^+$CD19$^+$) cells were analyzed by SNP-

arrays before and (if available) after serial transplantation in NSG mice. **B.** In patient 2 bulk,

stemB and ProB cells showed identical CNA. **C.** In Patient 3 animals transplanted with bulk,

stemB, and ProB cells showed identical CNA profiles: suggesting the existence of a single

dominant clone. * Indicate secondary transplants from a primary transplant for which SNP-

array data are unavailable due to insufficient numbers of cells for analysis. Secondary

transplants from Stem/B and bulk cells did not engraft.

Figure 4: Phenotypically, genetically and functionally distinct sub-clones co-exist in a

case of BCR-ABL1$^+$ ALL. A. FISH analysis of leukemic subpopulations at diagnosis and

after serial xenotransplantation in Patient 4 using a BCR/ABL-specific probe. To confirm the

leukemic status, sorted cells were analysed by FISH for the appropriate clonal marker.

Presence of the t(9;22) translocation is indicated by detection of one green, one red and two

red-green fusion signals. **B.** Competitive transplants of genetically and phenotypically distinct

BCR-ABL$^+$ ProB (CD34$^+$CD38$^+$CD19$^+$) and BCR-ABL$^-$ StemB (CD34$^+$CD38$^{-/low}$CD19$^+$) cells.

Input and output ratios of the two populations in individual mice are displayed. **C.** Cell cycle

analysis of leukemic ProB (upper panels) and StemB (lower panels) cells from competitively

transplanted NSG-mice with subpopulations from a BCR-ABL1$^+$ ALL patient. Percentages of

cells in G$_0$ (DAPI$^-$Ki67$^-$), G$_1$ (DAPI$^-$Ki67$^+$) and S/G$_{2/M}$ (DAPI$^+$Ki67$^+$) are shown.

Figure 5: Selective therapy resistance of CD34$^+$CD38$^{-/low}$CD19$^+$ Stem/B cells in high

risk BCR-ABL1$^+$ ALL. A. FACS analysis of BCR-ABL1 ALL subpopulations at diagnosis,

remission, and relapse (patient 4). Percentages of CD34$^+$CD38$^{-/low}$CD19$^-$ (HSC),

CD34$^+$CD38$^{-/low}$CD19$^+$ (StemB), CD34$^+$CD38$^+$CD19$^+$ (ProB) cells and CD34$^+$CD38$^+$CD19$^-$

(Progenitors) in total bone marrow (BM) mononuclear cells (MNC) are shown. The black

boxes show the gates used for frequency calculations. Not all collected events are shown. **B.**

FACS analysis of BM cells from serially transplanted NSG mice injected with sorted StemB

and ProB cells from patient 4 relapse. **C.** PCA analysis based on bulk RNA sequencing of

diagnostic and relapse cells harvested from primary and secondary xenografts. Six different

populations of cells were analysed. i) diagnostic stemB cells, ii) diagnostic proB cells iii)

relapse stemB cells harvested from secondary xenografts injected with stemB cells isolated

from primary xenografts iv) relapse proB cells harvested from secondary xenografts injected

with stemB cells isolated from primary xenografts, v) relapse proB cells harvested from

secondary xenografts injected with proB cells isolated from primary xenografts, and vi)

relapse stemB cells harvested from secondary xenografts injected with proB cells isolated

from primary xenografts. Arrows showing the contribution of different gene expression

signatures (calculated as per panel D) to the sample's clusters biology are shown. **D.** Gene

set enrichment analysis (GSEA) for a selection of MSigDB-50 Hallmark gene-sets and

published signatures associated with predefined differentiation stages of the B-cell

ontogeny[19]. ProB diagnostic cells are compared to the stemB diagnostic cells. Significance

thresholds: <0.05 after adjusting for multiple testing (using the Benjamini-Hochberg method).

Figure 6: Distinct sub-clones in the BCR-ABL1$^+$ ALL and the BCR-ABL$^-$ relapse clones

are genetically related (Patient 4). A. SNP array results of diagnostic stem/B, xenografted

stem/B, diagnostic proB, xenografted proB, diagnostic preB, relapse StemB, relapse ProB

and MRD-negative Remission sample are displayed. A chromosome 12 deletion was

present in all but the remission control sample illustrating the clonal relationship of all leukemic samples. CNA of diagnostic stem/B and ProB were identified and stable during serial transplantation with identification of some mosaicism suggesting sub-clones in each compartment. Relationships of transplants are indicated. Relapse stem/B and proB were genetically identical but stem/B showed 90% mosaicism of all clones suggesting the existence of a pre-leukemic clone. **B.** Detailed analyses SNP array revealed different sizes of some chromosome deletions at diagnosis, remission and relapse and suggest the existence of distinct pre-leukemic sub-clones for which a clonal hierarchy cand be re-constructed.

Figure 7: Pre-leukemic and leukemic hierarchy. Based on genetic mosaicism and different sizes of chromosome 7 and 14 deletions (Patient 4) a pre-leukemic clonal hierarchy could be constructed which showed clonal evolution at the pre-leukemic stage that gave rise to the diagnostic leukemic sub-clones as well as evolving into another pre-leukemic sub-clone responsible for relapse.

Supplementary Figures Legends

Figure S1: Flow cytometry sorting strategy. Representative gating strategy used for the analysis of immunophenotypically-distinct Down's syndrome ALL subpopulations before and after xenotransplantation. StemB (CD34$^+$CD38$^{-/low}$CD19$^+$) and proB (CD34$^+$CD38$^+$CD19$^+$) cells were sorted. **A-B.** Patient 2 and 3 leukaemia respectively.

Figure S2: Cell cycle analysis. A. Cell cycle analysis of representative xenografted stemB and ProB as compared to the **B.** patient relapse.

Figure S3: RNA sequencing analysis. A-C Gene Set Variation Analysis (GSVA) analysis of RNA sequencing data looking at the expression of cell cycle related gene signatures. Six different populations of cells were analysed: i) diagnostic stemB cells, ii) diagnostic proB cells iii) relapse stemB cells harvested from secondary xenografts injected with stemB cells isolated from primary xenografts iv) relapse proB cells harvested from secondary xenografts injected with stemB cells isolated from primary xenografts, v) relapse proB cells harvested from secondary xenografts injected with proB cells isolated from primary xenografts, and vi) relapse stemB cells harvested from secondary xenografts injected with proB cells isolated from primary xenografts. Within each heatmap samples are sorted by category and genes split into top 500 most variable vs all others. **A.** G1/S gene signature[36], **B.** G2/M gene signature[36], **C.** E2F targets gene signature. **C.** Gene set enrichment analysis (GSEA) for a selection of MSigDB-50 Hallmark gene-sets and published signatures associated with predefined differentiation stages of the B-cell ontogeny[19]. This analysis looks specifically at

676   the top 500 genes driving clustering along the PC2 component of the PCA analysis. **D.** PCA
677   analysis based on bulk RNA sequencing of diagnostic and relapse cells harvested from
678   primary and secondary xenografts. Arrows showing the contribution of gene expression
679   signatures involved in BCR signalling and B-cell differentiation are shown. Significance
680   thresholds: <0.05 after adjusting for multiple testing (using the Benjamini-Hochberg method).

Figure 1

Figure 2

A

Homozygous copy loss
copy loss
CnLOH
Chr 9

B

Connection present in
100%
80%
60%
40%
20%
of all graphs

Node present in
100%
80%
20%
of all graphs

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supplementary Figure 1

Supplementary Figure

A

BCR-ABL FISH analysis of sorted cells

* Ctrl: BCR-ABL status of non-Down's
associated ALL

	Ctrl *	Pt. 4
HSC	0	0
Stem/B	100	0
proB	100	100

100 cells/sample scored

Supplementary Figure 2

Cell cycle Analysis:

Supplementary Figure 3