1 FRONT MATTER

2 **Title**

3 **What do Functional Synchronization Networks Indicate About Underlying Structure and**
4 **System Dynamics? A network theory study of the islet.**
5

Short Title
Understanding Functional and Structural Networks

Authors
Jennifer K. Briggs[1], Vira Kravets[1,2], JaeAnn M. Dwulet[1], David J. Albers[1], Richard K. P. Benninger*[1,2]
1. Department of Bioengineering, University of Colorado Anschutz Medical Campus, Aurora,
Colorado
2. Barbara Davis Center for Childhood Diabetes, University of Colorado Anschutz Medical
Campus, Aurora, Colorado
* Corresponding author: richard.benninger@cuanschutz.edu
15

Abstract

Network theory provides tools for quantifying communication within heterogeneous biological systems. Functional and structural networks are common way to represent a biological system. The relationship between these two networks is not well quantified, leaving a knowledge gap for how to interpret functional networks in the context of structural changes. In diabetes, both structural and functional networks of the islet of Langerhans are altered, motivating the investigation as to how the two networks interact. Here, we analyze both experimental and computational models of the islet to quantify the relationship between the gap junction structural network and functional synchronization networks. We show that functional networks poorly reflect gap junction structure. Instead, they indicate intrinsic factors driving excitability, such as cellular metabolism. This surprising finding provides a new interpretation of functional networks within the islet of Langerhans and other biological systems.

Teaser

We use experimental and computational models to provide a new interpretation of intracellular synchronization in biological networks.

MAIN TEXT

Introduction

Our understanding of biological systems has benefitted from the application of network theory *(1–4)*, a
set of analysis techniques used to quantify the relationships between interacting parts of a system. In
network analysis, a system is represented by its entities (called nodes) and the interactions between those
entities (called edges). The topology and degree structure of the network **(Table 1)** can then provide
information about the biological system.

39   To apply network theory tools, one must decide how to represent the system. Common network
representations in biological tissues are based on either structure or function *(5–7)*. In both
representations, nodes generally correspond to physical neurons, cells, or regions of interest. The
difference between structure and function lies in how the edges are constructed. *Structural networks*
represent physical connections between nodes; therefore, edges correspond to physical connections, such
as synapses or gap junctions. However, in complex systems, such as most biological systems, the
structure network alone cannot explain how the system works *(5)*. Instead, functional behavior emerges
from complex interactions between individual cell dynamics and the underlying structure *(2, 8)*.
Therefore, functional networks are meant to represent the emergent system behavior, where edges
correspond to functionally similar cell pairs.

49   When deciding whether to use a functional or structural network, one must consider both the
research question they are investigating and the tools accessible to them. A *structural network* preserves
spatial information and should be used when asking questions concerning node influence through
physical communication or a physical change in the system. Obtaining a complete structural network of
a biological system requires imaging the entire 3D system with techniques that differentiate cells and
cell connections. Such network reconstruction may be an impossible task for some systems *(7)* and is
highly labor-intensive for others. *Functional networks* should be used when quantifying emergent system
behavior. Functional networks are often generated by drawing edges between pairs of neurons or other
cells that show highly correlated dynamics, for example, as observed through calcium-based imaging *(5,
8–10)*. Many network-informed biological studies utilize synchronization-based functional networks
because calcium-based imaging approaches are relatively common *(9, 11, 12)* and more accessible than
directly identifying structural connections. Since functional networks are more common and accessible
than structural networks in biology, it is relatively common for scientists to draw conclusions about the
systems' structure by using the structural network. This conclusion relies on the assumption that structure
and function have a 1-to-1 correspondence, which may not always be the case. In this paper, we
investigate the relationship between structural and functional networks to assess the validity of this
assumption.

66   Considered independently, structural and functional networks can solve interesting questions
regarding physical communication or emergent dynamics *(5)*. However, we are often concerned with the
intersection between these concepts: such as how physical changes in the structural network relate to
emergent dynamics in the functional network. Little is known about this interaction between functional
and structural networks. One system in which this is critically important is the pancreatic islet, which is
the focus of this paper. Pancreatic islets are comprised of insulin-producing, highly heterogeneous β-
cells that electrically communicate through connexin36 (Cx36) gap junctions *(13–15)*. This
communication allows for insulin release in coordinated pulses *(13, 16–18)*. Coordinated pulsatile

behavior is central to regulating the homeostasis of glucose and other nutrients *(19–21)* and is lost during
the development of type 1 and type 2 diabetes *(20, 22)*. Loss of intra-islet coordination in diabetes is
associated with changes in the *functional β-cell network*. However, it is unclear how disruption in the
functional network is related to changes in individual β-cell function and disruptions in the structural
network, as type 1 and some presentations of type 2 diabetes are also associated with dysfunction or
death of insulin-producing β-cells and loss of gap junction coupling *(22–26)*. Therefore, understanding
the interaction between structure, function, and system dynamics is important.

Studies have suggested that the functional β-cell network follows a scale-free-like distribution
(9) with a tendency toward high clustering *(27, 28)*, implying there exists a subset of highly connected
nodes called "β-cell hubs" *(6, 10)*. Note that mathematically scale free networks are rare in nature *(29)*,
so we use the term scale-free-like to describe a distribution with a roughly logarithmic distribution. Since
these hubs emerge from the functional network, it is unclear whether their high synchronization is due
to physical influence via higher than average structural coupling or intrinsic properties of the β-cells
themselves. Functional network studies suggest that these emergent β-cell hubs have a large influence
over islet dynamics *(10, 30, 31)*. Alternatively, proxies for structural networks, such as
electrophysiological experiments and computational studies measuring gap junction electrical coupling,
suggest it is not feasible for one cell or a small population of cells to have disproportionate influence
over the islet *(30, 32–34)*. These seemingly contradicting results have caused debate over the most
accurate topological description of the islet network *(35, 36)* (see *(37)* for a concise review of the β-cell
hub debate) and motivated our investigation of the relationship between structure and function of the
islet.

In this paper, we use experimental and computational techniques to investigate the relationship
between the structural and functional networks using three guiding questions (results are graphically
outlined in **figure 8)**. 1) Are subpopulations that emerge from the β-cell functional network
differentiated by a unique location in the structural network or by intrinsic properties of the cells? 2)
What does the islet functional network indicate about its underlying structure or intrinsic cell properties?
3) Can we fully explain the disruption in the functional network by the removal of nodes and edges in
the structural network, or are there other factors involved? Our data lead to the conclusion that the islet
functional network is more reflective of variability in cellular metabolic activity than gap junction
structural connections. These results provide fresh insight into interpreting the relationship between
structure, function, and intrinsic cell dynamics in the pancreatic islet.

Results

Cellular metabolism is more influential than gap junction coupling in determining highly synchronized cells in a simulated β-cell network.

Accurately analyzing the structural network or the intrinsic factors underlying cellular excitability is non-trivial to perform experimentally. Thus, to investigate the relationships between functional and structural networks, we first analyzed a simulated β-cell network. We used a multi-cellular coupled ODE model, which describes the electrophysiological properties of the β-cells. This model has been shown to accurately describe experimental findings concerning spatial-temporal dynamics *(38)*, the relationship between electrical coupling and metabolic heterogeneity *(39)*, the relationship between electrical heterogeneity and electrical activity *(40)*, and the influence of excitability parameters such as rate of glucose metabolism (k_{glyc}) and K_{ATP} channel opening kinetics on diabetes mutations *(39, 41)*. Our model included physiologically informed heterogeneity in multiple factors underlying cellular excitability and in gap junction electrical coupling (**Figure 1a**).

The first question we investigated was: Are subpopulations that emerge from the β-cell functional network differentiated by a unique location in the structural network or by intrinsic properties of the cells? Specifically, we first investigated whether β-cell hubs are differentiated by a unique location in the structural network or by the inherent properties of the cell? We first created the simulated islet's functional network of the intracellular free Ca^{2+} ($[Ca^{2+}]$) dynamics. We represented each cell with a node and drew an edge between any cell pair whose temporal calcium traces had a Pearson correlation coefficient (R_{ij}) greater than the threshold R_{th}. In a network, the degree is defined as the number of edges a given cell has, while the degree distribution describes the distribution of degrees throughout the network **(Table 1)**. We set the threshold to $R_{th} = 0.9995$, so the normalized degree distribution showed the presence of many cells with a relatively low degree and a few cells with a very high degree in the five simulated islets (**Figure 1b**), as indicated in previous studies *(9, 10)*. This power law-like distribution shows that the functional network acts as a scale-free-like distribution (Table 1), which implies that few cells in the islet are highly synchronized while most cells are not. One way to interpret a scale-free-like distribution is that highly synchronized cells are driving or coordinating the islet function.

We then identified a subpopulation of highly synchronized cells, referred to as β-cell hubs, as any cell whose degree (see **Table 1)** was greater than 60% of all other cells in the functional, as outlined in *(10)*. Within a simulated islet, multiple such β-cell hubs were present (**Figure 1c).** These β-cell hubs tended to depolarize before and repolarize after the islet average (**Figure 1d**), in agreement with previous studies *(10)*. We next analyzed the properties of these highly connected β-cell hubs. Johnston et al. showed that β-cell hubs have higher than average metabolic activity *(10, 41)*; therefore, we chose to investigate this parameter. Critical factors which influence β-cell excitability include the rate of glucose metabolism to generate ATP (k_{glyc}) and the activity of ATP-sensitive K^+ (K_{ATP}) channels. β-cell hubs had a significantly higher rate of glucose metabolism (k_{glyc}) than non-hubs ($p \leq 0.0001$) (**Figure 1e**) but were not significantly different in their maximum K_{ATP} conductance (g_{KATP}) ($p = 0.18$) (**Figure 1f**). These results indicate that β-cell hubs are differentiated by glucose metabolism rather than K_{ATP}-dictated cellular excitability. Interestingly, β-cell hubs had only a slightly higher gap junction conductance (g_{coup}) than non-hubs ($p = 0.032$) **(Figure 1g).** To ensure consistency in our findings, we repeated the following analysis with various thresholds **(Supplemental 1a)**. For all thresholds analyzed,

the significance between hubs and non-hubs was greater for k_{glyc} than g_{coup} **(Supplemental 1b)**. These
results indicate that both metabolic activity and gap junction coupling correlate with high cellular
synchronization, where the rate of metabolic activity has a stronger correlation.

<u>Elevated cellular metabolic activity, but not elevated gap junction permeability is observed</u>
<u>experimentally in highly connected cells</u>
To further investigate question 1 concerning subpopulations that emerge from the β-cell functional
network, we tested whether the properties of highly connected cells that we identified within the
simulated network occur experimentally within islets of Langerhans. We performed time-lapse imaging
of [Ca^{2+}] within islets isolated from Mip-CreER; Rosa-LSL-GCamp6s mice that express GCamP6s
specifically in β-cells (β-GCamP6s mice, see methods). At elevated glucose, islet β-cells showed
synchronous oscillations in [Ca^{2+}] **(Figure 2a,b)**. We performed Pearson-based functional network
analysis on these [Ca^{2+}] time-courses (see methods) **(Figure 2c)** and generated a normalized degree
distribution that describes this experimentally-derived functional network **(Figure 2d)**, which again
reflects a large number of cells with low degree and few cells with high degree, which can be interpreted
as a scale-free-like distribution (Table 1).
To test whether cellular metabolic activity differentiated high-degree cells from normal or low-
degree cells in the functional network, we performed two-photon excitation of NAD(P)H
autofluorescence of β-cells in conjunction with time-lapse imaging of Ca^{2+} dynamics at 2- and 11-mM
glucose levels **(Figure 2e)**. The metabolic activity of individual cells in the islet was calculated as a
difference between the NAD(P)H autofluorescence level at high glucose compared to low glucose
($NAD(P)H_{11-2}$). This was adjusted by the mean change in autofluorescence across all cells in the islet
($NAD(P)H_{11av-2av}$). Therefore, the metric was 0 if the NAD(P)H response of a cell was the similar to the
islet average, >0 if a cell showed an elevated metabolic response indicating it is more metabolically
active, and <0 if a cell showed a reduced metabolic response indicating it is less metabolically active.
The NAD(P)H response to glucose elevation decreased as the cell's normalized degree decreased
(Figure 2f). This indicates that highly synchronized cells are more metabolically active, as measured by
NAD(P)H autofluorescence, in agreement with findings from simulated islets. Furthermore, the
NAD(P)H response trended lower in cells that were functionally disconnected, as indicated by Ca^{2+}
oscillations lacking synchronization with any other cells, compared to the NAD(P)H response in
connected cells **(Figure 2g)**. Therefore, the functional network is strongly related to metabolism which
drives islet dynamics.
While it is not possible to experimentally measure the strength of every gap junction connection
across the islet, fluorescence recovery after photobleaching (FRAP) measurements of dye transfer
kinetics can measure the gap junction permeability within the islet with cellular resolution *(43)*. We
performed FRAP measurements of gap junction permeability using a Cx36 gap-junction-permeable
tracer Rhodamine-123 in conjunction with Ca^{2+} time-lapse imaging of GCamP6s, to map cellular gap
junction connections in the same cell layer as functional connections are calculated **(Figure 2h,i)**. We
found that the rate of recovery, which assesses gap junction permeability, lacked any trend with Ca^{2+}
synchronization such that cells with high degrees showed similar permeability as cells with low degrees
(Figure 2j). This indicates that cell synchronization was not strongly correlated with gap junction

permeability, as measured by FRAP, which is consistent with findings from the simulated islet.
Therefore, the functional network is not strongly related to the structural network.

<u>A synchronized cell pair is more likely to indicate shared metabolic properties between the cells than a</u>
<u>shared gap junction.</u>
Our findings so far indicate that highly synchronized cells within the functional network correlate more
strongly with intrinsic metabolic properties such as metabolic rate and activity than with gap junction
electrical connections. We then investigated the second question: What does the islet functional network
indicate about its underlying structure or intrinsic cell properties? We observed few instances in which
a cell pair was connected in both the functional and structural networks (**Figure 3a**), irrespective of a
cell's degree of synchronization (considering highly connected β-cell hubs, cells with average
synchronization, or cells with low synchronization**).** This indicates that cellular synchronization does not
imply the presence of a gap junction. To quantify this relationship, we calculated the conditional
probabilities of finding edges in the functional and/or structural networks. The probability that two cells
were synchronized in the functional network given that they shared a gap junction in the structural
network, $Pr(Sync|GJ) = 0.39$ (**Figure 3b**). Therefore, less than half of cell pairs that are electrically
coupled by gap junction connections are considered to be functionally coupled. The alternative
probability of a structural edge given a functional edge, $Pr(GJ|Sync) = 0.04$ (**Figure 3b**), indicating
that very few cells that are functionally coupled are directly connected by gap junction connections.
These findings indicate cells that are connected in the structural network (gap junction) and cells that are
connected in the functional network (synchronized $[Ca^{2+}]$ dynamics) are two distinct groups with little
overlap ($Pr(both) = 0.2$) (**Figure 3c**).
We again analyzed the sensitivity of these measures to synchronization threshold R_{th}. By nature
of the method used to create the functional network, as the threshold was increased, the number of cells
considered synchronized substantially decreased (**Supplemental 1d**). The probability that two cells were
gap junction coupled given that they were connected in the functional network greatly increased with
increasing correlation threshold (**Supplemental 1e**), such that when $R_{th} = 0.99995$, $Pr(GJ|Sync) \approx$
0.5. This was not due to an increasing "overlap" between the functional network and structural network,
but rather the decrease in the size of the functional network sample space (**Supplemental 1f**). Therefore,
our findings that the functional and structural networks are two distinct groups with little overlap is
consistent across thresholds.
These data corroborate with initial findings showing that function is not strongly correlated with
structure. To investigate whether metabolic properties are correlated with function, as shown in our
previous results, we created two new networks involving rate of glucose metabolism (k_{glyc}). In the first
network (i), we drew an edge between cells if they had similar rates of k_{glyc} and were separated by less
than 15 cells (see methods) **(Figure 3d light green).** In the second network (ii), we drew an edge between
cells if they had higher than average rates of k_{glyc} and were separated by less than 15 cells **(Figure 3d**
dark green). We chose these parameters such that there was no difference in the probability of
synchronization (edge in the functional network) given a gap junction or metabolic quality (**Figure 3e**).
The alternative probability of unique properties in glucose metabolism (similar or high k_{glyc}) given a
functional edge was significantly higher than that of structural connections ($p \approx 0.005$ for both networks
(Figure 3f). While total number of links was larger in the metabolic networks, it is unlikely that these

results were caused by a difference in sample size given the initial probabilities $\Pr(Sync|GJ)$ and
$\Pr(Sync|k_{glyc})$ were similar. Therefore, if two cells have synchronized Ca^{2+} oscillations, they are more
likely to show either similar metabolic activity or have higher than average metabolic activity, compared
to the two cells that are gap junction coupled. This is further indicated by the increased overlap between
the Ca^{2+} synchronization-derived functional network and the metabolic activity-derived network (**Figure**
3g). These results further indicate that metabolic activity, not gap junction connections, is a greater
driving factor for cells to show high Ca^{2+} synchronization and thus influence the functional network, in
agreement with our previous findings.

<u>Elevated cellular rate of glucose metabolism is a greater driver of long-range functional connections.</u>
We noted that in the simulated islet model, gap junction connections only exist between highly proximal
cells, in accordance with the islet cytoarchitecture. However, there was no distance constraint on $[Ca^{2+}]$
oscillation synchronization (**Figure 4a**). To determine whether this spatial constraint was responsible
for the low correspondence between the functional and structural networks, we asked whether a series
of highly conductive gap junction connections could predict long-range functional connections. Previous
work has shown that that long-range functional connections can arise without assuming long-range
structural connections *(44)*. We can assess the total gap junction conductance along a path connecting
two cells by weighting the structural network by gap junction conductance. We then calculated the path
which maximized conductance between every cell pair in the simulated islets (**Figure 4b**) (see methods:
shortest weighted path length), allowing us to quantify the chains of highly conductive gap junctions.
We then partitioned the cell pairs based on whether or not they were connected in the functional network.
We express our results as the total g_{coup} for different distances of cell separation. In some cases, the total
g_{coup} between a given cell and a cell it was synchronized with was greater than that between the given
cell and a non-synchronized cell of the same cell distance, **(Figure 4b top)** while in other cases the non-
synchronized cell showed a higher total g_{coup} **(Figure 4b bottom).** This indicates that not all
synchronized cell pairs have higher chains of gap junction conductance than non-synchronized cell pairs.
Further, the probability distributions of total conductance for synchronized cells highly overlapped with
those for non-synchronized cells. This was observed for cells that were directly gap junction coupled
(**Figure 4c**) and cells that showed longer-range functional connections (**Figure 4d**). We compared the
total conductance between all cells irrespective of separation distance by normalizing the total
conductance by separation distance. The normalized total conductance was slightly less for non-
synchronized cells than for synchronized cells (**Figure 4e**), although still with substantial overlap
(**Figure 4 c,d**). Thus, long-range functional connections traverse cells that are connected on average by
slightly higher gap junction conductance. However, the substantial overlap in total conductance between
synchronized and non-synchronized connections is consistent with the minor influence of gap junction
conductance in driving high synchronization between cells. Therefore, while gap junction conductance
influences long-range synchronization, it is not the driving factor and cannot be used to differentiate cell
pairs connected in the functional network.
To examine how cellular rate of glucose metabolism influenced long-range functional
connections, we repeated this procedure but weighted graph edges by k_{glyc} rather than coupling
conductance (g_{coup}). Weighting the network edges by k_{glyc} allows us to quantify whether highly
synchronized cells result from chains of cells with high rates of glucose metabolism. We present the total

k_{glyc} along the path which maximized k_{glyc} between two cell pairs for different distances of cell separation.
Again, the probability distributions of the total k_{glyc} for synchronized cells were highly overlapping with
that for non-synchronized cells; both for cells that were directly gap junction coupled (**Figure 4f**) and
cells that showed longer-range functional connections (**Figure 4g**). The total rate of glucose metabolism,
normalized by separation distance, was significantly less for non-synchronized cells than for
synchronized cells (**Figure 4h**), with little overlap. Thus, long-range functional connections on average
traverse cells with higher rates of glucose metabolism.

To test whether there was a spatial relationship for which rate of glucose metabolism or gap
junction conductance-controlled synchronization, we then analyzed the total conductance or total
cellular metabolic rate for different separation distances compared between synchronized cells or non-
synchronized cells. Cumulative distribution functions (CDFs) reveal the small difference in total
conductance (**Supplemental 1**) or total rate of glucose metabolism (**Supplemental 2**) for each cell
separation distance. Synchronized cells had significantly higher total cellular metabolic rate than non-
synchronized cells for distances 1 to 5 cells apart, with very high level of significance (see methods,
significance testing). Conversely gap junction conductance was significantly higher for only
synchronized cells 2, 4, 5, 6, and 7 distances apart, with a lower level of significance
(average significant $P_{k_{glyc}} = 0.0064$ and average significant $P_{g_{coup}} = 0.0194$). Thus, cells pairs with
similar metabolic rate can strongly influence longer-range function connections up to distances of 5 cells
apart. In contrast, cells with higher gap junction conductance have less influence over longer-range
functional connections. Nevertheless, gap junctions retain some influence over long distances.

<u>Individual gap junction connections cannot be predicted by the functional network with correlated gap</u>
<u>junction conductance and metabolic activity.</u>
Our computational model assumes that metabolic parameters and gap junction coupling are independent.
However, it has previously been suggested that cells with highly synchronized Ca^{2+} oscillations possess
both elevated metabolic activity (in agreement with our findings, **Figure 2**) *and* high levels of gap
junction conductance (unlike what we see in our experiment, **Figure 2**) *(10, 31)*. To address the
possibility that cells with high metabolic activity may also have higher levels of gap junction
conductance, we simulated an islet in which the rate of glucose metabolism positively correlated with
gap junction conductance (**Figure 5a**), similar to the distribution presented in *(30)*, and generated the
functional network (**Figure 5b**). There was no difference in the average degree of the functional network
(i.e., average islet synchronization) between the random distribution (**Figure 1**) and this proportional
distribution (**Figure 5c).** Highly synchronized β-cell hubs had a significantly higher metabolic rate
compared to non-hubs (**Figure 5d**). In this simulated islet the gap junction conductance was also
significantly higher in hubs (**Figure 5e**), as expected given that cells with higher k_{glyc} are programmed
to have higher g_{coup}. Again, there was no significant difference in g_{KATP} between β-cell hubs and non-
hubs (**Figure 5f**). As with a normal distribution, the probability that two cells were synchronized in the
functional network given that they shared a gap junction connection in the structural network
($Pr(Sync|GJ)$=.303) was much greater than the alternative probability of a structural edge given
functional edge ($Pr(GJ|Sync)$=0.027) (**Figure 5g**) Importantly, both $Pr(Sync|GJ)$ and $Pr(GJ|Sync)$
were smaller than the equivalent probabilities determined when k_{glyc} and g_{coup} are not correlated (**Figure**
5g, **Figure 3b**). Therefore, the structural network still could not be effectively derived from the

functional network **(Figure 5 g,h).** We calculated the total g_{coup} between synchronized and non-
synchronized cells along weighted networks using the previous methodology. The total g_{coup} was
significantly higher for cell distances from 1-9 and 14 cells apart **(Figure 5i)**. Therefore, when g_{coup} is
proportional to k_{glyc}, synchronization is more predictive of long-range connections mediated by high gap
junction conductance than of individual gap junction connections.

<u>Modulating the islet-wide structural network in Cx36 knockout islets</u>
Our final question was: Can we fully explain the disruption in the functional network by the removal of
nodes and edges in the structural network, or are there other factors involved? This question is
particularly relevant in pathophysiological conditions, where physical connections or cells are removed
from the system. Given that Cx36 forms gap junction channels that enable electrical communication
between β-cells, reducing or eliminating Cx36 will remove edges from the structural network. It is well
established that changes to gap junction conductance affect the emergent Ca^{2+} dynamics across the islet
(13). We performed functional network analysis on time-courses of oscillatory $[Ca^{2+}]$ dynamics in islets
from wild-type mice with standard gap junction coupling (Cx36$^{+/+}$), heterozygous Cx36 knockout mice
with approximately 50% reduced gap junction coupling (Cx36$^{+/-}$) and homozygous Cx36 knockout mice
with no gap junction coupling (Cx36$^{-/-}$) *(13)* **(Figure 6a-c)**. To ensure consistency across genotypes, we
chose a synchronization threshold R_{th} such that the degree distribution of the WT islets was similar to a
scale-free-like distribution *(9)* while the average network degree ranged from 5-15 (see methods).

Figure 6c shows representative degree distributions of the functional networks corresponding to
the islets shown in **Figure 6a**. Average histograms of the degree distribution for all islets are presented
in **Supplemental 4a**. Results separated by islets are shown in **Supplemental 5.** As expected, the
functional degree distributions for both Cx36$^{+/-}$ islets and Cx36$^{-/-}$ islets were increasingly right-skewed
(**Figure 6c, Supplemental 4a**), indicating a decrease in the average islet synchronization. Heterozygous
Cx36$^{+/-}$ islets demonstrated decreased average network degree (see **Table 1**) compared to wild-type
Cx36$^{+/+}$ islets (**Figure 6d**, **Figure 6a-c** middle vs. left), and homozygous Cx36$^{-/-}$ demonstrated further
decreased degree compared to both wild-type Cx36$^{+/+}$ islets (**Figure 6d**, **Figure 6a-c** right vs. left) and
Cx36$^{+/-}$ islets (**Figure 6d**, **Figure 6a-c** right vs. middle), further indicating that decreases in gap junction
conductance leads to decreases in $[Ca^{2+}]$ oscillation synchronization (**Supplemental 4b**). Consistent
with this, substantially more cells in Cx36$^{+/-}$ islets (74% of cells) and Cx36$^{-/-}$ islets (93% of cells) were
not functionally connected to any other cell in the islet, compared to in Cx36$^{+/+}$ islets (40% of cells)
(**Figure 6e**). These data show that edges in the functional network become highly sparse as the structural
network decreases. It unlikely that these findings result from differences in network size across
genotypes as islets were of similar size ranges in each case (**Supplemental 3c**).
We next sought to quantify how the removal of edges in the structural network influences the
functional network topology (**Table 1**). Network topology statistics give us insight into how the network
functions. For example, *clustering coefficient* (C_{avg}) is a local parameter that quantifies the percent of
nodes connected to a given node that are also connected to each other (**Table 1**). A local parameter is a
measure of the neighborhood of connections around an individual node. A high clustering coefficient in
the functional network would indicate that cells tend to synchronize with other cells that share some
property. The clustering coefficient of the functional networks decreased significantly as gap junction
coupling decreased, being significantly lower in Cx36$^{+/-}$ islets and Cx36$^{-/-}$ islets compared to Cx36$^{+/+}$

islets (**Figure 6f**). To statistically quantify whether the functional network topology is intentionally
organized or a result of random edge generation, we compared each metric with that determined from
one thousand random networks created with the same number of nodes, edges, and degree of
connectivity as the functional network of interest *(4, 7, 9)*. An example random graph is shown in **Table
1**. The relative overlap between the network characteristics from the islets of interest with those of a
random network describes the likelihood that the functional network of the islets could be created by
random chance, where random graphs have low clustering *(7)*. If the physical gap junctions are the
property that gives the functional network a high clustering coefficient, then reducing gap junction
coupling should result in a clustering coefficient closer to zero and be completely explained by the
random graph. For all islets, the clustering coefficient of the islet network was greater than that of a
random network $C_{avg} > C_{rand}$ (**Figure 6f,** box and whisker plot). Thus, in agreement with previous
results, this finding suggests that something other than the gap junction structural network contributes to
local properties of the functional network topology.

Average shortest path length (L_{avg}) is a global parameter defined as the average smallest number
of nodes required to traverse between any cell pair (**Table 1**). Global parameters define the connectivity
across the entire network. A random network or a regular network (see Table 1) with a few edges
randomly moved (**Table 1**) will have a small shortest path length. The normalized shortest path length
increased significantly as gap junction coupling decreased, and was significantly higher in Cx36[+/-] islets
and Cx36[-/-] islets compared to Cx36[+/+] islets (**Figure 6g**). For Cx36[+/+] islets, the measured shortest path
length was much greater than that of a random network ($L_{avg} > L_{rand}$). However, as gap junction
coupling decreased, the measured shortest path length showed a larger intersection with that of a random
network: for Cx36[-/-] islets all measured data points could be explained by random chance ($L_{avg} \approx$
L_{rand}). This finding suggests that the gap junction structural network contributes to global, long-range
properties of the functional network topology, similar to our findings in figure 4. *Global efficiency*, which
has previously been used to describe islet networks *(9)*, is related to the inverse of the shortest path length
(**Table 1**). Global efficiency significantly decreased with decreased gap junction coupling (**Figure 6h**).
Similar to path length, as gap junction coupling decreased, the measured global efficiency showed a
larger intersection with that of a random network.

The functional network in WT mice islets has been shown to contain *small-world qualities (9)*,
meaning that it exhibits the high clustering of a regular network and the high efficiency (short path
length) of a random network (**Table 1**). A small-word functional network is highly efficient to traverse
(along functional edges), but cells still tend to synchronize will other cells of similar attributes. From
such a network hubs arise. We sought to investigate whether these functional network small-world
qualities were diminished when structural edges were removed. Previously *(9)*, small-worldness was
quantified using random networks. However, as the functional network becomes more sparse, the
uncertainty in random network based approach increases (**Supplemental 4d,** seen more clearly in
Supplemental 6d). To remove this uncertainty, we used an alternative metric *(45)* which was formulated
specifically to analyze small world-ness in sparse networks. *Local efficiency* reveals how efficient the
communication is between neighbors of a node when that node is removed. Thus if both global efficiency
E_{global} and local efficiency E_{local} are large, the network has small world qualities *(45)*. There was a
significant decrease in local efficiency with reduced gap junction coupling, being significantly lower in
Cx36[+/-] islets and Cx36[-/-] islets compared to Cx36[+/+] islets (**Figure 6j).** These data show that as gap

junction coupling decreases, both global and local efficiency decrease and therefore small world-ness of
the functional network decreases. Therefore reducing gap junction coupling decreases (but does not
completely abolish) the communication efficiency of the islet. However, the fact that the random
networks cannot explain the clustering of Cx36$^{-/-}$ indicates that other attributes are driving cellular
synchronization, in agreement with our previous results.

Finally, to quantify the relationship between physical location and [Ca^{2+}] oscillation
synchronization, we calculated the ratio between the average distance separating synchronized cells and
the average distance separating all cells in the islet $\left(D = \frac{D_{sync}}{D_{all}}\right)$. If this ratio is less than one, it would
indicate that cells tended to synchronize preferentially based on location. The average distance between
synchronized cells compared to islet average was less than one for all but one Cx36$^{-/-}$ mouse **(Figure**
6k) and there was no significant difference between the three islet genotypes, suggesting that cells tend
to synchronize to nearby cells but this is not fully dependent on gap junction coupling. These results
together suggest that variations in physical gap junction coupling is not the only factor driving the
synchronization of oscillatory [Ca^{2+}] dynamics. In all cases, these functional network characteristics did
not show any significant mouse to mouse variability (**Supplemental 5**).

<u>Altering the gap junction structural network in the computational model agrees with experiments.</u>
Finally, we combined all three of our questions by investigating how changes in the structural network
affect the functional network topology and relationships between function, structure, and metabolism.
We simulated decreased structural edges by altering the average coupling conductance (g$_{coup}$) in our
simulated islet. As expected, islets with reduced average g$_{coup}$ showed reduced [Ca^{2+}] oscillation
synchronization: simulated islets with average g$_{coup}$ = 0.12 nS had higher synchronization than those
with average g$_{coup}$= 0.06 nS or 0.03 nS (**Figure 7a**). Similarly, following functional network analysis,
the total number of functional network edges decreased as gap junction coupling was decreased
(visualized in **Figure 7 a,b**), indicating that our simulated islet follows closely with experiments. The
probability that two cells were connected in the functional network given that they were gap junction
coupled (connected in the structural network, $\Pr(Sync|GJ)$) decreased significantly as gap junction
coupling was decreased (**Figure 7c**). This indicates that as the gap junction conductance decreases and
the network becomes sparser, the probability that two gap junction connected cells will be functionally
connected decreases. Similarly, the probability that two cells were connected in the functional network
given the cells had similar rates of glucose metabolism decreased as gap junction coupling decreased
(**Figure 7d**). This indicates that gap junctions do play a role in synchronizing metabolically similar cells.
Conversely, the probability of a gap junction connection given that two cells were synchronized in the
functional network ($\Pr(GJ|Sync)$) increased as gap junction coupling decreased (**Figure 7e**). The
probability of two cells showing similar rates of glucose metabolism given that two cells were connected
in the functional network also increased as gap junction coupling decreased **(Figure 7f)** and showed
higher probabilities ($0.2 - 0.3$ vs $0.1 - 0.2$) than that for $\Pr(Sync|GJ)$ in each case. These results
indicates that as gap junction coupling decreases, a synchronized cell is more likely to be connected via
a gap junction or metabolic similarity. However, this is not because the overlap between the two
networks is larger but because the number of synchronized cells is smaller (**Figure 7b).** Together, these

results show that gap junctions are important for facilitating islet synchronization, but metabolism drives
synchronization between individual cells given sufficient coupling.
We then quantified the three-dimensional stimulated functional network topology upon reduced
gap junction conductance and compared these with network topology calculated from the two-
dimensional calcium imaging data. The average degree significantly decreased as coupling decreased
(**Figure 7g).** The clustering coefficient C_{avg} progressively decreased with decreasing gap junction
conductance (**Figure 7e**). However, the clustering coefficient from the simulated islets was always
substantially greater than that determined from a random network, with no overlap (**Figure 7e**). Global
efficiency also progressively decreased with decreased gap junction conductance (**Figure 7i**). The global
efficiency from the simulated islets was always less than that determined from a random network with
no overlap (**Figure 7i**). These trends are similar to that of the experiments but the lack of overlap between
random and measured global efficiency is more pronounced, likely due to having more cells from a
three-dimensional view of the islet network. Also similar to our experimental observations the local
efficiency decreased with gap junction conductance (**Figure 7j**). The decreases in both global efficiency
and local efficiency with decreasing gap junction conductance indicate that the network strays away
from small world properties as gap junction conductance decreases. This follows our experimental
observations.
450

Discussion:

The old saying goes "cells that fire together, wire together." However, this has not been directly proven for every biological system. Further, it may be the case that cells that fire together have similar intrinsic excitatory characteristics that allows them to share dynamics. The primary aim of this study was to investigate the relationship between functional networks, structural networks, and underlying system dynamics of the β-cells in the pancreatic islet. We framed our study around three guiding questions. 1) Are subpopulations that emerge from the β-cells functional network differentiated by a unique location in the structural network or by intrinsic properties of the cells? 2) What does the islet functional network indicate about its underlying structure or intrinsic cell properties? 3) Can we fully explain the disruption in the functional network by the removal of nodes and edges in the structural network or are there other factors involved? **Figure 8** shows a graphical representation of the answers to these questions based on our data.

Calcium-based imaging is a common technique used to analyze cellular dynamics, identify cell subpopulations, and represent the system by functional networks. For this reason, most network-informed biological studies utilize synchronization-based functional networks to make conclusions about a system. This approach has yielded tremendous insight into how different subpopulations of cells can contribute to system dynamics *(9, 10, 27, 30, 32, 46)*. However, when discussing the loss of synchronization due to cell death or disruption in signaling, such as the case for many diseases, it is changes in the structural network (e.g., gap junctions, synapses, accumulation of plaque) that cause dysfunction. Therefore, it is important to understand the relationship between structural and functional networks. The islet of Langerhans is a relatively closed biological system with simple architecture that affords the opportunity for high fidelity experimental and computation analysis, making it an ideal system to answer biological network theory-based questions.

To make study more robust, we used both experimental and computational techniques. Computational modeling is beneficial as it allows us to carefully adjust and extract the exact system parameters, investigate the system explicitly, and quantify the entire three-dimensional network without noise. Experiments can then be used to verify model findings and trends on real-world data. This kind of joint analysis is important for robust, repeatable conclusions.

1. Subpopulations of β-cells in the functional network are driven by metabolism more than gap junctions

Our computational and experimental results show the existence of a subpopulation of highly synchronized cells within the islet, known as β-cell hubs *(10, 47)* **(Figure 1)**. β-cell hubs are identified as any cell whose degree from the functional network is >60% of the network maximum and they are characterized by a higher than average metabolic activity and the electrical signature of depolarizing before and repolarizing after the rest of the islet. The simulated β-cell hubs emerge from the network with the assumption that rate of glucose metabolism and gap junction coupling is independent of one another. Therefore, the dependency between rate of glucose metabolism and gap junction coupling seem to not be required for the existence of β-cell hubs. Interestingly, rate of glucose metabolism is associated with β-cell hubs more strongly than structural gap junction coupling. Computational studies showed a low correlation between gap junction coupling and synchronization, while experiments showed no trend. In contrast, experimental studies showed a clear correlation between cell degree in the functional

network and metabolic activity (reflected by NAD(P)H florescence) (**Figure 2**). Semi-quantitative
immunofluorescence has indicated that β-cell hubs may show elevated metabolic activity due to elevated
glucokinase protein levels *(10)*. However, to our knowledge real-time quantification of metabolic
activity has not been correlated to cellular synchronization. Our results were validated experimentally.
Fluorescence recovery after photobleaching (FRAP) *(48)* provides a measurement for gap junction
permeability. By analyzing islets using both Ca^{2+}-based functional analysis and FRAP, our data showed
that highly synchronized cells do not have higher gap junction permeability (**Figure 2**). *These*
experimental findings support our computational findings that gap junctions are not the primary
distinguishing factor in determining variations in cellular synchronization.

<u>2. Metabolic similarities drive the functional network</u>
Conditional probabilities quantify the relationship between the structural and functional networks at a
cell-to-cell basis. The probability of synchronization given gap junction connection (Pr(Sync|GJ))
addresses the question of whether cells connected an edge in the structural network are always connected
in the functional network. Unexpectedly, only ~40% of gap junction connected cells were considered
highly synchronized by functional network analysis (**Figure 3b)**. While it is a well-known fact that gap
junctions are required for islet-wide synchronization *(13)*, this finding suggests that other factors besides
just physical connections contribute to the high synchronization of individual cell pairs. The probability
of a gap junction given a synchronized cell pair (Pr(GJ|Sync)) addresses the assumption that
synchronization can be used to infer gap junctions. If this probability is high, then functional network
analysis can be used to directly infer the underlying gap junction structure. We found that Pr(Sync|GJ)
for five simulated islets was $< 5\%$, indicating that synchronization does not directly imply a gap junction
connection (**figure 3b)**. Interestingly, the probability that two synchronized cells had high average
metabolism or similar average metabolism was significantly larger than the probability of them sharing
a gap junction β-cell hubs(Figure 3e, f). This leads to the conclusion that elevated glucose metabolism,
rather than elevated gap junction coupling is driving high synchronization between individual cells. This
conclusion corresponds to our first finding that subpopulations of β-cells in the functional network are
driven by glucose metabolism to a greater degree than gap junction connections.
We also investigated whether a long-range connected cell pair in the functional network indicates
a chain of strongly coupled cells in the structural network (**Figure 4)**. While the electrical communication
between β-cells is not uni-directional, this analysis allows us to detect even the most minute influence
of gap junction paths on Ca^{2+} synchronization. Synchronized cells had a lower gap junction resistance
(or higher conductance) than non-synchronized cells however, the strong overlap in distributions
indicates that functionally connected long-range pair of cells do not always have a higher gap junction
conductance than a cell pair that does not share a functional connection. Interestingly, we also found that
longer-range cellular synchronization was more strongly correlated with the metabolic rate for up to 5
cells separation. These findings indicate that short to medium range synchronization is more reflective
of metabolic activity.
The surprising part of our findings is that gap junctions do not guarantee high synchronization
between a cell pair, as previously assumed *(28)*. We know that gap junctions facilitate ion passage
between cells to synchronize electrical oscillations. Therefore, it makes sense that long-range
synchronization requires a chain of highly conductive gap junctions, and that islet wide synchronization

is decreased when gap junctions are removed. However, we also know that in the case of electrically
active cells such as neurons or cells in the islet of Langerhans, gap junctions are not the primary factor
driving electrical activity. In β-cells, cell metabolism is responsible for driving electrical activity and
Ca^{2+} elevations, while gap junctions simply facilitate ion passage between cells to synchronize electrical
oscillations. Additionally, we know from electrophysiology that it is the individual cell's membrane
conductance, controlled in part by glucose metabolism, that determines the actual coupling current
strength *(30, 49)*. Prior experiments have also shown that cells that are preferentially activated by
optogenetic ChR2 stimulation were more metabolically active by NAD(P)H autofluorescence, but not
preferentially gap junction coupled *(38)*. Furthermore, gap junctions have a larger role in repolarizing
cells back to resting membrane potential compared to depolarizing cells to an elevated membrane
potential *(30)*; whereas metabolic activity is more influential in driving cellular depolarization.
Additionally, in a completely uncoupled network, cells with identical frequencies will still be
synchronized. Since the metabolic activity is highly influential in a cell's oscillatory frequency, it makes
sense that metabolic activity would be strongly influential in determining functional edges.

<u>3. Removal of edges in the structural network decreases functional efficiency and local connectivity</u>
While the functional network is not a direct replica of the structural network, previous studies show that
there is a relationship between the two *(13, 50)*. Genetically knocking out the Cx36 gap junction protein
in mice islets allows us to "remove" edges from the structural network. Previous studies have shown that
Cx36 deficiency in mice leads to slower calcium waves *(13)*, loss of calcium oscillations and
synchronization and loss of pulsatile insulin secretion *(15)*. In modeling studies, coupling strength was
shown to be anticorrelated with small worldness *(34)*.
In the computational model, neither clustering coefficient which reflects local properties of the
functional network, nor global efficiency which reflects global properties of the functional network could
be explained by randomness, even at the lowest g_{coup} β-cell hubs **(Figure 7)**. This result was similar to
experimental findings which *reflect* a single 2-dimensional plane whereas the computational model
reflects the entire 3-dimensional network. According to experiments by Watts and Strogatz *(4)*, only the
most ordered networks will show zero overlap with randomness between the clustering and efficiency
statistics. Our results are unexpected because if gap junctions were solely responsible for variations in
islet synchronization, a decrease in gap junction coupling would cause the functional network to appear
more random. Instead, this indicates that β-cells are preferentially, not randomly, synchronized even in
the presence of very low gap junction coupling. This synchronization is likely to be explained by other
intrinsic cellular properties of the β-cells, which supports our previous claim that gap junctions are not
the only factor important in determining variations in cellular synchronization. The high clustering
coefficient found experimentally in Cx36 knockout islets **(Figure 6)** and computationally with low g_{coup}
indicates that cells tend to be functionally connected (i.e., highly synchronized) independent of variations
in gap junctions. These results again contribute to the evidence that gap junction connections are not the
only driver of synchronization. Of note, the partial overlap between global efficiency between Cx36
knockout islets and a random network suggests that gap junctions play some role in maintaining long-
range functional connections. This is consistent with our computational measurements that show slightly
elevated gap junction conductance in long-range (>4 cells) connections between synchronized cells
compared to non-synchronized cells. Further, the significant decrease in both clustering coefficient and

global efficiency of the functional network with decreasing Cx36 expression and g_{coup} indicates that as
structural gap junction connections are removed, the islet loses efficient signal propagation and high
ability to synchronize, which likely has pathophysiological implications. For example, this decreased
functional small worldness likely contributes to the lack of islet coordination and pulsatile insulin, in
diseases such as diabetes, where gap junction coupling has been shown to decrease *(25, 26, 51)*. Islet
coordination and pulsatile insulin release are essential for proper glucose homeostasis *(19–21)*.

<u>Broader implications</u>
We have shown evidence that the functional network of the islet of Langerhans is influenced by its
underlying structure but is more reflective of intrinsic cellular metabolic activity than gap junction
connections. Studies have shown that the hyper-polarization of highly synchronized β-cell hubs
disproportionately disrupts islet calcium dynamics, but the actual removal of hubs does not *(30, 32)*. Our
new perspective that β-cell hubs are uniquely defined by metabolic activity rather than gap junction
coupling, helps explain these results. β-cell hubs drive synchronization across the islet because they are
more metabolically active, not because they are more gap junction coupled. This perspective could also
be utilized when studying calcium dynamics, with the new understanding that synchronization can give
insight into cellular properties that drive excitability of the cell, such as the metabolic activity of β-cells.
Further, perturbations in metabolism, such as metabolic stress that occurs in diabetes, may have a
significant influence on the synchronization of the islet. While this study was conducted on experimental
and computational representations of the islets of Langerhans, we suggest that these findings may be
extrapolated to larger systems.

<u>Potential Limitations</u>
The most obvious limitation of this study, as for any computational study, is that computational models
do not perfectly reflect reality. Our computational model is based on careful physiological recordings
and has been previously validated *(38–40)*. We further address this limitation through experimental
studies which agree well with results from these computational models. However, a strength in our use
of the computational islet model is that all our results are internally derived. That is, even if the
computational model is not fully reflective of the experimental islet, the functional and structural
networks were obtained from the same model, so the relationship between them is not dependent on the
models' ability to reflect islet dynamics (though we have shown that it does). This fact also allows our
methods to translate to any oscillatory network with similar characteristics.
The choice of threshold when creating the functional network is influential in the results
(**Supplemental 1**) and should be carefully considered in any use of synchronization to create the
functional network. We picked our threshold to match that of the prior experimental results *(9, 10, 32)*.
The computational threshold was very high ($R_{th} = 0.9995$) because, unlike experiments, the
computational model had no noise, so the synchronization was more acute. However, the trends in our
results were consistent across thresholds (**Supplemental 1)**.
Finally, in the islet computational model we analyze only fast electrical oscillations, whose
frequency are less than 1 minute *(52)*. A similar analysis should eventually be done considering slower
metabolic oscillations and factoring in the (currently unknown) mechanism by which these metabolic
oscillations may be synchronized across the islet. We also assume that gap junction connections between
cells are constant throughout the time of analysis. Finally, the islet also contains α, δ, γ, and other cell

types which have been shown to influence β-cell dynamics through a number of mechanisms *(53–55)*. This study only analyzes β-cells which is the common practice among other network studies which we compare to *(9, 10, 31)*. However, we cannot exclude long-range structural projections from δ cells *(56, 57)* and that δ cells may be gap junction coupled *(59)*. In the future, network analysis should be applied to include other cell types as it could give valuable insight how their interactions can shape the overall islet response to nutrient changes.

Conclusions:

In this study, we investigate the relationship between the synchronization-based functional network and the structural network formed by gap junctions, within the β-cells in the islets of Langerhans. We conclude with three major points. *First*, highly connected subpopulations of β-cells, including β-cell hubs are driven by metabolism more than gap junctions. *Second*, metabolic similarities rather than gap junctions drive edges in the functional network of the islet. *Third,* removal of edges in the structural network decreases global communication efficiency and local connectivity. Specifically, we show that decreasing gap junction conductance causes the functional network to become more sparse and less small world-like, which can have pathophysiologic implications. These results provide insight into understanding the relationship between function and structure in islet networks and biological networks as a whole.

Methods:

<u>Calcium imaging, FRAP, and NAD(P)H</u>

Animal care: Male and female mice were used under protocols approved by the University of Colorado Institutional Animal Care and Use Committee. β-cell-specific GCaMP6s expression (β-GCaMP6s) was achieved through crossing a MIP-CreER (The Jackson Laboratory) and a GCaMP6s line (The Jackson Laboratory)*(26)*. Genotype was verified through qPCR (Transetyx, Memphis, TN). Mice were held in a temperature-controlled environment with a 12 h light/dark cycle and given continuous access to food and water. CreER-mediated recombination was induced by 5 daily doses of tamoxifen (50mg/kg bw in corn oil) delivered IP.

Islet isolation and culture: Islets were isolated from mice under ketamine/xylazine anesthesia (80 and 16 mg/kg) by collagenase delivery into the pancreas via injection into the bile duct. The collagenase-inflated pancreas was surgically removed and digested. Islets were handpicked and planted into the glass-bottom dishes (MatTek) using CellTak cell tissue adhesive (Sigma-Aldrich). Islets were cultured in RPMI medium (Corning, Tewksbury, MA) containing 10% fetal bovine serum, 100 U/mL penicillin, and 100 mg/mL streptomycin. Islets were incubated at 37C, 5% CO2 for 24-72 h before imaging.

Imaging: An hour prior to imaging nutrition media from the isolated islets was replaced by an imaging solution (125 mM NaCl, 5.7 mM KCl, 2.5 mM CaCl2, 1.2 mM MgCl2, 10 mM HEPES, and 0.1% BSA, pH 7.4) containing 2 mM glucose. During imaging the glucose level was raised to 11mM. Islets were imaged using either a LSM780 system (Carl Zeiss, Oberkochen, Germany) with a 40x 1.2 NA objective or with an LSM800 system (Carl Zeiss) with 20x 0.8 NA PlanApochromat objective, or a 40x 1.2 NA objective, with samples held at 37C.

For $[Ca^{2+}]$ measurements GCaMP6s fluorescence was excited using a 488-nm laser. Images were acquired at 1 frame/s at 10-20 m depth from the bottom of the islet. Glucose was elevated 3 minutes after the start of recording, unless stated otherwise.

NAD(P)H autofluorescence and $[Ca^{2+}]$ dynamics were performed in the same z-position within the islet. NADH(P)H autofluorescence was imaged under two-photon excitation using a tunable mode-locked Ti:sapphire laser (Chameleon; Coherent, Santa Clara, CA) set to 710 nm. Fluorescence emission was detected at 400–450 nm using the internal detector. Z-stacks of 6–7 images were acquired spanning a depth of 5 μm. First the NAD(P)H was recorded at 2 mM glucose, then the $[Ca^{2+}]$ dynamics was recorder at 2 mM and during transition to 11 mM glucose. After the $[Ca^{2+}]$ wave was established, the NAD(P)H was recorded at 11 mM glucose.

Cx36 gap junction permeability and $[Ca^{2+}]$ dynamics were performed in the same z-position within the islet, with gap junction permeability measured using fluorescence recovery after photobleaching (FRAP), as previously described *(43)*. After $[Ca^{2+}]$ imaging, islets were loaded with 12 mM Rhodamine-123 for 30 min at 37C in imaging solution. Islets were then washed and FRAP performed at 11mM glucose at room temperature. Room temperature was used because at this temperature the Rhodamine-123 travels between the cells only through the Cx36 gap junctions, versus at 37 C it can permeate a cell membrane. Rhodamine-123 was excited using a 488-nm laser line, and fluorescence emission was detected at 500–580 nm. Three baseline intensity images were initially recorded. A region of interest was then photobleached achieving, on average, a 50% decrease in fluorescence, and images were then acquired every 5-15 s for 15-30 min.

Analysis of [Ca2+] dynamics: Pearson-product-based network analysis presented in Figure 2 was
performed as previously reported *(9)*. [Ca^{2+}] time courses were analyzed during the second-phase [Ca^{2+}]
response, when the slow calcium wave was established. The Pearson product for each cell pair in the
islet was calculated over each time point, and the time-average values were computed to construct a
correlation matrix. An adjacency matrix was calculated by applying a threshold to the correlation matrix.
The same threshold of 0.9 was applied to all islets. All cell pairs with a non-zero values in the adjacency
matrix were considered to have a functional edge. The percent of edges was calculated with respect to
the maximum number of edges per cell in each individual islet. For example, if a most connected cell
possessed max=10 edges, and other cells had 1, 3, …7 edges – then the % were: 10%, 30%, …70%.

<u>Prior calcium imaging first presented in (13)</u>

Islet isolation: Islets were isolated as described in Scharp et al. (60) and Stefan et al. (61) and maintained
in Roswell Park Memorial. Institute medium containing 10% fetal bovine serum, 11 mM glucose at 37C
under humidified 5% CO2 for 24–48 h before imaging.

Imaging islets: Isolated islets were stained with 4 mM Fluo-4 AM (Invitrogen, Carlsbad, CA) in imaging
medium (125 mM NaCl, 5.7 mM KCl, 2.5 CaCl2, 1.2 mM MgCl2, 10 mM HEPES, 2mM glucose, 0.1%
bovine serum albumin, pH 7.4) at room temperature for 1–3 h before imaging. Islets were imaged in a
polydimethylsiloxane (PDMS) microfluidic device, the fabrication of which has been previously
described in Rocheleau et al. , which holds the islet stable for imaging and allows rapid reagent change,
such as varying glucose stimulation or adding gap junction inhibitors. Fluo-4 fluorescence is imaged 15
min after a step increase in glucose from low (2 mM) to high (11 mM). High speed imaging is performed
on an LSM5Live with a 203 0.8 NA Fluar Objective (Zeiss, Jena, Germany) using a 488 nm diode laser
for excitation and a 495 nm long-pass filter to detect fluorescence emission. The microfluidic device is
held on the microscope stage in a humidified temperature-controlled chamber, maintained at 37 C.
Images were acquired at a rate of 4–6 frames per second, with average powers at the sample being
minimized to, 200 mW/cm2.

Analysis of Ca2+ Imaging data: We present data from 11 Islets from 6 WT mice, 11 Islets from 11
heterozygous Cx36 +/− knockout mice, and 14 Islets from 3 homozygous Cx36 -/- knockout mice. We
extracted cell calcium dynamics by visually identifying and circling all cells in the islet. We assumed
that the pixels within a cell should be well coordinated, so we removed any pixels whose dynamics were
not within 5-10 STD of the average. This usually resulted in the removal of 1-5 pixels on the edge of the
cell boundary.

Threshold: Previous studies have shown that wild type Islets should have a degree distribution that is
roughly linear when plotted on a log-log plot and average degree of either 8 or between 5-15 *(9, 10, 32,*
63) . To determine R_{th} that best satisfied both of these findings, we utilized constrained optimization
matlab algorithm fminsearchbnd*(64)* to find the $R_{th_{opt}}$ which maximized the goodness of fit to a power
law distribution, while forcing $5 \leq k_{avg} \leq 15$ for each WT Islet constrained optimization. The average
optimal threshold ($R_{th} = 0.90$).

Average distance between connected cells: The average distance between connected cells calculated the
total number of pixels between the center of two connected cells. The average distance was expressed
as the normalized distance between connected cells and the average distance between all cells in the Islet
to control for image and Islet size.

<u>Computational Model:</u>

This ordinary differential equation model has been described previously (30) and validated on
experimental studies (38–41)**.** It is based on a single β-cell electrophysiologic model (65), in which the
membrane potential (V_i) for β-cell$_i$ is solved for each time step using eq 1a. We created a 1000 β-cell
network and electrically coupled any cell pairs within a small distance from each other. We chose 1000
β-cell because most species (including human and mouse) contain on average 1000 β-cells in an islet
(66) and it is the number which has been validated experimentally. The coupling current, I_{coup} is
determined by the difference in voltage between the cell pair and the average coupling conductance
(g_{coup}) between them (1b). All code was written in C++ and run on the SUMMIT supercomputer
(University of Colorado Boulder). All simulations are run at 11mMol glucose. The membrane potential
(V_i) for β-cell$_i$ is solved for using the ODE:

$$C_m \frac{dV_i}{dt} = I_{Cav} + I_{TRPM} + I_{SOC} + I_{bNSC} + I_{KDr} + I_{KCa(SK)} + I_{K_{ATP}} + I_{NaCa} + I_{PMCA} + I_{NaCa} \qquad (1a)$$
$$+ I_{Coup}$$

$$I_{Coup} = \sum_i g_{coup}^{ij}(V_i - V_j) \qquad (1b)$$

The average number of gap junction connections per cell was 5.24. Cellular heterogeneity was
introduced by assigning randomized metabolic and electrical parameter values to each cell based on their
distributions previously determined by experimental studies. For example, coupling conductance (g_{coup})
was assigned so that the Islet average is $g_{coup} = 0.12nS$ and glucokinase (k_{glyc}), the rate-limiting step
in glycolysis, was assigned so that the average $k_{glyc} = 1.26 * 10^{-04} \ sec^{-1}$. We ran the simulation for
500 seconds and only the 2nd phase was analyzed, in concordance with *(10)*. All the results presented are
based on five different Islets with randomized parameter values. β-cell heterogeneity was simulated by
randomizing parameters including coupling conductance (g_{coup}) and metabolic rate (k_{glyc}) according
to experimental results. To explore the effects of coupling conductance on the functional network, we
altered Islet average coupling conductivity to $g_{coup} = 0.06 \ nS$ and $g_{coup} = 0.03 \ nS$, and then assigning
each cell a randomized g_{coup} based on the new target average. The gap junction conductance (g_{coup}) for
any cell was calculated as the sum of the average conductivity (g_{coup}) between the cell of interest and
any cells it shares a gap junction connection with:

$$g_{coup_j} = \Sigma_{i=1}^{m} \frac{1}{2} (g_{coup}^{i} + g_{coup}^{j})$$

Network Analysis:

Creating Functional Network from simulated Islet: The methodology was based on that previously defined in *(9)*. First, the correlation coefficient between each cell was calculated using corr() function in matlab, which follows equation: $R_{ij} = \frac{\sum[\bar{x}_i - x_i(t)][\bar{x}_j - x_j(t)]}{s_{x_i} s_{x_j}}$. Next, a threshold ($R_{th}$) was defined to determine whether each cell pair is 'synchronized' or 'not synchronized'. For computational experiments, the threshold was chosen such that the network roughly followed a power law distribution, as predicted in *(9, 10)*. Unless otherwise noted, $R_{th} = 0.9995$ for computational analysis.

Creating the Metabolic network: Because gap junctions enforce localization onto the analysis, we looked at cell pairs whose Pythagorean distance was $\leq 15\%$ of the Islet. Within this sample space, we looked at cell pairs whose average k$_{glyc}$ was higher than the Islet average (**Figure 4b**, green) ($0.5 * \left|k_{glyc_i} + k_{glyc_j}\right| > average(0.5 * \left|k_{glyc_i} + k_{glyc_j}\right| \forall_{ij})$), and cell pairs whose K$_{glyc}$ was more similar than the islet average ($\left|k_{glyc_i} - k_{glyc_j}\right| < average(\left|k_{glyc_i} - k_{glyc_j}\right| \forall_{ij})$). We chose these parameters such that the $Pr(Sync|GJ) = Pr(Sync|Met)$, which acts as a control allowing us to directly compare $Pr(GJ|Sync)$ to $Pr(Met|Sync)$.

Average degree: The degree k_i was calculated by counting the number of connections for $cell_i$ and averaging k over all cells in the Islet, then normalized to the Islet size to remove any size dependence ($\frac{k_{avg}}{n}$, $where\ n \equiv islet\ size$).

Degree Distribution: The degree distribution was calculated by first calculating the degree of each cell by taking the column sum of the adjacency matrix. Then each cell degree was normalized by dividing by the maximum degree in the Islet. The histogram was then calculated using GraphPad Prism.

Hub identification: In accordance with , any cell with more than 60% of the Islet's connections, calculated by the degree distribution, was considered a hub cell.

Probabilities:

We first created the functional network and structural (GJ related probabilities) or metabolic network (metabolism). We then calculated the probability that two cells were synchronized by:

$$Pr(sync) = \frac{\frac{m_{sync}}{(n-1)*n}}{2}$$

Where m_{sync} = number of edges in the synchronized network, and n = number of nodes. Similarly, we calculated the probability that two cells were either gap junction coupled or metabolically related using

$$Pr(GJ\ or\ k_{glyc}) = \frac{\frac{m_{GJ\ or\ k_{glyc}}}{(n-1)*n}}{2}$$

Where $m_{GJ\ or\ k_{glyc}}$ is the number of edges in the gap junction or metabolic network.

773 To find the probability that a cell pair was both synchronized and GJ or metabolically connected, we
774 calculated

$$\Pr(both) = \frac{\frac{m_{both}}{(n-1)*n}}{2}$$

776 Where m_{both} is any edge that exists in both matrices.
777 Finally, we calculated conditional probabilities by:

$$\Pr(sync|GJ) = \frac{\Pr(both)}{\Pr(GJ)}$$

$$\Pr(GJ|sync) = \frac{\Pr(both)}{\Pr(sync)}$$

780 These quantities were calculated separately for each Islet seed and then averaged.
781
<u>Network topology analysis:</u>
Shortest weighted path length: The gap junction network was weighted using the inverse of g_{coup} or k_{glyc}
between $Cell_i$ and $Cell_i$: $W_{ij} = \frac{1}{\frac{1}{2}(g_{coup_i} + g_{coup_j})}$ or $W_{ij} = \frac{1}{\frac{1}{2}(k_{glyc_i} + k_{glyc_j})}$. This is done because the
shortest path length algorithm views weights as "resistance" and finds the path of least resistance. The
shortest weighted path between every cell was calculated using Johnson's algorithm (67). Cell pairs were
then categorized as synchronized or not synchronized if their correlation coefficient was $< R_{th}$. The
average for synchronized and non-synchronized cells was calculated over each Islet for each distance.
To normalize over distance, each data point was divided by the average of the non-synchronized and
synchronized islets for the given distance.

Clustering Coefficient: The clustering coefficient represents the 'cliquishness' of the network . This is
defined by *(9)* as the "number of existing connections between all neighbors of a node, divided by the
number of all possible connections between them." This was calculated by making a subgraph of each
cell's connections and counting the number of connections between those cells. For example, if A is
connected to B, C, and D, and B and C are connected but D is not connected to any other cell (see
matrix). Then the clustering for cell A is $\frac{2}{3*2} = \frac{1}{3}$. Each node is a assigned a clustering coefficient C such
that:

Subgraph - A	B	C	D
B		1	0
C	1		0
D	0	0	

The average clustering coefficient is $C_{avg} = \frac{1}{n}\sum_i^n C_i$.
Average Shortest Path Length: Shortest path length was calculated with matlab function
graphallshortestpaths(). This function uses the Johnson's algorithm *(67)* to find the shortest path between
every pair of cell in the Islet. For example, the path length between $cell_i$ and $cell_j$ is 1 if they are directly
synchronized, or 2 if $cell_i$ is not synchronized with $cell_j$ but each is synchronized with $cell_k$. To
compensate for highly sparse network, any non-connected node was given a characteristic path length

of $n + 1$. Finally, the characteristic path length (L), or average path length, was expressed as the sum of
all path lengths normalized to total possible connections (size*(size-1)).
The normalized average shortest path length (Supplemental 2d) is therefore calculated as $L_{avg} =$
$\frac{1}{n}\frac{1}{n}\sum_{i=1}^{n}\sum_{j=1}^{n}L_{ij}$. To compensate for any non-connected cell, whose path length $L_{ix} = \infty$, where x is
any cell in the Islet, we set the path length for non-connected cells to $L_{ix} \rightarrow n + 1$ where n is the number
of cells in the Islet.

Global Efficiency: The global efficiency is related to the inverse of global path length *(45)*.

811
$$E_{\{global\}} = \frac{1}{Islet_{size} * (Islet_{size} - 1)}\sum_{jk}\frac{1}{L_{jk}}$$

length using $E_{global} = \frac{1}{n(n-1)}\sum_{i=1}^{n}\sum_{j=1}^{n}\frac{1}{L_{ij}}$. Because $E_{global} = 0$ for a non-connected cell, the
disconnected network is naturally compensated for.

Random Networks: Random networks were created using an Erodos Renyi approach (68). For each Islet,
we created 1000 random networks with equal number of nodes, edges, and average degree as the Islet
of interest. The probability that two cells were connected was given by $P = \frac{k_{avg}}{n}$. For each cell pair within
the Islet, a random number generator created a number from 0-1. If that number was below P, those cells
were connected by an edge.

Small World properties: For each Islet, small world parameters S_E and S_L were calculated 1000 times
based on the 1000 random networks.

822
$$S_L = \frac{\frac{C_{avg}}{C_{rand}}}{\frac{L_{avg}}{L_{rand}}}$$

823
$$S_E = \frac{\frac{C_{avg}}{C_{rand}}}{\frac{E_{rand}}{E_{avg}}}.$$

824 *Local Efficiency:* Local efficiency is calculated by first extracting a subgraph based on the connections
825 for an individual cell. Within that cell, the efficiency is calculated by:

826
$$E = \frac{1}{n_i * (n_i - 1)}\sum\frac{1}{L_{j,k}}$$

Where n_i is the number of nodes that an individual node is connected to. This is done for every node in
the network and all efficiencies are averaged to present the local efficiency of the network.

<u>Significance Testing:</u>
Significance tests are done in Prism (GraphPad).
The computational model is run on 5 seeds, therefore all significance tests are paired.
Figure 1e-g, Figure 4e&h, and Figure 5c-f are paired, two tailed t-tests.

Figure 3e,f are repeated measures paired one-way ANOVA with multiple comparison using Tukey's
multiple comparison.
Figure 4i,j are multiple paired t-tests with Bonferroni-Dunn adjustment.
All significance tests in figure 6 are unpaired ordinary one-way ANOVA tests.
Figure 7 c-j are ordinary paired one-way ANOVA with Tukey multiple comparison test.
Supplemental 1b and 2b, Figure 4 i and j and Figure 5i: To compare, we used a paired T-test with
Bonferroni multiple comparisons corrections. Since we there were 15 cell distances, we set the
significance threshold $\alpha = 0.003$. For convenience, we present asterisks next to significant P-values.
Supplemental 4b and c are two way ANOVA with Sidak's multiple comparisons
842

References

1. M. Koutrouli, E. Karatzas, D. Paez-Espino, G. A. Pavlopoulos, A guide to conquer the biological network era using graph theory. *Frontiers in bioengineering and biotechnology* **8**, 34 (2020).

2. S. H. Strogatz, Exploring complex networks. *Nature* **410**, 268–276 (2001).

3. M. E. Newman, The structure and function of complex networks. *SIAM review* **45**, 167–256 (2003).

4. D. J. Watts, S. H. Strogatz, Collective dynamics of 'small-world'networks. *nature* **393**, 440–442 (1998).

5. C. W. Lynn, D. S. Bassett, The physics of brain network structure, function and control. *Nature Reviews Physics* **1**, 318–332 (2019).

6. E. Bullmore, O. Sporns, Complex brain networks: graph theoretical analysis of structural and functional systems. *Nature reviews neuroscience* **10**, 186–198 (2009).

7. M. Newman, *Networks* (Oxford university press, 2018).

8. D. R. Chialvo, Emergent complex neural dynamics. *Nature physics* **6**, 744–750 (2010).

9. A. Stožer, M. Gosak, J. Dolenšek, M. Perc, M. Marhl, M. S. Rupnik, D. Korošak, Functional connectivity in islets of Langerhans from mouse pancreas tissue slices. *PLoS Comput Biol* **9**, e1002923 (2013).

10. N. R. Johnston, R. K. Mitchell, E. Haythorne, M. P. Pessoa, F. Semplici, J. Ferrer, L. Piemonti, P. Marchetti, M. Bugliani, D. Bosco, others, Beta cell hubs dictate pancreatic islet responses to glucose. *Cell metabolism* **24**, 389–401 (2016).

11. D. J. Hodson, M. Schaeffer, N. Romanò, P. Fontanaud, C. Lafont, J. Birkenstock, F. Molino, H. Christian, J. Lockey, D. Carmignac, others, Existence of long-lasting experience-dependent plasticity in endocrine cell networks. *Nature communications* **3**, 1–10 (2012).

12. C. Grienberger, A. Konnerth, Imaging calcium in neurons. *Neuron* **73**, 862–885 (2012).

13. R. K. Benninger, M. Zhang, W. S. Head, L. S. Satin, D. W. Piston, Gap junction coupling and calcium waves in the pancreatic islet. *Biophysical journal* **95**, 5048–5061 (2008).

14. V. Serre-Beinier, S. Le Gurun, N. Belluardo, A. Trovato-Salinaro, A. Charollais, J.-A. Haefliger, D. F. Condorelli, P. Meda, Cx36 preferentially connects beta-cells within pancreatic islets. *Diabetes* **49**, 727–734 (2000).

15. M. A. Ravier, M. Güldenagel, A. Charollais, A. Gjinovci, D. Caille, G. Söhl, C. B. Wollheim, K. Willecke, J.-C. Henquin, P. Meda, Loss of connexin36 channels alters β-cell coupling, islet synchronization of glucose-induced Ca2+ and insulin oscillations, and basal insulin release. *Diabetes* **54**, 1798–1807 (2005).

16. R. K. Benninger, V. Kravets, The physiological role of β-cell heterogeneity in pancreatic islet
function. *Nature Reviews Endocrinology* **18**, 9–22 (2022).

17. J. Weitz, D. Menegaz, A. Caicedo, Deciphering the Complex Communication Networks That
Orchestrate Pancreatic Islet Function. *Diabetes* **70**, 17–26 (2021).

18. W. S. Head, M. L. Orseth, C. S. Nunemaker, L. S. Satin, D. W. Piston, R. K. Benninger, Connexin-
36 gap junctions regulate in vivo first-and second-phase insulin secretion dynamics and glucose
tolerance in the conscious mouse. *Diabetes* **61**, 1700–1707 (2012).

19. N. Pørksen, M. Hollingdal, C. Juhl, P. Butler, J. D. Veldhuis, O. Schmitz, Pulsatile insulin
secretion: detection, regulation, and role in diabetes. *Diabetes* **51**, S245–S254 (2002).

20. A. V. Matveyenko, D. Liuwantara, T. Gurlo, D. Kirakossian, C. Dalla Man, C. Cobelli, M. F.
White, K. D. Copps, E. Volpi, S. Fujita, others, Pulsatile portal vein insulin delivery enhances hepatic
insulin action and signaling. *Diabetes* **61**, 2269–2279 (2012).

21. P. Bratusch-Marrain, M. Komjati, W. Waldhäusl, Efficacy of pulsatile versus continuous insulin
administration on hepatic glucose production and glucose utilization in type I diabetic humans.
Diabetes **35**, 922–926 (1986).

22. L. S. Satin, P. C. Butler, J. Ha, A. S. Sherman, Pulsatile insulin secretion, impaired glucose
tolerance and type 2 diabetes. *Mol Aspects Med* **42**, 61–77 (2015).

23. D. J. Hodson, R. K. Mitchell, E. A. Bellomo, G. Sun, L. Vinet, P. Meda, D. Li, W.-H. Li, M.
Bugliani, P. Marchetti, others, Lipotoxicity disrupts incretin-regulated human β cell connectivity. *The*
Journal of clinical investigation **123**, 4182–4194 (2013).

24. D. A. Lang, D. Matthews, M. Burnett, R. Turner, Brief, irregular oscillations of basal plasma
insulin and glucose concentrations in diabetic man. *Diabetes* **30**, 435–439 (1981).

25. J. R. St Clair, M. J. Westacott, N. L. Farnsworth, V. Kravets, W. E. Schleicher, J. G. Miranda, A.
Heintz, N. W. Ludin, R. K. Benninger, Restoring connexin-36 function in diabetogenic environments
precludes mouse and human islet dysfunction. *bioRxiv* (2020).

26. M. E. C. do Amaral, V. Kravets, J. M. Dwulet, N. L. Farnsworth, R. Piscopio, W. E. Schleicher, J.
G. Miranda, R. K. Benninger, Caloric Restriction recovers impaired β-cell-β-cell coupling, calcium
oscillation coordination and insulin secretion in prediabetic mice. *bioRxiv* (2020).

27. M. Gosak, R. Marković, J. Dolenšek, M. S. Rupnik, M. Marhl, A. Stožer, M. Perc, Network
science of biological systems at different scales: A review. *Physics of life reviews* **24**, 118–135 (2018).

28. R. Marković, A. Stožer, M. Gosak, J. Dolenšek, M. Marhl, M. S. Rupnik, Progressive glucose
stimulation of islet beta cells reveals a transition from segregated to integrated modular functional
connectivity patterns. *Sci Rep* **5**, 7845 (2015).

29. A. D. Broido, A. Clauset, Scale-free networks are rare. *Nature communications* **10**, 1–10 (2019).

30. J. M. Dwulet, J. K. Briggs, R. K. P. Benninger, Small subpopulations of beta-cells do not drive
islet oscillatory [Ca2+] dynamics via gap junction communication. *PLOS Computational Biology* **17**,
e1008948 (2021).

31. C.-L. Lei, J. A. Kellard, M. Hara, J. D. Johnson, B. Rodriguez, L. J. Briant, Beta-cell hubs maintain
Ca2+ oscillations in human and mouse islet simulations. *Islets* **10**, 151–167 (2018).

32. D. Korošak, M. Jusup, B. Podobnik, A. Stožer, J. Dolenšek, P. Holme, M. S. Rupnik, Autopoietic
influence hierarchies in pancreatic β-cells. *arXiv:2101.02255 [cond-mat, physics:physics]* (2021)
(available at http://arxiv.org/abs/2101.02255).

33. L. S. Satin, Q. Zhang, P. Rorsman, "Take Me To Your Leader": An Electrophysiological Appraisal
of the Role of Hub Cells in Pancreatic Islets. *Diabetes* **69**, 830–836 (2020).

34. J. P. Hogan, B. E. Peercy, Flipping the switch on the hub cell: Islet desynchronization through cell
silencing. *PloS one* **16**, e0248974 (2021).

35. G. A. Rutter, N. Ninov, V. Salem, D. J. Hodson, Comment on Satin et al."Take me to your leader":
an electrophysiological appraisal of the role of hub cells in pancreatic islets. Diabetes 2020; 69: 830–
836. *Diabetes* **69**, e10–e11 (2020).

36. L. S. Satin, P. Rorsman, Response to comment on satin et al."Take me to your leader": An
electrophysiological appraisal of the role of hub cells in pancreatic islets. Diabetes 2020; 69: 830–836.
Diabetes **69**, e12–e13 (2020).

37. B. E. Peercy, A. S. Sherman, Do oscillations in pancreatic islets require pacemaker cells? *Journal
of Biosciences* **47**, 1–11 (2022).

38. M. J. Westacott, N. W. Ludin, R. K. Benninger, Spatially organized β-cell subpopulations control
electrical dynamics across islets of Langerhans. *Biophysical journal* **113**, 1093–1108 (2017).

39. J. M. Dwulet, N. W. Ludin, R. A. Piscopio, W. E. Schleicher, O. Moua, M. J. Westacott, R. K.
Benninger, How heterogeneity in glucokinase and gap-junction coupling determines the islet [Ca2+]
response. *Biophysical journal* **117**, 2188–2203 (2019).

40. T. H. Hraha, M. J. Westacott, M. Pozzoli, A. M. Notary, P. M. McClatchey, R. K. Benninger,
Phase transitions in the multi-cellular regulatory behavior of pancreatic islet excitability. *PLoS
computational biology* **10**, e1003819 (2014).

41. A. M. Notary, M. J. Westacott, T. H. Hraha, M. Pozzoli, R. K. P. Benninger, Decreases in Gap
Junction Coupling Recovers Ca2+ and Insulin Secretion in Neonatal Diabetes Mellitus, Dependent on
Beta Cell Heterogeneity and Noise. *PLOS Computational Biology* **12**, e1005116 (2016).

42. V. Salem, L. D. Silva, K. Suba, E. Georgiadou, S. N. M. Gharavy, N. Akhtar, A. Martin-Alonso, D. C. Gaboriau, S. M. Rothery, T. Stylianides, others, Leader beta-cells coordinate Ca 2+ dynamics across pancreatic islets in vivo. *Nature Metabolism* **1**, 615–629 (2019).

43. N. L. Farnsworth, A. Hemmati, M. Pozzoli, R. K. Benninger, Fluorescence recovery after photobleaching reveals regulation and distribution of connexin36 gap junction coupling within mouse islets of Langerhans. *The Journal of physiology* **592**, 4431–4446 (2014).

44. G. Cappon, M. G. Pedersen, Heterogeneity and nearest-neighbor coupling can explain small-worldness and wave properties in pancreatic islets. *Chaos: An Interdisciplinary Journal of Nonlinear Science* **26**, 053103 (2016).

45. V. Latora, M. Marchiori, Efficient behavior of small-world networks. *Physical review letters* **87**, 198701 (2001).

46. M. T. Adams, J. M. Dwulet, J. K. Briggs, C. A. Reissaus, E. Jin, J. M. Szulczewski, M. R. Lyman, S. M. Sdao, V. Kravets, S. D. Nimkulrat, others, Reduced synchroneity of intra-islet Ca2+ oscillations in vivo in Robo-deficient β cells. *Elife* **10** (2021).

47. R. K. Benninger, D. J. Hodson, New understanding of β-cell heterogeneity and in situ islet function. *Diabetes* **67**, 537–547 (2018).

48. N. L. Farnsworth, R. K. Benninger, New insights into the role of connexins in pancreatic islet function and diabetes. *FEBS letters* **588**, 1278–1287 (2014).

49. R. Plonsey, R. C. Barr, A. Bioelectricity, *Quantitative Approach* (Springer, 2007).

50. A. Sherman, J. Rinzel, Model for synchronization of pancreatic beta-cells by gap junction coupling. *Biophysical journal* **59**, 547–559 (1991).

51. N. L. Farnsworth, R. L. Walter, A. Hemmati, M. J. Westacott, R. K. Benninger, Low level pro-inflammatory cytokines decrease connexin36 gap junction coupling in mouse and human islets through nitric oxide-mediated protein kinase Cδ. *Journal of Biological Chemistry* **291**, 3184–3196 (2016).

52. R. Bertram, A. Sherman, L. S. Satin, Electrical bursting, calcium oscillations, and synchronization of pancreatic islets. *The Islets of Langerhans* , 261–279 (2010).

53. J.-C. Henquin, Paracrine and autocrine control of insulin secretion in human islets: evidence and pending questions. *American Journal of Physiology-Endocrinology and Metabolism* **320**, E78–E86 (2021).

54. T. Moede, I. B. Leibiger, P.-O. Berggren, Alpha cell regulation of beta cell function. *Diabetologia* **63**, 2064–2075 (2020).

55. T. Van Der Meulen, C. J. Donaldson, E. Cáceres, A. E. Hunter, C. Cowing-Zitron, L. D. Pound, M.
W. Adams, A. Zembrzycki, K. L. Grove, M. O. Huising, Urocortin3 mediates somatostatin-dependent
negative feedback control of insulin secretion. *Nature medicine* **21**, 769–776 (2015).

56. R. Arrojo e Drigo, S. Jacob, C. F. García-Prieto, X. Zheng, M. Fukuda, H. T. T. Nhu, O.
Stelmashenko, F. L. M. Peçanha, R. Rodriguez-Diaz, E. Bushong, others, Structural basis for delta cell
paracrine regulation in pancreatic islets. *Nature communications* **10**, 1–12 (2019).

57. L. Briant, T. Reinbothe, I. Spiliotis, C. Miranda, B. Rodriguez, P. Rorsman, δ-cells and β-cells are
electrically coupled and regulate α-cell activity via somatostatin. *The Journal of physiology* **596**, 197–
215 (2018).

58. R. A. e Drigo, S. Jacob, C. F. García-Prieto, X. Zheng, M. Fukuda, H. T. T. Nhu, O. Stelmashenko,
F. L. M. Peçanha, R. Rodriguez-Diaz, E. Bushong, others, Structural basis for delta cell paracrine
regulation in pancreatic islets. *Nature communications* **10**, 1–12 (2019).

59. C. Miranda, M. Begum, E. Vergari, L. J. Briant, Gap junction coupling and islet delta-cell function
in health and disease. *Peptides* **147**, 170704 (2022).

60. D. W. SCHARP, C. B. KEMP, M. J. KNIGHT, W. F. BALLINGER, P. E. LACY, The use of
ficoll in the preparation of viable islets of langerhans from the rat pancreas. *Transplantation* **16**, 686–
688 (1973).

61. Y. Stefan, P. Meda, M. Neufeld, L. Orci, others, Stimulation of insulin secretion reveals
heterogeneity of pancreatic B cells in vivo. *The Journal of clinical investigation* **80**, 175–183 (1987).

62. J. V. Rocheleau, M. S. Remedi, B. Granada, W. S. Head, J. C. Koster, C. G. Nichols, D. W. Piston,
Critical role of gap junction coupled KATP channel activity for regulated insulin secretion. *PLoS*
biology **4**, e26 (2006).

63. J. Zmazek, M. S. Klemen, R. Markovič, J. Dolenšek, M. Marhl, A. Stožer, M. Gosak, Assessing
different temporal scales of calcium dynamics in networks of beta cell populations. *Frontiers in*
physiology **12**, 337 (2021).

64. J. D'Errico, fminsearchbnd, fminsearchcon
(https://www.mathworks.com/matlabcentral/fileexchange/8277-fminsearchbnd-fminsearchcon).
MATLAB Central File Exchange .

65. C. Y. Cha, Y. Nakamura, Y. Himeno, J. Wang, S. Fujimoto, N. Inagaki, Y. E. Earm, A. Noma,
Ionic mechanisms and Ca2+ dynamics underlying the glucose response of pancreatic beta cells: a
simulation study. *Journal of General Physiology* **138**, 21–37 (2011).

66. A. Pisania, G. C. Weir, J. J. O'neil, A. Omer, V. Tchipashvili, J. Lei, C. K. Colton, S. Bonner-
Weir, Quantitative analysis of cell composition and purity of human pancreatic islet preparations.
Laboratory investigation **90**, 1661–1675 (2010).

67. D. B. Johnson, Efficient algorithms for shortest paths in sparse networks. *Journal of the ACM (JACM)* **24**, 1–13 (1977).

68. P. Erd\Hos, A. Rényi, On the evolution of random graphs. *Publ. Math. Inst. Hung. Acad. Sci* **5**, 17–60 (1960).

Acknowledgements

Richard KP Benninger (University of Colorado) is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. We thank David W Piston (Washington University St Louis) with whom data presented in figure 6 was previously acquired. We thank Aaron Clauset for helpful conversations and advice. All authors acknowledge that no conflict of interest exists.

Funding:

National Institute of Health (NIH) grant R01 DK102950 (RKPB)

National Institute of Health (NIH) grant R01 DK106412 (RKPB)

National Science Foundation (NSF) Graduate Research Fellowship DGE-1938058_Briggs (JKB)

Juvenile Diabetes Research Foundation (JDRF) grant 3-PDF-2019-741-A-N (VK)

Beckman Research Institute-City of Hope (HIRN) grant 25B1104 (VK)

National Institute of Health (NIH) grant F31 DK126360 (JMD).

The authors are grateful for utilization of the SUMMIT supercomputer from the University of Colorado Boulder Research Computing Group, which is supported by the National Science Foundation (awards ACI-1532235 and ACI-1532236), the University of Colorado Boulder, and Colorado State University. Microscopy use was supported in part by NIH grant P30 DK116073 and the University of Colorado Neurotechnology center.

Author Contributions:

Conceptualization: JKB, RKPB, VK

Methodology: JKB, RKPB, VK, JMD

Investigation: JKB, VK

Visualization: JKB

Supervision: RKPB

Writing—original draft: JKB

Writing—review & editing: JKB, RKPB, VK, DJA, JMD

Competing interests: The funders had no role in the study design, data collection, and analysis, decisions to publish, or preparation of the manuscript.

Data and materials availability:

Raw microscopy imaging data to the EMBL-EBI-supported BioImage Archive. Analysis code, Model code, and Simulated Data will be available via GitHub at https://github.com/jenniferkbriggs/Functional_and_Structural_Networks.git

1047 **Tables**

1048

Table 1: Essential network statistics and types
Left: Five networks statistics used to quantify the network in this paper. Representative networks
show an example of the statistic in red and the rest of the network in blue. Right: Five network
types referred to in this paper. Regular, small world, and random networks are all made with the
same number of nodes and edges, but different configurations of edges. Scale free network
shows three "hub" nodes where node size is proportional to degree.
1055

1056   **Figures**

1057

Figure 1: Analysis of parameters underlying highly connected cells in a simulated islet
network.
a: schematic of 1000 β-cell computational model, with cells false-colored by heterogeneity in k_{glyc}
(left) and g_{coup} (right) parameter values. **b:** Distribution of functional connections (edges),

determined from functional network analysis. Colors show five seeds. Hub cells (red outline) are
any cell with $> 60\%$ of maximum number of links. The R^2 for linear fit was 0.559. **c:** 2-
dimensional slice of the simulated islet, with lines (edges) representing functional connections
between synchronized cells. Hub cells indicated in red. Slice is taken from middle of the Islet (see
inset). **d:** Representative [Ca^{2+}] time-course of a hub (red) and non-hub (blue). Inset shows zoomed
enlarged signal from $45 - 80$ seconds. **e:** Average rate of metabolism (k_{glyc}) parameter values
compared between hub and non-hub, retrospectively analyzed. **f:** as in e for maximum conductance
of ATP sensitive potassium channel (g$_{KATP}$). **g:** as in e for gap junction conductance
(g_{coup}). Significance in e-g was determined by paired Students t-test. $*P \leq 0.05$ and $***P \leq$
0.001
1072

1073

Figure 2: Comparison of the calcium-based network with the metabolic activity, NAD(P)H and coupling conductance

a: Mouse pancreatic islet expressing fluorescent calcium indicator GCaMP6s in β-cells. Glucose level 11 mM. **b:** GCamp6s time traces recorded at 2- and 11-mM glucose. Red curves represent dynamics of the most coordinated cells. These cells had highest number of edges, i.e., normalized degree >0.9. Green curves represent dynamics of the least coordinated cells, i.e., normalized degree <0.1, the rest of the cells are shown in grey. **c:** Ca^{2+} - based functional network of the islet shown in (e). Red dots represent most coordinated cells, and green dots – least coordinated cells which had at least 1 edge. **d:** Degree distribution of all eight analyzed islets. Threshold of R_{th}=0.9 was used to filter the Pearson-based coordination matrix and obtain the functional network. **e:** *Left:* Mouse pancreatic islet expressing fluorescent calcium indicator GCaMP6s. Glucose level 11 mM. *Middle:* NAD(P)H autofluorescence of the same islet and the same cell layer, recorded at 2 mM glucose. *Right: :* NAD(P)H autofluorescence recorded at 11 mM glucose. **f:** Change of the NAD(P)H levels in each cell in response to glucose elevation, with respect to the islet-average change. Metabolic activity here is compared for connected and disconnected cells. **g:** Change in NAD(P)H levels in response to glucose elevation, with respect to the islet-average change: here comparison is made for the most coordinated cells (normalized degree >0.9) with the less

coordinated cells (normalized degree >0.8, >0.7, >0.6, …<0.2, <0.1). f,g represent n=5 islets with

a total of 131 cells. **h:** Rhodamine123 fluorescence of the same islet and the same cell layer as in

a, recorded immediately after photobleaching of the bottom half of the islet **(left)** and recorded at

360 seconds after the photobleaching **(right)**. **i:** Time trace of the fluorescence recovery after

photobleaching (FRAP) of the lower half of the islet, shown in **h**. The rate of recovery is

proportional to the Cx36 gap junction permeability of each cell. **j:** FRAP recovery rate (sec^{-1}) in

each of the photobleached cells: here the comparison is made for the most coordinated cells

(normalized degree >90%) with the less coordinated cells (normalized degree >80, >70, >60,

…<20, <10%). j shows results from 4 islets and 202 cells). All the cells in this graph had a degree

larger than 0.01.

1101

1102

Figure 3: Comparison of functional connections, gap junction junctions, and glucose metabolism

a: Functional connections and structural connections for a representative highly connected β-cell hub, average cell, and low connected cell. Blue cells and edges indicate cells determined as "synchronized" via functional network analysis. Red cells are GJ coupled. Black cells are both GJ coupled and synchronized. **b:** Probability that a cell is synchronized given it is gap junction coupled, or vice versa, that a cell is gap junction coupled given it is synchronized. **c:** Ven diagram showing overlap between the synchronized cells and gap junction coupled cells within the simulated islet. **d:** Functional connections and metabolic connections for a β-cell hub and a cell with averaged synchronization. Blue cells and edges indicate cells determined as "synchronized" via functional network analysis. Green cells and edges indicate cells that have similar k_{glyc} and are within 15% of the islet from each other. Black cells and edges indicate both synchronized and similar k_{glyc}. **e:** Probability that a cell pair is synchronized given it shares a gap junction, has high average k_{glyc} or has similar k_{glyc}. **f:** Probability that the cell pair is gap junction coupled, has higher than average k_{glyc} and within 15% of islet distance ($Pr = 0.0050$), or has similar k_{glyc} and within 15% of islet distance ($Pr = 0.0052$), respectively, given the pair is synchronized. **g:** Ven diagram showing overlap between the synchronized cells and metabolically similar cells. Shaded area in c and g is proportional to indicated probability. Significance in e and f was determined by a repeated measures one-way ANOVA with Tukey's multiple comparisons **$P \leq 0.01$.

1122

1123

Figure 4: Comparison of long-range functional synchronization, gap junction network, and glucose metabolism.

a: Functional connections and gap junction connections for a representative cell (black). Synchronized cells in blue, gap junction coupled cells in red. Inset shows representative structural network (grey). **b:** Two representative cells (black) and the shortest path to a synchronized cell

(blue) and a non-synchronized cell (green). Path is weighted (shown by edge thickness) by g_{coup}.

For the cell on the top panel, the synchronized path has a higher cumulative gap junction

conductivity (0.68 nS) than the non synchronized path (0.43 nS). for the cell on the bottom panel,

the synchronized path has a lower cumulative gap junction conductance (0.74 nS) than the non

synchronized path (1.02 nS). **c:** Probability distribution of total g_{coup} for synchronized cells (blue)

and non-synchronized cells (green) that are directly connected by gap junctions. **d:** As in c for

cells pairs that are 7 cells apart. **e:** Comparison of the total g_{coup}, normalized by distance for

synchronized cells (blue) and non-synchronized cells (green). Each dot indicates the average

resistance for a single simulated islet. **f:** As in c but with connections weighted by rate of

metabolism. **g:** As in d but with connections weighted by the rate of glucose metabolism. **h:** As

in e but with connections weighted by the rate of glucose metabolism. **i:** Total g_{coup} for

synchronized and non synchronized cells organized by cell distance. **j:** As in i but for metabolically

weighted network. Significance in e,h was assessed by a two-tailed paired t-test, with P value

indicated. Significance in i and j was, assessed by paired t-tests with Bonferroni correction.

Reported P values are adjusted. $*P \leq 0.05$, $**P \leq 0.01$, $***P \leq 0.001$.

Figure 5

Figure 5: Assessment of functional and structural network overlap in a correlated distribution of glucose metabolism and gap junction conductance

a: Distribution where g_{coup} is correlated with k_{glyc}, such that cells with high metabolism also have high gap junction coupling. **b:** 2-dimensional slice of the simulated islet, with lines (edges) representing functional connections between synchronized cells. β-cell hubs are indicated in red. Slice is taken from middle of the Islet (see inset). **c:** Average degree comparing proportional distribution shown in a and normal distribution (Figure 1), where there is no relationship between k_{glyc} and g_{coup} **d:** Average rate of glucose metabolism (k_{glyc}) parameter values compared between hub and non-hub, retrospectively analyzed. **e:** as in d for gap junction conductance (g_{coup}). **f:** as

in d for maximum conductance of ATP sensitive potassium channel (g_{KATP}). **g:** probability of synchronization given a cell pair is gap junction coupled P(Sync|GJ), and probability that a cell pair which is gap junction coupled, is synchronized P(GJ|Sync). **h:** Venn diagram showing overlap between synchronized cell pairs and gap junction connected cell pairs. **i:** Total g_{coup} for synchronized and not synchronized cell pairs organized by cell distance. Shaded area in h is proportional to indicated probability. Significance in c-g was determined by 2-tailed paired Students t-test. Significance in i was assessed by paired t-tests with Bonferroni correction. *$P \leq$ 0.05, **$P \leq 0.01$, ***$P \leq 0.001$.

Figure 6

Figure 6: Experimental islet functional network is influenced by changes in the structural network

a: Representative images of Cx36$^{+/+}$ islet (left), heterozygous Cx36$^{+/-}$ islet (middle), and homozygous Cx36$^{-/-}$ islet (right), overlaid with a synchronization network. Dot signifies a cell, blue lines represent functional network edges that connect synchronized cells. Red cells are β-cell hubs. **b:** Oscillations in [Ca^{2+}] of corresponding islet in a. grey lines represent time-course for each cell, colored line represents mean islet time-course. **c:** Frequency histograms of connections for the corresponding Islet in a. Connections are calculated using functional network analysis and normalized to the maximum degree in the Islet. As such 100% links represents the cell with the greatest total number of connections. **d:** Normalized degree of functional network between cell pairs, averaged over islets of each Cx36$^{+/+}$, Cx36$^{+/-}$, Cx36$^{-/-}$ mouse. **e:** Percent of cells in the Islet which have are not connected to any other cell (eg. All $R_{ij} < R_{th}$). **f:** Average clustering coefficient. Overlaid is the box and whisker plot of 1000 random networks per Islet created with the same number of nodes, edges, and degree as the islet. **g:** Average Shortest path length. Overlaid is the box and whisker plot of 1000 random networks per islet, as in f. **h:** Average Global Efficiency. Overlaid is the box and whisker plot of 1000 random networks per islet, as in f. **i:** Local efficiency. **j:** Average number of pixels between synchronized cells normalized to the average number of pixels between all cells in the Islet. $D < 1$ indicates that cells preferentially synchronize to nearby cells. Significance in d-k determined by ordinary one-way ANOVA with Tukey multiple comparison test. Reported P values are adjusted. *$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$.

1186

Figure 7: Simulated islet functional network is influenced by changes in the structural
network

a: Representative cells within a simulated islet with high gap junction conductance ($g_{coup} =$
$0.12nS$) (left) and within an islet with gap junction low conductance ($g_{coup} = 0.03\ nS$)(right).
Connections between a representative cell are shown with synchronized cell pairs in blue, gap
junction cell pairs in red, and both in black. All data points represent the average statistic for each
mouse. **b:** Venn diagrams of the probability of synchronization and probability of gap junction

with the probability of both synchronization and gap junction as the overlapped area. **c:** Probability
that two cells are synchronized given they share a gap junction connection. **d:** Probability that two
cells were synchronized given they had similar rate of glucose metabolism and were within 15%
of the islet from each other. **e:** Probability that two cells shared a gap junction connection given
they were synchronized. **f:** Probability of similar rate of glucose metabolism given two cells were
synchronized. **g:** Global efficiency for $0.12nS, 0.06nS$, and $0.03nS$, for each islet seed. Overlaid
is the box and whisker plot of 1000 random networks per simulated islet seed. **h:** Clustering
coefficient for $0.12nS, 0.06nS$, and $0.03nS$, for each islet seed. Overlaid is the box and whisker
plot of 1000 random networks per simulated islet seed, as in g. **i:** Small world-ness of a network
calculated using global efficiency. 1000 random networks per islet were created to determine
uncertainty of each data point (shown in black). **j:** Local efficiency for $0.12nS, 0.06nS$, and
$0.03nS$, for each islet seed. Shaded area in b is proportional to indicated probability. Statistical
significance in c-j assessed by an ordinary one-way ANOVA with Tukey multiple comparison test.
Reported P values are adjusted. $*P \leq 0.05$, $**P \leq 0.01$, $***P \leq 0.001$.
1208

1209

Figure 8: Graphical summary of Results

Graphical summary of results from data. To answer question 1: we show that sub populations of β-cells in the functional network are driven by metabolism more than gap junction coupling (Top). To answer question 2: we show that metabolic similarities drive edges in the functional network (right). To answer question 3: we show that removal of edges in the structural network decreased global communication efficiency and local connectivity (bottom).

1217   SUPPLEMENTARY MATERIALS FOR

1218

1219   **What do Functional Synchronization Networks Indicate About Underlying Structure and**

1220   **System Dynamics? A network theory study of the islet.**

1221

1222   Jennifer K. Briggs, Vira Kravets, JaeAnn M. Dwulet, David J. Albers, Richard K. P. Benninger*

1223

1224

Fig. S1. Network sensitivity to threshold

a: Degree distribution of five simulated islets with five different thresholds. **b:** k_{glyc} comparing hubs and non hubs across five different thresholds. **c:** g_{coup} comparing hubs and non hubs across five different thresholds. Significance is assessed using Tukeys Multiple Comparisons. **d:** Islet with representative cell connections calculated by a lower threshold ($R_{th} = 0.999$) **(right)** and higher threshold ($R_{th} = 0.99995$) **(left)**. Synchronized cell pairs are shown in blue, gap junction pairs are shown in red and both are shown in black. **e:** Probability of a gap junction connection given synchronization for the five thresholds. **f**: Venn diagrams for the probability of synchronization (blue) and gap junction connection (red) with the overlapping area representing cell pairs that were both synchronized and gap junction coupled.

1236

Fig. S2: Paths that maximized gap junction conductivity.
a: Cumulative distribution functions of total conductance between two cells from 1-12 cells apart.
Synchronized CDF is shown in blue, non-synchronized CDF is shown in green. **b**: Bar charts comparing
the average conductance for synchronized and non-synchronized cells. P values are not corrected, but

1241   significance (denoted by asterisks) is determined using a Bonferroni-corrected paired T-test. Since we

1242   there were 15 cell distances, we set the significance threshold $\alpha = 0.003$.

1243

1244

Fig. S3. Paths that maximized metabolic rate.
a: Cumulative distribution functions of total k_{glyc} between two cells from 1-12 cells apart. Synchronized CDF is shown in blue, non-synchronized CDF is shown in green. **b:** Bar charts comparing the average conductance for synchronized and non-synchronized cells. P values are not corrected, but significance

1249   (denoted by asterisks) is determined using a Bonferroni-corrected paired T-test. Since we there were 15

1250   cell distances, we set the significance threshold $\alpha = 0.003$.

1251

1252

Fig. S4: Alternative Network Metrics for Calcium Islets from Cx36-KO mice
a: Averaged degree distribution histograms for all Islets analyzed. Error bars indicate standard deviation. Purple (**top**) histogram shows the distribution of connections for wild type (Cx36$^{+/+}$) islets. Orange (**middle**) shows the distribution of connections for heterozygous Cx36 knockout (Cx36$^{+/-}$) islets. Blue (**right**) shows the distribution of connections for homozygous Cx36 knockout (Cx36$^{-/-}$) islets. **b**: Average Pearson cross-correlation coefficient, between all cells in the islet. **c:** Number of cells in each islet. **d**: Small world-ness of a network calculated using global efficiency. 1000 random networks per islet were created to determine uncertainty of each data point (shown in black). Significance, or lack thereof is determined by an ordinary one-way ANOVA and Tukey Multiple comparison test. *$P \leq 0.05$, **$P \leq 0.01$

1264

Fig. S5: Network Metrics assessed by Islets

This figure shows similar information as that of Figure 7, but the data points are individual islets and the bars are those Islets grouped by mouse. Genotypes from top to bottom: homozygous Cx36 knockout (Cx36$^{-/-}$) islets, heterozygous Cx36 knockout (Cx36$^{+/-}$) islets, wild type (Cx36$^{+/+}$) islets. Random network distributions are overlayed as box-and-whisker plots.

1270

Fig. S6: Alternative Network Metrics for Simulated Islet with Gap Junction Coupling Change
a: Averaged degree distribution histograms for all Islets analyzed. Error bars indicate standard deviation. Purple (**top**) histogram shows the distribution of connections for fully coupled (0.12 nS) islets. Orange (**middle**) shows the distribution of connections for half coupled (0.06 nS) islets. Blue (**right**) shows the distribution of connections for homozygous quarter coupled (0.03 nS) islets. **b**: Average Pearson cross-correlation coefficient, between all cells in the islet. **c:** Small world-ness of a network calculated using global efficiency. 1000 random networks per islet were created to determine uncertainty of each data point (shown in black). Significance, or lack thereof is determined by an ordinary one-way ANOVA and Tukey Multiple comparison test . *$P \leq 0.05$, **$P \leq 0.01$

1280