Running Head: Can livestock be functional substitutes of native herbivores?

Functional substitutability of native herbivores by livestock for long-term soil carbon

sequestration depends on microbial decomposers.

Shamik Roy and Sumanta Bagchi[*]

Centre for Ecological Sciences, Indian Institute of Science, Bangalore – 560012, India

Corresponding author: Sumanta Bagchi, Centre for Ecological Sciences, Indian Institute of

Science, Bangalore – 560012, India; Phone number: 080-2293-3528; Email:

sbagchi@iisc.ac.in

ORCID: SR, 0000-0002-9916-5651; SB, 0000-0002-4841-6748.

Author contributions

S.R. and S.B. designed research; S.R. performed research; S.R. and S.B. analysed data; and

S.R. and S.B. wrote the paper.

Conflict of Interest: The authors declare that they have no conflict of interest.

Preprint server: bioRxiv, DOI: 10.1101/2022.02.07.479355

Classification: Biology, Ecology

Keywords: Carbon use efficiency, Land-use, Grassland, Soil microbial enzymes, Microbiome

rewilding, Structural equation modelling

This PDF file includes:

Main text

Figure 1 to 5

Abstract

The influence of native mammalian herbivores on ecosystems is declining as livestock have

displaced them to become the dominant grazers across the world. Alongside rewilding,

restoration, and conservation, it is now important to address whether livestock can assume the

functional roles of displaced native herbivores in grasslands, savannas, and steppes. If yes,

this has implications for the future of biodiversity, ecosystem functions and services, food-

security, and other social and environmental dimensions. We test competing alternative

hypotheses with a replicated, long-term, landscape-level experiment to ask whether livestock

in the Trans-Himalayan ecosystem can match decadal-scale (2006-17) soil-C sequestration

under native herbivores. Despite comparable settings, livestock were imperfect substitutes

and attained three-fourths of soil-C stocks under native herbivores. We evaluate multiple

lines of evidence surrounding this difference with a grazing-exclusion experiment (nutrient-

cycling, microbial metabolism, and e-DNA), and asses their inter-relationships. Livestock

and native herbivores differed in their effects on several soil microbial variables. Structural

equation models showed that livestock alter soil microbial communities towards lower fungal

abundance, which is detrimental for microbial carbon use efficiency (CUE), and ultimately

also for soil-C. Herbivore-identity influenced ecosystem functions even when total amount of

herbivory was comparable. Lags in soil-C storage can be addressed through microbial

restoration and rewilding of soil. Therefore, the idea of functional substitutability must be

leveraged alongside microbial rewilding to restore and conserve ecosystem functions.

Together they can reconcile conflicting demands from food-security and ecosystem services

in ways that have eluded conventional approaches to become an integral part of nature-based

climate solutions.

Significance Statement

Grazing by large mammalian herbivores affords an important natural climate solution via soil-C storage. But native grazers are progressively being replaced by domestic livestock to accommodate changing human diets. Even though livestock are now the dominant grazers around the world, it remains unclear whether they can be functional substitutes for native herbivores in sequestering soil-C. And if so, then to what extent? We find that livestock could only attain three-fourths of soil-C stocks under native herbivores, because they shift the microbial decomposer community in soil towards lower fungal abundance. This reduces microbial carbon use efficiency, and ultimately is detrimental for soil-C storage. So, microbial rewilding may help livestock production systems to emulate the ecosystem services provided by native herbivores.

Main text

Introduction

Large mammalian herbivores have undergone widespread declines in their abundance and distribution across the world's grasslands, steppes, and savannas. Pre-historic humans might have hunted many megafauna to extinction. More recently, humans have progressively replaced them with livestock across the world. Relatively intact assemblages of native herbivores are now increasingly confined to parks and reserves; elsewhere, livestock have not only become the most abundant herbivores, but they also represent the most geographically expansive land-use today (Asner et al. 2004, Bar-On et al. 2018). Further displacement of native herbivores seems inevitable due to steadily rising demands for livestock-products to accommodate evolving human-diets (Thornton 2010, Leclère et al. 2020). This severely restricts and alters the erstwhile influence of native herbivores on ecosystem structure and functions, their broad influence on global biogeochemical cycles, and their influence on soil-C in particular (Brodie et al. 2021, Roy and Bagchi 2021).

Increasingly, it is important to ask whether livestock can maintain the functional roles of the displaced native herbivores (Malhi et al. 2016, Cromsigt et al. 2018, Lundgren et al. 2020). If livestock can maintain ecological functions and services of their native counterparts, then such functional substitutability can have implications for the future of biodiversity, ecosystem functions and services, food-security, and other social and environmental dimensions (Sayer et al. 2013, Veblen et al. 2016, Willemen et al. 2020, Western et al. 2020, Delabre et al. 2021). The extent to which livestock can maintain ecosystem functions and services, particularly soil-C sequestration, becomes important to inform policies over sustainable use of grazing ecosystems and leverage their potential as natural climate solutions (Smith 2014, Cromsigt et al. 2018, Bossio et al. 2020).

Opportunities to restore and maintain ecosystem processes using livestock as

surrogates of displaced native herbivores arise from the mutual similarity in their functional

traits (Malhi et al. 2016, Lundgren et al. 2020, Schowanek et al. 2021). Indeed, traits are

better predictors of ecosystem functions than taxonomic identity of species (Hooper et al.

2005). Analyses of key traits, such as body size, diet-choice, and fermentation type,

hypothesize that livestock can be approximate surrogates for native herbivores, i.e., nearest

neighbors in trait-space (Malhi et al. 2016, Lundgren et al. 2020, Schowanek et al. 2021).

However, empirical evidence on ecosystem functions and services is needed before these

ideas can be implemented toward ecosystem management, and such evidence is lacking

(Reinhart et al. 2021).

From this functional-substitutability hypothesis one can ask (Betts et al. 2021) – if

there is sufficient overlap in trait-space between livestock and native herbivores, then to what

extent can they act as surrogates? It is challenging to address this question because many

modern livestock-production systems no longer have native counterparts within comparable

ecological settings that can allow a fair comparison. Unlike intensive livestock production

systems that are often associated with various forms of environmental degradation, traditional

pastoral production systems where both livestock and native herbivores occur in multiple-use

landscapes use (Asner et al. 2004, du Toit et al. 2009, Veblen et al. 2016, Western et al.

2020) can be testbeds to assess whether livestock are good substitutes, or not. While this

functional premise does not expect livestock to be perfect replacements for the displaced

native herbivores (Bagchi and Ritchie 2010, Sitters et al. 2020), it is important to estimate the

degree to which their ecological roles are matched to identify gaps and shortfalls as well as

the scope for improved livestock production systems (Smith 2014, Western et al. 2020,

Reinhart et al. 2021, Delabre et al. 2021).

Here we address impacts of livestock and native herbivores on soil-C sequestration as

grazing by large mammals is an important component of the global carbon cycle (Schmitz et

al. 2014, Roy and Bagchi 2021). We compare their ecological impacts in Spiti region of

northern India, which is part of the greater Trans-Himalayan ecosystem spread across the

Tibetan plateau and surrounding mountains. Soil-C sequestration in the Trans-Himalaya is

particularly important under human land-use for livestock production and under ongoing

climate change (Murthy and Bagchi 2018, Naidu and Bagchi 2021). We investigate long-

term (decadal-scale) net soil-C sequestration under livestock and native herbivores, and their

influence on key microbial decomposer functions related to the C-cycle (Falkowski et al.

2008, Sinsabaugh et al. 2016, Roy and Bagchi 2021). We take advantage of several key

features in the Trans-Himalaya: (a) the livestock are a multi-species assemblage that show

considerable trait-overlap with the native herbivores (Fig. 1), (b) livestock and native

herbivores occupy distinct but juxtaposed watersheds with comparable environmental

settings (Fig. S1-S4), (c) their abundances are comparable so that grazing pressure on

vegetation is also similar (see Methods, Fig. S7). Absent major confounding influences, the

livestock and native assemblages in Trans-Himalaya can help evaluate their functional

substitutability (Malhi et al. 2016, Cromsigt et al. 2018, Lundgren et al. 2020, Schowanek et

al. 2021). Specifically, we investigate (a) whether, and by how much, does grazing by

livestock and native herbivores differ in its influence on long-term (decadal-scale) soil-C

sequestration, and (b) whether any differences can be attributed to how each assemblage

influences microbial decomposer functions in soil.

Results

LONG-TERM TREND IN SOIL-CARBON

Over a decade (2006-17), soils under native herbivores stored 8.12±1.3 SE more carbon (mg

C cm^{-3}) than under livestock ($F_{1,27}$=9.86, P<0.01). Model comparison showed that spatio-

temporal heterogeneity was not sufficient to explain long-term variation in soil-C, and it

could instead be attributed to herbivore assemblage-type. There was stronger support for

M_A^{soilC} than for M_0^{soilC} (difference in AIC=−34.3, and difference in RMSE=−0.66, and

LR=36.2, P<0.001). Effectively, areas under livestock could realize up to 70% of the soil-C

found under native herbivores (Fig. 2).

GRAZER-EFFECTS ON SOIL MICROBES

Analysis of grazer-effects indicated that the observed difference in soil-C (Fig. 2) was related

to how soil microbes respond to grazing by livestock or native herbivores (Fig. 3). Livestock

and native herbivores had comparable effects on B, 16s rDNA abundance, MBC, MBN, LC,

LN, and qCO$_2$ (Fig. 3, Table S1). Grazing-intensity and grazer-effect on soil-C were also

comparable between livestock and native herbivores. For these nine variables, the competing

alternative models were not supported, and their variability could be attributed to spatio-

temporal heterogeneity rather than to herbivore assemblage-type (Fig. 3, Table S1). But

livestock and native herbivores differed in their influence on BR, SIR, F, ITS abundance, and

CUE. For these five variables, there was stronger support for the competing models than the

null models (Table S1), as assemblage-type was an important explanatory variable over and

above background spatio-temporal heterogeneity (Fig. 3). Grazing by livestock increased

both basal and potential microbial respiration (BR and SIR), compared to grazing by native

herbivores (Fig. 3). But grazing by livestock reduced both the fungal fraction of microbial

respiration and the abundance of fungal genetic markers (F and ITS). Respiration-based

estimates of fungal and bacterial fractions were consistent with DNA-based estimates (Fig.

3). Importantly, grazing by livestock reduced CUE, whereas grazing by native herbivores

increased it (Fig. 3).

INTER-RELATIONSHIPS BETWEEN SOIL MICROBIAL RESPONSES

To assess how differences in grazer-effects on one microbial variable may influence other

variables, we evaluated two competing SEM models. The null model was supported by the

data (SEM$_0$: Fisher's C=3.34, P=0.503, df=4, AIC=69.34). This showed, regardless of

herbivore assemblage-type, grazing influences microbial community composition (fungal

fraction) to alter CUE, and grazers also influence the labile-C pool via their effects on

microbial biomass and labile-N (Fig. 4). The alternative competing model was also supported

by the data (SEM$_A$: Fisher's C=2.108, P=0.716, df=4, AIC=88.11). It showed that change in

assemblage-type can influence CUE via its effect on microbial community composition (Fig.

4). Specifically, change in herbivore-assemblage from native to livestock has negative

influence on fungal abundance, which in turn is detrimental for CUE (Fig. 4). Both these

SEMs indicate that direction and magnitude of grazer-effects are consistent with higher CUE

under native herbivores improves net soil-C storage, compared to livestock (Fig. 2, Fig. 4).

LINKS BETWEEN MICROBIAL RESPONSES AND SOIL-C

We compared the role of herbivore assemblage-type ('identity') and grazing-intensity

('intensity') together, as a thought experiment to connect the inter-relationship between

different microbial responses with soil-C (Fig. 5). This model showed that changing

herbivore assemblage-type from native to livestock is detrimental to soil-C via the indirect

effects on microbial community composition (fungal fraction) and microbial CUE (Fig. 4-5).

In this way, it supported our expectation that effects arising from microbial processes at

monthly scale (seen in Fig. 4) can indeed accumulate over years and decades to influence

soil-C (Fisher's C=12.82, P=0.118, df=8, AIC=124.82). The links between microbial

community composition, CUE, and soil-C were not influenced by any background variation

in grazing-intensity (Fig. 3). Since grazing-intensity was comparable between livestock and

native herbivores, there were no paths connecting it to soil-C. In other words, herbivore

identity was important even if total amount of herbivory is comparable (Fig. 5).

Discussion

Previous studies from around the world have emphasized that grazing by large mammalian

herbivores leads to net soil-C storage (Conant and Paustian 2002, Witt et al. 2011, Abdalla et

al. 2018, Derner et al. 2019, Liu et al. 2020), and grazing ecosystems should be integral to

nature-based climate solutions (Smith 2014, Cromsigt et al. 2018, Willemen et al. 2020,

Bossio et al. 2020, Roy and Bagchi 2021). In parallel, studies have also identified the

limitations of nature-based climate solutions derived from soil, because soil-C tends to accrue

at very slow rates but can be lost quite rapidly (Smith 2014, Reinhart et al. 2021). Therefore,

preserving existing soil-C stocks is seen as an important and necessary step toward mitigating

land-atmosphere feedbacks alongside more ambitious targets of gaining soil-C (Smith 2014,

Naidu and Bagchi 2021). And we found that while soil-C has fluctuated, there has been no

net loss over decadal timescales (Fig. 2). But it has remained unclear whether this service can

be maintained when natural ecosystems are repurposed for livestock production (Reinhart et

al. 2021). And if so, then to what extent?

Our results address several inter-related aspects of the functional ecology of native

herbivores and livestock. We find that a multi-species herds of livestock can have remarkable

functional similarities with the native assemblages they displace (Fig. 1). But livestock do not

emerge as perfect substitutes since they achieve about three-fourths of the soil-C stock,

compared to the native herbivores (Fig. 2). Nevertheless, consistent with the functional

substitutability hypothesis, a key ecosystem service provided by native herbivores can also be

partly realized in livestock-production systems. This can have implications for how we

manage environmental impacts of livestock (Sayer et al. 2013, Smith 2014, Veblen et al.

2016, Willemen et al. 2020, Western et al. 2020, Delabre et al. 2021), and meet the rising

demand for livestock-products in the coming decades (Thornton 2010, Leclère et al. 2020).

To determine the scope of improved management, it is important to identify the

potential drivers and mechanisms that explain why livestock lag native herbivores in

maintaining soil-C. Prevailing management practices have largely focused on two aspects—

overgrazing and vegetation change—which result in altered C-input to soil from plants

(Bagchi and Ritchie 2010, Reinhart et al. 2021). However, this prevailing focus on C-input

alone can be simplistic and inadequate since it does not account for subsequent metabolism

of soil organic matter by microbes (Fontaine et al. 2004, Six et al. 2006, Falkowski et al.

2008, Sinsabaugh et al. 2013, Roy and Bagchi 2021). Our results from the grazer-exclusion

experiment identify soil microbial responses as the proximate explanation for why livestock

are not perfect substitutes for native herbivores (Fig. 3). Tests of the competing hypotheses

address whether the observed patterns in soil-C reflects idiosyncratic and/or pre-existing

differences between watersheds, or whether they are linked to the identity of the herbivores.

We attribute this missing soil-C to how microbial processes respond to grazing when native

herbivores are replaced by livestock (Fig. 4-5). However, this still leaves the possibility that

livestock have been historically overgrazed their watersheds, and their soils have not yet

recovered. Even though vegetation may recover from historical overgrazing, it may take

much longer for soil microbial functions to recover (López Zieher et al. 2020). Seen together,

these highlight the importance of soil microbial restoration in tandem with management of

grazing intensity and stocking rates. CUE is a fundamental microbial trait that controls

residence time of carbon in the soil, and it determines the flow of carbon through ecosystems

(Six et al. 2006, Falkowski et al. 2008, Sinsabaugh et al. 2013). The observed grazer-effects

on fungi (Fig. 3) are also known from other ecosystems (Bardgett et al. 1997, Zhang and Fu

2021). Direction and magnitude of difference in the microbial community under livestock

reduces CUE (Fig. 3-4), and this is consistent with lower soil-C (Fig. 5) because fungi are

known to have higher CUE than bacteria (Fontaine et al. 2004, Six et al. 2006).

Several overlapping factors can alter microbial communities towards lower fungi, and

their individual roles may not be easily disentangled from each other. For instance, variation

in vegetation composition can result in differences in the quality of litter-input to soil, and

this in turn can influence the rate and efficiency with which microbes metabolize soil organic

matter (Fontaine et al. 2004, Six et al. 2006, Sinsabaugh et al. 2013). Next, as discussed

above, microbial communities may have not yet recovered from any historical overgrazing in

the past (López Zieher et al. 2020), even though vegetation biomass does not indicate

overgrazing is currently a concern (Fig. 5). Additionally, veterinary care offered to the

livestock leads to inadvertent release of antibiotics and other drugs into soil from their dung

and urine, and this can alter microbial communities towards lower CUE (Wepking et al.

2019). Such overlapping factors need to be addressed together to improve soil-C

sequestration, rather than in isolation.

Our results from alternative competing SEMs offer insights into how net soil-C

sequestration can be improved (Fig. 4-5). Since replacing native herbivores with livestock is

detrimental for CUE, promoting wildlife conservation over livestock-production would favor

soil-C (Fig. 4-5). As that would compromise livelihoods and food-security, and it circles back

to the inconsistencies and conflicts between the goals of conservation and human land-use.

This conventional and exclusionary approach may no longer be tenable and is also

increasingly difficult to justify (Adams and Mulligan 2002); it is timely to assess alternative

ideas (Smith 2014). The null model (SEM_0, Fig. 4) can offer insights into ways to resolve

these conflicting demands on land-use. It suggests altered microbial communities

compromise CUE regardless of whether the community shift is a fallout of land-use, or not.

This opens opportunities for restoration of microbial communities to sustain ecosystem

services under human land-use (Paluch et al. 2013, Wubs et al. 2016, Mills et al. 2017).

Admittedly, there are many gaps in our current understanding of structure-function

relationships for hyper-diverse microbial communities (Bier et al. 2015). Yet, there are

encouraging signs that effective restoration can be achieved even with incomplete taxonomic

knowledge (Paluch et al. 2013, Wubs et al. 2016, Mills et al. 2017). Since livestock already

deliver 70% of the ecosystem service provided by native herbivores, further incorporating

microbial rewilding of soils into livestock-production may offset the 30% lag against native

herbivores—in the process, making livestock-production more environmentally responsible

(Smith 2014, Cromsigt et al. 2018, Willemen et al. 2020, Bossio et al. 2020).

In conclusion, our results highlight: (a) multi-species and functionally diverse

assemblages of livestock can partly approximate the ecological roles of native herbivores,

and (b) restoration and rewilding of soil microbial communities may enable livestock to

emulate the services provided by native herbivores. Alongside continued management of

stocking rates, functional substitutability and microbial restoration in-tandem may help

reconcile rising demands for livestock-products with conservation of ecosystem functions

and services.

Materials and methods

STUDY AREA

Traditional pastoral livestock production is the major land-use in Spiti region of the Trans-Himalayan grazing ecosystem. Watersheds near human settlements are used by a multi-species livestock assemblage, and these are juxtaposed with watersheds that retain native herbivores (Fig. 1, Fig. S1). Natural terrain and barriers (canyons, rivers, high ridges) that restrict frequent animal movement, maintain replicate watersheds under two alternative types of herbivore assemblages (Fig. S1, S2). Historical local extinctions of native herbivores and their replacement by livestock over the past decades remains a major conservation challenge in this region (Mishra et al. 2002, Namgail et al. 2013), and around the world (Veblen et al. 2016, Western et al. 2020). The extant native herbivores are ibex (*Capra sibirica*), bharal (*Pseudois nayaur*), and yak (*Bos grunniens grunniens*) that are related to wild yak (Bos grunniens mutus; also known as B. grunniens and B. mutus, respectively, 48). Currently, kiang (*Equus kiang*) are known from the fringes of the study area, and there are sporadic reports of argali (*Ovis ammon*). But Tibetan antelope (*Pantholops hodgsonii*) and Tibetan gazelle (*Procapra picticaudata*) are no longer found in the region. The livestock consist of goat, sheep, donkey, horse, cattle and yak-cattle hybrids (Fig. 1). The livestock and native herbivores (after including kiang and argali) show sufficiently high overlap in key traits (Lundgren et al. 2020), such as body mass, dietary guild and graminoid consumption, fermentation type, sexual-dimorphism, and limb morphology (Fig. 1). In recent decades, total animal biomass has remained comparable (6.9×10^4 kg for native and 7.2×10^4 kg for livestock; 23, 49), such that grazing pressure on vegetation has also remained similar for sufficiently long periods (Fig. S7).

The two types of watersheds are comparable in their abiotic conditions (Fig. S3–S4). Climate in Spiti is cold and semi-arid with temperature ranging from 25 °C in summer to

below -30 °C in winters (Fig. S5). Precipitation occurs as winter-snow (100-200 cm yr^{-1}) and

summer-rain (150-300 mm yr^{-1}), and vegetation growth-season is short (May-August, Fig.

S5). Vegetation in this treeless landscape consists of grasses (*Poa*, *Elymus*, *Festuca*), sedges

(*Carex*, *Kobresia*), forbs and shrubs (*Artemisia*, *Lindelofia*, *Caragana*). Both types of

watersheds have an overlapping pool of plant species, but show spatial variation in vegetation

composition. Livestock watersheds have high abundance of sedges and grasses, have higher

ground cover as well as higher aboveground plant production than grass-and-forb vegetation

in the native watersheds (23).

SOIL CARBON

Between 2006 and 2017, we measured soil-C at regular intervals in four watersheds with

native herbivores, and another four watersheds with livestock (Fig. S1, S2). In each

watershed, we set up 3-4 replicated permanent plots (10×10 m^2), for a total of n=15 plots

under native herbivores, and another n=15 plots under livestock. We collected soils with a 5

cm diameter and 20 cm deep cores, and measured carbon content (TruSpec, Leco, USA, and

Elementar Analysensysteme, GmbH, Germany) to compare long-term soil-C stocks (mg C

cm^{-3}) under livestock and native herbivores.

GRAZING-EXCLUSION EXPERIMENT

We distinguish whether any observed difference in soil-C between the watersheds is merely

an artefact of pre-existing site-specific conditions, or it can be attributed to grazing by the

alternative herbivore assemblages. For this, starting in 2005, we set up experimental grazer-

exclusion with a paired-adjacent fence at each location (also 10×10 m^2, each). In 2017, after

up to 12 years of herbivore-exclusion, we estimated grazer-effects on key soil microbial

functions in the paired-adjacent fenced and control plots.

GRAZING INTENSITY

336 We measured standing above-belowground biomass at the end of the growing season in

337 paired fenced and grazed plots as an indicator of grazing-intensity. We collected live shoot

338 biomass from 0.5×0.5 m^2 quadrats in each paired plot. Biomass samples were sun-dried, and

339 subsequently oven-dried to constant weight at 40 °C to obtain their dry-weight (g m^{-2}).

SOIL MICROBIAL FUNCTIONS

We estimated the response of key soil microbial functions to grazing, that are related to the

carbon cycle and net soil-C sequestration: Basal respiration (BR), Potential, or substrate-

induced respiration (SIR), Soil microbial community composition, broadly as fungal and

bacterial fractions, Microbial biomass C and N (MBC and MBN), labile Carbon (LC) and

labile Nitrogen (LN) fractions of soil organic matter, Metabolic quotient (qCO$_2$), and

Microbial carbon use efficiency (CUE). Since these microbial processes have seasonal

dynamics (Bagchi et al. 2017), we measured them at five time-points during the growing

season, between May and September. We followed standard laboratory protocols to measure

these microbial variables (Robertson et al. 1999, Bagchi et al. 2017, Roy and Bagchi 2021).

Briefly, for BR we used an alkali-trap where we incubated 4 g soil at 60% water holding

capacity and measured respired CO$_2$ (mg C g^{-1} soil day^{-1}). Similarly, we measured potential

microbial respiration (mg C g^{-1} soil day^{-1}) using substrate-induced respiration (SIR) after

adding 0.5% (w/v) of glucose to soil. We estimated broad microbial community composition

as fungal and bacterial contributions to potential respiration under selective inhibition using

Streptomycin (anti-bacterial) and Cycloheximide (anti-fungal). We used chloroform

fumigation-extraction to estimate MBC (mg C g^{-1} soil), and MBN (mg N g^{-1} soil). We

measured SOM from weight loss on ignition at 450 °C after 4 h. We measured labile

fractions (mg g^{-1} soil) of SOM using two-step hydrolysis with H$_2$SO$_4$. We estimated

metabolic quotient as the ratio of BR and MBC (qCO$_2$, respiration per unit biomass, hr^{-1}). We

estimated CUE from established relationships between key microbial extracellular enzymes

(carbon-acquiring Beta-glucosidase, alongside nitrogen-acquiring Leucine aminopeptidase

and *N*-acetyl glucosaminidase), MBC and MBN, and labile fractions of C and N in SOM

(Sinsabaugh et al. 2013, 2016). We measured activity of the three enzymes using their

specific fluorogenic substrates (4-MUB beta-D-glucopyranoside for Beta-glucosidase, L-

Leucine-7-amido-4-methylcoumarin hydrochloride for Leucine aminopeptidase, and 4-MUB

N-acetyl-beta-D-glucosaminide for N-acetylglucosamindase) (German et al. 2011) in soils

from growth season of 2017. These parameters influence microbial metabolism to determine

their investment in growth (G) and respiration (R), so that $CUE = \frac{G}{G+R}$ (Sinsabaugh et al.

2013, 2016). Theoretically CUE can range between 0 and 1, i.e., nil to perfect efficiency,

with higher values indicating greater potential for C-sequestration. But, CUE in nature

generally does not exceed 0.6 (Sinsabaugh et al. 2013, 2016).

SOIL MICROBIAL DNA

Based on patterns seen in fungal and bacterial contribution to soil respiration, we evaluated

community composition in more detail using their DNA markers in 2019 (Fierer et al. 2005,

Maestre et al. 2015). We collected soils from ten paired-adjacent plots at four time points

during the growing season and extracted microbial DNA using MOBIO Power Soil DNA

Extraction Kit following manufacturer's protocols. Next, we quantified the Internal

Transcribed Spacer (ITS-1 and ITS-2) regions of rDNA for fungal abundance, and region of

16s rDNA for bacterial abundance with qPCR (Fierer et al. 2005, Maestre et al. 2015). We

used these genetic data to confirm whether DNA-based estimates align with respiration-based

estimates of fungal and bacterial abundance. Additional details of laboratory analyses are in

Appendix.

DATA ANALYSIS

We evaluated competing hypotheses (Tredennick et al. 2021, Betts et al. 2021) to determine

whether the observed variation in the data merely represent site-specific and time-specific

heterogeneity between replicates, or it is necessary to invoke herbivore assemblage-type

(livestock or native) as an explanatory variable. We used a linear mixed-effects model to

evaluate long term trends in soil-C and grazer-exclusion effects on each microbial variable.

For soil-C, we used a null-model $\left(M_0^{soilC}\right)$ with time (years, 2006-17) and plot identity as

random-effects to account for background variation due to sampling locations and times. This

model was evaluated against the alternative competing model $\left(M_A^{soilC}\right)$ which included

assemblage-type as a fixed-effect, alongside both random-effects. We compared these models

based on parsimony, goodness-of-fit, and likelihood-ratio test (i.e., AIC, RMSE, and LR,

(Tredennick et al. 2021)).

We estimated grazer-effect on each microbial variable as the log-ratio of values in the

fenced and paired-adjacent control plots i.e., $\Delta = ln\left(\frac{\text{Grazing}_i}{\text{Fenced}_i}\right)$ for the i^{th} pair, and calculated

the mean and 95% CI of the log-ratio (Roy and Bagchi 2021). As earlier, we first evaluated a

null-model to assess whether the observed variation in grazer-effects can simply be attributed

to heterogeneity across sampling locations and times (months, May-September). As the null-

model for the j^{th} microbial variable $\left(M_0^j\right)$, we used plot identity and month as random-effects.

In the competing alternative model for each variable $\left(M_A^j\right)$, we included herbivore

assemblage-type as a fixed-effect alongside the random-effects. As before, we used AIC,

RMSE and LR for model comparison to judge whether herbivore assemblage-type is a

necessary explanatory variable for the data (Tredennick et al. 2021). We assessed whether the

data met statistical assumptions from observed and theoretical quantiles of residuals; the

variables required no further transformation (Δ is already in log-scale). We used 'nlme'

library in R (Pinheiro et al. 2020) to perform these analyses.

We assessed the inter-relationships between grazer-effects (Δ) on different microbial

variables with structural equations models (SEM). SEM help evaluate hypothesized pathways

over how change in one variable can influence another (Grace 2006). We identified SEM

paths from *a-priori* examples in the literature (Fig. 4-5). To avoid multicollinearity in SEM

arising from redundancy among variables, we removed those variables which were either

highly correlated with each other ($|r| > 0.49$), or were derived from each other (Fig. S8). So,

ΔSIR, ΔB, ΔMBN, and ΔqCO_2 were not included in this step, and we built SEM using

ΔMBC, ΔLN, ΔLC, ΔF, and ΔCUE. We did not use genetic data from ITS and 16s rDNA in

the SEM because these were collected from a subset of plots, were also not from the same

year as the other microbial variables, and because estimates from e-DNA broadly matched

respiration-based estimates of community composition. As earlier, we evaluated two

competing SEMs – with and without herbivore assemblage-type as an explanatory variable,

i.e., SEM_0 and SEM_A. We incorporated SEM paths as mixed-effects models for each variable

as piece-wise structural equations (Lefcheck 2016). We assessed goodness of fit with Fisher's

C statistic (Shipley 2009, Roy and Bagchi 2021) and report standardised path coefficients

alongside their statistical significance ($\alpha=0.05$). When SEM paths are supported by data, then

they offer a plausible explanation for an underlying process. When SEM paths are not

supported, the hypothesized process may not give rise to the observed data.

Since the temporal scale and span of the microbial variables (monthly dynamics) are

different from soil-C (decadal scale dynamics), SEM_0 and SEM_A by themselves cannot

directly evaluate how grazer-effects ultimately influence soil-C via microbes. While these

data have a common spatial dimension (n=30 replicate pairs), they differ in their temporal

dimensions. So, we evaluated their connections under the expectation that effects seen on

microbial processes (fast variables) can accumulate over years and eventually affect soil-C

(slow variables). For this, we first calculated the average (mean effect on j^{th} variable, $\overline{\Delta^J}$)

across all the months for each microbial variable, and across years for soil-C and grazing

intensity, for all replicates. Next, we evaluated SEM paths using these average values to

assess whether herbivore assemblage type and grazing intensity explain variation in soil-C

via grazer-effects on microbes. Here, plot identity was a random-effect in the SEM paths as

this allows the intercept to vary between samples that may have background difference in $\overline{\Delta^J}$

due to other factors such as grazing-intensity. Although grazing-intensity did not vary

systematically between the native herbivores and livestock (Fig. S7), we evaluated SEM with

both identity and intensity as explanatory variables to test for the relative influence of

herbivore identity and amount of herbivory.

Acknowledgements

CSIR-India provided graduate fellowship to SR; SB was supported by MHRD-India. DST-

SERB, DST-FIST, DBT-IISc, and MoEFCC provided funding support for fieldwork and lab-

analyses. Govt. of Himachal Pradesh has supported our long-term research in Spiti. We

received generous support from Abhilash Vijay Nair, Ananya Jana, Atish Roy Chowdhury,

Debarshi Dasgupta, Deepak Shetti, Dilip Naidu, Dipshikha Chakravortty, Dorje Chhering,

Dorje Chhewang, Ishika Pramanick, Kavita Isvaran, Praveen Karanth, Pronoy Baidya, Raju

Roy, and Ritika Chatterjee. Discussions with Ben Bond-Lamberty and Mayank Kohli helped

us prepare the manuscript.

References:

Abdalla, M., A. Hastings, D. R. Chadwick, D. L. Jones, C. D. Evans, M. B. Jones, R. M.
Rees, and P. Smith. 2018. Critical review of the impacts of grazing intensity on soil
organic carbon storage and other soil quality indicators in extensively managed
grasslands. Agriculture, Ecosystems and Environment 253:62–81.
Adams, W. M., and M. Mulligan. 2002. Decolonizing nature: Strategies for conservation in a
post-colonial era. Page (W. Adams and M. Mulligan, Eds.) Decolonizing Nature:
Strategies for Conservation in a Post-colonial Era. 1st edition. Routledge, London.
Asner, G. P., A. J. Elmore, L. P. Olander, R. E. Martin, and A. T. Harris. 2004. Grazing
systems, ecosystem responses, and global change. Annual Review of Environment and
Resources 29:261–299.
Bagchi, S., and M. E. Ritchie. 2010. Introduced grazers can restrict potential soil carbon
sequestration through impacts on plant community composition. Ecology Letters
13:959–968.
Bagchi, S., S. Roy, A. Maitra, and R. S. Sran. 2017. Herbivores suppress soil microbes to
influence carbon sequestration in the grazing ecosystem of the Trans-Himalaya.
Agriculture, Ecosystems and Environment 239:199–206.
Bardgett, R. D., D. K. Leemans, R. Cook, and P. J. Hobbs. 1997. Seasonality of the soil biota
of grazed and ungrazed hill grasslands. Soil Biology and Biochemistry 29:1285–1294.
Bar-On, Y. M., R. Phillips, and R. Milo. 2018. The biomass distribution on Earth.
Proceedings of the National Academy of Sciences 115:6506–6511.
Betts, M. G., A. S. Hadley, D. W. Frey, S. J. K. Frey, D. Gannon, S. H. Harris, H. Kim, U. G.
Kormann, K. Leimberger, K. Moriarty, J. M. Northrup, B. Phalan, J. S. Rousseau, T. D.
Stokely, J. J. Valente, C. Wolf, and D. Zárrate-Charry. 2021. When are hypotheses
useful in ecology and evolution? Ecology and Evolution 11:5762–5776.
Bier, R. L., E. S. Bernhardt, C. M. Boot, E. B. Graham, E. K. Hall, J. T. Lennon, D. R.
Nemergut, B. B. Osborne, C. Ruiz-González, J. P. Schimel, M. P. Waldrop, and M. D.
Wallenstein. 2015. Linking microbial community structure and microbial processes: an
empirical and conceptual overview. FEMS Microbiology Ecology 91:fiv113.
Bossio, D. A., S. C. Cook-Patton, P. W. Ellis, J. Fargione, J. Sanderman, P. Smith, S. Wood,
R. J. Zomer, M. von Unger, I. M. Emmer, and B. W. Griscom. 2020. The role of soil
carbon in natural climate solutions. Nature Sustainability 2020 3:5 3:391–398.
Brodie, J. F., S. Williams, and B. Garner. 2021. The decline of mammal functional and
evolutionary diversity worldwide. Proceedings of the National Academy of Sciences
118:e1921849118.
Conant, R. T., and K. Paustian. 2002. Potential soil carbon sequestration in overgrazed
grassland ecosystems. Global Biogeochemical Cycles 16:90-1-90–9.
Cromsigt, J. P. G. M., M. te Beest, G. I. H. Kerley, M. Landman, E. le Roux, and F. A.
Smith. 2018. Trophic rewilding as a climate change mitigation strategy? Philosophical
Transactions of the Royal Society B: Biological Sciences 373.
Delabre, I., L. O. Rodriguez, J. M. Smallwood, J. P. W. Scharlemann, J. Alcamo, A. S.
Antonarakis, P. Rowhani, R. J. Hazell, D. L. Aksnes, P. Balvanera, C. J. Lundquist, C.
Gresham, A. E. Alexander, and N. C. Stenseth. 2021. Actions on sustainable food
production and consumption for the post-2020 global biodiversity framework. Science
Advances 7:eabc8259.
Derner, J. D., D. J. Augustine, and D. A. Frank. 2019. Does Grazing Matter for Soil Organic
Carbon Sequestration in the Western North American Great Plains? Ecosystems
22:1088–1094.

Falkowski, P. G., T. Fenchel, and E. F. Delong. 2008. The Microbial Engines That Drive Earth's Biogeochemical Cycles. Science 320:1034–1039.

Fierer, N., J. A. Jackson, R. Vilgalys, and R. B. Jackson. 2005. Assessment of Soil Microbial Community Structure by Use of Taxon-Specific Quantitative PCR Assays. Applied and Environmental Microbiology 71:4117–4120.

Fontaine, S., G. Bardoux, L. Abbadie, and A. Mariotti. 2004. Carbon input to soil may decrease soil carbon content. Ecology Letters 7:314–320.

German, D. P., M. N. Weintraub, A. S. Grandy, C. L. Lauber, Z. L. Rinkes, and S. D. Allison. 2011. Optimization of hydrolytic and oxidative enzyme methods for ecosystem studies. Soil Biology and Biochemistry 43:1387–1397.

Grace, J. B. 2006. Structural Equation Modeling and Natural Systems. Cambridge University Press.

Hooper, D. U., F. S. Chapin, J. J. Ewel, A. Hector, P. Inchausti, S. Lavorel, J. H. Lawton, D. M. Lodge, M. Loreau, S. Naeem, B. Schmid, H. Setälä, A. J. Symstad, J. Vandermeer, and D. A. Wardle. 2005. Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. Ecological Monographs 75:3–35.

Leclère, D., M. Obersteiner, M. Barrett, S. H. M. Butchart, A. Chaudhary, A. De Palma, F. A. J. DeClerck, M. Di Marco, J. C. Doelman, M. Dürauer, R. Freeman, M. Harfoot, T. Hasegawa, S. Hellweg, J. P. Hilbers, S. L. L. Hill, F. Humpenöder, N. Jennings, T. Krisztin, G. M. Mace, H. Ohashi, A. Popp, A. Purvis, A. M. Schipper, A. Tabeau, H. Valin, H. van Meijl, W. J. van Zeist, P. Visconti, R. Alkemade, R. Almond, G. Bunting, N. D. Burgess, S. E. Cornell, F. Di Fulvio, S. Ferrier, S. Fritz, S. Fujimori, M. Grooten, T. Harwood, P. Havlík, M. Herrero, A. J. Hoskins, M. Jung, T. Kram, H. Lotze-Campen, T. Matsui, C. Meyer, D. Nel, T. Newbold, G. Schmidt-Traub, E. Stehfest, B. B. N. Strassburg, D. P. van Vuuren, C. Ware, J. E. M. Watson, W. Wu, and L. Young. 2020. Bending the curve of terrestrial biodiversity needs an integrated strategy. Nature 585:551–556.

Lefcheck, J. S. 2016. piecewiseSEM: Piecewise structural equation modelling in r for ecology, evolution, and systematics. Methods in Ecology and Evolution 7:573–579.

Leslie, D. M., and G. B. Schaller. 2009. Bos grunniens and Bos mutus (Artiodactyla: Bovidae). Mammalian Species 836:1–17.

Liu, X., H. Sheng, Z. Wang, Z. Ma, X. Huang, and L. Li. 2020. Does Grazing Exclusion Improve Soil Carbon and Nitrogen Stocks in Alpine Grasslands on the Qinghai-Tibetan Plateau? A Meta-Analysis. Sustainability 12:977.

López Zieher, X. M., L. Vivanco, and L. Yahdjian. 2020. Soil bacterial communities remain altered after 30 years of agriculture abandonment in Pampa grasslands. Oecologia 193:959–968.

Lundgren, E. J., D. Ramp, J. Rowan, O. Middleton, S. D. Schowanek, O. Sanisidro, S. P. Carroll, M. Davis, C. J. Sandom, J.-C. Svenning, and A. D. Wallach. 2020. Introduced herbivores restore Late Pleistocene ecological functions. Proceedings of the National Academy of Sciences 117:7871–7878.

Maestre, F. T., M. Delgado-Baquerizo, T. C. Jeffries, D. J. Eldridge, V. Ochoa, B. Gozalo, J. L. Quero, M. García-Gómez, A. Gallardo, W. Ulrich, M. A. Bowker, T. Arredondo, C. Barraza-Zepeda, D. Bran, A. Florentino, J. Gaitán, J. R. Gutiérrez, E. Huber-Sannwald, M. Jankju, R. L. Mau, M. Miriti, K. Naseri, A. Ospina, I. Stavi, D. Wang, N. N. Woods, X. Yuan, E. Zaady, and B. K. Singh. 2015. Increasing aridity reduces soil microbial diversity and abundance in global drylands. Proceedings of the National Academy of Sciences of the United States of America 112:15684–15689.

Malhi, Y., C. E. Doughty, M. Galetti, F. A. Smith, J.-C. Svenning, and J. W. Terborgh. 2016.
Megafauna and ecosystem function from the Pleistocene to the Anthropocene.
Proceedings of the National Academy of Sciences 113:838–846.

Mills, J. G., P. Weinstein, N. J. C. Gellie, L. S. Weyrich, A. J. Lowe, and M. F. Breed. 2017.
Urban habitat restoration provides a human health benefit through microbiome
rewilding: the Microbiome Rewilding Hypothesis. Restoration Ecology 25:866–872.

Mishra, C., S. E. Wieren, I. M. A. Heitkönig, and H. H. T. Prins. 2002. A theoretical analysis
of competitive exclusion in a Trans-Himalayan large-herbivore assemblage. Animal
Conservation 5:251–258.

Murthy, K., and S. Bagchi. 2018. Spatial patterns of long-term vegetation greening and
browning are consistent across multiple scales: Implications for monitoring land
degradation. Land Degradation & Development 29:2485–2495.

Naidu, D. G. T., and S. Bagchi. 2021. Greening of the earth does not compensate for rising
soil heterotrophic respiration under climate change. Global Change Biology 27:2029–
2038.

Namgail, T., S. E. van Wieren, and H. H. T. Prins. 2013. Distributional congruence of
mammalian herbivores in the Trans-Himalayan Mountains. Current Zoology 59:116–
124.

Paluch, E. C., M. A. Thomsen, and T. J. Volk. 2013. Effects of Resident Soil Fungi and Land
Use History Outweigh Those of Commercial Mycorrhizal Inocula: Testing a Restoration
Strategy in Unsterilized Soil. Restoration Ecology 21:380–389.

Pinheiro, J., D. Bates, S. DebRoy, D. Sarkar, and R. R Development Core Team. 2020. nlme:
Linear and Nonlinear Mixed Effects Models, R package version 3.1-148.

Reinhart, K. O., H. S. Sanni Worogo, and M. J. Rinella. 2021. Ruminating on the science of
carbon ranching. Journal of Applied Ecology.

Robertson, G. P., D. C. Coleman, C. S. Bledsoe, and P. Sollins. 1999. Standard soil methods
for long-term ecological research. Oxford University Press, New York, NY.

Roy, S., and S. Bagchi. 2021. Large Mammalian Herbivores and the Paradox of Soil Carbon
in Grazing Ecosystems: Role of Microbial Decomposers and Their Enzymes.
Ecosystems In press:1–13.

Sayer, J., T. Sunderland, J. Ghazoul, J.-L. Pfund, D. Sheil, E. Meijaard, M. Venter, A. K.
Boedhihartono, M. Day, C. Garcia, C. van Oosten, and L. E. Buck. 2013. Ten principles
for a landscape approach to reconciling agriculture, conservation, and other competing
land uses. Proceedings of the National Academy of Sciences 110:8349–8356.

Schmitz, O. J., P. A. Raymond, J. A. Estes, W. A. Kurz, G. W. Holtgrieve, M. E. Ritchie, D.
E. Schindler, A. C. Spivak, R. W. Wilson, M. A. Bradford, V. Christensen, L. Deegan,
V. Smetacek, M. J. Vanni, and C. C. Wilmers. 2014. Animating the Carbon Cycle.
Ecosystems 17:344–359.

Schowanek, S. D., M. Davis, E. J. Lundgren, O. Middleton, J. Rowan, R. Ø. Pedersen, D.
Ramp, C. J. Sandom, and J. Svenning. 2021. Reintroducing extirpated herbivores could
partially reverse the late Quaternary decline of large and grazing species. Global
Ecology and Biogeography 30:896–908.

Shipley, B. 2009. Confirmatory path analysis in a generalized multilevel context. Ecology
90:363–368.

Singh, R., R. K. Sharma, and S. Babu. 2015. Pastoralism in transition: Livestock abundance
and herd composition in Spiti, Trans-Himalaya. Human Ecology 43:799–810.

Sinsabaugh, R. L., S. Manzoni, D. L. Moorhead, and A. Richter. 2013. Carbon use efficiency
of microbial communities: stoichiometry, methodology and modelling. Ecology Letters
16:930–939.

Sinsabaugh, R. L., B. L. Turner, J. M. Talbot, B. G. Waring, J. S. Powers, C. R. Kuske, D. L.
Moorhead, and J. J. Follstad. Shah. 2016. Stoichiometry of microbial carbon use
efficiency in soils. Ecological Monographs 86:172–189.
Sitters, J., D. M. Kimuyu, T. P. Young, P. Claeys, and H. Olde Venterink. 2020. Negative
effects of cattle on soil carbon and nutrient pools reversed by megaherbivores. Nature
Sustainability 3:360–366.
Six, J., S. D. Frey, R. K. Thiet, and K. M. Batten. 2006. Bacterial and Fungal Contributions
to Carbon Sequestration in Agroecosystems. Soil Science Society of America Journal
70:555–569.
Smith, P. 2014. Do grasslands act as a perpetual sink for carbon? Global Change Biology
20:2708–2711.
Thornton, P. K. 2010. Livestock production: recent trends, future prospects. Philosophical
Transactions of the Royal Society B: Biological Sciences 365:2853–2867.
du Toit, J. T., R. Kock, and J. C. Deutsch. 2009. Wild Rangelands: Conserving Wildlife
While Maintaining Livestock in Semi-Arid Ecosystems. John Wiley and Sons Ltd.
Tredennick, A. T., G. Hooker, S. P. Ellner, and P. B. Adler. 2021. A practical guide to
selecting models for exploration, inference, and prediction in ecology. Ecology
102:e03336.
Veblen, K. E., L. M. Porensky, C. Riginos, and T. P. Young. 2016. Are cattle surrogate
wildlife? Savanna plant community composition explained by total herbivory more than
herbivore type. Ecological Applications 26:1610–1623.
Wepking, C., B. Badgley, J. E. Barrett, K. F. Knowlton, J. M. Lucas, K. J. Minick, P. P. Ray,
S. E. Shawver, and M. S. Strickland. 2019. Prolonged exposure to manure from
livestock-administered antibiotics decreases ecosystem carbon-use efficiency and alters
nitrogen cycling. Ecology Letters 22:2067–2076.
Western, D., P. Tyrrell, P. Brehony, S. Russell, G. Western, and J. Kamanga. 2020.
Conservation from the inside-out: Winning space and a place for wildlife in working
landscapes. People and Nature 2:279–291.
Willemen, L., N. N. Barger, B. ten Brink, M. Cantele, B. F. N. Erasmus, J. L. Fisher, T.
Gardner, T. G. Holland, F. Kohler, J. S. Kotiaho, G. P. von Maltitz, G. Nangendo, R.
Pandit, J. A. Parrotta, M. D. Potts, S. D. Prince, M. Sankaran, A. Brainich, L.
Montanarella, and R. Scholes. 2020. How to halt the global decline of lands. Nature
Sustainability 3:164–166.
Witt, G. B., M. V. Noël, M. I. Bird, R. J. S. Beeton, and N. W. Menzies. 2011. Carbon
sequestration and biodiversity restoration potential of semi-arid mulga lands of Australia
interpreted from long-term grazing exclosures. Agriculture, Ecosystems and
Environment 141:108–118.
Wubs, E. R. J., W. H. Van Der Putten, M. Bosch, and T. M. Bezemer. 2016. Soil inoculation
steers restoration of terrestrial Ecosystems. Nature Plants 2:1–5.
Zhang, H., and G. Fu. 2021. Responses of plant, soil bacterial and fungal communities to
grazing vary with pasture seasons and grassland types, northern Tibet. Land Degradation
& Development 32:1821–1832.
644
645

Figures

Fig. 1

Figure 1. Trait space of livestock and native herbivores in Spiti, India. We find there is

considerable overlap in multidimensional trait space between native herbivores (green) and

livestock (blue). Native herbivores in the study area are ibex, bharal, yak; asterix (kiang,

argali) indicates that these animals are found in nearby regions. The livestock consist of goat,

sheep, donkey, horse, cattle and yak-cattle hybrids. Six traits were used to construct the trait-

space: body mass, dietary guild, fermentation type, sexual-dimorphism, graminoid, and limb

morphology (Lundgren et al. 2020) with Principal Coordinates Analysis on the Gower's

distance matrix.

Fig. 2

Figure 2. Long-term change in soil-C (mean ± 95CI) under grazing by native herbivores and

livestock in Spiti, India. Areas under livestock attained 70% of soil-C compared to areas

under native herbivores. This difference (two-thirds) was consistent across more than a

decade. Data are from n=30 permanent plots in Spiti region of northern India.

Figure 3. Grazer-effect ($\Delta = ln\left(\frac{grazed}{fenced}\right)$), as mean $\pm$ 95CI) of livestock and native herbivores

on twelve soil microbial variables, alongside grazer-effects on aboveground biomass and

soil-C, in Spiti, India. These variables are basal microbial respiration (BR), potential

microbial respiration (SIR), Fungal fraction (F), ITS DNA-amplicon abundance (ITS),

Bacterial fraction (B), 16s DNA-amplicon abundance (16s), Microbial Biomass Carbon

(MBC), Microbial Biomass Nitrogen (MBN), Labile-C (LC), Labile-N (LN), Metabolic

quotient (qCO_2), and microbial carbon use efficiency (CUE) which were measured at

monthly intervals during the growth season. Grazing-intensity (ΔShoot: from difference in

peak aboveground biomass) and grazer-effect on soil-C (ΔsoilC) are from long-term decadal

measurements. Direction and magnitude of differences in BR, SIR, F, ITS abundance, and

CUE varied with herbivore-assemblage type. But the remaining variables were not influenced

by herbivore-assemblage type. Data are from n=30 permanent plots in Spiti region of

northern India (except e-DNA data, which are from a subset of 20 plots).

Fig. 4

Figure 4. Mixed-effects structural equation models to evaluate two competing hypotheses – with and without herbivore assemblage-type as an explanatory variable, i.e., SEM_A and SEM_0. For each variable, grazer-effect ($\Delta = ln\left(\frac{grazed}{fenced}\right)$) was calculated from grazed and ungrazed paired plots. The variables are ΔBR: basal respiration, ΔMBC: microbial biomass-C, ΔLC: labile-C, ΔLN: labile-N, ΔF: fungal biomass, ΔCUE: carbon use efficiency. In SEM_A, as herbivore-assemblage goes from native to livestock, a decrease in ΔF leads to a decrease in ΔCUE. Alternatively, in SEM_0, increase in ΔF increases ΔCUE, independent of herbivore-assemblage type. Green and red arrows represent significant positive and negative standardized path coefficients respectively, while grey arrows represent non-significant paths ($\alpha=0.05$). Marginal and conditional R^2 are reported, and asterix represent statistical significance (*** $P \leq 0.001$, ** $P \leq 0.01$, and * $P \leq 0.05$).

702 Fig. 5.

SEM: Fisher's C = 12.82, P=0.118, df = 8, AIC = 124.82

Fig. 5. Mixed-effects structural equation models to evaluate whether grazer-effects on

microbial responses explain variation in soil-C. Unlike Fig. 4, here $\bar{\Delta}$ is the mean of grazer-

effect averaged across time (months for microbial variables and years for soil-C and grazing-

intensity). The helps us evaluates the relative influence of herbivore-identity (assemblage

type), and total amount of herbivory (grazing-intensity, $\overline{\Delta Shoot}$) on variation in soil-C. This

model shows that replacing native herbivores with livestock alters soil microbial

communities to reduce their CUE, and this is ultimately also detrimental for soil-C storage.

Green and red arrows represent significant positive and negative standardized path

coefficients respectively, while grey arrows represent non-significant paths (α=0.05).

Marginal and conditional R^2 are reported, and asterix represent statistical significance (***

P≤0.001, ** P≤0.01, and * P≤0.05).