Running Head: Can livestock be substitutes for native herbivores?

Functional substitutability of native herbivores by livestock for soil carbon depends on

microbial decomposers

Shamik Roy [1], Dilip GT Naidu [1,2], Sumanta Bagchi [1,2,*]

[1]Centre for Ecological Sciences, Indian Institute of Science, Bangalore 560012, India

[2]Divecha Centre for Climate Change, Indian Institute of Science, Bangalore 560012, India

*Correspondence: Sumanta Bagchi, Centre for Ecological Sciences, Indian Institute of

Science, Bangalore – 560012, India; Phone number: 080-2293-3528; Email:

sbagchi@iisc.ac.in

ORCID: SR, 0000-0002-9916-5651; DGTN, 0000-0001-7966-4579; SB, 0000-0002-4841-

6748.

Author contributions

S.R. and S.B. designed research; S.R., DGTN, and SB performed research; S.R. and S.B.

analysed data; and S.R., DGTN, and S.B. wrote the paper.

Conflict of Interest: The authors declare that they have no conflict of interest.

Preprint server: bioRxiv, DOI: 10.1101/2022.02.07.479355

Abstract

Grazing by large mammalian herbivores impacts climate as it can favor the size and stability of a large carbon (C) pool in soil. As native herbivores in the world's grasslands, steppes, and savannas are progressively being displaced by livestock, it is important to ask whether livestock can emulate the functional roles of their native counterparts. While livestock and native herbivores can have remarkable similarity in their traits, they can differ greatly in their impacts on vegetation composition which can affect soil-C. It is uncertain how their similarities and differences impact soil-C via their influence on microbial decomposers. We test competing alternative hypotheses with a replicated, long-term, landscape-level, grazing-exclusion experiment to ask whether livestock in the Trans-Himalayan ecosystem of northern India can match decadal-scale (2005-2016) soil-C stocks under native herbivores. We evaluate multiple lines of evidence from 17 variables that influence soil-C (quantity and quality of C-input from plants, microbial biomass and metabolism, microbial community composition, veterinary antibiotics in soil), and asses their inter-relationships. Livestock and native herbivores differed in their effects on several soil microbial processes. Microbial carbon use efficiency (CUE) was 19% lower in soils under livestock. Compared to native herbivores, areas used by livestock contained 1.5 kg C m^{-2} less soil-C. Structural equation models showed that alongside effects arising from plants, livestock alter soil microbial communities which is detrimental for CUE, and ultimately also for soil-C. Supporting evidence pointed toward a link between veterinary antibiotics used on livestock, microbial communities, and soil-C. Overcoming the challenges of sequestrating antibiotics to minimize their potential impacts on climate, alongside microbial rewilding under livestock, may reconcile the conflicting demands from food-security and ecosystem services. Conservation of native herbivores and better management of livestock is crucial for soil-C stewardship to envision and achieve natural climate solutions.

Keywords: Grassland, Soil microbial enzymes, Microbiome rewilding, Restoration,

Ecosystem disservice, Tetracycline, Carbon use efficiency

Introduction

Grazing by large mammalian herbivores is integral to the zoogeochemistry of the global

carbon (C) cycle (Schmitz et al., 2014, 2018). Herbivores exert a strong influence on climate

via their impacts on a large soil-C pool—over 500 Pg C in the world's grasslands, steppes,

and savannas (Derner et al., 2019; Liu et al., 2020; Naidu et al., 2022; Roy & Bagchi, 2022;

Sitters et al., 2020; Witt et al., 2011). Long before the industrial era, humans started

influencing global climate via their impacts on the distribution and abundance of large

mammalian herbivores—prehistoric humans may have hunted many megafauna to extinction,

and modern humans are progressively replacing native herbivores with domestic livestock

across the world (Asner et al., 2004; Bar-On et al., 2018; Schowanek et al., 2021). Intact

assemblages of native herbivores are now increasingly confined to parks and reserves;

elsewhere, livestock have not only become the most abundant grazers, but they are also the

most expansive land-use. Today, livestock biomass (10^{11} kg) exceeds that of all other

mammals; they occur at high densities ($>10^3$ individuals km^{-2}) across more than 60 million

km^2 which contain over 295 Pg of soil-C (Bar-On et al., 2018; Brus et al., 2017; Gilbert et al.,

2018). Further displacement of native herbivores seems inevitable due to steadily rising

demands for livestock-products to accommodate evolving human-diets (Leclère et al., 2020;

Thornton, 2010). Increasingly, it is important to ask whether livestock can maintain the

functional roles of native herbivores they displace (Cromsigt et al., 2018; Lundgren et al.,

2020; Malhi et al., 2016) because grazing can favor soil-C to decarbonize the atmosphere and

offers an important natural climate solution (Naidu et al., 2022; Roy & Bagchi, 2022). If

livestock can maintain the ecological functions of their native counterparts, then such

substitutability can have implications for the future of biodiversity, ecosystem functions and

services, food-security, and other social and environmental dimensions (Delabre et al., 2021;

du Toit et al., 2012; Sayer et al., 2013; Veblen et al., 2016).

The extent to which livestock can approximate the functional influence of native

herbivores on soil-C is important to inform and steer grazing management as a natural

climate solution (Bossio et al., 2020; Cromsigt et al., 2018; Reinhart et al., 2022; P. Smith,

2014). On one hand, the opportunity for livestock to be functional surrogates of displaced

native herbivores arises from the mutual similarity in their traits (Lundgren et al., 2020;

Malhi et al., 2016; Schowanek et al., 2021). Analyses of key traits, such as body size, diet-

choice, and fermentation type, hypothesize that livestock can be approximate surrogates for

native herbivores, i.e., nearest neighbors in trait-space have similar ecological roles

(Lundgren et al., 2020; Malhi et al., 2016; Schowanek et al., 2021). On the other hand, in

many ecosystems, livestock differ from native herbivores in how they impact vegetation

composition (Bagchi et al., 2012; Bagchi & Ritchie, 2010b; Price et al., 2022; Ratajczak et

al., 2022; Wells et al., 2022). This alters the quantity and quality of C-input to soil from

plants by changing the underlying distribution of plant traits (e.g., root:shoot ratio, C:N

stoichiometry of leaf and litter, etc.) that is ultimately consequential for soil-C (Bagchi &

Ritchie, 2010b; Laliberté & Tylianakis, 2012). While there is clarity on how herbivore

assemblages influence vegetation communities (Augustine & McNaughton, 1998; Bagchi et

al., 2012; Bakker et al., 2006), we have a limited understanding of the downstream

consequences for soil-C that are ultimately controlled by microbial actions on soil organic

matter (Falkowski et al., 2008; Fontaine et al., 2004; Roy & Bagchi, 2022; Sinsabaugh et al.,

2013; Six et al., 2006). For instance, differences in aboveground responses may percolate

belowground to alter the labile and recalcitrant fractions of soil organic matter (Bardgett &

Wardle, 2003), and depending on the direction and magnitude, this can either promote or

deplete soil-C (Fontaine et al., 2004; Kuzyakov, 2010). Further, livestock also release

veterinary antibiotics into soil through their dung and urine which can alter soil microbial

communities (Lucas et al., 2021; Wepking et al., 2019). But we do not know whether soil

microbes respond differently to grazing by livestock and native herbivores, and how this

influences soil-C under the two types of grazers. In other words, though the promise of

functional substitutability between livestock and native herbivores might be constrained by

aboveground responses by plants (Bagchi et al., 2012; Bagchi & Ritchie, 2010b), it remains

tangled with belowground responses over how soil microbes respond to the two types of

grazers. This limits our ability to envision and derive natural climate solutions from livestock

(Bossio et al., 2020; Cromsigt et al., 2018; Reinhart et al., 2022; P. Smith, 2014) across a

large fraction of the earth's terrestrial surface (Asner et al., 2004; Bar-On et al., 2018).

Here we ask how livestock and native herbivores influence soil-C via their effects on

soil microbial processes in addition to their aboveground effects (Falkowski et al., 2008; Roy

& Bagchi, 2022; Sinsabaugh et al., 2016). We assess their functional substitutability in Spiti

region of northern India, which is part of the greater Trans-Himalayan ecosystem spread

across the Tibetan plateau and surrounding mountains (Fig. S1). We investigate long-term

(decadal-scale) soil-C stocks under livestock and native herbivores, and their influence on

key microbial decomposer functions related to C-cycling in soil. We take advantage of

several key features in the Trans-Himalaya: (a) the livestock are a multi-species assemblage

that show considerable trait-overlap with the native herbivores (Fig. 1), (b) livestock and

native herbivores occupy replicate juxtaposed watersheds with comparable environmental

settings (e.g., similarity in edaphic factors, climatic conditions, Fig. S1-S4), (c) their

abundances are comparable such that grazing pressure on vegetation is also similar (Fig. S5).

Absent major confounding influences, the livestock and native herbivore assemblages in

Trans-Himalaya can help evaluate unresolved questions over how they influence soil-C

(Cromsigt et al., 2018; Lundgren et al., 2020; Malhi et al., 2016; Schowanek et al., 2021).

Materials and methods

STUDY AREA

Climate in Spiti region of Trans-Himalaya is cold and semi-arid with temperature ranging

from 25 °C in summer to below –30 °C in winters (Fig. S1, S2). Precipitation occurs as

winter-snow (100-200 cm yr^{-1}) and summer-rain (150-300 mm yr^{-1}), and vegetation growth-

season is short (May-August, Fig. S2), peak biomass in July-August. Vegetation in this

treeless ecosystem consists of grasses (*Poa*, *Elymus*, *Festuca*), sedges (*Carex*, *Kobresia*),

forbs (*Lindelofia*, *Astragalus*) and shrubs (*Artemisia*, *Caragana*). Traditional pastoral

livestock production is the major land-use across much of the Trans-Himalaya. Watersheds

near village Kibber in Spiti are used by a multi-species livestock assemblage, and these are

juxtaposed with watersheds that retain native herbivores (Fig. 1, Fig. S1). Natural terrain and

barriers (canyons, escarpments, rivers, high ridges) restrict frequent animal movement and

maintain replicate watersheds under two alternative herbivore assemblages (Fig. S1)—some

used primarily by native herbivores and others by livestock (Bagchi & Ritchie, 2010b). Like

other parts of the world (Veblen et al., 2016; Western et al., 2020), local extinctions of native

herbivores and their replacement by livestock over the past decades remains a major

conservation challenge in the Trans-Himalaya (Mishra et al., 2002; Namgail et al., 2013).

The extant native herbivores are ibex (*Capra sibirica*), bharal (*Pseudois nayaur*), and yak

(*Bos grunniens grunniens*) that are related to wild yak (Bos grunniens mutus; also known as

B. grunniens and B. mutus, respectively; Leslie & Schaller, 2009). Currently, kiang (*Equus*

kiang) are known from the fringes of the study area (Fig. S1), and there are sporadic reports

of argali (*Ovis ammon*). But Tibetan antelope (*Pantholops hodgsonii*) and Tibetan gazelle

(*Procapra picticaudata*) are no longer found in the study area (Mishra et al., 2002; Namgail

et al., 2013). The livestock consist of yak-cattle hybrids, cattle goat, sheep, horse, and donkey

(Fig. 1).

NATIVE HERBIVORES AND LIVESTOCK OF SPITI

Cumulative biomass of native herbivores (c. 1.1×10^5 kg) has remained comparable to that of

livestock over the past few decades (c. 1.2×10^5 kg) in the watersheds covering c. 40 km^2 near

village Kibber (Bagchi & Ritchie, 2010b; Singh et al., 2015) (Fig. S1). The livestock and

native herbivores (after including kiang and argali across the broader region) show

sufficiently high overlap in key traits (Lundgren et al., 2020), such as their body mass, dietary

guild and graminoid consumption, fermentation type, sexual-dimorphism, and limb

morphology (Fig. 1). Since the native herbivores appear to have a livestock counterpart (Fig.

1), one expects the two assemblages to be functionally similar. Yet, consistent with other

parts of the world, the livestock and native herbivores differ in their impacts on vegetation

composition (Bagchi et al., 2012; Bagchi & Ritchie, 2010b; Price et al., 2022; Ratajczak et

al., 2022; Wells et al., 2022). Specifically, native herbivores lead to forb-and-grass dominated

vegetation whereas sedges dominate under livestock (Fig. 1). Such differences in vegetation

composition can be attributed to diet selectivity of the two herbivore types (Augustine &

McNaughton, 1998; Bagchi et al., 2012; Bagchi & Ritchie, 2010b; Ratajczak et al., 2022).

However, peak-season live biomass, both above and below-ground, are comparable (Fig. 1).

In addition, standing litter biomass, and microbial biomass in soil are also comparable (Fig.

1). The two types of watersheds are also similar in several key abiotic variables (Fig. S3-S4).

Soil pH is near-neutral to alkaline in both native and livestock watersheds; electrical

conductivity and bulk density are also comparable (Fig. S3). Soils in both types of

watersheds have sandy-loam texture (Fig. S4).

These similarities and differences can determine whether livestock can match long-

term soil-C stocks under native herbivores, or not. Specifically, this not only depends on how

the two types of herbivores influence C-input from plants to soil, and but also how they

influence microbial processes during decomposition of soil organic matter (Bardgett &

Wardle, 2003; Falkowski et al., 2008; Roy & Bagchi, 2022; Sinsabaugh et al., 2016).

EXPERIMENTAL DESIGN

Humans arrived in Spiti in the pre-historic age (Bellezza, 2017), and livestock replaced the

native herbivores from these watersheds few decades before our study began (Mishra et al.,

2002; Namgail et al., 2013). Thus, *sensu stricto*, a contemporary landscape-level comparison

of these watersheds cannot distinguish the effects of change in animal assemblage from any

pre-existing differences between them. Therefore, instead of relying solely on prevailing

differences in above- and belowground factors that influence soil-C, we determine whether

any observed difference in soil-C between the watersheds can be attributed to how plants and

microbes respond to grazing by the alternative herbivore assemblages. We use long-term

experimental herbivore-exclusion in the replicate watersheds to quantify grazer-effects on

different variables to investigate (a) whether grazing by livestock and native herbivores has

different influence on soil-C, and (b) whether any difference in soil-C can be attributed to

how plants and microbes respond to grazing by the two assemblages. This approach overlays

a manipulative experiment over the natural experiment of two types of watersheds (Bagchi &

Ritchie, 2010b). For this, starting in 2005, we set up experimental grazer-exclusion with

replicated paired-and-adjacent control-and-fenced plots (10×10 m^2, each). We used four

watersheds used primarily by native herbivores, and another four by livestock (Fig. S1). We

set up 3-4 paired control-and-fenced plots in each watershed (n=15 plots under native

herbivores, and another n=15 plots under livestock), with a total of 30 paired control-and-

fenced plots.

SOIL CARBON AND PLANT BIOMASS

Over the next 12 years, we measured soil-C eight times at roughly inter-annual intervals in

each paired plot across the different watersheds. We collected soils with a 5 cm diameter and

20 cm deep cores, and measured carbon content (TruSpec, Leco, USA) to estimate soil-C

stocks (kg C m^{-2} up to 20 cm depth) under livestock and native herbivores; soil-depth rarely

exceeds 20 cm in this ecosystem and in other similar mountainous landscapes (Bagchi &

Ritchie, 2010b). We measured the difference in standing above- and belowground biomass at

peak of growing season between paired fenced and grazed plots as an indicator of grazing

pressure (Fig. S6). We collected live shoot biomass from 0.5×0.5 m^2 quadrats in each paired

plot. Subsequently we first sun-dried and then oven-dried the biomass samples to constant

weight at 40 °C to obtain the dry-weights (g m^{-2}). We separated live roots and other

belowground structures such as rhizomes from the soil cores, and similarly obtained dry-

weights (g m^{-2} up to 20 cm depth) (Fig. S6).

SOIL MICROBIAL FUNCTIONS

In 2016, after up to 12 years of herbivore-exclusion, we estimated grazer-exclusion effects on

soil microbial processes in the paired-adjacent fenced and control plots to assess whether they

explain any observed differences in soil-C. These soil microbial variables represent different

processes that collectively influence nutrient cycling in soil (Table S1). Residence-time of

soil-C in such subhumid ecosystems is 10-30 years, whereas it can be in excess of 100 years

in swamps and wetlands (Carvalhais et al., 2014; Raich & Schlesinger, 1992). Thus, much of

the soil-C from the time of initial replacement of herbivore assemblages would have

undergone considerable turnover. So, we expect our long-term grazing-exclusion experiment

to reflect how the two types of herbivore assemblages influence soil-C at decadal timescales

that is relevant to their functional influence on climate (Naidu et al., 2022). In this way, the

grazing-exclusion experiment offers an opportunity to assess if differences in plant and

microbial responses to grazing by different types of herbivores can explain how they

influence soil-C.

In 2016, we estimated the response of key soil microbial functions to grazing: basal

respiration (BR), potential respiration or substrate-induced respiration (SIR), broad soil

microbial community profile as bacterial and fungal fractions (B and F), microbial biomass C

and N (i.e., MBC and MBN), labile carbon (LC), labile nitrogen (LN), and recalcitrant

carbon (RC) fractions of soil organic matter, microbial metabolic quotient (qCO_2), and

microbial carbon use efficiency (CUE). Since these microbial processes have seasonal

dynamics (Bagchi et al., 2017; Roy & Bagchi, 2022), we measured them at five time-points

during the growing season between May and September (Fig. S7). We followed standard

laboratory protocols to measure these microbial variables (Bagchi et al., 2017; Robertson et

al., 1999; Roy & Bagchi, 2022). Briefly, for BR we used an alkali-trap to incubate 4 g soil at

60% water holding capacity and measured respired CO_2 (mg C g^{-1} soil day^{-1}). Similarly, we

measured SIR (mg C g^{-1} soil day^{-1}) after adding 0.5% (w/v) of glucose to soil. We estimated

broad microbial community profile as fungal and bacterial contributions to potential

respiration under selective inhibition using Streptomycin (anti-bacterial) and Cycloheximide

(anti-fungal) with respect to controls. We used chloroform fumigation-extraction to estimate

MBC (mg C g^{-1} soil), and MBN (mg N g^{-1} soil). We measured labile and recalcitrant

fractions (mg g^{-1} soil) of soil organic matter (SOM) using two-step hydrolysis with H_2SO_4.

We estimated qCO_2 as the ratio of BR and MBC (respiration per unit biomass, hr^{-1}). We

estimated CUE from established relationships between key microbial extracellular enzymes

(carbon-acquiring β-glucosidase, alongside nitrogen-acquiring Leucine aminopeptidase and

N-acetyl-β-D-glucosaminidase), MBC and MBN, and labile fractions of C and N in SOM

(Sinsabaugh et al., 2013, 2016). We measured activity of the three enzymes using their

specific fluorogenic substrates (4-MUB-β-D-glucopyranoside for β-glucosidase, L-Leucine-

7-amido-4-methylcoumarin hydrochloride for Leucine aminopeptidase, and 4-MUB-*N*-

acetyl-β-D-glucosaminide for *N*-acetyl-β-D-glucosamindase) (German et al., 2011) in soils

(Fig. S8). These parameters influence microbial metabolism to determine their investment in

growth (*G*) and respiration (*R*), so that $CUE = \frac{G}{G+R}$ (Sinsabaugh et al., 2013, 2016).

Theoretically CUE can range between 0 and 1, i.e., nil to perfect efficiency, with higher

values indicating greater potential for C-sequestration. But, CUE in nature generally does not

exceed 0.6 (Sinsabaugh et al., 2013, 2016).

SOIL MICROBIAL DNA

Based on patterns seen in fungal and bacterial contribution to soil respiration, we evaluated

community composition in more detail using their DNA markers from soils collected in 2019

(Fierer et al., 2005; Maestre et al., 2015). We used soils from a subset of 20 paired-adjacent

plots and extracted microbial DNA (MOBIO Power Soil DNA Extraction Kit). Since fungal

and bacterial fractions can have seasonal dynamics (Bagchi et al., 2017), we covered four

time points during the growing season (May, July, August, and September, for a total of n =

160 samples). Next, we quantified the Internal Transcribed Spacer regions (ITS-1 and ITS-2)

of rDNA for fungal abundance, and region of 16s rDNA for bacterial abundance with

quantitative polymerase chain reaction, qPCR (Fierer et al., 2005; Maestre et al., 2015). We

used these qPCR data to confirm whether DNA-based estimates align with respiration-based

estimates of fungal and bacterial abundance. Additional details are in Appendix.

From these DNA extracts for a subset of ten paired plots, we estimated microbial

taxonomic diversity in soil from Operational Taxonomic Units (OTUs). To capture seasonal

community dynamics, we estimated OTUs at three time points during the growing season

(May, July, and September, for a total of n = 60 samples of community wide OTUs). We

amplified and sequenced the 16S rRNA hyper variable V3-V4 region in Illumina MiSeq

platform, followed by taxonomic assignment using the SILVA_v138 database. Additional

details are in Appendix.

VETERINARY ANTIBIOITICS IN SOIL

Herbivore-mediated changes in plant communities, with subsequent alteration in the

underlying distribution of plants traits, can influence the quantity and quality of C-input to

soil (Bagchi & Ritchie, 2010a). This can lead to differences in microbial communities

between livestock and native herbivores due to alteration in labile and recalcitrant fractions of

soil organic matter (Bagchi & Ritchie, 2010b; Laliberté & Tylianakis, 2012). A more direct

effect on microbial communities can arise from veterinary use of antibiotics on livestock that

eventually enter soil via dung and urine (Albero et al., 2018; Jechalke et al., 2014; Kemper,

2008; O'Connor & Aga, 2007; Wepking et al., 2019). Livestock in Spiti are frequently

treated with antibiotics such as tetracycline (therapeutic and sub-therapeutic dosage). Among

the native herbivores, veterinary care for yaks is comparatively rare, while it is practically

absent for ibex and bharal. So, based on the patterns seen in microbial OTUs, we quantified

tetracycline (μg kg^{-1} soil) in soil from all 30 paired plots in August 2022. We used enzyme-

linked immunosorbent assays (ELISA) to quantify water-extractable tetracycline levels in

soil with a detection limit of 3 ppb (Immunolab, GmbH, Germany), relative to standards and

controls (Aga et al., 2003; O'Connor & Aga, 2007).

DATA ANALYSIS

We evaluated competing hypotheses (Betts et al., 2021; Tredennick et al., 2021) to determine

whether the observed variation in the data merely represent spatio-temporal heterogeneity

between replicates, or it is necessary to invoke grazing (fenced or grazed) and herbivore

assemblage-type (livestock or native) as an explanatory variable. We used a linear mixed-

effects model to evaluate long term trends in soil-C and grazer-exclusion effects on each

plant and microbial variable. For soil-C we used a null-model $\left(M_{null}^{soilC}\right)$ with time (years,

2006-16) and plot identity as random-effects to account for background variation due to

sampling locations (plots) and times (year). This model was evaluated against the alternative

competing model $\left(M_{alt}^{soilC}\right)$ which included assemblage-type and grazing as fixed-effects,

alongside the random-effects. We compared these competing models based on parsimony,

goodness-of-fit, and likelihood-ratio test, i.e., AIC, RMSE, and LR (Tredennick et al., 2021).

We estimated grazer-effect on each microbial variable as the log-ratio of values in the

fenced and paired-adjacent control plots i.e., $\Delta= ln\left(\frac{Grazed_i}{Fenced_i}\right)$ for the i^{th} pair, and calculated

the mean and 95% CI of the log-ratio (Roy & Bagchi, 2022). As earlier, we first evaluated a

null-model to assess whether the observed variation in grazer-effects can simply be attributed

to spatio-temporal heterogeneity. For the null-model for the j^{th} microbial variable $\left(M_{null}^{j}\right)$, we

used plot identity and month (May-September) as random-effects. In the competing

alternative model for each variable $\left(M_{alt}^{j}\right)$, we included herbivore assemblage-type as a

fixed-effect alongside the random-effects. As before, we used AIC, RMSE and LR for model

comparison to judge whether herbivore assemblage-type is a necessary explanatory variable

for the data (Tredennick et al., 2021). We assessed whether the data met statistical

assumptions from observed and theoretical quantiles of residuals; the variables required no

further transformation (Δ is already in log-scale). We used 'nlme' library in R (Pinheiro et al.,

2020) to perform these analyses. As the data on antibiotics had many zeros (i.e., tetracycline

concentration in many samples was below the detectable limit of 3 ppb), we used a

generalized mixed-effects model with binomial distribution and logit-link.

We assessed the inter-relationships between grazer-effects (Δ) on different microbial

variables with structural equations models (SEM). SEM help evaluate hypothesized pathways

over how variation in one variable can influence another (Grace, 2006). We identified SEM

paths from *a-priori* examples in the literature (Table S2). To avoid multicollinearity in the

SEM arising from redundancy among the variables, we removed those variables which were

derived from one another (Fig. S9). So, ΔSIR, ΔBR, ΔRC and ΔqCO$_2$ were not included in

this step, and we built SEM using ΔMBC, ΔMBN, ΔLN, ΔLC, ΔF, ΔB, and ΔCUE. We

incorporated SEM paths as mixed-effects models for each variable as piece-wise structural

equations (Lefcheck, 2016). We used directed paths (e.g., ΔF$\rightarrow$$\Delta$CUE) as tests of

hypothesized causal relationships, and bidirectional paths (e.g., ΔMBC$\leftrightarrow$$\Delta$MBN) for

variables that are correlated or stoichiometrically coupled (Naidu et al., 2022). We did not

use genetic data from ITS and 16s rDNA in the SEM because these were collected from a

subset of plots, they were also not from the same year as the other microbial variables, and

because estimates from e-DNA broadly matched respiration-based estimates of community

composition. Similarly, we did not include tetracycline data in the SEM because when

tetracycline abundance was below the detection limit of ELISA (3 ppb), we considered it to

be zero—this makes Δ undefined (Δ becomes ∞). First, we evaluated inter-relationships

between grazer-effects (Δ) on different microbial variables with herbivore assemblage-type

as an explanatory variable. We assessed goodness of fit with Fisher's C statistic (Roy &

Bagchi, 2022; Shipley, 2009) and report standardised path coefficients alongside their

statistical significance. When SEM paths are supported by data, then they offer a plausible

explanation for an underlying process. When SEM paths are not supported, the hypothesized

process may not give rise to the observed data.

Next, since the temporal scale and span of the microbial variables (monthly

dynamics) are different from that of plant biomass and soil-C (annual or decadal dynamics),

the first SEM by itself cannot directly evaluate how grazer-effects ultimately influence soil-C

via microbes. While these different types of data differ in their temporal dimensions, they

have a common spatial dimension (n=30 replicate pairs). So, we evaluated their connections

under the expectation that effects seen on microbial processes could accumulate and

potentially explain variation in soil-C. For this, we first calculated the average (mean effect

on j^{th} variable, $\bar{\Delta}$) across all the months for each microbial variable, and across years for soil-

C and grazing intensity, for all replicates. In this second SEM, we evaluated paths using these

average values to assess whether herbivore assemblage type explains variation in soil-C via

grazer-effects on microbes, alongside the effects originating from plants. As before, plot

identity was a random-effect in the SEM paths as this allows the intercept to vary between

samples that may have unmeasured background difference in $\bar{\Delta}$ due to other factors such as

variation grazing pressure, or in edaphic conditions (Naidu et al., 2022; Roy & Bagchi,

2022). We summarized and visualized the variation in microbial community composition

with Principal Coordinates Analysis (PCoA) over Sorensen dissimilarity. We determined the

effect of herbivore assemblage type and grazing on microbial community composition using

999 randomized iterations of the dissimilarity matrix. All analyses were performed in R 4.1.0.

Results

LONG-TERM TREND IN SOIL-CARBON

Model comparison showed that spatio-temporal heterogeneity was not sufficient to explain

long-term variation in soil-C, and it could instead be attributed to herbivore assemblage-type.

There was stronger support for M_{alt}^{soilC} than for M_{null}^{soilC} (ΔAIC=-74.39, ΔRMSE=-0.662,

LR=80.386, P<0.001). However, soils under native herbivores stored 1.55±0.3 SE more

carbon (kg C m^{-2}) than under livestock ($F_{1,184}$=40.11, P<0.001). Effectively, areas under

livestock could realize up to 71% of the soil-C found under native herbivores (Fig. 2).

Grazer-exclusion, by itself ($F_{1,178}$=3.66, P=0.06) and as interaction ($F_{1,178}$=1.25, P=0.26), had

weak influence on soil-C (Fig. 2) even though the fenced plots received higher C-inputs from

plant biomass that was not consumed by grazers (Fig. 3).

GRAZER-EFFECTS ON PLANT AND SOIL VARIABLES

Analysis of grazer-effects (grazed/fenced comparison as log-ratio Δ) indicated that the

observed difference in soil-C (Fig. 2) was related to how soil microbes respond to difference

in herbivore assemblage-type (whether livestock or native herbivores, Fig. 3). Livestock and

native herbivores had comparable effects on B, 16s rDNA abundance, MBC, MBN, LC, LN,

RC, qCO$_2$, and shoot biomass (Fig. 3, Table S3). For these variables, the competing

alternative models were not supported, and their variability in grazer-effects could be

attributed to spatio-temporal heterogeneity rather than to herbivore assemblage-type (Fig. 3,

Table S3). Experimentally removing livestock and native herbivores had comparable effects

on soil-C as well (Fig. 3). But livestock and native herbivores differed in their influence on

BR, SIR, F, ITS rDNA abundance, below-ground biomass, and on CUE. For these six

variables, there was stronger support for the competing models than the null models (Table

S3), as assemblage-type was an important explanatory variable over and beyond background

spatio-temporal heterogeneity (Fig. 3). Grazing by livestock increased both basal and

potential microbial respiration (BR and SIR), compared to grazing by native herbivores (Fig.

3). But grazing by livestock reduced both the fungal fraction of microbial respiration and the

abundance of fungal genetic markers (F and ITS). Respiration-based estimates of fungal and

bacterial fractions were consistent with DNA-based estimates (Fig. 3). Importantly, grazing

by livestock reduced CUE, whereas grazing by native herbivores increased it (Fig. 3).

INTER-RELATIONSHIPS BETWEEN SOIL MICROBIAL RESPONSES

To assess how differences in grazer-effects on one microbial variable may influence other

variables, we used mixed-effects structural equation model. The model was supported by the

data (SEM: Fisher's C=4.409, P=0.633, df=6, AIC=66.41). This showed, that change in

herbivore assemblage-type influences microbial community composition to alter CUE

(Assemblage→ΔF→ΔCUE, Fig. 4). Specifically, change in herbivore assemblage from

native to livestock alters the soil microbial community, which in turn is detrimental for CUE

(Fig. 4). This SEM indicates that direction and magnitude of grazer-effects are consistent

with higher CUE under native herbivores suggesting higher potential for net soil-C storage,

compared to livestock (Fig. 2, Fig. 4).

INFLUENCE OF PLANT AND MICROBIAL RESPONSES ON SOIL-C

We considered the role of herbivore assemblage-type (identity of herbivores) and vegetation

biomass (shoot and root biomass), quantity and quality of C-input to soil (labile and

recalcitrant pools), together, as a thought experiment to connect the inter-relationship

between different microbial responses with soil-C (Fig. 5). This SEM showed that changing

herbivore assemblage-type from native to livestock is detrimental to soil-C via two paths.

First, herbivore assemblage-type affects soil-C through a decline in quantity of C-input to soil

particularly from belowground root biomass (Assemblage→$\overline{\Delta Root}$→$\overline{\Delta soilC}$, Fig. 5).

Second, it also affects soil-C through effects on microbial community composition and

microbial CUE (Assemblage→$\overline{\Delta F}$→$\overline{\Delta CUE}$→$\overline{\Delta soilC}$; Fig. 4-5). In this way, it supported our

expectation that effects arising from microbial processes at monthly scales (seen in Fig. 4)

could indeed explain variation in soil-C (Fisher's C=8.92, *P*=0.71, df=12, AIC=128.92).

MICROBIAL COMMUNITIES AND VETERINARY ANTIBIOTICS IN SOIL

Following the connections between herbivores, microbes, and soil-C (Fig. 5), we evaluated

microbial community composition in more detail using 16S rDNA sequencing and

classification. Overall, Actinobacteriota, Acidobacteriota, Chlorofelxi, Planctomycetota, and

Proteobacteria were the common and abundant taxa (Fig. S10). Microbial community

composition differed between native herbivores and livestock ($F_{1,56}$=8.03, P<0.001; Fig. 6a).

However, grazer-exclusion did not alter microbial communities regardless of herbivore

assemblage-type (Fig. 6a).

To explore whether the difference in microbial communities could be related to

veterinary use of antibiotics on livestock, we quantified tetracycline concentration in soil. As

expected, grazed plots under livestock (mean 6.44 ± 1.42 SE, µg kg^{-1} soil) contained nearly

three times more tetracycline residues than under native herbivores (2.59 ± 1.27 SE, µg kg^{-1},

Fig. 6b). Expectedly, grazer-exclusion resulted in depletion of antibiotics in the fenced plots

and this effect was stronger for livestock (t_{28}=-3.37, P=0.002; Fig. 6b).

Discussion

We assessed whether livestock could provide climate-mitigation services through soil-C to

the same extent as the native herbivores they displace (Asner et al., 2004; Bar-On et al.,

2018; du Toit et al., 2012; Schowanek et al., 2021), as this has implications for grazing

management (Bossio et al., 2020; Cromsigt et al., 2018; Reinhart et al., 2022; P. Smith,

2014). We find that:- (1) multi-species herds of livestock can have remarkable functional

similarities with the native assemblages they displace (Fig. 1), (2) but livestock do not

emerge as perfect substitutes since they store less soil-C, compared to the native herbivores

(Fig. 2), (3) differences in soil microbial responses can be the proximate explanation for why

livestock are not perfect substitutes, in addition to above-ground grazer-effects that are

propagated via plants (Fig. 3-5), and (4) there is supporting evidence that veterinary

antibiotics can restrict soil-C under livestock (Fig. 6). These results can have implications for

how we manage environmental impacts of livestock (Delabre et al., 2021; du Toit et al.,

2012; Sayer et al., 2013; P. Smith, 2014; Veblen et al., 2016), and meet the rising demand for

livestock-products in the coming decades (Leclère et al., 2020; Thornton, 2010).

To determine the scope of improved management of a large fraction of the earth's

terrestrial surface (Asner et al., 2004; Bar-On et al., 2018), it is important to identify the

potential drivers and mechanisms that explain why livestock lag behind native herbivores in

maintaining soil-C. Below we discuss explanations for the observed differences in soil-C

stocks and the candidate underlying mechanisms which can be subjects for further study.

PRE-EXISTING CONDITIONS—Since our study began after humans had already settled in

the Trans-Himalayan landscape and the livestock had already displaced the native herbivores,

sensu stricto, we cannot resolve whether livestock occupied the watersheds that were

inherently low in soil-C (i.e., $time_0$). Arguably, any pre-existing difference in soil-C can be

maintained over subsequent decades due to comparable grazing pressures and because

turnover of soil-C can occur over 10 to 30-year timescales. While pre-existing differences

would explain Fig. 2, but they do not explain why livestock and native herbivores differ in

their influence on soil microbial decomposers (Fig. 3). Pre-existing conditions also do not

explain why the direction and magnitude of microbial responses is related to soil-C (Fig. 4-

5). Alternatively, one can invoke a related argument that livestock have historically

overgrazed their watersheds, thus, depleting their soil-C stocks. If this was a primary reason,

then one expects soil-C to recover after cessation from grazing in the fenced plots. One also

expects appreciable recovery within the decadal timescale of our study (Jones & Schmitz,

2009). However, the fenced plots in livestock watersheds did not show such a response even

after 12 years of grazing-exclusion (Fig. 2). Instead, excluding native herbivores and

livestock led to comparable responses in soil-C (Fig. 3). This undermines the argument that

livestock watersheds are yet to recover from historical overgrazing (Fig. 2). Rather, we find

that under grazing by both types of herbivores, soils effectively retain a greater fraction of

their C-input from plants (Roy & Bagchi, 2022). Contrastingly in the absence of grazing,

while soils receive greater C-input from plants, but are "leaky" as they fail to store soil-C,

relative to the grazed plots. Such responses are known from many different ecosystems

around the world and are attributable to how microbes respond to grazing (Roy & Bagchi,

2022). Finally, antibiotic residues in soil (Fig. 6) also point away from the primacy of pre-

existing differences, and toward more recent factors that may have altered the microbial

processes affecting soil-C. Seen together, pre-existing conditions cannot be the only

explanation for the observed differences between livestock and native herbivores. Instead, it

is plausible that cascading effects of antibiotics on microbial community composition and

ultimately on microbial CUE play an important role, alongside the effects which originate

from plants (Fig. 4-6).

PLANT PATHWAYS—We attempt to find potential links between plant- and microbe-

centric processes, above- and below-ground, and how they influence soil-C (Fig. 5). This

recapitulates previous findings in terms of differences in the quantity of C-input from plants

(Bagchi & Ritchie, 2010b), and accommodates downstream consequences such as quality of

C-input due to changes in vegetation community composition that can alter labile and

recalcitrant pools of soil organic matter (Roy & Bagchi, 2022). In other words, while the

difference in vegetation composition is a determinant of the differences in soil-C, microbial

responses also play an equally important role (Fig. 4-5). Unfortunately, these microbial

responses do not erase the effects arising from plants and they prevent livestock from

becoming perfect surrogates of the native herbivores (Fig. 4-5).

SOIL MICROBIAL PATHWAYS—Livestock and native herbivores differ in several ways

in how they influence soil microbial processes (Fig. 3). These microbial responses lead to

lower CUE in soils under livestock, compared to the native herbivores (Fig. 4). CUE is a

fundamental microbial trait that determines the rate at which soil organic matter is

metabolized, therefore, how much C can be stored in soil (Fontaine et al., 2004; Sinsabaugh

et al., 2013; Six et al., 2006). Not surprisingly, microbial community composition plays an

important role because grazing by livestock is seen to reduce CUE, compared to the native

herbivores (Fig. 3-4), and this can be consequential for soil-C (Fig. 5). So, the question

becomes—why do livestock establish a different microbial community (Fig. 6)?

LIVESTOCK AND SOIL MICROBIAL COMMUNITIES–Several overlapping factors can

alter microbial communities under livestock, and their individual roles may not be easily

disentangled from each other especially if they act in the same direction. Once established,

the differences in microbial community composition may persist for long periods (López

Zieher et al., 2020) due to reorganization of competition-cooperation interactions (Lechón-

Alonso et al., 2021); indeed, we find them to be unresponsive to decadal-scale grazer-

exclusion (Fig. 6). Differences in vegetation composition, and subsequent alteration in the

quantity and quality of C-input from plants can result in microbial community differences.

Additionally, veterinary care offered to the livestock releases antibiotics into soil via their

dung and urine (Albero et al., 2018; Jechalke et al., 2014; Kemper, 2008; O'Connor & Aga,

2007). This can alter microbial communities towards lower CUE (Lucas et al., 2021;

Wepking et al., 2019). Annually, about 10^7 kg of antibiotics are used on livestock around the

world. Within a few hours of administration, these antibiotics can enter the environment

either unaltered, or as metabolites that continue to remain active (O'Connor & Aga, 2007).

Indeed, soils under livestock had higher tetracycline residues (Fig. 6b). Moreover, select

microbial phyla (e.g., Bacteriodota, Firmicutes, Myxococcota, Fig. S10) that are known to be

particularly susceptible to antibiotics in lab-incubations (Qian et al., 2016; Shawver et al.,

2021), indeed also appear to have different abundance under livestock, compared to native

herbivores. This raises two unanswered questions: (1) does susceptibility to antibiotics lead to

a difference in microbial community composition? And, (2) does the resultant difference in

microbial community composition impact CUE to become consequential for soil-C? There is

scattered evidence from laboratory cultures on how veterinary antibiotics alters soil microbial

communities (Qian et al., 2016; Shawver et al., 2021), and how various microbial taxa differ

in their CUE (T. P. Smith et al., 2021). However, we are yet to fully address how antibiotics

can influence microbial CUE in soil in ways that ultimately also affect the global C-cycle

(Lucas et al., 2021; Wepking et al., 2019). Seen together, our results identify microbial CUE

as a proximate explanation for why livestock are not perfect functional substitutes for native

herbivores (Fig. 3-6), to highlight the importance of soil microbial restoration in tandem with

sequestering their antibiotics. In other words, livestock management may offer improved soil-

C services (Bossio et al., 2020; Cromsigt et al., 2018; Reinhart et al., 2022; P. Smith, 2014) if

it is coupled with microbial restoration and rewilding (Averill et al., 2022; Mills et al., 2017;

Paluch et al., 2013; Wubs et al., 2016). But several questions remain unanswered on how to

achieve effective microbial rewilding and whether it can sustain ecosystem-level benefits,

and these can be subjected to further study.

MICROBIAL REWILDLING

One possible path to favor soil-C under livestock may lie in rewilding and restoring microbial

communities to achieve improved ecosystem functions and services (Averill et al., 2022;

Mills et al., 2017; Wubs et al., 2016). Although microbial restoration through soil-

inoculations is practiced in croplands (Verbruggen et al., 2013), whether target microbial

communities can establish and persist under livestock remains unknown because this problem

is compounded by veterinary antibiotics. It is also unclear how we can sequester antibiotics to

prevent their impact on soil-C and on global climate. Further studies on effective rewilding of

microbial communities and sequestering antibiotics are needed to make livestock-production

more environmentally responsible (Bossio et al., 2020; Cromsigt et al., 2018; du Toit et al.,

2012; P. Smith, 2014).

In conclusion, our results highlight that multi-species and functionally diverse

assemblages of livestock can partially approximate the ecological roles of native herbivores.

This supports continued conservation of native herbivores and calls for new ideas to improve

livestock management. Extensive therapeutic and sub-therapeutic use of antibiotics on

livestock may restrict soil-C by altering soil microbial CUE. Sequestering antibiotics,

alongside effective restoration and rewilding of soil microbial communities may enable

livestock to emulate the services provided by native herbivores to achieve nature-based

climate solutions. These may help reconcile the rising demands for livestock-products with

ecosystem functions and services.

Acknowledgements

CSIR-India provided graduate fellowship to SR; D.G.T.N. was supported by a graduate

fellowship from the Divecha Centre for Climate Change; SB was supported by MHRD-India.

DST-SERB, DST-FIST, DBT-IISc, ISRO-STC and MoEFCC provided funding support for

fieldwork and lab-analyses. Govt. of Himachal Pradesh has supported our long-term research

in Spiti. We thank Deepak Shetti, Dorje Chhering, Dorje Sheru Chhewang, Tandup Sushil

Dorje, and the people of Kibber for help in the field and in the laboratory. We received

generous support from Abhilash Vijay Nair, Ananya Jana, Atish Roy Chowdhury, Debarshi

Dasgupta, Dipshikha Chakravortty, Ishika Pramanick, Kavita Isvaran, Praveen Karanth,

Pronoy Baidya, Raju Roy, and Ritika Chatterjee. Discussions with Ben Bond-Lamberty,

Mayank Kohli, and Oswald Schmitz helped us prepare the manuscript.

References:

Aga, D. S., Goldfish, R., & Kulshrestha, P. (2003). Application of ELISA in determining the fate of tetracyclines in land-applied livestock wastes. *The Analyst*, *128*(6), 658. https://doi.org/10.1039/b301630g

Albero, B., Tadeo, J. L., Escario, M., Miguel, E., & Pérez, R. A. (2018). Persistence and availability of veterinary antibiotics in soil and soil-manure systems. *Science of The Total Environment*, *643*, 1562–1570. https://doi.org/10.1016/J.SCITOTENV.2018.06.314

Asner, G. P., Elmore, A. J., Olander, L. P., Martin, R. E., & Harris, A. T. (2004). Grazing systems, ecosystem responses, and global change. *Annual Review of Environment and Resources*, *29*(1), 261–299. https://doi.org/10.1146/annurev.energy.29.062403.102142

Augustine, D. J., & McNaughton, S. J. (1998). Ungulate Effects on the Functional Species Composition of Plant Communities: Herbivore Selectivity and Plant Tolerance. *The Journal of Wildlife Management*, *62*(4), 1165. https://doi.org/10.2307/3801981

Averill, C., Anthony, M. A., Baldrian, P., Finkbeiner, F., van den Hoogen, J., Kiers, T., Kohout, P., Hirt, E., Smith, G. R., & Crowther, T. W. (2022). Defending Earth's terrestrial microbiome. *Nature Microbiology*. https://doi.org/10.1038/s41564-022-01228-3

Bagchi, S., Bhatnagar, Y. v, & Ritchie, M. E. (2012). Comparing the effects of livestock and native herbivores on plant production and vegetation composition in the Trans-Himalayas. *Pastoralism*, *2*(1), 21. https://doi.org/10.1186/2041-7136-2-21

Bagchi, S., & Ritchie, M. E. (2010a). Herbivore effects on above- and belowground plant production and soil nitrogen availability in the Trans-Himalayan shrub-steppes. *Oecologia*, *164*(4), 1075–1082. https://doi.org/10.1007/s00442-010-1690-5

Bagchi, S., & Ritchie, M. E. (2010b). Introduced grazers can restrict potential soil carbon sequestration through impacts on plant community composition. *Ecology Letters*, *13*(8), 959–968. https://doi.org/10.1111/j.1461-0248.2010.01486.x

Bagchi, S., Roy, S., Maitra, A., & Sran, R. S. (2017). Herbivores suppress soil microbes to influence carbon sequestration in the grazing ecosystem of the Trans-Himalaya. *Agriculture, Ecosystems and Environment*, *239*, 199–206. https://doi.org/10.1016/j.agee.2017.01.033

Bakker, E. S., Ritchie, M. E., Olff, H., Milchunas, D. G., & Knops, J. M. H. (2006). Herbivore impact on grassland plant diversity depends on habitat productivity and herbivore size. *Ecology Letters*, *9*(7), 780–788. https://doi.org/10.1111/J.1461-0248.2006.00925.X

Bardgett, R. D., & Wardle, D. A. (2003). Herbivore-mediated linkages between aboveground and belowground communities. *Ecology*, *84*(9), 2258–2268. https://doi.org/10.1890/02-0274

Bar-On, Y. M., Phillips, R., & Milo, R. (2018). The biomass distribution on Earth. *Proceedings of the National Academy of Sciences*, *115*(25), 6506–6511. https://doi.org/10.1073/pnas.1711842115

Bellezza, J. V. (2017). The Rock Art of Spiti. A General Introduction. *Revue d'Etudes Tibétaines*, *41*, 56–85.

Betts, M. G., Hadley, A. S., Frey, D. W., Frey, S. J. K., Gannon, D., Harris, S. H., Kim, H., Kormann, U. G., Leimberger, K., Moriarty, K., Northrup, J. M., Phalan, B., Rousseau, J. S., Stokely, T. D., Valente, J. J., Wolf, C., & Zárrate-Charry, D. (2021). When are hypotheses useful in ecology and evolution? *Ecology and Evolution*, *11*(11), 5762–5776. https://doi.org/10.1002/ece3.7365

624   Bossio, D. A., Cook-Patton, S. C., Ellis, P. W., Fargione, J., Sanderman, J., Smith, P., Wood,
625   S., Zomer, R. J., von Unger, M., Emmer, I. M., & Griscom, B. W. (2020). The role of
626   soil carbon in natural climate solutions. *Nature Sustainability*, *3*(5), 391–398.
627   https://doi.org/10.1038/s41893-020-0491-z

628   Brus, D., Hengl, T., Heuvelink, G., Kempen, B., Mulder, T. V. L., Olmedo, G. F., Poggio, L.,
629   Ribeiro, E., Thine Omuto, C., Yigini, Y., Baritz, R., & Vargas, R. (2017). *Soil organic*
630   *carbon mapping: GSOC map cookbook manual*. FAO.
631   https://www.fao.org/3/bs901e/bs901e.pdf

632   Carvalhais, N., Forkel, M., Khomik, M., Bellarby, J., Jung, M., Migliavacca, M., Mu, M.,
633   Saatchi, S., Santoro, M., Thurner, M., Weber, U., Ahrens, B., Beer, C., Cescatti, A.,
634   Randerson, J. T., & Reichstein, M. (2014). Global covariation of carbon turnover times
635   with climate in terrestrial ecosystems. *Nature*, *514*(7521), 213–217.
636   https://doi.org/10.1038/nature13731

637   Cromsigt, J. P. G. M., te Beest, M., Kerley, G. I. H., Landman, M., le Roux, E., & Smith, F.
638   A. (2018). Trophic rewilding as a climate change mitigation strategy? *Philosophical*
639   *Transactions of the Royal Society B: Biological Sciences*, *373*(1761), 20170440.
640   https://doi.org/10.1098/rstb.2017.0440

641   Delabre, I., Rodriguez, L. O., Smallwood, J. M., Scharlemann, J. P. W., Alcamo, J.,
642   Antonarakis, A. S., Rowhani, P., Hazell, R. J., Aksnes, D. L., Balvanera, P., Lundquist,
643   C. J., Gresham, C., Alexander, A. E., & Stenseth, N. C. (2021). Actions on sustainable
644   food production and consumption for the post-2020 global biodiversity framework.
645   *Science Advances*, *7*(12), eabc8259. https://doi.org/10.1126/sciadv.abc8259

646   Derner, J. D., Augustine, D. J., & Frank, D. A. (2019). Does Grazing Matter for Soil Organic
647   Carbon Sequestration in the Western North American Great Plains? *Ecosystems*, *22*(5),
648   1088–1094. https://doi.org/10.1007/s10021-018-0324-3

649   du Toit, J. T., Kock, R., & Deutsch, J. (Eds.). (2012). *Wild Rangelands: Conserving Wildlife*
650   *While Maintaining Livestock in Semi-Arid Ecosystems*. John Wiley & Sons.

651   Falkowski, P. G., Fenchel, T., & Delong, E. F. (2008). The Microbial Engines That Drive
652   Earth's Biogeochemical Cycles. *Science*, *320*(5879), 1034–1039.
653   https://doi.org/10.1126/science.1153213

654   Fierer, N., Jackson, J. A., Vilgalys, R., & Jackson, R. B. (2005). Assessment of Soil
655   Microbial Community Structure by Use of Taxon-Specific Quantitative PCR Assays.
656   *Applied and Environmental Microbiology*, *71*(7), 4117–4120.
657   https://doi.org/10.1128/AEM.71.7.4117-4120.2005

658   Fontaine, S., Bardoux, G., Abbadie, L., & Mariotti, A. (2004). Carbon input to soil may
659   decrease soil carbon content. *Ecology Letters*, *7*(4), 314–320.
660   https://doi.org/10.1111/j.1461-0248.2004.00579.x

661   German, D. P., Weintraub, M. N., Grandy, A. S., Lauber, C. L., Rinkes, Z. L., & Allison, S.
662   D. (2011). Optimization of hydrolytic and oxidative enzyme methods for ecosystem
663   studies. *Soil Biology and Biochemistry*, *43*(7), 1387–1397.
664   https://doi.org/10.1016/j.soilbio.2011.03.017

665   Gilbert, M., Nicolas, G., Cinardi, G., van Boeckel, T. P., Vanwambeke, S. O., Wint, G. R.
666   W., & Robinson, T. P. (2018). Global distribution data for cattle, buffaloes, horses,
667   sheep, goats, pigs, chickens and ducks in 2010. *Scientific Data 2018 5:1*, *5*(1), 1–11.
668   https://doi.org/10.1038/sdata.2018.227

669   Grace, J. B. (2006). *Structural Equation Modeling and Natural Systems*. Cambridge
670   University Press.

671   Jechalke, S., Heuer, H., Siemens, J., Amelung, W., & Smalla, K. (2014). Fate and effects of
672   veterinary antibiotics in soil. *Trends in Microbiology*, *22*(9), 536–545.
673   https://doi.org/10.1016/J.TIM.2014.05.005

Jones, H. P., & Schmitz, O. J. (2009). Rapid Recovery of Damaged Ecosystems. *PLoS ONE*,
4(5), e5653. https://doi.org/10.1371/journal.pone.0005653
Kemper, N. (2008). Veterinary antibiotics in the aquatic and terrestrial environment.
Ecological Indicators, *8*(1), 1–13. https://doi.org/10.1016/j.ecolind.2007.06.002
Kuzyakov, Y. (2010). Priming effects: Interactions between living and dead organic matter.
Soil Biology and Biochemistry, *42*(9), 1363–1371.
https://doi.org/10.1016/J.SOILBIO.2010.04.003
Laliberté, E., & Tylianakis, J. M. (2012). Cascading effects of long-term land-use changes on
plant traits and ecosystem functioning. *Ecology*, *93*(1), 145–155.
https://doi.org/10.1890/11-0338.1
Lechón-Alonso, P., Clegg, T., Cook, J., Smith, T. P., & Pawar, S. (2021). The role of
competition versus cooperation in microbial community coalescence. *PLOS*
Computational Biology, *17*(11), e1009584. https://doi.org/10.1371/journal.pcbi.1009584
Leclère, D., Obersteiner, M., Barrett, M., Butchart, S. H. M., Chaudhary, A., De Palma, A.,
DeClerck, F. A. J., Di Marco, M., Doelman, J. C., Dürauer, M., Freeman, R., Harfoot,
M., Hasegawa, T., Hellweg, S., Hilbers, J. P., Hill, S. L. L., Humpenöder, F., Jennings,
N., Krisztin, T., … Young, L. (2020). Bending the curve of terrestrial biodiversity needs
an integrated strategy. *Nature*, *585*(7826), 551–556. https://doi.org/10.1038/s41586-020-
2705-y
Lefcheck, J. S. (2016). piecewiseSEM: Piecewise structural equation modelling in r for
ecology, evolution, and systematics. *Methods in Ecology and Evolution*, *7*(5), 573–579.
https://doi.org/10.1111/2041-210X.12512
Leslie, D. M., & Schaller, G. B. (2009). Bos grunniens and Bos mutus (Artiodactyla:
Bovidae). *Mammalian Species*, *836*, 1–17. https://doi.org/10.1644/836.1
Liu, X., Sheng, H., Wang, Z., Ma, Z., Huang, X., & Li, L. (2020). Does Grazing Exclusion
Improve Soil Carbon and Nitrogen Stocks in Alpine Grasslands on the Qinghai-Tibetan
Plateau? A Meta-Analysis. *Sustainability*, *12*(3), 977.
https://doi.org/10.3390/su12030977
López Zieher, X. M., Vivanco, L., & Yahdjian, L. (2020). Soil bacterial communities remain
altered after 30 years of agriculture abandonment in Pampa grasslands. *Oecologia*,
193(4), 959–968. https://doi.org/10.1007/s00442-020-04736-3
Lucas, J. M., Sone, B. M., Whitmore, D., & Strickland, M. S. (2021). Antibiotics and
temperature interact to disrupt soil communities and nutrient cycling. *Soil Biology and*
Biochemistry, *163*, 108437. https://doi.org/10.1016/j.soilbio.2021.108437
Lundgren, E. J., Ramp, D., Rowan, J., Middleton, O., Schowanek, S. D., Sanisidro, O.,
Carroll, S. P., Davis, M., Sandom, C. J., Svenning, J.-C., & Wallach, A. D. (2020).
Introduced herbivores restore Late Pleistocene ecological functions. *Proceedings of the*
National Academy of Sciences, *117*(14), 7871–7878.
https://doi.org/10.1073/pnas.1915769117
Maestre, F. T., Delgado-Baquerizo, M., Jeffries, T. C., Eldridge, D. J., Ochoa, V., Gozalo, B.,
Quero, J. L., García-Gómez, M., Gallardo, A., Ulrich, W., Bowker, M. A., Arredondo,
T., Barraza-Zepeda, C., Bran, D., Florentino, A., Gaitán, J., Gutiérrez, J. R., Huber-
Sannwald, E., Jankju, M., … Singh, B. K. (2015). Increasing aridity reduces soil
microbial diversity and abundance in global drylands. *Proceedings of the National*
Academy of Sciences of the United States of America, *112*(51), 15684–15689.
https://doi.org/10.1073/pnas.1516684112
Malhi, Y., Doughty, C. E., Galetti, M., Smith, F. A., Svenning, J.-C., & Terborgh, J. W.
(2016). Megafauna and ecosystem function from the Pleistocene to the Anthropocene.
Proceedings of the National Academy of Sciences, *113*(4), 838–846.
https://doi.org/10.1073/pnas.1502540113

Mills, J. G., Weinstein, P., Gellie, N. J. C., Weyrich, L. S., Lowe, A. J., & Breed, M. F.
(2017). Urban habitat restoration provides a human health benefit through microbiome
rewilding: the Microbiome Rewilding Hypothesis. *Restoration Ecology*, *25*(6), 866–872.
https://doi.org/10.1111/rec.12610

Mishra, C., Wieren, S. E., Heitkönig, I. M. A., & Prins, H. H. T. (2002). A theoretical
analysis of competitive exclusion in a Trans-Himalayan large-herbivore assemblage.
Animal Conservation, *5*(3), 251–258. https://doi.org/10.1017/S1367943002002305

Naidu, D. G. T., Roy, S., & Bagchi, S. (2022). Loss of grazing by large mammalian
herbivores can destabilize the soil carbon pool. *Proceedings of the National Academy of*
Sciences, *119*(43), 1–7. https://doi.org/10.1073/pnas.2211317119

Namgail, T., Wieren, S. E. van, & Prins, H. H. T. (2013). Distributional congruence of
mammalian herbivores in the Trans-Himalayan Mountains. *Current Zoology*, *59*(1),
116–124. https://doi.org/10.1093/czoolo/59.1.116

O'Connor, S., & Aga, D. S. (2007). Analysis of tetracycline antibiotics in soil: Advances in
extraction, clean-up, and quantification. *TrAC Trends in Analytical Chemistry*, *26*(6),
456–465. https://doi.org/10.1016/J.TRAC.2007.02.007

Paluch, E. C., Thomsen, M. A., & Volk, T. J. (2013). Effects of Resident Soil Fungi and
Land Use History Outweigh Those of Commercial Mycorrhizal Inocula: Testing a
Restoration Strategy in Unsterilized Soil. *Restoration Ecology*, *21*(3), 380–389.
https://doi.org/10.1111/J.1526-100X.2012.00894.X

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R Development Core Team, R. (2020).
nlme: Linear and Nonlinear Mixed Effects Models, R package version 3.1-148.

Price, J. N., Sitters, J., Ohlert, T., Tognetti, P. M., Brown, C. S., Seabloom, E. W., Borer, E.
T., Prober, S. M., Bakker, E. S., MacDougall, A. S., Yahdjian, L., Gruner, D. S., Olde
Venterink, H., Barrio, I. C., Graff, P., Bagchi, S., Arnillas, C. A., Bakker, J. D.,
Blumenthal, D. M., … Wardle, G. M. (2022). Evolutionary history of grazing and
resources determine herbivore exclusion effects on plant diversity. *Nature Ecology &*
Evolution, *6*(9), 1290–1298. https://doi.org/10.1038/s41559-022-01809-9

Qian, X., Sun, W., Gu, J., Wang, X. J., Sun, J. J., Yin, Y. N., & Duan, M. L. (2016). Variable
effects of oxytetracycline on antibiotic resistance gene abundance and the bacterial
community during aerobic composting of cow manure. *Journal of Hazardous Materials*,
315, 61–69. https://doi.org/10.1016/J.JHAZMAT.2016.05.002

Raich, J. W., & Schlesinger, W. H. (1992). The global carbon dioxide flux in soil respiration
and its relationship to vegetation and climate. *Tellus B*, *44*(2), 81–99.
https://doi.org/10.1034/j.1600-0889.1992.t01-1-00001.x

Ratajczak, Z., Collins, S. L., Blair, J. M., Koerner, S. E., Louthan, A. M., Smith, M. D.,
Taylor, J. H., & Nippert, J. B. (2022). Reintroducing bison results in long-running and
resilient increases in grassland diversity. *Proceedings of the National Academy of*
Sciences, *119*(36), e2210433119. https://doi.org/10.1073/pnas.2210433119

Reinhart, K. O., Sanni Worogo, H. S., & Rinella, M. J. (2022). Ruminating on the science of
carbon ranching. *Journal of Applied Ecology*, *59*(3), 642–648.
https://doi.org/10.1111/1365-2664.14100

Robertson, G. P., Coleman, D. C., Bledsoe, C. S., & Sollins, P. (1999). *Standard soil methods*
for long-term ecological research. Oxford University Press.
https://doi.org/10.1016/s0016-7061(01)00063-5

Roy, S., & Bagchi, S. (2022). Large Mammalian Herbivores and the Paradox of Soil Carbon
in Grazing Ecosystems: Role of Microbial Decomposers and Their Enzymes.
Ecosystems, *25*(4), 976–988. https://doi.org/10.1007/s10021-021-00696-8

Sayer, J., Sunderland, T., Ghazoul, J., Pfund, J.-L., Sheil, D., Meijaard, E., Venter, M.,
Boedhihartono, A. K., Day, M., Garcia, C., van Oosten, C., & Buck, L. E. (2013). Ten

principles for a landscape approach to reconciling agriculture, conservation, and other competing land uses. *Proceedings of the National Academy of Sciences*, *110*(21), 8349–8356. https://doi.org/10.1073/pnas.1210595110

Schmitz, O. J., Raymond, P. A., Estes, J. A., Kurz, W. A., Holtgrieve, G. W., Ritchie, M. E., Schindler, D. E., Spivak, A. C., Wilson, R. W., Bradford, M. A., Christensen, V., Deegan, L., Smetacek, V., Vanni, M. J., & Wilmers, C. C. (2014). Animating the Carbon Cycle. *Ecosystems*, *17*(2), 344–359. https://doi.org/10.1007/s10021-013-9715-7

Schmitz, O. J., Wilmers, C. C., Leroux, S. J., Doughty, C. E., Atwood, T. B., Galetti, M., Davies, A. B., & Goetz, S. J. (2018). Animals and the zoogeochemistry of the carbon cycle. *Science*, *362*(6419). https://doi.org/10.1126/science.aar3213

Schowanek, S. D., Davis, M., Lundgren, E. J., Middleton, O., Rowan, J., Pedersen, R. Ø., Ramp, D., Sandom, C. J., & Svenning, J. (2021). Reintroducing extirpated herbivores could partially reverse the late Quaternary decline of large and grazing species. *Global Ecology and Biogeography*, *30*(4), 896–908. https://doi.org/10.1111/geb.13264

Shawver, S., Wepking, C., Ishii, S., Strickland, M. S., & Badgley, B. D. (2021). Application of manure from cattle administered antibiotics has sustained multi-year impacts on soil resistome and microbial community structure. *Soil Biology and Biochemistry*, *157*, 108252. https://doi.org/10.1016/j.soilbio.2021.108252

Shipley, B. (2009). Confirmatory path analysis in a generalized multilevel context. *Ecology*, *90*(2), 363–368. https://doi.org/10.1890/08-1034.1

Singh, R., Sharma, R. K., & Babu, S. (2015). Pastoralism in transition: Livestock abundance and herd composition in Spiti, Trans-Himalaya. *Human Ecology*, *43*(6), 799–810. https://doi.org/10.1007/s10745-015-9789-2

Sinsabaugh, R. L., Manzoni, S., Moorhead, D. L., & Richter, A. (2013). Carbon use efficiency of microbial communities: stoichiometry, methodology and modelling. *Ecology Letters*, *16*(7), 930–939. https://doi.org/10.1111/ele.12113

Sinsabaugh, R. L., Turner, B. L., Talbot, J. M., Waring, B. G., Powers, J. S., Kuske, C. R., Moorhead, D. L., & Shah, J. J. Follstad. (2016). Stoichiometry of microbial carbon use efficiency in soils. *Ecological Monographs*, *86*(2), 172–189. https://doi.org/10.1890/15-2110.1

Sitters, J., Kimuyu, D. M., Young, T. P., Claeys, P., & Olde Venterink, H. (2020). Negative effects of cattle on soil carbon and nutrient pools reversed by megaherbivores. *Nature Sustainability*, *3*(5), 360–366. https://doi.org/10.1038/s41893-020-0490-0

Six, J., Frey, S. D., Thiet, R. K., & Batten, K. M. (2006). Bacterial and Fungal Contributions to Carbon Sequestration in Agroecosystems. *Soil Science Society of America Journal*, *70*(2), 555–569. https://doi.org/10.2136/sssaj2004.0347

Smith, P. (2014). Do grasslands act as a perpetual sink for carbon? *Global Change Biology*, *20*(9), 2708–2711. https://doi.org/10.1111/GCB.12561

Smith, T. P., Clegg, T., Bell, T., & Pawar, S. (2021). Systematic variation in the temperature dependence of bacterial carbon use efficiency. *Ecology Letters*, *24*(10), 2123–2133. https://doi.org/10.1111/ele.13840

Thornton, P. K. (2010). Livestock production: recent trends, future prospects. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1554), 2853–2867. https://doi.org/10.1098/rstb.2010.0134

Tredennick, A. T., Hooker, G., Ellner, S. P., & Adler, P. B. (2021). A practical guide to selecting models for exploration, inference, and prediction in ecology. *Ecology*, *102*(6), e03336. https://doi.org/10.1002/ecy.3336

Veblen, K. E., Porensky, L. M., Riginos, C., & Young, T. P. (2016). Are cattle surrogate wildlife? Savanna plant community composition explained by total herbivory more than

herbivore type. *Ecological Applications*, *26*(6), 1610–1623. https://doi.org/10.1890/15-
1367.1

Verbruggen, E., Heijden, M. G. A., Rillig, M. C., & Kiers, E. T. (2013). Mycorrhizal fungal
establishment in agricultural soils: factors determining inoculation success. *New*
Phytologist, *197*(4), 1104–1109. https://doi.org/10.1111/j.1469-8137.2012.04348.x

Wells, H. B. M., Porensky, L. M., Veblen, K. E., Riginos, C., Stringer, L. C., Dougill, A. J.,
Namoni, M., Ekadeli, J., & Young, T. P. (2022). At high stocking rates, cattle do not
functionally replace wild herbivores in shaping understory community composition.
Ecological Applications, *32*(3), e2520. https://doi.org/10.1002/eap.2520

Wepking, C., Badgley, B., Barrett, J. E., Knowlton, K. F., Lucas, J. M., Minick, K. J., Ray, P.
P., Shawver, S. E., & Strickland, M. S. (2019). Prolonged exposure to manure from
livestock-administered antibiotics decreases ecosystem carbon-use efficiency and alters
nitrogen cycling. *Ecology Letters*, *22*(12), 2067–2076. https://doi.org/10.1111/ele.13390

Western, D., Tyrrell, P., Brehony, P., Russell, S., Western, G., & Kamanga, J. (2020).
Conservation from the inside☐out: Winning space and a place for wildlife in working
landscapes. *People and Nature*, *2*(2), 279–291. https://doi.org/10.1002/pan3.10077

Witt, G. B., Noël, M. v., Bird, M. I., Beeton, R. J. S. (Bob), & Menzies, N. W. (2011).
Carbon sequestration and biodiversity restoration potential of semi-arid mulga lands of
Australia interpreted from long-term grazing exclosures. *Agriculture, Ecosystems &*
Environment, *141*(1–2), 108–118. https://doi.org/10.1016/j.agee.2011.02.020

Wubs, E. R. J., van der Putten, W. H., Bosch, M., & Bezemer, T. M. (2016). Soil inoculation
steers restoration of terrestrial ecosystems. *Nature Plants*, *2*(8), 16107.
https://doi.org/10.1038/nplants.2016.107

Fig. 1

Figure 1. Similarities and differences between livestock and native herbivores in Spiti, India.

Livestock (L: blue; yak-cattle hybrids, cattle, goat, sheep, horse, and donkey) and native

herbivores (N: green, yak ibex, bharal, kiang, and argali) show considerable overlap in

multidimensional trait space (a). Asterix indicates that kiang and argali are found nearby our

study area in Spiti. Trait-space consists of six key traits: body mass, dietary guild,

fermentation type, sexual-dimorphism, graminoid consumption, and limb morphology

(Lundgren et al., 2020) with Principal Coordinates Analysis on Gower's distance matrix

(PCoA). Vegetation composition summarised with NMDS ordination on Bray-Curtis

dissimilarity (b) shows that watersheds under native watersheds are dominated by grass,

whereas sedges dominate under livestock. Peak plant biomass (shoot, litter and root),

microbial biomass, and potential respiration (c) in native and livestock watersheds. Data are

from previous studies (Bagchi et al., 2017; Bagchi & Ritchie, 2010a).

Fig. 2

Figure 2. Long-term monitoring of soil-C (mean ± 95% CI) under grazed and fenced plots in

watersheds with native herbivores and livestock in Spiti, India. The difference in soil-C under

native herbivores and livestock was consistent for more than a decade and was not altered by

grazer-exclusion. Data are from n=30 paired control (grazed) and treatment (fenced) plots.

Fig. 3

Figure 3. Grazer-effect (—— , as mean ± 95% CI) of livestock and native

herbivores on 15 plant and soil microbial variables, alongside soil-C in Spiti, India. These

variables are basal microbial respiration (BR), potential microbial respiration (SIR), Fungal

fraction (F), ITS DNA-amplicon abundance (ITS), Bacterial fraction (B), 16s DNA-amplicon

abundance (16s), Microbial Biomass Carbon (MBC), Microbial Biomass Nitrogen (MBN),

Labile-C (LC) and Labile-N (LN) fractions of soil organic matter, Microbial metabolic

quotient (qCO_2), and microbial carbon use efficiency (CUE) which were measured at

monthly intervals during the growth season. Vegetation biomass (ΔShoot: from difference in

peak aboveground biomass and ΔRoot: from difference in peak belowground biomass) and

grazer-effect on soil-C (ΔsoilC) are from long-term decadal measurements. Direction and

magnitude of differences in BR, SIR, F, ITS abundance, CUE, and Root varied with

herbivore assemblage-type. But the remaining variables were not influenced by herbivore

assemblage-type. Data are from n=30 permanent plots in Spiti region of northern India

(except e-DNA data, which are from a subset of 20 plots). See companion Fig. S9.

Fig. 4

Figure 4. Mixed-effects structural equation model to evaluate inter-relationships between

grazer-effects (⎯⎯⎯⎯) on different microbial variables with herbivore assemblage-

type as an explanatory variable. The variables are ΔMBC: microbial biomass-C, ΔMBN:

microbial biomass-N, ΔLC: labile-C, ΔLN: labile-N, ΔF: fungal biomass, ΔB: bacterial

biomass, ΔCUE: carbon use efficiency. SEM paths show that as herbivore-assemblage goes

from native to livestock, a decrease in ΔCUE via ΔF. Unidirectional arrows represent

hypothesized causal paths, bidirectional arrows indicate correlated paths, and rounded arrows

indicate interactions. Standardized path coefficients and their statistical significance are

shown alongside marginal and conditional R^2 for the different variables.

Fig. 5.

SEM: Fisher's C = 9.46, P=0.663, df = 12, AIC = 107.46

Figure 5. Thought experiment with mixed-effects structural equations to explore the

influence of plant and microbial responses to grazing by native and livestock on soil-C.

Unlike Fig. 4, here   is the mean grazer-effect averaged across time (months for microbial

variables, and years for soil-C and vegetation biomass). This helps evaluate the relative

influence of herbivore assemblage-type, and vegetation (and   ), and microbial

variables, on variation in soil-C. SEM paths suggest that replacing native herbivores with

livestock affects soil-C in two ways: via effects originating from plants, and through a change

in soil microbial communities that reduces CUE. Unidirectional arrows represent

hypothesized causal paths, bidirectional arrows indicate correlated paths, and rounded arrows

indicate interactions. Standardized path coefficients and their statistical significance are

shown alongside marginal and conditional R^2 for the different variables. For clarity and to

reduce clutter, only a few unimportant non-significant paths are shown in grey (see

companion Fig. S11 for all modelled paths).

Fig 6.

Figure 6. Differences in microbial community composition and antibiotic concentration

across native and livestock watersheds. Difference in microbial community composition in

two types of watersheds summarized with Principal Coordinates Analysis on Sorensen

dissimilarity matrix of DNA-based microbial OTU data (a). Microbial communities varied

between native and livestock watersheds, and there was no difference between grazed and

fenced plots. Difference in antibiotic residues (tetracycline) in soil (b). Soils under livestock

contain more antibiotics than soils under native herbivores, and they decline after grazer-

exclusion.

926