Sexual coercion in a natural mandrill population

Authors

Nikolaos Smit[1,*], Alice Baniel[2,3], Berta Roura-Torres[4,5], Paul Amblard-Rambert[6], Marie J. E. Charpentier [1,α] and Elise Huchard[1,α]

Affiliations

[1]ISEM, UMR5554 - University of Montpellier/CNRS/IRD/EPHE, Place Eugène Bataillon (cc065), 34095 Montpellier, France

[2]Center for Evolution and Medicine, Arizona State University, Tempe, AZ, 85281, USA

[3]School of Life Sciences, Arizona State University, Tempe, AZ, 85287, USA

[4]Behavioral Ecology & Sociobiology Unit, German Primate Center, Leibniz Institute for Primate Research, Kellnerweg 4, 37077 Gottingen, Germany.

[5]Department of Sociobiology/Anthropology, Johann-Friedrich-Blumenbach, Institute of Zoology and Anthropology, Georg-August University Gottingen, Kellnerweg 6, 37077 Gottingen, Germany

[6]Projet Mandrillus, BP 52, Bakoumba, Gabon

[α]These authors contributed equally to this work

[*]Correspondence to:

Nikolaos Smit, ORCID: 0000-0003-0440-1998

Phone: +33(0)782382856; email: snikos@tutanota.com

Abstract

Increasing evidence indicates that sexual coercion is widespread. While some coercive strategies are conspicuous, such as forced copulation or sexual harassment, less is known about the ecology and evolution of intimidation, where repeated male aggression promotes future rather than immediate mating success with targeted females. Although known in humans, intimidation was recently reported in chimpanzees (*Pan troglodytes*) and chacma baboons (*Papio ursinus*), where males are regularly violent against females. Here, we investigate the nature of male coercive strategies in wild mandrills (*Mandrillus sphinx*), a primate living in large polygynandrous groups where severe male aggression towards females is rare and females can form coalitions against males. Yet, we found support for all three predictions of the sexual coercion hypothesis, namely that male aggression (1) specifically targets sexually receptive females, (2) inflicts costs to these females, and (3) increases male mating success in the long-term. These results hold true when considering only non-physical threats, or only severe aggression. Finally, we show that high-ranking females are most targeted by males, probably because of their higher reproductive performances, while high-ranking males are most coercive. These results indicate that sexual intimidation is widespread in sexually dimorphic and group-living mammals, and that males and females vary in their propensities to use, and to be exposed to sexual coercion, respectively.

1   Introduction

The diverging evolutionary interests of males and females often lead to sexual conflict. While female reproductive success is typically limited by the elevated costs of reproduction, e.g. gestation and lactation in mammals, male reproductive success is primarily determined by the number of mating partners [1]. In some species, males use sexual coercion towards females, defined as "the

use by a male of force, or threat of force, that functions to increase the chances that a female will mate with him at a time when she is likely to be fertile, and to decrease the chances that she will mate with other males, at some cost to the female" [2], to improve their mating success [2, 3].

Behavioural ecologists have traditionally documented coercive strategies that are immediately visible, such as forced copulation (when a female is physically restrained by a male to mate with him), sexual harassment (when aggression immediately precedes copulation and is directed until the female cedes; [2]) and coercive mate-guarding (when a male aggressively herds females and enforce close proximity to prevent them to copulate with rival males; [4, 5]). These forms of sexual coercion have been reported from insects [6, 7] to vertebrates [8, 9, 10, 11, 12]. In contrast, long-term forms of sexual coercion – when aggression does not translate immediately but subsequently into mating benefits for the aggressor – are more elusive and have been less studied outside of human societies. Sexual intimidation, when repeated male aggression aims at enforcing future female sexual compliance, has only been documented in two primate societies characterized by severe male aggression to females (chimpanzees (*Pan troglodytes*): [13]; chacma baboons (*Papio ursinus*): [14]). Similarly, males of different taxa (e.g. birds and primates including humans) can also punish females following copulations with rival males to prevent cuckoldry in the future [15, 16, 17, 18].

Sexual coercion is increasingly recognized as a driving force influencing the evolution of mating and social systems in animals [19, 2, 20], including humans [21, 22]. In mammals, male coercive tactics appear most common in polygynous and polygynandrous species where males compete intensively over mating opportunities and a substantial fraction of males fails to secure copulations, and where sexual size dimorphism is pronounced, allowing males to threaten or harass females at low costs [23, 24]. In these species, female impediment to male copulation attempts has been associated

with an increased risk of severe injury or even death [25]. The forms of coercion used by males
are then likely to vary according to the stability of male-female associations and male dominance
status. Short-term strategies such as sexual harassment and forced copulations may be frequently
used in solitary species, where males and females only encounter each other for mating [3]. By
contrast, long-term strategies, such as intimidation and punishment, are more likely to evolve in
species living in stable bisexual groups where males and females maintain medium- to long-term
social relationships. Furthermore, in polygynous groups, harassment and forced copulations might
be used more frequently by subordinate males that are excluded from mating opportunities [26, 27]
while long-term male coercive strategies might be used more often by dominant males to constrain
female promiscuity and impose closer proximity (e.g. [28]).

Primates are good candidates to study sexual coercion because the diversity of their social and
mating systems may promote various male and female sexual strategies, while their extensive cog-
nitive abilities, including individual recognition and long-term memory, may facilitate the use of
long-term male coercive strategies [22]. Such strategies are also promoted by the fact that many
primates live in stable bisexual groups where males and females maintain differentiated relation-
ships, and by a widespread male-biased sexual dimorphism associated with polygynous or some
polygynandrous mating systems.

In this study, we examine whether males exert sexual coercion in a large natural, polygynandrous
group of mandrills (Mandrillus sphinx), a primate from the Cercopithecidae family characterized
by an extreme sexual dimorphism in body size (males are 3.4 times heavier than females; [29]) and
canine length [30]. Mandrills are seasonal breeders and most males immigrate in the social group
at the onset of the mating season ([31]; which generally lasts every year from April to September

[32]), resulting in intense male-male mating competition [33]. Male reproductive skew is high, since the alpha male monopolizes 60-70% of reproductions [34, 35]. Female mandrills develop perineal swellings during fertility that grow in size as they approach ovulation and dominant males focus their mate-guarding efforts on maximally swollen females [36]. Yet, both sexes mate promiscuously and females may exhibit some forms of mate choice [37], for example by avoiding males' attempts to copulate or interrupting copulation before ejaculation (MJEC personal observation). Severe male aggression towards females occurs but appears relatively infrequent for human observers. Female relatives form tight social relationships [34], including aggressive coalitions against males that can, exceptionally, lead to male's death (in captivity: [38]). Studying male sexual coercion in this species, where most males are temporary residents in the group during the mating season, females can retaliate against males and severe male aggression against females is inconspicuous, appears thus highly relevant.

We test the three key predictions of the sexual coercion hypothesis [2], namely that male aggression (i) targets sexually receptive females more than females in other reproductive states, (ii) is costly to females in the form of a greater exposure to injuries, and (iii) increases male mating success with the victim. For this last prediction, we further investigate different forms of coercion by testing if aggression by a male towards a female increases his chances to mate with her within the following minutes (harassment) or within a longer time-window (intimidation). We also test whether a female that has just copulated with a given male receives immediate aggression from other male(s) as a punishment. We subsequently test whether higher-ranking males are more aggressive towards females during the mating season given the high reproductive skew in their favour. Finally, as an alternative hypothesis to sexual coercion, we test the "aggressive male phenotype" hypothesis, stating that the correlation between male aggression and mating is observed because females prefer

to copulate with aggressive males due to direct (e.g. better infanticide protection) or indirect (i.e. better genes for their offspring; [39]) fitness benefits of these male traits to females [40, 41].

2 Methods

2.1 Study system

We studied a natural population of mandrills established in 2002 by the release of 36 captive individuals followed by the release of another 29 individuals in 2006, in the Lékédi park, a private park located in Southern Gabon [42]. Starting in 2003, wild males joined the group to reproduce. In early 2012, the Mandrillus Project was set-up to study this population, benefiting from an initial habituation of these captive-born individuals to human presence. In early 2020, only 8 females from ca. 210 individuals were captive-born. All individuals were individually-recognized, daily monitored and censused.

2.2 Behavioural data

Trained observers, blind to the topic of this study, collected daily ad libitum behavioural observations and performed 5-min focal sampling on all study individuals [43]. In this study, we used 2182 hours of focal data collected on 81 adult females aged $\geq$4 yrs (mean±SD: 26.9±39.3h per female) and 670 hours collected on 34 subadult and adult males aged $\geq$9 yrs (19.7±29.2h per male), collected from August 2012 to March 2020. We included subadult males (aged 9-10 yrs) because they have usually reached their full adult body size [44] and have started competing with other males and mate with adult females [45]. During focal sampling, sexual and agonistic interactions between a focal individual and its groupmates were recorded. The observers sys-

tematically recorded copulations of males with females (n=275). Male aggressive events towards females included grasping/hitting (n=401), biting (n=18), chasing (n=65), lunging (n=383), slapping the ground (n=138) and head bobbing (n=567). For the analyses below, we ran the models including all these behaviours and we also replicated the analyses using only severe aggression (grasping/hitting, biting and chasing) or only threats (lunging, slapping the ground and head bobbing) because both categories produce different female behavioural reactions (see discussion). Dominance ranks were established separately for each sex (on a yearly basis for females and on a monthly basis for males) based on avoidance and displacements and calculated using normalized David's score ([46]; as per [47]). Female rank is maternally inherited and generally stable during a female's life [48]. Here, females were divided into three classes of equal size (high-, medium- and low-ranking) while male rank was considered as a binary variable (alpha versus non-alpha) because of the distinct behavioural characteristics of the alpha male, who monopolizes most swollen females and is relentlessly challenged by other males [49]. In the test for intimidation, in case the swollen period spanned over two consecutive months, a male was considered as alpha if he achieved the highest position for at least one of these two months.

2.3 Age and male immigration patterns

The exact date of birth was known for 25 individuals. For the remaining 90 individuals, the date of birth was estimated using body size, condition and patterns of tooth eruption and wear [50]. The error made when estimating the age of these 90 individuals was less than a year (50 individuals), two years (26 individuals), three years (13 individuals) or five years (1 individual). Long-term life-history and demographic data were also available from all individuals.

Census data allowed to reconstitute patterns of male residency in the group. Here, we considered

a male as resident in a given mating season when censused in the group late during the preceding birth season, between January and March. When censused for the first time during the mating season (which takes place once per year between April and September) we considered the male as immigrant. For immigrant males, the first census date was the "arrival date". Each year, the day of arrival of the first immigrant male in the group was considered as the onset of the mating season (figure S1).

2.4 Female reproductive state and sex ratio

During each female estrous cycle, the perineal swelling inflates for several days until reaching a maximal swelling size around ovulation. Swelling size remains maximal for a few days before deflating within a few days. We used a scale from 0 to 3 (by increments of 0.5) to evaluate the swelling size of each female on a near-daily basis. The reproductive state of each adult female was also recorded on a near-daily basis. Each female was classified as: "non-swollen" (i.e. non-fertile phase of the cycle that does not fall within the following three categories), "swollen" (i.e. with a perineal sexual swelling), "pregnant" (i.e. with a characteristic pregnancy swelling and/or if she gave birth 163-190 days afterwards (average gestation length: mean$\pm$SD: 175.0$\pm$4.7 days; [32]) or "lactating" (i.e. nursing a $\leq$6 month-old infant without having resumed cycling). Finally, females were considered as nulliparous until their first parturition, and parous afterwards. We calculated monthly adult group sex ratio (SR) or group operational sex ratio (OSR) as the number of females (for SR) or females with inflating sexual swelling or swelling of maximal size (for OSR) divided by the number of males aged 9 yrs and above that were censused in the group that month.

2.5 Injuries

We recorded the occurrence, type of wound, freshness and body location of any injury on a near-daily basis on all subjects [51]. A total of 90 injuries (limping n=15, puncture of the skin n=11, bleeding or swollen skin n=48, other n=16) were recorded on 43 females over the study period. For most injuries, we did not witness the interaction and the cause but in the three cases with a known context the injury was inflicted by a male. We never observed violent female-female aggression resulting in an injury.

2.6 Statistical Analyses

To test whether male aggression targets swollen females preferentially (first prediction), we ran a binomial generalized linear mixed models (GLMMs) with a logit link function to study the relationship between the probability that a female received aggression by any (adult or subadult) male during that female focal observation (0/1; response variable) and her reproductive state at the time of observation (non-swollen, swollen, pregnant and lactating; for sample sizes, see table S1). We further controlled for the following fixed effects: female dominance rank (high-, medium- or low-ranking) to test if higher-ranking females are preferentially targeted by males, parity (nulliparous or parous) to test if parous females are preferentially targeted by males, SR to test if the number of males in relation to females in the group influences the probability of occurrence of male aggression and the duration of focal observation ($\leq$5min) to control for the observation time. Female identity and the year of focal observation were fitted as random factors. Second, we ran a similar model (same structure of fixed and random effects) with the response variable corresponding to the probability that a female received aggression by groupmates other than adult or subadult males. By doing so, we tested if swollen females were generally more targeted than

any other female, regardless of the age-sex group of the aggressor.

To test whether swollen females were more injured than females in other states (second predic-tion), we ran a binomial GLMM with a logit link function to study the relationship between the probability that a female got injured (observed injured for first time) on a given day (0/1; response variable) and her reproductive state that same day. As above, we further controlled for the follow-ing variables: female dominance rank and parity, and SR. Female identity and the year of focal observation were fitted as random factors (table S1). The daily monitoring of the group allowed us to detect with accuracy the day of occurrence of each injury.

We then tested whether males who were more aggressive also had a higher mating probability with their victim (third prediction). To study intimidation, we performed a binomial GLMM with a logit link function to test whether the rate of aggression received by a female from a given male (continuous fixed effect) before the next estrous cycle of the female increased the probability of copulation of that heterosexual dyad during the female's swollen period (0/1; response variable). The "aggression window" before the swollen period was defined as the time elapsed between the onset of the mating season (for resident males) or a male's arrival in the group a given year (for immigrant males) and until the beginning of the swollen period of the female (spanning from the first day of a female's sexual swelling to the last day where swelling size was maximal: mean±SD: 10.6±5.1 days; figure S1). We pooled focal observations from females and males (table S1). We controlled for the following fixed effects in our model: female dominance rank and parity, OSR (since we focused only on swollen females for that prediction) in the month corresponding to the first day of maximal swelling, male dominance rank (alpha vs. non-alpha) that same month in interaction with the rate of male aggression (to test whether the aggression of alpha males had

a greater impact on their mating success than the aggression of subordinate males) and the total focal observation time of the studied heterosexual dyad (during the swollen period of the female) to control for the time of observation. Female identity, male identity and year of observation were fitted as random factors. We restricted our analyses to those heterosexual dyads that were observed for at least 30 minutes of focal time during the female swollen period to avoid biases due to under-sampling that would prevent us from estimating reliably mating probability. However. we validated that our results remained similar when we used slightly different thresholds (25 or 35 minutes) or no threshold at all. We further ran the same model but restricting the swollen period to the few days of the cycle during which the female was maximally swollen (i.e. where the probability of conception is the highest; mean±SD: 2.9±2.9 days). Finally, to test for immediate effects of male aggression, we ran the same model as above considering the rate of aggression received by a female from a given male during her swollen period only (figure S1, top line).

To test for sexual harassment, we assessed for each female and male focal observation during which an aggressive event was recorded from a male to a swollen female, whether a copulation occurred or not between that same heterosexual dyad in the 150 seconds following the aggression (see electronic supplementary material; figure S2). To test for male punishment, we assessed for each female and male focal observation during which a copulation event was recorded between a male and a swollen female, whether an aggression from a different male occurred towards the copulating female in the 150 following seconds (figure S2; table S1).

We further ran GLMM with a negative binomial distribution to test whether alpha males were more aggressive than subordinates during the mating season. We used as a response variable the number of aggression events a male directed towards all adult females during each month of the

mating season (April to September). We considered only aggression towards females that were potential mating partners for males: late lactating females (during the 5th and 6th month of lactation when some females have already resumed cycling; MJEC personal observation), "non-swollen", "swollen" and early pregnant females (during the first two months of pregnancy, since males may not be able to distinguish early pregnant from "non-swollen" females). We pooled focal observations from females and each given male (table S1). We included the following explanatory variables: male dominance rank (alpha vs. non-alpha) and age (to test if younger males are more aggressive) and the OSR (to test if males are more aggressive when there are few swollen females in comparison to the number of males in the group). The observation time of a given male and all the females was log-transformed, and fitted as an offset variable. Male identity and the year of observation were fitted as random factors.

We explored an alternative scenario to sexual coercion, the "aggressive male phenotype" hypothesis [39, 52], to test whether males with aggressive phenotypes have higher mating success than less aggressive males, potentially because aggression may act as a sexually selected trait and may be chosen by females. We reran the GLMM used for testing the occurrence of intimidation, including as an explanatory variable the overall rate of aggression directed by the focal male towards any groupmate (except for adult females) during the corresponding mating season.

We ran all the above statistical tests in R version 4.0.3. For generalized linear mixed models (GLMMs; summarized in table S1) we used the glmer function of the lme4 package [53] (binomial models) and glmmTMB from the package glmmTMB [54] (negative binomial model). Whenever a singular fit was observed, we reran the relevant model with the bglmer function of the blme package [55]. Whenever necessary we increased the number of iterations and/or we changed the

optimizer of the model to achieve model convergence of the model and improve its fit. We used the Anova function of the car package [56] to test for the significance of fixed factors and computed their 95% confidence intervals. We further used the vif function of the same package to detect multicollinearities. All VIFs were <2.5 indicating no serious multicollinearities [57]. For multilevel categorical factors such as reproductive state, we switched the reference category sequentially [58] in order to test for pairwise differences between categories. We explored the distribution of residuals to validate the models using the functions testDispersion and simulateResiduals from the DHARMa package [59].

3 Results

3.1 Prediction 1: Male aggression targets swollen females

Swollen females received significantly more aggression from males (mean±SD: 0.613±1.070 bouts per hour) than females in any other reproductive state (non-swollen: 0.331±0.661, pregnant: 0.309±0.528 and lactating: 0.288±0.562; figure 1a, table 1). Such pattern was found for both severe aggression (rate toward swollen females: 0.349±0.948 bouts/hour, Chisq=12.539, p-value=0.006) and threats (0.260±0.390 bouts/hour, Chisq=8.660, p-value=0.034). By contrast, swollen females were not significantly more targeted by other groupmates (figure S3, table S2). In addition, high-ranking females received more male aggression than lower-ranking females (high-ranking females: 0.461±0.328 bouts/hour, medium-ranking females: 0.216±0.240, low-ranking females: 0.148±0.149, table 1).

3.2 Prediction 2: Swollen females are more injured

Swollen females were, on average, about five times more likely to become injured (mean±SD: 0.005±0.016 injuries per day) than females in any other reproductive state (non-swollen: 0.001±0.004, pregnant: 0.001±0.002 and lactating: 0.001±0.002; figure 1b). None of the other fixed factors, including female rank, parity and the group sex-ratio were significantly correlated with the probability of injury (table 2).

3.3 Prediction 3: Aggressive males have higher mating success with their victim

We found support for sexual intimidation in mandrills: the rate of male aggression received by a female during the time window preceding her swollen period (starting at the onset of a given mating season for resident males, or at male's arrival date in the group for immigrant males) was significantly and positively correlated to the probability of copulation of the dyad during that swollen period (figure 1c, table 3). Namely, in dyads that did copulate, the rate of male-to-female aggression before the swollen period was 0.083±0.419 (mean±SD) times per hour, while in dyads that did not copulate, this rate fell to 0.030±0.110. Alpha males copulated more than subordinate males, while female rank, parity, OSR and the interaction between male rank and aggression (Chisq=0.030, p-value=0.862) were not significantly correlated with the probability of copulation (table 3). The correlation between male aggression and mating within dyads remained significant when restricting the swollen period to the few days where a female was maximally swollen (i.e. close to ovulation, Chisq=4.574, p-value=0.032). However, the rate of male aggression calculated during the swollen period of the female (instead of before) did not significantly predict the probability of copulation during that same swollen period (table S3a). This indicates that immediate

aggression (i.e. during the swollen period) did not clearly influence female mating pattern, while previous aggressive interactions over a longer period (i.e. before the swollen period) did. The pattern of correlation between aggression and subsequent mating holds when only including severe aggression (table S3b) and becomes marginally non-significant when only including threats (table S3c). Note that the rate of severe aggression and the rate of threats a female receives from a male were moderately correlated (Kendall's tau=0.28, p-value<0.001).

We did not find support for sexual harassment and punishment. Following aggression, females copulated immediately (i.e. within 150 seconds) with their aggressor in only three out of 38 total cases of aggression observed between a male and a swollen female. Similarly, males were never observed directing aggression to a female in the 150 seconds after she copulated with a rival male (out of 173 observed copulations). Those sample sizes precluded any further formal statistical testing of those hypotheses.

Alpha males were significantly more aggressive towards adult females. Indeed, an alpha male assaulted, on average, about 2 times more adult females (mean±SD: 0.05±0.07 bouts per hour) than a non-alpha male (0.03±0.06; figure S4; table 4). In addition, males were more aggressive (marginally significant effect; table 4) when there were more swollen females in the group in relation to males but male aggression did not depend on its age (table 4).

Lastly, we did not find evidence for a female preference for aggressive male phenotypes, as females were not more likely to mate with the most aggressive males of the group (see electronic supplementary material).

4 Discussion

We found support for all three core predictions of the sexual coercion hypothesis in mandrills. First, swollen females received significantly more male aggression than other females. Elevated aggression towards females around ovulation has been observed frequently in mammals, even in species where females dominate males socially (e.g. spotted hyena (*Crocuta crocuta*): [60]), suggesting that sexual coercion is widespread. Second, swollen female mandrills were significantly more injured than females in other reproductive states. Such injuries are most likely caused by males because aggression from other groupmates did not intensify during female sexual receptivity. Male aggression thus potentially causes important fitness costs in female mandrills, as shown in other mammals exhibiting sexual coercion (e.g. feral sheep (*Ovis aries*): [61]; bottlenose dolphins (*Tursiops cf. aduncus*): [62], chacma baboons: [14], chimpanzees: [63]). These fitness costs may push females to comply and copulate more with aggressive males to avoid conflict escalation and the associated risk of injury [64, 65]. Third, our analysis suggests that increased and repeated male aggression before the receptive period increases male mating success with the targeted female at times where she is most likely fertile. This correlation holds true both with severe aggression and non-physical threats, which are only moderately correlated. Most studies on sexual coercion have focused exclusively on severe aggression [14, 13] but our results indicate that male mandrills use a wide aggressive repertoire, including threats, to coerce females. In this species, male threats (such as head-bob or ground-slap) typically produce little immediate behavioural reactions in females, but could increase their sexual compliance with the aggressor when exerted repeatedly [28], especially when male-female power asymmetry is high, as in mandrills, which display one of the largest sexual dimorphism in primates.

The observed correlation between male aggression and mating success does not seem well-explained by alternative interpretations to sexual coercion, as we did not find evidence supporting a female preference for particularly aggressive males. Females could potentially use male aggression as a badge of status [13, 66] to infer male competitive abilities, which may provide females with direct or indirect benefits [40, 41]. However, in our data, variation in aggression rates among heterosexual dyads explain male mating success better than male general aggressiveness, suggesting that male mating success reflects relational properties more than male aggressive phenotype. It is further possible that male-female aggression rates directly reflect differences in male-female spatial proximity, where males would direct more aggression to females who would happen to stand around them. However, patterns of spatial ranging in social groups are far from random, and typically reflect the group social structure, in the form of differentiated relationships (e.g. spatial proximity is positively correlated to the strength of social bond in wild boars (*Sus scrofa*) [67]). In such context, male-female aggressive rates are more likely to reflect the existence of such differentiated social bonds between males and females than a scenario where a male would attack females who randomly happen to stand in their proximity. In line with this, recent studies in chimpanzees indicate that males preferably coerce their affiliated female partners [68], mirroring observations in humans where intimate partner violence is extensive [69].

Our analyses reveal important aspects of the ecology of sexual coercion in mandrill societies. While we did not find evidence for sexual harassment, our results suggest that repeated aggression over extended periods increases mating probability to aggressors once females become fertile, and may further encourage them to stay around males who mate-guard them, as observed in hamadryas baboons (*Papio hamadryas*; [28]). Sexual intimidation has previously been shown in chimpanzees and chacma baboons [13, 14], two species characterized by relatively high male violence towards

females. We found that male mandrills use severe aggression towards swollen females more often on average than chacma baboons (mean±SD: 0.350±0.950 vs 0.130±0.190 times per hr; [14]) and at a rate that lies high within the chimpanzee's reported range [13, 63]. Such frequent use of coercion by mandrill males may relate to the fact that - unlike chimpanzees and chacma baboons - they breed seasonally, thus have a limited time window to achieve mating. Yet, swollen female mandrills are injured ca. three times less on average than chacma baboons (mean±SD: 0.005±0.016 vs 0.014±0.022 injuries per day; [14]). Hence, although male to female aggression is more frequent in mandrills than in chacma baboons, violent aggression resulting in serious injuries is probably less common.

Moreover, the fact that we did not find any evidence of punishment, likely reflects the absence of exclusive mating bonds in mandrills (outside mate-guarding episodes) and the ability of females to sneakily escape male monopolization strategies in their dense habitat. Punishment by males in response to female sexual activity with a rival has, for instance, been reported in geladas (*Theropithecus gelada*) which live in more open habitat [17] and where one leader male can aggressively defend sexual access to females from his family unit [70]. To sum-up, our results are generally consistent with expectations based on the socio-ecology of mandrills, who (i) are highly dimorphic thus where males pay low costs of intersexual aggression, (ii) breed seasonally, and where males face high pressure to mate in a relatively short period, and (iii) live in a polygynandrous mating system, and where males and females form differentiated social bonds - allowing intimidation to function - but no exclusive mating bonds, preventing the use of punishment by males.

Male dominance status appeared influential in their coercive tendencies. Alpha male mandrills were more aggressive towards females during the mating season, and they copulated significantly

more with females than non-alpha males. Given the high reproductive skew in favour of alpha male mandrills [34, 35], this result suggests that sexual coercion is an effective male reproductive strategy, although more detailed analysis is necessary in order to confirm the relationship between male coercion and reproductive success. Dominant males in other primates similarly use long-term coercive strategies to constrain female promiscuity and impose closer proximity (e.g. hamadryas baboons [28]). However, in other species, such as orang-utans, subordinate males have been reported to be more coercive, and use forced copulations more often than dominant males [27]. The use of coercive strategies may be rendered more difficult for subordinate males in group-living species compared to solitary ones, such as orang-utans, if other group members, including the alpha male, occasionally step in to defend the victim.

Our analyses further highlight that all females are not equally targeted by males. High-ranking females specifically receive more male aggression than low-ranking females, which may reflect male mating preferences because dominant females show better reproductive performances than subordinates [48, 32]. Similarly, male hyenas mate preferentially with high-ranking females [71, 72] while male chimpanzees direct more aggression towards parous than nulliparous females [13] and prefer old females [73], who have a higher rank and reproductive success than younger ones [74]. This result indicates that the highest costs of coercion are born by the most attractive females, as found in chimpanzees [13].

An important question remains whether and how female mandrills may navigate such a coercive landscape while still possibly expressing some mate choice [33]. Chimpanzee studies have raised contrasting results, with sexual coercion in some populations [13, 63] versus female mate choice in other populations [75, 76]. It is possible that such conflicting results reflect differences across

populations, or alternatively methodological differences between studies, where studies of mate choice often measure female choice through differential rates of approaches of males by females [75], while studies of sexual coercion correlate aggression and mating rates [13, 14]. The growing body of work on sexual coercion generally casts doubts on inferring mate choice from rates of approaches [4], as such approaches, as well as any affiliative interaction, could instead reflect female attempts to appease coercive males (i.e. [65]). Alternatively, it's possible that sexual coercion can co-occur with female mate choice, as is the case in humans.

Our work underlines the existence of sexual coercion in mandrills while evidence for female choice remains scarce in this species [33]. It is therefore hard, at this stage, to evaluate the freedom left for females to express their own reproductive strategies. Nevertheless, several mechanisms may help females to mitigate the constraints set by male coercion. They may form alliances with other females to defend themselves [3, 77] or heterosexual bonds with males who protect them [78]. They may also appease male aggressors to limit the risk of escalation and injuries [28, 65], fight-back against aggressors, flee, hide or close their genitals [79, 80]. Female mandrills may use some of these strategies, as their behavioural repertoire includes avoiding male approaches, laying down when males attempt to copulate with them, refusing some mating attempts [33, 37], interrupting copulation by fleeing away, seeking support from subordinate males against dominant ones (MJEC personal observation) or even forming violent coalitions against high-ranking males ([38], NS personal observation). In addition, previous studies on primates have demonstrated that female reproductive synchrony and large group sizes limit female monopolization by males (across species: [81]; in mandrills: [35]) and increase the potential for females to express their strategies, including mate choice or promiscuity [82, 83]. Therefore, the extreme size of mandrill social groups along with female reproductive synchrony, may facilitate the expression of female reproductive strategies

and reduce male coercion.

Here we report new evidence for sexual intimidation in a species where males, despite being much larger than females, are not conspicuously aggressive towards them (at least from a human observer perspective). The temporal uncoupling between male aggression and copulation explains why sexual intimidation may have long been overlooked, while it increasingly appears influential at shaping the social structure and mating system of polygynandrous mammals [20].

Ethics: All applicable international, national, and/or institutional guidelines for the care and use of animals were followed. This study was approved by the CENAREST institute (permit number, AR003/20/MESRSTT/CENAREST/CG/CST/CSAR) and adhered to the legal requirements of Gabon for the ethical treatment of non-human primates.

Data accessibility: The datasets and scripts necessary to replicate analyses included in this paper are deposited in the public depository: https://gitlab.com/nksmt/mandrills

Authors' contributions: N.S., M.J.E.C., and E.H. designed the study; B.R.T. and P.A.R collected behavioural data; N.S. performed the statistical analyses; N.S., M.J.E.C., E.H. wrote the original draft and all authors critically contributed to the draft and approved submission.

Competing interests: The authors declare no competing interests.

Funding: Several grants have funded the long-term collection of the data used in this study: Deutsche Forschungsgemeinschaft (DFG, KA 1082-20-1), SEEG Lékédi (INEE-CNRS) and Agence Nationale de la Recherche (ANR SLEEP 17-CE02-0002) to M.J.E.C, and ANR ERS-17-CE02-0008 to E.H. N.S. was funded by the State Scholarships Foundation (IKY) under the scholarship program from the proceeds of the "Nic. D. Chrysovergis" bequest.

Acknowledgements: The present work would not have been possible without the work of past and present field assistants of the Mandrillus Project and the logistical support of SODEPAL-COMILOG society. This is a Project Mandrillus publication n^oxx and ISEM n^oxx.

References

[1] Bateman AJ. 1948 Intra-Sexual Selection in Drosophila. *Heredity* **2**, 349–368.

[2] Smuts BB, w. Smuts R. 1993 Male Aggression and Sexual Coercion of Females in Nonhuman Primates and Other Mammals: Evidence and Theoretical Implications. In *Advances in the Study of Behavior* vol. 22 pp. 1–63. Elsevier.

[3] Clutton-Brock T, Parker G. 1995 Sexual Coercion in Animal Societies. *Animal Behaviour* **49**, 1345–1365.

[4] Muller MN, Thompson ME, Kahlenberg SM, Wrangham RW. 2011 Sexual Coercion by Male Chimpanzees Shows That Female Choice May Be More Apparent than Real. *Behavioral Ecology and Sociobiology* **65**, 921–933.

[5] King SL, Allen SJ, Krützen M, Connor RC. 2019 Vocal Behaviour of Allied Male Dolphins during Cooperative Mate Guarding. *Animal Cognition* **22**, 991–1000.

[6] Arnqvist G. 1989 Multiple Mating in a Water Strider: Mutual Benefits or Intersexual Conflict?. *Animal Behaviour* **38**, 749–756.

[7] Parker G. 1979 Sexual selection and sexual conflict. In *Sexual Selection and Reproductive Competition in Insects* pp. 123–166. Elsevier.

[8] Head ML, Brooks R. 2006 Sexual Coercion and the Opportunity for Sexual Selection in Guppies. *Animal behaviour* **71**, 515–522.

[9] Bro-Jørgensen J. 2011 Intra- and Intersexual Conflicts and Cooperation in the Evolution of Mating Strategies: Lessons Learnt From Ungulates. *Evolutionary Biology* **38**, 28–41.

[10] McKinney F, Evarts S. 1998 Sexual Coercion in Waterfowl and Other Birds. *Ornithological Monographs* pp. 163–195.

[11] Galdikas BMF. 1985 Subadult Male Orangutan Sociality and Reproductive Behavior at Tanjung Puting. *American Journal of Primatology* **8**, 87–99.

[12] Connor RC, Vollmer N. 2009 Sexual Coercion in Dolphin Consortships: A Comparison with Chimpanzees. *Sexual coercion in primates: An evolutionary perspective on male aggression against females* pp. 218–243.

[13] Muller MN, Kahlenberg SM, Emery Thompson M, Wrangham RW. 2007 Male Coercion and the Costs of Promiscuous Mating for Female Chimpanzees. *Proceedings of the Royal Society B: Biological Sciences* **274**, 1009–1014.

[14] Baniel A, Cowlishaw G, Huchard E. 2017 Male Violence and Sexual Intimidation in a Wild Primate Society. *Current Biology* **27**, 2163–2168.e3.

[15] Clutton-Brock TH, Parker GA. 1995 Punishment in Animal Societies. *Nature* **373**, 209–216.

[16] Valera F, Hoi H, Krištín A. 2003 Male Shrikes Punish Unfaithful Females. *Behavioral Ecology* **14**, 403–408.

[17] le Roux A, Snyder-Mackler N, Roberts EK, Beehner JC, Bergman TJ. 2013 Evidence for Tactical Concealment in a Wild Primate. *Nature Communications* **4**, 1462.

[18] Rodseth L, Novak SA. 2009 The Political Significance of Gender Violence. *Sexual coercion in primates and humans* pp. 292–321.

[19] Thornhill R, Alcock J et al.. 1983 *The Evolution of Insect Mating Systems.* Harvard University Press.

[20] Clutton-Brock T. 2021 Social Evolution in Mammals. *Science* **373**, eabc9699.

[21] Smuts B. 1992 Male Aggression against Women: An Evolutionary Perspective. *Human Nature* **3**, 1–44.

[22] Stumpf RM, Martinez-Mota R, Milich KM, Righini N, Shattuck MR. 2011 Sexual Conflict in Primates. *Evolutionary Anthropology: Issues, News, and Reviews* **20**, 62–75.

[23] Nunn C. 2000 Social Evolution in Primates: The Relative Roles of Ecology and Intersexual Conflict. *Infanticide by Males* pp. 388–419.

[24] Cassini MH. 2021 Sexual Aggression in Mammals. *Mammal Review* **51**, 247–255.

[25] Cassini M. 2000 A Model on Female Breeding Dispersion and the Reproductive Systems of Pinnipeds. *Behavioural Processes* **51**, 93–99.

[26] Boeuf BJL, Mesnick S. 1991 Sexual Behavior of Male Northern Elephant Seals: I. Lethal Injuries to Adult Females. *Behaviour* **116**, 143–162.

[27] Kunz JA, Duvot GJ, Willems EP, Stickelberger J, Spillmann B, Utami Atmoko SS, van Noordwijk MA, van Schaik CP. 2021 The Context of Sexual Coercion in Orang-Utans: When Do Male and Female Mating Interests Collide?. *Animal Behaviour* **182**, 67–90.

[28] Swedell L, Schreier A. 2009 Male Aggression towards Females in Hamadryas Baboons: Conditioning, Coercion, and Control. *Sexual coercion in primates and humans: an evolutionary perspective on male aggression against females. Harvard University Press, Cambridge* pp. 244–268.

[29] Setchell JM, Lee PC, Wickings EJ, Dixson AF. 2001 Growth and Ontogeny of Sexual Size Dimorphism in the Mandrill (Mandrillus Sphinx). *American Journal of Physical Anthropology* **115**, 349–360.

[30] Leigh SR, Setchell JM, Charpentier M, Knapp LA, Wickings EJ. 2008 Canine Tooth Size and Fitness in Male Mandrills (Mandrillus Sphinx). *Journal of Human Evolution* **55**, 75–85.

[31] Brockmeyer T, Kappeler PM, Willaume E, Benoit L, Mboumba S, Charpentier MJ. 2015 Social Organization and Space Use of a Wild Mandrill (*Mandrillus Sphinx*) Group: Mandrill Social Organization and Space Use. *American Journal of Primatology* **77**, 1036–1048.

[32] Dezeure J, Charpentier MJ, Huchard E. 2022 Fitness Effects of Seasonal Birth Timing in a Long-Lived Social Primate Living in the Equatorial Forest. *Animal Behaviour* **185**, 113–126.

[33] Setchell JM. 2016 Sexual Selection and the Differences between the Sexes in Mandrills (Mandrillus Sphinx). *American Journal of Physical Anthropology* **159**, 105–129.

[34] Charpentier MJE, Harté M, Poirotte C, de Bellefon JM, Laubi B, Kappeler PM, Renoult JP. 2020 Same Father, Same Face: Deep Learning Reveals Selection for Signaling Kinship in a Wild Primate. *Science Advances* **6**, eaba3274.

[35] Charpentier M, Peignot P, Hossaert-McKey M, Gimenez O, Setchell JM, Wickings EJ. 2005 Constraints on Control: Factors Influencing Reproductive Success in Male Mandrills (Mandrillus Sphinx). *Behavioral Ecology* **16**, 614–623.

[36] Setchell JM, Charpentier M, Wickings EJ. 2005 Mate Guarding and Paternity in Mandrills: Factors Influencing Alpha Male Monopoly. *Animal Behaviour* **70**, 1105–1120.

[37] Setchell JM. 2005 Do Female Mandrills Prefer Brightly Colored Males?. *International Journal of Primatology* **26**, 715–735.

[38] Setchell JM, Knapp LA, Wickings EJ. 2006 Violent Coalitionary Attack by Female Mandrills against an Injured Alpha Male. *American Journal of Primatology* **68**, 411–418.

[39] Fisher RA. 1915 The Evolution of Sexual Preference. *The Eugenics Review* **7**, 184–192.

[40] Cordero C, Eberhard WG. 2003 Female Choice of Sexually Antagonistic Male Adaptations: A Critical Review of Some Current Research. *Journal of Evolutionary Biology* p. 6.

[41] Pizzari T, Snook RR. 2003 Perspective: Sexual Conflict and Sexual Selection: Chasing Away Paradigm Shifts. *Evolution* **57**, 1223–1236.

[42] Peignot P, Charpentier MJ, Bout N, Bourry O, Massima U, Dosimont O, Terramorsi R, Wickings EJ. 2008 Learning from the First Release Project of Captive-Bred Mandrills Mandrillus Sphinx in Gabon. *Oryx* **42**.

[43] Altmann J. 1974 Observational Study of Behavior: Sampling Methods. *Behaviour* **49**, 227–267.

[44] Setchell JM, Dixson AF. 2002 Developmental Variables and Dominance Rank in Adolescent Male Mandrills (Mandrillus Sphinx). *American Journal of Primatology* **56**, 9–25.

[45] Setchell JM, Charpentier M, Wickings EJ. 2005 Sexual Selection and Reproductive Careers in Mandrills (Mandrillus Sphinx). *Behavioral Ecology and Sociobiology* **58**, 474–485.

[46] David HA. 1987 Ranking from Unbalanced Paired-Comparison Data. *Biometrika* p. 5.

[47] Poirotte C, Massol F, Herbert A, Willaume E, Bomo PM, Kappeler PM, Charpentier MJE. 2017 Mandrills Use Olfaction to Socially Avoid Parasitized Conspecifics. *Science Advances* **3**, e1601721.

[48] Setchell JM, Lee PC, Wickings EJ, Dixson AF. 2002 Reproductive Parameters and Maternal Investment in Mandrills (Mandrillus Sphinx). *International Journal of Primatology* p. 18.

[49] Charpentier MJE, Givalois L, Faurie C, Soghessa O, Simon F, Kappeler PM. 2018 Seasonal Glucocorticoid Production Correlates with a Suite of Small-Magnitude Environmental, Demographic, and Physiological Effects in Mandrills. *American Journal of Physical Anthropology* **165**, 20–33.

[50] Galbany J, Romero A, Mayo-Alesón M, Itsoma F, Gamarra B, Pérez-Pérez A, Willaume E, Kappeler PM, Charpentier MJ. 2014 Age-Related Tooth Wear Differs between Forest and Savanna Primates. *PLoS One* **9**, e94938.

[51] Dibakou SE, Basset D, Souza A, Charpentier M, Huchard E. 2019 Determinants of Variations in Fecal Neopterin in Free-Ranging Mandrills. *Frontiers in Ecology and Evolution* **7**, 368.

[52] Huk T, Winkel W. 2007 Testing the Sexy Son Hypothesis–a Research Framework for Empirical Approaches. *Behavioral Ecology* **19**, 456–461.

[53] Bates D, Maechler M, Bolker [aut B, cre, Walker S, Christensen RHB, Singmann H, Dai B, Scheipl F, Grothendieck G, Green P, Fox J, Bauer A, copyright on simulate.formula) PNKs. 2020 *Lme4: Linear Mixed-Effects Models Using 'Eigen' and S4.*

[54] Magnusson A, Skaug H, Nielsen A, Berg C, Kristensen K, Maechler M, van Bentham K, Bolker B, Sadat N, Lüdecke D, Lenth R, O'Brien J, Geyer CJ, McGillycuddy M, Brooks M. 2021 *glmmTMB: Generalized Linear Mixed Models Using Template Model Builder.*

[55] Dorie V. 2015 *Blme: Bayesian Linear Mixed-Effects Models.*

[56] Fox J, Weisberg S, Price B, Adler D, Bates D, Baud-Bovy G, Bolker B, Ellison S, Firth D, Friendly M, Gorjanc G, Graves S, Heiberger R, Krivitsky P, Laboissiere R, Maechler M, Monette G, Murdoch D, Nilsson H, Ogle D, Ripley B, Venables W, Walker S, Winsemius D, Zeileis A, R-Core. 2020 *Car: Companion to Applied Regression.*

[57] Zuur AF, Ieno EN, Elphick CS. 2010 A Protocol for Data Exploration to Avoid Common Statistical Problems. *Methods in Ecology and Evolution* **1**, 3–14.

[58] Pinheiro J, Bates D. 2006 *Mixed-Effects Models in s and s-Plus.* Springer Science & Business Media.

[59] Hartig F, Lohse L. 2021 *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models.*

[60] Szykman M, Engh AL, Horn RCV, Boydston EE, Scribner KT, Holekamp KE. 2003 Rare Male Aggression Directed toward Females in a Female-Dominated Society: Baiting Behavior in the Spotted Hyena. *Aggressive Behaviour* p. 18.

[61] Réale D, Boussès P, Chapuis JL. 1996 Female-Biased Mortality Induced by Male Sexual Harassment in a Feral Sheep Population. *Canadian Journal of Zoology* **74**, 1812–1818.

[62] Wallen MM, Patterson EM, Krzyszczyk E, Mann J. 2016 The Ecological Costs to Females in a System with Allied Sexual Coercion. *Animal Behaviour* **115**, 227–236.

[63] Watts DP. 2022 Male Chimpanzee Sexual Coercion and Mating Success at Ngogo. *American Journal of Primatology* **n/a**, e23361.

[64] Cassini MH. 2020 A Mixed Model of the Evolution of Polygyny and Sexual Size Dimorphism in Mammals. *Mammal Review* **50**, 112–120.

[65] Baniel A, Webb CE, Cowlishaw G, Huchard E. 2021 The Submissive Pattern of Postconflict Affiliation in Asymmetric Relationships: A Test in Male and Sexually Coerced Female Baboons. *Animal Behaviour* **175**, 87–97.

[66] Setchell JM, Jean Wickings E. 2005 Dominance, Status Signals and Coloration in Male Mandrills (Mandrillus Sphinx). *Ethology* **111**, 25–50.

[67] Podgórski T, Lusseau D, Scandura M, Sönnichsen L, Jędrzejewska B. 2014 Long-Lasting, Kin-Directed Female Interactions in a Spatially Structured Wild Boar Social Network. *PLOS ONE* **9**, e99875.

[68] Reddy RB, Mitani JC. 2020 Adolescent and Young Adult Male Chimpanzees Form Affiliative, yet Aggressive, Relationships with Females. *Journal of Human Evolution* **144**, 102813.

[69] Basile KC. 2002 Prevalence of Wife Rape and Other Intimate Partner Sexual Coercion in a Nationally Representative Sample of Women. *Violence and Victims* **17**, 511–524.

[70] Snyder-Mackler N, Alberts SC, Bergman TJ. 2012 Concessions of an Alpha Male? Cooperative Defence and Shared Reproduction in Multi-Male Primate Groups. *Proceedings of the Royal Society B: Biological Sciences* **279**, 3788–3795.

[71] Szykman M, Engh AL, Van Horn RC, Funk SM, Scribner KT, Holekamp KE. 2001 Association Patterns among Male and Female Spotted Hyenas (Crocuta Crocuta) Reflect Male Mate Choice. *Behavioral Ecology and Sociobiology* **50**, 231–238.

[72] Keddy-Hector AC. 1992 Mate Choice in Non-Human Primates. *American Zoologist* **32**, 62–70.

[73] Muller MN, Thompson ME, Wrangham RW. 2006 Male Chimpanzees Prefer Mating with Old Females. *Current Biology* **16**, 2234–2238.

[74] Pusey A, Williams J, Goodall J. 1997 The Influence of Dominance Rank on the Reproductive Success of Female Chimpanzees. *Science* **277**, 828–831.

[75] Stumpf RM, Boesch C. 2006 The Efficacy of Female Choice in Chimpanzees of the Taï Forest, Côte d'Ivoire. *Behavioral Ecology and Sociobiology* **60**, 749–765.

[76] Kaburu SSK, Newton-Fisher NE. 2015 Trading or Coercion? Variation in Male Mating Strategies between Two Communities of East African Chimpanzees. *Behavioral Ecology and Sociobiology* **69**, 1039–1052.

[77] Paoli T. 2009 The Absence of Sexual Coercion in Bonobos. *Sexual coercion in primates and humans* pp. 410–423.

[78] Smuts B. 1995 The Evolutionary Origins of Patriarchy. *Human Nature* **6**, 1–32.

[79] Eberle M, Kappeler PM. 2004 Selected Polyandry: Female Choice and Inter-Sexual Conflict in a Small Nocturnal Solitary Primate (Microcebus Murinus). *Behavioral Ecology and Sociobiology* **57**, 91–100.

[80] Huchard E, Canale CI, Le Gros C, Perret M, Henry PY, Kappeler PM. 2012 Convenience Polyandry or Convenience Polygyny? Costly Sex under Female Control in a Promiscuous Primate. *Proceedings of the Royal Society B: Biological Sciences* **279**, 1371–1379.

[81] Ostner J, Nunn CL, Schülke O. 2008 Female Reproductive Synchrony Predicts Skewed Paternity across Primates. *Behavioral Ecology* **19**, 1150–1158.

[82] Ims RA. 1990 The Ecology and Evolution of Reproductive Synchrony. *Trends in Ecology & Evolution* **5**, 135–140.

[83] Fürtbauer I, Mundry R, Heistermann M, Schülke O, Ostner J. 2011 You Mate, I Mate: Macaque Females Synchronize Sex Not Cycles. *PLoS ONE* **6**, e26144.

Figure 1: Results of the tests of the three predictions of the sexual coercion hypothesis. (a) Predicted probability of male aggression received by females as a function of their reproductive state. (b) Predicted probability for females to be injured as a function of their reproductive state. (c) Rates of male aggression (number of events per hour) received by females before their swollen period for heterosexual dyads who mated versus dyads that did not mate during the swollen period. The fitted values of the GLMMs are shown on the y-axis of panels a and b. The violin plots show the predicted probabilities (for a and b) or the raw rates (for c). Pairwise comparisons across female reproductive states and corresponding p-values are shown. 'ns': not significant ($p>0.05$); *: $p<0.05$; **: $p<0.01$; ***: $p<0.001$.

Table 1: Male aggression in relation to female reproductive state (for sample sizes, see table S1). Significant p-values and confidence intervals that did not cross zero appear in bold. The significance of each variable was assessed using chi-square tests (Chisq), while the significance of each level of a categorical variable was evaluated against a reference level (noted 'Ref') according to whether their confidence intervals (CI) overlap or not.

	Response variable: Probability of receiving aggression from adult males (0/1)				
Fixed Factor	Level	Estimate	CI 95%	Chisq	P-value
Reproductive State	Swollen (Ref: Non-Swollen)	0.442	**[0.170;0.714]**	15.926	**0.001**
	Pregnant (Ref: Non-Swollen)	0.070	[-0.132;0.273]		
	Lactating (Ref: Non-Swollen)	-0.094	[-0.309;0.122]		
	Swollen (Ref: Lactating)	0.536	**[0.268;0.804]**		
	Pregnant (Ref: Lactating)	0.164	[-0.030;0.358]		
	Swollen (Ref: Pregnant)	0.372	**[0.116;0.628]**		
Female Rank	Low Rank (Ref: High Rank)	-0.718	**[-0.981;-0.456]**	31.124	**<0.001**
	Medium Rank (Ref: High Rank)	-0.554	**[-0.904;-0.203]**		
Female Parity	Parous (Ref: Nulliparous)	0.150	[-0.230;0.529]	0.599	0.439
Group Sex Ratio		-0.014	[-0.059;0.031]	0.375	0.54
Observation Time		-0.097	**[-0.167;-0.027]**	7.459	**0.006**

Table 2: Injuries in relation to female reproductive state (for sample sizes, see table S1). Significant p-values and confidence intervals that did not cross zero appear in bold. The significance of each variable was assessed using chi-square tests (Chisq), while the significance of each level of a categorical variable was evaluated against a reference level (noted 'Ref') according to whether their confidence intervals (CI) overlap or not.

	Response variable: Probability of having an injury (0/1)				
Fixed Factor	Level	Estimate	CI 95%	Chisq	P-value
Reproductive State	Swollen (Ref: Non-Swollen)	1.183	**[0.579;1.787]**	34.535	**<0.001**
	Pregnant (Ref: Non-Swollen)	-0.452	[-1.026;0.123]		
	Lactating (Ref: Non-Swollen)	-0.507	[-1.076;0.061]		
	Swollen (Ref: Lactating)	1.656	**[1.013;2.299]**		
	Pregnant (Ref: Lactating)	0.100	[-0.503;0.704]		
	Swollen (Ref: Pregnant)	1.556	**[0.943;2.169]**		
Female Rank	Low Rank (Ref: High Rank)	0.203	[-0.396;0.802]	2.812	0.245
	Medium Rank (Ref: High Rank)	-0.418	[-1.146;0.310]		
Female Parity	Parous (Ref: Nulliparous)	0.132	[-0.826;1.090]	0.073	0.787
Group Sex Ratio		-0.013	[-0.109;0.083]	0.071	0.789

Table 3: Male aggression and mating success (for sample sizes, see table S1). Probability of copulation of a heterosexual dyad during a female's swollen period in relation to the rate of male aggression received before that swollen period. Significant p-values and confidence intervals that did not cross zero appear in bold. The significance of each variable was assessed using chi-square tests (Chisq), while the significance of each level of a categorical variable was evaluated against a reference level (noted 'Ref') according to whether their confidence intervals (CI) overlap or not.

	Response variable: Mating during the swollen period (0/1)				
Fixed Factor	Level	Estimate	CI 95%	Chisq	P-value
Aggression Rate		1.591	**[0.115;3.067]**	4.466	**0.035**
Male Rank	Alpha (Ref: Non-alpha)	1.242	**[0.490;1.994]**	10.476	**0.001**
Female Rank	Low Rank (Ref: High Rank)	0.699	[-0.186;1.584]	2.664	0.264
	Medium Rank (Ref: High Rank)	0.715	[-0.645;2.075]		
Female Parity	Parous (Ref: Nulliparous)	-0.454	[-2.815;1.907]	0.142	0.706
Operational Sex Ratio		0.024	[-0.495;0.543]	0.008	0.928
Observation Time		0.548	**[0.221;0.875]**	10.807	**0.001**

Table 4: Male rank and aggression (for sample sizes, see table S1). Male aggression towards adult females in the months of the mating season in relation to male rank, age and sex ratio. Significant p-values and confidence intervals that did not cross zero appear in bold. The significance of each variable was assessed using chi-square tests (Chisq), while the significance of each level of a categorical variable was evaluated against a reference level (noted 'Ref') according to whether their confidence intervals (CI) overlap or not.

Response variable: Aggression during a month of the mating season					
Fixed Factor	Level	Estimate	CI 95%	Chisq	P-value
Male Rank	Alpha (Ref: Non-alpha)	0.610	**[0.050;1.171]**	4.552	**0.033**
Male age		0.050	[-0.067;0.167]	0.707	0.400
Operational Sex Ratio		0.315	[-0.005;0.634]	3.728	0.054