Imputation of 3D genome structure by genetic-epigenetic interaction modeling in mice

Lauren Kuffler, Daniel A. Skelly, Anne Czechanski, Haley J. Fortin, Steven C. Munger, Christopher L. Baker, Laura G. Reinholdt, Gregory W. Carter

The Jackson Laboratory, Bar Harbor, ME, USA

Corresponding author: Gregory W. Carter, gregory.carter@jax.org

Abstract

Gene expression is known to be affected by interactions between local genetic variation and DNA accessibility, with the latter organized into three-dimensional chromatin structures. Analyses of these interactions has previously been limited, obscuring their regulatory context, and the extent to which they occur throughout the genome. Here we undertake a genome-scale analysis of these interactions in a genetically diverse population to systematically identify global genetic-epigenetic interaction, and reveal constraints imposed by chromatin structure. We establish the extent and structure of genotype-by-epigenotype interaction using embryonic stem cells derived from Diversity Outbred mice. This mouse population segregates millions of variants from eight inbred founders, enabling precision genetic mapping with extensive genotypic and phenotypic diversity. With 176 samples profiled for genotype, gene expression, and open chromatin, we used regression modeling to infer genetic-epigenetic interactions on a genome-wide scale. Our results demonstrate that statistical interactions between genetic variants and chromatin accessibility are common throughout the genome. We found that these interactions occur within the local area of the affected gene, and that this locality corresponds to topologically associated domains (TADs). The likelihood of interaction was most strongly defined by the three-dimensional (3D) domain structure rather than linear DNA sequence. We show that stable 3D genome structure is an effective tool to guide searches for regulatory elements and, conversely, that regulatory elements in genetically diverse populations provide a means to infer 3D genome structure. We confirmed this finding with CTCF ChIP-seq that revealed strain-specific binding in the inbred founder mice. In stem cells, open chromatin participating in the most significant regression models demonstrated an enrichment for developmental genes and the TAD-forming CTCF binding complex, providing an opportunity for statistical inference of shifting TAD boundaries operating during early development. These findings provide evidence that genetic and epigenetic factors operate within the context of three-dimensional chromatin structure.

37   **Graphical Abstract**
38

39
40
41

Introduction

Local regulatory mechanisms within the genome and their interaction with chromatin structure give rise to subtle variations in gene expression. Yet the interacting effects that genetic and epigenetic factors produce on gene transcription are rarely studied at a genome-wide scale, leaving us without global information on a key step between the genetic code and the phenotype. Studies are generally limited to examination of individual regions or overlapping single-nucleotide polymorphisms (SNPs) and open chromatin peaks with limited investigation into how these regulatory elements combine to affect gene transcription [1,2]. Genetic analyses are a powerful approach that allow the study of these interactions. Conversely, phenotypic variation in genetically diverse populations is a result of both genetic and epigenetic factors operating in tandem. Understanding the scope and landscape of these interactions on a genome-wide scale is a vital step towards deciphering the genetic regulation of gene expression and, in turn, the mechanisms of non-coding variation on phenotypic outcomes.

Physical distance along a linear genome is a common metric for determining whether a putative regulatory element will affect a given gene's transcription. This has led to the concept of a "local area" in which most regulatory interactions take place. Many have used a 100 to 500 kb flanking window around a gene to encapsulate the landscape of local epigenetic effects [3–5]. This interval size is motivated more by convenience than rigorous evidence. Incidentally, the estimated scale of this local area is approximately equivalent to the size of a topologically associated domain (TAD).

TADs are chromatin loops with an average length of approximately 1.1 Mb that are joined via the architectural protein CTCF and the cohesin protein complex at known binding motifs [6,7]. TADs are believed to be mostly cell-type invariant and conserved across species [6,8,9]. The current hypothesis is that TADs provide physical boundaries for local regulatory function, but functional analysis of regulatory interactions within TADs has been limited by the lack of informative genetic variation in the studies that led to TAD discovery. Reverse genetic approaches like saturation mutagenesis can be used to introduce functionally informative genetic variation. But such screens are difficult in mammals, and even modern CRISPR-Cas9 based approaches are still limited in their scope and application outside of immortalized cell lines [10].

Samples from genetically diverse populations permit comprehensive study of complex genetic interaction and non-coding regulation on a genome-wide scale [11]. The Diversity Outbred mouse population (DO) is a heterogeneous stock composed of mice derived from eight founder strains, including three wild-derived strains [12]. The ongoing process of random outbreeding produces a mouse population segregating millions of precisely-mapped SNPs, with well-balanced allele frequencies, and increasingly small blocks of linkage disequilibrium that now approach the size of TADs [13,14]. Samples from the DO have been used to create extensive resources and data sets, including mouse embryonic stem cell (mESC) lines. A recent study used mESCs derived from generation 19 DO embryos, grown in the absence of ERK inhibition to study the gene

85 regulatory networks that stabilize pluripotent states *in vitro.* This revealed phenotypic variability
86 in features of ground state pluripotency [15].
87
Here we determine the extent to which genetic-epigenetic interactions between single
nucleotide polymorphisms (SNPs) and regions of open chromatin are present, and uncover the
biological basis of their distribution in three-dimensional (3D) genomic organization. Our results
show that genetic-epigenetic interactions were found across the genome and involve regulatory
elements that would not be identified with single-omics data. These interacting elements cluster
together within TADs bounded by previously identified active CTCF binding sites. We infer
chromatin structure from interaction data, analyze interaction contributions from the main and
interaction regression elements, identify potential regulatory functions underlying a set of
interactions, and correlate these interaction behaviors to CTCF binding differences in inbred
founder lines.
98

Methods

Sample production and initial processing

DO mESC production was performed by Predictive Biology. Cultures were grown with G3SK
inhibitor (1i medium). Bulk RNA and ATAC sequencing were performed and normalized as
previously described [15]. RNA-seq resulted in 6M-55M 2 × 75bp paired-end (PE) reads per
sample. ATAC-seq libraries were amplified for 9 cycles and purified with 1.7X AMPure beads.
Expression data was formatted and analyzed as log2-transformed transcripts per million (TPM).
ATAC-seq data was formatted as trimmed mean of M values (TMM) [16]. Genotyping was
performed by Giga Mouse Universal Genotyping Array (GigaMUGA), an Illumina Infinium array
of 143,000 SNP markers with a special focus on Diversity Outbred founder strains [17].
Aneuploidies were removed with the argyle R package, by identifying chromosome-level gene
expression differences [18]. QTL2 haplotype reconstruction, normalization and pseudoprobe
processing was carried out as previously described[15]. Samples with X0 genotypes were
removed, and the union of all samples with the required data types resulted in 176 samples.
115

116 Sample genotypes were estimated in reflection of linkage disequilibrium and haplotype
117 prediction confidence, employing a previously calculated grid of GigaMUGA pseudoprobes,
118 which were evenly spaced by genetic distance [19,20]. This produced a subset of 68,413 imputed
119 haplotype calls centered at specific loci which were most likely to match the actual haplotype in
120 each region (Supplementary Information).
121

122 *Regression modeling*

124 RNA-seq, ATAC-seq and haplotype data were processed into SQLite databases and fit to a
125 regression model using the stats::step() function in R. We modeled the expression of each gene
126 by linear regression with a genotype-by-ATAC non-additive interaction term:
127
128 $$y_i = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_1 x_2 + \varepsilon_i \qquad\qquad \text{(Equation 1)}$$
129
where $y_i =$RNA abundance (log$_2$TPM), $x_1 =$ genotype, and $x_2 =$ ATAC intensity (TMM).
Genotypes were taken from the haplotype estimations, and coded as 0 for reference, 1 for
heterozygote, and 2 for homozygote non-reference. Multi-allelic variants were not present in the
pseudoprobes and were therefore not accounted for in the method. Models that retained the
$\beta_3 x_1 x_2$ term by passing the default Akaike Information Criterion cutoff were deemed to be
interacting. These were used as the input for all further analyses.
136
This regression model was compared to a null model in which a genetic variant and open
chromatin element affect gene expression, but do so independently of each other ($\beta_3 = 0$ in
Equation 1). F statistics were calculated with the R function pf(). Given the possibility of
overfitting as the number of model terms increases and the high number of models tested (see
Results), a Bonferroni adjusted p value cutoff of p < 1x10^{-7} was selected (hereafter referred to
as "significant models"). Bonferroni corrections were calculated via stats::p.adjust().
143

144 *Topologically Association Domain, motif, and DNA binding analyses*

145
146 mESC TAD data was derived from previously published work [6]. in a liftover from mm9 to mm10
147 via UCSC genome browser [21]. Chromosome information for mm10 retrieved from the Integrative
148 Genomics Viewer [22].

149
150 DNA motif and binding site discovery was performed using the MEMEsuite tools MEME,
151 STREME, and Tomtom, all set to default parameters (See Supplemental Data) [23–26]. SNP data
152 were imputed from DO founder genomes referenced from release v3 (REL-1303-SNPs_Indels-
153 GRCm238) of the Mouse Genome Project through the Sanger Institute [27,28].

154
155 We downloaded the CTCF ChIP-seq in C57BL/6 Bruce4 and 129/Ola E14TG2a.4 mESCs from
156 the ENCODE portal[29] with the following identifiers: ENCSR000CCB and ENCSR362VNF [30].
157 These were also used to estimate the locations of Smad3 activity. CTCF and SMAD3 binding
158 motifs were retrieved from HOCOMOCO [31]. with the identifiers CTCF_MOUSE.H11MO.0.A and
159 SMAD3_MOUSE.H11MO.0.B. These were overlapped with our ATAC-seq dataset via FIMO [27].
160

161 *CTCF ChIP-seq*

162
163 We performed CTCF ChIP-seq following previously established protocols [32]. Three cell lines
164 were run for four isogenic mESC strains, C57BL/6J, CAST/EiJ, PWK/Ph, and WSB/EiJ.

165 Cultures were grown in 1i medium. One sequencing run failed, resulting in the loss of one PWK
166 and one WSB sample.
167
168 FastQC, Bowtie alignment and g2gtools liftover from imputed strain genomes to BL/6J were
169 performed using a modification of a now publicly available ATAC-seq pipeline, accessed in
170 development 15 December 2020[33]. Mitochondrial read filtering steps were removed. Results
171 were filtered to those ChIP-seq binding locations that appeared in two or more isogenic strains.
172
173 All study data and scripts are available at Synapse (https://doi.org/10.7303/syn26534443).

Results

Genetic-Epigenetic Interactions are Pervasive and enriched within TAD boundaries

Although there are numerous reports detailing examples of genetic variants and chromatin state
altering gene expression, the non-additive interactions between these features have yet to be
systematically explored. We aimed to determine the extent to which genetic-epigenetic
regulatory interactions occur on a systemic level, and whether these interactions exhibit any
bias in their location within the genome. To explore these interactions, we utilized paired
transcriptome and chromatin accessibility profiling of Diversity Outbred mESCs [15]. These
samples incorporate genetic backgrounds from eight well-characterized founder strains from
three *M. musculus* subspecies, randomly outbred for 19 generations. Thus, the resultant data
constitute a unique resource for addressing questions of genetic-epigenetic regulatory
interactions.
187
To search for non-additive interaction effects on a global scale, we developed a method of
analysis that could be broadly applied to, and could permit discrimination between, co-
occurrence and interaction effects. To that end, we combined regression model fitting (see
Methods, Equation 1) and model selection via information criteria in order to test whether gene
expression was affected by a genetic variant, an open chromatin region, both independently of
each other, or both working in concert.
194
195 To account for the resolution limits on genetic variant effects imposed by linkage disequilibrium,
196 we subsetted our genotype data to 68,413 high confidence haplotype calls evenly spaced by
197 genetic distance to reflect this resolution limit (Methods). We collated these with 102,173 ATAC-
198 seq peaks and 13,631 genes with RNA-seq coverage within our sample set.
199
With 95 trillion possible interactions genome-wide, testing had to be focused to an informative
and computationally feasible subset. To assess the significance and extent of non-additive
interaction models and identify target locations for more focused analyses, a subset of randomly
selected trios of intrachromosomal genes, genetic variants, and ATAC-seq peaks were tested
with the interaction model. A total of 102,104 ATAC-seq peaks, 68,413 variants, and 13,631
genes were used to model 39,021,625 combinations. This resulted in 27,509 significant models

(Bonferroni adjusted p < 1x10^{-7}), or 0.07% of all models, involving 19,145 ATAC-seq peaks,
14,476 variants, and 1,286 genes (18.75%, 21.16%, and 8.52% of the total input respectively).
Complete results are tallied in Tables S1 and S2.
209
If interactions were detected randomly, we would expect a null distribution with no enrichment of
low p-value models. Instead, the number of significant models was four orders of magnitude
greater than expected, thus indicating the presence of true associations (Fig. 1A). The locations
of significant associations indicated non-additive interactions were much more likely to occur
within the linkage disequilibrium block that contains the gene (Fig. 1B), matching prior work that
the majority of detectable genetic effects originate in local regulatory regions [34–36].
Approximately half (49%) of all significant interacting models included a genetic variant or ATAC
peak within 4 Mb of the gene they affected. Within this range, we found that less than 10% of
variants and ATAC peaks clustered closer to each other than they did to the gene they affected,
indicating that co-localization may not the driving factor in many interacting models (Table S3).
220
As a result of these findings, we considered potential chromatin features that might influence the
local area for genetic-epigenetic interactions. We hypothesized that TADs may play a role in
determining the prevalence of non-additive interactions, as they are a 3D genome structure
known to enable enhancer access to local genes. Therefore, we generated a second, TAD-
focused subset of regression models, testing all potential combinations between each gene and
every genetic variant and ATAC peak that fall within one TAD of its transcription start site in
either direction. TAD boundary locations were retrieved from publicly deposited Hi-C based
analyses of B6 embryonic stem cells [6].
229
230 Models with at least one interacting element within the same TAD as the gene were 3.7 times
231 more likely to reach our threshold, compared to models that did not (3.3% versus 0.9%). This
232 constituted 3,836 genetic variants and 3,725 ATAC-seq peaks affecting the expression of 896
233 genes in our initial exploratory subset, and 11,904 SNPs, 25,961 ATAC-seq peaks, and 1,099
234 genes in our equivalently-sized TAD-centric subset (Table S1-S2).
235
ATAC-seq peaks involved in non-additive interactions favored results between CTCF binding
sites associated with TAD formation (Fig. S1A). Regression models that did not identify
interaction involvement also show this local, TAD-bound organization of ATAC-seq peaks (Fig.
S1B). Interacting regulatory elements as defined by our model are thus clustered within the
same TAD as the gene they affect. This provides a possible structural explanation for previously
observed local regulatory ranges.
242
243 *Interacting elements in genetically diverse samples escape conventional discovery*
244 *methods*
245
246 We next evaluated whether genetic-epigenetic, non-additive interactions provide information
247 that genotyping or ATAC-seq alone cannot. To this end, we compared interacting models, found

248 to represent 32.29% of significant intra-TAD models (Fig. 1C), with models associated with
249 genetic variants or ATAC-seq peaks without interactions. We broke down the overlap between
250 intra-TAD regression model results by genetic variants and ATAC-seq peaks separately.
251
252 Previous studies have found correlation between local open chromatin and gene expression,
253 including within this DO mESC population [15]. Mediation analysis has indicated more complex
254 co-regulatory relationships are also extant, but their relationships have not been
255 comprehensively studied.
256
Each genetic or epigenetic regulatory feature input to our regression model can potentially
influence multiple genes, alongside multiple other regulators. Thus, each feature could be
engaged in many different regulatory relationships with its local genes, some of which may or
may not be detectable with correlation or mediation analyses. We wished to determine how
many regulatory elements were exclusively found in interacting regression models, and would
thus escape notice with conventional analyses.
263
We found that 26.35% of genetic variants are shared between non-additive interacting models
and additive models, and 23.12% are shared with models that contain no ATAC-seq peak
contribution. In contrast, few models contain *only* an ATAC-seq peak contribution, and 41.81%
of ATAC-seq peaks are exclusive to interacting models (Fig. 1D). This is also in contrast to
results from our randomly sampled gene-ATAC-genotype trios, which show a preference for
genotype-only models (Fig. S1C-D).
270
These results suggest that although genetic variants are the primary driver of variation in
expression, non-additive interactions with chromatin states further reveal the origins of
expression variation. Our findings also suggest that genetic variation and open chromatin data
cannot be used independently to capture these regulatory features. Conversely, this suggests
that ATAC-seq data alone is a less effective predictor of gene expression in a genetically
diverse population.
277

279
Figure 1. Interacting and additive models are abundant and favor local genes. **A.** Density of Bonferroni
Adjusted p-value distribution in randomly sampled intrachromosomal models. **B.** Density of non-additive
interacting models where adj. $p < 1\times10^{-7}$, by minimum distance between regulatory elements. **C.**
Distribution of model term retention across intra-TAD models. **D & E.** Euler plots of ATAC-seq peaks and
genetic variants, dividing them by their participation in single-term, additive, and non-additive interacting
models. ATAC-seq peaks and genetic variants involved in interacting models are highlighted, as they are
investigated further in this paper.

TAD boundaries limit genetic-epigenetic interactions

The specific preference for non-additive interactions within active CTCF binding sites warranted
further study. We sought to determine if active CTCF binding sites provided an appreciable
boundary to interactions (Fig. 2A), similar to reported segregation of enhancer elements [37] . We
carried out regression analyses across the genome for all possible models involving each gene-
genotype-ATAC combination within the gene's TAD and nearest flanking TADs or intra-TAD
regions. Although resolution of causal variant location limited by linkage disequilibrium, ATAC-
seq peaks that interact with variants could be confidently localized in relation to TAD
boundaries.

We found that TADs contained more ATAC peaks that interact with local genetic variants to
affect expression of a resident gene (Fig. 2B). With a window of 200 kb on either side of an
active CTCF boundary, we found 52,511 ATAC-seq peaks sharing a TAD with their interacting
gene, versus 24,229 peaks in different TADs or inter-TAD regions This constitutes a significant
enrichment of intra-TAD gene-peak interactions over expectations based on overall ATAC-seq
peak density (an odds ratio of 1.48, null expectation $p < 2.2 \times 10^{-16}$) (Fig. S2). We also identified
29 individual ATAC-seq peaks that were involved in more than 200 non-additive interactions
each, which are distinctly visible within the aggregate view (Fig. S3). These were exclusively
affecting genes within the same TAD as the interacting ATAC-seq peak, which suggests that
our results are not due to higher density of open chromatin within TAD structures. This is
consistent with previous findings that the effects of enhancers and other regulatory elements are
constrained by TADs [38].

Revising constraints on local regulator area using genetic data

Definitions of local regulatory area are used to establish the scope of analyses.
Many cell types and animal models do not have publicly available Hi-C or CTCF ChIA-PET
datasets, further limiting many researchers that may want to study genetic-epigenetic non-
additive interactions. Our data supports that TADs may act as a biologically-defined boundary.
Therefore, we wanted to determine how far the TAD-defined local area was likely to extend from
any given gene. We found that to capture 95% of inter-TAD genetic-epigenetic interactions at
adjusted $p < 1 \times 10^{-7}$, a window of 2,074,778 bp upstream and 2,575,581 bp downstream of the
gene transcription start site (TSS) was required (Fig. 2C). The density of results using linear
DNA sequence does not necessarily experience a linear drop-off over this distance, due to
variable TAD lengths (Fig. 3E-F).

Figure 2. ATAC-seq peaks that interact with genetic variants generally reside within the affected gene's TAD. **A.** Schematic of a TAD loop, including gene (purple) and density of interacting model elements (red). Loop interior is in blue, exterior DNA is gray, and CTCF binding sites are in yellow. **B.** Location of interacting ATAC-seq peaks relative to TAD boundary location, merged across all genes. TAD interior denotes the TAD in which the dependent gene was found. **C.** Interacting ATAC-seq peaks by distance from associated gene TSS. Local area cutoffs of 100 kb and 500 kb flanking regions are marked.

We next considered the location of these interacting elements with regards to genes and known regulatory elements. ATAC-seq peaks involved in non-additive genetic-epigenetic interactions were analyzed by overlap with mm10 NCBI RefSeq genes and selected Enhancer Atlas 2 datasets, as well as FANTOM5's list of mm10 promoters[39]. (Fig. S4A-B). 65.37% of peaks on each chromosome fell within gene bodies. Normalized for length, exons were especially enriched, with 34.41% of peaks falling within exons, half of which fell within the first (or only) exon. 24.06% of peaks overlap with promoters. 6.09% of peaks fall within enhancers curated for ESC R, R1, KH2 embryonic stem cell lines. There was no inverse relationship between gene

body vs. enhancer peak percentage (Fig. S4C), indicating that this was not simply reflecting the
relative density of genes within various TADs across the genome.

Figure 3. TADs provide context for interactions and increase interaction search efficacy. **A.** Counts of
intra-TAD ATAC-seq peaks involved in all non-additive interactive models, centered on the TSS of the
gene affected by the genotype-ATAC interaction. Coordinates transformed to a standard scale. **B.**
Example TAD, displaying interacting ATAC peak density and gene locations. Peak relevance generally
decays relative to intra-TAD distance rather than linear chromosomal distance. **C-F.** A comparison
between linear sequence-based and TAD-limited search methods for interacting ATAC-seq peaks. C and
D compare percentage of significantly interacting ATAC-seq peaks at each gene-relative locus. E and F
compare density of ATAC-seq peaks at each locus. TAD-based search shows a higher density of
interactions and places limits on search distance due to testing only TAD-internal ATAC-seq peaks.
356

357 *Density of interactions between genomic elements is defined by 3D context*
358
359 If TADs act as a constraint to local interaction, then their 3D looping structure should be
360 reflected in the regulatory patterns of genes found within them. At the most basic level of TAD
361 structural organization, the CTCF binding site brings distant areas of chromatin into close
362 physical association. We analyzed genome-wide distribution of genotype-interacting ATAC
363 peaks relative to genes within their respective TADs.
364
This revealed that the gene-centric increase in regulatory interaction density responded to the
3D context of the TAD loop, crossing CTCF boundaries irrespective of linear DNA sequence.
This creates a distribution of non-additive interactions centered at the gene promoter that
reaches its minimum halfway around the TAD loop from the gene (Fig. 3A). In fact, we found
local regulatory regions that displayed organization that spanned the TAD boundary sites,
creating interaction density gradients that appear non-linear when viewed without TAD data
(Fig. 3B).
372
We sought to determine whether gene-centered, TAD-internal searches for interacting elements
were potentially a more efficient method for searching for non-additive interactions, when
compared to local area searches along the linear DNA sequence. We compared the two
methods, running a pair of genome-wide, gene-centric analyses out to the limit of our TAD-
internal search (+/- ~2.5 Mb). We calculated the percentage of ATAC-seq peaks within this
region that participated in significant interactions. (Fig 3C-D). The overall density of ATAC-seq
peaks within these regions was also compared (Fig. E-F).
380
381 We found that TAD-based searches produce consistently higher percentages of interacting
382 ATAC-seq peaks across the analysis, and higher ATAC-seq peak density out to +/- ~1Mb. TAD-
383 based searches experience a progressive falloff in ATAC-seq peak density when compared to
384 linear sequence-based searches. This is largely due to dropout of smaller TADs from the
385 analysis. This results in more variable interaction percentages past ~1Mb from the gene.
386
Linear search results show a consistently lower percentage of interacting ATAC-seq peaks, and
a lower density of ATAC-seq peaks per locus. This was unexpected, as the immediate vicinity of
the gene was expected to contain a similar density and non-additive interaction potential of

ATAC-seq peaks when compared to the TAD-based search method. We therefore hypothesized
that TAD boundaries in close proximity to some genes might create this discrepancy. We found
that while the majority of genes were randomly distributed within their TADs, 12,437 or
24,2124.31% of genes were found to have a TAD boundary close upstream to their transcription
start site (Fig. S4D).
395
Interestingly, there is a depletion in the non-additive interaction rate of ATAC-seq peaks flanking
the gene (Fig. 3D), which is not noted in raw interaction counts (Fig 3A). This may be due to the
necessary presence of open chromatin at the promoter and the gene body during priming and
transcription regardless of interaction. There are also genes which are concomitantly active
within our samples, and do not show variation in expression, thus experiencing activity without
detectable regulatory interaction.
402
Our results indicate that the local area for non-additive genetic-epigenetic interaction is not only
constrained by TADs, but also shaped according to the overall 3D genome structure of the TAD
loop. Analytical methods which reflect this are more efficient at discovering interacting
regulatory elements.

Genetic-epigenetic interactions influence gene expression more than epigenetic factors
alone
The coefficients estimated by our regression model quantify the effects of the genetic and
epigenetic features on the expression of each gene (Equation 1). We postulated that our ATAC-
seq, genotypic variants, and non-additive interaction effects would display patterns that
corresponded to positive and negative regulatory roles with varying strengths.
413
In previous analyses of SNP-SNP interactions, the importance of relative magnitude of
interaction effects has been a subject of debate, as single effect terms ("main effects") are
normally of greater magnitude than interaction effects [40,41]. However, in the context of non-
additive genetic-epigenetic interactions we observed interaction effects of greater absolute
value than ATAC-seq effects 61.24% of the time, greater than genotypic variant coefficients
11.42% of the time, and greater than both main effects 8.52% of the time.
420
421 This suggests that ATAC-seq peaks are generally a modifier of underlying genetics, so the
422 majority of ATAC-seq peak effects are smaller than interaction effects. This was supported by
423 the relatively low contribution of ATAC-seq peaks to single-effect regression models (Fig 1D).
424 Their quantitative variation produces relatively subtle effects. This stands in comparison to the
425 binary presence or absence of variants, which also directly capture regulatory sequence.
426
The directionality of effects on gene expression specifies the positive or negative influences
from genetic variants, ATAC peaks, and their combinations. Positive and negative effects
represent correlation or anti-correlation of variant presence or ATAC-seq score with gene
expression. When all effects have the same sign (positive or negative), the total effect is
synergistic and the pairwise combination of non-reference genotype and ATAC peak alter

expression beyond the sum of the two. When the interaction effect has the opposite sign of the
main effects, this indicates redundancy or interference, based on the idea that redundant factors
create an "or" logic that yields a combined result that is less than the sum-based expectation.
Alternatively, mixed main effects with interactions can also signify a suppression of one
outcome in favor of the other, in which the sign of the interaction term serves to move the
additive expectation nearer to one of the two marginal outcomes.
438
We found that models indicating redundancy or interference were the most common overall,
totaling 40.53% 39.54% of all models (Fig. S5A). Synergistic effects were found in 17.02% of all
models. These two observations, suggest that a greater proportion of regulatory non-additive
interactions attenuate gene expression, rather than strengthening it. While ATAC-seq peaks are
often correlated with increased gene expression [42], we were surprised to find that an increase in
ATAC-seq signal had a negative effect on gene expression in 40.76% of models (Fig. S5A).
Due to the high proportion of ATAC-seq peaks found within gene bodies and the association
between open chromatin and gene transcription, this result warranted further investigation.
447

Motif Enrichment Analysis reveals CTCF complex participation in genetic-epigenetic
interactions
We looked next for potential binding sites or functional motifs underlying our results to provide
clues as to the mechanistic underpinnings of model effects. We were especially interested in the
subset of ATAC-seq peaks with negative effects on gene transcription, as these results were
counter to our expectations. We hypothesized that these areas of open chromatin might expose
binding sites of repressive regulatory factors.
455
456 To test this hypothesis, we tested for DNA motif enrichment analysis using MEMEsuite (Fig.
457 4A). We selected the subset of ATAC-seq peaks involved in 10 or more significant non-additive
458 interactions, at least 50% of which have negative ATAC-seq effects (negative effectors). This
459 subset was compared versus all ATAC peaks and against shuffled control sequences via
460 STREME analysis, which finds enriched ungapped motifs in provided sequences.
461
Results showed the negative effector subset was enriched for 49 motifs (Supplemental File 1),
including the CTCF binding site ($p < 2.0 \times 10^{-14}$) and SMAD3 binding site ($p < 1.5 \times 10^{-8}$). These
motifs were present, but less significantly enriched in positive effectors, or in all significant
ATAC-seq peaks. This suggested that a portion of negative ATAC-seq effects can be
functionally explained by altered behavior of CTCF binding sites carried by specific ancestries in
the DO. To complement this analysis, we also quantified the CTCF motif occupancy in negative
effectors versus other ATAC-seq peaks. FIMO motif scanning showed that 53.3% of top
negative ATAC-seq sequences had at least one CTCF binding motif in them, compared to
35.3% in ATAC-seq peaks with positive effectors on gene transcription (See deposited data).
471
We next examined whether these peak locations contain SNPs that might alter CTCF binding
potential. Imputing from the founder genomes of the DO population, we analyzed the locations

of SNPs in CTCF binding motifs associated with regression models versus all CTCF motifs. Out
of CTCF motifs within this negative subset, 96.1% of these were found to contain SNPs from the
non-reference DO founder strains, versus 18.0% for positive effect-associated CTCF motifs (p <
2.2×10^{-16}). Similar results were found for Smad3. We also found that across all motifs, the
density of SNPs favored the start and end of the binding sequence (Fig. 4B). However, in motifs
associated with regression models, the density of SNPs was approximately equal across all
bases (Fig. 4C). These bases have previously been identified as having high protein-DNA
binding energies with the canonical sequence [42,43].
482
Most genomic sequences matching the CTCF binding motif are not known to be bound,
according to ChIP-seq experiments [44]. We wanted to determine the overlap between CTCF
binding sites found in our data and known active binding sites in mESCs. Comparing to
available ENCODE CTCF ChIP-seq in C57BL/6 Bruce4 and 129/Ola E14TG2a.4 mESCs [30,45],
we found 2.04% and 1.87% of these overlapped with negative ATAC effect CTCF binding sites
that contained SNPs, versus 17.45% and 16.20% overlapping with positive effect CTCF binding
sites that contained SNPs. Expanding the scope to attempt to find flanking Smad3 regions
netted consistently low results, with a 1 kb flanking window returning between 1.20% and 0.41%
for Bruce4 and E14TG2a.4 respectively. These findings show that the majority of CTCF binding
sites found within our significant models are not captured in previous analyses of ESCs two
different *M. musculus musculus* strains.
494

495
Figure 4. Motif analysis identifies differences in interacting CTCF binding motifs. **A.** A schematic of our
motif analysis through MEMEsuite. FASTA files derived from interacting ATAC-seq peaks are used to
identify enriched motifs, identify protein binding sequences, and locate the sequences within the ATAC-
seq peaks. **B-C.** Binding sites found within significant motifs are less protected from genetic variation.
SNP counts are shown at each locus in the CTCF binding sequence, comparing motifs within interacting
ATAC-seq peaks versus all CTCF binding sites.

Putative developmental regulator Platr2 is regulated by multiple redundant elements

To provide an example of our analytical method and probe a gene previously proposed to be
important in stem cell development, we examined *Platr2* and its TAD. *Platr2*'s TAD contains a
high concentration of confidently called regression models in our analysis (Fig 3B), with 1,031
models of non-additive genotype-ATAC interaction reaching our significance cutoff for resident
genes, of which 179 were models of *Platr2* expression. It also contains 22 haplotype markers,
allowing a certain level of model variant localization. Previous studies have found a group of
genes regulated in *trans* by expression quantitative trait loci (eQTLs) mapped to *Platr2* [15]. These

target genes are associated with embryonic ectoderm, indicating *Platr2* may act as a regulator
of stem cell state. These factors made it a target of interest for further exploration.
512
When the direction of effects for *Platr2* were analyzed, the distribution showed a shift toward
models where ATAC and genotype effects agreed with each other, but not with the interaction
term (Fig. S5B). As discussed above, this potentially indicates functional redundancy between
haplotype and chromatin openness at these sites. Motif enrichment analysis of interacting
ATAC-seq peaks identified a sequence at 16 sites which contain Smad3 binding motifs, and
another sequence at 15 sites that contain Ctcf binding motifs (Fig. S5 C-D, Supplemental File
2), which may suggest modulation of CTCF binding strength. These results suggest that *Platr2*
may have differential regulation patterns governed by changes in TAD formation.
521

CTCF binding in inbred mESCs validates strain specific effects
CTCF binding to DNA is associated with open chromatin around the binding site, TAD
formation, and regulation of local gene expression [32,46–48]. Genetic variation within CTCF binding
sites can alter binding potential [42,43]. We hypothesized that our regression models indicated
areas of strain-specific CTCF binding due to polymorphisms in the DO haplotypes. To test this,
we performed CTCF ChIP-seq on mESCs derived from four of the eight DO founder strains,
including representatives from the three subspecies contributing to the DO population.
C57BL/6J was also included as the standard reference.
530
531 Sample ChIP-seq binding intensities clustered by strain in PCA, and were separated by
532 subspecies as expected (Fig. S6). Consensus peaks were identified from replicates of each
533 strain. Across all samples, 65,541 ChIP-seq peaks passed our significance threshold (found in
534 at least two samples in at least one strain), or 68.21% of results. 90.65% of these overlapped
535 with the publicly available TAD dataset we had employed for our previous analyses.
536
537 Our DO data suggested to us that there are genetically determined differential CTCF binding
538 sites within that population. If that is the case, then we would expect to see strain-specific
539 differences in CTCF peak occupancy. Consensus peaks were identified from replicates of each
540 strain. Fifty-two percent of CTCF ChIP-seq peaks were shared across all four strains, with the
541 remaining 48% found in three or fewer strains (Fig. 3.5A).
542
We found 23,887 of our significant CTCF ChIP-seq peaks these overlapped with CTCF binding
sites that had been previously identified, or 36.45% overlap with significant results. We also
found that 16,951 (25.86%) significant results overlap with the CTCF ChIP-seq data previously
retrieved from ENCODE, and among those, 7,880 (12.02%) that overlap with significant,
interacting ATAC-seq peaks in our DO mESC data. About 80% of these (~6,300) are found in
all four strains.
549
550 To further explore strain-specific differences in CTCF binding, we examined reported binding
551 intensity. Our CTCF ChIP-seq was performed as a bulk analysis, so binding intensity can show

552 the probability of finding a protein bound at a locus within a sample. If CTCF binding is disrupted
553 by strain-specific genetic factors, then its binding intensity will vary.
554
Our analysis found a range of binding intensities at individual loci, with 13% having a variance in
fold enrichment greater than 10 (Fig. 5B). This gave us confidence that there was truly
differential binding of CTCF in our samples. Given that we previously saw an enrichment of
CTCF binding sites under significantly interacting ATAC-seq peaks, this lends weight to our
hypothesis that we were seeing evidence of 3D chromatin structural differences in our
genetically diverse mESCs.
561
562 Fifty-two percent of CTCF ChIP-seq peaks were shared across all four strains, with the
563 remaining 48% found in three or fewer strains (Fig. 5A). A range of binding intensities was also
564 found at individual loci, with 13% having a variance in fold enrichment greater than 10. (Fig. 5B).
565

566
567 **Figure 5.** CTCF ChIP-seq analysis shows predictable strain-specific differences in binding
568 intensity. A. Percentage of ChIP-seq peaks in surveyed strains. B. Variance (log10) in binding
569 intensity fold enrichment for all ChIP-seq peaks. C. Percentage of significance in association
570 between DO genotype at CTCF peaks and CTCF binding intensity on inbred ChIP-seq samples,
571 in various subsets.

Non-additive interactions are predictive of CTCF binding patterns

With our ChIP-seq data indicating the presence of strain-specific CTCF binding, we
hypothesized this differential binding could be predicted from our regression models. We
anticipated that inbred founder strain CTCF ChIP-seq would have more genotype effects at
binding sites with open chromatin in non-additive interactions with local genotype, as opposed
to binding sites without interactions.

Both interacting and non-interacting areas contain genetic variation and ATAC-seq peaks, but
our models indicated interacting areas had more effects on local gene expression and genetic
variability in CTCF binding sites. More specifically, we expected negative effector ATAC-seq
peaks to have the greatest predictive power, as we believed their negative effects on local gene
expression arise from strain-specific differences in CTCF binding and loop formation, whereas
positive effects on gene expression may come from non-localized genetic variants in other
binding sites.

To test these hypotheses, we retrieved CTCF ChIP-seq locations and identified biallelic SNPs
found in the four inbred DO founder strains we used for our ChIP-seq experiments. The
correlation of SNP genotype to CTCF binding site occupancy was assayed using a paired
sample Student's t-test (Fig. 5C).

The proportion of results below $p < 0.05$ in all CTCF ChIP-seq peaks was 16%, establishing a
baseline of predictive power for SNPs genome-wide. Subsetting to those CTCF binding sites
found in ATAC-seq peaks increased the proportion of significant correlation to 31%. This was
expected, as open chromatin in the vicinity of CTCF binding sites is associated with binding site
occupancy [32,47].

To test whether our theory that non-additive interacting regression models held greater
predictive power of CTCF binding intensity, we subset CTCF binding sites in ATAC-seq peaks
with significant effects on local gene expression. Binding sites associated with non-additive
interacting models had 37% correlation. This outperformed additive models, which had 32%
correlation.

ATAC-seq peaks can be associated with significant effects on multiple genes, potentially in
combination with multiple genetic variants, resulting in some ATAC-seq peaks associating with
multiple significant regression. Interestingly, we found that among ATAC-seq peaks over CTCF
binding sites, those associated with additive models were a subset of ATAC-seq peaks with
non-additive interacting models. This means that any ATAC-seq peaks that had effects on local
gene expression and were localized to CTCF binding sites *always* had interaction effects with
the local genotype, and these models are more predictive of CTCF binding intensity.

These results matched our expectations. ATAC-seq peaks co-localized with a functional
polymorphism affecting a CTCF binding site would be more likely to affect gene expression in a
non-additive fashion, as the polymorphism would only affect CTCF binding based on chromatin
state at the binding site or any nearby priming factors. This is in contrast to additive models,
where genetic effects and ATAC-seq effects are predicted to be independent of each other and
are thus less likely to be co-localized.
619

620   Subsetting those ATAC-seq peaks that were negative effectors in interacting regression models
621   produced a 39% correlation. This outperformed ATAC-seq peaks with a positive effect, which
622   had 32% correlation. This was in line with our previously stated predictions and suggested that
623   non-additive interactions can be used to evaluate and predict local 3D chromatin structure.
624

Finally, we wanted to determine whether the areas where we identified the most DO regression
models supported our hypothesis, or if a larger amount of activity would result in a leveling
effect on DO genotype's predictive power on CTCF binding. We subset the results to those
regions associated with a greater than average number of regression models. Results showed
that the gap between negative and positive effector regions widened in these regions to 44%
versus 27% correlation. This indicates that our interacting regression model can provide
detailed and biologically data in regions of dense genetic-epigenetic interaction, and further
emphasizes our previous results.
633

Discussion

Integrating analysis of multi-omics data in a genetically diverse population allows for greater
specificity and modeling of complex regulatory interactions. Populations that segregate natural
genetic variation provide dense, randomized perturbations from which gene expression variance
can be modeled. This contrasts with experimental designs that rely on a limited number of
engineered perturbations in isogenic cells or animals. By collecting genetic, epigenetic, and
transcriptomic data from the same cellular panel, models can be inferred without confounding
factors such as different experimental protocols and environments. Our approach to integrating
genetic and functional genomics data from 176 Diversity Outbred mouse embryonic stem cell
lines allowed us to systematically probe how genetic and epigenetic variation interact to jointly
influence local gene expression.
646

We integrated ATAC-seq peaks and genotypic variants via an interactive regression model of
their effects on gene expression to determine how often these factors are independent of each
other, and how often they interact. From this, we were able to infer the wide-scale presence of
local interactions between these regulatory mechanisms. This method is well-suited for outbred
populations, which are not compatible with methods used for analyzing inbred cohorts, such as
differential expression or variance between samples. While similar interactions have been
observed in isolated contexts and in co-localized genetic variants and ATAC-seq peaks [38,49], this

marks the first time that the phenomenon has been observed genome-wide, with the genetic
mapping resolution provided by outbred, heterozygous samples.
656
From our analysis of genetic-epigenetic interactions we have put forth several key findings
regarding the interactivity, structure, and function in the regulation of gene expression. We
discovered that patterns of genetic-epigenetic interaction reflect the structure of topologically
associating domains. Our model inferred the presence of frequent interactions between
genotypic variation and open chromatin, with a strong preference for coordinates internal to
known TADs in mESCs. We demonstrated this inference in several ways, including the
clustering of highly interacting areas of open chromatin within TADs, and clustering of
interactions based on inter-TAD distance rather than linear DNA sequence.
665
Interacting ATAC-seq peaks were found to cluster within gene bodies, promoters, and
annotated enhancers. These findings confirm that TADs generally define the local area for
interactions. While open chromatin density is higher within TADs, we found that open chromatin
density is less important for interactivity than TAD boundaries are. Furthermore, linear proximity
cannot be used to identify association between genes and regulatory features. Genetic variants
or areas of open chromatin that are segregated by a TAD boundary may be incapable of
affecting expression of a nearby gene. Conversely, a distant genetic variant or ATAC peak may
be placed near a gene within a TAD loop.
674
Interaction effects were further classified based on the magnitude and direction of their effects
on gene expression. These models contained an unexpectedly high proportion of increased
ATAC-seq peak effects associated with reduced gene expression. Upon further investigation,
these were found to be enriched for CTCF binding sites, which were further enriched for genetic
variation in Diversity Outbred mice, particularly in the core sequence of CTCF binding sites (Fig.
4B-C). This indicates that these CTCF binding sites may be differentially bound in different
samples, resulting in downregulation of transcription, either by cutting off access to nearby
enhancers or by abolishing TAD structure.
683
We further analyzed the interplay between direction of effect for genetic variants, ATAC-seq
peaks, and their interactive components (as in Fig S4A), allowing us to make hypotheses about
the functional significance of various interactions. Models where all effects are positive or
negative suggest the open chromatin region and genetic variant enhance each other's
effectiveness in increasing or decreasing gene expression. These synergistic effects are
indicative of two regulatory factors working together to produce a greater change in gene
expression, beyond what either could produce independently. Functional redundancy or
interference can be inferred from models where the non-reference genotype and ATAC-seq
peak have a positive effect on gene expression and the interaction effect is negative, or vice
versa. Redundancy is rather common in our genetic-epigenetic interactions, while synergistic
effects are relatively rare. This appears to align with previous analyses of purely genetic
interactions in other mouse crosses [6,40]. Other models are more cryptic, but contain variation in
beta coefficients associated with these models (Fig. S6), suggesting that functional subtypes
that can be investigated within each group.

698
Our approach can be directly extended to specific gene targets. Mediation analysis in the DO by
Skelly et al. revealed several mESC genes that act as mediators of downstream gene networks
[15,50]. Our analysis identified several genetic-epigenetic interactions in the area of one mediator
gene, *Platr2*, and provided a targeted list of SNPs and ATAC-seq peaks that may influence the
gene's expression (Fig. 3B). ATAC-seq peaks were enriched for Smad3 binding sites, a
component of the CTCF binding complex. This provides further regulatory information and
potential targets for experimental manipulation.
706
707 Through DNA motif analysis, we identified distal regulatory activity via our interaction modeling.
708 This permits functional analysis of protein factors on gene activity that is otherwise undetected
709 by this method, such as regulatory proteins with no local QTL. This means that while interaction
710 models may achieve the best resolution in areas of low linkage disequilibrium, they can still be
711 used to identify and infer regulatory action of key conserved proteins in *trans*.
712
713 Genetically diverse populations in mice still contain significant linkage disequilibrium blocks
714 which affect genetic resolution. This limited our ability to identify the causal SNPs for the
715 purposes of this paper, but as demonstrated with our analysis of *Platr2*'s local regulatory area,
716 the variable genetic distance across the DO genome sometimes permits us sufficient resolution
717 to analyze whether interacting elements are colocalized or not.
718
719 Our CTCF ChIP-seq experiments identified strain-specific differences in CTCF binding, which
720 were most predictably affected by genetic variation in regions tied to non-additive interacting
721 regression models in DO mESCs. We also found that previously identified CTCF binding sites in
722 negative effector ATAC-seq peaks are more likely to contain SNPs that can predictably CTCF
723 binding potential.
724
725
Our results have several implications for genetic analysis of gene expression in genetically
diverse populations. Experiments performed in cell cultures or isogenic models often fail to
produce replicable results in other tissues or human trials, Genetic-epigenetic interactions may
underpin some of these failures, particularly those that have initially shown strong genotype-
dependent effects. Thus, it is important to consider generating datasets with a combination of
genetic mapping, gene expression, and appropriate epigenetic data, either on a local or
genome-wide scale. At present, publicly available data that matches these criteria is not
available. As research expands towards greater coverage in wild populations and humans,
accessibility is likely to increase. Furthermore, ATAC-seq could be substituted for other
experimental datasets, such as H3K4me3 ChIP-seq.
736
In the future, our study would benefit from the generation of more precise mapping of active
CTCF binding sites, across more genetic backgrounds. Our CTCF ChIP-seq experiments
expanded on previous findings that mouse strains have different strengths in CTCF binding [51].
Our results indicated that almost half of CTCF binding is strain-specific in mESCs, with imputed
links to genetic variants in the Diversity Outbred mouse population. We also found a link

between CTCF binding site occupancy and our imputed genetic-epigenetic interactions, which
could indicate 3D structural variants that have effects on local gene expression.
744
To confirm this, more experiments are needed. ChIP-seq can identify where CTCF is binding,
but not whether it has formed a TAD binding complex. Chromatin conformation experiments like
Hi-C and ChIA-PET in the DO founder strains would permit us to directly examine genome
structure. This could determine whether there are interactions that were obscured in some
samples by the blocking effect of a TAD boundary, and whether there are some TADs that are
more tolerant of boundary shifts than others.
751
Small changes in TAD boundary location are previously indicated to be involved in
developmental and differentiation processes [37,48,52], and other studies have shown that
chromatin rearrangement that alters TAD boundaries is linked to cancer development [6,40,53].
However, many publicly available datasets of CTCF binding activity have limited resolution
and/or coverage. With CTCF binding sites scattered across the genome, there could be further
subtle shifts in local regulatory areas that have previously gone unnoticed, particularly in
developmental contexts or across different tissues.
759
760 Overall, this study demonstrates that genetic-epigenetic interaction analysis can reveal 3D
761 genome structure through the positioning of interacting genome features. We see in this how
762 genetics and genome structure can inform each other. These findings imply that including TAD
763 boundaries and TAD loops in analyzing genomic features affecting gene expression, such as
764 chromatin states and genetic variants, can maximize and contextualize results.
765

Acknowledgments
We thank G. Churchill for helpful comments. This study was funded by the National Institute of
General Medical Sciences of the National Institutes of Health under award number
R01GM115518. C.L.B was supported by the National Institute of General Medical Sciences
(NIGMS) grant R35GM133724. The content is solely the responsibility of the authors and does
not necessarily represent the official views of the National Institutes of Health.

772
773
774 **References**
775

1.   Ackermann, A. M., Wang, Z., Schug, J., Naji, A. & Kaestner, K. H. Integration of ATAC-seq

and RNA-seq identifies human alpha cell and beta cell signature genes. *Molecular*

Metabolism **5**, 233–244 (2016).

2.   Wu, J. *et al.* The landscape of accessible chromatin in mammalian preimplantation

embryos. *Nature* **534**, 652–657 (2016).

3.   Veyrieras, J.-B. *et al.* High-Resolution Mapping of Expression-QTLs Yields Insight into

Human Gene Regulation. *PLoS Genet.* **4**, (2008).

4.   Stranger, B. E. *et al.* Population genomics of human gene expression. *Nat. Genet.* **39**, 1217

(2007).

5.   Dixon, A. L. *et al.* A genome-wide association study of global gene expression. *Nat. Genet.*

39, 1202 (2007).

6.   Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of

chromatin interactions. *Nature* **485**, 376–380 (2012).

7.   Pombo, A. & Dillon, N. Three-dimensional genome architecture: players and mechanisms.

Nat. Rev. Mol. Cell Biol. **16**, 245 (2015).

8.   Harmston, N. *et al.* Topologically associating domains are ancient features that coincide

with Metazoan clusters of extreme noncoding conservation. *Nat. Commun.* **8**, (2017).

9.   Yakushiji-Kaminatsui, N. *et al.* Similarities and differences in the regulation of HoxD genes

during chick and mouse limb development. *PLoS Biol.* **16**, (2018).

10. Serebrenik, Y. V. & Shalem, O. CRISPR mutagenesis screening of mice. *Nat. Cell Biol.* **20**,

1235 (2018).

11. Svenson, K. L. *et al.* High-Resolution Genetic Mapping Using the Mouse Diversity Outbred

Population. *Genetics* **190**, 437 (2012).

12. Churchill, G. A., Gatti, D. M., Munger, S. C. & Svenson, K. L. The Diversity Outbred Mouse

Population. *Mamm. Genome* **23**, 713 (2012).

13. Broman, K. W. Haplotype Probabilities in Advanced Intercross Populations. *G3:*

Genes|Genomes|Genetics **2**, 199 (2012).

14. Chesler, E. J. *et al.* Diversity Outbred Mice at 21: Maintaining Allelic Variation in the Face of

Selection. *G3: Genes|Genomes|Genetics* **6**, 3893 (2016).

15. Skelly, D. A. *et al.* Mapping the Effects of Genetic Variation on Chromatin State and Gene

Expression Reveals Loci That Control Ground State Pluripotency. *Cell Stem Cell* **27**,

(2020).

16. Robinson, M. D. & Oshlack, A. A scaling normalization method for differential expression

analysis of RNA-seq data. *Genome Biology* vol. 11 R25 Preprint at

https://doi.org/10.1186/gb-2010-11-3-r25 (2010).

17. Morgan, A. P. *et al.* The Mouse Universal Genotyping Array: From Substrains to

Subspecies. *G3: Genes, Genomes, Genetics* **6**, 263–279 (2016).

18. Morgan, A. P. argyle: An R Package for Analysis of Illumina Genotyping Arrays. *G3* **6**,

(2015).

19. Bubier, J. A. *et al.* Genetic variation regulates opioid-induced respiratory depression in

mice. *Sci. Rep.* **10**, 1–15 (2020).

20. Chick, J. M. *et al.* Defining the consequences of genetic variation on a proteome–wide

scale. *Nature* **534**, 500 (2016).

21. Kuhn, R. M., Haussler, D. & Kent, W. J. The UCSC genome browser and associated tools.

Briefings in Bioinformatics vol. 14 144–161 Preprint at https://doi.org/10.1093/bib/bbs038

(2013).

22. Robinson, J. T. *et al.* Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).

23. Bailey, T. L. *et al.* MEME Suite: tools for motif discovery and searching. *Nucleic Acids Res.*

37, W202–W208 (2009).

24. Bailey, T. L. & Elkan, C. Fitting a mixture model by expectation maximization to discover

motifs in biopolymers. *Proc. Int. Conf. Intell. Syst. Mol. Biol.* **2**, (1994).

25. Bailey, T. L. STREME: Accurate and versatile sequence motif discovery. *Cold Spring*

Harbor Laboratory 2020.11.23.394619 (2020) doi:10.1101/2020.11.23.394619.

26. Gupta, S., Stamatoyannopoulos, J. A., Bailey, T. L. & Noble, W. S. Quantifying similarity

between motifs. *Genome Biol.* **8**, 1–9 (2007).

27. Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given motif.

Bioinformatics vol. 27 1017–1018 Preprint at https://doi.org/10.1093/bioinformatics/btr064

(2011).

28. Keane, T. M. *et al.* Mouse genomic variation and its effect on phenotypes and gene

regulation. *Nature* **477**, 289.

29. Sloan, C. A. *et al.* ENCODE data at the ENCODE portal. *Nucleic Acids Res.* **44**, D726–

D732 (2015).

30. The ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the

human genome. *Nature* **489**, 57–74 (2012).

31. Kulakovskiy, I. V. *et al.* HOCOMOCO: towards a complete collection of transcription factor

binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids*

Res. **46**, D252–D259 (2017).

32. Oomen, M. E., Hansen, A. S., Liu, Y., Darzacq, X. & Dekker, J. CTCF sites display cell

cycle–dependent dynamics in factor binding and nucleosome positioning. *Genome Res.* **29**,

236–249 (2019).

33. GitHub - TheJacksonLaboratory/cs-nf-pipelines: The Jackson Laboratory Computational

Sciences Nextflow based analysis pipelines. *GitHub*

https://github.com/TheJacksonLaboratory/cs-nf-pipelines.

34. Ronald, J., Brem, R. B., Whittle, J. & Kruglyak, L. Local regulatory variation in

Saccharomyces cerevisiae. *PLoS Genet.* **1**, (2005).

35. Su, J., Teichmann, S. A. & Down, T. A. Assessing Computational Methods of Cis-

Regulatory Module Prediction. *PLoS Comput. Biol.* **6**, e1001020 (2010).

36. Nelson, C. E., Hersh, B. M. & Carroll, S. B. The regulatory content of intergenic DNA

shapes genome architecture. *Genome Biol.* **5**, 1–15 (2004).

37. Rodríguez-Carballo, E. *et al.* Chromatin topology and the timing of enhancer function at the

HoxD locus. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 31231–31241 (2020).

38.   Krijger, P. H. & de Laat, W. Regulation of disease-associated gene expression in the 3D

genome. *Nat. Rev. Mol. Cell Biol.* **17**, (2016).

39.   Lizio, M. *et al.* Gateways to the FANTOM5 promoter level mammalian expression atlas.

Genome Biol. **16**, 22 (2015).

40.   Tyler, A. L., Donahue, L. R., Churchill, G. A. & Carter, G. W. Weak Epistasis Generally

Stabilizes Phenotypes in a Mouse Intercross. *PLoS Genet.* **12**, e1005805 (2016).

41.   Andrew C. Leon, M. H. Sample Sizes Required to Detect Interactions between Two Binary

Fixed-Effects in a Mixed-Effects Linear Regression Model. *Comput. Stat. Data Anal.* **53**,

603 (2009).

42.   Cao, J. *et al.* Joint profiling of chromatin accessibility and gene expression in thousands of

single cells. *Science* **361**, 1380 (2018).

43.   Zuo, Z., Roy, B., Chang, Y. K., Granas, D. & Stormo, G. D. Measuring quantitative effects

of methylation on transcription factor–DNA binding affinity. *Science Advances* (2017)

doi:10.1126/sciadv.aao1799.

44.   Maurano, M. T. *et al.* Role of DNA Methylation in Modulating Transcription Factor

Occupancy. *Cell Rep.* **12**, (2015).

45.   Shen, Y. *et al.* A map of the cis-regulatory sequences in the mouse genome. *Nature* **488**,

(2012).

46.   On the choreography of genome folding: A grand pas de deux of cohesin and CTCF. *Curr.*

Opin. Cell Biol. **70**, 84–90 (2021).

47.   Li, Z. *et al.* Identification of transcription factor binding sites using ATAC-seq. *Genome Biol.*

20, 1–21 (2019).

48.   Andrey, G. *et al.* A Switch Between Topological Domains Underlies HoxD Genes

Collinearity in Mouse Limbs. *Science* **340**, (2013).

49. Kumasaka, N., Knights, A. J. & Gaffney, D. J. Fine-mapping cellular QTLs with RASQUAL

and ATAC-seq. *Nat. Genet.* **48**, 206–213 (2015).

50. Skelly, D. A. *et al.* Genetic variation influences pluripotent ground state stability in mouse

embryonic stem cells through a hierarchy of molecular phenotypes. *Cold Spring Harbor*

Laboratory 552059 (2019) doi:10.1101/552059.

51. Fasolino, M. *et al.* Genetic Variation in Type 1 Diabetes Reconfigures the 3D Chromatin

Organization of T Cells and Alters Gene Expression. *Immunity* **52**, 257–274.e11 (2020).

52. Takayama, N. *et al.* The Transition from Quiescent to Activated States in Human

Hematopoietic Stem Cells Is Governed by Dynamic 3D Genome Reorganization. *Cell Stem*

Cell **28**, (2021).

53. Menghi, F. *et al.* The Tandem Duplicator Phenotype Is a Prevalent Genome-Wide Cancer

Configuration Driven by Distinct Gene Mutations. *Cancer Cell* **34**, 197–210.e5 (2018).

893
894
895