Resistance bioassays and allele characterisation inform *Spodoptera frugiperda* (Lepidoptera:

Noctuidae) management and introduction pathways in Asia and Australia

Tay WT[1,3], Rane RV[2,3], James W[1], Gordon KHJ[1], Downes S[4], Kim J[5], Kuniata L[6], Walsh TK[1,3]

1. CSIRO Black Mountain Laboratories, Clunies Ross Street, ACT 2601, Australia

2. CSIRO, 343 Royal Parade, Parkville, VIC 3052, Australia

3. Applied BioSciences, Macquarie University, Sydney NSW 2100, Australia

4. CSIRO Mc Master Laboratories, New England Highway, Armidale NSW 2350, Australia

4. College of Agriculture and Life Science, Kangwon National University, Republic of Korea

5. Ramu Agri Industries Ltd., PNG

Correspondence: weetek.tay@csiro.au

Abstract

The fall armyworm *Spodoptera frugiperda* is present in over 70 countries in Africa, Asia, and Oceania. Its rapid

dispersal since 2016 since first reported in western Africa, and associated devastation to agricultural

productivity, highlight the challenges posed by this pest. Currently, its management largely relies on insecticide

sprays and transgenic *Bacillus thuringiensis* toxins, therefore understanding their responses to these agents

and characteristics of any resistance genes enables adaptive strategies. In Australia, *S. frugiperda* was reported

at the end of January 2020 in northern Queensland and by March 2020, also in northern Western Australia.

We undertook bioassays on two Australian populations, one each from these initial points of establishment. To

assist with resistance response interpretation, two related endemic noctuid pest species, *Helicoverpa armigera*

and *Spodoptera litura*, were concurrently screened to obtained LC50 estimates for larvae against a range of

insecticides. We characterised known resistance alleles from the VGSC, ACE-1, RyR, and ABCC2 genes for five

populations to compared with published allele frequencies and bioassay responses from native and invasive *S.*

frugiperda populations. An approximately 10x LC50 difference for indoxacarb was detected between

Australian populations, which was approximately 28x higher than an Indian population. Characterisation of

ACE-1 and VGSC alleles provided further evidence of multiple introductions in Asia, and multiple pathways

involving genetically distinct individuals into Australia. Bioassay results and resistance allele patterns from

invasive *S. frugiperda* populations therefore suggest multiple introduction events have contributed to the

pest's spread in the Old World, and challenge the axiom of its rapid 'west-to-east' spread.

Keywords

FAW, whole genome sequencing, agricultural biosecurity, population genomics, invasion biology

Introduction

The fall armyworm (FAW) *Spodoptera frugiperda* (J. E. Smith), is a noctuid moth pest made up of two morphologically indistinguishable strains (C- and R-strains) that is native to tropical and subtropical regions of the Americas. It is highly polyphagous and feeds on host species from at least 76 plant families, principally Poaceae (106 spp.), Asteraceae (31) and Fabaceae (31) (Montezano et al. 2018). In the Americas, it causes significant economic damage to maize, rice, sorghum, millet, soya bean, wheat, alfalfa, cotton, turf, and fodder crops (Nagoshi et al. 2019a). Recognised globally as one of the top 20 arthropod pests (Willis 2017), the FAW was first officially confirmed outside of its native range in Western Africa in early 2016 (Goergen et al. 2016) and then detected across virtually all of Sub-Saharan Africa by February 2018 (Nagoshi et al. 2018, Otim et al. 2018, Botha et al. 2019). In July 2018, it was confirmed in Yemen and India, and by December 2018 in Bangladesh, Sri Lanka, Myanmar and Thailand, followed closely by China in January 2019. Nepal confirmed its presence in May 2019, and in July 2019 it was also reported in Egypt, Indonesia, Laos, Cambodia, Malaysia, Vietnam, Taiwan, The Republic of Korea, and Japan (FAO 2021). The moth's strong flight ability (Rose et al. 1975, Sparks 1979, Jones et al. 2019), its potential to contaminate certain commodities, and ability to act as a 'hitchhiker' in trade (Early et al. 2018) contribute to its rapid spread. Whole genome analyses of invasive FAW populations from various African nations, India, China, Southeast Asia, and Australia also suggested that a high proportion of FAW individuals were hybrids of the C- and R-strains (Gui et al. 2020, Tay et al. 2020, Zhang et al. 2020, Schlum et al. 2021, Tay et al. 2021, Rane et al. 2022).

Much of South and Southeast Asia, and much of Northern Australia, are highly suitable climatically for the FAW year-round (du Plessis et al. 2018, Early et al. 2018). It was confirmed in Queensland (Qld; Bamaga Northern Peninsula Area), Australia following clearance of lure traps on 31 January 2020, followed by reports also in north-Western Australia (WA; Kununurra) and in the Northern Territory (NT) in March 2020. Within two weeks of the Bamaga detection, there were reports of FAW damaging maize fields in Strathmore (Qld; *ca*. 1,000 km from the initial Bamaga detection site), followed by successive southward detections of FAW in Qld, marking the FAW's south-eastward antipodean expansion into New South Wales (NSW) in September 2020. Through CLIMEX simulation analysis based on irrigation patterns and rainfall records, the FAW is expected to undertake seasonal migrations and may reach as far south as Victoria and Tasmania (du Plessis et al. 2018). Indeed, the FAW was reported in Victoria in December 2020 and in Tasmania in April, 2021. Concurrently, and from the Western Australia Kununurra detection, the FAW followed the CLIMEX simulated southward expansion patterns along the WA's western coast, with larval populations detected in the Gingin area (< 80 Km north of the city of Perth) by February 2021.

The incursion of FAW into Australia is believed to have occurred through natural dispersal across the Torres Strait/Timor Sea into northern Australia via a single-entry point (Jing et al. 2021, Qi et al. 2021). However, incursion pathways to Australia could also involve multiple entry points as reported for the biting midges *Culicoides bravitarsis* (Tay et al. 2016), other anthropogenic routes such as infested commodities in trade, as well as through seasonal or on-going new arrivals of individuals from populations that have successfully established in Asia and Southeast Asia (SEA). The eastward spread of FAW across sub-Saharan Africa, the

Middle East, the Indian sub-continent, SEA, China, and the Far East (South Korea, Japan) mostly followed
chronological detection and confirmation which led to a widely accepted belief that an invasive population
established in west Africa (Goergen et al. 2016) through a single founder, similar to the 'invasive bridgehead
effect' (Guillemaud et al. 2011), was the starting point for its subsequent global expansion. This hypothesis is
supported by an overwhelmingly homogeneous genetic signature from single gene markers (e.g., Cock et al.
2017, Nagoshi et al. 2018). Whole genome analysis and genome-wide single nucleotide polymorphic marker
analysis of invasive populations from Africa (Benin, Uganda, Malawi, Tanzania) and Asia (India, China) instead
suggest multiple founding events (Tay et al. 2020, Zhang et al. 2020, Schlum et al. 2021).

Distinguishing among single vs. multiple introductions within Australia (and elsewhere), and understanding the
resistance profiles of existing and any new incursions, is critical for informing the future management of this
pest. For example, this information will assist with forecasting likely resistance profiles of invasive populations
selected outside of Australia, and could prioritise efforts to prevent the introgression (potentially of multiple
separate introductions) of known insecticide resistance genes and alleles from endemic populations (e.g.,
Carvalho et al. 2013, Banerjee et al. 2017, Boaventura et al. 2020a, Guan et al. 2020) into invasive populations,
or through reciprocal introgression of newly selected/developed resistance traits from invasive populations to
native populations.

In this study, we report on bioassay experiments and resistance gene characterisation that aimed to
understand how the first invasive populations of *S. frugiperda* in Australia responded to selected approved
insecticide compounds and *Bacillus thuringiensis* (Bt) toxins in commercial transgenic plant varieties and
available as sprays for the grains and cotton industries. Since populations were presumed collected before
selection occurred in Australia these responses indicate selection against insecticides used elsewhere prior to
arrival. Comparing the response differences between two FAW populations from different northern Australia
regions indicated the possibility of multiple points of entry *cf.* a single entry and spread event as generally
assumed (e.g., Jing et al. 2021, Qi et al. 2021). To provide a first insight into the efficacy of key compounds and
toxins we compared the responses of FAW with two related major endemic crop pests in Australia,
Helicoverpa armigera (subspecies *conferta;* Hardwick 1965, Anderson et al. 2016, Pearce et al. 2017b, a, Zhang
et al. 2022) and *S. litura*, which are currently managed through adaptive resistance management plans. Finally,
we provided a critical review of resistance profiles in selected invasive populations to better understand the
invasion biology of FAW.

Material and Methods

Live insects

Live FAW populations were sourced from Queensland (Qld) and Western Australia (WA). This population
(CSIRO code: 'Sf20-1') consisted of 30 field-collected pupae for setting up the laboratory colony, was first
noticed around 19[th] Feb, 2020, was collected on 3[rd] March, 2020, and represented the first reported FAW
detected attacking maize in Australia's agricultural landscape. The sampling site for this population was from

the University of Queensland (UQ) field station (Rex Road, Walkamin Qld 4872) from Strathmore Station,

(Lat/Lon: -17.17892, 145.43359, Elevation 685m).

A second FAW population (CSIRO code: 'Sf20-4') was collected as larvae from a maize crop from the Northern

Australian Crop Research Alliance Pty Ltd, Kununurra, Western Australia. These larvae were collected on 17-

Aug, 2020 from the Kununurra Airport field trail block (off Victoria Highway; -15.10090, 128.81342) on V4-6

growth stage maize, and were all collected prior to applications of insecticides. A total of 30 larvae of different

instar stages arrived (= F_0), and 10 (six females and four males) survived to pupation to produce >200 F_1

individuals and to generate sufficient F_2 larvae to commence bioassays.

Two related endemic noctuid species, *S. litura* and *Helicoverpa armigera conferta* (hereafter as '*H. armigera*'),

that are pests of a range of broadacre cropping and horticulture industries (and especially grains and cotton)

were also included in the bioassay experiments to assist with interpretating results such as base line

susceptibility and tolerance levels to the chemicals and toxins tested. Approximately 30 *S. litura* larvae (from

Mareeba and Toowoomba, Qld) with unknown resistance profiles were used to start the laboratory colony.

The *H. armigera* colony (CSIRO general rearing 'GR' strain) was a long-established laboratory strain housed at

the Black Mountain Laboratory in the ACT, with individuals originally collected as eggs from cotton fields in

and around Narrabri NSW and were periodically replenished to sustain population genetic diversity and

minimise inbreeding. The *H. armigera* GR population was confirmed by bioassays as having no

Cry1Ac/Cry2Ab/VIP3A resistance alleles.

Colony maintenance

Helicoverpa armigera and *Spodoptera* species (*S. frugiperda* Sf20-1 and Sf20-4; *S. litura*) were reared by the

method of Teakle and Jensen (1985) at 25 °C, 50 ± 10% relative humidity, and with a light/dark cycle of 14:10

h. Cooked artificial diet consisted of 81 g soya flour, 37.5 g wheat germ, 33 g brewer's yeast, 2 g ascorbic acid,

2 g nipagen, 3 mL vegetable oil, and 13.5 g agar, and was made up with water to 1 litre. Diet was poured into

rearing cups or bioassay plates, allowed to cool and then stored in the refrigerator (4-8°C) for no longer than 3

days.

Bioassays

The bioassay experiments involved commercially available insecticidal compounds and Bt toxins (available in

transgenic varieties of cotton and as commercial sprays), as well as Bt toxins that were produced and purified

by CSIRO (Table 1). For all insecticides and Bt toxins, Sf20-1 and Sf20-4 FAW populations were tested. The *H.*

armigera 'GR' line was tested alongside the FAW against all chemicals and Bt toxins except Cry1F because it is

known to be insensitive to it. We included *S. litura* in all Bt toxin bioassays as it is regarded as a pest of interest

by the Australian cotton industry who depend on Bt varieties.

139   [Table 1]

All bioassays included a negative control for background mortality which was treated with the diluent of the

individual pesticide. An initial experiment using 10x serial dilutions was performed to narrow the range for the

detailed bioassay using 8-16 individuals per dose replicated once. The full bioassay involved 2x serial dilutions
at 6-7 doses with 16-24 individuals per dose replicated 3-5 times. The bioassays involved diet incorporation,
surface overlay, or topical application of the insecticidal compounds or Bt toxins (Table 1). The selection
among these exposure methods for an insecticide was determined by its mode of action (contact or ingestion)
and the experience of similar bioassays in related noctuid pests (i.e., *H. armigera* and *H. punctigera*; Bird
2015).

The surface overlay of Bt toxin assays followed the approach outlined in Mahon et al. (2007b), Mahon et al.
(2008), Mahon et al. (2010), Mahon et al. (2012) and Walsh et al. (2014). They were conducted in 96-well trays
where each well contained approximately 300µl of rearing diet with a surface area of around 0.567 cm^2 per
well. When the diet cooled, 20µl solution containing an appropriate concentration of toxin was added and
allowed to air dry. One neonate was added to each well and the tray was heat sealed with a perforated lidding
material.

For diet incorporation, commercial grade insecticides were diluted to the appropriate concentration with
water, added to 150ml of artificial diet and mixed well to produce a homogenous solution. Insecticide-
incorporated diet was then dispensed into 45-well bioassay trays, each with approximately 1.5 ml of diet-
insecticide mix. One late-second or early-third instar larva was added to each well and the tray was heat
sealed with a perforated lidding material (as per Bird 2015).

Topical bioassays of known concentrations were conducted for alpha-cypermethrin (group 3A) and methomyl
(group 1) and involved applying drops of pesticide as described in Bird (2018). Specifically, 1 µl of
acetone/insecticide solution was applied to the dorsal thorax of 3rd to 4th instar larvae using a 50 µl micro-
syringe. Bioassay trays were incubated at 25 ˚C, 45-55% RH, and a photoperiod 14:10 (L:D) h for 6 days (e.g.,
see Bird and Akhurst 2007, Downes et al. 2009). The numbers of dead (moribund; incapable of movement
when prodded) and alive larvae (capable of coordinated movement when prodded) were counted and the
instars of surviving larvae recorded. The LC$_{50}$ for each toxin was calculated from pooled raw data by probit
analysis using the POLO-PC program (LeOra Sorftware 1987).

FAW specimens for genomic analyses
Populations of *S. frugiperda* from South Korea (SK), Papua New Guinea (PNG), and Peru were included for
whole genome sequencing (Table 2). The methods for SK and PNG specimen preservation involved collection
from fields and placing larvae directly into high concentrations of ethanol (95-99.9%) to transport to the
laboratory where they were then stored at -20˚C and replaced with fresh ethanol after 24-48 hours. Peru
specimens were samples intercepted from Australian pre-border inspections of imported
agricultural/horticultural commodities between 2016 and 2019 (see Tay et al. 2020). Australian samples (Sf20-
1 from Qld, Sf20-4 from WA) were F$_0$ individuals and represented individuals obtained directly from fields.
Samples from SK, PNG, and Peru were sent to CSIRO and stored at -20˚C until DNA extraction.

DNA extraction and genome library preparation

DNA of individual FAW samples was extracted using the Qiagen DNA extraction kit and eluted in 200μL elution

buffer following protocols as provided by the manufacturer (Qiagen, Hilden Germany). Genomic DNA libraries

for individual samples were prepared, quantified and sent for commercial sequencing by the Australian

Genome Research Facility (AGRF) in Melbourne, Victoria, Australia.

Processing of genome sequences

Genome sequencing data for individuals were trimmed to remove adapter sequences using trim_galore (v

0.6.6; https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) and aligned to the *S. frugiperda* rice

genome (v1.0) (Gouin et al. 2017) using bwa_mem2 (v2) (Vasimuddin et al. 2019). Duplicate alignments were

removed using SAMBLASTER (v 0.1.26) (Faust and Hall 2014) and sorting completed using SAMtools (v1.9) (Li

et al. 2009).

188   [Table 2]

For identification of FAW R- or C-strain, the used the partial mtCOI gene sequences for the R-strain (GenBank

MF197867) and the C-strain (GenBank MF197868) of Otim et al. (2018) as reference sequence for mapping

against the whole genome sequence data for each individual FAW from Sf20-1, Sf20-4, SK, PNG, and Peru.

Mapping of partial mtCOI gene was carried out within Geneious v11.1.5 (Biomatters Ltd, Auckland, NZ) using

the Geneious Mapper program with assembly parameters specified to 'Low Sensitivity / Fastest', with no

trimming before mapping, and the 'Fine Tuning' option set to 'Iterate 2 times' to map reads to the consensus

from the previous iteration. Full mitochondrial DNA genomes of all individuals were also assembled following

the procedures as outlined in (Otim et al. 2018) and have been reported by Rane et al. (2022). For all

resistance alleles of interest (ABCC2, ACE-1, RyR, VGSC; see Table S1), base-sequence at the genomic location

for individuals listed in Table 1 was extracted from the alignment file using BCFTOOLS MPILEUP followed by

CALL, since BBMAP does not report 'non-variant' sites. Results were tabulated for inference.

Comparisons of published ACE-1 and VGSC resistance allele frequencies and published indoxacarb and

chlorantraniliprole LC$_{50}$ values

Three resistance loci for the organophosphate/carbamate ACE-1 gene have been reported to-date in FAW: (i)

A201S, (ii) G227A, and (iii) F290V. To understand the frequencies of these alleles, we surveyed specimens from

invasive (i.e., Australia, PNG, South Korea) and native (Peru) FAW populations, and combined this information

with reported allele frequencies for these three loci from other native (Brazil, French Guiana, Mexico

Guadeloupe, Puerto Rico, USA) and invasive populations (Benin, Uganda, Kenya, Tanzania, Zambia, Malawi;

India, Myanmar, Laos, Vietnam, Malaysia, Indonesia, China, Australia) (see Table S2 and references therein).

For the pyrethroid *para* sodium channel resistance gene VGSC, three loci have been identified to date in FAW:

(i) L1014F, (ii) L932F, and (iii) T929I (e.g., Carvalho et al. 2013, Guan et al. 2020, Yainna et al. 2021). Resistance

and susceptible allele frequencies in these three VGSC loci were also surveyed from published studies from

native and invasive FAW populations (see Table S2).

For comparisons of bioassay LC_{50} data for indoxacarb and chlorantraniliprole between published data and our
study, we considered only those that reported broadly similar methodologies (see Table 1) such as clearly
stated route of delivery (i.e., diet incorporation/ingestion) for the insecticidal compounds, and using
comparable developmental stages of larvae (i.e., 2^{nd}/early 3^{rd} instar) and scoring criteria.

Results

Bioassays of Individual Bt proteins

Cry1Ac was not highly effective against the *Spodoptera* species tested in this study (Figure S1; Table 3). Based
on the LC_{50}, and relative to *H. armigera*, FAW Sf20-1 was 174x less sensitive, FAW Sf20-4 was 99x less
sensitive, and *S. litura* was 120x less sensitive. This suggests that in *S. frugiperda*, Cry1Ac would give similar
control in Australia to that found in *S. litura* but far less than for *H. armigera*.

The *Spodoptera* species are less sensitive to Cry2Ab than *H. armigera* but the differences are not as striking as
for Cry1Ac (Figure S2; Table 3); FAW Sf20-1 was 13x less sensitive, FAW Sf20-4 was 4x less sensitive and *S.*
litura was 10x less sensitive. This suggests that in *S. frugiperda*, Cry2Ab would give similar control in Australia
to that found in *S. litura* which is unlikely to be different from *H. armigera* in terms of field control.

Cry1F is a Bt protein that has been deployed in certain genetically modified plants to target *Spodoptera*.
Though not effective against *H. armigera sensu lata* even at a relatively high dose, Cry1F was effective against
S. frugiperda at a much lower dose (Figure S3; Table 3), indicating that this species is much more sensitive to
this Bt protein than *H. armigera*. Relative to *S. litura,* the LC_{50} data shows that FAW Sf20-1 and FAW Sf20-4 are
2x less sensitive to Cry1F. In terms of field control there is unlikely to be any distinguishable difference in Cry1F
efficacy against FAW vs. *S. litura.*

FAW Sf20-1 and FAW Sf20-4 showed a similar tolerance to Vip3A as *S. litura* (~2.34x and 1.2x higher,
respectively) and all three populations were more sensitive to Vip3A than *H. armigera* (0.15x, 0.08x, and 0.07,
respectively). In terms of field control there is unlikely to be any distinguishable difference in efficacy of VIP3A
against FAW relative to *H. armigera* and *S. litura* (Figure S4; Table 3).

Based on published studies which show resistance ratios of >2,500 fold in *H. armigera* and FAW that are
homozygous for recessive resistance alleles against Bt toxins (e.g., Mahon et al. (2007a) - Cry2Ab, *H. armigera*;
Horikoshi et al. (2016), Cry1F, Cry1A, Vip3A, FAW) it is unlikely that either of the founding FAW populations in
Australia carried resistance alleles to Cry1Ac, Cry2Ab or Vip3A in homozygous states.

<u>Sprayable Bt</u>

The performance against FAW of the sprayable products containing multiple Bt toxins (XenTari and DIPEL)
showed both products to differ in efficacies against *H. armigera*, FAW, and *S. litura*. XenTari with the mixture
of Cry1 toxins including Cry1C was formulated to provide *Spodoptera* control ([valentbiosciences](), last accessed
13-Jan, 2022) but also for cabbage looper and the diamondback moth (DBM), while DIPEL was formulated for

broad spectrum caterpillar control ([valentbiosciences](), last accessed 13-Jan, 2022). Both DIPEL and XenTari are

reported to contain different toxin complements but the relative amounts are unclear. DIPEL contains Cry1Ac

and Cry2-type toxins and was less effective against both tested FAW populations than XenTari which contains

Cry1Aa, Cry1Ab, Cry1C and Cry1D. Relative to *H. armigera*, FAW Sf20-1 was 24x less sensitive, FAW Sf20-4 was

18x less sensitive and *S. litura* was 17x less sensitive to Dipel. XenTari toxicity ratios relative to *H. armigera*

ranged 2 to 4x for FAW Sf20-1, FAW Sf20-4 and *S. litura* (Figure S5 and Figure S6; Table 3).

<u>Conventional pesticides</u>

Cypermethrin is a pyrethroid pesticide and the bioassay results suggest that FAW Sf20-1 and FAW Sf20-4 are

much less sensitive to it than *H. armigera* (56x and 145x respectively: Figure S7; Table 4). Broadly speaking,

given that pyrethroid resistance is common in the field in *H. armigera* (Walsh et al. 2018), this result suggests

that it may be difficult to control *S. frugiperda* with pyrethroids. It also indicates some heterogeneity in the

response to pyrethroids in the first invasive FAW populations in Australia.

Bioassay results for the carbamate pesticide methomyl are not as repeatable as other toxins with considerable

variability between replicates (Figure S8; Table 4). However, overall FAW Sf20-4 shows 52x less sensitivity to

methomyl than *H. armigera* while FAW Sf20-1 shows a 4x lower sensitivity. This difference is consistent with

some level of heterogeneity between the first invasive FAW populations in Australia.

The dose response and LC_{50} for indoxacarb (Figure S9; Table 5) show that FAW Sf20-1 and FAW Sf20-4 are less

sensitive to this chemistry than *H. armigera* (22x and 208x respectively). This difference is consistent with

some level of heterogeneity between the first invasive FAW populations in Australia.

The dose response and LC_{50} for chlorantraniliprole show similar sensitivity for *H. armigera* and FAW Sf20-1,

with the latter showing a 3x difference in the LC_{50}. FAW Sf20-4 demonstrated a 15x reduction in sensitivity

compared to *H. armigera* (Figure S10; Table 5).

The dose response and LC_{50} data for emamectin benzoate and spinetoram (Figure S11, Figure S12; Table 5)

suggest that FAW Sf20-1 and FAW Sf20-4 are not significantly different in their level of sensitivity compared to

H. armigera.

[Table 3, Table 4, Table 5]

Resistance alleles by whole genome sequencing

The loci examined for potential resistance alleles were present in the data set with the appropriate level of

coverage to accurately call the genotype. Resistance alleles to carbamate/organophosphates (ACE-1) were the

only resistance alleles identified in the Kununnura, Strathmore, South Korea, Papua New Guinea, and Peru

populations. No resistance alleles were detected for pyrethroid (VGSC) or for diamide insecticides (RyR).

Resistance allele profile differences between the two Australian populations were evident between the A201S

and the F290V amino acid substitutions. Similar ACE-1 resistance allele profiles were detected between

Strathmore and South Korean populations, while the Papua New Guinea population shared ACE-1 allele

profiles with Strathmore population (A201S) and with Kununnura (F290V). The Peruvian FAW population was

the only population that have heterozygous individuals with resistance allele for G227A amino acid
substitution (Table 6).

284   [Table 6]

Resistance allele characterisation by sequencing approaches from this study and published studies for the ACE-
1 gene is summarised in (Figure 1, Table S2 and references therein). The most common resistance allele
detected in the invasive and native population (456 individuals examined in total) was the F290V mutation
with 66 homozygous and 222 heterozygous resistant genotypes detected. This mutation is a T to G single
nucleotide polymorphism (SNP) that changes the codon encoding the amino acid from TTT to GTT leading to a
phenylalanine (F) to valine (V) change in the protein sequence encoded by the ACE-1 gene. This mutation was
present at all locations and evenly distributed between the invasive and native populations. In Australia, of the
sequenced individuals (n = 146), 24.7% (n = 36) were heterozygous, and 8% (n = 12) were homozygous for the
resistance allele. Heterozygous and homozygous individuals were found in populations from all four states
(i.e., WA, NT, Queensland, and NSW) as well as from the single individual from Erub Island, suggesting it is
common across Australia.

The second most common allele detected was A201S (456 individuals examined) with 90 heterozygous
individuals detected. This mutation is a C to A SNP which changes the codon from GCG to TCG leading to an
alanine (A) to serine (S) amino acid change. While less common, this mutation also appears to be in both the
invasive and native range with no obvious pattern. In Australia, of the sequenced individuals (n = 146), 15.8%
(n = 23) were heterozygous, and 0 were homozygous for the resistance allele. Heterozygous individuals were
found in WA and Queensland, also suggesting it is common across Australia.

The G227A mutation was the least common (from 456 individuals) with 20 heterozygous individuals and 2
homozygous individuals detected. This mutation is a G to a C SNP which alters the codon from GGA to CGA
encoding a glycine (G) to alanine (A) amino acid change. Interestingly G227A was only present in individuals
from the native range (Brazil, USA, Puerto Rico, Peru) but absent in individuals from across the invasive range
that were surveyed in this study and from the related studies of Tay et al. (2020) and Tay et al. (2021). When
compiling allele frequencies for both the VGSC and ACE-1 resistant genes from this study and from published
whole genome sequencing and targeted PCR/Sanger sequencing data (Figure 1; see also Table S1), the rare
G227A resistance allele was present only in the Indonesian (Boaventura et al. 2020a) and Hubei populations
(Guo et al. 2020) but absent in African, Australian, and other Asian (e.g., Indian, Myanmar, etc.) populations,
including populations from six other Chinese provinces.

No alleles predicted to cause resistance to pyrethroids or the group 28 pesticides were identified in this work.
However, while the previously identified resistance alleles (Bolzan et al. 2019, Boaventura et al. 2020b) were
not detected, considerable variation was present in the RyR gene at the potential resistance loci. This should
be further investigated in conjunction with bioassays to establish whether any of the variants could contribute
to resistance.

ABCC2 resistance alleles in Australia, Papua New Guinea, South Korea, and Peru populations

None of the known and validated Cry1 resistance ABCC2 mutations were identified in the individuals
sequenced for this work however, some variation from the reference was observed (Figure S13). Numerous
synonymous mutations and a total of 31 non-synonymous mutations were observed in the coding sequence of
the ABCC2 gene in all sequenced individuals. Several of these non-synonymous mutations were associated
with a common deletion and insertion motif in the first exon where an 11bp deletion and a 2 bp insertion
maintain the reading frame but replace and change several amino acids. Most of the other mutations are the
result of one (i.e., single nucleotide variant; SNV) or two (i.e., multiple nucleotide variant; MNV) nucleotide
changes. Several of these mutations are found in other assemblies of *S. frugiperda* and likely reflect natural
variation. An alternative explanation for the variation is that they are associated with the corn strain which is
thought to make up at least some of the genome of the invasive populations. Only one mutation, a 2 bp
deletion in one individual collected from the Burdekin in Queensland was identified that might cause a frame
shift mutation as has been identified in Bt resistant individuals in other studies. It was present as a
heterozygote in the individual and has not been experimentally validated.

331   [Figure 1]

Discussion

In this study, we showed insecticide and Bt response differences in two of the first established FAW
populations in Australia, with the Queensland (Sf20-1) population being less tolerant to various insecticides
such as methomyl, chlorantraniliprole, and indoxacarb compared to the Western Australian (Sf20-4)
population. On the other hand, the Queensland population exhibited at least a two-fold higher tolerance to
Cry1Ac, Cry2Ab, Cry1F and Vip3A Bt toxins than the WA population. Characterisation of the ABCC2 resistance
gene identified one FAW individual from Queensland as potentially being heterozygous with a 2 bp deletion
that could underpin Cry1F resistance, although confirmation of the detected mutation and of the resistance
phenotype is required. This could be accomplished via PCR and Sanger sequencing (e.g., Guan et al. 2020) as
well as via the CRISPR/Cas9 gene editing approach (e.g., see Wang et al. 2017). The response differences to
chemical insecticides and Bt toxins of the two studied invasive FAW populations in Australia and their ACE-1
resistance allele profile differences (Table 6) may reflect the diverse genetic composition across the pest's
recent expanding range (Tay et al. 2020, Schlum et al. 2021, Rane et al. 2022), and suggest that separate
pathways were involved in establishment of these Queensland and Western Australian populations. This is
contrary to the current postulation of a single introduction pathway for the arrival of FAW to Australia based
on an assumption (Jing et al. 2021) or reverse trajectory simulation (Qi et al. 2021) and highlights the
importance of harmonising simulation studies with genomic and phenotypic evidence.

With significant economic impacts on agriculture from FAW reported in over 80 countries (excluding the New
World native range) from Africa, Middle East, Asia, Southeast Asia, and Oceania, its response to different
insecticidal and Bt control agents is increasingly investigated at recently impacted localities (e.g., Worku and
Ebabuye 2019, Deshmukh et al. 2020, Zhang et al. 2020, Kulye et al. 2021, Lv et al. 2021). However, the diverse
bioassay methods (approach, larval stage, scoring criteria) used in these studies complicate meaningfully
comparison of findings. Taking emamectin benzoate and indoxacarb as examples, Zhang et al. (2022) used a

topical application bioassay on 3^{rd} instar larvae, Deshmukh et al. (2020) used leaf-dip bioassays on 2^{nd} instar

larvae, while Hardke et al. (2011) used a diet-incorporation approach on 3^{rd} instar larvae. Responses have been

expressed as LD_{50} (e.g., Deshmukh et al. 2020) or as EC_{50} (e.g., Kulye et al. 2021), further making it difficult to

rigorously compare outcomes.

A further challenge for invasive FAW management is the lack of base-line values representing susceptible

responses in newly populated areas which makes it difficult to monitor changes through time in insecticide

efficacy due to resistance. Knowledge of whether the introduction occurred once (e.g., single 'invasive

bridgehead effect'; Guillemaud et al. 2011) vs. multiple times (e.g., multiple 'mass dispersal'; Wilson et al.

2009) and patterns of gene flow also impact the long-term monitoring of insecticide resistance evolution.

Using native FAW colonies established in 2005 from cotton fields in Louisiana, USA, Wilson et al. (2009)

undertook diet-incorporation bioassays (on 3^{rd} instar larvae), using insecticides including chlorantraniliprole,

indoxacarb, and spinetoram (see Table 7). Yu (1991) undertook topical application bioassays against methomyl

(on 4^{th} instar larvae) which included comparison with a susceptible population free from insecticide exposure

since 1975, and a resistant population collected from a maize field in Gainesville, Florida (see Table 8). In

comparing insecticide responses in Indian invasive FAW populations with diet incorporation assays (on 3^{rd}

instar larvae), Yu (1991) also included a native susceptible FAW population from Brazil to assist with

interpreting changes to insecticide responses (including chlorantraniliprole and spinetoram) at spatial and

temporal scales.

In the absence of native FAW populations in Australia, and in addition to the information collected herein for

H. armigera and *S. litura,* we used the information from these global studies on native and newly invaded

populations to assist interpretation of the early data collected in this study on invasive FAW populations in

Australia. These examples used broadly similar approaches, instars and scoring criteria to our study for

bioassays with specific insecticides. Relative to the indoxacarb results from Hardke et al. (2011) on native FAW

populations from the USA, our bioassay findings for the invasive Western Australian and Queensland

populations suggest a 28- and 3-folds difference, respectively (Table 7). For spinetoram, the responses of the

two Australia populations (which were like each other) is around 10 and 1.5 times higher than for the native

populations from Brazil and the USA respectively. In contrast the ratios for the invasive populations from India

was 1.3 and 0.3 relative to the Brazil and USA native populations. The differences are even more pronounced

for chlorantraniliprole where the Sf20-4 WA population exhibited ratios that were 32 and 2.4 times higher

than both the native Brazilian and American populations, which are at least around 3-fold higher than for the

invasive populations in South Africa and India (Table 7) and also for the invasive Indian population studied by

Deshmukh et al. (2020) using a leaf dip assay (and hence not reported in Table 7).

Methomyl resistance alleles have been reported in invasive populations from China, Indonesia, Africa, and in

this study. Comparisons between susceptible and resistant strains of native FAW populations from Florida USA

(Yu 1991) and with the South African invasive population of FAW (Eriksson 2019) showed that the Queensland

population was similar in its response as the susceptible Florida FAW, while the Western Australia population

was around 5 times more tolerant. This contrasted the South African population which was 110 times more

tolerant than the Florida susceptible FAW strain, and around 8 times more tolerant than the resistant FAW

strain from Florida (Table 8). Taken as a whole, similar bioassay results between native (i.e., Brazil, USA) and

various invasive populations suggested potential significant genetic diversity in introduced populations. This

concurs with population genomic analyses (Tay et al. 2020, Zhang et al. 2020, Schlum et al. 2021, Rane et al.

2022) that suggested multiple origins for the invasive African, Asian (Indian, Chinese), and Southeast Asian

(e.g., Malaysia) FAW populations.

It is possible that the differential response between the two Australian FAW populations to insecticides and Bt

toxins represents natural variation in the national population which has emerged from a single founding

incursion. It is unlikely that the differential responses within Australian populations represents local selection

pressures because the period between populations establishing and being collected for this study was

presumed to be too short to enable this opportunity. It could be that separate incursions involving different

source populations occurred in WA compared to the eastern states of Australia. Different selection pressures

on the global population which recently originated from multiple Asian, Southeast Asian, and African

incursions (Tay et al. 2020, Zhang et al. 2020, Schlum et al. 2021, Rane et al. 2022) may have driven different

phenotypes which entered the expanding ranges tested in our study and others reported herein. However,

Kulye et al. (2021) demonstrated in FAW populations collected in India during 2018, 2019, and 2020 that large

response changes such as those observed in South African (e.g., methomyl; Eriksson 2019) and Western

Australian (chlorantraniliprole, indoxacarb; this study) populations were unlikely considering the short time

frame since the very recent arrival of FAW especially assuming a west-to-east spread (Goergen et al. 2016,

Cock et al. 2017, Nagoshi et al. 2018, Nagoshi et al. 2019b). Another explanation for the differential responses

of the two Australian populations that we studied is that they have different fitness because of variation in

genetic diversity which reflects the number of individuals founding the populations. However, if this was the

case one might expect sensitivity levels to be consistently higher or lower to all of the agents that we tested

but this was not the case.

[Table 7, Table 8]

Alien invasive agricultural pests are increasingly being shown to carry novel insecticide resistance genes (e.g.,

Anderson et al. 2018, Walsh et al. 2018, Tay and Gordon 2019). This includes the FAW in which invasive

populations have been confirmed via whole genome sequence analyses (e.g., Guan et al. 2020, Zhang et al.

2020, Yainna et al. 2021) and molecular characterisation to harbour selected resistance genes (Boaventura et

al. 2020b, Boaventura et al. 2020a, Zhao et al. 2020). Our review of reported ACE-1 and VGSC resistance allele

frequency differences (Figure 1; Table S2 and references therein) suggests that the invasive FAW populations

within China (Guo et al. 2020), Indonesia (Boaventura et al. 2020a), Malaysia and Queensland, Australia (Tay et

al. 2021, Rane et al. 2022), and South Korea (this study) were genetically diverse and likely originated from

different native populations (Figure 1). This further supports the perceived rapid spread of FAW across Africa,

Asia, and Oceania as likely to also involve multiple independent introduction events.

Given the insecticide resistance allele frequency differences in FAW (e.g., Boaventura et al. 2020a, Guan et al.

2020, Lv et al. 2021, Yainna et al. 2021) and variation between the two populations sampled herein in bioassay

responses to some of the approved insecticides for broadacre cropping, effort in Australia and indeed, for
other invasive regions where possible, should now be directed to establishing baseline susceptibilities against
key chemistries for multiple populations across geographies. For instance, our comparisons with *H. armigera*
and *S. litura* suggest that pyrethroids like cypermethrin are unlikely to provide good control in Australia against
FAW but the large discrepancies between WA and Qld populations for methomyl and indoxacarb require study
of further populations before drawing firm conclusions. This information will be critical for on-going
monitoring of resistance allele frequencies to key chemistries and determining their field application
efficiencies, among geographies which is required to inform resistance management plans. It will also be
important to understand gene flow patterns between different populations. Early detection of potential future
introductions into Australia of novel resistance genes/alleles should also be a priority; for example, the VGSC
L1014 resistance allele and the ACE G227A resistance allele from Southeast Asia (Boaventura et al. 2020a) and
China (Guo et al. 2020), ryanodine receptor (RyR) resistance alleles from Brazil and China (Bolzan et al. 2019,
Boaventura et al. 2020b, Lv et al. 2021) and ABCC2 resistance alleles from the Americas (Banerjee et al. 2017,
Flagel et al. 2018, Guan et al. 2020, Yainna et al. 2021).

Being alert to new FAW incursions carrying novel resistance genes is relevant also for the Bt toxins. Our
bioassay findings suggest Australia's FAW populations likely do not carry Cry1F resistance alleles in
homozygous state known to exist in native range FAW populations (e.g., Banerjee et al. 2017, Flagel et al.
2018, Guan et al. 2020, Yainna et al. 2021), however the 2 bp deletion identified via whole genome sequencing
in a single heterozygous individual from Burdekin (Queensland) will require further confirmation. Field-
selected lines of VIP3A20 resistant FAW have been reported in Brazil and the USA where they occur as a
recessive trait (Yang et al. 2013, Bernardi et al. 2015, Bernardi et al. 2016, Yang et al. 2018, Yang et al. 2019).
The candidate resistance gene(s) and associated mutation(s) underpinning this resistance are yet to be
identified, therefore phenotypic bioassays are required to detect resistance to VIP3A. Using what was
essentially an F_0 screen of material we did not detect resistance to VIP3A in our field-derived laboratory-
maintained FAW colonies, however bioassays involving F_2-crosses (Andow and Alstad 1998) would be needed
to detect individuals that are heterozygous for a recessive form of VIP3A resistance allele. VIP3A remained
highly effective when tested against both populations, and should be a valuable tool in protecting the
Australian cotton industry against FAW due to the > 90% up-take of Bollgard III cotton containing Cry1Ac,
Cry2Ab and Vip3A proteins. To protect agriculture in Australia (and elsewhere) from an accidental introduction
of VIP3A resistant populations from the Americas (e.g., from Brazil, Bernardi et al. 2016) into global invasive
populations, national (e.g., pre-border) and industry biosecurity preparedness strategies must be co-ordinated
to prevent and to increase the chances of early detection of such novel introductions of new traits.

Finally, to enable meaningful comparisons of bioassay findings from across the FAW invasive range, there is a
need to globally standardise approaches for testing of insecticides and Bt toxins. Localised novel resistance
traits in populations of FAW would likely serve as a new resistance management challenge to other
neighbouring regions (Kalyebi 2020), and movements of FAW in the new invasive ranges could lead to as yet

unknown and complex gene flow patterns that could significantly hinder the development of suitable pest and
resistance management strategies for this global pest complex.

Acknowledgements

The *Spodoptera frugiperda* larvae used to establish the SF20-1 colony were provided by Prof. Myron Zalucki
(University of Queensland). Mr Jack Daniel (Northern Australian Crop Research Alliance Pty Ltd) provided the
S. frugiperda larvae used to establish SF20-4. Material from an established colony of *S. litura* was provided by
Dr Ian Newton and Dr Melina Miles (QDAF). Some of the results reported herein were collected for a project
led by CSIRO as a co-investment with the Grains Research and Development Corporation, the Australian
Centre for International Agricultural Research, the Cotton Research and Development Corporation, FMC
Australasia and Corteva Agriscience. GRDC manages the project on behalf of the co-investors.

References

Anderson, C. J., W. T. Tay, A. McGaughran, K. Gordon, and T. K. Walsh. 2016. Population structure and gene
flow in the global pest, *Helicoverpa armigera*. Mol Ecol 25: 5296-5311.

Anderson, C. J., J. G. Oakeshott, W. T. Tay, K. H. J. Gordon, A. Zwick, and T. K. Walsh. 2018. Hybridization and
gene flow in the mega-pest lineage of moth, *Helicoverpa*. Proc Natl Acad Sci U S A 115: 5034-5039.

Andow, D. A., and D. N. Alstad. 1998. F-2 screen for rare resistance alleles. Journal of Economic Entomology
91: 572-578.

Banerjee, R., J. Hasler, R. Meagher, R. Nagoshi, L. Hietala, F. Huang, K. Narva, and J. L. Jurat-Fuentes. 2017.
Mechanism and DNA-based detection of field-evolved resistance to transgenic Bt corn in fall armyworm
(*Spodoptera frugiperda*). Sci Rep 7: 10877.

Bernardi, O., D. Bernardi, R. S. Ribeiro, D. M. Okuma, E. Salmeron, J. Fatoretto, F. C. L. Medeiros, T. Burd,
and C. Omoto. 2015. Frequency of resistance to Vip3Aa20 toxin from Bacillus thuringiensis in *Spodoptera*
frugiperda (Lepidoptera: Noctuidae) populations in Brazil. Crop Prot 76: 7-14.

Bernardi, O., D. Bernardi, R. J. Horikoshi, D. M. Okuma, L. L. Miraldo, J. Fatoretto, F. C. L. Medeiros, T. Burd,
and C. Omoto. 2016. Selection and characterization of resistance to the Vip3Aa20 protein from Bacillus
thuringiensis in *Spodoptera frugiperda*. Pest Management Science 72: 1794-1802.

Bird, L. J. 2015. Baseline Susceptibility of *Helicoverpa armigera* (Lepidoptera: Noctuidae) to Indoxacarb,
Emamectin Benzoate, and Chlorantraniliprole in Australia. Journal of Economic Entomology 108: 294-300.

Bird, L. J. 2018. Pyrethroid and carbamate resistance in Australian *Helicoverpa armigera* (Lepidoptera:
Noctuidae) from 2008 to 2015: what has changed since the introduction of Bt cotton? B Entomol Res 108:
781-791.

Bird, L. J., and R. J. Akhurst. 2007. Variation in susceptibility of *Helicoverpa armigera* (Hubner) and Helicoverpa
punctigera (Wallengren) (Lepidoptera : Noctuidae) in Australia to two *Bacillus thuringiensis* toxins. J
Invertebr Pathol 94: 84-94.

Boaventura, D., M. Martin, A. Pozzebon, D. Mota-Sanchez, and R. Nauen. 2020a. Monitoring of Target-Site
Mutations Conferring Insecticide Resistance in *Spodoptera frugiperda*. Insects 11: 545.

Boaventura, D., A. Bolzan, F. E. O. Padovez, D. M. Okuma, C. Omoto, and R. Nauen. 2020b. Detection of a
ryanodine receptor target-site mutation in diamide insecticide resistant fall armyworm, *Spodoptera*
frugiperda. Pest Management Science 76: 47-54.

Bolzan, A., F. E. O. Padovez, A. R. B. Nascimento, I. S. Kaiser, E. C. Lira, F. S. A. Amaral, R. H. Kanno, J. B.
Malaquias, and C. Omoto. 2019. Selection and characterization of the inheritance of resistance of
Spodoptera frugiperda (Lepidoptera: Noctuidae) to chlorantraniliprole and cross-resistance to other
diamide insecticides. Pest Management Science 75: 2682-2689.

Botha, A. S., A. Erasmus, H. du Plessis, and J. Van den Berg. 2019. Efficacy of Bt Maize for Control of
Spodoptera frugiperda (Lepidoptera: Noctuidae) in South Africa. Journal of Economic Entomology 112:
1260-1266.

Carvalho, R. A., C. Omoto, L. M. Field, M. S. Williamson, and C. Bass. 2013. Investigating the Molecular
Mechanisms of Organophosphate and Pyrethroid Resistance in the Fall Armyworm *Spodoptera*
frugiperda. Plos One 8: e62268.

Cock, M. J. W., P. K. Beseh, A. G. Buddie, G. Cafa, and J. Crozier. 2017. Molecular methods to detect
Spodoptera frugiperda in Ghana, and implications for monitoring the spread of invasive species in
developing countries. Sci Rep 7: 4103.

Deshmukh, S., H. B. Pavithra, C. M. Kalleshwaraswamy, B. K. Shivanna, M. S. Maruthi, and D. Mota-Sanchez.
2020. Field efficacy of insecticides for management of invasive fall armyworm, *Spodoptera frugiperda* (J.
E. Smith) (Lepidoptera: Noctuidae) on maize in India. Fla Entomol 103: 221-227.

Downes, S., T. L. Parker, and R. J. Mahon. 2009. Frequency of Alleles Conferring Resistance to the *Bacillus*
thuringiensis Toxins Cry1Ac and Cry2Ab in Australian Populations of *Helicoverpa punctigera* (Lepidoptera:
Noctuidae) From 2002 to 2006. Journal of Economic Entomology 102: 733-742.

du Plessis, H., J. van den Berg, N. Ota, and D. J. Kriticos. Year. Published. *Spodoptera frugiperda* (Fall
Armyworm), p. 7. *In*, CSIRO-InSTePP Pest Geography, June, 2018 2018, Canberra, Australia. CSIRO.

Early, R., P. Gonzalez-Moreno, S. T. Murphy, and R. Day. 2018. Forecasting the global extent of invasion of the
cereal pest *Spodoptera frugiperda*, the fall armyworm. Neobiota: 25-50.

Eriksson, J. 2019. Toxicity bioassays with insecticide formulations used for control of *Spodoptera frugiperda*
(Lepidoptera: Noctuidae). Master of Science in Environmental Science with Integrated Pest Management,
North West University South Africa.

Faust, G. G., and I. M. Hall. 2014. SAMBLASTER: fast duplicate marking and structural variant read extraction.
Bioinformatics 30: 2503-2505.

Flagel, L., Y. W. Lee, H. Wanjugi, S. Swarup, A. Brown, J. L. Wang, E. Kraft, J. Greenplate, J. Simmons, N.
Adams, Y. F. Wang, S. Martinelli, J. A. Haas, A. Gowda, and G. Head. 2018. Mutational disruption of the
ABCC2 gene in fall armyworm, *Spodoptera frugiperda*, confers resistance to the Cry1Fa and Cry1A.105
insecticidal proteins. Sci Rep-Uk 8: 7255.

Goergen, G., P. L. Kumar, S. B. Sankung, A. Togola, and M. Tamo. 2016. First Report of Outbreaks of the Fall
Armyworm *Spodoptera frugiperda* (J E Smith) (Lepidoptera, Noctuidae), a New Alien Invasive Pest in West
and Central Africa. PLoS One 11: e0165632.

Gouin, A., A. Bretaudeau, K. Nam, S. Gimenez, J. M. Aury, B. Duvic, F. Hilliou, N. Durand, N. Montagne, I.
Darboux, S. Kuwar, T. Chertemps, D. Siaussat, A. Bretschneider, Y. Mone, S. J. Ahn, S. Hanniger, A. G.
Grenet, D. Neunemann, F. Maumus, I. Luyten, K. Labadie, W. Xu, F. Koutroumpa, J. M. Escoubas, A.
Llopis, M. Maibeche-Coisne, F. Salasc, A. Tomar, A. R. Anderson, S. A. Khan, P. Dumas, M. Orsucci, J.
Guy, C. Belser, A. Alberti, B. Noel, A. Couloux, J. Mercier, S. Nidelet, E. Dubois, N. Y. Liu, I. Boulogne, O.
Mirabeau, G. Le Goff, K. Gordon, J. Oakeshott, F. L. Consoli, A. N. Volkoff, H. W. Fescemyer, J. H.
Marden, D. S. Luthe, S. Herrero, D. G. Heckel, P. Wincker, G. J. Kergoat, J. Amselem, H. Quesneville, A.
T. Groot, E. Jacquin-Joly, N. Negre, C. Lemaitre, F. Legeai, E. d'Alencon, and P. Fournier. 2017. Two
genomes of highly polyphagous lepidopteran pests (*Spodoptera frugiperda*, Noctuidae) with different
host-plant ranges. Sci Rep 7: 11816.

Guan, F., J. Zhang, H. Shen, X. Wang, A. Padovan, T. K. Walsh, W. T. Tay, K. H. J. Gordon, W. James, C.
Czepak, M. H. Otim, D. Kachigamba, and Y. Wu. 2021. Whole-genome sequencing to detect mutations
associated with resistance to insecticides and Bt proteins in *Spodoptera frugiperda*. Insect Sci. 28: 627-
638.

Gui, F. R., T. M. Lan, Y. Zhao, W. Guo, Y. Dong, D. M. Fang, H. Liu, H. M. Li, H. L. Wang, R. S. Hao, X. F. Cheng,
Y. H. Li, P. C. Yang, S. K. Sahu, Y. P. Chen, L. Cheng, S. Q. He, P. Liu, G. Y. Fan, H. R. Lu, G. H. Hu, W. Dong,
B. Chen, Y. Jiang, Y. W. Zhang, H. H. Xu, F. Lin, B. Slipper, A. Postma, M. Jackson, B. A. Abate, K. Tesfaye,
A. L. Demie, M. D. Bayeleygne, D. T. Degefu, F. Chen, P. K. Kuria, Z. M. Kinyua, T. X. Liu, H. M. Yang, F.
N. Huang, X. Liu, J. Sheng, and L. Kang. 2020. Genomic and transcriptomic analysis unveils population
evolution and development of pesticide resistance in fall armyworm *Spodoptera frugiperda*. Protein Cell.
doi: 10.1007/s13238-020-00795-7.

Guillemaud, T., M. Ciosi, E. Lombaert, and A. Estoup. 2011. Biological invasions in agricultural settings:
Insights from evolutionary biology and population genetics. Cr Biol 334: 237-246.

Guo, J. M., X. L. Liu, S. R. Liu, Z. Q. Wei, W. K. Han, Y. Z. Guo, and S. L. Dong. 2020. Functional Characterization
of Sex Pheromone Receptors in the Fall Armyworm (*Spodoptera frugiperda*). Insects 11: 193.

Hardke, J. T., J. H. Temple, B. R. Leonard, and R. E. Jackson. 2011. Laboratory Toxicity and Field Efficacy of
Selected Insecticides against Fall Armyworm (Lepidoptera: Noctuidae). Fla Entomol 94: 272-278.

Hardwick, D. F. 1965. The Corn Earworm complex. Memoirs of the Entomological Society of Canada 40: 1-248.

Horikoshi, R. J., D. Bernardi, O. Bernardi, J. B. Malaquias, D. M. Okuma, L. L. Miraldo, F. S. D. E. Amaral, and
C. Omoto. 2016. Effective dominance of resistance of *Spodoptera frugiperda* to Bt maize and cotton
varieties: implications for resistance management. Sci Rep-Uk 6.

Jing, W., C. Huang, C. Y. Li, H. X. Zhou, Y. L. Ren, Z. Y. Li, L. S. Xing, B. Zhang, X. Qiao, B. Liu, C. H. Liu, Y. Xi, W.
X. Liu, W. K. Wang, W. Q. Qian, S. Mckirdy, and F. H. Wan. 2021. Biology, invasion and management of
the agricultural invader: Fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae). Journal of
Integrative Agriculture 20: 646-663.

Jones, C. M., H. Parry, W. T. Tay, D. R. Reynolds, and J. W. Chapman. 2019. Movement Ecology of Pest
Helicoverpa: Implications for Ongoing Spread. Annu Rev Entomol 64: 277-295.

Kalyebi, A. 2020. Documentation of farmer practices that have been useful in Africa for managing the fall
armyworm (*Spodoptera frugiperda*): A case study from Uganda. pp. 30. GRDC.

Kulye, M., S. Mehlhorn, D. Boaventura, N. Godley, S. K. Venkatesh, T. Rudrappa, T. Charan, D. Rathi, and R.
Nauen. 2021. Baseline Susceptibility of *Spodoptera frugiperda* Populations Collected in India towards
Different Chemical Classes of Insecticides. Insects 12: 758.

LeOra_Sorftware 1987. POLO-PC: A Users Guide to Probit or Logit Analysis computer program, version By
LeOra_Sorftware.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin, and S.
Genome Project Data Processing. 2009. The Sequence Alignment/Map format and SAMtools.
Bioinformatics 25: 2078-2079.

Lv, S. L., Y. Shi, J. C. Zhang, P. Liang, L. Zhang, and X. W. Gao. 2021. Detection of ryanodine receptor target-
site mutations in diamide insecticide-resistant *Spodoptera frugiperda* in China. Insect Science 28: 639-
648.

Mahon, R. J., K. M. Olsen, and S. Downes. 2008. Isolations of Cry2Ab resistance in Australian populations of
Helicoverpa armigera (Lepidoptera : Noctuidae) are allelic. Journal of Economic Entomology 101: 909-
914.

Mahon, R. J., S. J. Downes, and B. James. 2012. Vip3A Resistance Alleles Exist at High Levels in Australian
Targets before Release of Cotton Expressing This Toxin. Plos One 7.

Mahon, R. J., K. M. Olsen, K. A. Garsia, and S. R. Young. 2007a. Resistance to *Bacillus thuringiensis* toxin
Cry2Ab in a strain of *Helicoverpa armigera* (Lepidoptera : Noctuidae) in Australia. Journal of Economic
Entomology 100: 894-902.

Mahon, R. J., K. M. Olsen, S. Downes, and S. Addison. 2007b. Frequency of alleles conferring resistance to the
bt toxins Cry1Ac and Cry2Ab in Australian populations of *Helicoverpa armigera* (Lepidoptera : noctuidae).
Journal of Economic Entomology 100: 1844-1853.

Mahon, R. J., S. Downes, W. James, and T. Parker. 2010. Why Do F-1 Screens Estimate Higher Frequencies of

604   Cry2Ab Resistance in *Helicoverpa armigera* (Lepidoptera: Noctuidae) Than Do F-2 Screens? Journal of

605   Economic Entomology 103: 472-481.

606   **Montezano, D. G., A. Specht, D. R. Sosa-Gomez, V. F. Roque-Specht, J. C. Sousa-Silva, S. V. Paula-Moraes, J.**

607   **A. Peterson, and T. E. Hunt. 2018.** Host plants of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in the

608   Americas. Afr Entomol 26: 286-300.

609   **Nagoshi, R. N., G. Goergen, K. A. Tounou, K. Agboka, D. Koffi, and R. L. Meagher. 2018.** Analysis of strain

610   distribution, migratory potential, and invasion history of fall armyworm populations in northern Sub-

611   Saharan Africa. Sci Rep-Uk 8.

612   **Nagoshi, R. N., B. Y. Nagoshi, E. Canarte, B. Navarrete, R. Solorzano, and S. Garces-Carrera. 2019a.** Genetic

613   characterization of fall armyworm (*Spodoptera frugiperda*) in Ecuador and comparisons with regional

614   populations identify likely migratory relationships. PLoS One 14: e0222332.

615   **Nagoshi, R. N., I. Dhanani, R. Asokan, H. M. Mahadevaswamy, C. M. Kalleshwaraswamy, Sharanabasappa,**

616   **and R. L. Meagher. 2019b.** Genetic characterization of fall armyworm infesting South Africa and India

617   indicate recent introduction from a common source population. Plos One 14.

618   **Nguyen, D. T., Y. Z. Chen, and G. A. Herron. 2021.** Preliminary characterisation of known pesticide resistance

619   alleles in *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in its invasive Australian range. Austral Entomol

620   60: 782-790.

621   **Otim, M. H., W. T. Tay, T. K. Walsh, D. Kanyesigye, S. Adumo, J. Abongosi, S. Ochen, J. Sserumaga, S. Alibu,**

622   **G. Abalo, G. Asea, and A. Agona. 2018.** Detection of sister-species in invasive populations of the fall

623   armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) from Uganda. PLoS One 13: e0194571.

624   **Pearce, S. L., D. F. Clarke, P. D. East, S. Elfekih, K. H. J. Gordon, L. S. Jermiin, A. McGaughran, J. G. Oakeshott,**

625   **A. Papanicolaou, O. P. Perera, R. V. Rane, S. Richards, W. T. Tay, T. K. Walsh, A. Anderson, C. J.**

626   **Anderson, S. Asgari, P. G. Board, A. Bretschneider, P. M. Campbell, T. Chertemps, J. T. Christeller, C. W.**

627   **Coppin, S. J. Downes, G. Duan, C. A. Farnsworth, R. T. Good, L. B. Han, Y. C. Han, K. Hatje, I. Horne, Y. P.**

628   **Huang, D. S. T. Hughes, E. Jacquin-Joly, W. James, S. Jhangiani, M. Kollmar, S. S. Kuwar, S. Li, N. Y. Liu,**

629   **M. T. Maibeche, J. R. Miller, N. Montagne, T. Perry, J. Qu, S. V. Song, G. G. Sutton, H. Vogel, B. P.**

630   **Walenz, W. Xu, H. J. Zhang, Z. Zou, P. Batterham, O. R. Edwards, R. Feyereisen, R. A. Gibbs, D. G.**

631   **Heckel, A. McGrath, C. Robin, S. E. Scherer, K. C. Worley, and Y. D. Wu. 2017a.** Genomic innovations,

632   transcriptional plasticity and gene loss underlying the evolution and divergence of two highly

633   polyphagous and invasive *Helicoverpa* pest species. BMC Biol 15: 63.

634   **Pearce, S. L., D. F. Clarke, P. D. East, S. Elfekih, K. H. J. Gordon, L. S. Jermiin, A. McGaughran, J. G. Oakeshott,**

635   **A. Papanicolaou, O. P. Perera, R. V. Rane, S. Richards, W. T. Tay, T. K. Walsh, A. Anderson, C. J.**

636   **Anderson, S. Asgari, P. G. Board, A. Bretschneider, P. M. Campbell, T. Chertemps, J. T. Christeller, C. W.**

637   **Coppin, S. J. Downes, G. Duan, C. A. Farnsworth, R. T. Good, L. B. Han, Y. C. Han, K. Hatje, I. Horne, Y. P.**

638   **Huang, D. S. T. Hughes, E. Jacquin-Joly, W. James, S. Jhangiani, M. Kollmar, S. S. Kuwar, S. Li, N. Y. Liu,**

M. T. Maibeche, J. R. Miller, N. Montagne, T. Perry, J. Qu, S. V. Song, G. G. Sutton, H. Vogel, B. P.
Walenz, W. Xu, H. J. Zhang, Z. Zou, P. Batterham, O. R. Edwards, R. Feyereisen, R. A. Gibbs, D. G.
Heckel, A. McGrath, C. Robin, S. E. Scherer, K. C. Worley, and Y. D. Wu. 2017b. Erratum to: Genomic
innovations, transcriptional plasticity and gene loss underlying the evolution and divergence of two highly
polyphagous and invasive *Helicoverpa* pest species. BMC Biol 15: 69.

Qi, G. J., J. Ma, J. Wan, Y. L. Ren, S. McKirdy, G. Hu, and Z. F. Zhang. 2021. Source Regions of the First
Immigration of Fall Armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae) Invading Australia.
Insects 12: 1104.

Rane, R., T. K. Walsh, P. Lenancker, W. James, A. Gock, D. T. Hang, V. Nguyen. L, T. N. Khin, D. Amalin, K.
Chittarath, F. Muhammad, S. Annamalai, S. S. Thanarajoo, Y. A. Trisyono, K. Sahia, J. Kim, L. Kuniata, K.
Powell, K. H. J. Gordon, and W. T. Tay. 2022. Genome-wide SNP analysis suggests complex introductions
of *Spodoptera frugiperda* in Southeast Asia and Australia. eLife: submitted.

Rose, A. H., R. H. Silversides, and O. H. Lindquist. 1975. Migration Flight by an Aphid, *Rhopalosiphum maidis*
(Hemiptera-Aphididae), and a Noctuid, *Spodoptera frugiperda* (Lepidoptera-Noctuidae). Can Entomol
107: 567-576.

Schlum, K. A., K. Lamour, C. P. de Bortoli, R. Banerjee, R. Meagher, E. Pereira, M. G. Murua, G. A. Sword, A.
E. Tessnow, D. V. Dillon, A. M. L. Ramirez, K. S. Akutse, R. Schmidt-Jeffris, F. N. Huang, D. Reisig, S. J.
Emrich, and J. L. Jurat-Fuentes. 2021. Whole genome comparisons reveal panmixia among fall
armyworm (*Spodoptera frugiperda*) from diverse locations. BMC Genomics 22: 179.

Sparks, A. N. 1979. Review of the Biology of the Fall Armyworm (Lepidoptera, Noctuidae). Fla Entomol 62: 82-
87.

Tay, W. T., and K. H. J. Gordon. 2019. Going global - genomic insights into insect invasions. Curr Opin Insect Sci
31: 123-130.

Tay, W. T., P. J. Kerr, and L. S. Jermiin. 2016. Population Genetic Structure and Potential Incursion Pathways of
the Bluetongue Virus Vector *Culicoides brevitarsis* (Diptera: Ceratopogonidae) in Australia. Plos One 11:
e0146699.

Tay, W. T., R. Rane, B. James, A. Gock, D. T. Hang, V. L. Nguyen, T. N. Khin, D. Amalin, C. Khonesavanh, F.
Muhammad, S. Annamalai, and T. Walsh. 2021. Final report on Southeast Asian and Australian FAW
population genomics for biosecurity preparedness. Prevention and preparedness for fall armyworm
(*Spodoptera frugiperda*) - Output 2, pp. 25. CSIRO/GRDC.

Tay, W. T., R. Rane, A. Padovan, T. Walsh, S. Elfekih, S. Downes, K. Nam, E. d'Alencon, J. Zhang, Y. Wu, N.
Negre, D. Kunz, D. J. Kriticos, C. Czepak, M. Otim, and K. H. J. Gordon. 2020. Global FAW population
genomic signature supports complex introduction events across the Old World. bioRxiv
2020.06.12.147660.

Vasimuddin, M., S. Misra, H. Li, and S. Aluru. 2019. Efficient Architecture-Aware Acceleration of BWA-MEM
for Multicore Systems. Int Parall Distrib P: 314-324.

Walsh, T. K., S. J. Downes, J. Gascoyne, W. James, T. Parker, J. Armstrong, and R. J. Mahon. 2014. Dual
Cry2Ab and Vip3A Resistant Strains of *Helicoverpa armigera* and *Helicoverpa punctigera* (Lepidoptera:
Noctuidae); Testing Linkage Between Loci and Monitoring of Allele Frequencies. Journal of Economic
Entomology 107: 1610-1617.

Walsh, T. K., N. Joussen, K. Tian, A. McGaughran, C. J. Anderson, X. Qiu, S. J. Ahn, L. Bird, N. Pavlidi, J.
Vontas, J. Ryu, A. Rasool, I. Barony Macedo, W. T. Tay, Y. Zhang, M. E. A. Whitehouse, P. J. Silvie, S.
Downes, L. Nemec, and D. G. Heckel. 2018. Multiple recombination events between two cytochrome
P450 loci contribute to global pyrethroid resistance in *Helicoverpa armigera*. PLoS One 13: e0197760.

Wang, J., H. D. Wang, S. Y. Liu, L. P. Liu, W. T. Tay, T. K. Walsh, Y. H. Yang, and Y. D. Wu. 2017. CRISPR/Cas9
mediated genome editing of *Helicoverpa armigera* with mutations of an ABC transporter gene HaABCA2
confers resistance to *Bacillus thuringiensis* Cry2A toxins. Insect Biochem Molec 87: 147-153.

Willis, K. J. 2017. Sate of the World's Plants 2017. Report., pp. 100. *In* K. J. Killis [ed.]. Royal Botanic Gardens,
Kew.

Wilson, J. R. U., E. E. Dormontt, P. J. Prentis, A. J. Lowe, and D. M. Richardson. 2009. Something in the way
you move: dispersal pathways affect invasion success. Trends Ecol Evol 24: 136-144.

Worku, M., and Y. Ebabuye. 2019. Evaluation of efficacy of insecticides against the fall army worm *Spodoptera*
frugiperda. Indian Journal of Entomology 81: 13-15.

Xiao, H., X. Ye, H. Xu, Y. Mei, Y. Yang, X. Chen, Y. Yang, T. Liu, Y. Yu, W. Yang, Z. Lu, and F. Li. 2020. The
genetic adaptations of fall armyworm *Spodoptera frugiperda* facilitated its rapid global dispersal and
invasion. Mol Ecol Resour 20: 1050-1068.

Yainna, S., N. Negre, P. J. Silvie, T. Brevault, W. T. Tay, K. Gordon, E. dAlencon, T. Walsh, and K. Nam. 2021.
Geographic Monitoring of Insecticide Resistance Mutations in Native and Invasive Populations of the Fall
Armyworm. Insects 12: 468.

Yang, F., J. Williams, P. Porter, F. Huang, and D. L. Kerns. 2019. F-2 screen for resistance to *Bacillus*
thuringiensis Vip3Aa51 protein in field populations of *Spodoptera frugiperda* (Lepidoptera: Noctuidae)
from Texas, USA. Crop Prot 126.

Yang, F., F. N. Huang, J. A. Qureshi, B. R. Leonard, Y. Niu, L. P. Zhang, and D. S. Wangila. 2013. Susceptibility
of Louisiana and Florida populations of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) to transgenic
Agrisure (R) Viptera (TM) 3111 corn. Crop Prot 50: 37-39.

Yang, F., S. Morsello, G. P. Head, C. Sansone, F. N. Huang, R. T. Gilreath, and D. L. Kerns. 2018. F-2 screen,
inheritance and cross-resistance of field-derived Vip3A resistance in *Spodoptera frugiperda* (Lepidoptera:
Noctuidae) collected from Louisiana, USA. Pest Management Science 74: 1769-1778.

Yu, S. J. 1991. Insecticide Resistance in the Fall Armyworm, *Spodoptera frugiperda* (Smith,J.E.). Pestic Biochem
Phys 39: 84-91.

Zhang, J. P., F. Zhang, W. T. Tay, C. Robin, Y. Shi, F. Guan, Y. H. Yang, and Y. D. Wu. 2022. Population
genomics provides insights into lineage divergence and local adaptation within the cotton bollworm. Mol
Ecol Resour. https://doi.org/10.1111/1755-0998.13581.

Zhang, L., B. Liu, W. G. Zheng, C. H. Liu, D. N. Zhang, S. Y. Zhao, Z. Y. Li, P. J. Xu, K. Wilson, A. Withers, C. M.
Jones, J. A. Smith, G. Chipabika, D. L. Kachigamba, K. Nam, E. D'Alencon, B. Liu, X. Y. Liang, M. H. Jin, C.
Wu, S. Chakrabarty, X. M. Yang, Y. Y. Jiang, J. Liu, X. L. Liu, W. P. Quan, G. R. Wang, W. Fan, W. Q. Qian,
K. M. Wu, and Y. T. Xiao. 2020. Genetic structure and insecticide resistance characteristics of fall
armyworm populations invading China. Mol Ecol Resour. 20: 1682-1696.

Zhao, Y. X., J. M. Huang, H. Ni, D. Guo, F. X. Yang, X. Wang, S. F. Wu, and C. F. Gao. 2020. Susceptibility of fall
armyworm, *Spodoptera frugiperda* (J.E.Smmith), to eight insecticides in China, with special reference to
lambda-cyhalothrin. Pestic Biochem Phys. 168: 104623.

Table 1: List of insecticides and Bt toxins used in the bioassay experiments. All insecticides and Bt toxins were

tested on two populations of *Spodoptera frugiperda* (FAW) from Queensland (Sf20-1, from Walkamin) and

Western Australia (Sf20-4, from Kununurra). Commercial names of insecticides/Bt toxins, their manufactured

or expiry dates are provided where available. Bioassays involved either: (i) diet incorporation (DI); (ii) Topical

application (TA); or (iii) Surface treatment (ST) of diet to be consumed. The larval stages used in the bioassays

are specified. For topical application we used larvae from late 3^{rd} to early 4^{th} instar with each larva weighing

approximately 30mg. Species tested were: *Helicoverpa armigera, Spodoptera frugiperda* (FAW), and

Spodoptera litura.

Insecticides/Bt toxins	Commercial name	Date	Approach	Larval stage	Species tested		
					FAW	*H. armigera*	*S. litura*
Chlorantraniliprole (200g/L)	Coragen	Manufactured : 28/06/2019	DI	2^{nd}/early 3^{rd}	Yes	Yes	No
Indoxacarb (150g/L)	Steward 150	Manufactured : 30/03/2019	DI	2^{nd}/early 3^{rd}	Yes	Yes	No
Emamectin (17g/L)	Affirm 017 EC	Unspecified†	DI	2^{nd}/early 3^{rd}	Yes	Yes	No
Spinetoram (120g/L)	Dow AgroSciences Success® Neo	Manufactured : 02/2018	DI	2^{nd}/early 3^{rd}	Yes	Yes	No
Alpha cypermethrin (100g/L)	Dominex Duo	Manufactured : 30/04/2019	TA	Late 3^{rd}/ early 4^{th}	Yes	Yes	No
Methomyl		Expiry date: 02/2022	TA	Late 3^{rd}/ early 4^{th}	Yes	Yes	No
Advertised as: Cry2A, Cry2B, Cry1Ab, Cry1Aa, Cry1Ac	Yates Nature's Way DIPEL	Expiry date: 27/05/2022	ST	Neonates	Yes	Yes	Yes
Advertised as: Cry1D, Cry1C, Cry1Ab, Cry1Aa	XenTari WG Bta	Manufactured : 05/02/2019	ST	Neonates	Yes	Yes	Yes
Cry1Ac	CSIRO; from HD73 strain	NA	ST	Neonates	Yes	Yes	Yes
Cry2Ab	CSIRO; from Bt expressing Cry2Ab	NA	ST	Neonates	Yes	Yes	Yes
Cry1F	CSIRO; from *E. coli* expressing Cry1F	NA	ST	Neonates	Yes	Yes	Yes
VIP3A	CSIRO; from *E. coli* expressing Vip3A	NA	ST	Neonates	Yes	Yes	Yes

Note: † likely manufactured recently as this was requested from Syngenta Australia Pty Ltd and provided

directly for the bioassay purpose. N/A: not available. These Bt toxins were produced and purified from *B.*

thuringiensis cultures (Cry1Ac and Cry2Ab) or cloned into vectors in *Escherichia coli* (Cry1F and Vip3A) and

grown at the CSIRO Black Mountain Laboratories.

Table 2: *Spodoptera frugiperda* samples from Australian populations (Sf20-1, Qld; Sf20-4, WA), Papua New Guinea, South Korea, and Peru used in whole genome

sequencing for resistance gene characterisation. Number of specimens from each location is indicated by 'N'. Where available, GPS co-ordinates and elevation are

provided. The number of individuals belonging to the *S. frugiperda* R-strain or the C-strain mitochondrial DNA COI haplotypes is also provided.

Country	State/Province	Population	N	Date	Note	R : C
Australia	WA	Kununurra	9	16-Aug-20	'Sf20-4', represent the original WA field-collected individuals	9 : 0
	Qld	Strathmore Station, Walkamin	29	03-Mar-20	'Sf20-1', represent the original Qld field-collected individuals	29 : 0
South Korea	Milyang	MF	7	Feb-20	Milyang-si, Gyeongsangnam-do, G_4 lab, GPS: 35˚29.29 N, 128˚44.31 E	7 : 0
	Goryong	GR	1	Sep-19	Goryeong-gun, Gyeongsangbuk-do, Corn-field; GPS: 35˚ 64.04 N, 128˚39.08 E	1 : 0
	Haenam	MN	1	Aug-19	Haenam-gun, Jeollanam-do, Corn field; GPS: 34˚24.38 N, 126˚37.55 E	1 : 0
	Milyang	MY	2	Sep-19	Milyang-si, Gyeongsangnam-do, Corn field; GPS: 35˚29.29 N, 128˚44.31 E	0 : 2
	Muan	MA	1	Aug-19	Muan-gun, Jeollanam-do, Corn field; GPS: 34˚52.21 N, 126˚31.11 E	1 : 0
Papua New Guinea	Madang Province	Ramu Sugar Estate	16	15-17 Jun-2020	5°58.154 S, 145°53.252 E, Maize host	16 : 0
	Central Province	Yule Island Junction	1	5-Jun-20	corn host	1 : 0
Peru	n/a	Pre-border intercepted specimens on imported agricultural commodities	16	2016 - 2019	Samples from Tay et al. (2020) (Suppl. Table 1, individuals PE01 – PE16)	0 : 16

739 **Table 3:** Summary bioassay data on *Spodoptera frugiperda* populations from Walkamin, Queensland (Sf20-1) and Kununurra, Western Australia (Sf20-4), *S. litura* (Mareeba)

740 and *Helicoverpa armigera* (GR laboratory line) involving surface treatment of the diet with Bt toxins and products.

Pesticide	Populations/species	LC_{50}	95% CI	LC_{99}	Toxicity ratio (*H. armigera* = 1)	Slope
Cry1Ac	*H. armigera*	0.025	0.021 - 0.029	0.465	-	1.837 ±0.126
(μg/cm^2)	Sf20-1	4.34	1.81 - 8.15	2813.9	174	0.827 ±0.102
	Sf20-4	2.48	1.79 - 3.29	99.9	99	1.449 ±0.149
	S. litura	2.99	2.06 - 4.20	80.7	120	1.626 ±0.135
Cry2Ab	*H. armigera*	0.049	0.034 -0.068	1.74	-	1.501 ±0.143
(μg/cm^2)	Sf20-1	0.655	0.435 - 0.951	16.88	13	1.649 ±0.136
	Sf20-4	0.178	0.138 - 0.221	2.03	4	2.201 ±0.246
	S. litura	0.511	0.335 - 0.734	44.9	10	1.197 ±0.081
Cry1F	Sf20-1	0.025	0.015 - 0.038	0.838	-	1.531 ±0.115
(μl/cm^2)	Sf20-4	0.021	0.014 - 0.029	0.394	-	1.809 ±0.149
	S. litura	0.0088	0.0069 - 0.011	0.294	-	1.524 ±0.121
VIP3a	*H. armigera*	0.0062	0.0051 - 0.0075	0.23	-	1.486 ±0.093
(μl/cm^2)	Sf20-1	0.0021	0.0010 - 0.0031	0.035	0.152	1.881 ±0.225
	Sf20-4	0.0019	0.0016 - 0.0023	0.018	0.078	2.398 ±0.212
	S. litura	0.00065	0.0005 - 0.0008	0.015	0.065	1.693 ±0.129
Dipel	*H. armigera*	2.11	1.59 - 2.72	39.2	-	1.832 ±0.154
(IU/cm^2)	Sf20-1	52.02	36.55 - 70.05	2612.6	24	1.832 ±0.154
	Sf20-4	38.93	23.97 - 56.35	904.1	18	1.703 ±0.153
	S. litura	35.96	28.23 - 44.99	1540.8	17	1.426 ±0.098
XenTari	*H. armigera*	5.95	4.69 - 7.40	197.2	-	1.530 ±0.136
(DBM/cm^2)	Sf20-1	11.81	8.33 - 15.17	112.7	2	2.375 ±0.289
	Sf20-4	19.01	9.27 - 29.8	1058.9	3	1.333 ±0.180
	S. litura	24.86	17.08 - 31.14	206.9	4	2.375 ±0.289

743 **Table 4:** Summary bioassay data on *Spodoptera frugiperda* populations and *Helicoverpa armigera* (GR laboratory line) involving topical application to the insect with

744 contact insecticides. Sf20-1 = *S. frugiperda* from Walkamin Queensland; Sf20-4 = *S. frugiperda* from Kununurra Western Australia.

Toxin	Population/Species	LC$_{50}$	95% C.I.	LC$_{99}$	Toxicity ratio (*H. armigera* = 1)	Slope
Alpha cypermethrin	*H. armigera*	0.0036	0.0032 - 0.0041	0.023	-	2.849 ±0.140
µg/larvae	Sf20-1	0.201	0.171 - 0.239	1.88	56	2.399 ±0.154
	Sf20-4	0.523	0.427 - 0.641	6.06	145	2.186 ±0.132
Methomyl	*H. armigera*	0.057	0.031 - 0.097	43.12	-	0.809 ±0.059
µg/larvae	Sf20-1	0.254	0.177 - 0.356	39.2	4	1.064 ±0.076
	Sf20-4	2.96	1.87 - 4.66	1363.7	52	0.874 ±0.07

746 **Table 5:** Summary bioassay data on *Spodoptera frugiperda* populations (Sf20-1 from Walkamin Queensland; Sf20-4 from Kununurra Western Australia) and *Helicoverpa*

747 *armigera* (GR laboratory line) involving diet incorporation of insecticides.

Pesticide	Population/Species	LC$_{50}$	95% CI	LC$_{99}$	Toxicity ratio (*H. armigera* = 1)	Slope
Chlorantraniliprole	*H. armigera*	0.011	0.009 -0.013	0.056	-	3.199 ±0.234
(µg/ml diet)	Sf20-1	0.032	0.024 - 0.043	0.28	3	2.484 ±0.191
	Sf20-4	0.163	0.132 - 0.201	2.19	15	2.065 ±0.152
Indoxacarb	*H. armigera*	0.054	0.028 - 0.089	1.29	-	1.684 ±0.122
(µg/ml diet)	Sf20-1	1.203	1.031 - 1.398	11.66	22	2.359 ±0.159
	Sf20-4	11.206	9.254 - 13.654	213.85	208	1.817 ±0.130
Emamectin	*H. armigera*	0.0107	0.0087 - 0.0131	0.0945	-	2.465 ±0.164
(µg/ml diet)	Sf20-1	0.0158	0.013 - 0.019	0.124	1	2.606 ±0.175
	Sf20-4	0.0179	0.016 - 0.020	0.094	2	3.234 ±0.226
Spinetoram	*H. armigera*	0.086	0.023 - 0.157	1.977	-	1.711 ±0.173
(µg/ml diet)	Sf20-1	0.118	0.101 - 0.137	0.623	1	3.220 ±0.198
	Sf20-4	0.102	0.092 - 0.112	0.301	1	4.921 ±0.489

750 **Table 6:** Number of *Spodoptera frugiperda* individuals (N) from populations from Australia (Kununurra SF20-4; Strathmore SF20-1), South Korea, Papua New Guinea, and

751 Peru with homozygous susceptible (S/S), heterozygous (S/R) and homozygous resistance (R/R) profiles characterised via whole genome sequence data of the ACE-1 locus

752 involving the A2001S, G227A, and F290V amino acid substitutions.

Population	N	ACE-1 (A201S)			ACE-1 (G227A)			ACE-1 (F290V)		
		S/S	S/R	R/R	S/S	S/R	R/R	S/S	S/R	R/R
Kununnura	9	9	0	0	9	0	0	2	4	3
Strathmore	30	26	4	0	30	0	0	0	0	30
South Korea	12	11	1	0	12	0	0	0	0	12
Papua New Guinea	17	15	2	0	17	0	0	1	12	4
Peru	16	4	12	0	10	6	0	2	14	0

Table 7: Comparisons of bioassay results for selected insecticides fed to laboratory populations of *Spodoptera*
frugiperda via diet incorporation approaches. *S. frugiperda* populations were from the native ranges (USA,
Louisiana State University (LSU) culture; see Hardke et al. 2011); Brazil SUS-2005, (Kulye et al. 2021) and
introduced ranges from the original 2018 populations from Kanartaka, India (population I-2018; Kulye et al.
2021); a population from Groblersdal, Mpumalanga province, South Africa (see Discussion above, Eriksson
2019), and from our Sf20-1 and Sf20-4 laboratory populations from Queensland and Western Australia,
Australia, respectively. Potency ratios of the invasive populations were measured against the native Brazilian
(SUS-2005) population as reported by Kulye et al. (2021). Diet incorporation approach was for the Brazil, USA
(LSU), India I-2018, and Australia (Sf20-1; Sf20-4) populations.

Insecticides	Populations	Range	LC_{50} (ppm) / EC_{50} (µg/mL)	Fiducial limits† (95%)	Slope ± SE	Toxicity ratios
Spinetoram	Brazil (SUS-2005)[#]	native	0.010	0.009 - 0.012	2.35 ± 0.37	-
	USA (LSU)	native	0.066	0.053 - 0.081	2.54 ± 0.36	6.6
	India (I-2018)[#]	invasive	0.013	0.012 - 0.014	2.25 ± 0.13	1.3
	Sf20-1 (Qld)	invasive	0.118	0.101 - 0.137	3.220 ±0.198	11. 8
	Sf20-4 (WA)	invasive	0.102	0.092 - 0.112	4.921 ±0.489	10.2
Chlorantraniliprole	Brazil[#]	native	0.005	0.005 - 0.006	4.54 ± 0.76	-
	USA (LSU)	native	0.068	0.317 - 0.481	2.55 ± 0.23	13.6
	India (I-2018)[#]	invasive	0.009	0.008 - 0.009	2.39 ± 0.11	1.8
	South Africa	invasive	0.06	0.01 - 0.16	1.01 ± 0.42	12
	Sf20-1 (Qld)	invasive	0.032	0.024 - 0.043	2.484 ± 0.191	6.4
	Sf20-4 (WA)	invasive	0.163	0.132 - 0.201	2.065 ±0.152	32.6
Indoxacarb	USA (LSU)	native	0.392	0.317 - 0.481	2.35 ± 0.25	-
	Sf20-1 (Qld)	invasive	1.203	1.031 - 1.398	2.359 ±0.159	3.07
	Sf20-4 (WA)	invasive	11.206	9.254 - 13.654	1.817 ±0.130	28.6

Note: †Fiducial limits are similar to confidence limits but are based on logistic growth or S-shaped curve
distribution. #EC_{50} reported instead of LD_{50} by Kulye et al. (2021). In this comparison we assumed both EC_{50}
and LD_{50} estimates to be broadly similar.

Table 8: Comparisons of topically applied methomyl bioassay results on late 3rd and/or early 4th instar larvae of
FAW between resistant or susceptible populations from Gainesville Florida, USA (FL) (Yu 1991; see Discussion
above), and invasive populations from South Africa (Eriksson 2019; see Discussion above) and Queensland or
Western Australia, Australia.

Populations	LC$_{50}$ (ppm)	Fiducial limits (95%)	Slope ± SE	Toxicity ratios
USA (FL susceptible)	0.57	0.45 - 0.78	4.9	-
South Africa	62.98	17.22 – 52.12	11.14 ± 9.20	110.5
Sf20-1 (Qld)	0.254	0.177 - 0.356	1.064 ±0.076	0.45
Sf20-4 (WA)	2.96	1.87 - 4.66	0.874 ±0.07	5.2
USA (FL resistance)	8.23	4.86 - 11.6	2.9	-
South Africa	62.98	17.22 – 52.12	11.14 ± 9.20	7.65
Sf20-1 (Qld)	0.254	0.177 - 0.356	1.064 ±0.076	0.031
Sf20-4 (WA)	2.96	1.87 - 4.66	0.874 ±0.07	0.36

Figure 1 caption

Figure 1: Summary of Acytylcholinesterase (ACE-1) susceptible and resistance allele frequencies in *Spodoptera*

frugiperda populations from 17 countries (i.e., Benin, Uganda, Kenya, Tanzania, Zambia, Malawi, India,

Indonesia, Papua New Guinea (PNG), South Korea, China, Australia) across the invasive range. A total of 1,246

individuals representing 76 populations in the Table S2 were used to compile the data below. Population

identity is as provided in Table S2 that combined data from this study (Australia SF20-1 and SF20-4 (generation

0 representing field-collected individuals), PNG, South Korea) and from published studies (Boaventura et al.

2020a, Guan et al. 2020, Zhang et al. 2020, Zhao et al. 2020, Nguyen et al. 2021, Tay et al. 2021, Yainna et al.

2021, Rane et al. 2022). Susceptible and resistant alleles from the three previously reported loci (i.e., F290V,

G227A, A201S) from the ACE-1 gene provided evidence to support multiple independent introductions across

the invasive FAW populations, such as in Indonesia (#17; G227A), South Korea (#20; F290V), and China (Hubei

province (#10-13); G227A) as indicated by the red arrows. In Australia, newly established FAW populations

between Queensland (e.g., #27, #28 (Walkamin), #29 (Strathmore), #32 (Burdekin), and Western Australia (i.e.,

#2 (Kununurra)/Northern Territory (e.g., #21 (Bluey's Farm)) suggested this likely involved multiple

introductions from diverse populations from neighbouring countries and likely arrived via separate pathways

and entry points.

Figure 1