Resistance bioassays and allele characterisation inform analysis of *Spodoptera frugiperda*

(Lepidoptera: Noctuidae) introduction pathways in Asia and Australia

Tay WT[1,3], Rane RV[2,3], James W[1], Gordon KHJ[1], Downes S[4], Kim J[5], Kuniata L[6], Walsh TK[1,3]

1. CSIRO Black Mountain Laboratories, Clunies Ross Street, ACT 2601, Australia

2. CSIRO, 343 Royal Parade, Parkville, VIC 3052, Australia

3. Applied BioSciences, Macquarie University, Sydney NSW 2100, Australia

4. CSIRO McMaster Laboratories, New England Highway, Armidale NSW 2350, Australia

4. College of Agriculture and Life Science, Kangwon National University, Republic of Korea

5. Ramu Agri Industries Ltd., PNG

Correspondence: weetek.tay@csiro.au

Abstract

The fall armyworm (FAW) *Spodoptera frugiperda* is present in over 70 countries in Africa, Asia, and Oceania. Its

rapid dispersal since 2016 when it was first reported in western Africa, and associated devastation to

agricultural productivity, highlight the challenges posed by this pest. Currently, its management largely relies

on insecticide sprays and transgenic *Bacillus thuringiensis* toxins, therefore understanding their responses to

these agents and characteristics of any resistance genes enables adaptive strategies. In Australia, *S. frugiperda*

was reported at the end of January 2020 in northern Queensland and by March 2020, also in northern

Western Australia. As an urgent first response we undertook bioassays on two Australian populations, one

each from these initial points of establishment. To assist with preliminary sensitivity assessment, two endemic

noctuid pest species, *Helicoverpa armigera* and *Spodoptera litura*, were concurrently screened to obtain larval

LC50 estimates against various insecticides. We characterised known resistance alleles from the VGSC, ACE-1,

RyR, and ABCC2 genes to compare with published allele frequencies and bioassay responses from native and

invasive *S. frugiperda* populations. An approximately 10x LC50 difference for indoxacarb was detected

between Australian populations, which was approximately 28x higher than that reported from an Indian

population. Characterisation of ACE-1 and VGSC alleles provided further evidence of multiple introductions in

Asia, and multiple pathways involving genetically distinct individuals into Australia. The preliminary bioassay

results and resistance allele patterns from invasive *S. frugiperda* populations suggest multiple introductions

have contributed to the pest's spread and challenge the axiom of its rapid 'west-to-east' spread.

Keywords

FAW, whole genome sequencing, agricultural biosecurity, population genomics, invasion biology

Introduction

The fall armyworm (FAW) *Spodoptera frugiperda* (J. E. Smith), is a noctuid moth pest made up of two morphologically indistinguishable strains (C- and R-strains) that is native to tropical and subtropical regions of the Americas. It is highly polyphagous and feeds on host species from at least 76 plant families, principally Poaceae (106 spp.), Asteraceae (31) and Fabaceae (31) (Montezano et al. 2018). In the Americas, it causes significant economic damage to maize, rice, sorghum, millet, soya bean, wheat, alfalfa, cotton, turf, and fodder crops (Nagoshi et al. 2019a). Recognised globally as one of the top 20 arthropod pests (Willis 2017), *S. frugiperda* was first officially confirmed outside of its native range in Western Africa in early 2016 (Goergen et al. 2016) and then detected across virtually all of Sub-Saharan Africa by February 2018 (Nagoshi et al. 2018, Otim et al. 2018, Botha et al. 2019). In July 2018, it was confirmed in Yemen and India, and by December 2018 in Bangladesh, Sri Lanka, Myanmar and Thailand, followed closely by China in January 2019. Nepal confirmed its presence in May 2019, and in July 2019 it was also reported in Egypt, Indonesia, Laos, Cambodia, Malaysia, Vietnam, Taiwan, The Republic of Korea, and Japan (FAO 2021). The moth's strong flight ability (Rose et al. 1975, Sparks 1979, Jones et al. 2019), its potential to contaminate certain commodities, and ability to act as a 'hitchhiker' in trade (Early et al. 2018) contribute to its rapid spread. Whole genome analyses of invasive *S. frugiperda* populations from various African nations, India, China, Southeast Asia, and Australia also suggested that a high proportion of *S. frugiperda* individuals were hybrids of the C- and R-strains (Gui et al. 2020, Tay et al. 2020, Zhang et al. 2020, Schlum et al. 2021, Tay et al. 2021, Rane et al. 2022).

Much of South and Southeast Asia, and much of Northern Australia, are highly suitable climatically for *S. frugiperda* year-round (du Plessis et al. 2018, Early et al. 2018). It was confirmed in Queensland (Qld; Bamaga Northern Peninsula Area), Australia following clearance of lure traps on 31 January 2020, followed by reports also in north-Western Australia (WA; Kununurra) and in the Northern Territory (NT) in March 2020. Within two weeks of the Bamaga detection, there were reports of *S. frugiperda* damaging maize fields in Strathmore (Qld; *ca.* 1,000 km from the initial Bamaga detection site), followed by successive southward detections of *S. frugiperda* in Qld, marking the FAW's south-eastward antipodean expansion into New South Wales (NSW) in September 2020. Through CLIMEX simulation analysis based on irrigation patterns and rainfall records, *S. frugiperda* is expected to undertake seasonal migrations and may reach as far south as Victoria and Tasmania (du Plessis et al. 2018). Indeed, *S. frugiperda* was reported in Victoria in December 2020 and in Tasmania in April, 2021. Concurrently, and from the Western Australia Kununurra detection, *S. frugiperda* followed the CLIMEX simulated southward expansion patterns along the WA's western coast, with larval populations detected in the Gingin area (< 80 Km north of the city of Perth) by February 2021.

The incursion of *S. frugiperda* into Australia is believed to have occurred through natural dispersal across the Torres Strait/Timor Sea into northern Australia via a single-entry point (Jing et al. 2021, Qi et al. 2021). However, incursion pathways to Australia could also involve multiple entry points as reported for the biting midges *Culicoides bravitarsis* (Tay et al. 2016), other anthropogenic routes such as infested commodities in trade, as well as through seasonal or on-going new arrivals of individuals from populations that have successfully established in Asia and Southeast Asia (SEA). The eastward spread of *S. frugiperda* across sub-

Saharan Africa, the Middle East, the Indian sub-continent, SEA, China, and the Far East (South Korea, Japan)

mostly followed chronological detection and confirmation which led to a widely accepted belief that an

invasive population established in west Africa (Goergen et al. 2016) through a single founder, similar to the

'invasive bridgehead effect' (Guillemaud et al. 2011), was the starting point for its subsequent global

expansion. This hypothesis is supported by an overwhelmingly homogeneous genetic signature based on single

gene markers (e.g., Cock et al. 2017, Nagoshi et al. 2018).

Whole genome analysis and genome-wide single nucleotide polymorphic marker analysis of invasive

populations from Africa (Benin, Uganda, Malawi, Tanzania) and Asia (India, China) instead suggest multiple

founding events (Tay et al. 2020, Zhang et al. 2020, Schlum et al. 2021); a similar conclusion reached also from

analysis of multiple *S. frugiperda* populations from across China based on microsatellite DNA markers (Jiang et

al. 2022). Recently, population genomic analysis of *S. frugiperda* populations from SEA (i.e., Philippines,

Vietnam, Laos, Malaysia, Myanmar), from East Asia (i.e., South Korea), and Pacific/Oceania (i.e., Papua New

Guinea; Australia) identified significant genetic diversity in these invasive *S. frugiperda* populations, and

distinct population signatures between populations in close proximity (e.g., Kedah and Penang populations

from Malaysia, and between Qld, WA and NT populations in Australia (Rane et al. 2022); between populations

within e.g., Yunnan province (Tay et al. 2022), and populations from Anhui and Jiangsu (Jiang et al. 2022)).

Such distinct population genomic structure contradicted the expected gene flow signatures of a single

introduction and a west-to-east spread of this pest. Instead, they suggested relatively limited and localised

spread of populations while also identified the likely multiple independent introduction pathways of *S.*

frugiperda into the region including in SEA, Asia and Australia (Rane et al. 2022).

Distinguishing among single vs. multiple introductions within Australia (and elsewhere), and understanding the

resistance profiles of existing and any new incursions, is therefore critical for informing the future

management of this pest. For example, this information will assist with forecasting likely resistance profiles of

invasive populations selected outside of Australia, and could prioritise efforts to prevent the introgression

(potentially of multiple separate introductions) of known insecticide resistance genes and alleles from endemic

populations (e.g., Carvalho et al. 2013, Banerjee et al. 2017, Boaventura et al. 2020a, Guan et al. 2020) into

invasive populations, or through reciprocal introgression of newly selected/developed resistance traits from

invasive populations to native populations.

In this study, we report on the first bioassay experiments and resistance gene characterisation that aimed to

understand how the initial invasive populations of *S. frugiperda* in Australia responded to selected approved

insecticide compounds and *Bacillus thuringiensis* (Bt) toxins in commercial transgenic plant varieties and

available as sprays for the horticultural, grains and cotton industries. Since populations were presumed

collected before selection occurred in Australia, these responses served as the first indication of selection

against insecticides and Bt toxins used elsewhere (i.e., in the species' native range and in its recent invasive

ranges from Southeast Asia, Asia, and/or Africa (Schlum et al. 2021; Tay et al. 2022; Jiang et al. 2022; Rane et

al. 2022)) prior to arrival. While sampling from multiple Australian regions would be ideal (but impossible at

the time as our laboratory had only been provided with these two lines during the pandemic travel

restriction), comparing the response differences between these two *S. frugiperda* populations from different
northern Australia regions nevertheless could demonstrate the use of bioassay data to distinguish signatures
of possible multiple points of entry compared with a single entry and spread event as generally assumed (e.g.,
Jing et al. 2021, Qi et al. 2021). To provide a first insight into the efficacy of key compounds and toxins we
compared the responses of *S. frugiperda* with two related major endemic crop pests in Australia, *Helicoverpa*
armigera (subspecies *conferta;* Hardwick 1965, Anderson et al. 2016, Pearce et al. 2017b, a, Zhang et al. 2022)
and *S. litura*, which are currently managed through adaptive resistance management plans. Finally, we
provided a critical review of resistance profiles in selected invasive populations of *S. frugiperda* globally to
better understand its invasion biology, and consider our work alongside a subsequent more comprehensive
examination in Australia of insecticide resistance profiles (Bird et al. 2022). Our study is the only available
information for Australian populations on the vulnerability of new arrived *S. frugiperda* to relevant Bt toxins.

Material and Methods

Live insects

Live FAW populations were sourced from Queensland (Qld) and Western Australia (WA). This population
(CSIRO code: 'Sf20-1' from Qld; see Table 2) consisted of 30 field-collected pupae (of which 29 pupated and
emerged as adults) for setting up the laboratory colony, was first noticed around 19[th] Feb, 2020, was collected
on 3[rd] March, 2020, and represented the first reported FAW detected attacking maize in Australia's agricultural
landscape. The sampling site for this population was from the University of Queensland (UQ) field station (Rex
Road, Walkamin Qld 4872) Strathmore Station, (Lat/Lon: -17.17892, 145.43359, Elevation 685m).

A second FAW population (CSIRO code: 'Sf20-4' from WA; see Table 2) was collected as larvae from a maize
crop from the Northern Australian Crop Research Alliance Pty Ltd, Kununurra, Western Australia. These larvae
were collected on 17-Aug, 2020 from the Kununurra Airport field trail block (off Victoria Highway; -15.10090,
128.81342) on V4-6 growth stage maize, prior to applications of insecticides. A total of 30 larvae of different
instar stages arrived (= F_0), and 10 (six females and four males) survived to pupate and emerge as adult moths
to produce >200 F_1 individuals and to generate sufficient larvae to commence bioassays from F_4.

Two related endemic noctuid species, *S. litura* and *Helicoverpa armigera conferta* (hereafter '*H. armigera*'),
that are pests of a range of broadacre cropping and horticulture industries (and especially grains and cotton)
were also included in the bioassay experiments to assist with interpretating results such as base line
susceptibility and tolerance levels to the chemicals and toxins tested. Approximately 30 *S. litura* larvae with
unknown resistance profiles (from Mareeba and Toowoomba, Qld) pupated and were used to start the
laboratory colony. The *H. armigera* colony (CSIRO general rearing 'GR' strain) is a laboratory strain housed at
the Black Mountain Laboratory in the ACT that was established during the mid-1980's with individuals
originally collected as eggs from cotton fields around Narrabri NSW. Initially the GR colony was periodically
replenished to sustain population genetic diversity and minimise inbreeding but no new material has been
introduced for around two decades and the colony is susceptible to many chemical insecticides, including
organochlorines, organophosphates, carbamates and pyrethroids, as well as to the Cry1Ab, Cry1Ac, and

Cry2Ab Bt toxins (Pearce et al. 2017a, Pearce et al. 2017b). It therefore has tolerance levels that are not
dissimilar to those of a progenitor population created in 2011 that was deliberately maintained as susceptible
through screening (see Bird et al. 2021).The *H. armigera* GR population was confirmed by bioassays as having
no Cry1Ac/Cry2Ab/VIP3A resistance alleles. The field collected *S. frugiperda* strains were tested within 4
generations of establishment in the laboratory.

Colony maintenance

Helicoverpa armigera and *Spodoptera* species (*S. frugiperda* Sf20-1 and Sf20-4; *S. litura*) were reared by the
method of Teakle and Jensen (1985) at 25 °C, 50 ± 10% relative humidity, and with a light/dark cycle of 14:10
h. Artificial diet for *H. armigera* consisted of 81 g soya flour, 37.5 g wheat germ, 33 g brewer's yeast, 2 g
ascorbic acid, 2 g nipagen, 3 mL vegetable oil, and 13.5 g agar, which was processed in a microwave oven and
made up with water to 1 litre. Diet was poured into rearing cups or bioassay plates, allowed to cool and then
stored in the refrigerator (4-8°C) for no longer than 3 days. Diet for *S. frugiperda* was made to 1200mL with
water, and contained 100g navy bean flour, 25g soy flour, 60g wheat germ, 30g Brewer's yeast, 15g casein, 3g
Nipagen, 1.5g sorbic acid, 10g vitamin mixture, 4mL vegetable oil, and 18g agar. The mixture was processed in
a microwave oven prior to being poured into rearing cups/bioassay plates and allowed to cool as described
above for the *H. armigera* diet.

Bioassays

The bioassay experiments involved commercially available insecticidal compounds and Bt toxins (available in
transgenic varieties of cotton and as commercial sprays), as well as Bt toxins that were produced and purified
by CSIRO (Table 1). For all insecticides and Bt toxins, Sf20-1 and Sf20-4 *S. frugiperda* populations were tested.
The *H. armigera* 'GR' line was tested alongside *S. frugiperda* against all chemicals and Bt toxins except Cry1F
because it is known to be insensitive to it. We included *S. litura* in all Bt toxin bioassays as it is regarded as a
pest of interest by the Australian cotton industry who depend on Bt varieties.

171   [Table 1]

All bioassays included a negative control for background mortality (i.e., to enable natural mortality rates to be
adjusted accordingly) which was treated with the diluent of the individual pesticide. An initial experiment
using 10x serial dilutions was performed to narrow the range for the detailed bioassay using 8-16 individuals
per dose replicated once. The full bioassay involved 2x serial dilutions at 6-7 doses with 16-24 individuals per
dose replicated 3-5 times. The bioassays involved diet incorporation, surface overlay, or topical application of
the insecticidal compounds or Bt toxins (Table 1). The selection among these exposure methods for an
insecticide was determined by its mode of entry (contact or ingestion) and the experience of similar bioassays
in related noctuid pests (i.e., *H. armigera* and *H. punctigera*; Bird 2015; Bird & Walker 2019).

The surface overlay of Bt toxin assays followed the approach outlined in Mahon et al. (2007b), Mahon et al.
(2008), Mahon et al. (2010), Mahon et al. (2012) and Walsh et al. (2014). They were conducted in 96-well trays

where each well contained approximately 300µl of rearing diet with a surface area of around 0.567 cm^2 per

well. When the diet cooled, 20µl solution containing an appropriate concentration of toxin was added and

allowed to air dry. One neonate was added to each well and the tray was heat sealed with a perforated lidding

material.

For diet incorporation, commercial grade insecticides were diluted to the appropriate concentration with

water, added to 150ml of artificial diet and mixed well to produce a homogenous solution. Insecticide-

incorporated diet was then dispensed into 45-well bioassay trays, each with approximately 1.5 ml of diet-

insecticide mix. One late-second or early-third instar larva was added to each well and the tray was heat

sealed with a perforated lidding material (as per Bird 2015).

Topical bioassays of known concentrations were conducted for alpha-cypermethrin (group 3A) and methomyl

(group 1) and involved applying drops of pesticide as described in Bird (2018). Specifically, 1 µl of

acetone/insecticide solution was applied to the dorsal thorax of 3rd to 4th instar larvae (30-40g) using a 50 µl

micro-syringe. Bioassay trays were incubated at 25 ˚C, 45-55% RH, and a photoperiod 14:10 (L:D) h for 6 days

(e.g., see Bird and Akhurst 2007, Downes et al. 2009). The numbers of dead (moribund; incapable of

movement when prodded) and alive larvae (capable of coordinated movement when prodded) were counted

and the instars of surviving larvae recorded. The LC$_{50}$ for each toxin was calculated from pooled raw data by

probit analysis using the POLO-PC program (LeOra Sorftware 1987).

Spodoptera frugiperda specimens for genomic analyses

Populations of *S. frugiperda* from South Korea (SK), Papua New Guinea (PNG), and Peru were included for

whole genome sequencing (Table 2). The methods for SK and PNG specimen preservation involved collection

from fields and placing larvae directly into high concentrations of ethanol (95-99.9%) to transport to the

laboratory where they were then stored at -20˚C and replaced with fresh ethanol after 24-48 hours. Peru

specimens were samples intercepted from Australian pre-border inspections of imported

agricultural/horticultural commodities between 2016 and 2019 (see Tay et al. 2020). Australian samples (Sf20-

1 from Qld, Sf20-4 from WA) were F$_0$ individuals and represented individuals obtained directly from fields.

Samples from SK, PNG, and Peru were sent to CSIRO and stored at -20˚C until DNA extraction.

DNA extraction and genome library preparation

DNA of individual *S. frugiperda* samples was extracted using the Qiagen DNA extraction kit and eluted in

200µL elution buffer following protocols as provided by the manufacturer (Qiagen, Hilden Germany). Genomic

DNA libraries for individual samples were prepared, quantified and sent for commercial sequencing by the

Australian Genome Research Facility (AGRF) in Melbourne, Victoria, Australia.

Processing of genome sequences

Genome sequencing data for individuals were trimmed to remove adapter sequences using trim_galore (v

0.6.6; https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) and aligned to the *S. frugiperda* rice

genome (v1.0) (Gouin et al. 2017) using bwa_mem2 (v2) (Vasimuddin et al. 2019). Duplicate alignments were

removed using SAMBLASTER (v 0.1.26) (Faust and Hall 2014) and sorting completed using SAMtools (v1.9) (Li

et al. 2009).

[Table 2]

Strain identification and resistance alleles characterisation by whole genome sequencing

For identification of *S. frugiperda* R- or C-strain, the used the partial mtCOI gene sequences for the R-strain

(GenBank MF197867) and the C-strain (GenBank MF197868) of Otim et al. (2018) as reference sequence for

mapping against the whole genome sequence data for each individual *S. frugiperda* from Sf20-1, Sf20-4, SK,

PNG, and Peru. Mapping of partial mtCOI gene was carried out within Geneious v11.1.5 (Biomatters Ltd,

Auckland, NZ) using the Geneious Mapper program with assembly parameters specified to 'Low Sensitivity /

Fastest', with no trimming before mapping, and the 'Fine Tuning' option set to 'Iterate 2 times' to map reads

to the consensus from the previous iteration. Full mitochondrial DNA genomes of all individuals were also

assembled following the procedures as outlined in (Otim et al. 2018) and have been reported by Rane et al.

(2022). For all resistance alleles of interest (ABCC2, ACE-1, RyR, VGSC; see Table S1), base-sequence at the

genomic location for individuals listed in Table 1 was extracted from the alignment file using BCFTOOLS

MPILEUP followed by CALL, since BBMAP does not report 'non-variant' sites. Results were tabulated for

inference.

Comparisons of published ACE-1 and VGSC resistance allele frequencies and published indoxacarb and

chlorantraniliprole LC_{50} values

Three resistance loci for the organophosphate/carbamate ACE-1 gene have been reported to-date in *S*

frugiperda: (i) A201S, (ii) G227A, and (iii) F290V. To understand the frequencies of these alleles, we surveyed

specimens from invasive (i.e., Australia, PNG, South Korea) and native (Peru) *S. frugiperda* populations, and

combined this information with reported allele frequencies for these three loci from other native (Brazil,

French Guiana, Mexico Guadeloupe, Puerto Rico, USA) and invasive populations (Benin, Uganda, Kenya,

Tanzania, Zambia, Malawi; India, Myanmar, Laos, Vietnam, Malaysia, Indonesia, China, Australia) (see Table S2

and references therein). For the pyrethroid *para* sodium channel resistance gene VGSC, three loci have been

identified to date in *S. frugiperda*: (Carvalho et al. 2013, Guan et al. 2020, Yainna et al. 2021).⬚Carvalho et al.

2013, Guan et al. 2020, Yainna et al. 2021)⬚. Resistance and susceptible allele frequencies in these three VGSC

loci were also surveyed from published studies from native and invasive *S. frugiperda* populations (see Table

S2).

For comparisons of bioassay LC_{50} data for indoxacarb and chlorantraniliprole between published data and our

study, we considered only those that reported broadly similar methodologies (see Table 1) such as clearly

stated route of delivery (i.e., diet incorporation/ingestion) for the insecticidal compounds, and using

comparable developmental stages of larvae (i.e., 2nd/early 3rd instar) and scoring criteria.

Results

Bioassays of Individual Bt proteins

Cry1Ac was not highly effective against the *Spodoptera* species tested in this study (Fig. S1; Table 3). Based on the LC_{50}, and relative to *H. armigera*, *S. frugiperda* Sf20-1 was 174x less sensitive, *S. frugiperda* Sf20-4 was 99x less sensitive, and *S. litura* was 120x less sensitive. This suggests that in *S. frugiperda*, Cry1Ac would give similar control in Australia to that found in *S. litura* but far less than for *H. armigera*.

The *Spodoptera* species are less sensitive to Cry2Ab than *H. armigera* but the differences are not as striking as for Cry1Ac (Fig. S2; Table 3); FAW Sf20-1 was 13x less sensitive, *S. frugiperda* Sf20-4 was 4x less sensitive and *S. litura* was 10x less sensitive. This suggests that in *S. frugiperda*, Cry2Ab would give similar control in Australia to that found in *S. litura* which is unlikely to be different from *H. armigera* in terms of field control.

Cry1F is a Bt protein that has been deployed in certain genetically modified plants to target *Spodoptera*. Though not effective against *H. armigera sensu lato* even at a relatively high dose, Cry1F was effective against *S. frugiperda* at a much lower dose (Fig S3; Table 3), indicating that this species is much more sensitive to this Bt protein than *H. armigera*. Relative to *S. litura*, the LC_{50} data shows that *S. frugiperda* Sf20-1 and *S. frugiperda* Sf20-4 are 2x less sensitive to Cry1F. In terms of field control there is unlikely to be any distinguishable difference in Cry1F efficacy against *S. frugiperda* vs. *S. litura*.

FAW Sf20-1 and FAW Sf20-4 showed a similar tolerance to Vip3A as *S. litura* (~2.34x and 1.2x higher, respectively) and all three populations were more sensitive to Vip3A than *H. armigera* (0.15x, 0.08x, and 0.07, respectively). In terms of field control there is unlikely to be any distinguishable difference in efficacy of VIP3A against FAW relative to *H. armigera* and *S. litura* (Fig. S4; Table 3).

Based on published studies which show resistance ratios of >2,500 fold in *H. armigera* and FAW that are homozygous for recessive resistance alleles against Bt toxins (e.g., Mahon et al. (2007a) - Cry2Ab, *H. armigera*; Horikoshi et al. (2016), Cry1F, Cry1A, Vip3A, *S. frugiperda*) it is unlikely that either of the founding *S. frugiperda* populations in Australia carried resistance alleles to Cry1Ac, Cry2Ab or Vip3A in homozygous states.

Foliar Bt

The performance against FAW of the sprayable products containing multiple Bt toxins (XenTari and DIPEL) showed both products to differ in efficacies against *H. armigera*, *S. frugiperda*, and *S. litura*. XenTari with the mixture of Cry1 toxins including Cry1C was formulated to provide *Spodoptera* control (valentbiosciences, last accessed 13-Jan, 2022) but also for cabbage looper and the diamondback moth (DBM), while DIPEL was formulated for broad spectrum caterpillar control (valentbiosciences, last accessed 13-Jan, 2022). Both DIPEL and XenTari are reported to contain different toxin complements but the relative amounts are unclear. DIPEL contains Cry1Ac and Cry2-type toxins and was less effective against both tested FAW populations than XenTari which contains Cry1Aa, Cry1Ab, Cry1C and Cry1D. Relative to *H. armigera*, *S. frugiperda* Sf20-1 was 24x less sensitive, *S. frugiperda* Sf20-4 was 18x less sensitive and *S. litura* was 17x less sensitive to Dipel. XenTari

toxicity ratios relative to *H. armigera* ranged 2 to 4x for *S. frugiperda* Sf20-1, *S. frugiperda* Sf20-4 and *S. litura*

(Figs. S5 and S6; Table 3).

<u>Conventional pesticides</u>

Cypermethrin is a pyrethroid pesticide and the bioassay results suggest that *S. frugiperda* (i.e., the Sf20-1 and

Sf20-4 laboratory cultures) is less sensitive to it than the laboratory strain of *H. armigera* (56x and 145x

respectively: Fig. S7; Table 4).

Bioassay results for the carbamate pesticide methomyl are not as repeatable as other toxins with considerable

variability between replicates (Fig. S8; Table 4). However, overall *S. frugiperda* Sf20-4 shows 52x less sensitivity

to methomyl than *H. armigera* while *S. frugiperda* Sf20-1 shows a 4x lower sensitivity. This difference is

consistent with some level of heterogeneity between the first invasive *S. frugiperda* populations in Australia.

The dose response and LC_{50} for indoxacarb (Fig. S9; Table 5) show that *S. frugiperda* Sf20-1 and *S. frugiperda*

Sf20-4 are less sensitive to this chemistry than *H. armigera* (22x and 208x respectively). This difference is

consistent with some level of heterogeneity between the first invasive *S. frugiperda* populations in Australia.

The dose response and LC_{50} for chlorantraniliprole show similar sensitivity for *H. armigera* and *S. frugiperda*

Sf20-1, with the latter showing a 3x difference in the LC_{50}. Interestingly, the *S. frugiperda* Sf20-4 exhibited a

15x reduction in sensitivity compared to *H. armigera* (Fig. S10; Table 5), and a 5x response variation to

chlorantraniliprole when compared to the Sf20-1 line.

The dose response and LC_{50} data for emamectin benzoate and spinetoram (Figs. S11, S12; Table 5) suggest that

S. frugiperda Sf20-1 and FAW Sf20-4 are not significantly different in their level of sensitivity compared to *H.*

armigera.

<div align="center">[Table 3, Table 4, Table 5]</div>

Resistance alleles by whole genome sequencing

The loci examined for potential resistance alleles were present in the data set with the appropriate level of

coverage to accurately call the genotype. Resistance alleles to carbamate/organophosphates (ACE-1) were the

only resistance alleles identified in the Kununnura, Strathmore, South Korea, Papua New Guinea, and Peru

populations. No resistance alleles associated with target site mutations were detected for pyrethroid (VGSC) or

for diamide insecticides (RyR), although there are likely other genes (e.g., detoxification genes) that confer

resistance to synthetic pyrethroid in these laboratory lines of *S. frugiperda* (Bird et al. 2022). Resistance allele

profile differences between the two Australian populations were evident between the A201S and the F290V

amino acid substitutions. Similar ACE-1 resistance allele profiles were detected between Strathmore and South

Korean populations, while the Papua New Guinea population shared ACE-1 allele profiles with Strathmore

population (A201S) and with Kununnura (F290V). The Peruvian FAW population was the only population that

have heterozygous individuals with resistance allele for G227A amino acid substitution (Table 6).

<div align="center">[Table 6]</div>

Resistance allele characterisation by sequencing approaches from this study and published studies for the ACE-1 gene is summarised in (Fig. 1, Table S2 and references therein). The most common resistance allele detected in the invasive and native population (456 individuals examined in total) was the F290V mutation with 66 homozygous and 222 heterozygous resistant genotypes detected. This mutation is a T to G single nucleotide polymorphism (SNP) that changes the codon encoding the amino acid from TTT to GTT leading to a phenylalanine (F) to valine (V) change in the protein sequence encoded by the ACE-1 gene. This mutation was present at all locations and evenly distributed between the invasive and native populations. In Australia, of the sequenced individuals (n = 146), 24.7% (n = 36) were heterozygous, and 8% (n = 12) were homozygous for the resistance allele. Heterozygous and homozygous individuals were found in populations from all four states (i.e., WA, NT, Queensland, and NSW) as well as from the single individual from Erub Island, suggesting it is common across Australia.

The second most common allele detected was A201S (456 individuals examined) with 90 heterozygous individuals detected. This mutation is a C to A SNP which changes the codon from GCG to TCG leading to an alanine (A) to serine (S) amino acid change. While less common, this mutation also appears to be in both the invasive and native range with no obvious pattern. In Australia, of the sequenced individuals (n = 146), 15.8% (n = 23) were heterozygous, and 0 were homozygous for the resistance allele. Heterozygous individuals were found in WA and Queensland, also suggesting it is common across Australia.

The G227A mutation was the least common (from 456 individuals) with 20 heterozygous individuals and 2 homozygous individuals detected. This mutation is a G to a C SNP which alters the codon from GGA to CGA encoding a glycine (G) to alanine (A) amino acid change. Interestingly G227A was only present in individuals from the native range (Brazil, USA, Puerto Rico, Peru) but absent in individuals from across the invasive range that were surveyed in this study and from the related studies of Tay et al. (2020) and Tay et al. (2021). When compiling allele frequencies for both the VGSC and ACE-1 resistant genes from this study and from published whole genome sequencing and targeted PCR/Sanger sequencing data (Fig. 1; see also Table S1), the rare G227A resistance allele was present only in the Indonesian (Boaventura et al. 2020a) and Hubei populations (Guo et al. 2020) but absent in African, Australian, and other Asian (e.g., Indian, Myanmar, etc.) populations, including populations from six other Chinese provinces.

No target site mutation alleles predicted to cause resistance to pyrethroids or the group 28 pesticides were detected in this work. However, while the previously identified resistance alleles (Bolzan et al. 2019, Boaventura et al. 2020b) were not detected, considerable variation was present in the RyR gene at the potential resistance loci. This should be further investigated in conjunction with bioassays to establish whether any of the variants could contribute to resistance.

ABCC2 resistance alleles in Australia, Papua New Guinea, South Korea, and Peru populations

None of the known and validated Cry1 resistance ABCC2 mutations were identified in the individuals sequenced for this work however, some variation from the reference was observed (Fig. S13). Numerous synonymous mutations and a total of 31 non-synonymous mutations were observed in the coding sequence of

the ABCC2 gene in all sequenced individuals. Several of these non-synonymous mutations were associated
with a common deletion and insertion motif in the first exon where an 11bp deletion and a 2 bp insertion
maintain the reading frame but replace and change several amino acids. Most of the other mutations are the
result of one (i.e., single nucleotide variant; SNV) or two (i.e., multiple nucleotide variant; MNV) nucleotide
changes. Several of these mutations are found in other assemblies of *S. frugiperda* and likely reflect natural
variation. An alternative explanation for the variation is that they are associated with the c-strain which is
thought to make up at least some of the genome of the invasive populations. Only one mutation, a 2 bp
deletion in one individual collected from the Burdekin in Queensland was identified that might cause a frame
shift mutation (see Supplemental Figures S14 and S15) as has been identified in Bt resistant individuals in
other studies. It was present as a heterozygote in the individual and has not been experimentally validated.

367   [Figure 1]

Discussion

In this study, we showed insecticide and Bt response differences in two of the first reported *S. frugiperda*
populations in Queensland (i.e., SF20-1) and Western Australia (Sf20-4) in Australia, with the Queensland
(Sf20-1) population being less tolerant to various insecticides such as methomyl, chlorantraniliprole, and
indoxacarb compared to the Western Australian (Sf20-4) population. On the other hand, the Queensland
population exhibited at least a two-fold higher tolerance to Cry1Ac, Cry2Ab, Cry1F and Vip3A Bt toxins than
the WA population. Characterisation of the ABCC2 resistance gene identified one *S. frugiperda* individual from
Queensland as potentially being heterozygous with a 2 bp deletion that could underpin Cry1F resistance,
although confirmation of the detected mutation and of the resistance phenotype is required. This could be
accomplished via PCR and Sanger sequencing (e.g., Guan et al. 2020) as well as via the CRISPR/Cas9 gene
editing approach (e.g., see Wang et al. 2017). The response differences to chemical insecticides and Bt toxins
of the two studied invasive *S. frugiperda* populations in Australia and their ACE-1 resistance allele profile
differences (Table 6) may reflect the diverse genetic composition across the pest's recent expanding range
(Tay et al. 2020, Schlum et al. 2021, Rane et al. 2022), and suggest that separate pathways were involved in
establishment of these Queensland and Western Australian populations. This is contrary to the current
postulation of a single introduction pathway for the arrival of *S. frugiperda* to Australia based on an
assumption (Jing et al. 2021) or reverse trajectory simulation (Qi et al. 2021) and highlights the importance of
harmonising simulation studies with genomic and phenotypic evidence.

With significant economic impacts on agriculture from *S. frugiperda* reported in over 80 countries (excluding
the New World native range) from Africa, Middle East, Asia, Southeast Asia, and Oceania, its response to
different insecticidal and Bt control agents is increasingly investigated at recently impacted localities (e.g.,
Worku and Ebabuye 2019, Deshmukh et al. 2020, Zhang et al. 2020, Kulye et al. 2021, Lv et al. 2021). However,
the diverse bioassay methods (approach, larval stage, scoring criteria) used in these studies complicate
meaningfully comparison of findings. Taking emamectin benzoate and indoxacarb as examples, Zhang et al.
(2022) used a topical application bioassay on 3^{rd} instar larvae, Deshmukh et al. (2020) used leaf-dip bioassays
on 2^{nd} instar larvae, while Hardke et al. (2011) used a diet-incorporation approach on 3^{rd} instar larvae.

Responses have been expressed as LD_{50} (e.g., Deshmukh et al. 2020) or as EC_{50} (e.g., Kulye et al. 2021), further

making it difficult to rigorously compare outcomes.

It is important to stress the challenge and difficulty to compare bioassay results between studies due to the

different genetic background of the test samples (e.g., due to different number of individuals used to establish

test populations), methods and approaches between research groups, and the general different rearing

conditions of laboratory cultures that could contribute to varied response outcomes. A further challenge for

invasive *S. frugiperda* management is the lack of base-line values representing susceptible responses in newly

populated areas which makes it difficult to monitor changes through time in insecticide efficacy due to

resistance. Knowledge of whether the introduction occurred once (e.g., single 'invasive bridgehead effect';

Guillemaud et al. 2011) vs. multiple times (e.g., multiple 'mass dispersal'; Wilson et al. 2009) and patterns of

gene flow also impact the long-term monitoring of insecticide resistance evolution. Using native *S. frugiperda*

colonies established in 2005 from cotton fields in Louisiana, USA, Wilson et al. (2009) undertook diet-

incorporation bioassays (on 3[rd] instar larvae), using insecticides including chlorantraniliprole, indoxacarb, and

spinetoram (see Table 7). Yu (1991) undertook topical application bioassays against methomyl (on 4[th] instar

larvae) which included comparison with a susceptible population free from insecticide exposure since 1975,

and a resistant population collected from a maize field in Gainesville, Florida (see Table 8). In comparing

insecticide responses in Indian invasive *S. frugiperda* populations with diet incorporation assays (on 3[rd] instar

larvae), Yu (1991) also included a native susceptible *S, frugiperda* population from Brazil to assist with

interpreting changes to insecticide responses (including chlorantraniliprole and spinetoram) at spatial and

temporal scales.

In the absence of native *S. frugiperda* populations in Australia, and in addition to the information collected

herein for *H. armigera* and *S. litura,* we used the information from these global studies on native and newly

invaded populations to assist interpretation of the early data collected in this study on invasive *S. frugiperda*

populations in Australia. These examples used broadly similar approaches, instars and scoring criteria to our

study for bioassays with specific insecticides. Relative to the indoxacarb results from Hardke et al. (2011) on

native *S. frugiperda* populations from the USA, our bioassay findings for the invasive Western Australian and

Queensland populations suggest a 28- and 3-folds difference, respectively (Table 7). For spinetoram, the

responses of the two Australia populations (which were like each other) is around 10 and 1.5 times higher than

for the native populations from Brazil and the USA respectively. In contrast the ratios for the invasive

populations from India was 1.3 and 0.3 relative to the Brazil and USA native populations. The differences are

even more pronounced for chlorantraniliprole where the Sf20-4 WA population exhibited ratios that were 32

and 2.4 times higher than both the native Brazilian and American populations, which are at least around 3-fold

higher than for the invasive populations in South Africa and India (Table 7) and also for the invasive Indian

population studied by Deshmukh et al. (2020) using a leaf dip assay (and hence not reported in Table 7).

Methomyl resistance alleles have been reported in invasive populations from China, Indonesia, Africa, and in

this study. Comparisons between susceptible and resistant strains of native *S. frugiperda* populations from

Florida USA (Yu 1991) and with the South African invasive population of *S. frugiperda* (Eriksson 2019) showed

that the Queensland population was similar in its response as the susceptible Florida *S. frugiperda*, while the
Western Australia population was around 5 times more tolerant. This contrasted the South African population
which was 110 times more tolerant than the Florida susceptible *S. frugiperda* strain, and around 8 times more
tolerant than the resistant *S. frugiperda* strain from Florida (Table 8). Taken as a whole, similar bioassay results
between native (i.e., Brazil, USA) and various invasive populations suggested potential significant genetic
diversity in introduced populations. This concurs with population genomic and genetic analyses (Tay et al.
2020, Zhang et al. 2020, Schlum et al. 2021, Rane et al. 2022, Jiang et al. 2022) that suggested multiple origins
for the invasive African, Asian (Indian, Chinese), and Southeast Asian (e.g., Malaysia) FAW populations.

It is possible that the differential response between the two Australian *S. frugiperda* populations to
insecticides and Bt toxins represents natural variation in the national population which has emerged from a
single founding incursion. However, it is unlikely that the differential responses within Australian populations
represents local selection pressures because the period between populations establishing and being collected
for this study was presumed to be too short to enable this opportunity. It could be that separate incursions
involving different source populations occurred in WA compared to the eastern states of Australia, as
supported by genome-wide SNP marker population genomic studies (Rane et al. 2022). Different selection
pressures on the global population which recently originated from multiple Asian, Southeast Asian, and African
incursions (Tay et al. 2020, Zhang et al. 2020, Schlum et al. 2021, Rane et al. 2022) may have driven different
phenotypes which entered the expanding ranges tested in our study and others reported herein. However,
Kulye et al. (2021) demonstrated in *S. frugiperda* populations collected in India during 2018, 2019, and 2020
that large response changes such as those observed in South African (e.g., methomyl; Eriksson 2019) and
Western Australian (chlorantraniliprole, indoxacarb; this study) populations were unlikely considering the
short time frame since the very recent arrival of *S. frugiperda* especially assuming a west-to-east spread
(Goergen et al. 2016, Cock et al. 2017, Nagoshi et al. 2018, Nagoshi et al. 2019b). Another explanation for the
differential responses of the two Australian populations that we studied is that they have different fitness
because of variation in genetic diversity which reflects the number of individuals founding the populations.
However, if this was the case one might expect sensitivity levels to be consistently higher or lower to all of the
agents that we tested but this was not the case.

458   [Table 7, Table 8]

Alien invasive agricultural pests are increasingly being shown to carry novel insecticide resistance genes (e.g.,
Anderson et al. 2018, Walsh et al. 2018, Tay and Gordon 2019). This includes *S. frugiperda* in which invasive
populations have been confirmed via whole genome sequence analyses (e.g., Guan et al. 2020, Zhang et al.
2020, Yainna et al. 2021) and molecular characterisation to harbour selected resistance genes (Boaventura et
al. 2020b, Boaventura et al. 2020a, Zhao et al. 2020). Our review of reported ACE-1 and VGSC resistance allele
frequency differences (Fig, 1; Table S2 and references therein) suggests that the invasive *S. frugiperda*
populations within China (Guo et al. 2020), Indonesia (Boaventura et al. 2020a), Malaysia and Queensland,
Australia (Tay et al. 2021, Rane et al. 2022), and South Korea (this study) were genetically diverse and likely

originated from different native populations (Fig, 1). This further supports the perceived rapid spread of *S.*
frugiperda across Africa, Asia, and Oceania as likely to also involve multiple independent introduction events.

Given the insecticide resistance allele frequency differences in FAW *S. frugiperda* (e.g., Boaventura et al.
2020a, Guan et al. 2020, Lv et al. 2021, Yainna et al. 2021) and variation between the two populations sampled
herein in bioassay responses to some of the approved insecticides for broadacre cropping, effort in Australia
and indeed, for other invasive regions where possible, should now be directed to establishing baseline
susceptibilities against key chemistries for multiple populations across geographies. For instance, pyrethroid
resistance is common in the field in *H. armigera* (Walsh et al. 2018), and our comparisons with *H. armigera*
and *S. litura* suggest that pyrethroids like cypermethrin are unlikely to provide good control in Australia against
S. frugiperda. It also indicates some heterogeneity in the response to pyrethroids in the first invasive FAW
populations in Australia. The large discrepancies between WA and Qld populations for methomyl and
indoxacarb also require study of further populations before drawing firm conclusions. This information will be
critical for on-going monitoring of resistance allele frequencies to key chemistries and determining their field
application efficiencies, among geographies which is required to inform resistance management plans. It will
also be important to understand gene flow patterns between different populations. Early detection of
potential future introductions into Australia of novel resistance genes/alleles should also be a priority; for
example, the VGSC L1014 resistance allele and the ACE G227A resistance allele from Southeast Asia
(Boaventura et al. 2020a) and China (Guo et al. 2020), ryanodine receptor (RyR) resistance alleles from Brazil
and China (Bolzan et al. 2019, Boaventura et al. 2020b, Lv et al. 2021) and ABCC2 resistance alleles from the
Americas (Banerjee et al. 2017, Flagel et al. 2018, Guan et al. 2020, Yainna et al. 2021).

Being alert to new *S. frugiperda* incursions carrying novel resistance genes is relevant also for the Bt toxins.
Our bioassay findings suggest Australia's FAW *S. frugiperda* populations likely do not carry Cry1F resistance
alleles in homozygous state known to exist in native range *S. frugiperda* populations (e.g., Banerjee et al. 2017,
Flagel et al. 2018, Guan et al. 2020, Yainna et al. 2021), however the 2 bp deletion identified via whole genome
sequencing in a single heterozygous individual from Burdekin (Queensland) will require further confirmation.
Field-selected lines of VIP3A20 resistant *S. frugiperda* have been reported in Brazil and the USA where they
occur as a recessive trait (Yang et al. 2013, Bernardi et al. 2015, Bernardi et al. 2016, Yang et al. 2018, Yang et
al. 2019). The candidate resistance gene(s) and associated mutation(s) underpinning this resistance are yet to
be identified, therefore phenotypic bioassays are required to detect resistance to VIP3A. Using what was
essentially an F_0 screen of material we did not detect resistance to VIP3A in our field-derived laboratory-
maintained *S. frugiperda* colonies, however bioassays involving F_2-crosses ▯▯▯(Andow and Alstad 1998)▯
would be needed to VIP3A resistance allele.both populations, and should be a valuable tool in protecting the
Australian cotton industry against *S. frugiperda* due to the > 90% up-take of Bollgard III cotton containing
Cry1Ac, Cry2Ab and Vip3A proteins. To protect agriculture in Australia (and elsewhere) from an accidental
introduction of VIP3A resistant populations from the Americas (e.g., from Brazil, Bernardi et al. 2016) into
global invasive populations, national (e.g., pre-border) and industry biosecurity preparedness strategies must

be co-ordinated to prevent and to increase the chances of early detection of such novel introductions of new
traits.

Novel introductions leading to unique population structure has been reported for western African *S.*
frugiperda (Nagoshi et al. 2022), in populations from China (Jiang et al. 2022), in Africa (e.g., Benin vs. Malawi;
Tay et al. 2022), and in, e.g., Australia, Malaysia, and Myanmar vs. China populations (Rane et al. 2022),
suggesting that the widely anticipated long distance migration of *S. frugiperda* especially in the invasive range
could likely be less and may be impacted by localised ecological and climatic determinants. Factors impacting
the founding for the two Australian populations (e.g., founding number (i.e., involving one/few individuals vs.
many); gene flow dynamics at spatial and temporal scales, and frequencies and impact from potential novel
and on-going introductions) used in our bioassay studies is at present unknown. While bioassays testing of a
larger population pooled from several sites or several individual populations from multiple sites would be
especially relevant in a pest capable of high dispersal across the landscape, however the spread of *S.*
frugiperda in the invasive range is increasingly being recognised as may not be as rapid and widespread as
originally believed (e.g., Naghoshi et al. 2022; Rane et al. 2022; Tay et al. 2022; Jiang et al. 2022). As such, we
advocate cautionary approaches to avoid premature assumptions that extensive population admixture via
gene flow in Australia landscape would lead to homogenised populations and therefore the approach to
combine multiple distantly sampled populations in bioassay studies (e.g., Bird et al. 2022).

Finally, to enable meaningful comparisons of bioassay findings from across the *S. frugiperda* invasive range,
there is a need to globally standardise approaches for testing of insecticides and Bt toxins. Localised novel
resistance traits in populations of *S. frugiperda* would likely serve as a new resistance management challenge
to other neighbouring regions (Kalyebi 2020), and movements of *S. frugiperda* in the new invasive ranges
could lead to as yet unknown and complex gene flow patterns that could significantly hinder the development
of suitable pest and resistance management strategies for this global pest complex.

Acknowledgements

The *Spodoptera frugiperda* larvae used to establish the SF20-1 colony were provided by Prof. Myron Zalucki
(University of Queensland). Mr Jack Daniel (Northern Australian Crop Research Alliance Pty Ltd) provided the
S. frugiperda larvae used to establish SF20-4. Material from an established colony of *S. litura* was provided by
Dr Ian Newton and Dr Melina Miles (QDAF). Some of the results reported herein were collected for a project
led by CSIRO as a co-investment with the Grains Research and Development Corporation, the Australian
Centre for International Agricultural Research, the Cotton Research and Development Corporation, FMC
Australasia and Corteva Agriscience. GRDC manages the project on behalf of the co-investors.

References

Anderson, C. J., W. T. Tay, A. McGaughran, K. Gordon, and T. K. Walsh. 2016. Population structure and gene
flow in the global pest, *Helicoverpa armigera*. Mol Ecol 25: 5296-5311.

Anderson, C. J., J. G. Oakeshott, W. T. Tay, K. H. J. Gordon, A. Zwick, and T. K. Walsh. 2018. Hybridization and
gene flow in the mega-pest lineage of moth, *Helicoverpa*. Proc Natl Acad Sci U S A 115: 5034-5039.

Andow, D. A., and D. N. Alstad. 1998. F-2 screen for rare resistance alleles. Journal of Economic Entomology
91: 572-578.

Banerjee, R., J. Hasler, R. Meagher, R. Nagoshi, L. Hietala, F. Huang, K. Narva, and J. L. Jurat-Fuentes. 2017.
Mechanism and DNA-based detection of field-evolved resistance to transgenic Bt corn in fall armyworm
(*Spodoptera frugiperda*). Sci Rep 7: 10877.

Bernardi, O., D. Bernardi, R. S. Ribeiro, D. M. Okuma, E. Salmeron, J. Fatoretto, F. C. L. Medeiros, T. Burd,
and C. Omoto. 2015. Frequency of resistance to Vip3Aa20 toxin from Bacillus thuringiensis in *Spodoptera*
frugiperda (Lepidoptera: Noctuidae) populations in Brazil. Crop Prot 76: 7-14.

Bernardi, O., D. Bernardi, R. J. Horikoshi, D. M. Okuma, L. L. Miraldo, J. Fatoretto, F. C. L. Medeiros, T. Burd,
and C. Omoto. 2016. Selection and characterization of resistance to the Vip3Aa20 protein from Bacillus
thuringiensis in *Spodoptera frugiperda*. Pest Management Science 72: 1794-1802.

Bird, L. J. 2015. Baseline Susceptibility of *Helicoverpa armigera* (Lepidoptera: Noctuidae) to Indoxacarb,
Emamectin Benzoate, and Chlorantraniliprole in Australia. Journal of Economic Entomology 108: 294-300.

Bird, L. J. 2018. Pyrethroid and carbamate resistance in Australian *Helicoverpa armigera* (Lepidoptera:
Noctuidae) from 2008 to 2015: what has changed since the introduction of Bt cotton? B Entomol Res 108:
781-791.

Bird, L. J., and R. J. Akhurst. 2007. Variation in susceptibility of *Helicoverpa armigera* (Hubner) and Helicoverpa
punctigera (Wallengren) (Lepidoptera : Noctuidae) in Australia to two *Bacillus thuringiensis* toxins. J
Invertebr Pathol 94: 84-94.

Bird, L. J., and P. W. Walker. 2019. Baseline susceptibility of *Helicoverpa punctigera* (Lepidoptera: Noctuidae)
to indoxacarb, emamectin benzoate, and chlorantraniliprole. Journal of Economic Entomology 112: 818–
826.

Bird, L., M. Miles, A. Quade, and H. Spafford. 2022. Insecticide resistance in Australian Spodoptera frugiperda
(J.E. Smith) and development of testing procedures for resistance surveillance. PLoS ONE
doi.org/10.1371/journal.pone.0263677.

Boaventura, D., M. Martin, A. Pozzebon, D. Mota-Sanchez, and R. Nauen. 2020a. Monitoring of Target-Site
Mutations Conferring Insecticide Resistance in *Spodoptera frugiperda*. Insects 11: 545.

Boaventura, D., A. Bolzan, F. E. O. Padovez, D. M. Okuma, C. Omoto, and R. Nauen. 2020b. Detection of a
ryanodine receptor target-site mutation in diamide insecticide resistant fall armyworm, *Spodoptera*
frugiperda. Pest Management Science 76: 47-54.

Bolzan, A., F. E. O. Padovez, A. R. B. Nascimento, I. S. Kaiser, E. C. Lira, F. S. A. Amaral, R. H. Kanno, J. B.
Malaquias, and C. Omoto. 2019. Selection and characterization of the inheritance of resistance of
Spodoptera frugiperda (Lepidoptera: Noctuidae) to chlorantraniliprole and cross-resistance to other
diamide insecticides. Pest Management Science 75: 2682-2689.

Botha, A. S., A. Erasmus, H. du Plessis, and J. Van den Berg. 2019. Efficacy of Bt Maize for Control of
Spodoptera frugiperda (Lepidoptera: Noctuidae) in South Africa. Journal of Economic Entomology 112:
1260-1266.

Carvalho, R. A., C. Omoto, L. M. Field, M. S. Williamson, and C. Bass. 2013. Investigating the Molecular
Mechanisms of Organophosphate and Pyrethroid Resistance in the Fall Armyworm *Spodoptera*
frugiperda. Plos One 8: e62268.

Cock, M. J. W., P. K. Beseh, A. G. Buddie, G. Cafa, and J. Crozier. 2017. Molecular methods to detect
Spodoptera frugiperda in Ghana, and implications for monitoring the spread of invasive species in
developing countries. Sci Rep 7: 4103.

Deshmukh, S., H. B. Pavithra, C. M. Kalleshwaraswamy, B. K. Shivanna, M. S. Maruthi, and D. Mota-Sanchez.
2020. Field efficacy of insecticides for management of invasive fall armyworm, *Spodoptera frugiperda* (J.
E. Smith) (Lepidoptera: Noctuidae) on maize in India. Fla Entomol 103: 221-227.

Downes, S., T. L. Parker, and R. J. Mahon. 2009. Frequency of Alleles Conferring Resistance to the *Bacillus*
thuringiensis Toxins Cry1Ac and Cry2Ab in Australian Populations of *Helicoverpa punctigera* (Lepidoptera:
Noctuidae) From 2002 to 2006. Journal of Economic Entomology 102: 733-742.

du Plessis, H., J. van den Berg, N. Ota, and D. J. Kriticos. Year. Published. *Spodoptera frugiperda* (Fall
Armyworm), p. 7. *In*, CSIRO-InSTePP Pest Geography, June, 2018 2018, Canberra, Australia. CSIRO.

Early, R., P. Gonzalez-Moreno, S. T. Murphy, and R. Day. 2018. Forecasting the global extent of invasion of the
cereal pest *Spodoptera frugiperda,* the fall armyworm. Neobiota: 25-50.

Eriksson, J. 2019. Toxicity bioassays with insecticide formulations used for control of *Spodoptera frugiperda*
(Lepidoptera: Noctuidae). Master of Science in Environmental Science with Integrated Pest Management,
North West University South Africa.

Faust, G. G., and I. M. Hall. 2014. SAMBLASTER: fast duplicate marking and structural variant read extraction.
Bioinformatics 30: 2503-2505.

Flagel, L., Y. W. Lee, H. Wanjugi, S. Swarup, A. Brown, J. L. Wang, E. Kraft, J. Greenplate, J. Simmons, N.
Adams, Y. F. Wang, S. Martinelli, J. A. Haas, A. Gowda, and G. Head. 2018. Mutational disruption of the
ABCC2 gene in fall armyworm, *Spodoptera frugiperda,* confers resistance to the Cry1Fa and Cry1A.105
insecticidal proteins. Sci Rep-Uk 8: 7255.

Goergen, G., P. L. Kumar, S. B. Sankung, A. Togola, and M. Tamo. 2016. First Report of Outbreaks of the Fall
Armyworm *Spodoptera frugiperda* (J E Smith) (Lepidoptera, Noctuidae), a New Alien Invasive Pest in West
and Central Africa. PLoS One 11: e0165632.

Gouin, A., A. Bretaudeau, K. Nam, S. Gimenez, J. M. Aury, B. Duvic, F. Hilliou, N. Durand, N. Montagne, I.
Darboux, S. Kuwar, T. Chertemps, D. Siaussat, A. Bretschneider, Y. Mone, S. J. Ahn, S. Hanniger, A. G.
Grenet, D. Neunemann, F. Maumus, I. Luyten, K. Labadie, W. Xu, F. Koutroumpa, J. M. Escoubas, A.
Llopis, M. Maibeche-Coisne, F. Salasc, A. Tomar, A. R. Anderson, S. A. Khan, P. Dumas, M. Orsucci, J.
Guy, C. Belser, A. Alberti, B. Noel, A. Couloux, J. Mercier, S. Nidelet, E. Dubois, N. Y. Liu, I. Boulogne, O.
Mirabeau, G. Le Goff, K. Gordon, J. Oakeshott, F. L. Consoli, A. N. Volkoff, H. W. Fescemyer, J. H.
Marden, D. S. Luthe, S. Herrero, D. G. Heckel, P. Wincker, G. J. Kergoat, J. Amselem, H. Quesneville, A.
T. Groot, E. Jacquin-Joly, N. Negre, C. Lemaitre, F. Legeai, E. d'Alencon, and P. Fournier. 2017. Two
genomes of highly polyphagous lepidopteran pests (*Spodoptera frugiperda,* Noctuidae) with different
host-plant ranges. Sci Rep 7: 11816.

Guan, F., J. Zhang, H. Shen, X. Wang, A. Padovan, T. K. Walsh, W. T. Tay, K. H. J. Gordon, W. James, C.
Czepak, M. H. Otim, D. Kachigamba, and Y. Wu. 2021. Whole-genome sequencing to detect mutations
associated with resistance to insecticides and Bt proteins in *Spodoptera frugiperda*. Insect Sci. 28: 627-
638.

Gui, F. R., T. M. Lan, Y. Zhao, W. Guo, Y. Dong, D. M. Fang, H. Liu, H. M. Li, H. L. Wang, R. S. Hao, X. F. Cheng,
Y. H. Li, P. C. Yang, S. K. Sahu, Y. P. Chen, L. Cheng, S. Q. He, P. Liu, G. Y. Fan, H. R. Lu, G. H. Hu, W. Dong,
B. Chen, Y. Jiang, Y. W. Zhang, H. H. Xu, F. Lin, B. Slipper, A. Postma, M. Jackson, B. A. Abate, K. Tesfaye,
A. L. Demie, M. D. Bayeleygne, D. T. Degefu, F. Chen, P. K. Kuria, Z. M. Kinyua, T. X. Liu, H. M. Yang, F.
N. Huang, X. Liu, J. Sheng, and L. Kang. 2020. Genomic and transcriptomic analysis unveils population
evolution and development of pesticide resistance in fall armyworm *Spodoptera frugiperda*. Protein Cell.
doi: 10.1007/s13238-020-00795-7.

Guillemaud, T., M. Ciosi, E. Lombaert, and A. Estoup. 2011. Biological invasions in agricultural settings:
Insights from evolutionary biology and population genetics. Cr Biol 334: 237-246.

Guo, J. M., X. L. Liu, S. R. Liu, Z. Q. Wei, W. K. Han, Y. Z. Guo, and S. L. Dong. 2020. Functional Characterization
of Sex Pheromone Receptors in the Fall Armyworm (*Spodoptera frugiperda*). Insects 11: 193.

Hardke, J. T., J. H. Temple, B. R. Leonard, and R. E. Jackson. 2011. Laboratory Toxicity and Field Efficacy of
Selected Insecticides against Fall Armyworm (Lepidoptera: Noctuidae). Fla Entomol 94: 272-278.

Hardwick, D. F. 1965. The Corn Earworm complex. Memoirs of the Entomological Society of Canada 40: 1-248.

Horikoshi, R. J., D. Bernardi, O. Bernardi, J. B. Malaquias, D. M. Okuma, L. L. Miraldo, F. S. D. E. Amaral, and
C. Omoto. 2016. Effective dominance of resistance of *Spodoptera frugiperda* to Bt maize and cotton
varieties: implications for resistance management. Sci Rep-Uk 6.

Jiang, Y. Y., Y. Y. Zhang, X. Y. Zhou, X Y. Hong, and L. Chen. 2022. Population genetics reveal multiple
independent invasions of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in China. Bull Entomol Res, 1-
11 doi: 10.1017/S0007485322000190.

Jing, W., C. Huang, C. Y. Li, H. X. Zhou, Y. L. Ren, Z. Y. Li, L. S. Xing, B. Zhang, X. Qiao, B. Liu, C. H. Liu, Y. Xi, W.
X. Liu, W. K. Wang, W. Q. Qian, S. Mckirdy, and F. H. Wan. 2021. Biology, invasion and management of
the agricultural invader: Fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae). Journal of
Integrative Agriculture 20: 646-663.

Jones, C. M., H. Parry, W. T. Tay, D. R. Reynolds, and J. W. Chapman. 2019. Movement Ecology of Pest
Helicoverpa: Implications for Ongoing Spread. Annu Rev Entomol 64: 277-295.

Kalyebi, A. 2020. Documentation of farmer practices that have been useful in Africa for managing the fall
armyworm (*Spodoptera frugiperda*): A case study from Uganda. pp. 30. GRDC.

Kulye, M., S. Mehlhorn, D. Boaventura, N. Godley, S. K. Venkatesh, T. Rudrappa, T. Charan, D. Rathi, and R.
Nauen. 2021. Baseline Susceptibility of *Spodoptera frugiperda* Populations Collected in India towards
Different Chemical Classes of Insecticides. Insects 12: 758.

LeOra_Sorftware 1987. POLO-PC: A Users Guide to Probit or Logit Analysis computer program, version By
LeOra_Sorftware.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin, and S.
Genome Project Data Processing. 2009. The Sequence Alignment/Map format and SAMtools.
Bioinformatics 25: 2078-2079.

Lv, S. L., Y. Shi, J. C. Zhang, P. Liang, L. Zhang, and X. W. Gao. 2021. Detection of ryanodine receptor target-
site mutations in diamide insecticide-resistant *Spodoptera frugiperda* in China. Insect Science 28: 639-
648.

Mahon, R. J., K. M. Olsen, and S. Downes. 2008. Isolations of Cry2Ab resistance in Australian populations of
Helicoverpa armigera (Lepidoptera : Noctuidae) are allelic. Journal of Economic Entomology 101: 909-
914.

Mahon, R. J., S. J. Downes, and B. James. 2012. Vip3A Resistance Alleles Exist at High Levels in Australian
Targets before Release of Cotton Expressing This Toxin. Plos One 7.

Mahon, R. J., K. M. Olsen, K. A. Garsia, and S. R. Young. 2007a. Resistance to *Bacillus thuringiensis* toxin
Cry2Ab in a strain of *Helicoverpa armigera* (Lepidoptera : Noctuidae) in Australia. Journal of Economic
Entomology 100: 894-902.

Mahon, R. J., K. M. Olsen, S. Downes, and S. Addison. 2007b. Frequency of alleles conferring resistance to the
bt toxins Cry1Ac and Cry2Ab in Australian populations of *Helicoverpa armigera* (Lepidoptera : noctuidae).
Journal of Economic Entomology 100: 1844-1853.

Mahon, R. J., S. Downes, W. James, and T. Parker. 2010. Why Do F-1 Screens Estimate Higher Frequencies of
Cry2Ab Resistance in *Helicoverpa armigera* (Lepidoptera: Noctuidae) Than Do F-2 Screens? Journal of
Economic Entomology 103: 472-481.

Montezano, D. G., A. Specht, D. R. Sosa-Gomez, V. F. Roque-Specht, J. C. Sousa-Silva, S. V. Paula-Moraes, J.
A. Peterson, and T. E. Hunt. 2018. Host plants of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in the
Americas. Afr Entomol 26: 286-300.

Nagoshi, R. N., G. Goergen, K. A. Tounou, K. Agboka, D. Koffi, and R. L. Meagher. 2018. Analysis of strain
distribution, migratory potential, and invasion history of fall armyworm populations in northern Sub-
Saharan Africa. Sci Rep-Uk 8.

Nagoshi, R. N., B. Y. Nagoshi, E. Canarte, B. Navarrete, R. Solorzano, and S. Garces-Carrera. 2019a. Genetic
characterization of fall armyworm (*Spodoptera frugiperda*) in Ecuador and comparisons with regional
populations identify likely migratory relationships. PLoS One 14: e0222332.

Nagoshi, R. N., I. Dhanani, R. Asokan, H. M. Mahadevaswamy, C. M. Kalleshwaraswamy, Sharanabasappa,
and R. L. Meagher. 2019b. Genetic characterization of fall armyworm infesting South Africa and India
indicate recent introduction from a common source population. Plos One 14.

Nagoshi, R. N., G. Goergen, D. Koffi, et al. 2022. Genetic studies of fall armyworm indicate a new introduction
into Africa and identify limits to its micratory behaviour. Sci. Rep. 12, 1941 .
https://doi.org/10.1038/s41598-022-05781-z

Nguyen, D. T., Y. Z. Chen, and G. A. Herron. 2021. Preliminary characterisation of known pesticide resistance
alleles in *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in its invasive Australian range. Austral Entomol
60: 782-790.

Otim, M. H., W. T. Tay, T. K. Walsh, D. Kanyesigye, S. Adumo, J. Abongosi, S. Ochen, J. Sserumaga, S. Alibu,
G. Abalo, G. Asea, and A. Agona. 2018. Detection of sister-species in invasive populations of the fall
armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) from Uganda. PLoS One 13: e0194571.

Pearce, S. L., D. F. Clarke, P. D. East, S. Elfekih, K. H. J. Gordon, L. S. Jermiin, A. McGaughran, J. G. Oakeshott,
A. Papanicolaou, O. P. Perera, R. V. Rane, S. Richards, W. T. Tay, T. K. Walsh, A. Anderson, C. J.
Anderson, S. Asgari, P. G. Board, A. Bretschneider, P. M. Campbell, T. Chertemps, J. T. Christeller, C. W.
Coppin, S. J. Downes, G. Duan, C. A. Farnsworth, R. T. Good, L. B. Han, Y. C. Han, K. Hatje, I. Horne, Y. P.
Huang, D. S. T. Hughes, E. Jacquin-Joly, W. James, S. Jhangiani, M. Kollmar, S. S. Kuwar, S. Li, N. Y. Liu,
M. T. Maibeche, J. R. Miller, N. Montagne, T. Perry, J. Qu, S. V. Song, G. G. Sutton, H. Vogel, B. P.
Walenz, W. Xu, H. J. Zhang, Z. Zou, P. Batterham, O. R. Edwards, R. Feyereisen, R. A. Gibbs, D. G.
Heckel, A. McGrath, C. Robin, S. E. Scherer, K. C. Worley, and Y. D. Wu. 2017a. Genomic innovations,
transcriptional plasticity and gene loss underlying the evolution and divergence of two highly
polyphagous and invasive *Helicoverpa* pest species. BMC Biol 15: 63.

Pearce, S. L., D. F. Clarke, P. D. East, S. Elfekih, K. H. J. Gordon, L. S. Jermiin, A. McGaughran, J. G. Oakeshott,
A. Papanicolaou, O. P. Perera, R. V. Rane, S. Richards, W. T. Tay, T. K. Walsh, A. Anderson, C. J.
Anderson, S. Asgari, P. G. Board, A. Bretschneider, P. M. Campbell, T. Chertemps, J. T. Christeller, C. W.
Coppin, S. J. Downes, G. Duan, C. A. Farnsworth, R. T. Good, L. B. Han, Y. C. Han, K. Hatje, I. Horne, Y. P.
Huang, D. S. T. Hughes, E. Jacquin-Joly, W. James, S. Jhangiani, M. Kollmar, S. S. Kuwar, S. Li, N. Y. Liu,
M. T. Maibeche, J. R. Miller, N. Montagne, T. Perry, J. Qu, S. V. Song, G. G. Sutton, H. Vogel, B. P.
Walenz, W. Xu, H. J. Zhang, Z. Zou, P. Batterham, O. R. Edwards, R. Feyereisen, R. A. Gibbs, D. G.
Heckel, A. McGrath, C. Robin, S. E. Scherer, K. C. Worley, and Y. D. Wu. 2017b. Erratum to: Genomic
innovations, transcriptional plasticity and gene loss underlying the evolution and divergence of two highly
polyphagous and invasive *Helicoverpa* pest species. BMC Biol 15: 69.

Qi, G. J., J. Ma, J. Wan, Y. L. Ren, S. McKirdy, G. Hu, and Z. F. Zhang. 2021. Source Regions of the First
Immigration of Fall Armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae) Invading Australia.
Insects 12: 1104.

Rane, R., T. K. Walsh, P. Lenancker, W. James, A. Gock, D. T. Hang, V. Nguyen. L, T. N. Khin, D. Amalin, K.
Chittarath, F. Muhammad, S. Annamalai, S. S. Thanarajoo, Y. A. Trisyono, K. Sahia, J. Kim, L. Kuniata, K.
Powell, K. H. J. Gordon, and W. T. Tay. 2022. Genome-wide SNP analysis suggests complex introductions
of *Spodoptera frugiperda* in Southeast Asia and Australia. eLife: submitted.

Rose, A. H., R. H. Silversides, and O. H. Lindquist. 1975. Migration Flight by an Aphid, *Rhopalosiphum maidis*
(Hemiptera-Aphididae), and a Noctuid, *Spodoptera frugiperda* (Lepidoptera-Noctuidae). Can Entomol
107: 567-576.

Schlum, K. A., K. Lamour, C. P. de Bortoli, R. Banerjee, R. Meagher, E. Pereira, M. G. Murua, G. A. Sword, A.
E. Tessnow, D. V. Dillon, A. M. L. Ramirez, K. S. Akutse, R. Schmidt-Jeffris, F. N. Huang, D. Reisig, S. J.
Emrich, and J. L. Jurat-Fuentes. 2021. Whole genome comparisons reveal panmixia among fall
armyworm (*Spodoptera frugiperda*) from diverse locations. BMC Genomics 22: 179.

Sparks, A. N. 1979. Review of the Biology of the Fall Armyworm (Lepidoptera, Noctuidae). Fla Entomol 62: 82-
87.

Tay, W. T., and K. H. J. Gordon. 2019. Going global - genomic insights into insect invasions. Curr Opin Insect Sci
31: 123-130.

Tay, W. T., P. J. Kerr, and L. S. Jermiin. 2016. Population Genetic Structure and Potential Incursion Pathways of
the Bluetongue Virus Vector *Culicoides brevitarsis* (Diptera: Ceratopogonidae) in Australia. Plos One 11:
e0146699.

Tay, W. T., R. Rane, B. James, A. Gock, D. T. Hang, V. L. Nguyen, T. N. Khin, D. Amalin, C. Khonesavanh, F.
Muhammad, S. Annamalai, and T. Walsh. 2021. Final report on Southeast Asian and Australian FAW
population genomics for biosecurity preparedness. Prevention and preparedness for fall armyworm
(*Spodoptera frugiperda*) - Output 2, pp. 25. CSIRO/GRDC.

Tay, W. T., R. Rane, A. Padovan, T. Walsh, S. Elfekih, S. Downes, K. Nam, E. d'Alencon, J. Zhang, Y. Wu, N.
Negre, D. Kunz, D. J. Kriticos, C. Czepak, M. Otim, and K. H. J. Gordon. 2020. Global FAW population
genomic signature supports complex introduction events across the Old World. bioRxiv
2020.06.12.147660.

Vasimuddin, M., S. Misra, H. Li, and S. Aluru. 2019. Efficient Architecture-Aware Acceleration of BWA-MEM
for Multicore Systems. Int Parall Distrib P: 314-324.

Walsh, T. K., S. J. Downes, J. Gascoyne, W. James, T. Parker, J. Armstrong, and R. J. Mahon. 2014. Dual
Cry2Ab and Vip3A Resistant Strains of *Helicoverpa armigera* and *Helicoverpa punctigera* (Lepidoptera:
Noctuidae); Testing Linkage Between Loci and Monitoring of Allele Frequencies. Journal of Economic
Entomology 107: 1610-1617.

Walsh, T. K., N. Joussen, K. Tian, A. McGaughran, C. J. Anderson, X. Qiu, S. J. Ahn, L. Bird, N. Pavlidi, J.
Vontas, J. Ryu, A. Rasool, I. Barony Macedo, W. T. Tay, Y. Zhang, M. E. A. Whitehouse, P. J. Silvie, S.
Downes, L. Nemec, and D. G. Heckel. 2018. Multiple recombination events between two cytochrome
P450 loci contribute to global pyrethroid resistance in *Helicoverpa armigera*. PLoS One 13: e0197760.

Wang, J., H. D. Wang, S. Y. Liu, L. P. Liu, W. T. Tay, T. K. Walsh, Y. H. Yang, and Y. D. Wu. 2017. CRISPR/Cas9
mediated genome editing of *Helicoverpa armigera* with mutations of an ABC transporter gene HaABCA2
confers resistance to *Bacillus thuringiensis* Cry2A toxins. Insect Biochem Molec 87: 147-153.

Willis, K. J. 2017. Sate of the World's Plants 2017. Report., pp. 100. *In* K. J. Killis [ed.]. Royal Botanic Gardens,
Kew.

Wilson, J. R. U., E. E. Dormontt, P. J. Prentis, A. J. Lowe, and D. M. Richardson. 2009. Something in the way
you move: dispersal pathways affect invasion success. Trends Ecol Evol 24: 136-144.

Worku, M., and Y. Ebabuye. 2019. Evaluation of efficacy of insecticides against the fall army worm *Spodoptera*
frugiperda. Indian Journal of Entomology 81: 13-15.

Xiao, H., X. Ye, H. Xu, Y. Mei, Y. Yang, X. Chen, Y. Yang, T. Liu, Y. Yu, W. Yang, Z. Lu, and F. Li. 2020. The
genetic adaptations of fall armyworm *Spodoptera frugiperda* facilitated its rapid global dispersal and
invasion. Mol Ecol Resour 20: 1050-1068.

Yainna, S., N. Negre, P. J. Silvie, T. Brevault, W. T. Tay, K. Gordon, E. dAlencon, T. Walsh, and K. Nam. 2021.
Geographic Monitoring of Insecticide Resistance Mutations in Native and Invasive Populations of the Fall
Armyworm. Insects 12: 468.

Yang, F., J. Williams, P. Porter, F. Huang, and D. L. Kerns. 2019. F-2 screen for resistance to *Bacillus*
thuringiensis Vip3Aa51 protein in field populations of *Spodoptera frugiperda* (Lepidoptera: Noctuidae)
from Texas, USA. Crop Prot 126.

Yang, F., F. N. Huang, J. A. Qureshi, B. R. Leonard, Y. Niu, L. P. Zhang, and D. S. Wangila. 2013. Susceptibility
of Louisiana and Florida populations of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) to transgenic
Agrisure (R) Viptera (TM) 3111 corn. Crop Prot 50: 37-39.

Yang, F., S. Morsello, G. P. Head, C. Sansone, F. N. Huang, R. T. Gilreath, and D. L. Kerns. 2018. F-2 screen,
inheritance and cross-resistance of field-derived Vip3A resistance in *Spodoptera frugiperda* (Lepidoptera:
Noctuidae) collected from Louisiana, USA. Pest Management Science 74: 1769-1778.

Yu, S. J. 1991. Insecticide Resistance in the Fall Armyworm, *Spodoptera frugiperda* (Smith,J.E.). Pestic Biochem
Phys 39: 84-91.

Zhang, J. P., F. Zhang, W. T. Tay, C. Robin, Y. Shi, F. Guan, Y. H. Yang, and Y. D. Wu. 2022. Population
genomics provides insights into lineage divergence and local adaptation within the cotton bollworm. Mol
Ecol Resour. https://doi.org/10.1111/1755-0998.13581.

Zhang, L., B. Liu, W. G. Zheng, C. H. Liu, D. N. Zhang, S. Y. Zhao, Z. Y. Li, P. J. Xu, K. Wilson, A. Withers, C. M.
Jones, J. A. Smith, G. Chipabika, D. L. Kachigamba, K. Nam, E. D'Alencon, B. Liu, X. Y. Liang, M. H. Jin, C.
Wu, S. Chakrabarty, X. M. Yang, Y. Y. Jiang, J. Liu, X. L. Liu, W. P. Quan, G. R. Wang, W. Fan, W. Q. Qian,
K. M. Wu, and Y. T. Xiao. 2020. Genetic structure and insecticide resistance characteristics of fall
armyworm populations invading China. Mol Ecol Resour. 20: 1682-1696.

Zhao, Y. X., J. M. Huang, H. Ni, D. Guo, F. X. Yang, X. Wang, S. F. Wu, and C. F. Gao. 2020. Susceptibility of fall
armyworm, *Spodoptera frugiperda* (J.E.Smmith), to eight insecticides in China, with special reference to
lambda-cyhalothrin. Pestic Biochem Phys. 168: 104623.

Table 1: List of insecticides and Bt toxins used in the bioassay experiments. All insecticides and Bt toxins were
tested on two populations of *Spodoptera frugiperda* from Queensland (Sf20-1, from Walkamin) and Western
Australia (Sf20-4, from Kununurra). Bioassays involved either: (i) diet incorporation (DI); (ii) Topical application
(TA); or (iii) Surface treatment (ST) of diet to be consumed. The larval stages used in the bioassays are
specified. For topical application we used larvae from late 3^{rd} to early 4^{th} instar with each larva weighing
approximately 30mg. Species tested were: *Spodoptera frugiperda*, *Helicoverpa armigera,* and *Spodoptera*
litura.

Insecticides/Bt toxins	Commercial name	Date	Approach	Larval stage	Species tested		
					S. frugiperda	*H. armigera*	*S. litura*
Chlorantraniliprole (200g/L)	Coragen	Manufactured: 28/06/2019	DI	2^{nd}/early 3^{rd}	Yes	Yes	No
Indoxacarb (150g/L)	Steward 150	Manufactured: 30/03/2019	DI	2^{nd}/early 3^{rd}	Yes	Yes	No
Emamectin (17g/L)	Affirm 017 EC	Unspecified†	DI	2^{nd}/early 3^{rd}	Yes	Yes	No
Spinetoram (120g/L)	Dow AgroSciences Success® Neo	Manufactured: 02/2018	DI	2^{nd}/early 3^{rd}	Yes	Yes	No
Alpha cypermethrin (100g/L)	Dominex Duo	Manufactured: 30/04/2019	TA	Late 3^{rd}/ early 4^{th}	Yes	Yes	No
Methomyl		Expiry date: 02/2022	TA	Late 3^{rd}/ early 4^{th}	Yes	Yes	No
Advertised as: Cry2A, Cry2B, Cry1Ab, Cry1Aa, Cry1Ac	Yates Nature's Way DIPEL	Expiry date: 27/05/2022	ST	Neonates	Yes	Yes	Yes
Advertised as: Cry1D, Cry1C, Cry1Ab, Cry1Aa	XenTari WG Bta	Manufactured: 05/02/2019	ST	Neonates	Yes	Yes	Yes
Cry1Ac	CSIRO; from HD73 strain	NA	ST	Neonates	Yes	Yes	Yes
Cry2Ab	CSIRO; from Bt expressing Cry2Ab	NA	ST	Neonates	Yes	Yes	Yes
Cry1F	CSIRO; from *E. coli* expressing Cry1F	NA	ST	Neonates	Yes	Yes	Yes
VIP3A	CSIRO; from *E. coli* expressing Vip3A	NA	ST	Neonates	Yes	Yes	Yes

Note: Commercial names of insecticides/Bt toxins, their manufactured or expiry dates are provided where
available. † likely manufactured recently as this was requested from Syngenta Australia Pty Ltd and provided
directly for the bioassay purpose. N/A: not available. These Bt toxins were produced and purified from *B.*

thuringiensis cultures (Cry1Ac and Cry2Ab) or cloned into vectors in *Escherichia coli* (Cry1F and Vip3A) and

grown at the CSIRO Black Mountain Laboratories.

Table 2: *Spodoptera frugiperda* samples from Australian populations (Sf20-1, Qld; Sf20-4, WA), Papua New Guinea, South Korea, and Peru used in whole genome
sequencing for resistance gene characterisation. Number of specimens from each location is indicated by 'N'. Where available, GPS co-ordinates and elevation are
provided. The number of individuals belonging to the *S. frugiperda* R-strain or the C-strain mitochondrial DNA COI haplotypes is also provided.

Country	State/Province	Population	N	Date	Note	R : C
Australia	WA	Kununurra	9	16-Aug-20	'Sf20-4', represent the original WA field-collected individuals	9 : 0
	Qld	Strathmore Station, Walkamin	29	03-Mar-20	'Sf20-1', represent the original Qld field-collected individuals	29 : 0
South Korea	Milyang	MF	7	Feb-20	Milyang-si, Gyeongsangnam-do, G_4 lab, GPS: 35°29.29 N, 128°44.31 E	7 : 0
	Goryong	GR	1	Sep-19	Goryeong-gun, Gyeongsangbuk-do, Corn-field; GPS: 35° 64.04 N, 128°39.08 E	1 : 0
	Haenam	MN	1	Aug-19	Haenam-gun, Jeollanam-do, Corn field; GPS: 34°24.38 N, 126°37.55 E	1 : 0
	Milyang	MY	2	Sep-19	Milyang-si, Gyeongsangnam-do, Corn field; GPS: 35°29.29 N, 128°44.31 E	0 : 2
	Muan	MA	1	Aug-19	Muan-gun, Jeollanam-do, Corn field; GPS: 34°52.21 N, 126°31.11 E	1 : 0
Papua New Guinea	Madang Province	Ramu Sugar Estate	16	15-17 Jun-2020	5°58.154 S, 145°53.252 E, Maize host	16 : 0
	Central Province	Yule Island Junction	1	5-Jun-20	corn host	1 : 0
Peru	n/a	Pre-border intercepted specimens on imported agricultural commodities	16	2016 - 2019	Samples from Tay et al. (2020) (Suppl. Table 1, individuals PE01 – PE16)	0 : 16

810 **Table 3:** Summary bioassay data on *Spodoptera frugiperda* populations from Walkamin, Queensland (Sf20-1) and Kununurra, Western Australia (Sf20-4), *S. litura*
811 (Mareeba) and *Helicoverpa armigera* (GR laboratory line) involving surface treatment of the diet with Bt toxins and products.

Pesticide	Populations/species	N	Slope	LC_{50}	95% CI	LC_{99}	95% CI	χ2	Toxicity ratio (*H. armigera* = 1)
Cry1Ac	*H. armigera*	741	1.837 ±0.126	0.025	0.021 - 0.029	0.465	0.316 - 0.773	31.66	-
($\mu g/cm^2$)	Sf20-1	504	0.827 ±0.102	4.34	1.81 - 8.15	80.6	36.85 - 249.17	18.38	174
	Sf20-4	575	1.449 ±0.149	2.48	1.79 - 3.29	85.34	53.32 - 159.68	17.77	99
	S. litura	501	1.626 ±0.135	2.99	2.06 - 4.20	80.7	38.60 – 284.54	49.14	120
Cry2Ab	*H. armigera*	476	1.501 ±0.143	0.049	0.034 -0.068	1.74	0.86 - 6.57	31.834	-
($\mu g/cm^2$)	Sf20-1	575	1.649 ±0.136	0.655	0.435 - 0.951	16.88	7.24.- 86.01	86.222	13
	Sf20-4	551	2.201 ±0.246	0.178	0.138 - 0.221	2.03	1.33 - 4.94	39.076	4
	S. litura	859	1.197 ±0.081	0.511	0.335 - 0.734	44.9	20.179 - 154.39	91.946	10
Cry1F	Sf20-1	576	1.531 ±0.115	0.025	0.015 - 0.038	0.838	0.365 - 3.949	19.947	-
($\mu l/cm^2$)	Sf20-4	575	1.809 ±0.149	0.021	0.014 - 0.029	0.394	0.211 - 1.138	87.309	-
	S. litura	574	1.524 ±0.121	0.0088	0.0069 - 0.011	0.294	0.187 - 0.535	58.968	-
VIP3a	*H. armigera*	763	1.486 ±0.093	0.0062	0.0051 - 0.0075	0.230	0.140 - 0.400	33.738	-
($\mu l/cm^2$)	Sf20-1	619	1.881 ±0.225	0.0021	0.0010 - 0.0031	0.049	0.021 - 0.257	68.846	0.152
	Sf20-4	599	2.398 ±0.212	0.0019	0.0016 - 0.0023	0.018	0.013 - 0.028	20.934	0.078

	S. litura	575	1.693 ±0.129	0.00065	0.0005 - 0.0008	0.015	0.088 - 0.035	39.108	0.065
Dipel	*H. armigera*	526	1.832 ±0.154	2.11	1.59 - 2.72	39.2	22.43 - 92.76	33.878	-
(IU/cm^2)	Sf20-1	644	1.832 ±0.154	52.02	36.55 - 70.05	2612.6	1379.89 - 6752.99	29.258	24
	Sf20-4	574	1.703 ±0.153	38.93	23.97 - 56.35	904.1	469.80 - 2779.63	53.767	18
	S. litura	788	1.426 ±0.098	35.96	28.23 - 44.99	1540.8	916.66 - 3110.88	40.916	17
XenTari	*H. armigera*	620	1.530 ±0.136	5.95	4.69 - 7.40	197.2	118.7 - 398.3	23.06	-
(DBM/cm^2)	Sf20-1	647	2.375 ±0.289	11.81	8.33 - 15.17	112.7	69.8 - 270.1	39.18	2
	Sf20-4	549	1.333 ±0.180	19.01	9.27 - 29.8	1058.9	379.5 - 10739.7	43.25	3
	S. litura	741	2.375 ±0.289	24.86	17.08 - 31.14	206.9	126.2 - 597.5	47.92	4

814 **Table 4:** Summary bioassay data on *Spodoptera frugiperda* populations and *Helicoverpa armigera* (GR laboratory line) involving topical application to the insect with

815 contact insecticides. Sf20-1 = *S. frugiperda* from Walkamin Queensland; Sf20-4 = *S. frugiperda* from Kununurra Western Australia.

Toxin	Population/Species	N	Slope	LC_{50}	95% C.I.	LC_{99}	95%CI	χ2	Toxicity ratio (*H. armigera* = 1)
Alpha cypermethrin	*H. armigera*	1328	2.849 ±0.140	0.0036	0.0032 - 0.0041	0.023	0.018 - 0.032	109.84	-
µg/larvae	Sf20-1	675	2.399 ±0.154	0.201	0.171 - 0.239	1.88	1.31 - 3.06	43.903	56
	Sf20-4	766	2.186 ±0.132	0.523	0.427 - 0.641	6.06	3.99 - 10.85	67.208	145
Methomyl	*H. armigera*	858	0.809 ±0.059	0.057	0.031 - 0.097	43.12	10.27 - 53.93	135.89	-
µg/larvae	Sf20-1	765	1.064 ±0.076	0.254	0.177 - 0.356	39.2	15.81 - 156.30	72.309	4
	Sf20-4	631	0.874 ±0.07	2.96	1.87 - 4.66	1363.7	380.11 - 10950.47	57.160	52

818 **Table 5:** Summary bioassay data on *Spodoptera frugiperda* populations (Sf20-1 from Walkamin Queensland; Sf20-4 from Kununurra Western Australia) and *Helicoverpa*

819 *armigera* (GR laboratory line) involving diet incorporation of insecticides.

Pesticide	Population/Species	N	Slope	LC_{50}	95% CI	LC_{99}	95%CI	χ2	Toxicity ratio (*H. armigera* = 1)
Chlorantraniliprole	*H. armigera*	540	3.199 ±0.234	0.011	0.009 -0.013	0.056	0.038 - 0.109	72.27	-
(µg/ml diet)	Sf20-1	613	2.484 ±0.191	0.032	0.024 - 0.043	0.28	0.162 - 0.777	99.976	3
	Sf20-4	897	2.065 ±0.152	0.163	0.132 - 0.201	2.19	1.40 - 4.10	99.895	15
Indoxacarb	*H. armigera*	653	1.684 ±0.122	0.054	0.028 - 0.089	1.29	0.49 - 12.59	226.42	-
(µg/ml diet)	Sf20-1	697	2.359 ±0.159	1.203	1.031 - 1.398	11.66	8.50 - 17.62	34.789	22
	Sf20-4	541	1.817 ±0.130	11.206	9.254 - 13.654	213.85	136.20 - 391.41	24.511	208
Emamectin	*H. armigera*	810	2.465 ±0.164	0.0107	0.0087 - 0.0131	0.0945	0.065 - 1.607	75.940	-
(µg/ml diet)	Sf20-1	631	2.606 ±0.175	0.0158	0.013 - 0.019	0.124	0.082 - 2.292	67.73	1
	Sf20-4	720	3.234 ±0.226	0.0179	0.016 - 0.020	0.094	0.723 - 1.331	40.00	2
Spinetoram	*H. armigera*	472	1.711 ±0.173	0.086	0.023 - 0.157	1.977	0.686 - 8.409	154.67	-
(µg/ml diet)	Sf20-1	831	3.220 ±0.198	0.118	0.101 - 0.137	0.623	0.457 - 0.990	94.168	1
	Sf20-4	542	4.921 ±0.489	0.102	0.092 - 0.112	0.301	0.247 - 0.400	13.119	1

Table 6: Number of *Spodoptera frugiperda* individuals (N) from populations from Australia (Kununurra SF20-4; Strathmore SF20-1), South Korea, Papua New Guinea, and Peru with homozygous susceptible (S/S), heterozygous (S/R) and homozygous resistance (R/R) profiles characterised via whole genome sequence data of the ACE-1 locus involving the A2001S, G227A, and F290V amino acid substitutions.

Population	N	ACE-1 (A201S)			ACE-1 (G227A)			ACE-1 (F290V)		
		S/S	S/R	R/R	S/S	S/R	R/R	S/S	S/R	R/R
Kununnura	9	9	0	0	9	0	0	2	4	3
Strathmore	30	26	4	0	30	0	0	0	0	30
South Korea	12	11	1	0	12	0	0	0	0	12
Papua New Guinea	17	15	2	0	17	0	0	1	12	4
Peru	16	4	12	0	10	6	0	2	14	0

Table 7: Comparisons of bioassay results for selected insecticides fed to laboratory populations of *Spodoptera frugiperda* via diet incorporation approaches. *S. frugiperda* populations were from the native ranges (USA, Louisiana State University (LSU) culture; see Hardke et al. 2011); Brazil SUS-2005, (Kulye et al. 2021) and introduced ranges from the original 2018 populations from Kanartaka, India (population I-2018; Kulye et al. 2021); a population from Groblersdal, Mpumalanga province, South Africa (see Discussion above, Eriksson 2019), and from our Sf20-1 and Sf20-4 laboratory populations from Queensland and Western Australia, Australia, respectively. Potency ratios of the invasive populations were measured against the native Brazilian (SUS-2005) population as reported by Kulye et al. (2021). Diet incorporation approach was for the Brazil, USA (LSU), India I-2018, and Australia (Sf20-1; Sf20-4) populations.

Insecticides	Populations	Range	LC$_{50}$ (ppm) / EC$_{50}$ (μg/mL)	Fiducial limits† (95%)	Slope ± SE	Toxicity ratios
Spinetoram	Brazil (SUS-2005)[#]	native	0.010	0.009 - 0.012	2.35 ± 0.37	-
	USA (LSU)	native	0.066	0.053 - 0.081	2.54 ± 0.36	6.6
	India (I-2018)[#]	invasive	0.013	0.012 - 0.014	2.25 ± 0.13	1.3
	Sf20-1 (Qld)	invasive	0.118	0.101 - 0.137	3.220 ±0.198	11. 8
	Sf20-4 (WA)	invasive	0.102	0.092 - 0.112	4.921 ±0.489	10.2
Chlorantraniliprole	Brazil[#]	native	0.005	0.005 - 0.006	4.54 ± 0.76	-
	USA (LSU)	native	0.068	0.317 - 0.481	2.55 ± 0.23	13.6
	India (I-2018)[#]	invasive	0.009	0.008 - 0.009	2.39 ± 0.11	1.8
	South Africa	invasive	0.06	0.01 - 0.16	1.01 ± 0.42	12
	Sf20-1 (Qld)	invasive	0.032	0.024 - 0.043	2.484 ± 0.191	6.4
	Sf20-4 (WA)	invasive	0.163	0.132 - 0.201	2.065 ±0.152	32.6
Indoxacarb	USA (LSU)	native	0.392	0.317 - 0.481	2.35 ± 0.25	-
	Sf20-1 (Qld)	invasive	1.203	1.031 - 1.398	2.359 ±0.159	3.07
	Sf20-4 (WA)	invasive	11.206	9.254 - 13.654	1.817 ±0.130	28.6

Note: †Fiducial limits are similar to confidence limits but are based on logistic growth or S-shaped curve distribution. #EC$_{50}$ reported instead of LD$_{50}$ by Kulye et al. (2021). In this comparison we assumed both EC$_{50}$ and LD$_{50}$ estimates to be broadly similar.

Table 8: Comparisons of topically applied methomyl bioassay results on late 3[rd] and/or early 4[th] instar larvae of *S. frugiperda* between resistant or susceptible populations from Gainesville Florida, USA (FL) (Yu 1991; see Discussion above), and invasive populations from South Africa (Eriksson 2019; see Discussion above) and Queensland or Western Australia, Australia.

Populations	LC$_{50}$ (ppm)	Fiducial limits (95%)	Slope ± SE	Toxicity ratios
USA (FL susceptible)	0.57	0.45 - 0.78	4.9	-
South Africa	62.98	17.22 – 52.12	11.14 ± 9.20	110.5
Sf20-1 (Qld)	0.254	0.177 - 0.356	1.064 ±0.076	0.45
Sf20-4 (WA)	2.96	1.87 - 4.66	0.874 ±0.07	5.2
USA (FL resistance)	8.23	4.86 - 11.6	2.9	-
South Africa	62.98	17.22 – 52.12	11.14 ± 9.20	7.65
Sf20-1 (Qld)	0.254	0.177 - 0.356	1.064 ±0.076	0.031
Sf20-4 (WA)	2.96	1.87 - 4.66	0.874 ±0.07	0.36

Figure 1 Caption

Figure 1: Summary of Acytylcholinesterase (ACE-1) susceptible and resistance allele frequencies in *Spodoptera frugiperda* populations from 17 countries (i.e., Benin, Uganda, Kenya, Tanzania, Zambia, Malawi, India, Indonesia, Papua New Guinea (PNG), South Korea, China, Australia) across the invasive range. A total of 1,246 individuals representing 76 populations in the Table S2 were used to compile the data below. Population identity is as provided in Table S2 that combined data from this study (Australia SF20-1 and SF20-4 (generation 0 representing field-collected individuals), PNG, South Korea) and from published studies (Boaventura et al. 2020a, Guan et al. 2020, Zhang et al. 2020, Zhao et al. 2020, Nguyen et al. 2021, Tay et al. 2021, Yainna et al. 2021, Rane et al. 2022). Susceptible and resistant alleles from the three previously reported loci (i.e., F290V, G227A, A201S) from the ACE-1 gene provided evidence to support multiple independent introductions across the invasive *S. frugiperda* populations, such as in Indonesia (#17; G227A), South Korea (#20; F290V), and China (Hubei province (#10-13); G227A) as indicated by the red arrows. In Australia, newly established *S. frugiperda* populations between Queensland (e.g., #27, #28 (Walkamin), #29 (Strathmore), #32 (Burdekin), and Western Australia (i.e., #2 (Kununurra)/Northern Territory (e.g., #21 (Bluey's Farm)) suggested this likely involved multiple introductions from diverse populations from neighbouring countries and likely arrived via separate pathways and entry points (see also Rane et al. 2022).

Figure 1