A single cell atlas of the cycling murine ovary

Morris ME[1#*]**,** Meinsohn MC[2,3*], Chauvin M[2,3], , Saatcioglu HD[2,3], Kashiwagi A.[2,3], Sicher NA. [2,3], Nguyen

NMP[2,3], Yuan S[2,3], Rhian Stavely[2,3], Hyun M[4,] Donahoe PK[2,3], Sabatini B[4], Pépin D[2,3#]

[1]Department of Gynecology and Reproductive Biology, Massachusetts General Hospital, Boston/MA,

[2]Pediatric Surgical Research Laboratories, Massachusetts General Hospital, Boston/MA,[3]Department of

Surgery, Harvard Medical School, Boston/MA, [4]Howard Hughes Medical Institute, Department of

Neurobiology, Harvard Medical School, Boston/MA

Co-corresponding

* Contributed equally

Abstract

The estrous cycle is regulated by rhythmic endocrine interactions of the nervous and reproductive systems,

which coordinate the hormonal and ovulatory functions of the ovary. Folliculogenesis and follicle

progression require the orchestrated response of a variety of cell types to allow the maturation of the follicle

and its sequela, ovulation, corpus luteum (CL) formation, and ovulatory wound repair. Little is known about

the cell state dynamics of the ovary during the estrous cycle, and the paracrine factors that help coordinate

this process. Herein we used single-cell RNA sequencing to evaluate the transcriptome of > 34,000 cells

of the adult mouse ovary and describe the transcriptional changes that occur across the normal estrous

cycle and other reproductive states to build a comprehensive dynamic atlas of murine ovarian cell types

and states.

Introduction

The ovary is composed of a variety of cell types that govern its dynamic functions as both an endocrine

organ capable of producing hormones such as sex steroids, and a reproductive organ orchestrating the

development of follicles, a structure defined by an oocyte surrounded by supporting somatic cells such as

granulosa cells and theca cells. Most follicles in the ovary are quiescent primordial follicles, representing

the ovarian reserve. Once activated, a primordial follicle grows in size and complexity as it progresses to

primary, preantral, and antral stages, adding layers of granulosa and theca cells and forming an antral

cavity, until it ultimately ejects the oocyte-cumulus complex at ovulation while the follicular remnants

undergo terminal differentiation to form the corpus luteum (1). This process necessitates precise

coordination of germ cells and several somatic cell types, including granulosa cells, thecal cells, vascular

cells, and other stromal cells of the ovary to support the growth of the oocyte until its ovulation or follicular

atresia. In addition to supporting germ cells, ovarian somatic cells must produce the necessary hormonal

cues, as well as coordinate the profound tissue remodeling, necessary to accommodate these dynamic

developing structures. For reproductive success to occur the state of each of these cells must change in a

coordinated fashion over the course of the estrous cycle; this allows waves of follicle to grow and mature,

ovulation to be triggered precisely, and provides the hormonal support necessary for pregnancy.

Single cell RNA sequencing (scRNAseq) has been used in a variety of tissues to obtain an in-depth

understanding of gene expression and cellular diversity. In the ovary, this technique has allowed us, and

others, to explore various physiologic processes during early ovarian development and ovarian aging (2–

10). For example, Fan et al. catalogued the transcriptomic changes that occur during follicular development

and regression and mapped the cell types of the human ovary using surgical specimens (9). A primate

model has been used to investigate changes in cell types and states that occur in the ovary with aging (10).

Zhao et al. looked at the development of the follicle during early embryonic ovarian development to discern

the relationship of oocytes to their support cells in formation of follicles (2). We have used scRNAseq to

identify inhibitory pathways regulated by AMH during the first wave of follicular growth in the murine ovary

(8). While all these studies have helped establish a static framework to understand the major cell types in

the ovary, they fail to describe the dynamic nature of cell states across the reproductive cycle, known as

the estrous cycle. The estrous cycle in mice is analogous to the human menstrual cycle, which both reflect

follicle development in the ovary. In mice this cycle lasts four to five days and is composed of 4 different

phases known as proestrus, estrus, metestrus and diestrus. The murine proestrus is analogous to the

human follicular stage and leads to ovulation at estrus. Metestrus and diestrus are analogous to early and

late secretory stages of the reproductive cycle in humans, which are orchestrated by secretion of

progesterone by the corpus luteum (11).

To understand more fully the dynamic effects of cyclic endocrine, autocrine, and paracrine signals on

ovarian cell states we performed high-throughput single cell RNA sequencing of ovaries from adult mice

across a physiological spectrum of reproductive states. Ovaries were harvested from mice in the four

phases of the normal estrous cycle: proestrus, estrus, metestrus, and diestrus. Additionally, ovaries were

evaluated from mice that were lactating or non-lactating 10 days postpartum mice, and from randomly

cycling adult mice to increase the diversity of cell states. Herein, we describe the previously unrecognized

complexity in the ovarian cellular subtypes and their cyclic expression states during the estrous cycle and

identify secreted factors that represent potential marker for staging.

Results:

scRNA-seq of adult mouse ovaries across reproductive states

To survey the dynamic transcriptional landscape of ovaries at the single cell level across a range of

physiological reproductive states in sexually mature female mice, we isolated the ovaries (4 mice per group)

at each stage of estrous cycling (proestrus, estrus, metestrus, diestrus), post-partum non-lactating (day 10

post-partum, with pups removed on the day they were born), post-partum lactating (day 10 postpartum,

actively lactating with pups), and non-monitored mice to increase sample diversity and cell counts.

Following enzymatic digestion of the ovaries, we generated single cell suspensions and sorted them by

microfluidics using the inDROP methodology (15), targeting 1500 cells per animal. Resulting libraries were

indexed and combined for sequencing (**Figure 1A**).

Following dimensionality reduction and clustering using the Seurat algorithm we identified multiple clusters

which could be combined to represent the major cell categories of the ovary (**Figure 1B**). To assign cell

type identity we used cluster-specific markers which were previously described in other studies or newly

identified makers later validated by RNA in situ (**Supplementary File 2**). The largest groups of clusters

consisted of granulosa cells (N=17627 cells) and mesenchymal cells of the ovarian stroma (N=10825 cells).

Other minor cell types were identified including endothelial cells (N= 3501 cells), ovarian surface epithelial

cells (N=1088 cells), immune cells (N=1649 cells), and oocytes (N=22 cells), altogether recapitulating all

the major cell types of the ovary (Figure 1 – figure supplement 1A). Oocytes were poorly represented in the

dataset due to cell size limitations of inDROP, likely restricting our sampling to small oocytes of primordial

follicles (**Figure 1B**). To better characterize the transcriptional signatures of the identified cell types we

evaluated a heatmap of marker gene expression across the major categories of cell types and states

(**Figure 1C)**. Cells were also classified depending on the stage of the estrous cycle or lactating states in

which the ovaries were collected (**Fig1B)**. Morphological differences between the stages of proestrus,

estrus, metestrus, diestrus, and also post-partum lactating and non-lactating, were documented in **Figure**

1- figure supplement 1B. The granulosa, mesenchyme, and epithelium clusters were isolated and

reanalyzed to identify subclusters.

Single cell sequencing reveals heterogeneity within granulosa and mesenchymal cell clusters

Cellular diversity of Mesenchymal cells

The mesenchymal cluster was the second largest cluster identified in our analysis. Based on prior studies

and conserved marker expression (9,10) we were able to identify subclusters within mesenchymal cells,

and their relative abundance (percentage) as follows : early theca (16.8%), which formed the theca interna

of preantral follicles, steroidogenic theca (13.2%), which formed the theca interna of antral follicles, smooth

muscle cells (10.2%), which were part of the theca externa of both antral and preantral follicles, pericytes

(6.2%), which surrounded the vasculature, and two interstitial stromal cell clusters, one composed of

steroidogenic cells (28.7%) and the other of fibroblast-like cells (24.9%), which together constituted the bulk

of the ovarian volume (outside of follicles). These subclusters can be seen in **Figure 2A,** with the top 5

expressed markers of each subcluster described in the **Figure 2B** heatmap and the top 10 listed in

Supplementary File 3.

Distinct transcriptional signatures were identified in each of these mesenchymal subclusters (**Figure 2B**);

to confirm the presumed identity and histology of these cell types (detailed in **Figure 1- supplement 1A**) ,

we validated markers prioritized by highest fold-change expression, highest differential percent expression,

and lowest P-value **(Figure 2C)**.

For the theca interna, the two clusters identified reflected the stage of development of the follicle: early

thecal cells could be defined by their expression of Hedgehog Interacting Protein (*Hhip)* and were

histologically associated with preantral follicles*;* meanwhile, the steroidogenic theca cells were identified by

their expression of Cytochrome P450 Family 17 Subfamily A Member 1 *(Cyp17a1),* an essential enzyme

for androgen biosynthesis (20)*,* and were found in antral follicles (**Figure 2C)**. The theca externa is a

connective tissue rich in extracellular matrix situated on the outermost layer of the follicle (**Figure 1-**

supplement 1A), constituted of fibroblasts, macrophages, blood vessels, and abundant smooth muscle

cells which we identified based on their expression of Microfibril Associated Protein 5 (*Mfap5)* by RNA in

situ hybridization (**Fig 2C**). To validate the identity and histology of these smooth muscle cells, we

performed RNAish/IHC co-localization of *Mfap5* and Actin Alpha 2 (*Acta2),* another marker of smooth

muscle, which confirmed their position within the theca externa. In contrast, *Hhip,* which was expressed in

theca interna (both immature and steroidogenic), didn't colocalize with Acta2 (**Figure 2- supplement 1A,**

B, C). These results suggest Mfap5+ exquisitely labels smooth muscle cells of the theca externa, which are

thought to perform a contractile function during ovulation (21).

Lastly, the bulk of the ovarian interstitial stromal space was made up of two closely related cell types which

could not be differentiated by specific dichotomous markers, but rather distinguished based on relative

expression of Ectonucleotide Pyrophosphatase/Phosphoiestrase 2 (*Enpp2*) (Fig 2C). While Enpp2+ cells

represented fibroblast-like stromal cell, Enpp2- interstitial cells were enriched for expression of genes such

as Patch1 (Ptch1), a member of the hedgehog signaling pathway, known as an important regulator of

ovarian steroidogenesis (22), suggesting these represented steroidogenic stromal cells. Indeed, the

steroidogenic activity of this stromal cell cluster was further confirmed by its high relative expression of

other genes associated with steroidogenesis including Cytochrome P450 Family 11 Subfamily A Member

1 (*Cyp11a1*), Hydroxy-Delta-5-Steroid Dehydrogenase, 3 Beta- And Steroid Delta-Isomerase 1 (*Hsd3b1*),

Cytochrome P450 Family 17 Subfamily A Member 1 (*Cyp17a1),* Steroid 5 Alpha-Reductase 1

(*Srd5a1),* along with other markers such as Potassium Two Pore Domain Channel Subfamily K Member 2

(*Kcnk2*) **(Figure 2- supplement 1E,F)**. In contrast the fibroblast stromal cluster had enriched expression of

many extracellular matrix genes such as Collagen Type I Alpha 1 Chain (*Col1a1*), Collagen Type V Alpha

1 Chain (*Col5a1*), Lumican (*Lum*), Lysyl Oxidase Like 1(*Loxl1*) (23) along with other known markers of

fibroblasts such as C-X-C Motif Chemokine Ligand 14 (*Cxcl14*) (24) and WT1 Transcription Factor (*Wt1*)

(25) **(Figure 2-supplement 1E,F)**.

The identities of all above-described mesenchymal clusters matched previously reported markers such as

Desmin (Des) for pericytes (26), Steroidogenic Acute Regulatory protein (Star) for the steroidogenic theca

(27), Cellular Communication Network Factor 1 (Ccn1) for the smooth muscle (28), Receptor Activity

Modifying Protein 2 (Ramp2) for the early theca cells surrounding preantral follicles (29) and finally C-X-C

motif chemokine ligand 12 (Cxcl12) for both stroma clusters (30) as illustrated in **Fig2 – supplement 1D)**.

Cellular diversity of Granulosa cells

To further explore the cellular heterogeneity within developing follicles (listed in **Figure1-supplement 1A**),

we investigated the sub-clustering of granulosa cells and their transcriptional profile. Consistent with

previous reports, we could distinguish discrete granulosa cell states in follicles based on their stage of

development (2,9,31). Granulosa cells could be subdivided into 8 main categories: Preantral-Cumulus

(27.3%), Antral-Mural (21.8%), Luteinizing mural (4.8%), Atretic (22.6%), Mitotic (14.4%), Regressing

corpus luteum (CL) (3.7%), Active CL (5.4%), (**Figure 3A). Supplementary file 4** lists the top ten markers

for each of these clusters. Distinctive gene expression programs were identified in the granulosa cell

subclusters, as visualized in the heatmap (**Figure 3B**), from which we selected potential markers for

validation.

Early pre-antral granulosa cells, and those constituting the cumulus of antral follicles, could be identified by

their shared expression of markers such as potassium channel tetramerization domain (*Kctd14*) (**Figure**

3C), which we had previously shown to be expressed by pre-antral follicles (8). In contrast, mural granulosa

cells of antral follicles expressed distinct markers (**Supplementary file 4**) such as *Male-specific*

Transcription in the developing Reproductive Organs (*Mro*) (**Figure 3C**). Luteinizing mural granulosa cells

could be identified by the expression of previously established markers (**Supplementary File 4**) and

Oxytocin Receptor gene (*Oxtr*) which we propose as a highly specific marker for this cell type, the likely

target of the surge in oxytocin during estrus (32)(**Figure 3C**). Furthermore, we identified two different

clusters that we hypothesize represent cell states of the corpus luteum, either active or regressing, which

both expressed Nuclear Paraspeckle Assembly Transcript 1 (*Neat1*), a known marker of *CLs* (33)*.* To

confirm the active and regressing CL cell states we investigated the expression of Top2a, a mitotic marker

(34), which was enriched in the active CL cluster, and Cdkn1a, a cell cycle exit and senescence marker

(35), which was enriched in the regressing cluster (**Figure 3- supplement 1B, C**). Moreover, when

identifying the composition of clusters depending on the reproductive stage, the regressing corpus luteum

cluster was found to be composed mostly of cells derived from the post-partum non-lactating (PPNL)

samples (**Figure 3- supplement 1E**), and overexpressed markers related to corpus luteum regression (36)

(**Figure 3 – supplement 1F**), consistent with a post-partum anestrous state in these mice. Finally, two

relatively abundant granulosa cell states could be identified based on marker expression: mitotic granulosa

cells could be found in both preantral and antral follicles, and were defined by their expression of *Top2a*,

and atretic granulosa cells, which expressed markers consistent with follicular atresia and apoptosis such

as Phosphoinositide-3-Kinase Interacting Protein 1 (*Pik3ip1*), Nuclear Protein 1, Transcriptional Regulator

(*Nupr1*), Growth Arrest And DNA Damage Inducible Alpha (*Gadd45a*), Vesicle Amine Transport 1 (*Vat1*),

Transgelin (*Tagln*), Melanocyte Inducing Transcription Factor (*Mitf*) (37)(**Figure 3C, Figure 3 –**

supplement 1A, Supplementary File 4). Furthermore, we propose growth hormone receptor (*Ghr*), which

was highly specific to this cluster, as a specific marker of atretic follicles, which warrant further investigation

of the role of growth hormone in this process **(Figure 3C)**.

Cellular states in the ovarian surface epithelium

The epithelial cluster was composed of 1088 ovarian surface epithelium cells, which could be further

subdivided into two clusters (**Figure 4A**), the larger one composed of non-dividing epithelium cells (96%)

and a smaller cluster (4%) composed of mitotic epithelium characterized by proliferation markers such as

Thymidine Kinase 1 (*Tk1*) (38), Rac GTPase-activating protein 1 (*Racgap1*)(39), Top2a, Casein kinase 1

(*Ck1*)(40), Protein Regulator Of Cytokinesis 1 (*Prc1*)(41), Ubiquitin Conjugating Enzyme E2 C (*Ube2c*)(42),

Baculoviral IAP Repeat Containing 5 (*Birc5*)(43) (**Figure 4A, B**). Interestingly, the proliferating subcluster

of ovarian surface epithelium was almost exclusively composed of cells in the estrus stage (**Figure 4C,D**),

consistent with their transient amplification during ovulatory wound closure (44).

Granulosa cell transcriptome is most dynamic during the proestrus/estrus transition

To identify changes in cell states associated with the stages of the estrous cycle we focused on the

granulosa cell subclusters, given the importance of follicular maturation in coordinating this process

(illustrated in **Figure 1 – supplement 1B**). When comparing the composition of granulosa cell subclusters

by estrous stage, we found that some clusters were dominated by cells from either the proestrous and

estrous samples, particularly the clusters corresponding to "antral/mural" and "periovulatory" clusters

respectively (**Figure 5A,B**). A volcano plot analysis confirmed that the transition between these two stages

was characterized by 24 significantly upregulated and 10 significantly downregulated markers (**Figure 5C**),

which together with the transition from estrus to metestrus, represents the largest change in gene

expression. In contrast, few genes were found to significantly change during the transition from metestrus

to diestrus, or diestrus to proestrus (**Figure 5 – supplement 1A**). Gene ontology analysis revealed that

the most significantly differentially regulated pathways between the proestrous and estrous phases were

related to ovarian matrix remodeling and steroidogenesis and hormones production (**Figure 5 –**

supplement 1B). To validate the genes with significant changes in expression identified within the single

cell sequencing dataset, we performed qPCR on whole-ovary samples at the proestrus to estrus transition,

including the steroid biosynthesis markers Cytochrome P450 Family 19 Subfamily A Member 1 *(Cyp19a1*,

p=0.0029, proestrus to estrus*)*, Steroidogenic Acute Regulatory Protein (*Star,* p=0.0187, proestrus to

estrus), Serum- and glucocorticoid-inducible kinase-1 (*Sgk1*, p=0.0056, proestrus to metestrus)*,* as well as

matrix remodeling genes such as Regulator Of Cell Cycle (*Rgcc*, p=0.0441, proestrus to estrus), Tribbles

Pseudokinase 2 (*Trib2*, p=0.0023, proestrus to estrus) (**Figure 5D,E**), and immediate early genes, Fos

Proto-Oncogene (*Fos*), Jun Proto-Oncogene (*Jun*, p=0.0022, proestrus to estrus), Jun Proto-Oncogene B

(*Junb*, p=0.0069, proestrus to diestrus) and Early Growth Response 1 (*Egr1*, p=0.0504 estrus to diestrus),

which represent a family of genes thought to be involved in wound repair, a sequela of ovulation (45–48)

(Figure 5 – supplement 1C). Transcriptional gene expression changes were found to be concordant

between the scRNAseq data and whole-ovary transcripts quantified by qPCR.

Identification and validation of secreted biomarkers varying throughout the estrous cycle

To identify new biomarkers that vary as a function of the estrous cycle and that could be used clinically for

staging in reproductive medicine, we screened for differentially expressed secreted factors (DAVID

Bioinformatics Resources)(49,50), which would therefore be potentially measurable in the blood.

Furthermore, to ensure specificity, we prioritized genes expressed specifically in the granulosa or ovarian

mesenchymal clusters and not highly expressed in other tissues based on GTEX profile

(51)(**Supplementary file 5**). As a primary screen, we first validated our ability to detect gene expression

changes by estrous stage using whole-ovary qPCR analysis in a sperate set of staged mice (N=4 per

group). Whole ovary qPCR successfully detected expression changes of estrous cycle markers such as

luteinizing hormone/choriogonadotropin receptor (52)(*Lhcgr*, p=0.0281 estrus to metestrus) and

progesterone receptor (*Pgr,* p=0.0096, proestrus to estrus) (53) (**Figure 6B**). Using this method, we

validated a set of significantly upregulated secreted markers in the proestrous to estrous transition, most

prominent of which were natriuretic peptide C (*Nppc,* p=0.0022 proestrus to estrus) and inhibin subunit

beta-A *(Inhba,* p= 0.0067, proestrus to estrus*)* (**Figure 6A,B**). Similarly, Tubulointerstitial Nephritis Antigen

Like 1 (*Tinagl1*) and Serine Protease 35 (*Prss35*) were secreted markers significantly upregulated in estrus

compared to their level of transcription in proestrus in the scRNAseq dataset (**Figure 6A**), and by qPCR

(*Tinagl1* p=0.0081 proestrus to estrus, *Prss35* p=0.0008 proestrus to estrus) (**Figure 6B**). In situ RNA

hybridization showed that, as expected, these markers were mostly expressed in mural granulosa cells of

antral follicles, while *Nppc* was expressed in both mural and cumulus cells (**Figure 6C**).

To evaluate the feasibility of measuring the secreted PRSS35, NPPCC, TINAGL1 and Activin A proteins in

the serum for staging we performed ELISA in mice at each stage of the estrous cycle (**Figure 6D**). We

found that the Activin A concentration in the serum was significantly increased between the diestrous and

proestrous stages (p=0.0312) and peaked at the proestrous stage (**Figure 6D**). The *Inhba* transcript, which

encodes for the Activin and Inhibin beta-A subunit, had a similar temporal expression profile (**Figure 6B**).

Circulating PRSS35 levels were lowest at the metestrous stage and were significantly increased during the

transition to diestrus (p=0.0009) and remained significantly elevated until the proestrus (**Figure 6D**). In

contrast, the Prss35 transcript was significantly induced earlier at estrus (**Figure 6B**). The serum

concentrations of TINAGL1, which was lowest at the diestrous and proestrous stages, was significantly

increased during the transition between proestrus and metestrus, peaking in estrus (p=0.0142) (**Figure**

6D)*.* This temporal pattern of expression was recapitulated at the transcriptional level by qPCR and

scRNAseq (**Figure 6A,B**). Finally, we observed a trend for serum protein concentrations of NPPC to be

lowest at the proestrous and estrous stages and increase during the metestrous and diestrous stages

(**Figure 6D**), although the differences were not statistically significant (p=0.0889, estrus to metestrus).

Importantly, these data provide a proof of concept that four markers could be used to monitor estrous cycle

progression when measured in conjunction in the blood (**Figure 6E**).

Discussion

Single cell RNA sequencing has been used to catalog the transcriptomes of a variety of tissues in several

species, across different physiological states (54). Herein, we used scRNAseq to survey the cellular

diversity and the dynamic cell states of the mouse ovary across the estrous cycle and other reproductive

states such as post-partum lactating and post-partum non-lactating.

The most significant changes in composition and cell states were associated granulosa cells, particularly

as they cycled through the estrous stages, reflective of their important role in cyclic follicular maturation and

hormone production. Early pre-antral follicle numbers are tought to be relatively stable across the estrous

cycle (55), given that they are largely unresponsive to gonadotropins (56), in contrast antral follicles, whose

numbers and size are more variable (55). Indeed, while subclusters such as "pre-antral granulosa cells"

were equally represented in samples from proestrus, metestrus, and diestrus, others, such as the

"luteinizing mural" cluster, were dominated by cells derived from one stage (in this case "estrus"). Genes

enriched in this cluster had been previously reported to be involved in the ovulatory process and regulated

by the LH surge, including markers of terminal differentiation and steroidogenesis such as *Smarca1* (57),

Cyp11a1 (58), Metallothionein 1 (*Mt1*), and Metallothionein *2* (*Mt2*) (59) (**Supplementary file 4**). Other

genes enriched in this subcluster include Prss35 (60) and Adamts1 (61,62), which had previously been

identified as playing a role in the follicular rupture necessary for ovulation.

Interestingly, we found that granulosa cells of preantral follicles and cumulus cells of antral follicles clustered

together and shared markers that distinguished them from mural granulosa cells. For example, Kctd14, a

member of the Potassium Channel Tetramerisation Domain Containing family, was expressed in granulosa

cells during the initial growth of early pre-antral follicles, but also specifically expressed only in cumulus

cells, but not mural granulosa cells, of larger antral follicles. Intriguingly, we've previously shown that anti-

Müllerian hormone (AMH, a.k.a. MIS), which is specifically expressed by cumulus cells (63) in antral

follicles, regulates KCTD14 expression in preantral follicles (8). This conservation of cellular state and

marker expression from preantral granulosa cells to cumulus cells of antral follicles suggests a continuous

lineage, potentially defined and maintained by the close interaction with the oocyte (63). This interpretation

is consistent with the presence of a differentiation fork in the granulosa cell lineage during antrum formation,

which would give rise to a distinct mural granulosa cell fate poised to respond to the LH surge. Indeed, the

peri-ovulatory granulosa cell state was identified based on its expression of genes regulated by LH such as

Smarca1 (57), and we propose Oxtr as a specific marker for these cells (**Figure 3- supplement 1D**). Oxtr

expression was found only in the mural cells of large Graafian follicles, suggesting it indeed corresponds

to an LH-stimulated mural granulosa cell state.

After ovulation, these LH-stimulated mural granulosa cells, along with the steroidogenic theca cells,

terminally differentiate into luteal cells and form the corpus luteum. The corpus luteum is a transient

structure with highly active steroid biosynthesis, providing the progesterone to maintain pregnancy (64). In

absence of implantation the CL degenerates (65). We found this progression of the corpus luteum to be

recapitulated at the transcriptional level, leading to two luteal subclusters: Active CL and Regressing CL.

While the Active CL was characterized by expression of proliferation markers (Top2a) in addition to

steroidogenic enzymes, the Regressing CL expressed the cell cycle inhibitor and senescence maker

Cdkn1a, along with luteolysis markers such as Syndecan 4 (Sdc4), Claudin Domain Containing 1 (Cldnd1),

and BTG Anti-Proliferation Factor 1 (Btg1) (36,66). The distinct expression signatures observed in these

two clusters may provide insights into the molecular basis of luteolysis and warrant further investigation.

The mesenchymal cluster was also surprisingly complex and variable across the estrous cycle, reflecting

its remodeling and physiological functions during follicle growth, steroid hormone production, and ovulation.

For example, during follicle maturation, the ovarian stroma adjacent to the developing follicle differentiates

into the theca, which is ultimately responsible for steroid hormone biosynthesis and therefore underlies the

cyclic hormone production of the ovary (67). Herein we identified two thecal clusters, designated as early

and steroidogenic theca. Early theca, was defined by markers such as Hhip (20,68), Mesoderm Specific

Transcript (Mest) (9), and Patched 1 (Ptch1) (9,20), and given its association with small follicles, presumed

to be immature and the precursor to steroidogenic theca. As the follicle matures and the antrum forms, this

layer becomes more vascularized and differentiates into theca interna, which is steroidogenic. This

steroidogenic theca cluster, was readily identifiable through its expression of steroidogenic enzymes such

as Hydroxy-Delta-5-Steroid Dehydrogenase, 3 Beta- and Steroid Delta-Isomerase 1 (Hsd3b1), Cyp17a1,

Cyp11a1, and also well-established markers such as Ferredoxin-1 (Fdx1) and prolactin receptor (Prlr)

(9,69). Interestingly, we confirmed the presence of steroidogenic interstitial stromal cells which also likely

contribute to sex steroid production in the ovary. Indeed, such cells likely represent the precursors of the

theca interna (70,71). Smooth muscle cells, which are part of the theca externa, were identified by their

expression of structural proteins such as Mfap5, Myosin Heavy Chain 11 (Myh11), Transgelin (Tagln), and

Smooth Muscle Actin (Sma or Acta2) (2). In contrast to mice, human smooth muscle cells are thought to

express high levels of collagen (9), which we did not observe. Another species difference between mice

and humans was expression of Aldehyde Dehydrogenase 1 Family Member A1 (Aldh1a1) which we found

primarily in the steroidogenic theca cluster, while it is presumably enriched in the theca externa in humans

(9).

Ovulation is associated with a dramatic remodeling of the ovary, including the subsequent ovulatory wound

repair. We identified fibroblast-like cells in the ovarian stroma expressing many of the extracellular matrix

protein known to play a role in these processes (44,72). Another important player in ovulatory would repair

is the ovarian surface epithelium (OSE), a simple mesothelial cell layer that covers the surface of the ovary,

and must dynamically expand to cover wound (73,74). The OSE cluster could be identified based on well-

established markers such as keratin (Krt) 7, 8 and 18 (75) and was represented by only 3% of all cells in

our dataset, which could be further sub-divided in proliferative and non-proliferative states. As expected

from their function in ovulatory wound repair, dividing OSE where enriched during estrus. furthermore,

genes associated to wound healing such as Galectin 1 (*Lgals1*) (76), were also significantly upregulated in

estrus. Similarly, the expression of the immediate-early genes Fos, Jun, Junb, and Egr1 was variable during

the estrous cycle, following a common pattern of strong downregulation at estrous compared to the other

stages, consistent with their temporal expression during the repair of other tissues such as the cornea (77).

Finally, to take advantage of this rich dataset, we sought to identify secreted markers which varied in

abundance during the estrous cycle and could thus be used as staging biomarkers in assisted reproduction.

We identified and prioritized four secreted biomarkers, expressed in mouse but also human ovaries, which

varied significantly during different transitions of the estrous cycle, namely *Inhba* (78), *Prss35* (60,79), *Nppc*

(80,81), and *Tinagl1* (82,83).

Activin A is a secreted protein homodimer translated from the *Inhba* transcript that is a crucial modulator of

diverse ovarian functions including pituitary feedback, whose expression level depends highly on the stage

of the estrous cycle (84). Quantification of Activin A protein in the serum by ELISA revealed elevated levels

in the blood during both proestrus and estrus, which is consistent with studies in other species such as

ewes (85). Importantly, the protein product of *Inhba*, the inhibin beta-A subunit, can be incorporated into

other protein dimers, such as Activin BA, and Inhibin A, which were not measured in this study and may

also represent cycling biomarkers.

The serine protease 35 transcript was expressed in the theca layers of preantral follicles and induced in

granulosa cells of preovulatory follicles and all stages of the corpora lutea, peaking at the estrous stage

according to qPCR, leading us to speculate that it may be involved tissue remodeling during ovulation and

corpus luteum formation (60). In contrast, the PRSS35 protein levels were highest in the diestrus and

proestrus stages as determined by ELISA, suggesting other tissue sources of PRSS35 or again an offset

in peak protein levels due to delays in accumulation of the protein in the circulation.

The natriuretic peptide precursor C (NPPC) protein is a peptide hormone encoded by the *Npcc* gene. *Nppc*

has been reported to be expressed by mural granulosa cells while its receptor Npr2 is expressed by

cumulus cells(80). The pair acts on developing follicles by increasing the production of intracellular cyclic

guanosine monophosphate and maintains oocyte meiotic arrest during maturation; upon downregulation of

this pathway, the oocyte can escape meiotic arrest and ovulate (86). This close relationship with the

ovulatory process makes *Nppc* an attractive marker to predict ovulation. Herein, qPCR analysis revealed

that *Nppc* was highest in the ovary at proestrus, and was quickly and significantly downregulated at estrous,

probably in response to the increased levels of LH which in turn inhibit the Nppc/Npr2 system (87). In

contrast there was a trend for the circulating NPCC peptide to be highest in metestrus and diestrus, albeit

not in a statistically significant way.

Finally, we evaluated the level of transcription and protein expression of the matricellular factor Tinagl1. We

found both the *Tinagl1* transcript and the circulating TINAGL1 protein in the blood to be highest during

estrous, thus coinciding with ovulation, with a pattern of expression consistent with expression by mural

granulosa cells of antral follicles. While the role of TINAGL1 in the ovary has not been extensively

investigated, it has been associated with delayed ovarian collagen deposition and increased ovulation in

aging Tinagl1 knock-out mice (82).

These four potential cyclic biomarkers provide a proof of concept that a deeper understanding of changes

at the single cell transcriptomics may translate into useful applications in assisted reproduction. It will be of

interest to follow up the findings of cyclic expression of Activin A, PRSS35, NPPC, and TINAGL1,

particularly in combination as an index, for use in humans and other species for the purpose of staging and

predicting ovulation timing.

In summary, this study outlines the dynamic transcriptome of murine ovaries at the single cell level and

across the estrous cycle and other reproductive states and extends our understanding of the diversity of

cell types in the adult ovary. We identified herein novel biomarkers of the estrous cycle that can be readily

measured in the blood and may have utility in predicting staging for assisted reproduction. This rich dataset,

and extensive validation of new molecular markers of cell types of the ovary, will provide a hypothesis-

generating framework of cell states across the cycle with which to elucidate the complex cellular interactions

that are required for ovarian homeostasis.

Materials and Methods

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
genetic reagent (*Mus musculus*)	C57BL/6-Tg(UBC-GFP)30Scha/J	Jackson Laboratory	stock #004353	
Antibody	Smooth muscle alpha action (SMA) (Rabbit polyclonal)	Abcam	#5694	Dilution: 1:300
Commercial assay or kit	ACTIVIN A commercial ELISA	RnD systems	#DAC00B	
Commercial assay or kit	NPPC commercial ELISA	Novus Bio	#NBP2-75790	
Commercial assay or kit	Tinagl1 commercial ELISA	LS-Bio	#LS-F49684	
Commercial assay or kit	PRSS35 commercial ELISA	Mybiosource	#MBS9717242	
Commercial assay or kit	RNA scope 2.5 HD Duplex detection kit	ACD bio	#322500	

Commercial assay or kit	RNA scope 2.5 HD red detection kit	ACD bio	#322360	
Commercial assay or kit	the target retrieval and protease plus reagents	ACD bio	#322330	
Other	Cdkn1a (M.musculus) NM_007669.4	ACD bio	# 408551	RNAscope probe
Other	Cxcl14 (M.musculus) NM_019568.2	ACD bio	#459741	RNAscope probe
Other	Cyp11a1 (M. musculus) NM_019779.4	ACD bio	#809181	RNAscope probe
Other	Cyp17a1 (M. musculus) NM_007809.3	ACD bio	#522611	RNAscope probe
Other	Ghr (M. musculus) NM_010284.3	ACD bio	#464951	RNAscope probe
Other	Hhip (M.musculus) NM_020259.4	ACD bio	#448441	RNAscope probe
Other	Inhba (M.musculus) NM_008380.1	ACD bio	# 455871	RNAscope probe

Other	Kcnk2 (M.musculus) NM_0011598 50.1	ACD bio	#440421	RNAscope probe
Other	Kctd14 (M.musculus) NM_0011362 35.1	ACD bio	#517811	RNAscope probe
Other	Mfap5 (M.musculus) NM_015776.2	ACD bio	#490211	RNAscope probe
Other	Mro (M.musculus) NM_0013058 82.1	ACD bio	#491181	RNAscope probe
Other	Neat1 (M.musculus) NR_003513.2	ACD bio	#440351	RNAscope probe
Other	Nppc (M.musculus) NM_010933.5	ACD bio	# 493291	RNAscope probe
Other	Onecut2 (M.musculus) NM_194268.2	ACD bio	#520541	RNAscope probe
Other	Oxtr (M.musculus) NM_0010811 47.1	ACD bio	#412171	RNAscope probe
Other	Prss35 (M.musculus) NM_178738.3	ACD bio	#492611	RNAscope probe
Other	Runx1 (M. musculus) NM_0011110 21.1	ACD bio	#406671	RNAscope probe

Other	Tinagl1 (M.musculus) NM_0011683 33	ACD bio	#312621	RNAscope probe
Other	Top2a (M.musculus) NM_011623.2	ACD bio	# 491221	RNAscope probe
Other	Wt1 (M.musculus) NM_144783.2	ACD bio	#432711	RNAscope probe
Software, algoritm	R version 4.1.3	R Project for Statistical Computing	https://scicrunch.org/resolver/SCR_001905	
Software, algoritm	Seurat package 4.1.0	R toolkit for single cell genomics	https://satijalab.org/seurat/articles/install.html	
Software, algoritm	BZ-X800 analysis software	Keyence	https://www.keyence.com/landing/microscope/lp_fluorescence.jsp	
Software, algoritm	GraphPad Prism, version 9.2.0	Graphpad		

Mice

Animal experiments were carried out in 6-8 weeks old C57BL/6 mice obtained from Charles River

Laboratory, approved by the National Institute of Health and Harvard Medical School Institutional Animal

Care and Use Committee, and performed in accordance with experimental protocols 2009N000033 and

2014N000275 approved by the Massachusetts General Hospital Institutional Animal Care and Use

Committee.

For the analysis of transcriptional changes in ovaries of cycling mice, animals were housed in standard

conditions (12/12 hours light/dark non-inverting cycle with food and water *ad libitum*) in groups of 5 with

added bedding from a cage that previously housed an adult male mouse to encourage cycling. Estrous

stage was determined by observation of the vaginal opening and by vaginal swabs done at the same time

daily, as previously described (12). Each mouse was monitored for a minimum of 2 weeks to ensure its

cyclicity. Four mice were sacrificed in each of the 4 phases of the estrous cycle and labeled as being from

experimental batch "cycling". An additional 8 mice were included in the analysis and labeled as being from

experimental batch "lactating". Four of these mice were lactating at day 10 postpartum and four were 10

days postpartum with pups removed at delivery. Four additional mice were not monitored for cycling and

included to increase sample diversity.

Additional mice were monitored throughout the estrous cycle, to collect ovaries at each stage (groups of

N=5 for proestrus, estrus, metestrus, diestrus) for gene validation. Paired ovaries were collected from each

staged mouse: one was used to extract mRNA for qPCR, while the other was fixed in 4% paraformaldehyde

for RNAish (RNAscope™) or immunohistochemistry to validate gene expression.

Superovulation

To stimulate superovulation, mature female mice (6-9 weeks C57BL/6) were injected IP with 5 IU of

pregnant mare serum gonadotropin (PMSG) (Calbiochem, San Diego, CA), followed 48 hours later by 5 IU

of human chorionic gonadotropin (hCG; Millipore Sigma, St. Louis, MO). The mice were euthanized 8 hours

after hCG treatment and ovaries harvested.

Staging of estrous cycle by vaginal cytology

As previously described (12,13), staging of mice was performed using a wet cotton swab, introduced into

the vaginal orifice then smeared onto a glass slide which was air-dried, stained with Giemsa, and scored

for cytology by two independent observers. Briefly, proestrus was determined if the smear showed a

preponderance of nucleated epithelial cells as well as leukocytes. Estrous was marked by an abundance

of cornified epithelial cells, while metestrous smears contained a mixture of cornified epithelial cells and

leukocytes. Finally, diestrus was characterized by abundant leukocytes with low numbers of cornified

epithelium or nucleated epithelial cells.

Generation of single cell suspension

Single cell suspension from mouse ovaries were obtained as previously described with uterine enzymatic

dissociation (14). Briefly, ovaries were incubated for 30min at 34C in dissociation medium (82 mM Na_2SO_4,

30 mM K_2SO_4, 10 mM Glucose, 10 mM HEPES, and 5 mM $MgCL_2$, pH 7.4) containing 15 mg of Protease

XXIII (Worthington), 100 U Papain, with 5 mm L-Cysteine, 2.5 mM EDTA (Worthington) and 1333 U of

DNase 1 (Worthington). The reaction was then stopped in cold medium, and samples were mechanically

dissociated, filtrated, and spun down three times before being resuspended to a concentration of 150,000

cells/mL in 20% Optiprep (Sigma) for inDrop sorting.

Single-cell RNA sequencing (inDrops)

Fluidic sorting was performed using the inDrop platform at the Single Cell Core facility at Harvard Medical

School as previously described (15,16). We generated libraries of approximately 1500 cells per animal

which were sequenced using the NextSeq500 (Illumina) platform. Transcripts were processed according to

a previously published pipeline (15) used to build a custom transcriptome from the Ensemble GRCm38

genome and GRCm38.84 annotation using Bowtie 1.1.1. Unique molecular identifiers (UMIs) were used to

reference sequence reads back to individual captured molecules, referred to as UMIFM counts. All steps

of the pipeline were run using default parameters unless explicitly specified.

Single-cell RNAseq data analysis

Data processing

The initial Seurat object was created using thresholds to identify putative cells (unique cell barcodes) with

the following parameters: 1000-20000 unique molecular identifiers, 500-5000 genes, and less than 15%

mitochondrial genes. The final merged dataset contained ~70000 cells which were clustered based on

expression of marker genes. These were further processed in several ways to exclude low-quality data

and potential doublets. Visualization of single-cell data was performed using a nonlinear dimensionality-

reduction technique, uniform manifold approximation and projection (UMAP). Markers for each level of

cluster were identified using MAST in Seurat (R version 4.1.3 - Seurat version 4.1.0). Following identification

of the main clusters (granulosa, mesenchyme, endothelium, immune, epithelium, and oocyte) we

reanalyzed each cluster population to perform subclustering. Briefly, the granulosa, mesenchyme, and

epithelium clusters were extracted from the integrated dataset by the Subset function. The isolated cluster

was then divided into several subclusters following normalization, scale, principal component analysis, and

dimensionality reductions as previously described (17).

Volcano Plots

Highly differentially expressed genes between different estrous cycles were identified using the function

FindMarkers in Seurat. Volcano plots were generated using ggplot2 package in R.

Pathway Enrichment Analysis

Differentially expressed genes with at least two-fold changes between contiguous estrous stages were used

as input for Gene Ontology Enrichment Analysis by clusterProfiler. Enrichplot package was used for

visualization. Biological process sub-ontology was chosen for this analysis.

Principal component analysis (PCA)

PCA was used to identify common patterns of gene expression across stages of the cycle. For each

Level0 cluster object, cycling cells were extracted and genes that were expressed in more than 5% of

cells were identified. The expression of these genes in the cycling cells were scaled (set to mean zero,

standard deviation 1) and averaged across each of the 4 cycle stages. PCA was run (prcomp) on the

average scaled expression data.

Data availability

The scRNAseq count matrix dataset was deposited in OSF (18), and at the Broad Institute Single Cell

Portal under study number SCP1914 (19).

In situ hybridization and immunohistochemistry

In situ hybridizations were performed using ACDBio kits as per manufacturer's protocol, and as previously

described (14) Briefly, RNAish was developed using the RNAscope® 2.5 HD Reagent Kit (RED and Duplex,

ACD Bio). Following deparaffinization in xylene, dehydration, peroxidase blocking, and heat-induced

epitope retrieval by the target retrieval and protease plus reagents (ACD bio), tissue sections were

hybridized with probes for the target genes (see Key resources material table for accession number and

catalog number of each gene) in the HybEZ hybridization oven (ACD Bio) for 2 hours at 40°C. The slides

were then processed for standard signal amplification steps, and chromogen development. Slides were

counterstained in 50% hematoxylin (Dako), air dried, and cover-slipped with EcoMount™. In addition to

cycling and noncycling mice, superovulated mice were used to validate markers from follicles associated

with LH surge response in ovulatory follicles at the estrous stage for more precise timing of collection.

For colocalization of RNAish staining with immunohistochemistry, we first processed the tissue section for

RNAscope™ as described above, including deparaffinization, antigen retrieval, hybridization, and

chromogen development. Sections were then blocked in 3% bovine serum albumin (BSA) in Tris-buffered

solution (TBS) for 1 hr. Following 3 washes with TBS, the sections were incubated with the primary antibody

(smooth muscle actin primary antibody (SMA); 1:300, Abcam) overnight at 4°C and developed with Dako

EnVision + System horseradish peroxidase (HRP). Labeled Polymer Anti-Rabbit was used as the

secondary antibody, and the HRP signal was detected using the Dako detection system. Slides were then

counterstained in hematoxylin and mounted as described above.

RT-qPCR

Mice were monitored through the estrous cycle and sacrificed at specific stage/time-points as described

above. Ovaries were dissected and total RNA was extracted using the Qiagen RNA extraction kit (Qiagen).

A cDNA library was synthesized from 500 ng total RNA using SuperScript III First-Strand Synthesis System

for RT-PCR using manufacturer's instructions with random hexamers (Invitrogen). The primers used for

this study are described in **Supplementary File 1**. Expression levels were normalized to the Gapdh

transcript using cycle threshold (Ct) values logarithmically transformed by the $2-\Delta Ct$ function.

ELISA

Blood was collected from mice by facial vein puncture, incubated at RT until spontaneously clotted,

centrifuged at 8000rpm for 5min to collect the serum layer, and diluted 1/10 in each ELISA kit according to

the manufacturing protocol; Mouse CNP/NPPC ELISA kit; Mouse serine protease inactive 35 (PRSS35)

ELISA kit; Mouse TINAGL1 / Lipocalin 7 ELISA kit; and Human/Mouse/Rat Activin A Quantikine ELISA Kit

(see Key resources material table).

Legends

Figure 1 - Single cell RNA sequencing of cycling mouse ovaries. A. Schematic of the single cell

sequencing pipeline. B. UMAP plot featuring the different clusters of the ovary and their composition by

stage of the estrus cycle, lactating stage, or unmonitored. C. Heatmap of the top 10 markers of each cluster.

Figure 2 - Identification of the different cell types of the mesenchyme cluster. A. UMAP plot featuring

the different cell subclusters belonging to the mesenchyme cluster. B. Heatmap of the top 5 markers of

each subcluster. C. Validation of the identitity of mesenchyme subcluster by UMAP-plots and RNA in situ

hybridization.

Figure 3 - Identification of the different cell types in the granulosa cluster. A. UMAP plot featuring the

different cell subclusters belonging to the granulosa cluster. B. Heatmap of the top 5 markers of each

cluster. C. Validation of the identitity of granulosa subclusters by UMAP-plots RNA in situ hybridization

Figure 4 – Identification of epithelial subclusters. A. UMAP plot of the epithelium cluster showing two

subclusters: epithelium and proliferating epithelium. B. Heatmap of proliferation markers expressed in the

proliferating epithelium cluster. C. UMAP plot of the cellular composition of the epithelium subclusters by

reproductive state. D. Expression of proliferation markers depending on the phase of the estrous cycle.

Figure 5 – Gene expression in granulosa cells by estrous stage. A. UMAP plot featuring estrous cycle

stages in the granulosa cell cluster. B. UMAP plots featuring each of the estrous cycle phases individually.

C. Volcano plot of genes differentially expressed between proestrus and estrous stages. D. DotPlot of

differentially expressed markers between proestrus and estrus. E. QPCR validation of differentially

expressed genes involved in extracellular matrix remodeling, and steroidogenesis markers (n=5 per group,

mean ± SEM, *$P<0.05$, **$P<0.01$, ***$P<0.005$, ****$P<0.001$).

Figure 6 – Identification and validation of new secreted estrous staging markers. A. Expression of

granulosa cell transcripts varying by estrous cycle stage. B. Validation of significantly up- and

downregulated transcripts of secreted estrous staging markers by qPCR (n=5 per group, mean ± SEM,

*P<0.05, **P<0.01, ***P<0.005). C. Localization of estrous staging markers by in situ hybridization

(RNAscope) in ovarian sections. D. Quantification of circulating estrous staging markers proteins in the

blood by ELISA (n=5 per group, mean ± SEM, *P<0.05, ***P<0.005). E. Summary of the timing of

expression of estrous staging markers in the blood.

Figure 1 – supplement 1 – Ovarian morphology by reproductive state. A. Illustration of the different

cell types of the ovary. B. Representative micrographs of sections of ovaries at each stage of the estrous

cycle, post-partum lactating and post-partum non-lactating reproductive states.

Figure 2 – supplement 1 – Characterization of mesenchymal cell clusters. A. Co-expression of Acta2

and Mfap5 or Acta2 and Hhip in UMAP plots. Co-localization of B. Acta2 and Mfap5 or C. Acta2 and Hhip

in ovarian tissue sections stained by IHC and RNA in situ hybridization. D. Expression of mesenchymal

markers by cluster in DotPlot. E. Expression of steroidogenesic and fibroblast markers in DotPlot that differ

between the two interstitial stromal cell clusters. F. RNA in situ hybridization of different markers

representative of the steroidogenic stroma (Cyp11a1, Ptch1) and the fibroblast-like stroma cell clusters

(Kcnk2, Cxcl14, and Col1a1).

Figure 3- supplement 1 – Characterization of apoptotic and corpus luteum clusters. A. UMAP plots

of different apoptotic markers that colocalize with *Ghr* in the apoptotic cluster. B. Colocalization of Top2a

and Cdkn1a in UMAP plots of corpus luteum clusters. C. RNA in situ hybridization of Top2a and Cdkn1a in

ovarian sections. Representative active (Top2a+) and regressing (Cdkn1a+) corpora lutea. D. Examples of

luteinizing mural markers in DotPlot that differ from antral mural. E. UMAP plot of cellular distribution of

post-partum lactating (PPL) cells across the active corpus luteum cluster and post-partum non-lactating

(PPNL) cells across the regressing corpus luteum cluster. F. Heatmap of representative markers of active

and regressing corpora lutea.

Figure 5 – supplement 1 – Characterization of the granulosa cell transcriptome across the estrous

cycle. A. Volcano plot of genes differentially expressed between the estrous/metestrous,

metestrous/diestrous and diestrous/proestrous stages. B. Pathway analysis of differentially expressed

genes (DEG) between the different stages of the estrous cycle. The x-axis shows the gene ratio (the

percentage of total DEGs in the given GO terms). "Count", reflected by the dot size, represents the number

of genes enriched in a GO term and dot color represents the adjusted p values. . C. QPCR validation of

ovarian surface epithelium-expressed markers (n=5 per group, mean ± SEM, *P<0.05, **P<0.01,

****P<0.001).

Supplementary File 1- List of primers used for qPCR experiments

Supplementary File 2 – Top 10 markers expressed in each ovary cluster

Supplementary File 3 - Top 10 markers from each mesenchyme subclusters

Supplementary File 4 - Top 10 markers from each granulosa subclusters

Supplementary File 5 - Secreted markers expressed in granulosa cells varying with the estrous

cycle

Source data 1 – qPCR experiments individual p values

Source data 2 – ELISA experiments individual p values

Acknowledgements

We thank LiHua Zhang, Bugra Uluyurt, Phoebe A. May, Caroline Coletti, and Sarah Mustafa Eisa for

technical help. This study was supported by the National Institute for Child Health and Human Development

to D.P. (1R01HD102014-01), the Huiying Fellowship (H.D.S.), Sudna Gar Fellowship (D.P.),

Massachusetts General Hospital Executive Committee on Research (D.P. and P.K.D.), and royalties

(P.K.D.) from the use of the MIS ELISA in infertility clinics.

Bibliography

556 1. **Dunlop CE, Anderson RA.** The regulation and assessment of follicular growth. *Scand J Clin Lab Invest*

557   *Suppl* 2014;244:13–17; discussion 17.

558 2. **Zhao Z-H, Ma J-Y, Meng T-G, Wang Z-B, Yue W, Zhou Q, Li S, Feng X, Hou Y, Schatten H, Ou X-H,**

559   **Sun Q-Y.** Single-cell RNA sequencing reveals the landscape of early female germ cell development.

560   *FASEB J* 2020;34(9):12634–12645.

561 3. **Stévant I, Kühne F, Greenfield A, Chaboissier M-C, Dermitzakis ET, Nef S.** Dissecting Cell Lineage

562   Specification and Sex Fate Determination in Gonadal Somatic Cells Using Single-Cell Transcriptomics.

563   *Cell Rep* 2019;26(12):3272-3283.e3.

564 4. **Wagner M, Yoshihara M, Douagi I, Damdimopoulos A, Panula S, Petropoulos S, Lu H, Pettersson K,**

565   **Palm K, Katayama S, Hovatta O, Kere J, Lanner F, Damdimopoulou P.** Single-cell analysis of human

566   ovarian cortex identifies distinct cell populations but no oogonial stem cells. *Nat Commun*

567   2020;11(1):1147.

568 5. **Niu W, Spradling AC.** Two distinct pathways of pregranulosa cell differentiation support follicle

569   formation in the mouse ovary. *Proc Natl Acad Sci U S A* 2020;117(33):20015–20026.

570 6. **Jevitt A, Chatterjee D, Xie G, Wang X-F, Otwell T, Huang Y-C, Deng W-M.** A single-cell atlas of adult

571   Drosophila ovary identifies transcriptional programs and somatic cell lineage regulating oogenesis.

572   *PLoS Biol* 2020;18(4):e3000538.

573 7. **Man L, Lustgarten-Guahmich N, Kallinos E, Redhead-Laconte Z, Liu S, Schattman B, Redmond D,**

574   **Hancock K, Zaninovic N, Schattman G, Rosenwaks Z, James D.** Comparison of Human Antral

575   Follicles of Xenograft versus Ovarian Origin Reveals Disparate Molecular Signatures. *Cell Rep*

576   2020;32(6):108027.

577 8. **Meinsohn M-C, Saatcioglu HD, Wei L, Li Y, Horn H, Chauvin M, Kano M, Nguyen NMP, Nagykery N,**

578   **Kashiwagi A, Samore WR, Wang D, Oliva E, Gao G, Morris ME, Donahoe PK, Pépin D.** Single-cell

sequencing reveals suppressive transcriptional programs regulated by MIS/AMH in neonatal ovaries.

Proc Natl Acad Sci U S A 2021;118(20). doi:10.1073/pnas.2100920118.

9.　**Fan X, Bialecka M, Moustakas I, Lam E, Torrens-Juaneda V, Borggreven NV, Trouw L, Louwe LA,**

Pilgram GSK, Mei H, van der Westerlaken L, Chuva de Sousa Lopes SM. Single-cell reconstruction of

follicular remodeling in the human adult ovary. *Nat Commun* 2019;10(1):3164.

10.　**Wang S, Zheng Y, Li J, Yu Y, Zhang W, Song M, Liu Z, Min Z, Hu H, Jing Y, He X, Sun L, Ma L, Esteban**

CR, Chan P, Qiao J, Zhou Q, Izpisua Belmonte JC, Qu J, Tang F, Liu G-H. Single-Cell Transcriptomic

Atlas of Primate Ovarian Aging. *Cell* 2020;180(3):585-600.e19.

11.　**Ajayi AF, Akhigbe RE.** Staging of the estrous cycle and induction of estrus in experimental rodents:

an update. *Fertil Res Pract* 2020;6:5.

12.　**Kano M, Sosulski AE, Zhang L, Saatcioglu HD, Wang D, Nagykery N, Sabatini ME, Gao G, Donahoe**

PK, Pépin D. AMH/MIS as a contraceptive that protects the ovarian reserve during chemotherapy.

Proc Natl Acad Sci U S A 2017;114(9):E1688–E1697.

13.　**Byers SL, Wiles MV, Dunn SL, Taft RA.** Mouse estrous cycle identification tool and images. *PLoS One*

2012;7(4):e35538.

14.　**Saatcioglu HD, Kano M, Horn H, Zhang L, Samore W, Nagykery N, Meinsohn M-C, Hyun M, Suliman**

R, Poulo J, Hsu J, Sacha C, Wang D, Gao G, Lage K, Oliva E, Morris Sabatini ME, Donahoe PK, Pépin

D. Single-cell sequencing of neonatal uterus reveals an Misr2+ endometrial progenitor indispensable

for fertility. *Elife* 2019;8. doi:10.7554/eLife.46349.

15.　**Klein AM, Mazutis L, Akartuna I, Tallapragada N, Veres A, Li V, Peshkin L, Weitz DA, Kirschner MW.**

Droplet barcoding for single-cell transcriptomics applied to embryonic stem cells. *Cell*

2015;161(5):1187–1201.

16.　**Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas AR, Kamitaki N,**

Martersteck EM, Trombetta JJ, Weitz DA, Sanes JR, Shalek AK, Regev A, McCarroll SA. Highly

Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell*

2015;161(5):1202–1214.

17. **Niu W, Spradling AC.** Two distinct pathways of pregranulosa cell differentiation support follicle

formation in the mouse ovary. *Proc. Natl. Acad. Sci. U.S.A.* 2020. doi:10.1073/pnas.2005570117.

18. **Pepin.** "Single Cell Sequencing of the Mouse Ovary in Diverse Reproductive States." OSF. 2022.

19. **Single Cell Portal.** Available at: https://singlecell.broadinstitute.org/single_cell/study/SCP1914/a-

single-cell-atlas-of-the-cycling-murine-ovary. Accessed July 7, 2022.

20. **Richards JS, Ren YA, Candelaria N, Adams JE, Rajkovic A.** Ovarian Follicular Theca Cell Recruitment,

Differentiation, and Impact on Fertility: 2017 Update. *Endocr Rev* 2018;39(1):1–20.

21. **Young JM, McNeilly AS.** Theca: the forgotten cell of the ovarian follicle. *Reproduction*

2010;140(4):489–504.

22. **Spicer LJ, Sudo S, Aad PY, Wang LS, Chun S-Y, Ben-Shlomo I, Klein C, Hsueh AJW.** The hedgehog-

patched signaling pathway and function in the mammalian ovary: a novel role for hedgehog

proteins in stimulating proliferation and steroidogenesis of theca cells. *Reproduction*

2009;138(2):329–339.

23. **Muhl L, Genové G, Leptidis S, Liu J, He L, Mocci G, Sun Y, Gustafsson S, Buyandelger B, Chivukula**

IV, Segerstolpe Å, Raschperger E, Hansson EM, Björkegren JLM, Peng X-R, Vanlandewijck M,

Lendahl U, Betsholtz C. Single-cell analysis uncovers fibroblast heterogeneity and criteria for

fibroblast and mural cell identification and discrimination. *Nat Commun* 2020;11(1):3953.

24. **Lu J, Chatterjee M, Schmid H, Beck S, Gawaz M.** CXCL14 as an emerging immune and inflammatory

modulator. *J Inflamm (Lond)* 2016;13:1.

25. **He H, Luthringer DJ, Hui P, Lau SK, Weiss LM, Chu PG.** Expression of CD56 and WT1 in ovarian

stroma and ovarian stromal tumors. *Am J Surg Pathol* 2008;32(6):884–890.

26. **Hughes S, Chan-Ling T.** Characterization of smooth muscle cell and pericyte differentiation in the rat

retina in vivo. *Invest Ophthalmol Vis Sci* 2004;45(8):2795–2806.

27. **Kiriakidou M, McAllister JM, Sugawara T, Strauss JF 3rd.** Expression of steroidogenic acute

regulatory protein (StAR) in the human ovary. *J Clin Endocrinol Metab* 1996;81(11):4122–4128.

28. **Yang R, Chen Y, Chen D.** Biological functions and role of CCN1/Cyr61 in embryogenesis and

tumorigenesis in the female reproductive system (Review). *Mol Med Rep* 2018;17(1):3–10.

29. **Hatzirodos N, Hummitzsch K, Irving-Rodgers HF, Rodgers RJ.** Transcriptome comparisons identify

new cell markers for theca interna and granulosa cells from small and large antral ovarian follicles.

PLoS One 2015;10(3):e0119800.

30. **Porcile C, Bajetto A, Barbieri F, Barbero S, Bonavia R, Biglieri M, Pirani P, Florio T, Schettini G.**

Stromal cell-derived factor-1alpha (SDF-1alpha/CXCL12) stimulates ovarian cancer cell growth

through the EGF receptor transactivation. *Exp Cell Res* 2005;308(2):241–253.

31. **Gallardo TD, John GB, Shirley L, Contreras CM, Akbay EA, Haynie JM, Ward SE, Shidler MJ,**

Castrillon DH. Genomewide discovery and classification of candidate ovarian fertility genes in the

mouse. *Genetics* 2007;177(1):179–194.

32. **Ho ML, Lee JN.** Ovarian and circulating levels of oxytocin and arginine vasopressin during the

estrous cycle in the rat. *Acta Endocrinol (Copenh)* 1992;126(6):530–534.

33. **Nakagawa S, Shimada M, Yanaka K, Mito M, Arai T, Takahashi E, Fujita Y, Fujimori T, Standaert L,**

Marine J-C, Hirose T. The lncRNA Neat1 is required for corpus luteum formation and the

establishment of pregnancy in a subpopulation of mice. *Development* 2014;141(23):4618–4627.

34. **Donadeu FX, Fahiminiya S, Esteves CL, Nadaf J, Miedzinska K, McNeilly AS, Waddington D, Gérard**

N. Transcriptome profiling of granulosa and theca cells during dominant follicle development in the

horse. *Biol Reprod* 2014;91(5):111.

35. **Ock S-A, Knott JG, Choi I.** Involvement of CDKN1A (p21) in cellular senescence in response to heat

and irradiation stress during preimplantation development. *Cell Stress Chaperones* 2020;25(3):503–

508.

36. **Talbott H, Hou X, Qiu F, Zhang P, Guda C, Yu F, Cushman RA, Wood JR, Wang C, Cupp AS, Davis JS.**

Early transcriptome responses of the bovine midcycle corpus luteum to prostaglandin F2α includes

cytokine signaling. *Mol Cell Endocrinol* 2017;452:93–109.

37. **Terenina E, Fabre S, Bonnet A, Monniaux D, Robert-Granié C, SanCristobal M, Sarry J, Vignoles F,**

Gondret F, Monget P, Tosser-Klopp G. Differentially expressed genes and gene networks involved in

pig ovarian follicular atresia. *Physiol Genomics* 2017;49(2):67–80.

38. **Liu C, Wang J, Zhao L, He H, Zhao P, Peng Z, Liu F, Chen J, Wu W, Wang G, Dong F.** Knockdown of

Thymidine Kinase 1 Suppresses Cell Proliferation, Invasion, Migration, and Epithelial-Mesenchymal

Transition in Thyroid Carcinoma Cells. *Front Oncol* 2019;9:1475.

39. **Yang X-M, Cao X-Y, He P, Li J, Feng M-X, Zhang Y-L, Zhang X-L, Wang Y-H, Yang Q, Zhu L, Nie H-Z,**

Jiang S-H, Tian G-A, Zhang X-X, Liu Q, Ji J, Zhu X, Xia Q, Zhang Z-G. Overexpression of Rac GTPase

Activating Protein 1 Contributes to Proliferation of Cancer Cells by Reducing Hippo Signaling to

Promote Cytokinesis. *Gastroenterology* 2018;155(4):1233-1249.e22.

40. **Gao H-S, Lin S-Y, Han X, Xu H-Z, Gao Y-L, Qin Z-Y.** Casein kinase 1 (CK1) promotes the proliferation

and metastasis of glioma cells via the phosphatidylinositol 3 kinase-matrix metalloproteinase 2

(AKT-MMP2) pathway. *Ann Transl Med* 2021;9(8):659.

41. **Liang Z, Li X, Chen J, Cai H, Zhang L, Li C, Tong J, Hu W.** PRC1 promotes cell proliferation and cell

cycle progression by regulating p21/p27-pRB family molecules and FAK-paxillin pathway in non-

small cell lung cancer. *Transl Cancer Res* 2019;8(5):2059–2072.

42. **Xiong Y, Lu J, Fang Q, Lu Y, Xie C, Wu H, Yin Z.** UBE2C functions as a potential oncogene by

enhancing cell proliferation, migration, invasion, and drug resistance in hepatocellular carcinoma

cells. *Biosci Rep* 2019;39(4):BSR20182384.

43. **Xu L, Yu W, Xiao H, Lin K.** BIRC5 is a prognostic biomarker associated with tumor immune cell

infiltration. *Sci Rep* 2021;11(1):390.

44. **Mara JN, Zhou LT, Larmore M, Johnson B, Ayiku R, Amargant F, Pritchard MT, Duncan FE.**

Ovulation and ovarian wound healing are impaired with advanced reproductive age. *Aging (Albany*

NY) 2020;12(10):9686–9713.

45. **Florin L, Knebel J, Zigrino P, Vonderstrass B, Mauch C, Schorpp-Kistner M, Szabowski A, Angel P.**

Delayed wound healing and epidermal hyperproliferation in mice lacking JunB in the skin. *J Invest*

Dermatol 2006;126(4):902–911.

46. **Wu M, Melichian DS, de la Garza M, Gruner K, Bhattacharyya S, Barr L, Nair A, Shahrara S, Sporn**

PHS, Mustoe TA, Tourtellotte WG, Varga J. Essential roles for early growth response transcription

factor Egr-1 in tissue fibrosis and wound healing. *Am J Pathol* 2009;175(3):1041–1055.

47. **Martin P, Nobes CD.** An early molecular component of the wound healing response in rat embryos--

induction of c-fos protein in cells at the epidermal wound margin. *Mech Dev* 1992;38(3):209–215.

48. **Yue C, Guo Z, Luo Y, Yuan J, Wan X, Mo Z.** c-Jun Overexpression Accelerates Wound Healing in

Diabetic Rats by Human Umbilical Cord-Derived Mesenchymal Stem Cells. *Stem Cells Int*

2020;2020:7430968.

49. **Sherman BT, Hao M, Qiu J, Jiao X, Baseler MW, Lane HC, Imamichi T, Chang W.** DAVID: a web

server for functional enrichment analysis and functional annotation of gene lists (2021 update).

Nucleic Acids Res 2022;50(W1):W216-221.

50. **Huang DW, Sherman BT, Lempicki RA.** Systematic and integrative analysis of large gene lists using

DAVID bioinformatics resources. *Nat Protoc* 2009;4(1):44–57.

51. **The Genotype-Tissue Expression (GTEx) project.** *Nat Genet* 2013;45(6):580–585.

52. **Toms D, Pan B, Li J.** Endocrine Regulation in the Ovary by MicroRNA during the Estrous Cycle. *Front*

Endocrinol (Lausanne) 2017;8:378.

53. **Kubota K, Cui W, Dhakal P, Wolfe MW, Rumi MAK, Vivian JL, Roby KF, Soares MJ.** Rethinking

progesterone regulation of female reproductive cyclicity. *Proc Natl Acad Sci U S A*

2016;113(15):4212–4217.

54. **Hwang B, Lee JH, Bang D.** Single-cell RNA sequencing technologies and bioinformatics pipelines. *Exp*

Mol Med 2018;50(8):1–14.

55. **Deb S, Campbell BK, Clewes JS, Pincott-Allen C, Raine-Fenning NJ.** Intracycle variation in number of

antral follicles stratified by size and in endocrine markers of ovarian reserve in women with normal

ovulatory menstrual cycles. *Ultrasound Obstet Gynecol* 2013;41(2):216–222.

56. **Richards JS.** Maturation of ovarian follicles: actions and interactions of pituitary and ovarian

hormones on follicular cell differentiation. *Physiol Rev* 1980;60(1):51–89.

57. **Lazzaro MA, Pépin D, Pescador N, Murphy BD, Vanderhyden BC, Picketts DJ.** The imitation switch

protein SNF2L regulates steroidogenic acute regulatory protein expression during terminal

differentiation of ovarian granulosa cells. *Mol Endocrinol* 2006;20(10):2406–2417.

58. **Irving-Rodgers HF, Harland ML, Sullivan TR, Rodgers RJ.** Studies of granulosa cell maturation in

dominant and subordinate bovine follicles: novel extracellular matrix focimatrix is co-ordinately

regulated with cholesterol side-chain cleavage CYP11A1. *Reproduction* 2009;137(5):825–834.

59. **Wang S, Liu W, Pang X, Dai S, Liu G.** The Mechanism of Melatonin and Its Receptor MT2 Involved in

the Development of Bovine Granulosa Cells. *Int J Mol Sci* 2018;19(7). doi:10.3390/ijms19072028.

60. **Wahlberg P, Nylander A, Ahlskog N, Liu K, Ny T.** Expression and localization of the serine proteases

high-temperature requirement factor A1, serine protease 23, and serine protease 35 in the mouse

ovary. *Endocrinology* 2008;149(10):5070–5077.

61. **Lussier JG, Diouf MN, Lévesque V, Sirois J, Ndiaye K.** Gene expression profiling of upregulated

mRNAs in granulosa cells of bovine ovulatory follicles following stimulation with hCG. *Reprod Biol*

Endocrinol 2017;15(1):88.

62. **Sayasith K, Lussier J, Sirois J.** Molecular characterization and transcriptional regulation of a

disintegrin and metalloproteinase with thrombospondin motif 1 (ADAMTS1) in bovine preovulatory

follicles. *Endocrinology* 2013;154(8):2857–2869.

63. **Diaz FJ, Wigglesworth K, Eppig JJ.** Oocytes determine cumulus cell lineage in mouse ovarian

follicles. *Journal of Cell Science* 2007;120(8):1330–1340.

64. **Duncan WC.** The inadequate corpus luteum. *Reprod Fertil* 2021;2(1):C1–C7.

65. **Noguchi M, Hirata M, Kawaguchi H, Tanimoto A.** Corpus luteum Regression Induced by

Prostaglandin F(2α) in Microminipigs During the Normal Estrous Cycle. *In Vivo* 2017;31(6):1097–

1101.

66. **Zhu R, Zou S-T, Wan J-M, Li W, Li X-L, Zhu W.** BTG1 inhibits breast cancer cell growth through

induction of cell cycle arrest and apoptosis. *Oncol Rep* 2013;30(5):2137–2144.

67. **RYAN KJ, PETRO Z.** Steroid Biosynthesis by Human Ovarian Granulosa and Thecal Cells. *The Journal*

of Clinical Endocrinology & Metabolism 1966;26(1):46–52.

68. **Hummitzsch K, Hatzirodos N, Macpherson AM, Schwartz J, Rodgers RJ, Irving-Rodgers HF.**

Transcriptome analyses of ovarian stroma: tunica albuginea, interstitium and theca interna.

Reproduction 2019;157(6):545–565.

69. **Grosdemouge I, Bachelot A, Lucas A, Baran N, Kelly PA, Binart N.** Effects of deletion of the

prolactin receptor on ovarian gene expression. *Reprod Biol Endocrinol* 2003;1:12.

70. **Sheng X, Zhou J, Kang N, Liu W, Yu L, Zhang Z, Zhang Y, Yue Q, Yang Q, Zhang X, Li C, Yan G, Sun H.**

Temporal and spatial dynamics mapping reveals follicle development regulated by different stromal

cell populations. *bioRxiv* 2022. doi:10.1101/2022.03.04.480328.

71. **Kinnear HM, Tomaszewski CE, Chang FL, Moravek MB, Xu M, Padmanabhan V, Shikanov A.** The

ovarian stroma as a new frontier. *Reproduction* 2020;160(3):R25–R39.

72. **Duffy DM, Ko C, Jo M, Brannstrom M, Curry TE.** Ovulation: Parallels With Inflammatory Processes.

Endocr Rev 2019;40(2):369–416.

73. **Hartanti MD, Hummitzsch K, Irving-Rodgers HF, Bonner WM, Copping KJ, Anderson RA, McMillen**

IC, Perry VEA, Rodgers RJ. Morphometric and gene expression analyses of stromal expansion during

development of the bovine fetal ovary. *Reprod Fertil Dev* 2019;31(3):482–495.

74. **Xu F, Stouffer RL, Müller J, Hennebold JD, Wright JW, Bahar A, Leder G, Peters M, Thorne M, Sims**

M, Wintermantel T, Lindenthal B. Dynamics of the transcriptome in the primate ovulatory follicle.

Mol Hum Reprod 2011;17(3):152–165.

75. **Kenngott RA-M, Sauer U, Vermehren M, Sinowatz F.** Expression of Intermediate Filaments and

Germ Cell Markers in the Developing Bovine Ovary: An Immunohistochemical and Laser-Assisted

Microdissection Study. *Cells Tissues Organs* 2014;200(2):153–170.

76. **Lin Y-T, Chen J-S, Wu M-H, Hsieh I-S, Liang C-H, Hsu C-L, Hong T-M, Chen Y-L.** Galectin-1 accelerates

wound healing by regulating the neuropilin-1/Smad3/NOX4 pathway and ROS production in

myofibroblasts. *J Invest Dermatol* 2015;135(1):258–268.

77. **Okada Y, Saika S, Hashizume N, Kobata S, Yamanaka O, Ohnishi Y, Senba E.** Expression of fos family

and jun family proto-oncogenes during corneal epithelial wound healing. *Curr Eye Res*

1996;15(8):824–832.

78. **Wijayarathna R, de Kretser DM.** Activins in reproductive biology and beyond. *Hum Reprod Update*

2016;22(3):342–357.

79. **Li S-H, Lin M-H, Hwu Y-M, Lu C-H, Yeh L-Y, Chen Y-J, Lee RK-K.** Correlation of cumulus gene

expression of GJA1, PRSS35, PTX3, and SERPINE2 with oocyte maturation, fertilization, and embryo

development. *Reprod Biol Endocrinol* 2015;13:93.

80. **Zhang M, Su Y-Q, Sugiura K, Xia G, Eppig JJ.** Granulosa cell ligand NPPC and its receptor NPR2

maintain meiotic arrest in mouse oocytes. *Science* 2010;330(6002):366–369.

81. **Xi G, An L, Wang W, Hao J, Yang Q, Ma L, Lu J, Wang Y, Wang W, Zhao W, Liu J, Yang M, Wang X,**

Zhang Z, Zhang C, Chu M, Yue Y, Yao F, Zhang M, Tian J. The mRNA-destabilizing protein

Tristetraprolin targets "meiosis arrester" Nppc mRNA in mammalian preovulatory follicles. *Proc Natl*

Acad Sci U S A 2021;118(22). doi:10.1073/pnas.2018345118.

82. **Akaiwa M, Fukui E, Matsumoto H.** Tubulointerstitial nephritis antigen-like 1 deficiency alleviates

age-dependent depressed ovulation associated with ovarian collagen deposition in mice. *Reprod*

Med Biol 2020;19(1):50–57.

83. **Kim Y-S, Hwan JD, Bae S, Bae D-H, Shick WA.** Identification of differentially expressed genes using

an annealing control primer system in stage III serous ovarian carcinoma. *BMC Cancer* 2010;10:576.

84. **Chang H-M, Leung PCK.** Physiological roles of activins in the human ovary. *Journal of Bio-X Research*

2018;1(3). Available at:

https://journals.lww.com/jbioxresearch/Fulltext/2018/12000/Physiological_roles_of_activins_in_th

e_human_ovary.2.aspx.

85. **O'Connell AR, McNatty KP, Hurst PR, Spencer TE, Bazer FW, Reader KL, Johnstone PD, Davis GH,**

Juengel JL. Activin A and follistatin during the oestrous cycle and early pregnancy in ewes. *J*

Endocrinol 2016;228(3):193–203.

86. **Celik O, Celik N, Ugur K, Hatirnaz S, Celik S, Muderris II, Yavuzkir S, Sahin İ, Yardim M, Aydin S.**

Nppc/Npr2/cGMP signaling cascade maintains oocyte developmental capacity. *Cell Mol Biol (Noisy-*

le-grand) 2019;65(4):83–89.

87. **Celik O, Celik N, Gungor S, Haberal ET, Aydin S.** Selective Regulation of Oocyte Meiotic Events

Enhances Progress in Fertility Preservation Methods. *Biochem Insights* 2015;8:11–21.

88. **Meinsohn M-C, Morin F, Bertolin K, Duggavathi R, Schoonjans K, Murphy BD.** The Orphan Nuclear

Receptor Liver Homolog Receptor-1 (Nr5a2) Regulates Ovarian Granulosa Cell Proliferation. *J Endocr*

Soc 2018;2(1):24–41.

89. **Hodgkinson K, Forrest LA, Vuong N, Garson K, Djordjevic B, Vanderhyden BC.** GREB1 is an estrogen

receptor-regulated tumour promoter that is frequently expressed in ovarian cancer. *Oncogene*

2018;37(44):5873–5886.

90. **Terman BI, Dougher-Vermazen M, Carrion ME, Dimitrov D, Armellino DC, Gospodarowicz D,**

Böhlen P. Identification of the KDR tyrosine kinase as a receptor for vascular endothelial cell growth

factor. *Biochem Biophys Res Commun* 1992;187(3):1579–1586.

91. **Parikh A, Lee C, Joseph P, Marchini S, Baccarini A, Kolev V, Romualdi C, Fruscio R, Shah H, Wang F,**

Mullokandov G, Fishman D, D'Incalci M, Rahaman J, Kalir T, Redline RW, Brown BD, Narla G, DiFeo

A. microRNA-181a has a critical role in ovarian cancer progression through the regulation of the

epithelial-mesenchymal transition. *Nat Commun* 2014;5:2977.

92. **Galvagni F, Nardi F, Spiga O, Trezza A, Tarticchio G, Pellicani R, Andreuzzi E, Caldi E, Toti P, Tosi**

GM, Santucci A, Iozzo RV, Mongiat M, Orlandini M. Dissecting the CD93-Multimerin 2 interaction

involved in cell adhesion and migration of the activated endothelium. *Matrix Biol* 2017;64:112–127.

93. **Richards JS, Fan H-Y, Liu Z, Tsoi M, Laguë M-N, Boyer A, Boerboom D.** Either Kras activation or Pten

loss similarly enhance the dominant-stable CTNNB1-induced genetic program to promote granulosa

cell tumor development in the ovary and testis. *Oncogene* 2012;31(12):1504–1520.

94. **Zhou Q, Wan M, Wei Q, Song Q, Xiong L, Huo J, Huang J.** Expression, Regulation, and Functional

Characterization of FST Gene in Porcine Granulosa Cells. *Anim Biotechnol* 2016;27(4):295–302.

95. **Bambino K, Lacko LA, Hajjar KA, Stuhlmann H.** Epidermal growth factor-like domain 7 is a marker of

the endothelial lineage and active angiogenesis. *Genesis* 2014;52(7):657–670.

96. **He W, Gauri M, Li T, Wang R, Lin S-X.** Current knowledge of the multifunctional 17β-hydroxysteroid

dehydrogenase type 1 (HSD17B1). *Gene* 2016;588(1):54–61.

97. **Oksjoki S, Sallinen S, Vuorio E, Anttila L.** Cyclic expression of mRNA transcripts for connective tissue

components in the mouse ovary. *Mol Hum Reprod* 1999;5(9):803–808.

98. **Park S, DiMaio TA, Scheef EA, Sorenson CM, Sheibani N.** PECAM-1 regulates proangiogenic

properties of endothelial cells through modulation of cell-cell and cell-matrix interactions. *Am J*

Physiol Cell Physiol 2010;299(6):C1468-1484.

99. **Wigglesworth K, Lee K-B, Emori C, Sugiura K, Eppig JJ.** Transcriptomic diversification of developing

cumulus and mural granulosa cells in mouse ovarian follicles. *Biol Reprod* 2015;92(1):23.

100.   **Varankar SS, More M, Abraham A, Pansare K, Kumar B, Narayanan NJ, Jolly MK, Mali AM,**

Bapat SA. Functional balance between Tcf21-Slug defines cellular plasticity and migratory

modalities in high grade serous ovarian cancer cell lines. *Carcinogenesis* 2020;41(4):515–526.

101.   **Herr D, Fraser HM, Konrad R, Holzheu I, Kreienberg R, Wulff C.** Human chorionic gonadotropin

controls luteal vascular permeability via vascular endothelial growth factor by down-regulation of a

cascade of adhesion proteins. *Fertil Steril* 2013;99(6):1749–1758.

102.   **Angelos MG, Abrahante JE, Blum RH, Kaufman DS.** Single Cell Resolution of Human

Hematoendothelial Cells Defines Transcriptional Signatures of Hemogenic Endothelium. *Stem Cells*

2018;36(2):206–217.

103.   **Bédard J, Brûlé S, Price CA, Silversides DW, Lussier JG.** Serine protease inhibitor-E2 (SERPINE2)

is differentially expressed in granulosa cells of dominant follicle in cattle. *Mol Reprod Dev*

2003;64(2):152–165.

104.   **Sleer LS, Taylor CC.** Cell-type localization of platelet-derived growth factors and receptors in the

postnatal rat ovary and follicle. *Biol Reprod* 2007;76(3):379–390.

105.   **de Vries C, Escobedo JA, Ueno H, Houck K, Ferrara N, Williams LT.** The fms-like tyrosine kinase,

a receptor for vascular endothelial growth factor. *Science* 1992;255(5047):989–991.

106.   **Wu Y, Lin J, Li X, Han B, Wang L, Liu M, Huang J.** Transcriptome profile of one-month-old lambs'

granulosa cells after superstimulation. *Asian-Australas J Anim Sci* 2017;30(1):20–33.

107.   **Galvagni F, Nardi F, Maida M, Bernardini G, Vannuccini S, Petraglia F, Santucci A, Orlandini M.**

CD93 and dystroglycan cooperation in human endothelial cell adhesion and migration adhesion and

migration. *Oncotarget* 2016;7(9):10090–10103.

108.   **Sterzyńska K, Klejewski A, Wojtowicz K, Świerczewska M, Andrzejewska M, Rusek D,**

Sobkowski M, Kędzia W, Brązert J, Nowicki M, Januchowski R. The Role of Matrix Gla Protein

(MGP) Expression in Paclitaxel and Topotecan Resistant Ovarian Cancer Cell Lines. *Int J Mol Sci*

2018;19(10). doi:10.3390/ijms19102901.

109.   **Kalucka J, de Rooij LPMH, Goveia J, Rohlenova K, Dumas SJ, Meta E, Conchinha NV, Taverna F,**

Teuwen L-A, Veys K, García-Caballero M, Khan S, Geldhof V, Sokol L, Chen R, Treps L, Borri M, de

Zeeuw P, Dubois C, Karakach TK, Falkenberg KD, Parys M, Yin X, Vinckier S, Du Y, Fenton RA,

Schoonjans L, Dewerchin M, Eelen G, Thienpont B, Lin L, Bolund L, Li X, Luo Y, Carmeliet P. Single-

Cell Transcriptome Atlas of Murine Endothelial Cells. *Cell* 2020;180(4):764-779.e20.

110.   **Cochain C, Vafadarnejad E, Arampatzi P, Pelisek J, Winkels H, Ley K, Wolf D, Saliba A-E,**

Zernecke A. Single-Cell RNA-Seq Reveals the Transcriptional Landscape and Heterogeneity of Aortic

Macrophages in Murine Atherosclerosis. *Circ Res* 2018;122(12):1661–1674.

111.   **Carpenter AR, Becknell MB, Ching CB, Cuaresma EJ, Chen X, Hains DS, McHugh KM.** Uroplakin

1b is critical in urinary tract development and urothelial differentiation and homeostasis. *Kidney Int*

2016;89(3):612–624.

112.   **Glowacka WK, Alberts P, Ouchida R, Wang J-Y, Rotin D.** LAPTM5 protein is a positive regulator

of proinflammatory signaling pathways in macrophages. *J Biol Chem* 2012;287(33):27691–27702.

113.   **Paulini F, Melo EO.** The role of oocyte-secreted factors GDF9 and BMP15 in follicular

861   development and oogenesis. *Reprod Domest Anim* 2011;46(2):354–361.

862   114.   **Rudat C, Grieskamp T, Röhr C, Airik R, Wrede C, Hegermann J, Herrmann BG, Schuster-Gossler**

863   **K, Kispert A.** Upk3b is dispensable for development and integrity of urothelium and mesothelium.

864   *PLoS One* 2014;9(11):e112112.

865   115.   **Yang P, Wu Q, Sun L, Fang P, Liu L, Ji Y, Park J-Y, Qin X, Yang X, Wang H.** Adaptive Immune

866   Response Signaling Is Suppressed in Ly6C(high) Monocyte but Upregulated in Monocyte Subsets of

867   ApoE (-/-) Mice - Functional Implication in Atherosclerosis. *Front Immunol* 2021;12:809208.

868   116.   **Tanaka M, Kihara M, Hennebold JD, Eppig JJ, Viveiros MM, Emery BR, Carrell DT, Kirkman NJ,**

869   **Meczekalski B, Zhou J, Bondy CA, Becker M, Schultz RM, Misteli T, De La Fuente R, King GJ, Adashi**

870   **EY.** H1FOO is coupled to the initiation of oocytic growth. *Biol Reprod* 2005;72(1):135–142.

871   117.   **Sewgobind NV, Albers S, Pieters RJ.** Functions and Inhibition of Galectin-7, an Emerging Target

872   in Cellular Pathophysiology. *Biomolecules* 2021;11(11). doi:10.3390/biom11111720.

873   118.   **Liu X, Morency E, Li T, Qin H, Zhang X, Zhang X, Coonrod S.** Role for PADI6 in securing the

874   mRNA-MSY2 complex to the oocyte cytoplasmic lattices. *Cell Cycle* 2017;16(4):360–366.

875   119.   **Auersperg N.** The stem-cell profile of ovarian surface epithelium is reproduced in the oviductal

876   fimbriae, with increased stem-cell marker density in distal parts of the fimbriae. *Int J Gynecol Pathol*

877   2013;32(5):444–453.

878   120.   **Sweet RA, Nickerson KM, Cullen JL, Wang Y, Shlomchik MJ.** B Cell-Extrinsic Myd88 and Fcer1g

879   Negatively Regulate Autoreactive and Normal B Cell Immune Responses. *J Immunol*

880   2017;199(3):885–893.

881   121.   **Bebbere D, Masala L, Albertini DF, Ledda S.** The subcortical maternal complex: multiple

882   functions for one biological structure? *J Assist Reprod Genet* 2016;33(11):1431–1438.

122.   **Iwata M, Takebayashi T, Ohta H, Alcalde RE, Itano Y, Matsumura T.** Zinc accumulation and

metallothionein gene expression in the proliferating epidermis during wound healing in mouse skin.

Histochem Cell Biol 1999;112(4):283–290.

123.   **Gava N, L Clarke C, Bye C, Byth K, deFazio A.** Global gene expression profiles of ovarian surface

epithelial cells in vivo. *J Mol Endocrinol* 2008;40(6):281–296.

124.   **Paillisson A, Dadé S, Callebaut I, Bontoux M, Dalbiès-Tran R, Vaiman D, Monget P.**

Identification, characterization and metagenome analysis of oocyte-specific genes organized in

clusters in the mouse genome. *BMC Genomics* 2005;6:76.

125.   **Higashi T, Tokuda S, Kitajiri S, Masuda S, Nakamura H, Oda Y, Furuse M.** Analysis of the

"angulin" proteins LSR, ILDR1 and ILDR2--tricellulin recruitment, epithelial barrier function and

implication in deafness pathogenesis. *J Cell Sci* 2013;126(Pt 4):966–977.

126.   **Stables MJ, Shah S, Camon EB, Lovering RC, Newson J, Bystrom J, Farrow S, Gilroy DW.**

Transcriptomic analyses of murine resolution-phase macrophages. *Blood* 2011;118(26):e192-208.

127.   **Rajkovic A, Lee JH, Yan C, Matzuk MM.** The ret finger protein-like 4 gene, Rfpl4, encodes a

putative E3 ubiquitin-protein ligase expressed in adult germ cells. *Mech Dev* 2002;112(1–2):173–

177.

128.   **Sontheimer RD, Racila E, Racila DM.** C1q: its functions within the innate and adaptive immune

responses and its role in lupus autoimmunity. *J Invest Dermatol* 2005;125(1):14–23.

129.   **Cai C, Tamai K, Molyneaux K.** KHDC1B is a novel CPEB binding partner specifically expressed in

mouse oocytes and early embryos. *Mol Biol Cell* 2010;21(18):3137–3148.

130.   **Suto F, Murakami Y, Nakamura F, Goshima Y, Fujisawa H.** Identification and characterization of

a novel mouse plexin, plexin-A4. *Mech Dev* 2003;120(3):385–396.

131.   **Choi Y, Yuan D, Rajkovic A.** Germ cell-specific transcriptional regulator sohlh2 is essential for

906   early mouse folliculogenesis and oocyte-specific gene expression. *Biol Reprod* 2008;79(6):1176–

907   1182.

908   132.   **Sakai N, Bain G, Furuichi K, Iwata Y, Nakamura M, Hara A, Kitajima S, Sagara A, Miyake T,**

909   **Toyama T, Sato K, Nakagawa S, Shimizu M, Kaneko S, Wada T.** The involvement of autotaxin in

910   renal interstitial fibrosis through regulation of fibroblast functions and induction of vascular leakage.

911   *Sci Rep* 2019;9(1):7414.

912   133.   **Puttabyatappa M, Guo X, Dou J, Dumesic D, Bakulski KM, Padmanabhan V.** Developmental

913   Programming: Sheep Granulosa and Theca Cell-Specific Transcriptional Regulation by Prenatal

914   Testosterone. *Endocrinology* 2020;161(8). doi:10.1210/endocr/bqaa094.

915   134.   **Singh H, Li Y, Fuller PJ, Harrison C, Rao J, Stephens AN, Nie G.** HtrA3 Is Downregulated in Cancer

916   Cell Lines and Significantly Reduced in Primary Serous and Granulosa Cell Ovarian Tumors. *J Cancer*

917   2013;4(2):152–164.

918   135.   **Luo F, Jia R, Ying S, Wang Z, Wang F.** Analysis of genes that influence sheep follicular

919   development by different nutrition levels during the luteal phase using expression profiling. *Anim*

920   *Genet* 2016;47(3):354–364.

921   136.   **Cheng Y, Kawamura K, Deguchi M, Takae S, Mulders SM, Hsueh AJW.** Intraovarian thrombin

922   and activated protein C signaling system regulates steroidogenesis during the periovulatory period.

923   *Mol Endocrinol* 2012;26(2):331–340.

924   137.   **Hatzirodos N, Glister C, Hummitzsch K, Irving-Rodgers HF, Knight PG, Rodgers RJ.**

925   Transcriptomal profiling of bovine ovarian granulosa and theca interna cells in primary culture in

926   comparison with their in vivo counterparts. *PLoS One* 2017;12(3):e0173391.

927   138.   **Lawrence JB, Oxvig C, Overgaard MT, Sottrup-Jensen L, Gleich GJ, Hays LG, Yates JR 3rd,**

928   **Conover CA.** The insulin-like growth factor (IGF)-dependent IGF binding protein-4 protease secreted

929 by human fibroblasts is pregnancy-associated plasma protein-A. *Proc Natl Acad Sci U S A*

930 1999;96(6):3149–3153.

931 139. **Hu S, Tamada K, Ni J, Vincenz C, Chen L.** Characterization of TNFRSF19, a novel member of the

932 tumor necrosis factor receptor superfamily. *Genomics* 1999;62(1):103–107.

933 140. **Reverchon M, Bertoldo MJ, Ramé C, Froment P, Dupont J.** CHEMERIN (RARRES2) decreases in

934 vitro granulosa cell steroidogenesis and blocks oocyte meiotic progression in bovine species. *Biol*

935 *Reprod* 2014;90(5):102.

936 141. **Guerrero-Juarez CF, Dedhia PH, Jin S, Ruiz-Vega R, Ma D, Liu Y, Yamaga K, Shestova O, Gay DL,**

937 **Yang Z, Kessenbrock K, Nie Q, Pear WS, Cotsarelis G, Plikus MV.** Single-cell analysis reveals

938 fibroblast heterogeneity and myeloid-derived adipocyte progenitors in murine skin wounds. *Nat*

939 *Commun* 2019;10(1):650.

940 142. **Veldhuis JD.** Interactions among endocrine control systems in the regulation of ovarian

941 function. *Clin Biochem* 1981;14(5):252–257.

942 143. **Xie T, Wang Y, Deng N, Huang G, Taghavifar F, Geng Y, Liu N, Kulur V, Yao C, Chen P, Liu Z,**

943 **Stripp B, Tang J, Liang J, Noble PW, Jiang D.** Single-Cell Deconvolution of Fibroblast Heterogeneity

944 in Mouse Pulmonary Fibrosis. *Cell Rep* 2018;22(13):3625–3640.

945 144. **Kim J-M, Song K-S, Xu B, Wang T.** Role of potassium channels in female reproductive system.

946 *Obstet Gynecol Sci* 2020;63(5):565–576.

947 145. **Katoh D, Kozuka Y, Noro A, Ogawa T, Imanaka-Yoshida K, Yoshida T.** Tenascin-C Induces

948 Phenotypic Changes in Fibroblasts to Myofibroblasts with High Contractility through the Integrin

949 αvβ1/Transforming Growth Factor β/SMAD Signaling Axis in Human Breast Cancer. *Am J Pathol*

950 2020;190(10):2123–2135.

146.   **Machelon V, Gaudin F, Camilleri-Broët S, Nasreddine S, Bouchet-Delbos L, Pujade-Lauraine E,**

Alexandre J, Gladieff L, Arenzana-Seisdedos F, Emilie D, Prévot S, Broët P, Balabanian K. CXCL12

expression by healthy and malignant ovarian epithelial cells. *BMC Cancer* 2011;11:97.

147.   **Popple A, Durrant LG, Spendlove I, Rolland P, Scott IV, Deen S, Ramage JM.** The chemokine,

CXCL12, is an independent predictor of poor survival in ovarian cancer. *Br J Cancer*

2012;106(7):1306–1313.

148.   **Mitchell TS, Bradley J, Robinson GS, Shima DT, Ng Y-S.** RGS5 expression is a quantitative

measure of pericyte coverage of blood vessels. *Angiogenesis* 2008;11(2):141–151.

149.   **Nagaraja AK, Middlebrook BS, Rajanahally S, Myers M, Li Q, Matzuk MM, Pangas SA.** Defective

gonadotropin-dependent ovarian folliculogenesis and granulosa cell gene expression in inhibin-

deficient mice. *Endocrinology* 2010;151(10):4994–5006.

150.   **Kofler NM, Cuervo H, Uh MK, Murtomäki A, Kitajewski J.** Combined deficiency of Notch1 and

Notch3 causes pericyte dysfunction, models CADASIL, and results in arteriovenous malformations.

Sci Rep 2015;5:16449.

151.   **Maguchi M, Nishida W, Kohara K, Kuwano A, Kondo I, Hiwada K.** Molecular cloning and gene

mapping of human basic and acidic calponins. *Biochem Biophys Res Commun* 1995;217(1):238–244.

152.   **Camoretti-Mercado B, Forsythe SM, LeBeau MM, Espinosa R 3rd, Vieira JE, Halayko AJ,**

Willadsen S, Kurtz B, Ober C, Evans GA, Thweatt R, Shapiro S, Niu Q, Qin Y, Padrid PA, Solway J.

Expression and cytogenetic localization of the human SM22 gene (TAGLN). *Genomics*

1998;49(3):452–457.

153.   **Palandri A, L'hôte D, Cohen-Tannoudji J, Tricoire H, Monnier V.** Frataxin inactivation leads to

steroid deficiency in flies and human ovarian cells. *Hum Mol Genet* 2015;24(9):2615–2626.

154.   **Krawczyk KK, Skovsted GF, Perisic L, Dreier R, Berg JO, Hedin U, Rippe C, Swärd K.** Expression

of endothelin type B receptors (EDNRB) on smooth muscle cells is controlled by MKL2, ternary

complex factors, and actin dynamics. *Am J Physiol Cell Physiol* 2018;315(6):C873–C884.

155.   **Conley CA, Fritz-Six KL, Almenar-Queralt A, Fowler VM.** Leiomodins: larger members of the

tropomodulin (Tmod) gene family. *Genomics* 2001;73(2):127–139.

156.   **O'Shannessy DJ, Somers EB, Wang L-C, Wang H, Hsu R.** Expression of folate receptors alpha and

beta in normal and cancerous gynecologic tissues: correlation of expression of the beta isoform with

macrophage markers. *J Ovarian Res* 2015;8:29.

157.   **Weng J, Liao M, Zou S, Bao J, Zhou J, Qu L, Feng R, Feng X, Zhao Z, Jing Z.** Downregulation of

FHL1 expression in thoracic aortic dissection: implications in aortic wall remodeling and

pathogenesis of thoracic aortic dissection. *Ann Vasc Surg* 2011;25(2):240–247.

158.   **Hayashi K-G, Ushizawa K, Hosoe M, Takahashi T.** Differential gene expression of serine

protease inhibitors in bovine ovarian follicle: possible involvement in follicular growth and atresia.

Reprod Biol Endocrinol 2011;9:72.

159.   **Li L, Mo H, Zhang J, Zhou Y, Peng X, Luo X.** The Role of Heat Shock Protein 90B1 in Patients with

Polycystic Ovary Syndrome. *PLoS One* 2016;11(4):e0152837.

160.   **Chen AQ, Wang ZG, Xu ZR, Yu SD, Yang ZG.** Analysis of gene expression in granulosa cells of

ovine antral growing follicles using suppressive subtractive hybridization. *Anim Reprod Sci*

2009;115(1–4):39–48.

161.   **Lee JH, Berger JM.** Cell Cycle-Dependent Control and Roles of DNA Topoisomerase II. *Genes*

(Basel) 2019;10(11). doi:10.3390/genes10110859.

162.   **Tsuji T, Kiyosu C, Akiyama K, Kunieda T.** CNP/NPR2 signaling maintains oocyte meiotic arrest in

early antral follicles and is suppressed by EGFR-mediated signaling in preovulatory follicles. *Mol*

Reprod Dev 2012;79(11):795–802.

163.   **Xi G, Wang W, Fazlani SA, Yao F, Yang M, Hao J, An L, Tian J.** C-type natriuretic peptide

998   enhances mouse preantral follicle growth. *Reproduction* 2019;157(5):445–455.

999   164.   **Piersanti RL, Santos JEP, Sheldon IM, Bromfield JJ.** Lipopolysaccharide and tumor necrosis

1000   factor-alpha alter gene expression of oocytes and cumulus cells during bovine in vitro maturation.

1001   *Mol Reprod Dev* 2019;86(12):1909–1920.

1002   165.   **Alfieri C, Chang L, Zhang Z, Yang J, Maslen S, Skehel M, Barford D.** Molecular basis of APC/C

1003   regulation by the spindle assembly checkpoint. *Nature* 2016;536(7617):431–436.

1004   166.   **Rovani MT, Gasperin BG, Ilha GF, Ferreira R, Bohrer RC, Duggavathi R, Bordignon V, Gonçalves

1005   PBD.** Expression and molecular consequences of inhibition of estrogen receptors in granulosa cells

1006   of bovine follicles. *J Ovarian Res* 2014;7:96.

1007   167.   **Hatzirodos N, Irving-Rodgers HF, Hummitzsch K, Rodgers RJ.** Transcriptome profiling of the

1008   theca interna from bovine ovarian follicles during atresia. *PLoS One* 2014;9(6):e99706.

1009   168.   **Landry DA, Rossi-Perazza L, Lafontaine S, Sirard M-A.** Expression of atresia biomarkers in

1010   granulosa cells after ovarian stimulation in heifers. *Reproduction* 2018;156(3):239–248.

1011   169.   **Diaz FJ, Sugiura K, Eppig JJ.** Regulation of Pcsk6 expression during the preantral to antral follicle

1012   transition in mice: opposing roles of FSH and oocytes. *Biol Reprod* 2008;78(1):176–183.

1013   170.   **Blanchard JM.** Cyclin A2 transcriptional regulation: modulation of cell cycle control at the G1/S

1014   transition by peripheral cues. *Biochem Pharmacol* 2000;60(8):1179–1184.

1015   171.   **Gong D, Ferrell JEJ.** The roles of cyclin A2, B1, and B2 in early and late mitotic events. *Mol Biol

1016   Cell* 2010;21(18):3149–3161.

1017   172.   **Luo Y, Qiao X, Ma Y, Deng H, Xu CC, Xu L.** Irisin deletion induces a decrease in growth and

1018   fertility in mice. *Reprod Biol Endocrinol* 2021;19(1):22.

1019   173.   **Pallier C, Scaffidi P, Chopineau-Proust S, Agresti A, Nordmann P, Bianchi ME, Marechal V.**

1020   Association of chromatin proteins high mobility group box (HMGB) 1 and HMGB2 with mitotic

1021   chromosomes. *Mol Biol Cell* 2003;14(8):3414–3426.

1022   174.   **Chun SY, McGee EA, Hsu SY, Minami S, LaPolt PS, Yao HH, Bahr JM, Gougeon A, Schomberg**

1023   **DW, Hsueh AJ.** Restricted expression of WT1 messenger ribonucleic acid in immature ovarian

1024   follicles: uniformity in mammalian and avian species and maintenance during reproductive

1025   senescence. *Biol Reprod* 1999;60(2):365–373.

1026   175.   **Gao F, Zhang J, Wang X, Yang J, Chen D, Huff V, Liu Y-X.** Wt1 functions in ovarian follicle

1027   development by regulating granulosa cell differentiation. *Hum Mol Genet* 2014;23(2):333–341.

1028   176.   **Park M, Choi Y, Choi H, Roh J.** Wilms' tumor suppressor gene (WT1) suppresses apoptosis by

1029   transcriptionally downregulating BAX expression in immature rat granulosa cells. *J Ovarian Res*

1030   2014;7:118.

1031   177.   **Gassmann R, Carvalho A, Henzing AJ, Ruchaud S, Hudson DF, Honda R, Nigg EA, Gerloff DL,**

1032   **Earnshaw WC.** Borealin: a novel chromosomal passenger required for stability of the bipolar mitotic

1033   spindle. *J Cell Biol* 2004;166(2):179–191.

1034   178.   **Bertolin K, Meinsohn M-C, Suzuki J, Gossen J, Schoonjans K, Duggavathi R, Murphy BD.** Ovary-

1035   specific depletion of the nuclear receptor Nr5a2 compromises expansion of the cumulus oophorus

1036   but not fertilization by intracytoplasmic sperm injection. *Biol Reprod* 2017;96(6):1231–1243.

1037   179.   **Jones MC, Zha J, Humphries MJ.** Connections between the cell cycle, cell adhesion and the

1038   cytoskeleton. *Philos Trans R Soc Lond B Biol Sci* 2019;374(1779):20180227.

1039   180.   **Bellanger S, de Gramont A, Sobczak-Thépot J.** Cyclin B2 suppresses mitotic failure and DNA re-

1040   replication in human somatic cells knocked down for both cyclins B1 and B2. *Oncogene*

1041   2007;26(51):7175–7184.

1042   181.   **Huang Y, Sramkoski RM, Jacobberger JW.** The kinetics of G2 and M transitions regulated by B

1043   cyclins. *PLoS One* 2013;8(12):e80861.

1044   182.   **Berisha B, Rodler D, Schams D, Sinowatz F, Pfaffl MW.** Prostaglandins in Superovulation

1045   Induced Bovine Follicles During the Preovulatory Period and Early Corpus Luteum. *Front Endocrinol*

1046   *(Lausanne)* 2019;10:467.

1047   183.   **Kfir S, Basavaraja R, Wigoda N, Ben-Dor S, Orr I, Meidan R.** Genomic profiling of bovine corpus

1048   luteum maturation. *PLoS One* 2018;13(3):e0194456.

1049   184.   **Tamura K, Matsushita M, Endo A, Kutsukake M, Kogo H.** Effect of insulin-like growth factor-

1050   binding protein 7 on steroidogenesis in granulosa cells derived from equine chorionic gonadotropin-

1051   primed immature rat ovaries. *Biol Reprod* 2007;77(3):485–491.

1052   185.   **Duan WR, Parmer TG, Albarracin CT, Zhong L, Gibori G.** PRAP, a prolactin receptor associated

1053   protein: its gene expression and regulation in the corpus luteum. *Endocrinology* 1997;138(8):3216–

1054   3221.

1055   186.   **Hanaue M, Miwa N, Takamatsu K.** Immunohistochemical Characterization of S100A6 in the

1056   Murine Ovary. *Acta Histochem Cytochem* 2012;45(1):9–14.

1057   187.   **Hernandez-Gonzalez I, Gonzalez-Robayna I, Shimada M, Wayne CM, Ochsner SA, White L,**

1058   **Richards JS.** Gene expression profiles of cumulus cell oocyte complexes during ovulation reveal

1059   cumulus cells express neuronal and immune-related genes: does this expand their role in the

1060   ovulation process? *Mol Endocrinol* 2006;20(6):1300–1321.

1061   188.   **Zamberlam G, Lapointe E, Abedini A, Rico C, Godin P, Paquet M, DeMayo FJ, Boerboom D.**

1062   SFRP4 Is a Negative Regulator of Ovarian Follicle Development and Female Fertility. *Endocrinology*

1063   2019;160(7):1561–1572.

1064   189.   **Wang SJ, Liu WJ, Wu CJ, Ma FH, Ahmad S, Liu BR, Han L, Jiang XP, Zhang SJ, Yang LG.** Melatonin

1065   suppresses apoptosis and stimulates progesterone production by bovine granulosa cells via its

1066   receptors (MT1 and MT2). *Theriogenology* 2012;78(7):1517–1526.

1067   190.   **Lee-Thacker S, Choi Y, Taniuchi I, Takarada T, Yoneda Y, Ko C, Jo M.** Core Binding Factor β

1068   Expression in Ovarian Granulosa Cells Is Essential for Female Fertility. *Endocrinology*

1069   2018;159(5):2094–2109.

1070   191.   **Park JY, Jang H, Curry TE, Sakamoto A, Jo M.** Prostate androgen-regulated mucin-like protein 1:

1071   a novel regulator of progesterone metabolism. *Mol Endocrinol* 2013;27(11):1871–1886.

1072   192.   **Baufeld A, Koczan D, Vanselow J.** Induction of altered gene expression profiles in cultured

1073   bovine granulosa cells at high cell density. *Reprod Biol Endocrinol* 2017;15(1):3.

1074   193.   **Kapfhamer J, Waite C, Ascoli M.** The Gα(q/11)-provoked induction of Akr1c18 in murine luteal

1075   cells is mediated by phospholipase C. *Mol Cell Endocrinol* 2018;470:179–187.

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

Figure 1

Figure 2

A.

B.

C.

Smooth muscle-*Mfap5*

Fibroblast-like stroma - *Enpp2*

Steroidogenic theca - *Cyp17a1*

Early theca - *Hhip*

Steroidogenic Stroma - *Ptch1*

Figure 3

A.

Preantral – Cumulus (4807 cells - 27.3%)
Atretic (3984 cells - 22.6%)
Antral – Mural (3846 cells - 21.8%)
Mitotic (2539 cells - 14.4%)
Active CL (951 cells - 5.4%)
Luteinizing mural (843 cells -4.8%)
Regressing CL (657 cells - 3.7%)

B.

C.

Atretic- *Ghr*

Preantral/Cumulus- *Kctd14*

Luteinizing mural-*Oxtr*

CL- *Neat1*

Antral/Mural- *Mro*

Mitotic- *Top2a*

Figure 4

Figure 5

Figure 6