Partitioning variance in reproductive success, within years and across lifetimes

Robin S. Waples

[1]Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, WA, 98112 USA

Running head: Partitioning variance in reproductive success

Key words: population genetics, computer simulations, reproductive skew, effective population size

Corresponding author:
Robin Waples (robin.waples@noaa.gov)
Northwest Fisheries Science Center
2725 Montlake Blvd. East
Seattle, WA, 98112 USA
(206) 860-3254 (voice); (206) 860-3335 (FAX)

February 2022

Abstract

Variance in reproductive success (σ_k^2, with k = number of offspring)—is the stuff of evolution. Evolutionary change by natural selection requires variation in offspring number, and the ratio $\phi = \sigma_k^2 / u_k$ is the most important factor that reduces effective size (N_e) compared to census size (N). Various frameworks have been proposed to parse factors that contribute to σ_k^2, but none has focused on age-specific values of ϕ, which indicate the degree to which reproductive skew is overdispersed among individuals of the same age and sex. Here I show how an ANOVA framework can be used to partition σ_k^2 into between-age and within-age components. For seasonal reproduction, the between-age effect depends on mean age-specific fecundity (b_x, available in many published life tables), but few empirical data are available on the within-age effect, which depends on ϕ_x. Analysis of variance in lifetime reproductive success ($\sigma_{k\bullet}^2$) is more complicated; however, by using age at death rather than age as the treatment, the same ANOVA framework can be used to partition overall $\sigma_{k\bullet}^2$ into three components: between-treatment, within-treatment, and longevity. For both seasonal and lifetime reproduction, I also show how to quantify the expected contribution of random demographic stochasticity to these variance components. Analyses of empirical and simulated data illustrate the new methods and show that the largely-neglected within-age effect a) typically represents a substantial component of the overall variance in offspring number (even under a null model of random reproductive success), and b) can dominate the overall variance when $\phi_x > 1$.

Introduction

Variance in reproductive success—number of offspring, k, contributed to the next generation—is the stuff of evolution. Evolution that results from natural selection acting on heritable genetic variation requires variation in offspring number among individuals. Crow (1958) defined what has come to be known as the Opportunity for Selection as $I = \sigma_k^2/(u_k)^2$, where u_k and σ_k^2 are the mean and variance in offspring number: for a given mean fitness, selection has more scope within which to operate when σ_k^2 is large. In all finite populations (which is to say, all real populations in nature), genetic drift also contributes a random component to evolution that is inversely proportional to the effective population size, N_e. N_e is generally less (sometimes a great deal less) than the census size, N, and the most important factor that reduces N_e compared to N is the ratio of variance to mean offspring number, $\phi = \sigma_k^2/u_k$. This term appears in the denominator of demographic formulas for calculating N_e (see next section), so N_e is reduced when σ_k^2 is large compared to u_k (i.e., when variance is overdispersed compared to the random Poisson expectation).

Although discrete-generation models are the simplest and most tractable, most real species are age or stage structured, so it is important to consider two different frameworks for measuring reproductive success: within time periods, and across lifetimes (quantified as lifetime reproductive success, or *LRS;* aka lifetime reproductive output, van Daalen and Caswell 2017). Many species have strongly seasonal (birth-pulse) reproduction, in which case all reproduction can be considered to occur at integer ages around individual birthdays (Felsenstein 1971; Caswell 2001), and that is the model considered here. Every time period (hereafter assumed to be years), all mature adults have an opportunity to reproduce. In this case, one can measure the mean and variance among all adults reproducing in one year and use those values to calculate annual *I* and annual effective size, which in this case would be the effective number of breeders, N_b. For seasonal reproduction, two systematic components contribute to the overall variance, σ_k^2: an among-age effect, and a within-age effect. The among-age effect depends on how expected reproductive success varies with age, which is reflected in the vector of age-specific expected fecundity (b_x) values from a standard life table. The within-age effect depends on $\phi_x = \sigma_{k,x}^2/u_{k,x}$ = the ratio of variance to mean offspring number for individuals of age *x*.

For lifetime reproduction, the relevant metric is variance in *LRS*, $\sigma_{k\bullet}^2$ (Hill 1972; Brown 1988; Tuljapurkar et al. 2020). Lifetime $\sigma_{k\bullet}^2$ is affected by these within- and among-age effects, as well as another factor: longevity. All else being equal, individuals that live longer have more opportunities to reproduce, and this increases disparity in lifetime offspring number among individuals and increases $\sigma_{k\bullet}^2$. Caswell (2011) and van Daalen and Caswell (2017) use the term "Markov chains with rewards" to describe the random aspects of this process of accumulating *LRS*, and variation in longevity can be the dominant factor contributing to variance in *LRS* (e.g., Newton 1989). Furthermore, in many species (especially ectotherms with indeterminate growth) fecundity increases with age, in which case individuals receive an additional bonus for surviving to older ages.

In the AgeNe model, Waples et al. (2011) showed how to calculate expected lifetime $\sigma_{k\bullet}^2$ and generational N_e under Hill's (1972) model, which assumes stable age structure and independence of survival and reproduction over time. AgeNe relies on age-specific survivals and fecundities from a life table, and the vector of ϕ_x values that capture the within-age component of variation in reproductive success. However, two obstacles currently limit the ability to fully utilize this approach. First, although published life tables are available for

numerous species from a wide range of taxa (Fung and Waples 2017 used data for 78 animal and
plant species with birth-pulse reproduction), data for the key age-specific parameter ϕ_x are rarely
published. As a consequence, researchers generally must either a) assume that $\phi_x = 1$ for each
age and sex, which is equivalent to assuming random reproductive success among individuals of
the same age and sex (e.g., Wang et al. 2010; Waples et al. 2013), or b) explore the consequences
of assuming a range of hypothetical values for ϕ_x (e.g., Waples et al. 2018a). General patterns of
how ϕ_x varies among species and among ages within species are largely unknown.
The second obstacle is that no method has been developed that would allow researchers
to systematically explore the relative importance of the two factors (the within-age and among-
age effects) that determine the seasonal variance in reproductive success or the three factors
(these two, plus longevity) that determine lifetime variance in reproductive success. Over the
past several decades, several frameworks have been proposed for partitioning *LRS* into different
components (Arnold and Wade 1984; Brown 1988; Broekman et al. 2021), but none of these has
focused on the component that derives from variance in offspring number among individuals of
the same age and sex. This is an important limitation; as shown below, this component of
reproductive variance is generally a substantial fraction of the overall variance even when it is
caused entirely by random stochasticity, and it can dominate the overall variance when same-age
individuals have different expectations of reproductive success.
This paper has two main objectives, designed to address these two major obstacles. First,
I describe a simple ANOVA-like analytical framework within which researchers can partition
variance in seasonal reproductive success into within-age and between age effects, and can
partition variance in *LRS* into within-age, between-age, and longevity effects. In addition,
analytical methods are developed that allow researchers to calculate what fraction of these
variance components can be attributed to random stochasticity in reproduction and survival.
The second objective is to highlight the importance of the neglected within-age
component to the overall variance. Modern molecular tools and improved methods for parentage
analysis make it easier than ever to collect empirical data on variance in offspring number.
Understanding the relative magnitude of between-age and within-age effects should lead to
richer insights into mating systems and reproductive biology, as well as an increased ability to
predict evolutionary responses to environmental changes that can affect fitness.
96

Methods

98   See Table 1 for notation and definitions of terms. The treatments here assume a single,
99 closed population and, except as noted, deal with population parameters (u_k, σ_k^2) and reflect an
100 exhaustive inventory of the population.
101

DEMOGRAPHIC MODEL

The focal population is assumed to be iteroparous, with separate sexes. Analogous
methods apply to both males and females; here for simplicity I consider data for a single sex,
nominally female. In the discrete-time, seasonal reproduction model, age is indexed by *x*. At
age *x*, each individual produces on average b_x offspring and then survives to age *x*+1 with
probability v_x. Age at maturity (defined as the first age with $b_x>0$) occurs at age α and maximum
age is ω. Newborns (age 0) do not reproduce, so b_x is scaled to production of offspring that
survive to age 1, at which point they can be enumerated. Cumulative survival from age 1
through age *x* is $l_x = \prod_{i=2}^{x} v_{i-1}$, with $l_1=1$. In a constant population with each birth cohort
consisting of N_1 yearlings, the expected number of individuals of age *x* alive at any given time is

$E(N_x) = N_1 l_x$, and expected total census size is given by $E(N_T) = \sum E(N_x) = N_1 \Sigma l_x$. Because we
are interested in reproductive success, the focus here is on the adult population size, N_A
$=\sum_{x=\alpha}^{\omega} N_x$.
115

EFFECTIVE POPULATION SIZE

117   Mean and variance in offspring number are the major factors that determine effective
118   size. For seasonal reproduction in an age-structured species, the most relevant effective-size
119   metric is the effective number of breeders per year (N_b) because it relates directly to the annual
120   number of adults (N_A). For each sex, N_b can be calculated using the standard discrete-generation
121   formula for inbreeding effective size (Crow and Denniston 1988; Caballero 1994):

122
$$N_b = \frac{u_k N_A - 1}{u_k - 1 + \sigma_k^2 / u_k} \quad . \tag{1}$$

N_b can be calculated separately for males and females and an overall N_b obtained using Wright's
(1938) sex-ratio adjustment. Examples below show how to separate the within-age and between-
age contributions to reduction of the key ratio N_b/N_A.
N_b relates directly to all adults breeding in a single year, but most evolutionary theory
depends on effective size per generation (N_e). Calculating N_e in iteroparous species is
complicated by the necessity of integrating information across multiple episodes of reproduction.
Hill (1972) provided the most general solution:

130
$$N_e \approx \frac{4 N_1 T}{\sigma_{k_\bullet}^2 + 2} \quad , \tag{2}$$

131   where $T = \Sigma x l_x b_x / \Sigma l_x b_x$ is generation length and $\sigma_{k_\bullet}^2$ is the variance in lifetime number of
132   offspring produced by the N_1 individuals in a cohort.
133

VARIANCE COMPONENTS ANALYSIS
Data for seasonal reproduction
The underlying model for a simple one-way analysis of variance with one treatment is the
following:
$k_{i,x} = \mu + \tau_x + \varepsilon_{i,x}$,
where u is the parametric mean, τ_x is the effect of treatment x ($x = 1,n$), $k_{i,x}$ is the value of the
response variable for individual i in treatment x, and $\varepsilon_{i,x}$ is a random error term. In the ANOVA
sums of squares table, the sources of variation are partitioned into three components (SST = total
sum of squares of deviations, SSB = between-treatment sum of squares, and SSE = error or
within-treatment sums of squares), as follows:
144

145      $$SST = \sum_{i=1}^{N_x} \sum_{x=1}^{n} (k_{i,x} - \mu_k)^2 \tag{3a}$$
146      $$SSB = \sum_{k=1}^{n} N_x (\mu_{k,x} - \mu_k)^2 \tag{3b}$$
147      $$SSE = \sum_{i=1}^{N_x} \sum_{x=1}^{n} (k_{i,x} - \mu_{k,x})^2 \;, \tag{3c}$$
148

where n is the number of treatments, N_x is the number of measurements for treatment x and $\mu_{k,x}$
and μ_k are treatment and overall means, respectively. In the ANOVA model, the vector of
samples sizes (N_x) is considered fixed and the sums of squares are additive, such that $SST = SSB$
$+ SSE$.
In the version of the model used here, the treatments are ages of individuals, $k_{i,x}$ is the
number of offspring produced by parent i of age x, and N_x is the number of potential parents of

age x. To allow for delayed age at maturity, we define a new adult-lifetime parameter, which
represents the number of treatments (adult age classes) as $AL = \omega-\alpha+1$. Our focus is on
partitioning the overall variance in offspring number ($\sigma_k^2 = SST/N_A$) into within-age and between-
age components. This can be done by weighting the age-specific variance components by the
relative number of parents in each age class (i.e., the weights are the vector N_x/N_A).

The weighted variance in offspring number among all individuals of age x is

$$\sigma_{k,x}^2 = (N_x/N_A) \sum_{i=1}^{N_x}(k_{i,x} - \mu_{k,x})^2/N_x = \sum_{i=1}^{N_x}(k_{i,x} - \mu_{k,x})^2/N_A = (N_x/N_A)\phi_x\mu_{k,x} \,,$$

and summed across all ages, the within-age component of σ_k^2 is

$$\sigma_{k(Within)}^2 = \sum_{x=\alpha}^{\omega}\sigma_{k,x}^2 = \sum_{x=\alpha}^{\omega}\sum_{i=1}^{N_x}(k_{i,x} - \mu_{k,x})^2/N_A = SSE/N_A. \qquad (4)$$

Thus, the residual (within-age) variance component is the sum of the product of the age-specific
vectors for ϕ and mean offspring number, and it can be computed as the within-treatment sum of
squares divided by the total number of parents.
In an analogous fashion, each age-specific term in the between-age component of σ_k^2 is
given by $(N_x/N_T)(\mu_{k,x} - \mu_k)^2$, so the overall between-age component is

$$\sigma_{k(Between)}^2 = \sum_{x=\alpha}^{\omega}(N_x/N_A) (\mu_{k,x} - \mu_k)^2 = SSB/N_A. \qquad (5)$$

As with the sums of squares, the components of overall σ_k^2 are additive, such that $\sigma_{k(Within)}^2 +$
$\sigma_{k(Between)}^2 = \sigma_k^2 = SST/N_A$. The relative contributions of the two variance components to
overall σ_k^2, which are identical to the relative contributions to SST, are:
Between ages: $\sigma_{k(Between)}^2/\sigma_k^2$
Within ages: $\sigma_{k(Within)}^2/\sigma_k^2 = 1 - \sigma_{k(Between)}^2.$

In the ANOVA framework, random variation among individual datapoints affects both
SSE and *SSB*. In ecology and evolutionary biology, it is often of interest to determine whether
variation in reproductive success is greater than the random expectation, because this determines
a) the scope for the action of natural selection, and b) the degree to which effective size is less
than census size. Tuljapurkar et al. (2020) called this random source of variation "dynamic
heterogeneity," while Caswell (2009) and van Daalen and Caswell (2017) prefer the term
"individual stochasticity." The Wright-Fisher model of random reproductive success is a
cornerstone of theoretical and empirical population genetics, so it is an appropriate null model to
use as a reference point for seasonal reproduction. Wright-Fisher reproduction involves a
multinomial sampling process, with expected variance in offspring number across adults of all
ages of $E(\sigma_{k(Random)}^2) = \mu_k(N_A-1)/N_A$. The expected relative contributions of the two variance
components to this total are N_A-AL for the within-age random effect and AL-1 for the between-
age random effect (see Supporting Information and Equations S7 for details). Note that these
relative weights are the same as the degrees of freedom for *SSE* and *SSB* (respectively) from an
ANOVA table. These equations accurately predicted the actual values of these random variance
components in simulated populations (Table S1).

This variance-partitioning approach makes no assumptions about the distribution of the
response variable (offspring number) and can be applied generally to widely-available data for
age-specific vital rates from a life table. For consistency and to facilitate comparisons across
species, the life table should be standardized to values of survival and fecundity that produce a
stable population.   For each age, the within-age and between-age variance components can be
calculated as described above and summed to obtain components $\sigma^2_{k(Within)}$ and $\sigma^2_{k(Between)}$.
The resulting estimates would apply to a hypothetical population whose realized data exactly
agreed with expectations from the life table. A potential limitation of this approach is that it
requires age-specific estimates of ϕ, which most standard life tables do not provide; in that event,
the user could conduct a sensitivity analysis to examine how different plausible assumptions
about ϕ affect the results.
The variance-partitioning method also can be applied to empirical age-specific vital rates.
Because the variance-to-mean ratio (ϕ) depends on mean offspring number in a sample ($\bar{k}$), it is
best to rescale ϕ to its expected value for a stable population, which can be done using a method
proposed by Crow and Morton (1955):

213
$$\phi_2 = 1 + \frac{\bar{k}_2}{\bar{k}_1}\left(\phi_1 - 1\right) \ . \tag{6}$$

214   In Equation 6, $\bar{k}_1$ is mean offspring number in the sample, $\bar{k}_2$ is the target mean offspring
215   number (the respective b_x value from a life table scaled to constant N), ϕ_1 is computed from the
216   sample, and ϕ_2 is the rescaled variance-to-mean ratio. After this rescaling is done for each age
217   and sex, analysis can the proceed as described above for data from a life table.
218
Data for lifetime reproductive success
Whereas seasonal reproduction involves individuals of different ages reproducing at the
same time, analysis of *LRS* requires longitudinal tracking of multiple episodes of reproduction
over time by the same individuals. We are interested in partitioning the variance in *LRS*, each
term of which involves a summation across an adult lifetime. Computing the expected variance
of *LRS* thus involves computing the variance of the sums of multiple vectors, which requires one
to consider all the pairwise covariance terms involving reproductive output of the same
individuals at different ages. This makes an analytical solution challenging, especially for long-
lived species, and in any case it would be rare to have the data to adequately parameterize this
full model.
Fortunately, a simple modification provides traction in the analysis of *LRS*. The most
widely-used models for N_e when generations overlap (Felsenstein 1971; Hill 1972) assume that
1) survival from age x to age $x+1$ is independent of an individual's reproductive output in
previous years; and 2) an individual's reproductive output is uncorrelated over time. Given these
assumptions, it is possible to calculate the expected value of lifetime $\sigma^2_{k\bullet}$ and hence N_e per
generation, using the vital rates from a standard life table, plus the age-specific vector ϕ. The
program AgeNe (Waples et al. 2011) does this by grouping individuals by age at death
(*AgeDeath*). Each individual has a single *AgeDeath* and a single *LRS*, so by using age at death as
the treatment, lifetime reproductive output can be analyzed with the same ANOVA framework
used for seasonal reproduction. Individuals that die at the same age all had the same number of
seasonal opportunities to reproduce and the same expectation for *LRS*. In the ANOVA
formulation for *LRS*, therefore, age at death rather than age is the treatment.

Calculation of the variance components for *LRS* from a life table proceeds as follows.
[For simplicity in notation, here I assume that age at maturity is 1, so that age at death (denoted
by q) takes the range 1-ω, which is the same range as x = age]. The number of individuals in
each treatment is $D_q = N_q - N_{q+1}$, with $D_\omega = N_\omega$ because it is assumed no individuals live beyond
age ω. D_q thus is the number of individuals that were alive at age q and had an opportunity to
reproduce but died before reaching age $q+1$. Calculation of *LRS* is done separately for each birth
cohort. Whereas $\Sigma N_x = N_A$ is the total number of adults alive at any given time, $\Sigma D_q = N_1$ is the
number of individuals in each cohort. As with the analysis of seasonal reproduction, we assume
that the population parameters exactly follow the expectations from the life table. Because of the
independence assumptions, the expected means and variances of *LRS* for each age at death q are
additive and given by $E(LRS_q) = \sum_{x=1}^{q} b_x$ and $E[var(LRS_q)] = \sum_{x=1}^{q} \sigma_{k,x}^2 = \sum_{x=1}^{q} b_x \, \phi_x$. The
within-treatment sums of squares are these treatment-specific variances multiplied by the number
of individuals in each treatment, so the overall within-treatment sums of squares for *LRS* is
254
255       $SSE \bullet = \sum_{q=1}^{\omega} D_q \sum_{x=1}^{q} b_x \, \phi_x.$ (7)
256
257   The between-treatment sums of squares ($SSB\bullet$) has ω terms, each of which is
258 proportional to the squared difference between treatment-specific $\overline{LRS}_q$ and the overall mean
259 $\overline{LRS}$:
260       $SSB \bullet = \sum_{q=1}^{\omega} D_q (\overline{LRS_q} - \overline{LRS})^2.$
261
262 Because we assume a stable population (and equal adult sex ratio), $\overline{LRS} = 2$, so this becomes
263
264       $SSB \bullet = \sum_{q=1}^{\omega} D_q (\overline{LRS_q} - 2)^2,$ (8)
265
with $SST \bullet = SSE \bullet + SSB \bullet$.
Longitudinal studies integrate reproductive success across multiple time periods, and as a
consequence they introduce two additional factors that contribute to *var*(*LRS*): differences in
longevity, and (potentially) non-zero correlations in individual reproductive success over time.
As shown below, in the two-part ANOVA framework, these additional factors get subsumed into
the between- and within-treatment effects, respectively. Our simplified *LRS* model assumes
temporally-spaced reproductive events are uncorrelated, so the second factor can be ignored (this
assumption is later relaxed). The effect of differences in longevity can be quantified by
removing any differences in age-specific fecundity (i.e., by assuming constant b_x with age). In
that case, the between-treatment effect only reflects differences in age at death = longevity. In
Results this partitioning of the between-treatment effect is illustrated with a worked example.
In an empirical dataset with longitudinal data for *LRS*, the raw data are in a matrix with
N_1 rows (one for each member of the cohort) and *AL* columns (one for each age in the maximum
adult lifespan) (see Table S2 for an example). In each cell, $k_{i,x}$ gives the number of offspring (that
survive to age at enumeration) produced by individual i at age x. The value $k_{i,x} = 0$ is used if
individual i is alive at age x but produces no offspring, but $k_{i,x}$ = NA is used for individuals that
die before reaching age x. The total matrix has $AL*N_1$ cells, of which ΣN_x are not 'NA', so ΣN_x
represents the total number of potential reproductive events (individuals alive at specific ages)
for the cohort. LRS_i for each individual is the sum of its numbers of offspring across its lifetime,
$\overline{LRS}$ is the mean *LRS* for the cohort (which might not exactly equal 2 in an empirical dataset),

and $\sigma_{k\bullet}^2$ is the variance of *LRS*. Parsing this overall variance into its components proceeds by
sequentially eliminating one component from the data, as illustrated with a worked example in
Results.
In the life-table analyses of *LRS*, it was assumed there were no temporal correlations in
reproductive success. In real populations, positive or negative correlations are likely in many
datasets, and to the extent that these correlations occur they will affect the within-treatment sums
of squares. Methods to evaluate these effects are illustrated in Results.
We can also calculate the expected random contribution to $SST\bullet$ and $\sigma_{k\bullet}^2$ under a null
model that randomly assigns each of the total *LRS* offspring to a member of the cohort. Under
the null model, all reproductive events have the same expectation (denoted below by $E(k)$), so
expected *LRS* for each individual is directly proportional to its age at death. For example, under
the null model with constant fecundity, an individual with *AgeDeath* = 5 has an expected *LRS* 5
times as large as another member of the cohort with *AgeDeath* = 1. More generally, in the null
model $E(LRS_q) = q * E(k)$. Furthermore, in the null model population size is constant so
overall $\overline{LRS} = 2$. Given these equalities, the random expectations for the within-treatment and
between-treatment sums of squares are given by (see Supporting Information for details):

$$E(SSE\bullet_{random}) = E(k) \sum_{q=1}^{\omega} (D_q - 1) * q; \qquad (9)$$

$$E(SSB\bullet_{random}) = E(k) * \sum_{q=1}^{\omega} q * \frac{(D_q - 1)}{D_q} + \sum_{q=1}^{\omega} D_q * (q * E(k) - 2)^2 . \qquad (10)$$

The random between-treatment sum of squares has two components (Equation 10): The first
term accounts for random differences in mean age-specific reproductive success, and the second
term accounts for random differences in age at death (longevity). As with seasonal reproduction,
simulations showed that Equations 9 and 10 accurately predict the random components of
variance in lifetime reproductive success (Table S3).
309

SIMULATIONS

Simulations were run to confirm the accuracy of analytical results and to illustrate effects
of random stochasticity on variance partitioning. All simulations were done in R (R Core Team
2021), using code provided in the SI. Numbers in each age group were fixed across replicates, as
were total numbers of offspring produced in each time period. Numbers of offspring produced
by each age group varied randomly in each replicate, with the constraint that the overall total was
fixed. This allowed the realized, age-specific mean offspring number ($\mu_{k,x}$) to vary randomly.
Overdispersion was determined by the age-specific parameter $\phi_x = \sigma_{k,x}^2/b_x$, with $\phi_x = 1$ indicating
Poisson variance. To simulate the desired level of overdispersion, vectors of parental weights ***w***
were generated as described in Waples (2022), where w_i is the relative probability that individual
i will be the parent of a given offspring. In modeling lifetime reproductive output, parental
weights were either shuffled randomly each year (to model independence of relative reproductive
success over time) or retained across individual lifetimes (to model persistent individual
differences), again as described in Waples (2022).
324

Results

The variance components analysis is illustrated with worked examples using empirical, hypothetical, and simulated data.

DATA FOR SEASONAL REPRODUCTION

Empirical life table data

Application to empirical data is illustrated using age-specific vital rates (Table 2) for male black bears in Michigan. Male black bears have a maximum longevity of at least 20 years, but data for bears older than 10 are sparse so the last age class (11+) is a plus group that includes all older males. This example assumes annual production of 800 yearlings, of which half are males (primary sex ratio is equal in this population). The best fit for the empirical age-at-death data indicated a lower survival prior to age 3, after which it was constant at 0.67/year. The N_x column in Table 2 shows the expected numbers in each age class, rounded to the nearest integer for purposes of this example. Age-specific fecundity (b_x) and the variance-to-mean ratio (ϕ_x)— both based on genetic parentage analysis and ageing from teeth—have been scaled to values expected for a stable population (see Table S4A in Waples et al. 2018b for details). Age-specific $\sigma^2_{k,x}$ is the product of b_x and ϕ_x, the column B_x shows the total expected number of offspring produced by each age class, and these sum to the total cohort size of 800. Because yearling males do not reproduce, we focus our analyses on the 698 adults age 2 and above, and adult lifespan consists of $AL = 10$ age classes, from 2 to 11+. Each year these males produce a cohort of 800 offspring, so an average adult produces $\mu_k = 800/698 = 1.15$ offspring that survive to age 1. Note that in this population, expected male fecundity rises sharply through age 10 before declining in older adults. Furthermore, the distribution of reproductive success is skewed, and strongly so in males older than 5, for which ϕ_x is >10.

The next sets of columns illustrate calculations for the variance components analyses. The "sums of squares" columns show age-specific values for the within-treatment (*SSE*) and between-treatment (*SSB*) components of an ANOVA table, with age as the treatment, calculated using Equations 3a-b. The "variance" columns show contributions of the within-age ($\sigma^2_{k(Within)}$) and between-age ($\sigma^2_{k(Between)}$) components to the overall variance in offspring number, using Equations 4 and 5.

For this life table, the between-treatment sums of squares (*SSB*, arising from systematic changes in fecundity with age) total 959, but the within-treatment sums of squares (*SSE*) are much larger (5473), so $SST = SSB + SSE = 6432$. Thus, the within-age component is responsible for the fraction 5473/6432, or 85.1%, of the total sums of squares, with the remainder attributable to the treatment (changes in fecundity with age). Because the components of σ^2_k are all directly proportion to their respective sums of squares, proportional partitioning of σ^2_k into within-age and between-age effects leads to the same result.

It also is of interest to ask how much of this overall variation in offspring number can be attributed to chance. As shown in Table 2, under the null model of random reproductive success, the expected value of *SST*$_{random}$ is 798.9, or 12.4% of the total *SST*. In the ANOVA framework, the vast majority (>98%) of this stochasticity shows up under *SSE*$_{random}$. If we subtract the expected magnitudes of random effects, we get an estimate of the net (non-random) within-age and between-age effects. These amount to 14.7% of the overall sums of squares for *SSB* and 72.8% for SSE, and these proportional results are the same for partitioning the random, within-

age, and between-age effects of σ_k^2. These results can be neatly illustrated with a pie chart
(Figure 1), which also shows results for female black bears. In females, changes in fecundity
with age are more modest and ϕ is also generally lower (3.21 for all ages; see Table S4 for
details). As a consequence, the random component is relatively more important for females
(29% overall compared to 12.4% for males), and within-age effects still dominate the non-
random components (Figure 1).
The variance-partitioning results for black bears also can be used to assess the relative
importance of within-age and between-age effects in determining the ratio of effective number of
breeders (N_b) to number of mature individuals (N_A). For males, inserting values for μ_k (1.15), σ_k^2
(9.21), and N_A (698) into Equation 1 produces $N_b = 98$ and $N_b/N_A = 0.14$—an overall reduction of
86% compared the expected result if seasonal reproduction exactly followed the WF model.
Although the variance components are additive, because of the inverse relationship between σ_k^2
and N_b their effects in reducing the N_b/N_A ratio are not additive. Nevertheless, we can partition
the overall reduction in N_b/N_A into within-age and between-age components based on their
relative contributions to σ_k^2 (from Table 2 and Figure 1). Because the Wright-Fisher result that
$E(N_b/N_A) = 1$ already accounts for random demographic stochasticity in σ_k^2, we use the net,
greater-than-random variance components, which for males are (from Table 2) total $\sigma_{k(net)}^2 =$
8.07, with the within- and between-age components being 6.71 and 1.36, respectively. Of the
86% total reduction in N_b/N_A in male black bears, therefore, we can attribute the fraction
6.71/8.07, or 71%, to within-age effects and the remaining 15% to between-age effects (Figure
2). For females, σ_k^2 is lower (3.51) so the reduction in effective size is not as strong ($N_b/N_A =$
0.29, a 71% reduction), but this reduction is also dominated by within-age effects (Figure 2).
392
Synthetic data
Computer simulations based on vital rates for hypothetical species were used to explore
variance partitioning and sensitivity under a wider range of scenarios. Table 3 shows a life table
for one sex (nominally females) for a hypothetical species with maximum lifespan of 5 years,
age at maturity = 1 year, annual survival constant at $v_x = 0.5$, expected fecundity (b_x) that is
proportional to age, and expected variance in reproductive success that is 5 times the mean ($\phi_x =$
5 for all ages). For purposes of this example, we assume that the actual data perfectly reflect
these expectations.
Each year the 775 females produce 800 offspring (both sexes combined), so overall μ_k is
800/775 = 1.032. From Methods, the between-age sum of squared deviations is $SSB = \sum N_x(\mu_{k,x} -$
$\mu_k)^2$. The table shows these age-specific products, which sum to 285.2. The residual (within-
treatment) sum of squares is $SSE = \sum(N_x*\phi_x*\mu_{k,x})$, whose terms are also shown in Table 3 and
sum to 4,000, so the total sum of squares is $SST = SSB + SSE = 4,285.2$. The among-ages
relative variance component therefore is 285.2/4285.2 = 0.067, so even though fecundity
increases in proportion to age, this accounts for only 6.7% of the overall variance in offspring
number.
To further partition the overall *SST*, we first calculate the expected random contributions
to the residual sum of squares, using Equations S2, S3, and S6 and using the overall $\mu_k = 1.032$
as $E(k)$. These expected random components are [799, 4.1, 794.8] for SST_{random}, SSB_{random}, and
SSE_{random}, respectively. Therefore, random (~Poisson) variance contributes 100*799/4285 =
18.6% to the overall *SST*, and the overdispersed (> Poisson) component of the within-age effect
(3205 = 4000-795) accounts for almost ¾ of the total variance in offspring number.
415

DATA FOR LIFETIME REPRODUCTION

Life table data

Analysis of *LRS* data using a life table is illustrated in Table 4, which uses the same vital rates as Table 3, except that ϕ is 3 at each age instead of 5. Based on the numbers-alive-at-age values (N_x), the numbers that die at ages $q=1\text{-}5$ are $D_q = [200,100,50,25,25]$, with $\Sigma D_q = 400$ being the annual cohort size of females. Under the independence assumptions of our simple model, expected LRS_q for each treatment (age at death) is simply the cumulative b_x through age at death, and with constant ϕ the expected variance among individuals that die at the same age is $E(LRS_q)*\phi$. The sums of squares are then computed for each treatment using Equations 7 and 8, with the overall result that $SSE\bullet = 2400$, $SSB\bullet = 1874.8$, and $SST\bullet = 4274.8$. The overall variance in *LRS* is $\sigma^2_{k\bullet} = SST\bullet/N_1 = 4274.8/400 = 10.69$. A bit over half (56.1%) of the overall sums of squares is due to within-treatment effects and the remainder to between-treatment effects.

The between-treatment $SSB\bullet$ can be further partitioned into between-age and longevity effects as demonstrated in Table S5. In the second panel of Table S5, the analysis is repeated after removing any age-related differences in fecundity. This is accomplished by replacing all the b_x values in the life table by the single fixed value that produces a stable population ($\mu_k = 1.032$). When this is done, $SSE\bullet$ remains unchanged but $SSB\bullet$ drops to 611, which represents the magnitude of the effect due to random differences in longevity. The difference between $SSB\bullet$ with and without age-related changes in fecundity (1874.8-611 = 1263.8) therefore represents the among-age component to $SSB\bullet$. Thus, age-related differences in fecundity account for about two thirds of $SSB\bullet$ and random variation in fecundity about one third.

For this life table, random effects account for almost one third (32.9%) of the $SST\bullet$, and within-treatment effects account for the majority of the remainder. In this partitioning, essentially all of the effects of longevity are absorbed into the random component, so the net (non-random) contribution to $SSB\bullet$ comes from changes in fecundity with age (see Figure 3).

This example assumed that reproduction and survival were independent across time. To evaluate consequences of violating this assumption, simulations were conducted using the vital rates in Table 4 but under a scenario that generated positive correlations in reproductive success for the same individuals over time. This was accomplished by allowing each surviving individual to retain the parental weight it had in year 1. An index of temporal correlation was calculated as $cor(k_{i,1}, k_{i,2\text{-}5})$, which is the correlation between an individual's reproductive output in year 1 and the total number of offspring it produced for the rest of its life. When parental weights were randomly shuffled each year, total $SST\bullet$ varied randomly around a value a bit over 4200 and $SSE\bullet$ was consistently about a third higher than $SSB\bullet$ (Figure 4, left side)—all as expected from Table 4—and $|cor(k_{i,1}, k_{i,2\text{-}5})|$ was close to zero (<0.001). When parental weights were retained across individual lifetimes, $SSB\bullet$ was not affected, but $SSE\bullet$ more than doubled so $SST\bullet$ increased as well (Figure 4, right), and mean $cor(k_{i,1}, k_{i,2\text{-}5})$ jumped to 0.36. This result demonstrates that 1) persistent differences in individual reproductive success can substantially increase the variance in *LRS*, and that 2) these correlations affect only the within-treatment component of $\sigma^2_{k\bullet}$.

 Empirical longitudinal data

The steps in partitioning variance in *LRS* for empirical longitudinal data are best
illustrated with a simple example, based on a population with vital rates as shown in Table 4.
The left panel of Table S2 shows longitudinal data over the 5-year lifespan for a sample of 40
randomly-generated individuals. The right panel summarizes these data by treatment = age at
death. Calculation of the within-treatment and between-treatment sums of squares follows
Equations 7 and 8. Mean *LRS* = (total number of offspring)/(sample size of parents) is $\overline{LRS}$ =
78/40 = 1.95. Two ways of computing variance in *LRS* ($V_1 = SST \bullet / \Sigma D_x$ and $V_2 = var(LRS)$)
produce the same result (11.9) when V_2 is computed as the population variance. In this example,
the between-treatment $SSB \bullet$ (310.6) is almost twice as large as the within-treatment $SSE \bullet$
(165.3).
471

Discussion

OTHER WAYS TO PARTITION VARIATION IN REPRODUCTIVE SUCCESS
The number of components of reproductive success that potentially could be identified is
essentially unlimited [e.g., van Noordwijk and van Balden (1988) list data for 25 *LRS*
components for great tits (*Parus major*) from the Netherlands], but Brown (1988) focused on
four that are commonly used: survival to breeding age; adult lifespan; age-specific fecundity; and
offspring survival to age at maturity. In their analysis of empirical data for a water-strider,
Ferguson and Fairbairn (2001) focused on a similar partitioning: pre-adult survival, adult
lifespan, and daily fecundity. Arnold and Wade (1984) partitioned reproductive success and
Crow's Opportunity for Selection into sequential episodes of selection and showed that the
components are additive only if one includes a residual factor that accounts for all the covariance
terms. More recently, several authors interested in distinguishing 'pluck' from 'luck' (Snyder
and Elner 2018) have focused on parsing out the component of *LRS* that can be attributed to
random demographic stochasticity in survival and reproduction (van Daalen and Caswell 2017;
Tuljapurkar et al. 2020; Broekman et al. 2020; Lee et al. 2020). All of these approaches can
provide useful insights, depending on one's objectives and the kinds of data that are available.
The methods outlined here differ from these previous efforts in two major ways. First,
this paper focuses on the largely-neglected component of reproductive variance that arises from
differences in realized reproductive success among individuals of the same age and sex. Second,
whereas most previous approaches to partitioning focus primarily or entirely on *LRS*, here equal
treatment is given to partitioning reproductive variance for individual years or seasons. Analysis
of seasonal data is important because any long-term, longitudinal study begins with data
collected for individual years or seasons, and many empirical studies never encompass entire
lifespans for the focal species (Nishida 1989). Furthermore, the distribution of seasonal
reproductive success provides key insights into mating systems, and the most direct way to study
sexual selection is by collecting data on all mature individuals that co-occur at the same time and
place. That is not the case when analyzing *LRS*.
500

501 **THE NEGLECTED AGE-SPECIFIC PARAMETER ϕ**
502   The variance partitioning analysis emphasizes why it is important to pay more attention
503 to the within-age component of variance in offspring number. For example, even though mean
504 fecundity varies substantially with age in both male and female black bears (Tables 2 and S1),
505 the within-ages component of σ_k^2 is much larger than the between-ages component in both sexes

(Figure 1). Furthermore, just the random component of within-age variance in reproductive success is almost as large as the sum of the random and non-random between-age components in males, and much larger than the between-age components in females (Figure S1). This indicates that a full understanding of reproductive variance cannot be achieved without considering the within-age component, even when the within-age variance is not overdispersed ($\phi = 1$).

NULL MODELS

Accuracy of the analytical results for the random expectations for σ_k^2 and $\sigma_{k\bullet}^2$ were confirmed by simulations. These results apply to a null model where a) each year or season, all adults behave like a Wright-Fisher ideal population with equal expectations of reproductive success, and b) for each individual in a cohort, survival to year i and reproduction in year i do not depend on realized reproductive success in any previous year. For seasonal reproduction, it was shown (Equations S7) that the relative magnitudes of the within-age and between-age random variance components are directly proportional to the degrees of freedom from an ANOVA table. This explains why the proportion of overall random variance in σ_k^2 due to within-age effects $[(N_T - n)/(N_T - 1)]$ generally will be much larger than the proportion due to between-age effects $[(n - 1)/(N_T - 1)]$, where n is the number of treatments (ages in the adult lifespan).

For analysis of LRS, a third random component (longevity) is important to consider. In this case the treatments are formed by grouping individuals by age at death, and random variation in longevity appears in the between-treatment component of random $\sigma_{k\bullet}^2$ (Equation S9). The other between-treatment component results from age-related changes in expected fecundity. These two components can be isolated by setting all b_x to the single value that will produce a stable population. This removes the between-age effect, so the remainder can be attributed to random variation in longevity.

These are not the only possible null models, and other random factors can contribute to variation in offspring number—for example, chance variation in mating success can depend on the operational sex ratio (Jennions et al. 2012), and parental coordination in provisioning offspring can arise from shared environments (Ihle et al. 2019). It is tempting to attribute family-correlated mortality to viability selection, but this phenomenon also can simply reflect bad luck. For example, nest predation in birds can eliminate entire clutches and greatly increase variance in offspring number—but how much of that family-correlated mortality is due to poor parental genes (and hence indicative of natural selection), and how much simply reflects a chance event that affects entire families only because the juvenile offspring co-occur in space and time? In the latter case, although mortality can be considered random, fates of individuals are not independent with respect to family, and this causes random variation in offspring number to exceed the Poisson expectation. Researchers should keep these issues in mind when trying to parse stochastic and deterministic factors contributing to variance in reproductive success.

TEMPORAL AUTOCORRELATIONS AND LIFETIME REPRODUCTIVE SUCCESS

The effects on $\sigma_{k\bullet}^2$ of random variation in survival (and hence random variation in the number of opportunities to reproduce) can be assessed as described in the previous section. Variance in *LRS* can also be strongly affected by correlations in individual reproductive success over time, and these correlations (if any) contribute to the within-treatment component of $\sigma_{k\bullet}^2$. Consistent positive correlations (e.g., some individuals consistently above or below average at producing offspring; see Figure 4) increase $\sigma_{k\bullet}^2$ (Lee et al. 2011). In contrast, negative correlations (often caused by skip breeding) generally reduce $\sigma_{k\bullet}^2$ because this life history

552 reduces the opportunity for certain individuals to consistently dominate production of offspring
553 (Waples and Antao 2014). This can be thought of as a type of variance-buffering 'portfolio'
554 effect.
555
LIMITATIONS
Some potential limitations of the analytical framework used here are important to keep in
mind. First, the demographic model is age-based and discrete-time and assumes birth-pulse
reproduction, which applies to a wide range of (but certainly not all) species. Modification for
continuous reproduction should be possible but is not attempted here. Many species' life
histories are more easily analyzed using stage-based models. Several recent authors (e.g., van
Daalen and Caswell 2017, Tuljapurkar et al. 2020, and colleagues) have used stage-based models
to separate random and deterministic components to variance in *LRS*. None of these, however,
has focused on the within-age component of $\sigma^2_{k\bullet}$.
Second, the current analyses follow the ANOVA model in treating the sample sizes—
numbers alive at age (N_x) for seasonal reproduction and numbers that die at each age (D_q) for
lifetime reproduction—as fixed. Which specific individuals survive is random, and the
proportion of offspring contributed by adults of each age is a random variable, subject to the
constraint that the total cohort size is constant, as is population size. Based on results of similar
models that have allowed random variation in the numbers surviving to each age (Waples et al.
2011; 2014), I expect that incorporating this feature would not appreciably change the results.
However, some species experience large swings in abundance in response to environmental
fluctuations, and in some cases vital rates can change with population density (Myhre et al. 2017;
Lee et al. 2020). Caution should be used in applying results obtained here to species
experiencing these kinds of population dynamic processes.
Finally, the ANOVA framework used here makes it easy to calculate the *F* statistic as the
ratio of two scaled sums of squares, in which case users might be tempted to use this statistic to
test the significance of the treatment effect (age for seasonal reproduction, age-at-death for *LRS*).
However, it should be clear from results presented here that the assumption of equal variances
often will be violated. Options for evaluating significance of the variance components for
empirical data include 1) using Monte-Carlo methods with an appropriate null model (as done
here and by van Daalen and Caswell (2017) and Tuljapurkar et al. (2020)); and 2) using
generalized least-squares approaches that allow one to model the within-treatment variance itself
as a function of some covariate (e.g., Zuur et al. 2009).
585
586 **Data availability statement**
587   This study did not generate any new empirical data, except by simulation. Code to
588 conduct the simulations and analyses described here is available in Supporting Information. The
589 author declares no conflict of interest.

REFERENCES

Arnold, S.J. and Wade, M.J. 1984. On the measurement of natural selection and sexual selection: theory. *Evolution*, **38**: 709–719.

Broekman, M.J., Jongejans, E. and Tuljapurkar, S., 2020. Relative contributions of fixed and dynamic heterogeneity to variation in lifetime reproductive success in kestrels (Falco tinnunculus). *Population Ecology*, *62*(4), pp.408-424.

Brown, D. 1988. Components of lifetime reproductive success. Pp. 439-453 in: T. Clutton-Brock, ed. *Reproductive success: studies of individual variation in contrasting breeding systems*. University of Chicago Press.

Caballero, A. (1994). Developments in the prediction of effective population size. *Heredity,* 73, 657–679.

Caswell, H. (2001). Matrix Population Models: Construction, Analysis, and Interpretation, 2nd edn. Sinauer Associates, Sunderland, MA.

Caswell, H. (2009). Stage, age and individual stochasticity in demography. Oikos, 118, 1763–1782.

Caswell, H. (2011). Beyond R0: demographic models for variability of life-time reproductive output. PLoS ONE, 6, e20809.

Crow, J. F. 1958. Some possibilities for measuring selection intensities in man. Human Biology 30:1:13.

Crow, J. F., and N. E. Morton. 1955. Measurement of gene frequency drift in small populations. Evolution 9:202-214.

Crow, J.F., and C. Denniston, 1988 Inbreeding and variance effective population numbers. *Evolution* 42: 482-495.

Felsenstein, J. (1971). Inbreeding and variance effective numbers in populations with overlapping generations. *Genetics,* 68, 581–597.

Ferguson, I.M. and Fairbairn, D.J., 2001. Is selection ready when opportunity knocks?. *Evolutionary Ecology Research*, *3*(2), pp.199-207.

Fung, HC, and RS Waples. 2017. Performance of IUCN proxies for generation length. Conservation Biology 31:883-893.

Hill, W.G. (1972). Effective size of population with overlapping generations. *Theoretical Population Biology* 3, 278–289.

Ihle, M., Pick, J.L., Winney, I.S., Nakagawa, S. and Burke, T., 2019. Measuring up to reality: null models and analysis simulations to study parental coordination over provisioning offspring. *Frontiers in Ecology and Evolution*, *7*, p.142.

Jennions, M.D., Kokko, H. and Klug, H., 2012. The opportunity to be misled in studies of sexual selection. *Journal of evolutionary biology*, *25*(3), pp.591-598.

Lee, A.M., S. Engen, & B.-E. Sæther, (2011) The influence of persistent individual differences and age at maturity on effective population size. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 278, 3303-3312.

Lee, A.M., Myhre, A.M., Markussen, S.S., Engen, S., Solberg, E.J., Haanes, H., Røed, K., Herfindal, I., Heim, M. and Sæther, B.E., 2020. Decomposing demographic contributions to the effective population size with moose as a case study. *Molecular ecology*, *29*(1), pp.56-70.

Myhre, A. M., Engen, S., & Sæther, B.-E. (2017). Effective size of density-dependent two-sex populations: The effect of mating systems. *Journal of Evolutionary Biology*, *30*(8), 1561–1575.

Newton, I. 1989. *Lifetime reproduction in birds*. Academic Press, London, 479 p.

Nishida, T., 1989. Is lifetime data always necessary for evaluating the "intensity" of selection?. *Evolution*, *43*(8), pp.1826-1827.

R Core Team (2021). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/

Snyder, R.E. and Ellner, S.P., 2018. Pluck or luck: does trait variation or chance drive variation in lifetime reproductive success?. *The American Naturalist*, *191*(4), pp.E90-E107.

Tuljapurkar, S., Zuo, W., Coulson, T., Horvitz, C. and Gaillard, J.M., 2020. Skewed distributions of lifetime reproductive success: beyond mean and variance. *Ecology letters*, *23*(4), pp.748-756.

van Daalen, S.F. & Caswell, H. (2017). Lifetime reproductive output: individual stochasticity, variance, and sensitivity analysis. Theor. Ecol., 10, 355–374.

van Noordwijk AJ and JH van Balen. 1988. The great tit, *Parus major*. Pp. 119-135 in: T. Clutton-Brock, ed. *Reproductive success: studies of individual variation in contrasting breeding systems*. University of Chicago Press.

Wang, J., Brekke, P., Huchard, E., Knapp, L.A., & Cowlishaw, G. (2010). Estimation of parameters of inbreeding and genetic drift in populations with overlapping generations. *Evolution*, *64*, 1704-1718.

Waples, RS. 2022. *TheWeight*: A simple and flexible algorithm for simulating non-ideal, age-structured populations. https://www.biorxiv.org/content/10.1101/2022.01.05.475135v1.

Waples, R.S., C. Do, and J. Chopelet. 2011. Calculating N_e and N_e/N in age-structured populations: a hybrid Felsenstein-Hill approach. *Ecology* 92:1513-1522.

Waples, R.S., G. Luikart, J.R. Faulkner, D.A. Tallmon. 2013. Simple life history traits explain key effective population size ratios across diverse taxa. Proc. Royal Society London, Ser. B. 280: 20131339.

Waples, R.S., T. Antao, and G. Luikart. 2014. Effects of overlapping generations on linkage disequilibrium estimates of effective population size. Genetics 197:769-780.

Waples, R.S., and T. Antao. 2014. Intermittent breeding and constraints on litter size: consequences for effective population size per generation (N_e) and per reproductive cycle (N_b). Evolution 68:1722-1734 (doi:10.1111/evo.12384).

Waples RS, Grewe PG, Bravington MV, Hillary R, Feutry P. 2018a. Robust estimates of a high N_e/N ratio in a top marine predator, southern bluefin tuna. Science Advances 4:eaar7759.

Waples RS, Scribner K, Moore J, Draheim H, Etter D, Boersen, M. 2018b. Accounting for age structure and spatial structure in eco-evolutionary analyses of a large, mobile vertebrate. Journal of Heredity 109:709-723.

Wright, S., 1938. Size of population and breeding structure in relation to evolution. *Science*, *87*, pp.430-431.

Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A. and Smith, G.M., 2009. *Mixed effects models and extensions in ecology with R* (Vol. 574). New York: Springer.

Table 1. Notation

x	Age (in years)
α	Age at maturity
ω	Maximum age
q	Age at death
v_x	Probability of surviving from age x to age $x+1$
l_x	Cumulative probability of surviving to age x
N_1	Cohort size = number of offspring produced per time period that survive to age 1
N_x	Number of individuals of age x alive at any given time; $N_x = N_1 l_x$
D_q	Number of individuals that died after reaching age q; $D_q = N_q - N_{q+1}$
N_T	Total number of individuals of all ages alive at any given time; $N_T = \Sigma N_x$
N_A	Number of adults with age$\geq\alpha$ alive at any given time
N_e	Effective population size per generation
N_b	Effective number of breeders in one time period
b_x	Expected number of offspring for parents of age x in one time period
k_i	Actual number of offspring produced by parent i in one time period
μ_k	Parametric mean k_i for all parents in one time period
σ_k^2	Parametric variance of k_i for all parents in one time period
$\sigma_{k,x}^2$	Variance of k for adults of age x
ϕ_x	Ratio of variance to mean reproductive success in one time period for adults of age x; $\phi_x = \sigma_{k,x}^2/b_x$
LRS	Lifetime reproductive success = total number of offspring produced across an individual's lifetime
$\overline{LRS_q}$	Mean LRS for individuals that died at age q
$\sigma_{k\bullet}^2$	Variance in LRS among the N_1 individuals in a cohort
SST	Total sum of squared deviations across all treatments, as in an ANOVA table
SSB	The between-treatment sum of squares
SSE	The error or within-treatment sum of squares
w_i	Relative probability that adult i will be the parent of a given offspring

Table 2. Variance partitioning analysis for seasonal reproduction based on a life table for male black bears from Michigan. These vital rates produce a constant cohort size of 800 age-1 offspring, of which half are male, and an adult male population size of 698 age 2+ individuals. In the top panel, columns 1-8 show vital rates from Waples et al. (2018b). The next two columns show the age-specific within-age and between-age sums of squares, and the last two columns show the same components for σ_k^2. The bottom panel shows results combining data for all ages. See Table 1 for notation, Table S4 for comparable data for females, and text for explanation of the calculations.

								Sums of squares		Variance	
Age	v_x	l_x	N_x	b_x	B_x	ϕ_x	$\sigma_{k,x}^2$	SSE_x	SSB_x	$\sigma_{k,x(W)}^2$	$\sigma_{k,x(B)}^2$
1	0.612	1	400	0	0	1.00	0.00	-	-	-	-
2	0.612	0.612	245	0.577	141	1.1	0.63	155.2	79.4	0.22	0.11
3	0.670	0.375	150	0.577	87	1.93	1.12	167.3	48.6	0.24	0.07
4	0.670	0.251	100	0.577	58	2.77	1.60	159.8	32.4	0.23	0.05
5	0.670	0.168	67	1.180	79	3.61	4.26	285.2	0.1	0.41	0.00
6	0.670	0.113	45	2.321	104	10.81	25.08	1128.8	62.1	1.62	0.09
7	0.670	0.076	30	2.984	90	10.81	32.24	967.3	101.3	1.39	0.15
8	0.670	0.051	20	3.739	75	10.81	40.41	808.1	134.5	1.16	0.19
9	0.670	0.034	14	4.515	63	10.81	48.80	683.2	158.9	0.98	0.23
10	0.670	0.023	9	7.131	64	10.81	77.06	693.6	322.3	0.99	0.46
11	0.670	0.015	18	2.182	39	10.81	23.58	424.5	19.3	0.61	0.03
		11+	18								
	Adults	698			800			5473	959	7.84	1.37

	μ_k	1.15				
			%	E_{random}	Net	%
SSE		5473	85.1	787	4686	83.2
SSB		959	14.9	12	947	16.8
SST		6432		799	5633	
$\sigma_{k(W)}^2$		7.84	85.1	1.13	6.71	83.2
$\sigma_{k(B)}^2$		1.37	14.9	0.02	1.35	16.8
σ_k^2		9.21		1.14	8.07	

Component	%
Random	799	12.4
Within	4686	72.8
Between	948	14.7
Total	6432

Table 3.   Variance components analysis for seasonal reproduction in a hypothetical species with a 5-year lifespan, fecundity that increases with age, and age-specific variance in reproductive success 5 times the mean. The top panel shows age-specific data and the bottom panel shows results combining data for all ages. See text for discussion.

Age	l_x	N_x	b_x	ϕ_x	$\sigma^2_{k,x}$	SSB_x	SSE_x
1	1	400	0.561	5	2.807	88.7	1122.8
2	0.5	200	1.123	5	5.614	1.6	1122.8
3	0.25	100	1.684	5	8.421	42.5	842.1
4	0.125	50	2.246	5	11.228	73.6	561.4
5	0.0625	25	2.807	5	14.035	78.7	350.9
	N_A	775				285.2	4000.0

μ_k	1.032
σ^2_k	5.529
ϕ	5.356

		Fraction	E_{Random}	Net	Component	%
SST	4285.2		799.0	3486.2	Random	18.6
SSB	285.2	0.067	4.1	281.1	Between	6.6
SSE	4000.0	0.933	794.8	3205.2	Within	74.8

Table 4. Variance components analysis for lifetime reproduction in a hypothetical species with vital rates as those in Table 3, but with $\phi = 3$ at each age. Long = longevity. The top panel shows age-specific data and the bottom panel shows results combining data for all ages. See text for discussion.

								Overall Sums of Squares				Random Expectations			
Age	N_x	b_x	ϕ_x	$\sigma^2_{k,x}$	D_x	$E(LRS_x)$	$\sigma^2_{k,x\bullet}$	$SSE\bullet$	$SSD\bullet$	Age	Long	$SSE\bullet$	$SSD\bullet$	Age	Long
1	400	0.561	3	1.68	200	0.56	1.68	336.8	413.9	226.6	187.3	111.7	188.4	1.0	187.4
2	200	1.123	3	3.37	100	1.68	5.05	505.3	10.0	9.6	0.4	166.7	2.5	2.0	0.4
3	100	1.684	3	5.05	50	3.37	10.11	505.3	93.6	33.5	60.1	165.1	63.1	3.0	60.1
4	50	2.246	3	6.74	25	5.61	16.84	421.1	326.5	213.2	113.3	134.7	117.2	4.0	113.2
5	25	2.807	3	8.42	25	8.42	25.26	631.6	1030.7	780.9	249.8	202.1	254.6	5.0	249.6
	775				400			2400.0	1874.8	1263.8	611.0	780.4	625.7	15.0	610.7

$E(LRS)$ 2.0

$\sigma^2_{k\bullet}$ 10.69

		%	E_{Random}	Net	Component	%
$SST\bullet$	4274.8		1406.1	2868.7	Random	32.9
$SSE\bullet$	2400.0	56.1	780.4	1619.6	Within	37.9
$SSB\bullet$	1874.8	43.9	625.7	1249.0	Between	29.2
Age	1263.8	29.6	15.0	1248.8	Age	29.2
Longevity	611.0	14.3	610.7	0.3	Longevity	<0.1

Figure 1. Relative contributions of random (denoted by initial 'R'), net within-age, and net between-age effects to seasonal variance in offspring number (σ_k^2) in black bears from Michigan. Results for males are from Table 2 and those for females are from Table S4.

Figure 2. Relative contributions of within-age and between-age components of seasonal variance in offspring number (σ_k^2) to the N_b/N ratio for black bears from Michigan. Random variation in reproductive success already is accounted for in reducing the N_b/N ratio to 1.0, so results depicted apply to the net (>random) variance components. Across all adults breeding in a given year, σ_k^2 is higher in males, so N_b/N is lower (0.14) than in females (0.29). Based on the variance partitioning results in Figure 1, for both sexes most of the total reduction in N_b/N (86% for males and 71% for females) can be attributed to variance in offspring number among individuals of the same age.

Figure 3. Results of partitioning lifetime variance in offspring number, using data from Table 4. Of the overall $\sigma^2_{k\bullet}$, 56% (reddish slices) can be attributed to within-treatment effects and 44% (bluish slices) to between-treatment effects. Of the total, 32% (paler slices) can be attributed to random stochasticity, leaving 38% and 29% (bolder colors) for the net (non-random) within-treatment and between-treatment effects, respectively. Most of the stochastic contributions are attributable to random variation in reproductive success among individuals that die at the same age (RWithin) and random variation in longevity (RLongevity); random between-age effects (RAge) account for <1% of the total $\sigma^2_{k\bullet}$.

Figure 4. Results of simulations of *LRS* using the vital rates shown in Table 4. The *Y* axis shows the sum of squares for the three variance components, with age at death being the treatment. On the *X* axis, in replicates 1-500 parental weights were shuffled each year, so there were no persistent individual differences in reproductive success. In replicates 501-1000, the same parental weights were retained across individual lifetimes, which introduced positive correlations in realized reproductive success across time; this had no effect on *SSB* but sharply increased *SSE* and hence *SST*.