Title: Pairing-dependent plasticity in a dissected fly brain is input-specific and

requires synaptic CaMKII enrichment and nighttime sleep

Abbreviated title: Associative memory formation in a dish

Mohamed Adel, Nannan Chen, Yunpeng Zhang, Martha L. Reed, Christina Quasney,

and Leslie C. Griffith

Department of Biology and Volen National Center for Complex Systems, Brandeis University,

Waltham, MA 02454-9110, USA

Corresponding author:

Leslie C. Griffith

Dept. of Biology MS008

Brandeis University

415 South St.

Waltham, MA 02454-9110

Tel: 781 736 3125

FAX: 781 736 3107

Email: griffith@brandeis.edu

Number of Pages: 42 pages

Number of Figures: 7 Figures

Abstract: 245 words

Introduction: 650 words

Discussion: 1151 words

Conflict of Interest Statement: The authors declare no competing financial interests.

Acknowledgments: This work was funded by NIH R01 MH067284 to LCG. Stocks

obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537) were

used in this study. The authors thank Dr. Kohei Ueno (Tokyo Metropolitan Institute of

Medical Science) for support, encouragement, and excellent advice during the

development of this project.

Abstract

In *Drosophila*, *in vivo* functional imaging studies revealed that associative memory formation is coupled to a cascade of neural plasticity events in distinct compartments of the mushroom body (MB). In-depth investigation of the circuit dynamics, however, will require an *ex vivo* model that faithfully mirrors these events to allow direct manipulations of circuit elements that are inaccessible in the intact fly. The current *ex vivo* models have been able to reproduce the fundamental plasticity of aversive short-term memory, a potentiation of the MB intrinsic neurons (Kenyon cells; KCs) responses after artificial learning *ex vivo*. However, this potentiation showed different localization and encoding properties from those reported *in vivo* and failed to generate the previously reported suppression plasticity in the mushroom body output neurons (MBONs). Here, we develop an *ex vivo* model that recapitulates behaviorally evoked plasticity in the KCs and MBONs. We demonstrate that this plasticity accurately localizes to the MB $\alpha'3$ compartment and is encoded by a coincidence between KCs activation and dopaminergic input. The formed plasticity is input-specific, requiring pairing of the conditioned stimulus (CS) and unconditioned stimulus (US) pathways; hence we name it pairing-dependent plasticity (PDP). PDP formation requires an intact *CaMKII* gene and is blocked by previous-night sleep deprivation but is rescued by rebound sleep. In conclusion, we show that our *ex vivo* preparation recapitulates behavioral and imaging results from intact animals and can provide new insights into mechanisms of memory formation at the level of molecules, circuits, and brain state.

Significance Statement

1
The mammalian *ex vivo* LTP model enabled in-depth investigation of the

hippocampal memory circuit. We develop a parallel model to study the *Drosophila*

mushroom body (MB) memory circuit. Pairing activation of the conditioned stimulus and

unconditioned stimulus pathways in dissected brains induces a potentiation pairing-

dependent plasticity (PDP) in the axons of $\alpha'\beta'$ Kenyon cells and a suppression PDP in

the dendrites of their postsynaptic MB output neurons, localized in the MB $\alpha'3$

compartment. This PDP is input-specific and requires the 3' untranslated region of

CaMKII. Interestingly, *ex vivo* PDP carries information about the animal's experience

before dissection; brains from sleep deprived animals fail to form PDP while those from

animals who recovered 2 hours of their lost sleep form PDP.

Introduction

A neutral experience (conditioned stimulus or CS) can be remembered as positive or negative if closely followed by rewarding or punishing reinforcement (unconditioned stimulus or US). The ability to form this type of "associative" memory is phylogenetically conserved; *Drosophila* form robust associative memories (Tully and Quinn, 1985), most of which are encoded and stored in the mushroom body (MB) (de Belle and Heisenberg, 1994). The MB is a higher brain structure made of 15 distinct compartments. Each compartment is built on a scaffold of axons of one of the three main types of Kenyon cells (KCs; $\alpha\beta$, $\alpha'\beta'$, and γ). The KCs connect to mushroom body output neurons (MBONs) which project out of the MB to bias behavior (Aso et al., 2014a; Aso et al., 2014b; Li et al., 2020). The KC$\rightarrow$MBON synapses are modulated by dopaminergic neurons.

During aversive olfactory associative learning, an odor (the CS) activates a sparse group of KCs, such that this odor identity is represented across all MB compartments (Turner et al., 2008; Lin et al., 2014). Simultaneously, dopaminergic neurons from the protocerebral posterior lateral (PPL1) cluster are activated by the US, encoding negative prediction errors in MB compartments (Riemensperger et al., 2005; Schroll et al., 2006; Claridge-Chang et al., 2009; Mao and Davis, 2009; Aso et al., 2012). When a coincidence occurs between the KCs activation and the dopaminergic signal within a compartment, the KC$\rightarrow$MBON synapses in that compartment are depressed, biasing the circuit output to aversion (Sejourne et al., 2011; Cohn et al., 2015; Hige et al., 2015; Owald et al., 2015; Owald and Waddell, 2015).

Many studies have investigated the properties of this circuitry using *in vivo* calcium imaging in intact animals (for review, see Adel and Griffith, 2021). In contrast, explanted brains have been used mostly for establishing connectivity between neurons or interrogating a specific biochemical pathway; only a few studies have attempted to understand memory circuit logic *ex vivo* (Wang et al., 2008; Ueno et al., 2013; 17; Suzuki-Sawano et al., 2017). In the best-developed paradigm, Ueno et al., 2013 observed a potentiation of KC responses in the tips of the MB vertical lobes which they termed "long-term enhancement" (LTE). This paradigm laid the groundwork for developing *ex vivo* models of this circuit, but there were major differences between LTE and associative memory observed in intact animals. The most significant were that the plasticity was not specific to the $\alpha'\beta'$ KCs and that dopamine release by the US was not observed; it was only seen after CS+US coincidence (Ueno et al., 2013; 2017).

In this study, we establish an *ex vivo* paradigm that resolves these discrepancies and exhibits the cardinal features of associative learning. We show that pairing odor and punishment pathway activation in dissected brains results in a localized potentiation of the $\alpha'\beta'$ KCs and suppression of their postsynaptic MBONs in the MB $\alpha'3$ compartment. Because both KC potentiation and MBON suppression are strictly dependent on temporal coincidence of the CS and US, we term this paradigm "pairing-dependent plasticity" (PDP). We show that like the CS-specificity of associative memories, PDP is specific to the subset of odor-representing projection neurons activated during the artificial training. We also provide evidence that dopamine is released by activation of the US pathway and does not require CS+US coincidence.

This *ex vivo* paradigm can be used for obtaining new mechanistic insight into

memory formation at the molecular and circuit levels. We present data indicating that

the 3'UTR of the *CaMKII* gene is critical for STM formation and that the primacy of α'

compartment plasticity in learning is due to differences in input/response relationships

between α and α'. Finally, we demonstrate that the ability of the *ex vivo* brain to be

plastic can be influenced by prior *in vivo* experience, as we report that brains of sleep-

deprived flies fail to form PDP, but as little as 2 hours of recovery sleep rescues this

learning impairment.

Materials and Methods

Fly husbandry. All fly stocks were cultured on standard food at room

temperature. Experimental flies were kept at 25°C and 70% relative humidity on a 12

hours light, 12 hours dark period. Fly lines used in this study include *VT030559-GAL4,*

MB027B split-GAL4 (Aso et al., 2014a), *GH146-GAL4, UAS-GCaMP6f, 20x UAS-*

GCaMP6f, UAS-jRCaMP1a, LexAop-P2X$_2$. *TH-lexA* was gifted to us from the Davis lab

(Berry et al., 2015) and *UAS-GRAB^{DA2m}* was gifted from the Li lab (Sun et al., 2020).

CaMKII^{UDel} flies are described in (Chen et al., 2021).

Ex vivo imaging and electrical stimulations. Brains from 4-8 days old female flies

were dissected in ice cold HL3 medium (NaCl, 70 mM; sucrose, 115 mM; KCl, 5 mM;

MgCl2, 20 mM; CaCl2, 1.8 mM; NaHCO3, 10 mM; trehalose, 5 mM; Hepes, 5 mM;

osmolarity: 395.4 mOsm; pH 7.3) (Stewart et al., 1994). Brains were then transferred to

an imaging chamber containing fresh HL3 saline and immobilized using tungsten pins

over the optic lobes. In the case of AL + AFV induction experiment, the brain was

dissected with the ventral nerve cord attached. The ventral nerve cord was then cut

using sharp scissors near its base, leaving one end of the AFV free. Glass suction

microelectrodes were used to apply the electrical stimulation to either the AL or the

AFV. Brains were always perfused with fresh saline throughout the experiment with a

flow rate of ~2 drops per second.

In pre-induction and post-induction AL stimulations, 5 trains of 20 pulses at

100Hz were applied. Pulse width is 1 millisecond and inter-pulse interval is 9

milliseconds. Inter-train interval is 15 seconds. Stimulation strength is 100 (low input

stimulation) or 200 (high input stimulation) μAmps. During induction, 12 AL stimulation

trains were applied with 5 seconds inter-train interval. AFV stimulation was similar to AL

stimulation, but AFV stimulation strength was adjusted to be above the threshold of KCs

responses; it varied between 500 to 1000 μAmps. The diameter of the AFV stimulation

electrode was large enough to suck in the free end of the AFV. Regarding the AL

electrode size, we noticed a relationship between the diameter of the AL electrode tip

and the input current such that applying a 100 μAmps input stimulation via a large

electrode tip diameter has a similar effect on baseline KCs calcium responses and on

the resultant plasticity as applying a 200 μAmps via a small electrode tip diameter. For

example, with a 100 μAmps input stimulation, PDP can be formed in the MB α3

compartment if the diameter of the AL electrode is significantly enlarged. Therefore, to

minimize variability within the same dataset, all electrodes used in any experiment in

this study were made at the same time using a p-97 micropipette puller (Sutter

Instruments) before data collection. To minimize variability across the different experiments, we kept the AL electrode size approximately one fourth of the AL size.

Images were captured using an ORCA Flash4.0 V3 sCMOS camera at 200 frames per second (except in the MBON imaging experiment using *MB027B split-GAL4>20x-UAS-GCaMP6f*, frame rate was reduced to 2 frames per second) with a 40x water immersion lens on an Olympus upright microscope BX50W1. Images were collected as 512x512 resolution and a binning factor of 2x2. Imaging was done using the HCImage Live software. Excitation of the used florescent sensors was done using the CoolLED pE-4000 LED source. For GCaMP and $GRAB_{DA2m}$, the 470 nm LED channel was used with an excitation filter Chroma 450/50 and emission filter FF01-525/45-25. jRCaMP excitation was done using the 550 nm LED channel and the excitation filter FF01-530/43-25 and an emission filter FF01-607/36-25. Calcium traces at every frame were calculated as $\Delta F/F_0 = (F-F_0)/F_0$ where F is the florescence value at a given frame and F_0 is the florescence value at baseline. Peak response to a stimulation was calculated by subtracting the average $\Delta F/F_0$ during the last second before a stimulation from the peak $\Delta F/F_0$ during stimulation. The responses to AL stimulations before and 15 minutes after induction were averaged to calculate the pre and post responses, respectively. PDP values or $\Delta F/F$ relative change was calculated as (post-pre)/pre.

The isolated pulse stimulator model 2100 and the perfusion system ValveLink 8.2 were triggered using a custom program written and controlled by the pClamp 11 software and the Digidata1550A digitizer.

Behavioral experiments. Aversive learning experiments were performed in an environmental room in red light at 25°C with 65% humidity. Flies were between 4-14 days old. Flies were given at least 10 minutes acclimation period in room before training or testing. Data for each experiment was pooled from at least 3 independent experimental days. The learning assays were performed as described by the Quinn lab (Tully and Quinn, 1985). The US was provided as 12 1-second 90- or 60-volts shocks during the 1-min CS-US pairing. 10% 4-methylcyclohexanol (MCH) and 3-octanol OCT were used as the CS odors. Flies were then given a 2-minute rest. Testing involved 2 minutes of simultaneous exposure to CS odors, after which flies choosing either odor were counted. A performance index (PI) was calculated for each trial as (number of flies choosing the conditioned odor) - (number of flies choosing the not-conditioned odor) / (total number of flies). This PI was averaged between reciprocal trials where one of the odors was conditioned in one trial and the other odor was conditioned in the other to calculate the Learning Index (LI). To confirm the *CaMKII^{UDel}* flies sensitivity to electric shock, 2-minute preference tests were performed during which flies chose between a stimulus vial (24 spaced 1-s 90 V shocks) and a neutral vial.

Sleep Assay. Sleep deprivation was done in 25°C incubators on a 12 h light, 12 h dark cycles. Flies were individually loaded into glass sleep tubes containing a food mixture of 5% sucrose and 2% agar. Drosophila Activity monitors (DAM) system (TriKinetics, Waltham) was used to measure sleep. Sleep was defined as inactivity bouts of 5 or more minutes (Hendricks et al., 2000; Shaw et al., 2000). Flies were sleep deprived by turning on the shaker between ZT12-ZT24. DAM data was analyzed using a custom MATLAB program (Donelson et al., 2012).

Immunostaining. Adult fly brains were dissected in cold Schneider's Insect

Medium (Sigma, S0146), and then fixed in 4% PFA solution for 30 mins at room

temperature. Fixed brains were washed 3x30 mins in 0.5% Triton-PBS (PBST) solution,

blocked in 10% normal goat serum solution for 1 hour, and incubated in mouse anti-

CaMKII antibody (1:10,000, Cosmo) for 3 days. CaMKII antibody solutions were

removed, and samples washed in PBST solution for 3x30 mins. Samples were then

incubated in Alexa Fluor 633 anti-mouse antibody (Invitrogen) overnight, then washed

3x30 mins in PBST solution and mounted in the Vectashield mounting medium. Images

were taken under a 20x objective lens with the same settings using a Leica SP5

confocal microscope. The images were analyzed by ImageJ software. For the intensity

of MB regions, the middle slices were selected as the representative pictures, and mean

intensity of all MB lobes were quantified.

Immunoblotting. 100 5 day old *Canton-S* wildtype (WT) and *CaMKII^{UDel}* flies were

frozen on dry ice and vortexed to remove heads. Fly heads were separated from the fly

bodies using a sieve. The heads were then homogenized in loading buffer (4X Bolt

LDS, Invitrogen, Novex with 5% β-mercaptoethanol added) and heated for 10 min.

Proteins were separated by SDS-PAGE (Bolt, Bis-Tris Protein Gels, Invitrogen) and

transferred to a nitrocellulose membrane (GE Healthcare). Membrane was blocked

(Blocking Buffer for Fluorescent Western Blotting, Rockland Immunochemicals) and

then incubated with Anti-dCaMKII Clone 18 (1:1000, CosmoBio) and anti-actin mAb C4

(1:1000, Millipore). The secondary antibody used was, DyLight 680 mouse. Membrane

was imaged using ChemiDoc system from Bio-Rad. Intensity of bands was calculated

using Adj. Volume (Int) within the ImageLab 6.0 software. Intensity of the CaMKII band

was normalized to that of actin in the same lane.

Experimental Design and Statistical Analysis. Female flies were used in all

imaging experiments due to their larger size. Both males and females aged 4 to 8 days

were used in other experiments unless noted otherwise. All statistical analyses were

performed in Prism 9 software. All tests were two tailed and confidence levels were set

at α = 0.05. Parametric tests were used for all experiments except the immunoblotting

experiment where non-parametric tests were used. The statistical tests, p-values, and

sample size for each experiment are listed in figure legends.

Results

Artificial aversive training induces a potentiation PDP in KCs

Previous studies had demonstrated that enhanced calcium responses similar to

those reported in the MB *in vivo* after aversive training can be achieved in dissected

brains (Wang et al., 2008; Tomchik and Davis, 2009; Ueno et al., 2013). To investigate

whether this plasticity recapitulates the formation of aversive memory, and to optimize

our protocol, we dissected the central nervous system (brain with attached ventral nerve

cord) of 4-8 days old mated female flies expressing the Ca^{2+} indicator GCaMP6f in the

MBs using the KCs driver (*VT030559-GAL4*). Dissected brains were pinned in an

imaging bath chamber and the cervical connective towards the base of the ventral nerve

cord was cut to allow electrical stimulation of the ascending fibers of the ventral nerve

cord (AFV) connecting the ventral nerve cord to the brain (Figure 1A). We used glass

suction electrodes to stimulate the antennal lobe projection neurons and the AFV to

activate the odor pathway and the electric shock pathway, respectively (Figure 1B).

Given that *in vivo* aversive STM formation is correlated with enhanced calcium

responses in the α'3 MB compartment but not in the α3 compartment (Krashes et al.,

2007; Wang et al., 2008; Cervantes-Sandoval et al., 2013; Zhang et al., 2019), we

decided to focus our calcium imaging on these two regions (Figure 1C). We found that

both α'3 and α3 compartments respond only to ipsilateral AL stimulation, but AFV

stimulation generated a calcium response in both compartments of both MBs.

Therefore, we imaged the MB ipsilateral to the stimulated AL.

To induce the formation of an aversive STM trace, we paired 12 trains of

stimulation of both the AL electrode and the AFV electrode, thus activating the CS and

the US pathways at the same time (paired induction). As controls, we repeated the

same induction paradigm but separated the AL stimulation from the AFV stimulation by

30 seconds (unpaired induction) or omitted the AFV stimulation (AL alone induction) or

omitted the AL stimulation (AFV alone induction) or simply allowed the brain to rest for

15 mins (no induction). We compared the change in calcium response to AL stimulation

before induction and 15 mins post induction (Figure 1D). We predicted that if this *ex*

vivo model truly recapitulated short-term aversive training, only the paired induction

should produce an enhancement of the calcium responses in the α'3 and none of the

stimuli should potentiate α3 (Krashes et al., 2007; Wang et al., 2008; Davis, 2011;

Cervantes-Sandoval et al., 2013). Indeed, this was the result we obtained (Figure 1E,F).

Interestingly, we noticed that the repetitive unpaired activation of KCs results in

suppression of their calcium responses (Figure 1F). This is reminiscent of

desensitization or habituation; a non-associative plasticity/memory described as a

decrement in neural responses to uninteresting, frequently-encountered stimuli whose

predictive value is negligible (Wilson and Linster, 2008; Wilson, 2009). These non-

associative processes have been documented in insects (Cho et al., 2004; Das et al.,

2011; Semelidou et al., 2018), rodents (Wilson, 1998; Best and Wilson, 2004), and

humans (Ferdenzi et al., 2014; Pellegrino et al., 2017).

Dopamine replaces the artificial US stimulus but does not replace the CS

Multiple behavioral and *in vivo* imaging studies have shown that dopaminergic

neurons encode the US valence information in the MB, with PPL1s providing aversive

reinforcement and PAMs providing appetitive reinforcement (Schwaerzel et al., 2003;

Riemensperger et al., 2005; Schroll et al., 2006; Kim et al., 2007; Claridge-Chang et al.,

2009; Mao and Davis, 2009; Aso et al., 2010; Aso et al., 2012; Pech et al., 2013; Aso et

al., 2014b; Yamagata et al., 2016; Handler et al., 2019). These studies suggest that the

US alone is sufficient to evoke dopamine release. However, mechanistic studies

employing the LTE paradigm reported no dopamine release after US pathway

stimulation alone. Strong dopamine release was only seen after coincident activation of

both the CS and the US pathways, and it was concluded that dopamine release is

downstream of the CS+US coincidence and does not encode the primary US

information (Ueno et al., 2017; Ueno et al., 2020). To directly address this discrepancy,

we expressed a G-protein-coupled receptor-activation-based dopamine sensor

($GRAB_{DA}$) (Sun et al., 2020) in the KCs. We then dissected the fly's central nervous

system and activated the US pathway by stimulating the AFV. We also expressed the

calcium indicator jRCaMP1a in the KCs to use as a reference for the strength of

activation (Figure 2A,B). We observed a robust dopamine release in the α'3 and a much weaker release onto the α3 compartment in response to the same AFV stimulation used in our induction experiments (Figure 2C,D). This result shows that dopamine release occurs in response to the US stimulus alone and does not require CS+US coincidence.

We next asked if dopamine perfusion can replace electrical stimulation of the AFV. We paired electrical stimulation of the AL with perfusion of different concentrations (1, 5 and 10 μM) dopamine and analyzed the KC response to AL stimulation before and after the pairing (Figure 2E,F). Similar to the plasticity seen after AL+AFV induction, we found that only the paired AL+ DA (5 or 10μM) induction induced an enhancement PDP in the α'3; pairing with 1 μM DA was not sufficient to induce PDP (Figure 2G). Also, in agreement with our observation that dopamine release in α3 was very weak compared to the α'3 compartment upon AFV stimulation, our protocol did not induce *ex vivo* PDP in the α3 compartment (Figure 2G). We also found that PDP (induced by AL + 10 μM DA) lasts for at least one hour after induction (Figure 2H).

Although these results supported a PDP model with direct dopamine modulation of KCs, we were concerned with the lack of synaptic specificity in the AL+DA induction, as dopamine was perfused onto the whole preparation. To address this concern, we repeated the experiment but replaced dopamine perfusion with activation of the dopaminergic neurons in the PPL1 cluster, which carry the negative valence US information to the MB (Schroll et al., 2006; Claridge-Chang et al., 2009; Mao and Davis, 2009; Aso et al., 2010; Aso et al., 2012; Burke et al., 2012; Aso and Rubin, 2016). We expressed the ATP-gated channel $P2X_2$ (Yao et al., 2012) in the PPL1 dopaminergic

cluster using the *TH-LexA* driver. We paired AL stimulation with application of 2.5 mM

ATP to activate PPL1s (Figure 2J). Again, we observed an enhancement PDP in the α'3

but not in the α3 compartment (Figure 2K).

Taken together, our findings agree with previous behavioral studies and *in vivo*

imaging studies and show that dopamine is released in response to the US alone and

carries US valence information to the MB to allow associative learning. Potentiation

PDP induced in our *ex vivo* paradigm also localizes to α' cells only, congruent with the

memory traces reported *in vivo* with aversive short-term memory.

PDP *ex vivo* plasticity is input specific

Associative memory is CS-specific. A fly that is aversively trained against an odor

shows aversive behavior to that odor and closely similar odors only, but does not

generalize the aversion to all odors (Barth et al., 2014). Olfactory information is sparsely

transmitted to KC dendrites, with the identity of the odor being determined by population

coding (Marin et al., 2002; Perez-Orive et al., 2002; Wong et al., 2002; Murthy et al.,

2008; Turner et al., 2008; Honegger et al., 2011). To mimic the application of 2 different

odors to the fly brain, we placed two suction glass electrodes onto the same antennal

lobe and spaced them as far as possible from each other to maximize the probability

that each electrode stimulates a distinct subset of projection neurons. Theoretically, a

certain degree of overlap between the two subsets could be allowed as long as the two

activated populations are sufficiently distinct. To test if the two electrodes activated

distinct populations of projection neurons, we expressed the calcium indicator

(GCaMP6f) in the projection neurons and divided the antennal lobe into multiple regions

of interest asking if the pattern of responses were equivalent for the two electrodes

(Figure 3-1 A,B in extended data). We found that spacing resulted in activation of

distinct populations of projection neurons, allowing us to provide two hypothetical odors

to the MB (Figure 3-1 C,D in extended data).

We then asked whether the observed *ex vivo* PDP is specific to the olfactory

input activated during induction. To challenge the olfactory-input specificity of our

preparation even more, we used the less specific induction method, AL+DA, in which

dopamine is impartially perfused to the whole brain to eliminate any specificity on the

US pathway side (Figure 3A). We recorded KC calcium responses to both electrode

stimulations before induction (pre). Then, we delivered a train of stimulations through

electrode #1, followed by a train of stimulations through electrode #2. Dopamine

perfusion started 5 seconds before the stimulation train for only one of the two

electrodes (paired) and stopped at the last pulse of the train, while normal HL3 saline

was perfused during the activation of the other electrode (unpaired). We then allowed

the brain to rest for 15 mins and tested the KC response to electrode #2 first then to

electrode #1 to eliminate any bias due to the order of stimulation (Figure 3B). An

increased calcium response was observed in α'3 after activation of the paired subset

compared to the unpaired one (Figure 3C,D). This result demonstrates that even with

the broader bath-dopamine induction paradigm, the plasticity achieved in our paradigm

is specific to the CS input activated during *ex vivo* training.

Artificial aversive training induces a suppression PDP in α'3 MBONs

While the α' branches of the KCs are potentiated in response to an aversively

trained odor in intact animals, responses in their postsynaptic MBONs are suppressed

(Sejourne et al., 2011; Owald et al., 2015; Owald and Waddell, 2015; Zhang et al.,

2019). To eliminate the possibility that our paradigm potentiates memory-relevant KCs

via a memory-irrelevant epiphenomenon, we needed to demonstrate that our *ex vivo*

training could produce a suppression PDP in MBONs. Therefore, we repeated the same

AL+DA induction described previously (Figure 2F), but this time we expressed

20xGCaMP6f with the *MB027b split-GAL4* driver to examine the responses in the $\alpha'3$

MBONs (MBON-α′3ap and MBON-α′3m) (Tanaka et al., 2008; Sejourne et al., 2011;

Aso et al., 2014a) (Figure 4A). We found that the $\alpha'3$ MBON response to AL stimulation

is suppressed after our artificial learning paradigm (Figure 4 B,C), agreeing with *in vivo*

results. These findings show that both the enhancement and suppression PDPs

achieved in our paradigm are learning-specific and are encoded as enhancement in the

presynaptic KCs and a suppression in the postsynaptic MBONs.

The 3'UTR mRNA of *CaMKII* is important for *ex vivo* PDP

We next turned our attention to exploring the utility of this paradigm for

understanding memory formation, choosing problems at several different levels of

analysis: molecular components of the memory machinery, organizational principles of

the circuitry and the role of brain state in gating plasticity. At the molecular level,

previous studies have demonstrated that CaMKII is important for synaptic plasticity and

memory formation in many species (Kelleher et al., 2004; Giese and Mizuno, 2013)

including *Drosophila melanogaster* (Griffith et al., 1993; Koh et al., 1999; Ashraf et al.,

2006; Malik et al., 2013; Mitchell et al., 2021). Recently, we found that the long 3'UTR

region of *CaMKII* mRNA is responsible for the activity-dependent synthesis of CaMKII in

presynaptic terminals at the larval neuromuscular junction (Kuklin et al., 2017) and for

the basal accumulation of axonal CaMKII protein in MB (Chen et al., 2021). We

hypothesized that the loss of the 3'UTR would impair associative plasticity by

decreasing synaptic CaMKII protein levels and disrupting the signaling machinery that is

triggered by the CS+US coincidence. We tested this idea and used our *ex vivo* PDP

preparation to gain insight into whether this effect is upstream or downstream of the

CS+US coincidence.

First, we asked if loss of the 3'UTR of *CaMKII* mRNA affects CaMKII levels. We

used animals in which the *CaMKII* gene was engineered using CRISPR/Cas9 to lack

the 3'UTR (*CaMKII*^Udel^, Chen et al., 2021) (Figure 5A). Immunostaining to quantify

CaMKII protein levels in the MB neuropil in both wildtype and *CaMKII*^UDel^ showed that

CaMKII levels are specifically decreased in synaptic regions (Figure 5B,C). We also

used western blotting to quantify total CaMKII levels and found a substantial decrease,

normalized to actin, in *CaMKII*^UDel^ flies compared to wildtype (Figure 5-1A,B in extended

data). This is in agreement with a previous study which found that mice expressing a

mutant form of *CaMKII* lacking the 3'UTR show decreased levels of *CaMKII* mRNA in

the dendritic, but not the somatic, region of hippocampal neurons (Miller et al., 2002).

We then tested the impact of the 3'UTR deletion on aversive STM. We found that

CaMKII^UDel^ flies showed a significant impairment in immediate memory performance

compared to wildtype flies (Figure 5D). This impairment was not due to sensorimotor

dysfunctionalities as both wildtype and *CaMKII*^UDel^ flies respond to electric shock and

odor (Figure 5-1C in extended data). We then tested the effect of the *CaMKII* mRNA

3'UTR deletion on potentiation PDP in KCs. We used the previously described AL+DA

induction protocol and examined KC responses to AL stimulation both before and 15

mins after either paired or unpaired induction. In the α'3 compartment, although both

wildtype and *CaMKII^UDel* genotypes showed statistically significant PDP relative to the

unpaired induction, the PDP in the wildtype flies was more than 5 times stronger than in

the *CaMKII^UDel* flies (Figure 5E). We also noticed that the change in signal in the α3

compartment in wildtype flies after paired induction was statistically different from the

unpaired induction. However, this does not translate into formation of PDP in that

compartment as this change was not different from zero (one-sample t-test, p-value

0.947). This is likely due to a technical difference as the AL input stimulation in this

experiment was stronger (200 μAmps vs. 100 μAmp in previous experiments) which led

to a stronger suppression after the unpaired AL stimulation in the unpaired induction

(see also below). These results show that the 3'UTR mRNA of *CaMKII* is important for

memory and its loss impairs learning-induced plasticity. It also indicates that the effect

on the learning circuit is downstream of the CS+US coincidence and does not block the

CS+US coincidence detection machinery itself.

Stronger inputs to the MB can recruit the α3 compartment into the short-term

memory circuit

At the circuit level, it has been hypothesized that memory in flies, like humans

(McClelland et al., 1995; Dudai, 2012), undergoes systems consolidation: initial

potentiation of the α'β' lobes with a time-dependent transfer of potentiation to $\alpha\beta$. The

MB α'3 and α3 compartments are adjacent to each other. Both respond to odor (Turner

et al., 2008) and AL stimulation (Figure 6B), and both use the same coincidence

detector, *rutabaga* (Livingstone et al., 1984; Levin et al., 1992; Mao et al., 2004; Gervasi

et al., 2010). This raises the question of how the two compartments are able to play

different roles in the memory circuit and why only the α'β' cells show immediate PDP

both *ex vivo* and *in vivo*. To determine if this might be due to differences in the intrinsic

properties of the two classes of cells, we examined the basal responses of both

compartments to a ramp of AL stimulations without any pairings (Figure 6A). We found

that α'3 axons are recruited first, with very weak AL stimulation, while α3 only starts

responding at higher stimulation strengths (Figure 6B). There was also a distinct

difference in the maximum response levels: α'3 compartment responses plateau at a

much lower level than α3 (Figure 6B), likely explained by the fact that the α3

compartment receives more axons than α'3 (Li et al., 2020). Previous studies reported

that odors activate a small percentage of KCs (~5-12%), and that the odor responses in

α'β' cells are stronger than those of αβ cells (Turner et al., 2008; Inada et al., 2017).

This suggests that actual odor encoding is similar to the weak AL stimulation used in

our experiments (100 μAmp), recruiting only α'3 but insufficient to recruit α3. We

hypothesized that an increase in the AL stimulation intensity during *ex vivo* induction

might be sufficient to recruit α3 and allow PDP in that compartment as well. We

repeated both AL+AFV and AL+DA inductions, but this time with a 200 μAmp AL

stimulation intensity. Indeed, under these conditions we found that PDP occurred in

both the α'3 and α3 compartments (Figure 6C,D).

This result agrees with predictions from our previously published theoretical

model of the associative learning circuit in which the CS+US coincidence triggers a

recurrent loop between KCs and dopaminergic neurons which increases dopamine

release only onto cells receiving both inputs together (Adel and Griffith, 2021). The

weak response of the α3 compartment to weak AL stimulation is subthreshold for the

MB coincidence detector, and insufficient for forming PDP in α3. Stronger AL

stimulation puts α3 over threshold and triggers the coincidence detector in α axons,

thus increasing dopamine gain and forming PDP/memory.

Our model further predicts that there should be a certain dopamine concentration

above which dopamine alone is sufficient to potentiate both α'3 and α3 compartments

without any need for pairing, as this hypothetical high dopamine concentration bypasses

the need for CS+US coincidence detector activation of the gain control machinery that

increases local dopamine release. To test this prediction, we applied either 10 μM

dopamine (used in our previous experiments) or a high dopamine concentration (200

μM) to the dissected brains without pairing with AL stimulation. Indeed, 200 μM

dopamine alone was sufficient to potentiate both α'3 and α3 responses to AL

stimulation (100 μAmps) 15 minutes post dopamine application (Figure 6E). A similar

effect of high dopamine was reported by Ueno et al., 2013, but interpreted differently.

We show that pairing-independent plasticity resulting from very high dopamine, while

possibly implemented by the same machinery responsible for associative learning, does

not truly represent a memory trace as it lacks specificity to CS-activated synapses, likely

representing a generalized potentiated state of the MB. In contrast, the PDP seen with

lower dopamine concentrations relies on an interplay between the CS and US signals to

gate dopamine gain only at the synapses activated by the CS during training.

PDP is blocked by sleep deprivation and rescued by rebound sleep

Many of the most interesting questions about memory formation are ones that

involve the interaction of plasticity with the animal's internal state. To test the ability of

this preparation to retain traces of previous experience and allow interrogation of

memory mechanisms at this level, we looked at the effect of sleep on subsequent *ex*

vivo plasticity. Sleep is linked to memory across phyla. In *Drosophila*, perturbations of

sleep have been shown to impair both memory formation (Bushey et al., 2007; Seugnet

et al., 2008; Li et al., 2009; Seugnet et al., 2009; Donlea et al., 2014; Seidner et al.,

2015) and consolidation (Ganguly-Fitzgerald et al., 2006; Donlea et al., 2009; Bushey et

al., 2011; Dag et al., 2019); reviewed in (Goel et al., 2009; Diekelmann and Born, 2010;

Dissel et al., 2015; Donlea, 2019). To determine whether the plasticity we observe in the

MB is sleep-dependent, we mechanically sleep deprived entrained (12:12 light:dark)

flies for 12 hours during the ZT12-24 night period. Control flies were housed on a

different shaker in the same incubator, but the shaker was turned off. A third group of

flies received shaking for only the last 2 hours of the night (ZT22-24) to control for the

acute physical stress that may result from the shaker. The next morning (starting at

ZT0), we used the AL+DA paradigm to induce the artificial memory (Figure 7A). We

found that sleep-deprived flies did not exhibit KC enhancement PDP, while both the

sleep control group and the stress control group had normal PDP (Figure 7C). We then

asked if allowing sleep-deprived flies a period of recovery sleep could restore their

ability to learn. As little as 2 hours of rebound sleep was sufficient to restore the ability

to induce the same level of α'3 PDP observed in the control groups (Figure 7C).

Interestingly, this 2 hours minimum of rebound sleep that we found in our experiment to

be required for rescuing the *ex vivo* plasticity is similar to the period reported *in vivo* to

be required for sleep-deprived flies to restore the ability to form short-term memory (Li

et al., 2009). These findings demonstrate that our *ex vivo* paradigm not only forms the

plasticity that is correlated with short-term memory, but also accurately recapitulates the

dynamics of the learning memory circuit and its interplay with the sleep circuit. These

data also suggest that the effect of sleep-deprivation on the memory circuit is

downstream to sensory processing of both the CS and the US information and CS+US

coincidence. This supports previous indications that sleep-deprivation likely impairs

memory formation by downregulating dopamine receptors or other downstream

molecules in the dopamine signaling pathway (Seugnet et al., 2008).

Discussion

Drosophila neural circuits are traditionally studied by relating *in vivo* genetic and

chemical manipulations with their consequent behavioral outcomes, from which circuit

information can then be inferred (Olsen and Wilson, 2008; Simpson, 2009). More

recently, the advent of *in vivo* calcium imaging allowed for tracing neural activity in

actively behaving flies. Over more than a decade of such *in vivo* studies, the general

circuit mechanisms of associative memory have been discovered, but there are

limitations imposed by imaging the brain of an active intact fly (for review, see Adel and

Griffith, 2021). These include the relatively low signal to noise ratio, the inaccessibility of

multiple brain regions due to restrictions on imaging angles, the difficulty of doing acute

pharmacological studies, and the possible confounds of studying the brain of a

movement-restricted fly experiencing on-going stress. Taking inspiration from the way

the LTP hippocampal slice model revolutionized our understanding of mammalian

memory (Bliss and Lomo, 1973), we provide here an *ex vivo* model of *Drosophila*

memory which can overcome these limitations and offer a powerful preparation for

studying *Drosophila* memory circuits. Importantly, this model provides a framework for

investigating the dynamics of neural circuits in the fly brain.

Most of the previous studies investigating the associative learning circuit *ex vivo*

have focused on mapping connectivity (Cohn et al., 2015; Barnstedt et al., 2016;

Felsenberg et al., 2017; Felsenberg et al., 2018; Zhao et al., 2018) or characterizing a

specific biochemical pathway (Tomchik and Davis, 2009; Handler et al., 2019; Ueno et

al., 2020). Only a few *ex vivo* studies (Wang et al., 2008; Ueno et al., 2013; Suzuki-

Sawano et al., 2017; Ueno et al., 2017) have focused on understanding MB circuit logic.

In the LTE model developed by Ueno et al (2013), pairing a stimulation of the CS and

US pathways induced a potentiation of KC responses in the tips of the MB vertical

lobes, but LTE did not fully recapitulate other characteristics of associative memory

observed in intact flies. Here we develop a modified *ex vivo* model that resolves these

discrepancies, showing that the paired activation of odor and punishment pathways

induces appropriate plasticity at multiple nodes in the circuit: potentiation of KCs and

suppression of MBONs. PDP is also localized to the MB α'3 compartment and not seen

in α3, in alignment with most imaging studies in intact flies. Importantly, in this *ex vivo*

preparation, punishment information is relayed to the MB through dopaminergic release

from the PPL1 subset. Bath application of dopamine in our preparation does not

interfere with the specificity of associative learning since PDP is exclusively formed in

the cells that were active during the dopamine application. These data settle several

inconsistencies between previous *ex vivo* studies (Ueno et al., 2013; Ueno et al., 2017)

and the majority of *in vivo* reports (Schwaerzel et al., 2003; Riemensperger et al., 2005;

Schroll et al., 2006; Kim et al., 2007; Claridge-Chang et al., 2009; Mao and Davis, 2009; Aso et al., 2010; Aso et al., 2012; Pech et al., 2013; Aso et al., 2014b; Yamagata et al., 2016; Cognigni et al., 2018; Handler et al., 2019). We suggest that the genesis of the discrepancies was not due to any inherent difference between intact and *ex vivo* brains but was rather a consequence of technical considerations including stimulation strength, dopamine concentration and the sensor tools employed (see Adel and Griffith, 2021 for a more complete discussion).

An *ex vivo* preparation that recapitulates the cardinal features of the circuits underlying associative memory formation should be useful for mechanistic studies at the molecular, cellular and systems levels. We used our model to ask a new question about the innerworkings of the circuit at each of these levels. At the molecular level we demonstrated the importance of normal levels of CaMKII by manipulating the 3'UTR of *CaMKII* mRNA. Deletion of this region of the *CaMKII* gene drastically reduces the amount of CaMKII protein in synaptic regions and blunts the ability to form STM and to generate a potentiation PDP in KC axons. Our data argues that the role of this molecule is downstream of the CS+US coincidence detector, as we observe a much weaker PDP in *CaMKII*[Udel] flies. Whether the behavioral defect is due solely to the KC PDP defect is not completely clear since CaMKII likely has active roles at other circuit nodes (Mitchell et al., 2021).

At the cellular level we asked why STM and PDP form in the α'3 but not the nearby α3 compartment when both compartments respond to odors (Turner et al., 2008) and AL stimulation, and both receive dopaminergic input from the same PPL1 cluster (Masek et al., 2015). Previous work found that real odors cause activity in only 5-

12% of KCs and elicit a much higher spike rate in the $\alpha'\beta'$ KCs than in the $\alpha\beta$ KCs. We

found that low-intensity AL stimulation (100 μAmps) elicit a stronger response in the $\alpha'3$

than in the $\alpha3$ compartment, while high-intensity AL-stimulation (200 μAmps) causes

strong responses in the $\alpha3$ compartment and recruits it to the learning circuit. Coupling

this with our observation of lower dopamine release in $\alpha3$, suggests a model in which

odor presentation during associative learning causes subthreshold responses in $\alpha\beta$

cells such that the CS+US coincidence detector is not triggered, while the stronger

responses in the $\alpha'\beta'$ cells bypass this threshold, allowing plasticity the $\alpha'\beta'$ cells only.

This notion is in alignment with the previous finding that $\alpha'\beta'$ cells have a lower firing

threshold than $\alpha\beta$ cells (Inada et al., 2017). Further, It is possible that long-term memory

and the enhancement memory trace in the $\alpha\beta$ KCs after repetitive space training (Yu et

al., 2006) require a gradual potentiation of the $\alpha\beta$ KCs responses with every training

session such that the responses bypass the coincidence detection threshold after

several training sessions. Whether repetition of AL+DA pairings recruits PDP in the $\alpha3$

compartment remains unclear. It is also yet to be determined whether shortcutting the

circuit and recruiting $\alpha\beta$ cells in the first training session reduces the need for multiple

spaced training sessions in long-term memory formation.

 Lastly, we looked at the ability of the effects of prior experience, or brain state, on

the memory circuit to be retained in the *ex vivo* preparation. Excitingly, we found that

sleep deprived flies could not form PDP, but that as little as 2 hours of rest before

dissection allowed the brain to recover PDP formation. The complete abolition of PDP in

sleep deprived flies at first and the gradual recovery in plasticity afterwards (Figure 7D)

suggests that sleep converges on the memory circuit upstream of the CS+US

coincidence detector. Whether this involves regulation of dopamine receptors in the MB

during sleep remains to be determined. The ability to retain in some functional way the

internal state of the brain will allow this preparation to be used to understand how

memory formation is altered by global system alterations.

References

Adel M, Griffith LC (2021) The Role of Dopamine in Associative Learning in Drosophila: An Updated Unified Model. Neuroscience Bulletin 37:831-852.

Ashraf SI, McLoon AL, Sclarsic SM, Kunes S (2006) Synaptic protein synthesis associated with memory is regulated by the RISC pathway in Drosophila. Cell 124:191-205.

Aso Y, Rubin GM (2016) Dopaminergic neurons write and update memories with cell-type-specific rules. Elife 5.

Aso Y, Siwanowicz I, Bracker L, Ito K, Kitamoto T, Tanimoto H (2010) Specific dopaminergic neurons for the formation of labile aversive memory. Curr Biol 20:1445-1451.

Aso Y, Herb A, Ogueta M, Siwanowicz I, Templier T, Friedrich AB, Ito K, Scholz H, Tanimoto H (2012) Three dopamine pathways induce aversive odor memories with different stability. PLoS Genet 8:e1002768.

Aso Y, Hattori D, Yu Y, Johnston RM, Iyer NA, Ngo TT, Dionne H, Abbott LF, Axel R, Tanimoto H, Rubin GM (2014a) The neuronal architecture of the mushroom body provides a logic for associative learning. Elife 3:e04577.

Aso Y et al. (2014b) Mushroom body output neurons encode valence and guide memory-based action selection in Drosophila. Elife 3:e04580.

Barnstedt O, Owald D, Felsenberg J, Brain R, Moszynski JP, Talbot CB, Perrat PN, Waddell S (2016) Memory-Relevant Mushroom Body Output Synapses Are Cholinergic. Neuron 89:1237-1247.

Barth J, Dipt S, Pech U, Hermann M, Riemensperger T, Fiala A (2014) Differential associative training enhances olfactory acuity in Drosophila melanogaster. J Neurosci 34:1819-1837.

Berry JA, Cervantes-Sandoval I, Chakraborty M, Davis RL (2015) Sleep Facilitates Memory by Blocking Dopamine Neuron-Mediated Forgetting. Cell 161:1656-1667.

Best AR, Wilson DA (2004) Coordinate synaptic mechanisms contributing to olfactory cortical adaptation. J Neurosci 24:652-660.

Bliss TV, Lomo T (1973) Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. J Physiol 232:331-356.

Burke CJ, Huetteroth W, Owald D, Perisse E, Krashes MJ, Das G, Gohl D, Silies M, Certel S, Waddell S (2012) Layered reward signalling through octopamine and dopamine in Drosophila. Nature 492:433-437.

Bushey D, Tononi G, Cirelli C (2011) Sleep and synaptic homeostasis: structural evidence in Drosophila. Science 332:1576-1581.

Bushey D, Huber R, Tononi G, Cirelli C (2007) Drosophila Hyperkinetic mutants have reduced sleep and impaired memory. J Neurosci 27:5384-5393.

Cervantes-Sandoval I, Martin-Pena A, Berry JA, Davis RL (2013) System-like consolidation of olfactory memories in Drosophila. J Neurosci 33:9846-9854.

Chen N, Zhang Y, Adel M, Kuklin EA, Reed ML, Mardovin JD, Bakthavachalu B, VijayRaghavan K, Ramaswami M, Griffith LC (2021) Specialized presynaptic proteomes required for associative memory formation depend on enhanced axonal translation of pre-localized mRNA. Submitted.

Cho W, Heberlein U, Wolf FW (2004) Habituation of an odorant-induced startle response in Drosophila. Genes Brain Behav 3:127-137.

Claridge-Chang A, Roorda RD, Vrontou E, Sjulson L, Li H, Hirsh J, Miesenbock G (2009) Writing memories with light-addressable reinforcement circuitry. Cell 139:405-415.

Cognigni P, Felsenberg J, Waddell S (2018) Do the right thing: neural network mechanisms of memory formation, expression and update in Drosophila. Curr Opin Neurobiol 49:51-58.

Cohn R, Morantte I, Ruta V (2015) Coordinated and Compartmentalized Neuromodulation Shapes Sensory Processing in Drosophila. Cell 163:1742-1755.

Dag U, Lei Z, Le JQ, Wong A, Bushey D, Keleman K (2019) Neuronal reactivation during post-learning sleep consolidates long-term memory in Drosophila. Elife 8.

Das S, Sadanandappa MK, Dervan A, Larkin A, Lee JA, Sudhakaran IP, Priya R, Heidari R, Holohan EE, Pimentel A, Gandhi A, Ito K, Sanyal S, Wang JW, Rodrigues V, Ramaswami M (2011) Plasticity of local GABAergic interneurons drives olfactory habituation. Proc Natl Acad Sci U S A 108:E646-654.

Davis RL (2011) Traces of Drosophila memory. Neuron 70:8-19.

de Belle JS, Heisenberg M (1994) Associative odor learning in Drosophila abolished by chemical ablation of mushroom bodies. Science 263:692-695.

Diekelmann S, Born J (2010) The memory function of sleep. Nat Rev Neurosci 11:114-126.

Dissel S, Melnattur K, Shaw PJ (2015) Sleep, Performance, and Memory in Flies. Curr Sleep Med Rep 1:47-54.

Donelson NC, Kim EZ, Slawson JB, Vecsey CG, Huber R, Griffith LC (2012) High-resolution positional tracking for long-term analysis of Drosophila sleep and locomotion using the "tracker" program. PLoS One 7:e37250.

Donlea JM (2019) Roles for sleep in memory: insights from the fly. Curr Opin Neurobiol 54:120-126.

Donlea JM, Ramanan N, Shaw PJ (2009) Use-dependent plasticity in clock neurons regulates sleep need in Drosophila. Science 324:105-108.

Donlea JM, Pimentel D, Miesenbock G (2014) Neuronal Machinery of Sleep Homeostasis in Drosophila. Neuron 81:1442.

Dudai Y (2012) The restless engram: consolidations never end. Annu Rev Neurosci 35:227-247.

Felsenberg J, Barnstedt O, Cognigni P, Lin S, Waddell S (2017) Re-evaluation of learned information in Drosophila. Nature 544:240-244.

Felsenberg J, Jacob PF, Walker T, Barnstedt O, Edmondson-Stait AJ, Pleijzier MW, Otto N, Schlegel P, Sharifi N, Perisse E, Smith CS, Lauritzen JS, Costa M, Jefferis G, Bock DD, Waddell S (2018) Integration of Parallel Opposing Memories Underlies Memory Extinction. Cell 175:709-722 e715.

Ferdenzi C, Poncelet J, Rouby C, Bensafi M (2014) Repeated exposure to odors induces affective habituation of perception and sniffing. Front Behav Neurosci 8:119.

Ganguly-Fitzgerald I, Donlea J, Shaw PJ (2006) Waking experience affects sleep need in Drosophila. Science 313:1775-1781.

Gervasi N, Tchenio P, Preat T (2010) PKA dynamics in a Drosophila learning center: coincidence detection by rutabaga adenylyl cyclase and spatial regulation by dunce phosphodiesterase. Neuron 65:516-529.

Giese KP, Mizuno K (2013) The roles of protein kinases in learning and memory. Learn Mem 20:540-552.

Goel N, Rao H, Durmer JS, Dinges DF (2009) Neurocognitive consequences of sleep deprivation. Semin Neurol 29:320-339.

Griffith LC, Verselis LM, Aitken KM, Kyriacou CP, Danho W, Greenspan RJ (1993) Inhibition of calcium/calmodulin-dependent protein kinase in Drosophila disrupts behavioral plasticity. Neuron 10:501-509.

Handler A, Graham TGW, Cohn R, Morantte I, Siliciano AF, Zeng J, Li Y, Ruta V (2019) Distinct Dopamine Receptor Pathways Underlie the Temporal Sensitivity of Associative Learning. Cell 178:60-75 e19.

Hendricks JC, Finn SM, Panckeri KA, Chavkin J, Williams JA, Sehgal A, Pack AI (2000) Rest in Drosophila is a sleep-like state. Neuron 25:129-138.

Hige T, Aso Y, Modi MN, Rubin GM, Turner GC (2015) Heterosynaptic Plasticity Underlies Aversive Olfactory Learning in Drosophila. Neuron 88:985-998.

Honegger KS, Campbell RA, Turner GC (2011) Cellular-resolution population imaging reveals robust sparse coding in the Drosophila mushroom body. J Neurosci 31:11772-11785.

Inada K, Tsuchimoto Y, Kazama H (2017) Origins of Cell-Type-Specific Olfactory Processing in the Drosophila Mushroom Body Circuit. Neuron 95:357-367 e354.

Kelleher RJ, 3rd, Govindarajan A, Tonegawa S (2004) Translational regulatory mechanisms in persistent forms of synaptic plasticity. Neuron 44:59-73.

Kim YC, Lee HG, Han KA (2007) D1 dopamine receptor dDA1 is required in the mushroom body neurons for aversive and appetitive learning in Drosophila. J Neurosci 27:7640-7647.

Koh YH, Popova E, Thomas U, Griffith LC, Budnik V (1999) Regulation of DLG localization at synapses by CaMKII-dependent phosphorylation. Cell 98:353-363.

Krashes MJ, Keene AC, Leung B, Armstrong JD, Waddell S (2007) Sequential use of mushroom body neuron subsets during drosophila odor memory processing. Neuron 53:103-115.

Kuklin EA, Alkins S, Bakthavachalu B, Genco MC, Sudhakaran I, Raghavan KV, Ramaswami M, Griffith LC (2017) The Long 3'UTR mRNA of CaMKII Is Essential

for Translation-Dependent Plasticity of Spontaneous Release in Drosophila melanogaster. J Neurosci 37:10554-10566.

Levin LR, Han PL, Hwang PM, Feinstein PG, Davis RL, Reed RR (1992) The Drosophila learning and memory gene rutabaga encodes a Ca2+/Calmodulin-responsive adenylyl cyclase. Cell 68:479-489.

Li F et al. (2020) The connectome of the adult Drosophila mushroom body provides insights into function. Elife 9.

Li X, Yu F, Guo A (2009) Sleep deprivation specifically impairs short-term olfactory memory in Drosophila. Sleep 32:1417-1424.

Lin AC, Bygrave AM, de Calignon A, Lee T, Miesenbock G (2014) Sparse, decorrelated odor coding in the mushroom body enhances learned odor discrimination. Nat Neurosci 17:559-568.

Livingstone MS, Sziber PP, Quinn WG (1984) Loss of calcium/calmodulin responsiveness in adenylate cyclase of rutabaga, a Drosophila learning mutant. Cell 37:205-215.

Malik BR, Gillespie JM, Hodge JJ (2013) CASK and CaMKII function in the mushroom body alpha'/beta' neurons during Drosophila memory formation. Front Neural Circuits 7:52.

Mao Z, Davis RL (2009) Eight different types of dopaminergic neurons innervate the Drosophila mushroom body neuropil: anatomical and physiological heterogeneity. Front Neural Circuits 3:5.

Mao Z, Roman G, Zong L, Davis RL (2004) Pharmacogenetic rescue in time and space of the rutabaga memory impairment by using Gene-Switch. Proc Natl Acad Sci U S A 101:198-203.

Marin EC, Jefferis GS, Komiyama T, Zhu H, Luo L (2002) Representation of the glomerular olfactory map in the Drosophila brain. Cell 109:243-255.

Masek P, Worden K, Aso Y, Rubin GM, Keene AC (2015) A dopamine-modulated neural circuit regulating aversive taste memory in Drosophila. Curr Biol 25:1535-1541.

McClelland JL, McNaughton BL, O'Reilly RC (1995) Why there are complementary learning systems in the hippocampus and neocortex: insights from the successes and failures of connectionist models of learning and memory. Psychol Rev 102:419-457.

Miller S, Yasuda M, Coats JK, Jones Y, Martone ME, Mayford M (2002) Disruption of dendritic translation of CaMKIIalpha impairs stabilization of synaptic plasticity and memory consolidation. Neuron 36:507-519.

Mitchell J, Smith CS, Titlow J, Otto N, van Velde P, Booth M, Davis I, Waddell S (2021) Selective dendritic localization of mRNA in Drosophila mushroom body output neurons. Elife 10.

Murthy M, Fiete I, Laurent G (2008) Testing odor response stereotypy in the Drosophila mushroom body. Neuron 59:1009-1023.

Olsen SR, Wilson RI (2008) Cracking neural circuits in a tiny brain: new approaches for understanding the neural circuitry of Drosophila. Trends Neurosci 31:512-520.

Owald D, Waddell S (2015) Olfactory learning skews mushroom body output pathways to steer behavioral choice in Drosophila. Curr Opin Neurobiol 35:178-184.

Owald D, Felsenberg J, Talbot CB, Das G, Perisse E, Huetteroth W, Waddell S (2015) Activity of defined mushroom body output neurons underlies learned olfactory behavior in Drosophila. Neuron 86:417-427.

Pech U, Pooryasin A, Birman S, Fiala A (2013) Localization of the contacts between Kenyon cells and aminergic neurons in the Drosophila melanogaster brain using SplitGFP reconstitution. J Comp Neurol 521:3992-4026.

Pellegrino R, Sinding C, de Wijk RA, Hummel T (2017) Habituation and adaptation to odors in humans. Physiol Behav 177:13-19.

Perez-Orive J, Mazor O, Turner GC, Cassenaer S, Wilson RI, Laurent G (2002) Oscillations and sparsening of odor representations in the mushroom body. Science 297:359-365.

Riemensperger T, Voller T, Stock P, Buchner E, Fiala A (2005) Punishment prediction by dopaminergic neurons in Drosophila. Curr Biol 15:1953-1960.

Schroll C, Riemensperger T, Bucher D, Ehmer J, Voller T, Erbguth K, Gerber B, Hendel T, Nagel G, Buchner E, Fiala A (2006) Light-induced activation of distinct modulatory neurons triggers appetitive or aversive learning in Drosophila larvae. Curr Biol 16:1741-1747.

Schwaerzel M, Monastirioti M, Scholz H, Friggi-Grelin F, Birman S, Heisenberg M (2003) Dopamine and octopamine differentiate between aversive and appetitive olfactory memories in Drosophila. J Neurosci 23:10495-10502.

Seidner G, Robinson JE, Wu M, Worden K, Masek P, Roberts SW, Keene AC, Joiner WJ (2015) Identification of Neurons with a Privileged Role in Sleep Homeostasis in Drosophila melanogaster. Curr Biol 25:2928-2938.

Sejourne J, Placais PY, Aso Y, Siwanowicz I, Trannoy S, Thoma V, Tedjakumala SR, Rubin GM, Tchenio P, Ito K, Isabel G, Tanimoto H, Preat T (2011) Mushroom body efferent neurons responsible for aversive olfactory memory retrieval in Drosophila. Nat Neurosci 14:903-910.

Semelidou O, Acevedo SF, Skoulakis EM (2018) Temporally specific engagement of distinct neuronal circuits regulating olfactory habituation in Drosophila. Elife 7.

Seugnet L, Suzuki Y, Vine L, Gottschalk L, Shaw PJ (2008) D1 receptor activation in the mushroom bodies rescues sleep-loss-induced learning impairments in Drosophila. Curr Biol 18:1110-1117.

Seugnet L, Suzuki Y, Thimgan M, Donlea J, Gimbel SI, Gottschalk L, Duntley SP, Shaw PJ (2009) Identifying sleep regulatory genes using a Drosophila model of insomnia. J Neurosci 29:7148-7157.

Shaw PJ, Cirelli C, Greenspan RJ, Tononi G (2000) Correlates of sleep and waking in Drosophila melanogaster. Science 287:1834-1837.

Simpson JH (2009) Mapping and manipulating neural circuits in the fly brain. Adv Genet 65:79-143.

Stewart BA, Atwood HL, Renger JJ, Wang J, Wu CF (1994) Improved stability of Drosophila larval neuromuscular preparations in haemolymph-like physiological solutions. J Comp Physiol A 175:179-191.

Sun F, Zhou J, Dai B, Qian T, Zeng J, Li X, Zhuo Y, Zhang Y, Wang Y, Qian C, Tan K, Feng J, Dong H, Lin D, Cui G, Li Y (2020) Next-generation GRAB sensors for monitoring dopaminergic activity in vivo. Nature methods 17:1156-1166.

Suzuki-Sawano E, Ueno K, Naganos S, Sawano Y, Horiuchi J, Saitoe M (2017) A Drosophila ex vivo model of olfactory appetitive learning. Sci Rep 7:17725.

Tanaka NK, Tanimoto H, Ito K (2008) Neuronal assemblies of the Drosophila mushroom body. J Comp Neurol 508:711-755.

Tomchik SM, Davis RL (2009) Dynamics of learning-related cAMP signaling and stimulus integration in the Drosophila olfactory pathway. Neuron 64:510-521.

Tully T, Quinn WG (1985) Classical conditioning and retention in normal and mutant Drosophila melanogaster. J Comp Physiol A 157:263-277.

Turner GC, Bazhenov M, Laurent G (2008) Olfactory representations by Drosophila mushroom body neurons. J Neurophysiol 99:734-746.

Ueno K, Naganos S, Hirano Y, Horiuchi J, Saitoe M (2013) Long-term enhancement of synaptic transmission between antennal lobe and mushroom body in cultured Drosophila brain. J Physiol 591:287-302.

Ueno K, Suzuki E, Naganos S, Ofusa K, Horiuchi J, Saitoe M (2017) Coincident postsynaptic activity gates presynaptic dopamine release to induce plasticity in Drosophila mushroom bodies. Elife 6.

Ueno K, Morstein J, Ofusa K, Naganos S, Suzuki-Sawano E, Minegishi S, Rezgui SP, Kitagishi H, Michel BW, Chang CJ, Horiuchi J, Saitoe M (2020) Carbon Monoxide, a Retrograde Messenger Generated in Postsynaptic Mushroom Body Neurons, Evokes Noncanonical Dopamine Release. J Neurosci 40:3533-3548.

Wang Y, Mamiya A, Chiang AS, Zhong Y (2008) Imaging of an early memory trace in the Drosophila mushroom body. J Neurosci 28:4368-4376.

Wilson DA (1998) Habituation of odor responses in the rat anterior piriform cortex. J Neurophysiol 79:1425-1440.

Wilson DA (2009) Olfaction as a model system for the neurobiology of mammalian short-term habituation. Neurobiol Learn Mem 92:199-205.

Wilson DA, Linster C (2008) Neurobiology of a simple memory. J Neurophysiol 100:2-7.

Wong AM, Wang JW, Axel R (2002) Spatial representation of the glomerular map in the Drosophila protocerebrum. Cell 109:229-241.

Yamagata N, Hiroi M, Kondo S, Abe A, Tanimoto H (2016) Suppression of Dopamine Neurons Mediates Reward. PLoS Biol 14:e1002586.

Yao Z, Macara AM, Lelito KR, Minosyan TY, Shafer OT (2012) Analysis of functional neuronal connectivity in the Drosophila brain. J Neurophysiol 108:684-696.

Yu D, Akalal DB, Davis RL (2006) Drosophila alpha/beta mushroom body neurons form a branch-specific, long-term cellular memory trace after spaced olfactory conditioning. Neuron 52:845-855.

Zhang X, Noyes NC, Zeng J, Li Y, Davis RL (2019) Aversive Training Induces Both Presynaptic and Postsynaptic Suppression in Drosophila. J Neurosci 39:9164-9172.

Zhao X, Lenek D, Dag U, Dickson BJ, Keleman K (2018) Persistent activity in a recurrent circuit underlies courtship memory in Drosophila. Elife 7.

Figure Legends

Figure 1. Simultaneous activation of the odor and the electric shock pathways induces an enhancement PDP in the KCs in the MB α'3 compartment. *A*,

Schematic of the dissected adult fly's central nervous system showing the MBs and the

ALs in the central brain and the ventral nerve cord. The ventral nerve cord (VNC) is then

cut at the base of the cervical connective to free the afferent fibers of the VNC (AFV). **B,**

Schematic of the imaging chamber showing the placement of the first electrode on the

AL, and the second electrode on the free end of the AFV. **C,** Representative image of

the MB ipsilateral to the stimulated AL. The calcium indicator GCaMP6f is expressed in

the KCs (driven by *VT030559-GAL4*). The circles show the analyzed ROIs surrounding

the α'3 and the α3 compartments. Scale bar is 25 μm. **D,** Paired and unpaired induction

protocols. Top, in the paired induction, the ipsilateral AL is activated by 5 stimulation

trains with 15 seconds intertrain interval followed by 1 minute rest (pre-induction). 12

trains of stimulations are then delivered to both the AL and the AFV simultaneously

(induction). The brain is then rested for 15 minutes before being tested by 5 trains of AL

stimulations like those applied during the pre-induction (post-induction). Bottom, the

unpaired induction: same as paired induction except that the AL and the AFV

stimulations are separated by 30 seconds during the induction stage. The stimulation

train is 20 pulses at 100 Hz; each pulse is 1 millisecond with 9 milliseconds inter-pulse

interval. AL stimulation strength: 100 μAmps. AFV stimulation strength: 0.5-1 mAmps.

E, Example of the KC calcium response in the α'3 compartment during pre (grey) and

post (black) stimulations. **F,** Mean relative change of the calcium responses calculated

as $(mean\ post\ responses - mean\ pre\ responses)/(mean\ pre\ responses)$ in the α'3 (left)

and the α3 (right) compartments after no induction (white), AL activation alone (light

grey), AFV activation alone (dark grey), unpaired AL+AFV induction (green), and paired

AL+AFV induction (blue). Mean ±SEM. Two-way ANOVA with Šídák post-hoc tests; α =

0.05; n = 8 in each condition. In the α'3 compartment: A vs. A, $p > 0.05$; B vs. A, $p \leq$

0.05 (0.015, < 0.0001, 0.0002, and <0.0001 in paired vs. no induction, paired vs. AL

alone, paired vs. AFV alone, and paired vs. unpaired, respectively). No statistical

significance across conditions in the α3 compartment (p-value > 0.05).

Figure 2. Electric shock pathway activation can be replaced by either dopamine

perfusion or activation of dopaminergic neurons. *A,* Schematic of the dissected

adult central nervous system showing the placement of one electrode to activate the

AFV. ***B,*** Representative images showing the simultaneous expression of *UAS-GRAB*

DA2m (left; green) and *UAS-jRCaMP1a* (right; red) in the same neurons in the MB

(driven by *VT030559-GAL4*). Scale bar is 25 μm. ***C,*** Time course of GRAB DA2m

responses in the α'3 and α3 compartments upon AFV stimulation (green) or no stimulus

(grey). Traces show mean ΔF/F ±SEM across the 6 flies tested. In each compartment, a

representative jRCaMP1a response is shown (red) as a reference for neural activity

upon AFV stimulation. ***D,*** Quantification of the GRAB DA2m responses in ***C***. Two-way

ANOVA with Šídák post-hoc tests; ****p<0.0001. ***E,*** Schematic of the AL+DA induction

protocol, showing electrode placement above the AL, and perfusion of dopamine. ***F,***

Paired and unpaired induction protocols. Top, in the paired induction, the ipsilateral AL

is activated by 5 stimulation trains with 15 seconds intertrain interval followed by resting

for 1 minute (pre-induction). 12 trains of stimulations are then applied via the AL

electrode, coincident with 60 seconds of dopamine perfusion. The brain is then rested

for 15 minutes before being tested by 5 trains of AL stimulations like those applied

during the pre-induction (post-induction). Bottom, the unpaired induction: same as

paired induction except that the AL stimulation and dopamine perfusion are separated

by 30 seconds during the induction stage. The stimulation train is 20 pulses at 100 Hz;

each pulse is 1 millisecond with 9 milliseconds inter-pulse interval. AL stimulation

strength: 100 μAmps. **G,** Mean relative change of the calcium responses in the α'3 (left)

and the α3 (right) compartments after DA perfusion alone (grey), unpaired AL+DA

induction (green), and paired AL+DA induction (blue). Three different DA concentrations

were used: 1 μM (left), 5 μM (middle), and 10 μM (right). Mean ±SEM. Two-way ANOVA

with Šídák post-hoc tests; n = 6 in each condition. In the α'3 compartment: A vs. A, p >

0.05; B vs. A, p ≤ 0.05 (<0.0001 when the paired induction is compared to the unpaired

induction in the 5 μM DA and 10 μM DA conditions). No statistical significance across

conditions in the α3 compartment (p-value > 0.05). **H,** Mean relative change of the

calcium responses in the α'3 (left) and the α3 (right) compartments pre induction and 30

or 60 minutes after paired AL+ 10 μM DA induction (blue). Mean ±SEM. Repeated

Measures ANOVA with Tukey post-hoc tests (statistical analysis is applied to the raw

responses before normalization); n = 4 in each condition. In the α'3 compartment: A vs.

A, p > 0.05; B vs. A, p ≤ 0.05 (0.041 and 0.0416 when pre is compared to 30 mins and

to 60 mins after induction, respectively). No statistical significance across conditions in

the α3 compartment (p-value > 0.05). **I,** Schematic of the AL+PPL1s induction protocol,

showing electrode placement above the AL, and perfusion of 2.5 mM ATP to activate

the $P2X_2$ channels expressed in the PPL1 dopaminergic neurons (driven by *TH-LexA*).

J, Paired and unpaired induction protocols. Top, in the paired induction, the ipsilateral

AL is activated by 5 stimulation trains with 15 seconds intertrain interval followed by

resting for 1 minute (pre-induction). We then apply 12 trains of stimulations to the AL

electrode, coincident with 60 seconds of 2.5 mM ATP perfusion. The brain is then

rested for 15 minutes before being tested by 5 trains of AL stimulations like those

applied during the pre-induction (post-induction). Bottom, the unpaired induction: same

as paired induction except that the AL stimulation and ATP perfusion are separated by

30 seconds during the induction stage. The stimulation train is 20 pulses at 100 Hz;

each pulse is 1 millisecond with 9 milliseconds inter-pulse interval. AL stimulation

strength: 100 μAmps. **K,** Mean relative change of the calcium responses in the α'3 (left)

and the α3 (right) compartments after unpaired AL+ATP induction (green), and paired

AL+ATP induction (blue). Mean ±SEM. Two-way ANOVA with Šídák post-hoc tests; n =

8 in each condition. **p = 0.0036. No statistical significance across conditions in the α3

compartment (p-value = 0.786).

Figure 3. PDP is CS-pathway input-specific. A, Schematic of the dissected adult

brain showing the placement of two electrodes in the same AL. Figure 3-1 in extended

data demonstrates that the two electrodes activate different subsets of projection

neurons. **B,** induction protocol. The ipsilateral AL is activated by 3 stimulation trains with

15 seconds intertrain interval followed by resting for 1 minute (pre-induction). 12 trains

of stimulations to electrode 1 (E1) are then delivered, followed by a 30-second rest, then

12 strains of stimulations to electrode 2 (E2). Dopamine [10 μM] is perfused in the

chamber for 60 seconds paired with either E1 or E2 stimulation. The brain is then rested

for 15 minutes before being tested by 3 trains of E2 stimulation followed by 3 trains of

E1 stimulation (post-induction). **C,** Representative trace of a prep in which dopamine

perfusion is paired with E1 stimulations. **D,** Mean relative change of the calcium

responses in the α'3 compartment. Mean ±SEM. Unpaired t-test, two tailed; n = 8. ***p

= 0.0005.

Figure 3-1. Different electrodes in the same AL activate different subsets of

Projection Neurons. A, Schematic showing the placement of electrode 1 (E1) and

electrode 2 (E2) in the same ALs. *B,* Basal calcium signals expressed in the AL

projection neurons (genotype: *w-;GH146-GAL4/+;UAS-GCaMP6f/+*) showing two ROIs

highlighted in blue (ROI near E1) and red (ROI near E2). *C,* calcium responses in each

ROIs to E1 (blue trace) and E2 (red trace) stimulation. *D,* Quantification of *C*. Dashed

line is plotted at ΔF/F = 0.05**.**

Figure 4. α'3 MBONs dendrites show a suppression PDP. *A,* Representative image

of the GCaMP6f signal expressed in the α'3 MBONs using the *MB027B* splitGAL4 line.

The MB is outlined in white, showing the localization of the analyzed dendritic signal in

the α'3 compartment. Scale bar is 25 μm. *B,* Representative trace of the α'3 MBONs

responses pre and post the AL+DA induction described in Figure 2F. *C,* Mean relative

change of the calcium responses in the MBONs' dendrites in the α'3 compartment.

Mean ±SEM. Unpaired t-test, two tailed; n = 6. **p = 0.0058.

Figure 5. The 3'UTR region of *CaMKII* is important for PDP formation. *A,* Schematic

of wildtype (top) and *CaMKII^{Udel}* where the 3'UTR is flipped out (bottom). *B,*

Representative immunostaining images showing CaMKII protein levels in the MB in

wildtype *CaMKII* flies (left) and in *CaMKII^{Udel}* flies (right). Scale bar is 20 μm. *C,*

Quantification of *B*. Mean±SEM. Unpaired t-test, two tailed; n = 20 wildtype flies and 22

CaMKII^{Udel} flies. ****p < 0.0001. *D,* Learning index of wildtype and *CaMKII^{Udel}* flies after

training with 90 volts or 60 volts. Mean ±SEM. Two-way ANOVA with Šídák post-hoc

tests; n = 16, 15, 18, and 12 (wildtype 90v, UDEL 90v, wildtype 60v, and Udel 60v,

respectively). ****p < 0.0001. Figure 5-1 in extended data shows total CaMKII protein

levels in wildtype and *CaMKII^{Udel}* flies and illustrates that both groups are similarly

responsive to electric shock. *E,* Mean relative change of the calcium responses in the

α'3 and the α3 compartments after the unpaired AL+DA induction (green), and paired

AL+DA induction (blue) as described in figure 2F. Mean ±SEM. Two-way ANOVA with

Šídák post-hoc tests; n = 6 in each condition. In the α'3 compartment: A vs. A, p > 0.05;

B vs. A, p ≤ 0.0001; C vs. A, p=0.0048; C vs. B, p=0.0336. In the α3 compartment: A vs.

A, p > 0.05; B vs. A, p ≤0.05. Stripe bars indicate the *CaMKII*^{Udel} genotype.

Figure 5-1. Total CaMKII protein in *CaMKII*^{Udel} flies and their effect on sensitivity to

electric shock. *A,* Western blot showing CaMKII and actin levels in wildtype or

CaMKII^{Udel} adult female brains and in purified CaMKII samples. *B,* Quantification of *A*.

Mean ±SEM. Mann-Whitney test; n = 4; *p=0.0286. *C,* Aversion of wildtype and

CaMKII^{Udel} flies to electric shock. Mean ±SEM. Unpaired t-test; data collected across 11

trials of the wildtype group and 20 trials of the *CaMKII*^{Udel} group; number of flies per

group ranged between 21 to 47 with an average of 29 flies per group; p=0.8518.

Figure 6. Increasing CS or US pathway activation recruits the α3 compartment to

the PDP circuit. *A,* Design of the AL ramp. AL input stimulation is increased by 30

μAmps every 30 seconds. *B,* Unpaired calcium responses to AL stimulations in the α'3

and α3 compartments showing mean±SEM. Unpaired t-test; n=6; *p<0.05. ***C and D,***

Mean relative change of the calcium responses in the α'3 and the α3 compartments

after the unpaired (green) or paired (blue) AL+AFV induction in ***C*** and AL+DA induction

in ***D*** when the AL input stimulation is increased to 200 μAmps. Mean ±SEM. Two-way

ANOVA with Šídák post-hoc tests; n = 8 in each condition. *p<0.05, **P<0.01,

****P<0.0001. *E,* Mean relative change of the calcium responses in the α'3 and the α3

compartments after unpaired 10μM or 200μM DA perfusion alone, stimulation at 100

μAmps. Two-way ANOVA with Šídák post-hoc tests; n = 6 in each condition. *p<0.05,

**P<0.01.

Figure 7. Sleep deprivation blocks PDP and rebound sleep recovers it. *A,*

Schematic of the different sleep groups: Sleep Control, flies are undisturbed during

night and tested between ZT0-ZT4 in the next day; Sleep Deprived, flies are kept on the

shaker during between ZT12-ZT24 then tested between ZT0-ZT2 in the next day; Sleep

Deprived + Recovery, same as Sleep Deprived but flies were allowed to recover sleep

between ZT0-ZT2 then tested between ZT2-ZT4; Stress Control: flies are undisturbed

during the night between ZT12-ZT22 and then the shaker was turned on between ZT22-

ZT24 before testing between ZT0-ZT2 in the next day. Testing was done by the AL+DA

induction described in Figure 2. *B,* Total sleep of the different groups at the different ZT

windows. Mean ±SEM. One-way ANOVA with Tukey's multiple comparisons. In the

ZT12:22 comparison, B vs. A, p<0.0001; A vs A, p=0.4916. In the ZT22:24 comparison,

B vs. A, p<0.0001; B vs B, p=0.7136. In the ZT0-2 window, B vs. A, p<0.0001. *C,* Mean

relative change of the calcium responses in the α'3 (left) and the α3 (right)

compartments after paired AL+DA induction. Mean ±SEM. Two-way ANOVA with Šídák

post-hoc tests; n = 6 in each condition. In the α'3 compartment: A vs. A, p > 0.05; B vs.

A, p ≤ 0.0001. No statistical significance across conditions in the α3 compartment (p>

0.05). *D,* PDP (mean relative change of dF/F) from individual animals plotted against

the animal's time of dissection. White circles indicate control animals (allowed to sleep

between ZT12-ZT24) and pink circles indicate sleep deprived animals (on the shaker

between ZT12-ZT24).

A

B   imaging chamber

perfusion inlet

suction

C   α'3   α3

KC>GCaMP6f   25µm

D
         pre   paired
                  induction   post
AL   ┤┤┤┤┤ ┤┤┤┤┤┤┤┤┤┤   ┤┤┤┤┤
                              15 mins
AFV ───── ┤┤┤┤┤┤┤┤┤┤ incubation ─────

         pre   unpaired induction   post
AL   ┤┤┤┤┤ ┤┤┤┤┤┤   ┤┤┤┤┤┤┤┤┤┤   ┤┤┤┤┤
                                        15 mins
AFV ───── ┤┤┤┤┤┤┤┤┤┤ ─────── incubation ─────

E   a'3 paired
           pre
           post

     0.2ΔF/F
        0.5 sec

F

no induction
AL (100µA) alone
AFV alone
unpaired AL (100µA) + AFV
paired AL (100µA) + AFV

A

E1　　　E2

DA

B

AL (E1)　　pre　paired
　　　　　　　induction　　　　　　　　post

　　　　　　　　　　　　　　　　15 mins
AL (E2)　　　　　　　　　　　　incubation

DA　　OR
[10μM]

C

representative trace (E1 paired)

ΔF/F

pre　　　　　post

response　response　response　response
to E1　　to E2　　to E2　　to E1

15 mins

1 min

D

ΔF/F Relative Change

***

unpaired　paired

A

MB027B > 20x GCaMP6f

25μm

B

pre

post

0.2 ΔF/F

1 sec

C

**

0.5

0.0

−0.5

−1.0

ΔF/F Relative Change

unpaired AL (100μA)
+ DA (10μM)

paired AL (100μA)
+ DA (10μM)

A

wildtype CaMKII

5' ▭ ▭ ATG ▮▮▮▮ 3'UTR 3'

CaMKII^UDel

5' ▭ ▭ ATG ▮▮▮▮ 3P3-RFP 3'

B

20 uM   20 uM

C

****

Mean Intensity (A.U.)

wildtype   *CaMKII^Udel*

D

****   ********

Learning Index

wildtype 90v   Udel 90v   wildtype 60v   Udel 60v

E

A   B   A   C   A   B   A   A

ΔF/F Relative Change

wildtype   *CaMKII^Udel*   wildtype   *CaMKII^Udel*

α'3   α3

☐ paired AL (200μA) + DA (10μM)

☐ unpaired AL (200μA) + DA (10μM)

A

	ZT0	Day 1 ZT22	ZT24/0	Day 2 ZT2	ZT4

LIGHTS OFF   LIGHTS ON

Sleep Control — Undisturbed — Test

Sleep Deprived — Shaker on — Test

Sleep Deprived + Recovery — Shaker on — Recovery — Test

Stress Control — Shaker off — Shaker on — Test

B

Day 1 ZT12-22   Day 1 ZT22-24   Day 2 ZT0-2

Total Sleep (mins)

A B A   A B B   A B

□ control
■ SD
▨ stress control
⬤ SD + recovery sleep

C

ΔF/F Relative Change

A B A A   ns

α'3   α3

D

PDP

time of dissection (minutes after ZT0)

30 60 90 120 150 180