Effects of substrate on shrimp growth, water quality and bacterial community in

the biofloc system nursing *Penaeus vannamei* postlarvae

Hai-Hong Huang [a*], Chao-Yun Li [b†], Yan-Ju Lei [a,c], Wei-Qi Kuang [a], Bo-Lan Zhou[a],

Wan-Sheng Zou [a,d], Pin-Hong Yang [a,e]

[a] *College of life and environmental sciences, Hunan University of Arts and Science,*

Changde Hunan 415000, China

[b] *Shandong Vocational Animal Science and Veterinary College*, *Weifang Shandong*

261061, *China*

[c] *Hunan Provincial Collaborative Innovation Center for Efficient and Health*

Production of Fisheries, Changde Hunan 415000, China

[d] *Hunan Provincial Key Laboratory for Health Aquaculture and Product Processing in*

Dongting Lake Area, Changde Hunan 415000, China

[e] *Hunan Engineering Research Center of Aquatic Organism Resources and*

Environmental Ecology, Changde Hunan 415000, China

Abstract

This study aimed to investigate the effects of substrate on water quality, shrimp growth

and bacterial community in the biofloc system with a salinity of 5‰. Two treatments,

biofloc system with (sB) or without (nB) addition of elastic solid packing filler (nylon)

as substrate, were set up. *Penaeus vannamei* postlarvae (PL, ~ stage 15) were stocked at

* Corresponding author. *E-mail address*: shinkanh@nwsuaf.edu.cn, shinkanh@huas.edu.cn.

† Co-first author.

a density of 4000 PL m^{-3} in each treatment with triplicates for a 28-days culture

experiment, taking glucose as carbon source (C:N 15:1). Results showed that the

survival rate (96.3±3.6%), FCR (0.76±0.06) and productivity (1.54±0.12 kg m^{-3}) in sB

treatment were significantly better than those in nB treatment (81.0±7.1%, 0.98±0.08

and 1.14±0.09 kg m^{-3}, $P<0.05$). All water parameters were in the recommended ranges.

Substrate showed significant effect on TAN, TSS, turbidity, biofloc volume, pH and

carbonate alkalinity ($P < 0.05$). Actinobacteria (4.0-22.7%), Bacteroidetes

(10.4-33.5%), Firmicutes (0.2-11.2%), Planctomycetes (4.0-14.9%) and Proteobacteria

(29.4-59.0%) were the most dominant phyla for both treatments. However, the bacterial

community in sB treatment showed to be significantly different from that in nB

treatment (Jaccard distance 0.94±0.01, $P=0.001$). Substrate showed significant effects

on Shannon, Heip, Pielou and Simpson index, as well as relative abundances of

Actinobacteria, Bacteroidetes and Planctomycetes ($P < 0.05$). The results suggested

that addition of substrate affected the shrimp growth, water quality and bacterial

community in the biofloc system nursing *P. vannamei* PL with a 5‰ salinity.

KEYWORDS

Penaeus vannamei; Biofloc technology; Low salinity; Substrate; Bacterial community

1 INTRODUCTION

Crustacean is the most valuable group in fish trade in the world, accounting for 23.0%

in terms of value with an 8.3% share in culture production, compared to the 65.4% Vs.

79.8% of fish, and the 11.6% Vs. 11.9% of mollusks and other aquatic invertebrates

(FAO, 2018). Among the cultured crustaceans, *Penaeus vannamei* (Boone, 1931) is the

most important species whose production accounted for 53.0% of the total crustaceans

production (FAO, 2018).

Prenursery of postlarvae (PL) is a very important step during culture of *P.*

vannamei in many places (Mishra et al., 2008; Rezende et al., 2018). In the

conventional prenursery system, PL are intensively stocked at a high density (up to 60

PL L^{-1}) with small-scale water volume and carefully nursed for 15-60 days (Schveitzer

et al., 2017), to save water, land and energy resources, and to deliver large individuals

that are resistant to environmental conditions to increase the survival and growth

performance during the grow-out phase (Mishra et al., 2008; Rezende et al., 2018).

However, in this high-stocking-density system, serious cannibalism behavior,

competition for space and degradation of water quality would affect growth and

survival of PL (Esparza-Leal et al., 2015). The use of artificial substrate is an effective

approach to supply additional surface and refugee for shrimp during prenursery

(Tierney et al., 2020). Additionally, biofloc technology (BFT) has also been tried to be

used to prenursery of *P. vannamei* PL (Khanjani et al., 2017; Schveitzer et al., 2017), to

control the problems usually appeared in the traditional prenursery system, such as

biosecurity and toxic ammonia and nitrite (Samocha, 2010), due to the advantages of

this technology on nitrogen assimilation *in situ* and pathogen control under minimal or

zero water exchange conditions (Avnimelech, 2015; Huang, 2020). Furthermore,

several studies have used substrates in biofloc systems to reinforce the controlling of

water quality and the growth performance of PL by supplying a natural food through

the periphyton growing on the substrate during the prenursery stage (Peiro-Alcantar et

al., 2019; Rezende et al., 2018; Tierney et al., 2020).

P. vannamei is an euryhaline species, and can be cultured at low salinity conditions

less than 1.0‰, which is a trend that will continue to grow globally (Roy et al., 2010).

Osmoregulation capability of *P. vannamei* develops gradually in the postlarvae stages

and can be acclimated to salinities as low as 0.5‰ by stage 12 (Van Wyk et al., 1999),

indicating that PL could be nursed under low salinity conditions. As well, biofloc

technology has been successfully used for prenursery of *P. vannamei* PL under salinity

conditions of 8-16‰ (Esparza-Leal et al., 2016; Luo et al., 2019). However, application

of substrate in prenursery of *P. vannamei* PL in the biofloc system under a low-salinity

condition has been not documented yet.

In this study, it is aimed to estimate the effects of substrate on shrimp growth

performance and water quality in the biofloc system nursing *P. vannamei* postlarvae

(PL) under a salinity condition as low as about 5‰. Furthermore, the effects of

substrate on the bacterial community diversity and composition in this system were

also investigated.

2   MATERIAL AND METHODS

2.1   Ethic statement

The culture experiments were carried out at a local farm of Bifuteng eco-agriculture

development Co., Ltd. (BEAD Co., Ltd., Changde, China, Lat. 28°53'57.88" N, Long.

111°38'3.08" E), with indoor tanks, under principles in good laboratory animal care,

according to the national standard of China (GB/T 35892-2018), 'Laboratory

animal-Guideline for ethical review of animal welfare'. The manuscript does not

require ethical approval.

2.2   Experimental design

There two treatments with three replicates for each were set up in the present study,

biofloc system with (sB) and without (nB) addition of substrate. In the sB treatment,

52.4% of the tank internal surface was equally coated with elastic solid packing filler

(nylon, with a diameter of 15 cm).

Before the beginning of the experiment, tanks (width $\times$ length $\times$ depth $= 2 \times 2.5$

$\times$ 1.3 m, 6.5 m^3 in volume) connected with a Roots blower air pump (CRS-100,

Changsheng mechanical and electrical equipment Co., Ltd., Zhangqiu, Shandong,

China) were filled 5.0 m^3 (1.0 m in depth) culture water with a salinity of about 5.0‰

per tank. The water for culture experiment was prepared according to the method of

Ray and Lotz (2017), with some modifications. Briefly, artificial sea salt powder

(Qianglong corporation, Tianjin, China), and food-grade chemical reagents of KCl,

$MgCl_2$, $CaCl_2$ and $NaHCO_3$ were added to tap water, to manage an initial salinity of

5.02±0.08‰, with K^+, Mg^{2+} and Ca^{2+} concentrations of 301.3±5.9, 925.2±12.1 and

306.7±6.9 mg L^{-1}, respectively, and a pH value of about 8.0. After that, sterilization

with 10.0 mg L^{-1} of chlorinedioxide followed by neutralization with 1.0 mg L^{-1} of

ascorbic acid was executed according to the processes of previous studies (Gaona et al.,

2017; Lara et al., 2017).

At the beginning of the experiment, *P. vannamei* PL was randomly assigned to each

tank at a density of 4000 PL per m^3. Those PL (~PL15, 2.5±0.4 mg) were kindly

supplied from BEAD Co., Ltd. and had been treated with a desalinating and

acclimation procedure to adapt the experimental conditions before being assigned (Luo

et al., 2019; Van Wyk et al., 1999). Next, during the 28-days experimental period, PL

were fed with a commercial formulated shrimp diet (crude protein 40.0%, crude lipid

5.0%, crude fibber 5.0%, crude ash 15.0%, moisture 12.0%, Alpha corporation,

Jiangmen, Guangdong, China), with a frequency of four times (6:00, 12:00, 18:00,

24:00) a day, basing on the feeding table and the estimated total biomass according to

methods of Van Wyk et al. (1999). Besides, glucose (food grade, carbohydrate content

90.0%, Fufeng biotechnology Co., Ltd., Hohhot, Inner Mongolia Autonomous, China)

was added as exogenous carbon source to the experimental tanks, according to an

inputted carbon to nitrogen ratio (C:N) of 15:1 (Huang et al., 2021). The inputted C:N

was the C:N contained in the inputted materials (feed and carbon source), and

determined basing on the carbon contents contained in carbon source and feed (Kumar

et al., 2017, Tinh et al. 2021) and the conception that 25% of the feed nitrogen is

theoretically converted as shrimp biomass (Piedrahita, 2003). Throughout the whole

experimental period, no water exchange was operated, but evaporating loss was

complemented with dechlorinated tap water per week.

During 14-28 d of the experiment, partial biofloc was removed by using a

side-stream settling chamber equipped a submersible pump (JGP-2500L, SENSEN

Group Co., Ltd., Zhoushan, China) which ran for 6 h every day at a rate of 2.25 m^3 per

h to maintain a biofloc volume < 15 mL L^{-1}.

2.3 Determination of water parameters

Water temperature, dissolved oxygen and pH were detected each day by using electric

analyzers (YSI-ProPlus, Yellow Springs Instruments Inc., OH, USA; pH-100 meter,

LICHEN Sci-Tech, Co., Ltd., Shanghai, China). Water samples were collected once a

week and passed thought a 0.22-μm pore size microfilter (Xinya purification equipment

Co., Ltd, Shanghai, China). Then, the dissolved total nitrogen in water (DTN), total

ammonia nitrogen (TAN), nitrite, nitrate, carbonate alkalinity, and total suspended

solid (TSS), were determined according to the standard methods (APHA, 1995).

Additionally, the whole total nitrogen (WTN) in water body of each treatment was also

determined every week with water sample without filtration (APHA, 1995). And the

total nitrogen contained in biofloc (BTN) was defined as the total nitrogen content

contained in biofloc in per liter water and was determined by the difference between

WTN and DTN. With an Imhoff cone, 1-liter water sample was settled for 15 min after

which the sediment volume was read as the biofloc volume (settleable solids, BFV)

(Avnimelech, 2015), and the supernatant was used to measure the turbidity (APHA,

1995).

2.4 Growth performance measurements

Thirty PL were selected randomly and individually weighed to the nearest 0.1 mg with

an electric balance (AUX220, Shimadzu, Japan) each week. Survival rate, weekly

increment of body weight (wiW), specific growth rate (SGR), feed conversion rate

(FCR) and productivity were calculated according the following formulates:

$$\text{Survival rate (\%)} = [(\text{survival shrimp counts})/\text{stocked shrimp counts}] \times 100$$

$$\text{wiW (g week}^{-1}) = (\text{fbw} - \text{ibw})/\text{culture weeks}$$

$$\text{SGR (\% d}^{-1}) = [(\ln \text{fbw} - \ln \text{ibw})/\text{culture days}] \times 100$$

$$\text{FCR} = \text{total mass of feed given}/\text{gained shrimp biomass}$$

$$\text{Productivity (kg m}^{-3}) = \text{harvested shrimp biomass}/\text{culture water volume}$$

Wherein, fbw and ibw represented the final and initial mean body weight of PL,

respectively.

2.5   High-throughput sequencing of 16S rRNA gene and data processing

Along the experimental period, water was sampled every week. Briefly, 50 ml water

was collected from each tank of a treatment according to the method of

Martínez-Córdova et al. (2018), pooled as one sample and filtered through a 0.22-μm

pore size membrane. Then, the membrane was stored at -80 $^{\circ}$C for extraction of

bacterial DNA genome with an E.Z.N.A$^{\text{TM}}$ Mag-Bind Soil DNA Kit (OMEGA

Bio-Tek, Inc., GA, USA), according to the manufacturers' instructions.

The V3-V4 region of 16S rRNA gene of bacterial DNA genome was amplified by

using the universal primers, 341F: 5'-CCTACGGGNGGCWGCAG-3' and 805R:

5'-GACTACHVGGGTATCTAATCC-3', in a 30-μl mixtures containing microbial

DNA (10 ng/µl) 2 µl, forward primer (10 µM) 1 µl, reverse primer (10 µM) 1 µl, and 2X

KAPA HiFi Hot Start Ready Mix 15 µl (TaKaRa Bio Inc., Japan), via a procedure with

a thermal instrument (Applied Biosystems 9700, USA): 1 cycle of denaturing at 95°C

for 3 min, first 5 cycles of denaturing at 95°C for 30 s, annealing at 45°C for 30 s,

elongation at 72°C for 30 s, then 20 cycles of denaturing at 95°C for 30 s, annealing at

55°C for 30 s, elongation at 72°C for 30 s and a final extension at 72°C for 5 min. The

PCR product was purified and recovered with MagicPure Size Selection DNA Beads

(TransGen Biotech Co., Ltd., Beijing, China). And then, the purified PCR product was

quantified and normalized with a Qubit ssDNA Assay Kit (Life Technologies, USA), to

construct the 16S gene library for high-throughput sequencing on the Miseq platform

according to the standard protocol (Illumina, USA) by Sangon Biotech (Shanghai) Co.,

Ltd. The raw data produced from high-throughput sequencing has been deposited at

NCBI with accession number of PRJNA646765.

Data processing for the high-throughput sequencing data was carried out under the

QIIME 2 (Quantitative Insights Into Microbial Ecology, Version 2019.10) framework

(Bolyen et al., 2019). In brief, ambiguous nucleotides, adapter sequences and primers

contained in reads, and short reads with length less than 30 bp were removed with the

cutadapt plugin (Martin, 2011). After that, bases in the two ends of reads with quality

score lower than 25 were trimmed. Thereafter, chimeras were filtered, and pair-ended

reads were joined, dereplicated, to obtain high-quality reads which were clustered to

operational taxonomic units (OTUs) with an identity of 0.97 by using the Vsearch tool

(Rognes et al., 2016). Thereafter, the counts of OTUs were normalized by 16S rRNA

gene copy number based on rrnDB database (version 5.6) with the QIIME 2 plugin of

q2-gcn-norm.

2.6   Bacterial diversity analysis

Bacterial community diversity was analyzed with Plugin diversity of QIIME 2 basing

on the normalized counts of OTUs. Metrics regard to alpha diversity, such as Shannon

index, richness indexes of Margalef, dominance indexes of Berger-Parker and Simpson,

and evenness indexes of Heip and Pielou, were computed with the diversity plugin of

QIIME 2. Besides, beta Jaccard index (distance) were also determined with this plugin,

and then significantly tested by using PERMANOVA with 999 permutations

(Anderson, 2001). Basing on the OTUs composition of samples, each sample was

plotted into a PCA diagram with STAMP (Parks et al., 2014).

2.7   Bacterial composition analysis

OTUs were annotated by referring to the GreenGene database 13.8, collapsed at

phylum, class, order, family and genus levels, respectively, with a QIIME 2 Plugin of

taxa. At each level, relative frequencies of species were calculated with the normalized

counts of OTUs. Differential analysis for phylum proportion between both treatments

was conducted with STAMP (Parks et al., 2014) and LEfSe (Segata et al., 2011).

2.8   Statistical analyses

Data was statistically analyzed with the SPSS platform for windows (version 22.0, IBM

Co., NY, USA). Student's T Test were executed for comparison of growth parameters

between both treatment, as soon as normality distribution and homogeneity of data

were proved with Shapiro-Wilk's test and Levene's test, respectively. Or else,

non-parametric Mann-Whitney U test was conducted. Percentage data was submitted to

arcsine transformation. Furthermore, during each stage, ANOVA with repeated

measurements was used to test the significance of main effects and interactions of

substrate and time on water parameters, bacterial diversity indices and bacterial

composition data under Mauchly's test of Sphericity assuming condition, or else, the P

value was corrected by using Greenhouse-Geisser test. Differences were considered

significant at $P < 0.05$.

3 RESULTS

3.1 Water quality

The levels of dissolved oxygen and temperature in the present study were above 5.0 mg

L^{-1} and 26.0 $^{\circ}$C, respectively (Table 1). The effects of time on both parameters were

significant ($P < 0.05$, Table 1). Whereas, substrate showed no significant effect on the

levels of dissolved oxygen and temperature ($P > 0.05$, Table 1).

The carbonate alkalinity in sB treatment was maintained at a high level of

306.5±51.7 mg L^{-1} CaCO$_3$, although it was lower than that in nB treatment (373.7±12.4

mg L^{-1} CaCO$_3$, Table 1). The pH values in nB treatment and sB treatment were

7.20±0.14 and 7.11±0.03, respectively (Table 1). Both parameters were significantly

affected by substrate and time, as well as their interactions ($P < 0.05$, Table 1).

The mean concentrations of the three inorganic nitrogen compounds were at low

levels in the present study (< 1.0 mg L^{-1}, Table 1). Those compounds were not

significantly affected by substrate, time and their interaction ($P > 0.05$), with

exceptions of the substrate main effect and the interaction of substrate and time on TAN

($P < 0.05$, Table 1).

No significant substrate effect on WTN (whole total nitrogen in waterbody), BTN

(total nitrogen contained in biofloc in per liter water) and DTN (dissolved total nitrogen

in waterbody) was found ($P > 0.05$, Table 1). Whereas, the effects of time on those

parameters were significant, as well as the interactions of substrate and time on WTN

and DTN ($P < 0.05$, Table 1).

The biofloc volume (BFV) levels in nB treatment and sB treatment were similar,

with a value of 8.8±2.3 and 8.3±2.3 mL L^{-1}, respectively (Table 1). However, the total

suspended solids (TSS) and turbidity in sB treatment (491.3±150.2 mg L^{-1}, and

299.5±92.0 nephelometric turbidity units, NTU) were higher than those in nB treatment

(148.5±31.3 mg L^{-1} and 111.9±56.3 NTU, Table 1). Substrate showed significant main

effects on those three parameters ($P < 0.05$, Table 1). The effects of time on BFV and

turbidity were also significantly, as well as the interaction of substrate and time on BFV

($P > 0.05$, Table 1).

3.2 Growth performance

During the 28-days culture experiment, the average body weights of shrimp in sB

treatment were higher than those of nB treatment (Fig. 1). At the end, although the final

body weight in sB treatment (0.40±0.03 g) was not significantly different from that of B

treatment (0.36±0.04 g, P = 0.596, Table 2), the survival rate (96.3±3.6%) and

productivity (1.54±0.12 kg m^{-3}) of the former treatment were significantly higher than

those of the latter (81.0±7.1% and 1.14±0.09 kg m^{-3}, P < 0.05, Table 2). The feed

conversion rate (FCR) in sB treatment (0.76±0.06) was significantly lower than that in

nB treatment (0.98±0.08, P = 0.044, Table 2). There no significant difference was

observed between both treatments for the weekly increment of body weight (wiW) and

specific growth rate (SGR) of shrimp during the culture experiment (P > 0.05, Table 2).

3.3 Bacterial diversity

The Shannon index for bacterial community in sB treatment (6.77±0.18) was

higher than that in nB treatment (6.14±0.07, Table 3). Substrate, time and their

interaction showed significant effects on this index (P < 0.05, Table 3). The Shannon

index in sB treatment obtained the highest value at 7 d and then slightly decreased, but

that in nB treatment peaked at 21 d (Fig. 2 a).

Higher OTU counts (6238.3±353.8) and Margalef index (387.7±28.0) was

observed in sB treatment, when compared to that in nB treatment (5713.5±368.2 and

343.9±22.0, Table 3). Both indexes significantly affected by time (P < 0.05), but not by

substrate (P > 0.05, Table 3). Those two indexes showed similar changing trends with

the Shannon index in each treatment (Fig. 2 b and c).

The evenness indexes, Heip index and Pielou index, in sB treatment (0.018±0.001

and 0.538±0.003) and nB treatment (0.013±0.001 and 0.493±0.01, Table 3) peaked at

14 d and 21 d, respectively (Fig. 2 d and e). The main effects and interactions of

substrate and time on both indexes were significant ($P < 0.05$), with exception of the

interaction on Heip index ($P = 0.544$, Table 3).

The Simpson index in sB treatment (0.955±0.003) was higher than that in nB

treatment (0.927±0.010, Table 3). This index in sB treatment displayed a peak value at

14 d, but that in nB treatment showed a decreasing trend during experiment (Fig. 2 f).

Substrate and time significantly affected this index ($P < 0.05$, Table 3). Whereas, the

Berger-Parker index in sB treatment (0.152±0.014) was lower than that in nB treatment

(0.193±0.020, Table 3). Substrate was not significantly affected this index ($P > 0.05$,

Table 3). The change trend of Berger-Parker index in the present study was contrary to

that of the Simpson index in each treatment (Fig. 2 g).

The bacterial community distances within nB treatment and sB treatment were

0.76±0.36 and 0.77±0.37, respectively (Table 4). PCA analysis also showed that water

samples collected from the same treatment were closer than those from the other

treatment, basing on the OTUs composition (Fig. 3). In addition, the bacterial beta

diversities between both treatments were significantly different with a Jaccard distance

of 0.94±0.01 ($P = 0.001$, PERMANOVA, Table 4).

3.4 Bacterial composition profile

283 Thirty phyla were assigned for the OTUs in the present study. The most dominant phyla

284 in both treatments were Actinobacteria (4.0-22.7%), Bacteroidetes (10.4-33.5%),

285 Firmicutes (0.2-11.2%), Planctomycetes (4.0-14.9%) and Proteobacteria (29.4-59.0%)

286 (Fig. 4 a). Besides, Chloroflexi and Verrucomicrobia were one of the dominant phyla in

287 nB treatment (0.1-11.4%) and sB treatment (0.3-16.2%), respectively (Fig. 4a). Among

288 those seven phyla mentioned above, the proportions of Chloroflexi, Planctomycetes

289 and Verrucomicrobia were significantly different between both treatments from a point

290 of view of the whole experimental period ($P < 0.05$, Fig. 4 b). The main effects and

291 interactions of substrate and time on proportions of those seven phyla were significant

292 ($P < 0.05$), except the substrate effects on Firmicutes and Proteobacteria, and the

293 interaction of substrate and time on Planctomycetes ($P > 0.05$, Table 5).

294 The bacterial composition profiles on class, order, family and genus levels were

295 showed in Fig. 1S-Fig. 4S, respectively.

296 The LEfSe analysis showed that an unassigned genus and the phylum

297 Verrucomicrobia were the most significant biomarkers for nB and sB, respectively (Fig.

298 5).

299 **4 DISCUSSION**

300 **4.1 Effects of substrate on water quality**

301 All the water parameters determined in the present study were within the acceptable

302 range for culturing *P. vannamei* (Furtado et al., 2014; Hargreaves, 2013; Schveitzer et

303 al., 2013a; Van Wyk et al., 1999; Xu et al., 2012; Zhang et al., 2017). Addition of

substrate showed significantly effect on TSS in the present study ($P < 0.05$), with

higher level in sB treatment than that in nB treatment. Consistently, Schveitzer et al.

(2013b) also observed significant effect of substrate on TSS with higher levels in the

biofloc systems with substrate. Whereas, in another previous study, the TSS levels in

the biofloc systems with substrate were lower than that in the system without substrate

(Rezende et al., 2018). It is proposed here that substrates with different shape have

different effects on TSS. For example, nylon or polyester fiber net with a small mesh

size (1-2 mm) has the ability to filter the water and capture biofloc, thereby reducing the

amount of solids in the rearing water, leading to the lower TSS value (Rezende et al.,

2018). On the contrary, the substrate with a larger mesh size, such as 2.5 cm (Tierney et

al., 2020), would filter both water and biofloc. Similarly, in the present study, elastic

solid packing filler (nylon) was used as substrate which is a long open space composed

with nylon yarns radially fixed to a central rope. Thus, the flocs would be not retained

within this open space of substrate, especially under the rough aeration condition of the

biofloc system, leading to an increasing of TSS level in water body.

In the present study, it was found that the turbidity levels in both treatments were

increased continuously. This might be contributed to the removal operation for biofloc

during 14-28 d of the experiment in the present study. The turbidity in the present study

was determined on water sample without big-size biofloc which settled by using an

Imhoff cone for 15 min, indicating that this parameter was correlated to the content of

small-size biofloc. Previous studies have showed that the bacterial groups are different

between small- and big-size bioflocs (Chen et al., 2019; Huang et al., 2020). Therefore,

the removal operation in this study might make settleable solids (big-size biofloc) as

well as bacterial groups attached be removed from the water column, which would

improve the growth of bacteria adhering on small-size biofloc retained in the water

body, by reducing the competition from bacteria relating with big-size biofloc, and in

turn, prompting formation of small-size suspended biofloc and increasing the turbidity.

Substrate showed significant effect on turbidity in the present study ($P = 0.002$). It

was thought that the addition of artificial substrates may influence the water circulation

in the tanks, leading to smaller turbulence in the water, and in turn to facilitating

sedimentation or particle aggregation and formation of large flocs, which reduces the

suspended solids and thus the turbidity in the water column (Ferreira et al., 2016;

Fleckenstein et al., 2020). However, in the present study, the turbidity level in sB

treatment was found to be higher than that in nB treatment. It was speculated that

substrate might play a positive role in the process of turbidity increasing caused by the

removal operation for big-size biofloc discussed above.

Substrate also showed significant effect on carbonate alkalinity in the current study

($P < 0.001$). And the carbonate alkalinity in sB treatment was lower than that in nB

treatment in the current study, indicating that more alkalinity was consumed in biofloc

system with substrate. This might be contributed to the different bacterial communities

which led to different bacterial activities consuming HCO_3^- (Ebeling et al., 2006).

Corroboratively, more diverse bacterial community in sB treatment (Shannon index

6.77±0.18) than that in nB treatment (Shannon index 6.14±0.07), and different bacterial

communities between both treatments (Jaccard distance 0.94±0.01, $P < 0.001$) were

observed in the present study. Nevertheless, the inferring function profile of bacterial

community should be investigated to better understand the difference on HCO_3^-

consumption between both treatments next.

The results of ANOVA with repeated measurements showed that addition of

substrate showed significantly effects on TAN, biofloc volume (BFV), and pH in the

present study ($P < 0.05$). Meanwhile, the interactions of substrate and time on most

water parameters, such as whole total nitrogen in waterbody (WTN), total dissolved

nitrogen (DTN), TAN, BFV, carbonate alkalinity and pH, were significant in the

present study, indicating that the changing styles of these parameters along with time

elapsing were different between the biofloc systems with and without addition of

substrate. Rezende et al. (2018) also observed that some water parameters were

significantly affected by the interactions of the rearing day and treatment (substrate).

4.2   Effects of substrate on shrimp growth

The body weight of shrimp in treatment with substrate would be significantly prompt,

when compared to that without substrate (Olier et al., 2020; Schveitzer et al., 2013b),

via forming periphytons on substrate as supplementary natural food which could be fed

by shrimp for 24 h a day (Anand et al., 2019; Asaduzzaman et al., 2008; Kumar et al.,

2017). In the present study, the final body weight, weekly increment of body weight

(wiW) and specific growth rate (SGR) of shrimp in sB treatment were elevated by

10.0-11.1%, compared to those in nB treatment, which were similar with the results

(8.4-11.0%) observed by Tierney et al. (2020). Previous studies showed that the raw

material (Mani et al., 2021; Peiro-Alcantar et al., 2019; Rezende et al., 2018) and shape

(Fleckenstein et al., 2020) of substrates could affect its elevating efficiency on growth

performance of shrimp. For example, periphyton nutritional values were shown to be

dependent on substrate type (Tinh et al., 2021). And natural or vegetable substrates was

found to be more beneficial to growth of shrimp than artificial substrates (Mani et al.,

2021). It is also thought that the substrate composed of individual sheets with a large

opening or mesh size and a low overall surface area would show less improving effect

on shrimp production metrics. For instance, Ferreira et al. (2016) found that the

addition of substrate with an amount equivalent 400% of the lateral area of the tanks

significantly elevated the body weight of shrimp, but that with 200% did not. Similarly,

Fleckenstein et al. (2020) and Tierney et al. (2020) also observed that addition of

substrate only increased the total in-tank surface area by 13.4% and 21%, respectively,

did not lead to significant effect on any shrimp production metrics. In this study, the

substrate used was a long open space composed with nylon yarns radially fixed to a

central rope, covering only 52.4% of the tank internal surface, which might be

contributed to the non-significant benefit for the final body weight, wiW and SGR of

shrimp in sB treatment, compared to those in nB treatment ($P > 0.05$).

Nevertheless, in the present study, the survival rate and productivity in sB

treatment significantly increased when compared with those in nB treatment ($P < 0.05$).

Substrate could act as a refuge to add space and reduce the cannibalism behavior,

leading to more survival. And this increased space by addition of substrate reduced the

relative stocking density, which is beneficial for shrimp growth. An inverse

relationship between stocking density and growth performance of shrimp was found in

previous studies (Esparza-Leal et al., 2015; Esparza-Leal et al., 2020 ; Liu et al., 2017;

Panigrahi et al., 2019; Peixoto et al., 2018). Moreover, the feed conversion rate (FCR)

in sB treatment significantly decreased, compared with that in nB treatment ($P < 0.05$).

Bioflocs which mostly grew on tank wall or substrate as periphyton could be grazed by

shrimp as natural supplementary food (Tinh et al., 2021), reducing input of artificial

feeds, and consequently decreasing the FCR (Avnimelech, 2015). Undoubtedly,

substrate addition facilitates this feeding behavior of shrimp by supplying a niche for

preying on biofloc (Anand et al., 2019; Asaduzzaman et al., 2008).

4.3 Effects of substrate on diversity of bacterial diversity

Generally, highly diverse bacterial community including 18-40 phyla were observed in

waterbody of biofloc systems raising *P. vannamei* (Baliga et al., 2021; Ferreira et al.,

2021; Huerta-Rabago et al., 2019; Vargas-Albores et al., 2019), with a Shannon index

as high as 5.31-7.23 (Addo et al., 2021; Chen et al., 2019; Gao et al., 2019; Huang et al.,

2021; Jiang et al., 2020; Kim et al., 2021; Pilotto et al., 2018). Similarly, in the present

study, 30 phyla were observed in both treatments with a Shannon index of 6.14-6.77.

In the biofloc system, the environment rich in organic matter could form several trophic

levels which facilitate the growth of a number of bacterial phyla (Fleckenstein et al.,

2020), resulting in a high bacterial diversity.

Between both treatments in the present study, the beta diversity Jaccard index was

significantly different, indicating different bacterial communities in both treatments.

Corroboratively, from a point of view of the whole experimental period, higher

Shannon index, OTU counts, Margalef index, Heip index, Pielou index and Simpson

index, but lower Berger-Parker index of bacterial community in sB treatment were

observed when compared to those in nB treatment, indicating higher bacterial diversity,

richness and evenness, but lower dominance of predominant species (Magurran, 2004),

in the biofloc system with addition of substrate (sB treatment).

The results of ANOVA with repeated measurements revealed that substrate showed

significant effects on some alpha diversity indexes, such as Shannon index, Heip index,

Pielou index and Simpson index ($P < 0.05$). Meanwhile, the interactions of substrate

and time on Shannon index and Pielou index were also significant ($P < 0.05$), indicating

that the changing styles of those indexes over time were different between treatments in

the present study. Consistently, the bacterial community in sB treatment obtained the

most diverse situation at 7 d with the highest Shannon index, but in nB treatment, the

Shannon index increased gradually and peaked at 21 d, indicating that substrate

addition accelerated the formation of high-diverse bacterial community in the biofloc

system.

All alpha diversity indexes were significantly affected by time ($P < 0.05$),

indicating a time-dependent succession pattern of bacterial diversity in biofloc systems

regardless of addition of substrate, which was also observed in other studies (Jiang et al.,

2020; Kim et al., 2021; Wang et al., 2020; Xu et al., 2019).

4.4   Effects of substrate on bacterial composition

In general, Proteobacteria and Bacteroidetes were found to be the most dominant phyla

in marine biofloc systems rearing *P. vannamei* (Ferreira et al., 2021; Huerta-Rabago et

al., 2019; Jiang et al., 2020; Pilotto et al., 2018; Martínez-Córdova et al., 2018;

Vargas-Albores et al., 2019), as well as in biofloc systems with low salinities (Addo et

al., 2021; Huang et al., 2021). Similarly, in the present study, those two phyla together

accounted for 41.33-82.44% and 55.56-74.69% in nB treatment and sB treatment,

respectively. Proteobacteria have a diverse phenotypic and phylogenetic lineage and

widely dispersed in the environment rich in organic compounds (Kersters et al., 2006;

Kirchman, 2002; Rud et al., 2017), such as biofloc systems. Bacteroidetes serve as a

degrader of polymeric organic matter and are found in a range of habitats (Thomas et al.,

2011; Woebken et al., 2007). Additionally, Actinobacteria, Chloroflexi, Firmicutes,

Planctomycetes and Verrucomicrobia were important phyla in both treatments of the

present study, which were also commonly detected in biofloc systems stocking

white-leg shrimp (Deng et al., 2018; Ferreira et al., 2021; Huang et al., 2021;

Huerta-Rabago et al., 2019; Jiang et al., 2020; Luis-Villaseñor et al., 2016;

Martínez-Córdova et al., 2018; Martins et al., 2020; Schveitzer et al., 2020).

Substrate showed significantly main effects on the proportions of most of the seven

predominant phyla mentioned above in this current study. It is considered that substrate

could supply attaching material for species which grow adhesively, and tend to adsorb

the bacteria with particle-attaching or biopolymer-degrading capability (Huang et al.,

2020). For example, the filamentous Chloroflexi grow on other microorganisms and

substrate (Kragelund et al., 2007; Miura et al., 2007); and Bacteroidetes prefer for

polymers and are frequently found attaching to surfaces, colonizing on particles (Deng

et al., 2019; Fernández-Gómez et al., 2013; Kirchman, 2002; Morgan-Sagastume et al.,

2008). Although Planctomycetes grow as single free-swimming cells, they also appear

as single cells within flocs (Morgan-Sagastume et al., 2008), and are usually found to

adhere to the surfaces of invertebrates (Fuerst et al., 1997), detrital aggregates (Crump

et al., 1999; Delong et al., 1993), and flocs (Huang et al., 2020). Actinobacteria are

chemoorganotrophic bacteria (saprophytes) with branched filaments (Madigan et al.,

2015) and have been described as small free-living cells (Ghai et al., 2013), leading to a

higher proportion in free-living fraction than that in particle-attached fraction

(Bachmann et al., 2018; Huang et al., 2020; Mestre et al., 2017). Therefore, the free- or

attaching-living characteristic of those phyla might contribute to the variations of their

abundance between both treatments in the present study.

Although Firmicutes are microorganisms to grow as single free-swimming cells

making them more likely to remain in suspension (Morgan-Sagastume et al., 2008).

And Proteobacteria survive as free-living to live suspended in the water column,

although some species might attach to particles, surfaces, or cells (Fernández-Gómez et

al., 2013; Morgan-Sagastume et al., 2008; Rud et al., 2017; Schveitzer et al., 2020),

substrate did not show significant effect on those two phyla in this present study. And

the reason should be investigated next.

The dominant phyla observed in the present study have variable functions. For

instance, Proteobacteria play important roles in the process of nutrient cycling and the

mineralization of organic compounds (Kersters et al., 2006; Kirchman, 2002; Rud et al.,

2017); Bacteroidetes produce extracellular enzymes with degradative capabilities to

serve as degraders of polymeric organic matter (Kirchman, 2002; Thomas et al., 2011;

Woebken et al., 2007); Chloroflexi reductively dehalogenate organic chlorinated

compounds to obtain energy (Gupta, 2013); Actinobacteria are important in the nutrient

and biomaterial recycling (Goodfellow and Williams, 1983), responsible for degrading

nucleic acids and proteins (Madigan et al., 2015), and for the production of many

hydrolytic enzymes (chitinase, chitinolytic, fatty acid, cellulolytic, nitrile hydrolyzing,

amylolytic, proteolytic, and lipolytic) (Das et al., 2008); Planctomycetes have function

to degrade the biopolymers of adsorbate (Woebken et al., 2007). Therefore, the

different bacterial community profiles between both treatments in the current study

might contribute to different activities of nitrogen and carbon metabolism in their water

body. However, the inferring functions of bacterial community should be induced next

to directly estimate the effects of addition of substrate on the bacterial activities.

The interactions of substrate and time on proportions of most of those phyla (except

Planctomycetes) also significant in the present study, indicating different development

styles of bacterial composition profile between both treatments. Furthermore, all the

seven dominant phyla in the present study were significantly affected by time ($P <$

0.05), indicating a time-dependent dynamics of bacterial composition in biofloc

systems, similar with that observed in other studies (Baliga et al., 2021; Gao et al., 2019;

Jiang et al., 2020; Martínez-Córdova et al., 2018; Xu et al., 2019).

5 CONCLUSION

In the present study, addition of elastic solid packing filler (nylon, with a diameter

of 15 cm) as substrate that coated 52.4% of the tank internal surface in the biofloc

system nursing *P. vannamei* PL with a salinity of 5‰ showed positive effects on

shrimp growth performance. Furthermore, substrate also showed significant main

effects or interactions with time on most water parameters as well as bacterial diversity

and composition.

Acknowledgements This work is financially supported by the scientific research

program of the Education Department of Hunan province (18B394), the open projects

of Hunan Provincial Key Laboratory for Molecular Immunity Technology of Aquatic

Animal Diseases (2021KJ001), Changde Research Center for Agricultural

Biomacromolecule (2020AB08).

REFERENCES

Addo, F.G., Zhang, S., Manirakiza, B., Ohore, O.E., Shudong, Y., 2021. The impacts

of straw substrate on biofloc formation, bacterial community and nutrient

removal in shrimp ponds. Bioresour. Technol. 326, 124727.

Anand, P.S.S., Balasubramanian, C.P., Christina, L., Kumar, S., Biswas, G., De, D.,

Ghoshal, T.K., Vijayan, K.K., 2019. Substrate based black tiger shrimp, *Penaeus*

monodon culture: Stocking density, aeration and their effect on growth

performance, water quality and periphyton development. Aquaculture. 507,

411-418.

Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of

variance. Austral Ecol. 26, 32-46.

APHA, 1995. Standard Methods for the Examination of Water and Wastewater.

American public health association, Washington, DC.

Asaduzzaman, M., Wahab, M.A., Mcj, V., Huque, S., Salam, M.A., Azim, M.E., 2008.

C/N ratio control and substrate addition for periphyton development jointly

enhance freshwater prawn *Macrobrachium rosenbergii* production in ponds.

Aquaculture. 280, 117-123.

Avnimelech, Y., 2015. Biofloc technology: a practical hand book, 3rd ed. The world

aquaculture society, Baton Rouge, Louisiana, EUA.

Bachmann, J., Heimbach, T., Hassenrück, C., Kopprio, G.A., Iversen, M.H., Grossart,

H.P., Gärdes, A., 2018. Environmental drivers of free-living vs. particle-attached

bacterial community composition in the Mauritania upwelling system. Front.

Microbiol. 9, 2836.

Baliga, P., Goolappa, P.T., Shekar, M., Girisha, S.K., Ramesh, K.S., Udyavara, V.,

Venugopal, M.N., 2021. Bacterial communities associated with *Penaeus*

vannamei Boone, 1931 surface and its rearing water in biofloc culture system.

Asian Fish. Sci. 34, 34-46.

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith,

G.A., Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J.E.,

Bittinger, K., Brejnrod, A., Brislawn, C.J., Brown, C.T., Callahan, B.J.,

Caraballo-Rodríguez, A.M., Chase, J., Cope, E.K., Da Silva, R., Diener, C.,

Dorrestein, P.C., Douglas, G.M., Durall, D.M., Duvallet, C., Edwardson, C.F.,

Ernst, M., Estaki, M., Fouquier, J., Gauglitz, J.M., Gibbons, S.M., Gibson, D.L.,

Gonzalez, A., Gorlick, K., Guo, J., Hillmann, B., Holmes, S., Holste, H.,

Huttenhower, C., Huttley, G.A., Janssen, S., Jarmusch, A.K., Jiang, L., Kaehler,

B.D., Kang, K.B., Keefe, C.R., Keim, P., Kelley, S.T., Knights, D., Koester, I.,

Kosciolek, T., Kreps, J., Langille, M.G.I., Lee, J., Ley, R., Liu, Y.-X., Loftfield,

E., Lozupone, C., Maher, M., Marotz, C., Martin, B.D., McDonald, D., McIver,

L.J., Melnik, A.V., Metcalf, J.L., Morgan, S.C., Morton, J.T., Naimey, A.T.,

Navas-Molina, J.A., Nothias, L.F., Orchanian, S.B., Pearson, T., Peoples, S.L.,

Petras, D., Preuss, M.L., Pruesse, E., Rasmussen, L.B., Rivers, A., Robeson, M.S.,

Rosenthal, P., Segata, N., Shaffer, M., Shiffer, A., Sinha, R., Song, S.J., Spear,

J.R., Swafford, A.D., Thompson, L.R., Torres, P.J., Trinh, P., Tripathi, A.,

Turnbaugh, P.J., Ul-Hasan, S., van der Hooft, J.J.J., Vargas, F., Vázquez-Baeza, Y.,

Vogtmann, E., von Hippel, M., Walters, W., Wan, Y., Wang, M., Warren, J., Weber,

K.C., Williamson, C.H.D., Willis, A.D., Xu, Z.Z., Zaneveld, J.R., Zhang, Y., Zhu,

Q., Knight, R., Caporaso, J.G., 2019. Reproducible, interactive, scalable and

extensible microbiome data science using QIIME 2. Nat. Biotechnol. 37,

852-857.

Chen, X., Luo, G.-z., Meng, H., Tan, F.Y., 2019. Effect of the particle size on the

ammonia removal rate and the bacterial community composition of bioflocs.

Aquacult. Eng. 86, 1-8.

Crump, B.C., Armbrust, E.V., Baross, J.A., 1999. Phylogenetic analysis of

particle-attached and free-living bacterial communities in the Columbia River, its

estuary, and the adjacent coastal ocean. Appl. Environ. Microbiol. 65, 3192-3204.

Das, S., Ward, L.R., Burke, C., 2008. Prospects of using marine actinobacteria as

probiotics in aquaculture. Appl. Microbiol. Biotechnol. 81, 419-429.

Delong, E.F., Franks, D.G., Alldredge, A.L., 1993. Phylogenetic diversity of

aggregate-attached vs. free-living marine bacterial assemblages. Limnol. &

Oceanogr. 38, 924-934.

Deng, M., Chen, J., Hou, J., Li, D., He, X., 2018. The effect of different carbon

sources on water quality, microbial community and structure of biofloc systems.

Aquaculture. 482, 103-110.

Deng, Y., Xu, X., Yin, X., Lu, H., Chen, G., Yu, J., Ruan, Y., 2019. Effect of stock

density on the microbial community in biofloc water and Pacific white shrimp

(*Litopenaeus vannamei*) gut microbiota. Appl. Microbiol. Biotechnol. 103,

4241-4252.

Ebeling, J.M., Timmons, M.B., Bisogni, J.J., 2006. Engineering analysis of the

stoichiometry of photoautotrophic, autotrophic, and heterotrophic removal of

ammonia–nitrogen in aquaculture systems. Aquaculture. 257, 346-358.

Esparza-Leal, H.M., Cardozo, A.P., Wasielesky, W.Jr., 2015. Performance of

Litopenaeus vannamei postlarvae reared in indoor nursery tanks at high stocking

density in clear-water versus biofloc system. Aquacult. Eng. 68, 28-34.

Esparza-Leal, H.M., Amaral Xavier, J.A., Wasielesky, W.Jr., 2016. Performance of

Litopenaeus vannamei postlarvae reared in indoor nursery tanks under biofloc

conditions at different salinities and zero-water exchange. Aquacult. Int. 24,

1435-1447.

Esparza-Leal, H.M., Ponce-Palafox, J.T., Alvarez-Ruiz, P., Lopez-Alvarez, E.S.,

Vazquez-Montoya, N., Lopez-Espinoza, M., Mejia, M.M., Gomez-Peraza, R.L.,

Nava-Perez, E., 2020. Effect of stocking density and water exchange on

performance and stress tolerance to low and high salinity by *Litopenaeus*

vannamei postlarvae reared with biofloc in intensive nursery phase. Aquacult. Int.

28, 1473-1483.

FAO, 2018. The state of world fisheries and aquaculture - Meeting the sustainable

development goals. Licence: CC BY-NC-SA 3.0 IGO, Rome.

Fernández-Gómez, B., Richter, M., Schüler, M., Pinhassi, J., Acinas, S.G., González,

J.M., Pedrós-Alió, C., 2013. Ecology of marine Bacteroidetes: a comparative

genomics approach. The ISME Journal. 7, 1026-1037.

Ferreira, G.S., Santos, D., Schmachtl, F., Machado, C., Fernandes, V., Boegner, M.,

Schleder, D.D., Seiffert, W.Q., Vieira, F.N., 2021. Heterotrophic,

chemoautotrophic and mature approaches in biofloc system for Pacific white

shrimp. Aquaculture. 533, 736099.

Ferreira, L.M.H., Lara, G., Wasielesky, W.Jr., Abreu, P.C., 2016. Biofilm versus

biofloc: Are artificial substrates for biofilm production necessary in the BFT

system? Aquacult. Int. 24, 921-930.

Fleckenstein, L.J., Kring, N.A., Tierney, T.W., Fisk, J.C., Lawson, B.C., Ray, A.J.,

2020. The effects of artificial substrate and stocking density on Pacific white

shrimp (*Litopenaeus vannamei*) performance and water quality dynamics in high

tunnel-based biofloc systems. Aquacult. Eng. 90, 102093.

Fuerst, J.A., Gwilliam, H.G., Lindsay, M., Lichanska, A., Belcher, C., Vickers, J.E.,

Hugenholtz, P., 1997. Isolation and molecular identification of planctomycete

bacteria from postlarvae of the giant tiger prawn, *Penaeus monodon*. Appl.

Environ. Microbiol. 63, 254-262.

Furtado, P.S., Poersch, L.H., Wasielesky, W.Jr., 2015. The effect of different alkalinity

levels on *Litopenaeus vannamei* reared with biofloc technology (BFT). Aquacult.

Int. 23, 345-358.

Furtado, P.S., Gaona, C.A.P., Poersch, L.H., Wasielesky, W.Jr., 2014. Application of

different doses of calcium hydroxide in the farming shrimp *Litopenaeus*

vannamei with the biofloc technology (BFT). Aquacult. Int. 22, 1009-1023.

Gao, F., Liao, S., Liu, D., Bai, H., Wang, A., Ye, J., 2019. The combination use of

Candida tropicalis HH8 and *Pseudomonas stutzeri* LZX301 on nitrogen removal,

biofloc formation and microbial communities in aquaculture. Aquaculture. 500,

50-56.

Gaona, C.A.P., Almeida, M.S.d., Viau, V., Poersch, L.H., Wasielesky, W.Jr., 2017.

Effect of different total suspended solids levels on a *Litopenaeus vannamei*

(Boone, 1931) BFT culture system during biofloc formation. Aquacult. Res. 48,

1070-1079.

Ghai, R., Mizuno, C.M., Picazo, A., Camacho, A., Rodriguez-Valera, F., 2013.

Metagenomics uncovers a new group of low GC and ultra-small marine

Actinobacteria. Sci. Rep. 3, 2471.

Goodfellow, M., Williams, S.T., 1983. Ecology of Actinomycetes. Annu. Rev.

Microbiol. 37, 189-216.

Gupta, R.S., 2013. Molecular Markers for Photosynthetic Bacteria and Insights into

the Origin and Spread of Photosynthesis. in: Beatty, J.T. (Ed.), Advances in

Botanical Research. Academic Press.

Hargreaves, J., 2013. Biofloc production systems for aquaculture. Southern Regional

Aquaculture Center, United States, Department of Agriculture, National Institute

of Food and Agriculture.

Huang, H.-H., 2020. Novel biofloc technology (BFT) for ammonia assimilation and

reuse in aquaculture in situ. in: Lu, Q., Serajuddin, M. (Eds.), Emerging

technologies and research for eco-friendly aquaculture. IntechOpen, London.

Huang, H.-H., Luo, T., Lei, Y.-J., Kuang, W.-Q., Zou, W.-S., Yang, P.-H., 2021. Water

quality, shrimp growth performance and bacterial community in a reusing-water

biofloc system for nursery of *Penaeus vananmei* rearing under a low salinity

condition. Aquacult. Rep. 21, 100894.

Huang, L., Guo, H., Chen, C., Huang, X., Chen, W., Bao, F., Liu, W., Wang, S., Zhang,

D., 2020. The bacteria from large-sized bioflocs are more associated with the

shrimp gut microbiota in culture system. Aquaculture. 523, 735159.

Huerta-Rabago, J.A., Martinez-Porchas, M., Miranda-Baeza, A., Nieves-Soto, M.,

Rivas-Vega, M.E., Martinez-Cordova, L.R., 2019. Addition of commercial

probiotic in a biofloc shrimp farm of *Litopenaeus vannamei* during the nursery

phase: Effect on bacterial diversity using massive sequencing 16S rRNA.

Aquaculture. 502, 391-399.

Jiang, W., Ren, W., Li, L., Dong, S., Tian, X., 2020. Light and carbon sources addition

alter microbial community in biofloc-based *Litopenaeus vannamei* culture

systems. Aquaculture. 515, 734572.

Kersters, K., Vos, P.D., Gillis, M., Swings, J., Stackebrandt, E., 2006. Introduction to

the Proteobacteria. in: Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.H.,

Stackebrandt, E. (Eds.), The Prokaryotes. Springer, New York.

Khanjani, M.H., Sajjadi, M.M., Alizadeh, M., Sourinejad, I., 2017. Nursery

performance of Pacific white shrimp (*Litopenaeus vannamei* Boone, 1931)

cultivated in a biofloc system: the effect of adding different carbon sources.

Aquacult. Res. 48, 1491-1501.

Kim, Y.-S., Kim, S.-E., Kim, S.-J., Jung, H.-K., Park, J., Jeon, Y.J., Kim, D.-H., Kang,

J.-H., Kim, K.-H., 2021. Effects of wheat flour and culture period on bacterial

community composition in digestive tracts of *Litopenaeus vannamei* and rearing

water in biofloc aquaculture system. Aquaculture. 531, 735908.

Kirchman, D.L., 2002. The ecology of Cytophaga–Flavobacteria in aquatic

environments. FEMS Microbiol. Ecol. 39, 91-100.

Kragelund, C., Caterina, L., Borger, A., Thelen, K., Eikelboom, D., Tandoi, V., Kong,

Y., Van Der Waarde, J., Krooneman, J., Rossetti, S., Thomsen, T.R., Nielsen, P.H.,

2007. Identity, abundance and ecophysiology of filamentous Chloroflexi species

present in activated sludge treatment plants. FEMS Microbiol. Ecol. 59, 671-682.

Kumar, S., Anand, P.S.S., De, D., Deo, A.D., Ghoshal, T.K., Sundaray, J.K., Ponniah,

A.G., Jithendran, K.P., Raja, R.A., Biswas, G., Lalitha, N., 2017. Effects of

biofloc under different carbon sources and protein levels on water quality, growth

performance and immune responses in black tiger shrimp *Penaeus monodon*

(Fabricius, 1978). Aquacult. Res. 48, 1168-1182.

Kumar, V.S., Pandey, P.K., Anand, T., Bhuvaneswari, R., Kumar, S., 2017. Effect of

periphyton (aquamat) on water quality, nitrogen budget, microbial ecology, and

growth parameters of *Litopenaeus vannamei* in a semi-intensive culture system.

Aquaculture. 479, 240-249.

Lara, G., Krummenauer, D., Abreu, P.C., Poersch, L.H., Wasielesky, W.Jr., 2017. The

use of different aerators on *Litopenaeus vannamei* biofloc culture system: effects

on water quality, shrimp growth and biofloc composition. Aquacult. Int. 25,

147-162.

Liu, G., Zhu, S., Liu, D., Guo, X., Ye, Z., 2017. Effects of stocking density of the

white shrimp *Litopenaeus vannamei* (Boone) on immunities, antioxidant status,

and resistance against Vibrio harveyi in a biofloc system. Fish Shellfish Immunol.

67, 19-26.

Luis-Villaseñor, I.E., Voltolina, D., Audelo-Naranjo, J.M., Pacheco-Marges, M.R.,

Herrera-Espericueta, V.E., Romero-Beltrán, E., 2016. Effects of biofloc

promotion on water quality, growth, biomass yield and heterotrophic community

in *Litopenaeus vannamei* (Boone, 1931) experimental intensive culture. Ital. J.

Anim. Sci. 14, 3726.

Luo, G., Liu, Z., Shan, L., Tan, H., 2019. Using poly-beta-hydroxybutyric as an

additional carbohydrate for biofloc in a shrimp *Litopenaeus vannamei* bioflocs

nursery system with brackish water. Aquaculture. 506, 181-187.

Madigan, M.T., Martinko, J.M., Bender, K.S., Buckley, D.H., Stahl, D.A., 2015.

Brock biology of microorganisms, 14 ed. Pearson, IL.

Magurran, A.E., 2004. Measuring biological diversity. Blackwell Publishing, Oxford.

Mani, S., Ramasamy, S.M., Chakrapani, S., Krishna, A., Anand, P.S.S., Lalramchhani,

C., Antony, J., Panigrahi, A., 2021. The effect of natural and artificial periphytic

substrates with biofloc system on shrimp *Penaeus vannamei* (Boone 1931)

culture: growth and immune response. Aquacult. Int. 29, 651-668.

Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput

sequencing reads. Embnet Journal. 17, 10-12.

Martínez-Córdova, L.R., Francisco, V.A., Estefanía, G.V., Ortíz-Estrada, á.M.,

Porchas-Cornejo, M.A., Asunción, L.L., Marcel, M.P., 2018. Amaranth and wheat

grains tested as nucleation sites of microbial communities to produce bioflocs

used for shrimp culture. Aquaculture. 497, 503-509.

Martins, M.A., Poli, M.A., Legarda, E.C., Pinheiro, I.C., Carneiro, R.F.S., Pereira,

S.A., Martins, M.L., Goncalves, P., Schleder, D.D., Vieira, F.d.N., 2020.

Heterotrophic and mature biofloc systems in the integrated culture of Pacific

white shrimp and Nile tilapia. Aquaculture. 514, 734517.

Mestre, M., Borrull, E., Sala, M.M., Gasol, J.M., 2017. Patterns of bacterial diversity

in the marine planktonic particulate matter continuum. The ISME Journal. 11,

999-1010.

Mishra, J.K., Samocha, T.M., Patnaik, S., Speed, M., Gandy, R.L., Ali, A.-M., 2008.

Performance of an intensive nursery system for the Pacific white shrimp,

Litopenaeus vannamei, under limited discharge condition. Aquacult. Eng. 38,

2-15.

Miura, Y., Hiraiwa, M.N., Ito, T., Itonaga, T., Watanabe, Y., Okabe, S., 2007. Bacterial

community structures in MBRs treating municipal wastewater: Relationship

between community stability and reactor performance. Water Res. 41, 627-637.

Morgan-Sagastume, F., Larsen, P., Nielsen, J.L., Nielsen, P.H., 2008. Characterization

of the loosely attached fraction of activated sludge bacteria. Water Res. 42,

843-854.

Olier, B.S., Tubin, J.S.B., Mello, G.L.d., Martinez-Porchas, M., Emerenciano, M.G.C.,

2020. Does vertical substrate could influence the dietary protein level and

zootechnical performance of the Pacific white shrimp *Litopenaeus vannamei*

reared in a biofloc system? Aquacult. Int. 28, 1227-1241.

Panigrahi, A., Sundaram, M., Saranya, C., Swain, S., Dash, R.R., Dayal, J.S., 2019.

Carbohydrate sources deferentially influence growth performances, microbial

dynamics and immunomodulation in Pacific white shrimp (*Litopenaeus vannamei*)

under biofloc system. Fish Shellfish Immunol. 86, 1207-1216.

Parks, D.H., Tyson, G.W., Hugenholtz, P., Beiko, R.G., 2014. STAMP: statistical

analysis of taxonomic and functional profiles. Bioinformatics. 30, 3123-3124.

Peiro-Alcantar, C., Rivas-Vega, M.E., Martinez-Porchas, M., Lizarraga-Armenta, J.A.,

Miranda-Baeza, A., Martinez-Cordova, L.R., 2019. Effect of adding vegetable

substrates on *Penaeus vannamei* pre-grown in biofloc system on shrimp

performance, water quality and biofloc composition. Lat. Am. J. Aquat. Res. 47,

784-790.

Peixoto, S., Silva, E., Costa, C.B., Nery, R.C., Soares, R., 2018. Effect of feeding

frequency on growth and enzymatic activity of Litopenaeus vannamei during

nursery phase in biofloc system. Aquacult. Nutr. 24, 579-585.

Piedrahita, R.H., 2003. Reducing the potential environmental impact of tank

aquaculture effluents through intensification and recirculation. Aquaculture. 226,

35-44.

Pierri, V., Valter-Severino, D., Goulart-de-Oliveira, K., Manoel-do-Esprito-Santo, C.,

Nascimento-Vieira, F., Quadros-Seiffert, W., 2015. Cultivation of marine shrimp

in biofloc technology (BFT) system under different water alkalinities. Braz. J.

Biol. 75, 558-564.

Pilotto, M.R., Gonçalves, A., N.A., Vieira, F.N., Seiffert, W.Q., Bachère, E., Rosa,

R.D. Perazzolo, L.M., 2018. Exploring the impact of the biofloc rearing system

and an oral WSSV challenge on the intestinal bacteriome of *Litopenaeus*

vannamei. Microorganisms. 6, 1-16.

Ray, A.J., Lotz, J.M., 2017. Comparing salinities of 10, 20, and 30‰ in intensive,

commercial-scale biofloc shrimp (*Litopenaeus vannamei*) production systems.

Aquaculture. 476, 29-36.

Rezende, P.C., Schleder, D.D., Silva, H.V.d., Henriques, F.M., Lorenzo, M.A.d.,

Seiffert, W.Q., Andreatta, E.R., Vieira, F.d.N., 2018. Prenursery of the Pacific

white shrimp in a biofloc system using different artificial substrates. Aquacult.

Eng. 82, 25-30.

Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a versatile

open source tool for metagenomics, PeerJ, e2584.

Roy, L.A., Davis, D.A., Saoud, I.P., Boyd, C.A., Pine, H.J., Boyd, C.E., 2010. Shrimp

culture in inland low salinity waters. Rev. Aquacult. 2, 191-208.

Rud, I., Kolarevic, J., Holan, A.B., Berget, I., Calabrese, S., Terjesen, B.F., 2017.

Deep-sequencing of the bacterial microbiota in commercial-scale recirculating

and semi-closed aquaculture systems for Atlantic salmon post-smolt production.

Aquacult. Eng. 78, 50-62.

Samocha, T.M., 2010. Use of intensive and super-intensive nursery systems.

Nottingham University Press, Nottingham.

Schveitzer, R., Arantes, R., Costodio, P.F.S., Santo, C.M.d.E., Arana, L.V., Seiffert,

W.Q., Andreatta, E.R., 2013a. Effect of different biofloc levels on microbial

activity, water quality and performance of *Litopenaeus vannamei* in a tank system

operated with no water exchange. Aquacult. Eng. 56, 59-70.

Schveitzer, R., Arantes, R., Baloi, M.F., Costodio, P.F.S., Arana, L.V., Seiffert, W.Q.,

Andreatta, E.R., 2013b. Use of artificial substrates in the culture of *Litopenaeus*

vannamei (Biofloc System) at different stocking densities: Effects on microbial

activity, water quality and production rates. Aquacult. Eng. 54, 93-103.

Schveitzer, R., Lorenzo, M.A.D., Felipe, D.N.V., Pereira, S.A., Mourino, J.L.P.,

Seiffert, W.Q., Andreatta, E.R., 2017. Nursery of young *Litopenaeus vannamei*

post-larvae reared in biofloc- and microalgae-based systems. Aquacult. Eng. 78,

140-145.

Schveitzer, R., Fonseca, G., Orteney, N., Temistocles Menezes, F.C., Thompson, F.L.,

Thompson, C.C., Gregoracci, G.B., 2020. The role of sedimentation in the

structuring of microbial communities in biofloc-dominated aquaculture tanks.

Aquaculture. 514, 734493.

Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S.,

Huttenhower, C., 2011. Metagenomic biomarker discovery and explanation.

Genome Biol. 12, R60.

Thomas, F., Hehemann, J.-H., Rebuffet, E., Czjzek, M., Michel, G., 2011.

Environmental and gut Bacteroidetes: The food connection. Front. Microbiol. 2,

93.

Tierney, T.W., Fleckenstein, L.J., Ray, A.J., 2020. The effects of density and artificial

substrate on intensive shrimp *Litopenaeus vannamei* nursery production.

Aquacult. Eng. 89, 102063.

Tinh, T.H., Koppenol, T., Ngoc, H.T., Verreth, J.A.J., Verdegem, M.C.J., 2021. Effects

of carbohydrate sources on a biofloc nursery system for whiteleg shrimp

(*Litopenaeus vannamei*). Aquaculture. 531.

Van Wyk, P., Davishodgkins, M., Laramore, R., Main, K.L., Mountain, J., Scarpa, J.,

1999. Farming marine shrimp in recirculating freshwater systems. Florida

department of agriculture and consumer services, Tallahassee.

Vargas-Albores, F., Martínez-Córdova, L.R., Gollas-Galván, T., Garibay-Valdez, E.,

Emerenciano, M.G.C., Lago-Leston, A., Mazorra-Manzano, M.,
Martínez-Porchas, M., 2019. Inferring the functional properties of bacterial
communities in shrimp-culture bioflocs produced with amaranth and wheat seeds
as fouler promoters. Aquaculture. 500, 107-117.

Wang, R., Guo, Z., Tang, Y., Kuang, J., Duan, Y., Lin, H., Jiang, S., Shu, H., Huang, J.,
2020. Effects on development and microbial community of shrimp *Litopenaeus*
vannamei larvae with probiotics treatment. Amb Express. 10, 109.

Woebken, D., Fuchs, B., Kuypers, M., Amann, R., 2007. Potential interactions of
particle-associated anammox bacteria with bacterial and Archaeal partners in the
Namibian upwelling system. Appl. Environ. Microbiol. 73, 4648-4657.

Xu, W., Xu, Y., Huang, X., Hu, X., Xu, Y., Su, H., Li, Z., Yang, K., Wen, G., Cao, Y.,
2019. Addition of algicidal bacterium CZBC1 and molasses to inhibit
cyanobacteria and improve microbial communities, water quality and shrimp
performance in culture systems. Aquaculture. 502, 303-311.

Xu, W.-J., Pan, L.-Q., Zhao, D.-H., Huang, J., 2012. Preliminary investigation into the
contribution of bioflocs on protein nutrition of *Litopenaeus vannamei* fed with
different dietary protein levels in zero-water exchange culture tanks. Aquaculture.
350, 147-153.

Zhang, K., Pan, L., Chen, W., Wang, C., 2017. Effect of using sodium bicarbonate to
adjust the pH to different levels on water quality, the growth and the immune
response of shrimp Litopenaeus vannamei reared in zero-water exchange
biofloc-based culture tanks. Aquacult. Res. 48, 1194-1208.

Table 1 Water parameters in the biofloc systems with (sB) and without (nB) addition of elastic solid packing filler (nylon) as substrate, during the 28-days experiment culturing *Penaeus vannamei* at a salinity of 5.0‰

parameters	Treatments		P-value		
	nB	sB	Substrate	Time	Interaction
TSS (mg L^{-1})	148.5±31.3	491.3±150.2	0.009	0.116	0.594
WTN (mg L^{-1})	76.8±11.2	76.3±9.6	0.057	<0.001	0.013
DTN (mg L^{-1})	13.0±2.6	14.7±3.6	0.890	<0.001	<0.001
BTN (mg L^{-1})	63.8±9.3	61.6±12.5	0.052	<0.001	0.085
TAN (mg L^{-1})	0.55±0.25	0.43±0.06	0.010	0.227	0.002
Nitrite (mg L^{-1})	0.37±0.28	0.10±0.03	0.352	0.348	0.342
Nitrate (mg L^{-1})	0.51±0.14	0.56±0.08	0.924	0.274	0.252
Biofloc volume (mL L^{-1})	8.8±2.3	8.3±2.3	0.044	<0.001	<0.001
Turbidity (NTU)	111.9±56.3	299.5±92.0	0.002	0.005	0.059
CAK (mg L^{-1} CaCO$_3$)	373.7±12.4	306.5±51.7	<0.001	<0.001	<0.001
Temperature (^{o}C)	26.9±0.7	27.1±0.6	0.656	<0.001	0.012
pH	7.20±0.14	7.11±0.03	0.001	0.001	<0.001
Dissolved oxygen (mg L^{-1})	5.85±0.32	5.63±0.59	0.215	<0.001	0.764

Note: BTN, the total nitrogen contained in biofloc; CAK, carbonate alkalinity; DTN, the dissolved total nitrogen in water; NTU, nephelometric turbidity units; TAN, total ammonia nitrogen; TSS, total suspended solids; WTN, the whole total nitrogen in water.

Table 2 Growth performance of *Penaeus vannamei* postlarvae cultured in the biofloc systems with (sB) and without (nB) addition of elastic solid packing filler (nylon) as substrate, during the 28-days experiment at a salinity of 5.0‰

parameters	B	sB	*P*-value
Final mean body weight (g)	0.36 ± 0.04	0.40 ± 0.03	0.596
Survival rate (%)	81.0 ± 7.1	96.3 ± 3.6	0.011
FCR	0.98 ± 0.08	0.76 ± 0.06	0.044
wiW (g week^{-1})	0.090 ± 0.014	0.099 ± 0.008	0.652
SGR (% d^{-1})	17.7 ± 0.5	18.1 ± 0.4	0.581
Productivity (kg m^{-3})	1.14 ± 0.09	1.54 ± 0.12	0.029

Note: FCR, feed conversion rate; SGR, specific growth rate; wiW, weekly increment of body weight.

Table 3 Bacterial alpha diversity indexes in the biofloc systems with (sB) and without (nB) addition of elastic solid packing filler (nylon) as substrate, during the 28-days experiment culturing *Penaeus vannamei* at a salinity of 5.0‰

Indexes	Treatments		P-value		
	B	sB	Substrate	Time	Interaction
Shannon	6.14±0.18	6.77±0.07	0.002	<0.001	<0.001
OTU counts	5713.5±368.2	6238.3±353.8	0.427	0.024	0.572
Margalef	343.9±22.0	387.7±28.0	0.367	0.007	0.471
Heip	0.013±0.001	0.018±0.001	0.026	0.012	0.544
Pielou	0.493±0.012	0.538±0.003	0.024	<0.001	0.016
B-P	0.193±0.020	0.152±0.014	0.098	0.002	0.676
Simpson	0.927±0.010	0.955±0.003	0.043	<0.001	0.127

OTU, operational taxonomic unit; B-P, Berger-Parker index.

Table 4 Bacterial community distance between the biofloc systems with (sB) and without (nB) addition of elastic solid packing filler (nylon) as substrate, during the 28-days experiment culturing *Penaeus vannamei* at a salinity of 5.0‰

Group 1	Group 2	Jaccard distance	Permutations	pseudo-*F*	*P*-value
B	B	0.761 ± 0.361	-	-	-
sB	sB	0.768 ± 0.365	-	-	-
B	sB	0.941 ± 0.007	999	3.96	0.001

Table 5 Proportions (%) of dominant phyla in the biofloc systems with (sB) and without (nB) addition of elastic solid packing filler (nylon) as substrate, during the 28-days experiment culturing *Penaeus vannamei* at a salinity of 5.0‰

Phylum	Treatments		P-value		
	nB	sB	Substrate	Time	Interaction
Actinobacteria	13.67±3.34	10.00±4.26	<0.001	<0.001	<0.001
Bacteroidetes	23.18±4.01	16.87±5.57	<0.001	<0.001	<0.001
Chloroflexi	3.79±2.57	0.63±0.22	<0.001	0.001	0.002
Firmicutes	4.53±2.09	5.92±1.79	0.087	0.001	0.010
Planctomycetes	8.57±2.10	12.49±1.00	0.006	0.035	0.486
Proteobacteria	43.82±6.10	45.90±2.46	0.405	<0.001	0.038
Verrucomicrobia	0.63±0.18	6.06±3.42	<0.001	<0.001	<0.001

Fig. 1 Mean body weight of *Penaeus vannamei* postlarvae cultured in the biofloc systems with (sB) and without (nB) addition of elastic solid packing filler (nylon) as substrate, with time elapsing during the 28-days experiment at a salinity of 5.0‰. Error bar indicates ± standard deviation (SD).

Fig. 2 Alpha diversity indexes of bacterial communities in the biofloc systems with (sB) and without (nB) addition of elastic solid packing filler (nylon) as substrate, with time elapsing during the 28-days experiment culturing *Penaeus vannamei* at a salinity of 5.0‰.

Fig. 3 PCA diagram plotting samples basing on their OTU composition profiles. Samples were collected from the biofloc systems with (sB) and without (nB) addition of elastic solid packing filler (nylon) as substrate at time points of 7, 14, 21 and 28 d during experiment culturing *Penaeus vannamei* at a salinity of 5.0‰.

Fig. 4 Heatmap displaying bacterial composition profiles at phylum level (a) and differential phyla (b) in the biofloc systems with (sB) and without (nB) addition of elastic solid packing filler (nylon) as substrate, with time elapsing during the 28-days experiment culturing *Penaeus vannamei* at a salinity of 5.0‰.

Fig. 5 Cladogram (a) and Linear discriminant analysis (LDA) scores of LEfSe (linear discriminant analysis effect size) analysis showing biomarkers in the biofloc systems with (sB) and without (nB) addition of elastic solid packing filler (nylon) as substrate, with time elapsing during the 28-days experiment culturing *Penaeus vannamei* at a salinity of 5.0‰.

a

nB
sB

b

a: f_Ellin6075
b: o_RB41
c: c_Chloracidobacteria
d: p_Acidobacteria
e: g_Arthrobacter
f: g_Unassigned_1
g: g_Leadbetterella
h: g_Fluviicola
i: f_Saprospiraceae
j: p_GN02
k: g_Lutibacterium
l: g_Unassigned_2
m: f_Bacteriovoracaceae
n: g_Marinobacter
o: g_HTCC
p: f_HTCC2188
q: g_Unassigned_2
r: f_Verrucomicrobiaceae
s: o_Verrucomicrobiales
t: c_Verrucomicrobiae
u: p_Verrucomicrobia

−4