SPRI: Spatial Pattern Recognition using Information based method for spatial gene expression data

Jin-Xian Hu [1]*, Zhi-Rui Hu* [2]*, Ye Yuan* [1,*]*, and Hong-Bin Shen* [1,*]

[1]Institute of Image Processing and Pattern Recognition, Shanghai Jiao Tong University, and Key Laboratory of System Control and Information Processing, Ministry of Education of China, Shanghai 200240, China

[2]Gladstone Institute of Data Science and Biotechnology, San Francisco, CA, USA

*Corresponding author
Address correspondence to Y.Y at: yuanye_auto@sjtu.edu.cn, H.B. Shen at: hbshen@sjtu.edu.cn

Abstract

The rapid development of spatially resolved transcriptomics has made it possible to analyze spatial gene expression patterns in complex biological tissues. To identify spatially differential expressed genes, we propose a novel and robust nonparametric information-based approach, SPRI. SPRI converts the problem of identifying spatial gene expression patterns into the detection of dependencies between spatial coordinates with observed frequencies measured by read counts. It directly models spatial transcriptome raw count without assuming a parametric model. SPRI was applied to spatial datasets with different resolutions, suggesting that SPRI outperforms previous methods, by robustly detecting more genes with significant spatial expression patterns, and revealing biological insights that cannot be identified by other methods.

Keywords: Spatial transcriptomics, Spatial gene expression pattern, Information theory

Background

Recently the rapid development of high-throughput spatial resolved transcriptome (SRT) technologies, such as 10x Visium [1] and Stereo-seq [2] enables the understanding of spatially resolved gene expression patterns in complicated tissues [3-5]. This technology first partitions tissue into small regions (spots) and labels all

transcripts within one spot with known spatial coordinate barcodes, then sequences

them to capture the expression levels of thousands of genes in the spot. Such

technology provides an efficient spatial approach for new biological discoveries and

understanding of multiple biological processes in diseases [6, 7].

Identification of genes with spatial expression patterns (SE genes) is an essential

step in the analysis of SRT data. For this task, several existing methods can be divided

into two groups: normalized data-based method and raw count data-based method.

Trendsceek [8] based on marked point processes to identify genes with spatial

expression trends. It assigns points to describe the spatial locations of cells, and gene

expression levels are described by marks on each point. SpatialDE [9] constructs

multidimensional Gaussian distribution for normalized gene expression. MERINGUE

[10] is based on spatial autocorrelation and cross-correlation for normalized gene

expression. Giotto [11] is based on enrichment analysis of spatial network neighbors

in binarized high gene expression state. Without normalization, SPARK [12] uses a

generalized linear spatial model with a series of custom spatial kernel functions to

describe the raw count data using Poisson distribution. However, these existing

methods still have limitations: 1) Most of them are based on normalized gene

expression data, thus would fail to consider the variance in raw counts. For example,

Trendsceek focuses on modeling two points in space; Giotto and MERINGUE focus

on modeling spatial neighbors. These algorithms assume that differences between

neighbors are comparable, requiring analysis on the normalized data. 2) Majority of

previous methods are based on certain parametric models that will limit their abilities

to identify various possible spatial distribution patterns when the parametric

assumption is violated. For example, SpatialDE assumes that the data obeys Gaussian

distribution; and SPARK requires settings of specific spatial kernel functions. 3) They

generally output a significant score to rank genes, however low significant scores, e.g.

P or Q values, do not necessarily mean real spatial patterns [13].

In this work, we propose nonparametric **S**patial **P**attern **R**ecognition using

Information based method, SPRI, which models raw count data directly without

restrictive model assumptions to identify spatial gene expression patterns. SPRI firstly

converts the spatial gene pattern problem into an association detection problem

between (x, y) coordinate values with observed raw count data, and then estimates

associations using an information-based method, TIC (total information coefficient)

[14, 15], which calculates the total mutual information with all possible x-y-grids.

Without assuming that data is generated from a parametric model, SPRI can detect

more types of SE gene patterns with higher accuracy.

Results

Overview of SPRI and results on the simulation dataset.

Fig.1 | Overview of the SPRI method and results on the simulation dataset. a,

Overview of SPRI algorithm. SPRI converts the spatial gene pattern problem to

association detection problem between coordinates values of (x, y) with read count of

each gene as the observed frequencies and then calculates their TIC using all possible

x-y-grids. Background correction was used to remove the effect of the shape of the

tissue. Then permutation test was used to identify statistically significant SE genes. **b**,

Proportion of true positives for all four different simulation patterns, including

Hotspot, Streak, Step gradient and Linear gradient, which display the proportion of

true positive samples among genes (*Y*-axis) detected by the compared methods at

different FDRs (*X*-axis).

The overview of SPRI is shown in **Fig. 1a**. SPRI firstly converts the spatial gene

pattern problem to association detection problem between coordinates values of (x, y)

with read count of each gene as the observed frequencies and then calculates their TIC

using all possible x-y-grids. Background correction was used to remove the effect of

the shape of the tissue. Then permutation test was used to identify statistically

significant SE genes. Different from existing approaches of Trendsceek, SpatialDE,

MERINGUE and Giotto, which are based on normalized gene expression data with

assumption that the sum of RNA transcripts of each cell is equal, SPRI directly

models the raw count data. Unlike SPARK, which is based on statistical hypothesis of

Gaussian distribution and certain spatial gene pattern kernel assumptions, SPRI

converts the spatial gene pattern problem to association detection problem between

coordinates values of (x, y) using observed count data as observed frequencies, and it

then estimates the association using the information-based approach TIC to calculate

the total mutual information with all possible x-y-grids. To evaluate the performance

of SPRI, we compared it with these five recently developed methods with simulations,

including SPARK, SpatialDE, Trendsceek, MERINGUE and Giotto on four simulated

patterns (**Fig. 1b, Supplementary Fig. 1**). Without restrictive assumptions about data

distribution, simulation results indicate that SPRI can identify higher proportion of

true positives while controlling false discovery rate.

Following comparison strategy in ref. [8, 12], the simulated patterns are set as

Hotspot, Streak, Step gradient and Linear gradient respectively. To explore the

robustness of these methods, we also tried different parameters for the simulation data

and the variation of performance is shown as the shaded area in **Fig 1b**. See the

details in **Supplementary Notes 1**. As can be seen, for all four simulation patterns,

SPRI outperforms all other previous methods on the task of identifying SE genes.

Among these methods, SPARK, MERINGUE and Giotto are the second best followed

by SpatialDE and Trendsceek on different simulated patterns respectively, which is

consistent with previous studies [10-12]. More results of different tissue shape can be

seen in **Supplementary Fig. 2.** The spatial coordinates of real tissues were adopted,

and real gene read counts were used to generate the same four simulated patterns.

Different-BG methods were used for background correction. SPRI can also identify

higher proportion of true positives for real tissue shape while controlling false

discovery rate.

Human breast cancer data

Fig.2 | Results of the human breast cancer data (Breast Cancer Layer 2) with

Different-BG. a, Line plot that displays the number of genes with SE patterns (*Y*-axis)

detected by six different methods at different FDRs (*X*-axis), respectively. **b,** Boxplot

of expression levels of SE genes identified by SPRI, SPARK, SpatialDE,

MERINGUE and Giotto in Breast Cancer Layer 2 data. **c,** Percentage (pink/blue) of

SPRI/SPARK SE genes that are overlapped with SPARK/SPRI top-ranked SE genes.

d, Enrichment of SE genes in the breast cancer marker genes provided by CancerMine

database. The significant level of enrichment was quantified by Fisher's exact test. **e,**

Visualization of gene spatial expression patterns for five genes that are detected by

SPRI only. The first one is hematoxylin & eosin stained brightfield image of Breast

Cancer Layer 2 from ref. [16], with dark staining indicates potential region of tumor. **f,**

GO enrichment analysis on top 100 SE genes by SPRI, SPARK, SpatialDE,

MERINGUE and Giotto respectively. **g**, Bubble plot of enriched GO terms and

KEGG pathways (purple) on the whole SPRI SE genes at FDR = 0.05.

We first analyzed a human breast cancer data (breast cancer layer 2) [16]

generated by the ST platform, which has 14,789 genes measured on 250 spots. The

resolution of this dataset is 100 mm. **Fig. 2** shows the result for breast cancer layer 2.

Permutation test was performed to test the significance of the genes and then FDR

correction was used to compare with five existing methods, including SPARK,

SpatialDE, Trendsceek, MERINGUE and Giotto. Following SPARK paper, we

named the genes with FDR cutoff of 0.05 as SE genes. As can be seen in **Fig. 2a,**

SPRI can identify more SE genes. Totally, SPRI identified 580 SE genes, while

SPARK identified 290 SE genes (overlap with SPRI = 157; **Supplementary Fig. 3a**),

SpatialDE identified 115 SE genes (overlap with SPRI = 45), Trendsceek only

identified 13 SE genes (overlap with SPRI = 13), MERINGUE identified 207 SE

genes (overlap with SPRI = 142), Giotto identified 146 SE genes (overlap with SPRI

= 69).

Firstly, the comparison of expression levels for SE genes shows that SPRI can

detect more highly expressed genes than existing methods. As shown in **Fig. 2b**, the

expression level uniquely detected by SpatialDE was very low, and the level uniquely

detected by SPARK, MERINGUE and Giotto are comparable. In contrast, the

expression level of SE genes identified only by SPRI is the closest to that of SE genes

found by all five methods, which have the highest expression level. Secondly, the

proportion of top SPARK-ranked SE genes that were also identified by SPRI as SE

genes and the proportion of top SPRI-ranked genes that were also identified by

SPARK as SE genes were compared (**Fig. 2c**). The results showed that SE genes

identified by SPRI can cover more top SPARK ranked genes. The comparison with

other methods can be found in **Supplementary Fig. 3d**. Thirdly, to further validate

our method, we compared the SE genes identified by SPRI with known marker genes

related to human breast cancer which was downloaded from CancerMine database

[17]. As shown in **Fig. 2d**, SPRI demonstrates higher enrichment of known marker

genes than other methods. Top SPRI-ranked genes only identified by SPRI were also

listed (**Fig. 2e**) to visually evaluate the spatial patterns of SE genes detected by SPRI.

These genes are clearly distributed in candidate tumor regions.

We next explore the biological insights of the SE genes found by SPRI. Manual

inspection of the top five SE genes uniquely identified by SPRI and other methods

(**Supplementary Fig. 4)** indicates that SPRI genes are more spatially variable, and

four of them are found associated with breast cancer, supported by literature,

including *TMSB10, PABPC1, ACTG1* and *RPS27A. TMSB10* was upregulated in

breast cancer tissues and its overexpression promotes invasion, proliferation and

migration of breast cancer cells [18]; The *PABPC1* gene was a downstream target of

SNHG14 and mediates *SNHG14-induced* oncogenesis in breast cancer [19]; *ACTG1*, a

cytoskeletal protein, is thought to be a component of the cell migration machinery,

and when destabilized is able to inhibit the migration of cancer cells [20]; RPS27A is

reported to be overexpressed in breast cancers [21].

In addition, functional enrichment analysis of SE genes detected by SPRI,

SPARK, SpatialDE, MERINGUE and Giotto was also performed (**Methods**). We

firstly compared the top 10 Gene Ontology (GO) terms found by these five methods

for the same number of genes (top 100 (**Fig. 2f**), top 150 and 200 in **Supplementary**

Fig. 3e). The result indicates that SPRI obtains much more significant GO terms than

other methods.

Next, functional enrichment analysis was performed on all SE genes (**Fig. 2g,**

Supplementary Fig. 3b and Fig. 3c). Totally, 690 GO terms and 32 Kyoto

Encyclopedia of Genes and Genomes (KEGG) pathways were enriched in the SE

genes identified by SPRI, while SPARK had 532 enriched GO terms and 32 KEGG

pathways, SpatialDE had only 192 enriched GO terms and 6 KEGG pathways,

MERINGUE had 239 enriched GO terms and 9 KEGG pathways and Giotto had 281

enriched GO terms and 11 KEGG pathways. The results show that many enriched GO

terms detected by SPRI only are associated with immune responses, such as activation

of immune response (GO:0002253; SPRI P value= 1.58×10^{-5}, Giotto P value= 0.019,

while SPARK, SpatialDE and MERINGUE did not have this enriched GO term). In

addition, some KEGG pathways identified by SPRI only are directly relevant to

cancer, such as Proteoglycans in cancer (hsa05205; SPRI P value = 3.15×10^{-8},

SPARK P value = 1.46×10^{-3}, SpatialDE P value = 1.83×10^{-2}, Giotto P value = $2.92\times$

10^{-4}, while MERINGUE did not have this enriched KEGG pathways). The results of

alternative background filling methods Statistical-mean-BG can be found in the

Supplementary materials (**Supplementary Fig. 5, 6, 7**).

LIBD human dorsolateral prefrontal cortex (DLFPC) data

Fig.3 | Results of the DLPFC data (slice 151507) with Different-BG. a, Line plot

that displays the number of genes with SE patterns (*Y*-axis) detected by six different

methods at different FDRs (*X*-axis), respectively. **b**, Enrichment of SE genes in the

"Allen Brain Atlas Adult Human Brain Tissue gene expression profiles" gene list of

Harmonizome database [5]. **c,** Percentage (pink/blue) of SPRI/SPARK SE genes that

are overlapped in SPARK/SPRI top-ranked SE genes. **d**, Boxplot of expression levels

of SE genes identified by SPRI, SPARK, SpatialDE and Giotto in tissue slice 151507.

e, Visualization of gene spatial expression patterns for top four genes that are detected

by SPRI. **f**, GO enrichment analysis on top 500 SE genes by SPRI, SPARK,

SpatialDE and Giotto respectively. **g**, Bubble plot of enriched GO terms and KEGG

pathways (purple) on the whole SPRI SE genes at FDR = 0.05.

To show the generalization ability of SPRI to more complex tissues of higher

resolution, LIBD human dorsolateral prefrontal cortex 10x Visium (DLPFC) data was

analyzed. The datasets generated by 10x Visium platform with a resolution of 55 μm.

In this study, four slices of human brain across six neuronal layers and white matter of

the DLPFC were analyzed. **Fig. 3** shows the result for slice 151507, which has 33,538

genes measured on 4,226 spots. **Fig. 3a** shows that for tissue slice 151507, SPRI

identified more SE genes than all the five compared methods. Totally, SPRI identified

3,893 SE genes, while SPARK identified 1,488 SE genes (overlap with SPRI = 737;

Supplementary Fig. 8a), SpatialDE identified 1,258 SE genes (overlap with SPRI =

1,795), Trendsceek and MERINGUE did not find any SE genes, Giotto identified

1,104 SE genes (overlap with SPRI = 745). We also compared the SE genes identified

by SPRI with the "Allen Brain Atlas Adult Human Brain Tissue gene expression

profiles" gene list in Harmonizome database [5] (**Fig. 3b**). SPRI demonstrates higher

enrichment than other methods. Similar to previous analysis, SPRI can recover most

top ranked SE genes identified by other methods (**Fig. 3c, Supplementary Fig. 8d**)

and detect more highly expressed SE genes (**Fig. 3d**).

After visualization of SE genes detected by SPRI only (**Fig. 3e**), we evaluate the

biological insights found by SPRI. Manual inspection of the top five SE genes

uniquely identified by SPRI (**Supplementary Fig.9)** indicates that three of them have

been found associated with brain activity, including *PEBP1*, *PSAP,* and *GPX4*. For

instance, studies have shown that downregulation of PEBP1 leads to Alzheimer's

disease [22]; studies have shown that PSAP plays an important role in neurotrophy

and neuroprotection as a multifunctional protein and that deletion of saposin leads to

age-dependent neurodegenerative diseases [23]; Gpx4 is essential for maintaining

mitochondrial integrity and neuronal survival, and deficiency of Gpx4 leads to

neurodegeneration in brain [24]. For functional enrichment analysis, the top 10 GO

terms found by SPRI, SPARK, SpatialDE, Trendsceek and Giotto for the same

number of genes (top 100**,** 200, 300, 400 in **Supplementary Fig. 8e** and top 500 in

Fig. 3f) were compared firstly. Then, functional enrichment analyses were performed

on all SE genes for SPRI (**Fig. 3g**), SPARK, SpatialDE and Giotto (**Supplementary**

Fig. 8b, c). SPRI has more significant GO terms than other methods, and these

enriched GO terms and KEGG pathway shows highly associations with brain

activities and neural diseases.

The same analysis was applied to tissue slice 151671, 151672 and 151676

(**Supplementary Fig. 14, 21, 24**). Still, SPRI is found to be able to identify more SE

genes (**Supplementary Fig. 14a, 21a, 24a**) at FDR=0.05. SPRI also demonstrates a

higher enrichment score than other methods in the same human brain marker gene list.

(**Supplementary Fig. 14b, 21b, 24b**). Further analysis shows that SPRI can recover

most top ranked SE genes identified by other methods (**Supplementary Fig. 14c, 21c,**

24c) and is capable of detecting more highly expressed SE genes (**Supplementary**

Fig. 14d, 21d, 24d).

For functional enrichment analyses, top 10 GO terms for the same number of

genes (top 500; **Supplementary Fig. 14f, 21f, 24f**) were compared. Then, functional

enrichment analyses were performed on all SE genes (**Supplementary Fig. 14g, 21g,**

24g). All the above results suggest that SE genes identified by SPRI are significantly

enriched in more GO terms than those of other methods. The results of the other

background filling methods, i.e., Statistical-mean-BG and no background correction

can be found in the Supplementary materials (**Supplementary Fig.10, 11, 12, 13**).

Mouse olfactory bulb data

Fig.4 | Results of the mouse olfactory bulb data (Stereo-seq) with Different-BG. a,

Line plot that displays the number of genes with SE patterns (*Y*-axis) detected by six

different methods respectively. **b,** Enrichment of SE genes that are verified in the

study of Tepe et al.[25]. **c,** Percentage (pink/blue) of SPRI/SPARK SE genes that are

verified in SPARK/SPRI top-ranked SE genes. **d,** Expression levels of top SE genes

identified by top SPRI, SPARK, Giotto genes and union set of the three. **e,**

Visualization of gene spatial expression patterns from Stereo-seq dataset for

representative marker genes. **f,** *In situ* hybridization obtained from the Mouse Brain

Atlas of the Allen Brain Atlas for three marker genes (*Cck*, *Ptgds*, and *Mbp*). **g,** GO

enrichment analysis on top 100 SE genes by SPRI, SPARK, SpatialDE and Giotto

respectively. **h,** Bubble plot of enriched GO terms and KEGG pathways (purple) on

the whole SPRI SE genes.

We also identified genes with spatial expression patterns on higher resolution

mouse olfactory bulb (MOB) data. Stereo-seq is a newly published spatial technique

that achieves subcellular spatial resolution. The data used in this work were binned

into cellular level resolution (~14 μm). **Fig. 4** shows the result for mouse olfactory

bulb of Stereo-seq [2], which has 27,106 genes measured on 19,527 cells. SPRI can

identify more SE genes (**Fig. 4a**). Totally, SPRI identified 525 genes, while SPARK

identified 377 genes (overlap with SPRI = 326; **Supplementary Fig. 27a**), SpatialDE,

Trendsceek and MERINGUE did not identify any SE genes, and Giotto identified 88

genes (overlap with SPRI = 30).

We also compared the SE genes identified by SPRI with a cell type-specific

marker gene list which was downloaded from a recent single-cell RNA sequencing

research of olfactory bulbs [25], to further validate our method (**Fig. 4b**). Secondly,

SPRI can recover most top SPARK SE genes (**Fig. 4c**), and more results can be found

in **Supplementary Fig. 27d**. Still, the expression comparison of SE genes shows that

SPRI can detect more highly expressed genes than existing methods (**Fig. 4d**).

To visually evaluate the SE genes detected by SPRI, we also clustered the SE

genes identified by SPRI and obtained seven major spatial patterns (**Supplementary**

Fig. 27e). The three patterns correspond to three cell layers of mouse olfactory layer

respectively: glomerular layer (pattern IV), mitral cell layer (pattern V), and the

granular cell layer (pattern VI). Six marker genes were selected to visualize these

three spatial patterns (**Fig. 4e**). The *in situ* hybridization images from the Allen Brain

Atlas further cross-validated these genes exhibiting spatial expression patterns (**Fig.**

4f).

We next explore the biological insights found by SPRI. Manual inspection of the

top five SE genes uniquely identified by SPRI (**Supplementary Fig. 28**) indicates

that four of them are found associated with brain functions, including *Ckb*, *Cadm2*,

Hsp90ab1, *Cox6c*. For example, *Ckb* is reported to be upregulated in the Alzheimer's

disease brain and is important for cellular energetics [26]. *Cadm2* is significantly

downregulated in human glioma tissue and may be a new potential therapeutic target

for human gliomas [27]. *Hsp90* family chaperones, particularly *Hsp90ab1*, is closely

associated with astrocytes, suggesting it's possible association with Alzheimer's

disease mediated by the accumulation of intracellular hyperphosphorylated tau

proteins and extracellular amyloid β peptide [28]. The decreased expression and

activity of Cox6c in the temporal and parietal cortex of patients with Alzheimer's

disease suggests that it may be a potential biomarker for the diagnosis of brain

diseases [29]. In addition, functional enrichment analyses were performed. We firstly

compared the top 10 Gene Ontology (GO) terms found by these methods for the same

number of genes (**Fig. 4g**; **Supplementary Fig. 27g**), which indicates that SPRI

obtains more significant GO terms than other methods. Then, functional enrichment

analyses were performed on all SE genes (**Fig. 4h**, **Supplementary Figs. 27b** and **c**).

The results of the other background filling methods, i.e., Statistical-mean-BG and no

background correction can be found in the Supplementary materials (**Supplementary**

Figs. 29, 30).

Adult mouse brain data

Fig.5 | Results of the Adult mouse brain data with Different-BG. a, Line plot that

displays the number of genes with SE patterns (*Y*-axis) detected by six different

methods at different FDRs (*X*-axis), respectively. **b**, Enrichment of SE genes that are

verified in the "Allen Brain Atlas Adult Mouse Brain Tissue gene expression profiles"

gene list of Harmonizome database. **c,** Percentage (pink/blue) of SPRI/SPARK SE

genes that are verified in SPARK/SPRI top-ranked SE genes. **d**, Boxplot of

expression levels of SE genes identified only by SPRI, SPARK, SpatialDE and Giotto

in Adult Mouse Brain data. **e**, Visualization of gene spatial expression patterns for six

genes that are detected by SPRI. **f**, GO enrichment analysis on top 500 SE genes by

SPRI, SPARK, SpatialDE, Trendsceek and Giotto respectively. **g**, Bubble plot of

enriched GO terms and KEGG pathways (purple) on the whole SPRI SE genes.

In order to validate the applicability of SPRI on sections containing more

biologically complex tissues, whole brain data were analyzed. For Adult Mouse Brain

from the 10x genomics platform which has 19,565 genes measured on 2,264 spots,

SPRI can identify more SE genes (**Fig. 5a**). And SE genes identified by SPRI have

high enrichment in the "Allen Brain Atlas Adult Mouse Brain Tissue gene expression

profiles" marker gene list from Harmonizome database (**Fig. 5b**). In addition, SPRI

can recover most top ranked SE genes on most cases (**Fig. 5c, Supplementary Fig.**

31d) and detect more highly expressed SE genes (**Fig. 5d**). After visualization of SE

genes detected by SPRI (**Fig. 5e**), we evaluate the biological insights found by SPRI.

Manual inspection of the top five SE genes uniquely identified by SPRI

(**Supplementary Fig. 32)** indicates that four of them have been found associated with

brain activity, including *Eif3c*, *Ctbp1*, *Mff* and *Ube2d3*. Loss of function of the *Eif3c*

subunit leads to neural tube defects and affects the development of the nervous system

[30]. *Ctbp1* helps *Fbxo32* promote epithelial–mesenchymal transition during brain

development [31]. Loss of *Mff* leads to disturbances in mitochondrial and peroxisome

dynamics, ultimately causing peripheral neuropathy [32]. *Ube2d3* promotes glioma

proliferation by regulating the STAT3 signaling pathway [33]. For functional

enrichment analyses, top 10 GO terms found by SPRI, SPARK, SpatialDE,

Trendsceek and Giotto for the same number of genes (top 100, 200, 300, 400 in

Supplementary Fig. 31e and top 500 in **Fig. 5f**) were compared firstly. Then,

functional enrichment analyses were performed on all SE genes at 0.05 FDR cutoff

for SPRI (**Fig. 5g**), SPARK, SpatialDE, Trendsceek and Giotto (**Supplementary**

Fig.31b and Fig. 31c). The results showed that SPRI SE genes have much higher

significant GO terms correlated with brain structure.

Discussion and conclusion

The recent rapid development of high throughput spatially resolved

transcriptomics technology opens a new door to understand the spatial resolved

biological behaviors of genes and cells. One essential and initial step of such analysis

is to detect genes with spatial expression patterns.

In this work, we propose a novel information-based spatial pattern gene

identification method, SPRI, to model spatial raw count data directly. It converts the

SE gene detection problem to a dependencies mining problem between spatial

coordinate pairs with raw gene read count as the observed frequencies. Such strategy

distinguishes SPRI from existing SE methods relying on certain model or data

assumptions. For example, methods based on normalized data assume implicitly that

the total number of RNA transcripts in each cell is identical, which is not always true

[34]. Other methods modeling raw count, like SPARK, also rely on certain parametric

statistical models or specific kernel functions, which still limits the ability to identify

various possible spatial distribution patterns.

To evaluate SPRI's performance, we compared it with five existing methods on

seven publicly available datasets comprehensively. The results consistently indicate

that SPRI can robustly identify more genes with true spatial expression patterns

validated by *In situ* hybridization experiments, and that SE genes identified by SPRI

are more spatially variable and some of them are supported by recent studies.

Methods

SPRI: model and algorithm

In SPRI, firstly we convert the problem of identifying genes with spatial

expression patterns into the problem of identifying dependencies on (X, Y) coordinate

observations based on the raw count expression of genes. To encapsulate the

relationship between two coordinate variables, observed data is partitioned by

drawing a grid on the scatter plot, and all the possible grids up to a maximum grid

resolution were explored to find the one that best encapsulates this relationship.

Specifically, we use a dynamic programming algorithm **(Supplementary Notes 2)** to

detect the highest mutual information (MI) achievable using a grid with k columns

and l rows. The mutual information under different grid divisions is normalized to

enable fair comparisons in a characteristic matrix. TIC is the sum of the characteristic

matrix. Background correction was applied to remove the effect of the shape of the

tissue. Next, genes with spatial pattern were identified by permutation test. A gene is

determined as spatially variable if there is significant difference between the observed

TIC value and the TIC values after shuffling the read count on spatial spots.

Computing the TIC for each gene.

TIC [15] belongs to a family of maximal information-based nonparametric

exploratory statistics (MINE) [14]. To calculate the statistics in the MINE family for a

set of two jointly observed variables, MINE explores all possible k-by-l grids and

calculates the maximum possible mutual information for each k-by-l combination. In

information theory, MI quantifies the mutual dependence between two random

variables, since it measures the amount of information about one random variable that

can be obtained by observing another one variable. Specifically, given a set of two

jointly observed data (x, y) for variables (X, Y), the MI $I\big((X, Y), k, l\big)$ is computed

under all k-by-l grids:

$$I\big((X, Y), k, l\big) = \sum_{y \in Y} \sum_{x \in X} p(x, y) \log \left(\frac{p(x,y)}{p(x)p(y)} \right) \tag{1}.$$

where $p(x, y)$ is the joint probability of x and y, and $p(x)$ and $p(y)$ are the marginal

probability of x and y respectively, and the number of grids $kl < B$, where B is an

upper limit on the grid size to be searched. In other words, $I(k,l)$ denotes the MI of the

probability distribution estimated by the k-by-l grids, where the probability of a bin is

proportional to the number of points that fall in the grid.

For each k-by-l grid, the maximum mutual information is retained:

$$I^*\big((X,Y),k,l\big) = \max_{G \in G(k,l)} I\big((X,Y)|_G\big) \qquad (2).$$

For a fair comparison under different grid divisions, the maximum mutual information

$m_{k,l}$ under each grid G are normalized to between 0 and 1, constituting characteristic

matrix $M = (m_{k,l})$.

$$\widehat{M}(D)_{k,l} = \frac{I^*(D,k,l)}{\log \min\{k,l\}} \qquad (3).$$

That is, $m_{k,l}$ is the largest normalized mutual information achieved by all grids

with size of k-by-l. TIC is the sum of all entries in characteristic matrix M. TIC is able

to obtain a small bias and promising power by summing over all entries in the

independent test. In other words, TIC captures the presence or absence of

dependencies between two variables. The definition of TIC is following:

$$TIC(D) = \sum_{kl \le B(n)} \widehat{M}(D)_{k,l} \qquad (4).$$

The choice of the right B is important: setting B too large will result in non-zero

scores even for independent data, since each data points has its own cell, while setting

B too small will lose power in detecting dependency. According to the theoretical

analysis and experiments of [14], a default settings $B = n^{\alpha}$ was adopted where n is the

number of samples and α is 0.6.

We apply TIC to determine whether there is a dependency relationship between

two-variable (X, Y) to identify genes with spatial expression patterns, and the higher

the TIC value, the more likely the gene has a spatial expression pattern. By using a

permutation test (shown below), we could obtain a *P* value for each gene. We defined

a SE gene with FDR < 0.05 and ranked them based on TIC values.

TIC can be estimated in linear time using TIC_e algorithm (**Supplementary Notes**

2) with α set as 0.4. The most time-consuming step in calculating TIC is the

construction of the characteristic matrix. After calculating the characteristic matrix,

the time to calculate the TIC is trivial. The ideal characteristic matrix *M* is optimized

for all possible grids. For computational efficiency, APPROX-TIC [14] algorithm

(**Supplementary Notes 2**) was applied in practice. APPROX-TIC using a dynamic

programming algorithm that optimizes on a subset of the possible grids to calculate

the heuristic approximation of the characteristic matrix. The runtime complexity of

the search procedure for the APPROX-TIC algorithm used in practice to calculate the

approximate MIC is $O(n^{4\alpha})$. For large data sets, TIC_e can further reduce the

computation time and achieve comparable results by finding only the optimal

one-dimensional partitions with a time complexity of $O(n^{5\alpha/2})$ [35]. In this work, we

use the APPROX-TIC for all dataset analysis, with α is set as 0.6. For larger data like

10x Visium dataset, additional experiments were also performed using TIC_e, which

can achieve comparable results (**Supplementary Fig. 36**) with a lower running time.

Background correction.

For genes without spatial pattern, TIC can still be nonzero because of the

boundary of the sample. To remove the effect of the shape of the tissue, we added

background spot locations (**Fig. 1a**) based on its original spatial coordinates. There

are two ways to fill the values of the background coordinates which we called

Different-BG and Statistical-mean-BG respectively. Different-BG randomly samples

count from each gene's read count distribution for each background spot

independently. For human breast cancer data, the expression of background was

randomly sampled from the entire distribution of read count of all genes due to the

very small number of spots. The Statistical-mean-BG is set as the trimean value [36]

of the gene over all spots as following:

$$mean = \frac{1}{2}Q_2 + \frac{1}{4}(Q_1 + Q_3)$$ (5).

where Q_1, Q_2, and Q_3 are the first, second and third quartiles of current gene

expression respectively. Trimean is less sensitive to outliers compared to a simple

average.

Identifying statistically significant SE genes with permutation test.

We defined background distribution as the distribution of the gene expression not

being associated with the spatial coordinates; in other words, the read count of a gene

is independently and identically distributed over the grids. In our method, a gene is

determined as SE genes if there is significant difference between the observed TIC

value and the TIC values under background distribution. We shuffled the read counts

on spatial spots to remove any spatial dependency while keeping the marginal

distribution of read counts intact. Specifically, in the permutation test, we keep the

spot locations fixed, randomly shuffle gene count expression, and then recalculate the

TIC values. The *P* value of each gene is computed as following:

$$P\ value = \frac{\#\{i|TIC_i > TIC_{obs}, i=1,2,3\dots,N\}}{N}$$ (6).

where TIC_i is the TIC value for the *i-th* permutation, and TIC_{obs} is the original TIC

value of gene expression; *N* is the number of total permutation times, we set *N* to

10,000 for small ST data and 1,000 for other datasets in our experiment. After getting

the *P*-value for each gene, we used Benjamini-Yekutieli approach in the python

package of "statsmodels" to control FDR. The genes with FDR < 0.05 are considered

as significant SE genes.

Gene sets, cluster, visualization and functional enrichment analysis

For these SRT datasets, we downloaded lists of known genes to validate the SE

genes recognized by different methods. For the human breast cancer data, we

downloaded a list of genes associated with breast cancer from the CancerMine

database [37]. These breast cancer related genes are composed of three parts, namely

cancer drivers, oncogenes, and tumor suppressors. For the mouse olfactory bulb data,

we downloaded a list from of 2,030 cell type-specific marker genes identified in

recent single-cell RNA sequencing studies in the olfactory bulb [25]. For the DLPFC

and Adult Mouse Brain, we downloaded two brain gene lists of human and mouse

from Harmonizome database [5], the "Allen Brain Atalas Adult Human Brain Tissue

gene expression profiles" and the "Allen Brain Atalas Adult Mouse brain Tissue gene

expression profiles", respectively.

Besides, we follow the clustering algorithm in SPARK, and cluster top SE genes

of Stereo-seq dataset (FDR<0.05) identified by SPRI into seven clusters according to

manual annotation. In the visualization of SPRK, variance-stabilizing transformation

was performed on the raw count data and the log-scale total counts was adjusted to

get a relative gene-expression. We follow the visualization strategy of gene spatial

expression patterns proposed by SPARK and the raw count was directly visualized in

Supplementary Figures. Finally, the same number of top SE genes and all SE genes at

a 0.05 FDR cutoff identified by SPRI, SPARK, SpatialDE, Trendsceek, MERINGUE

and Giotto were used for functional enrichment analysis including GO terms analysis

and KEGG pathways analysis. Following the SPARK paper, we adopted the R

package of "clusterProfiler (v3.18.1)" to perform all functional enrichment analysis.

In the package, we set the P value correction method as the default 'BH' and the cutoff

of FDR as 0.05.

Data sets

In this work, seven SRT datasets of three different platform, i.e., ST, 10x Visium

and Stereo-seq, were analyzed. One ST datasets, human breast cancer data [16], were

downloaded from Spatial Research [38]. Five 10x Visium datasets include four slices

from the LIBD human dorsolateral prefrontal cortex data [39], and one adult mouse

brain data was downloaded from 10x Genomics [40]. One mouse olfactory bulb data

of Stereo-seq [2]. These SRT datasets consist of two components: the spatial locations

(spots) and the gene expression (read counts) observed at these spatial locations.

Following the SPARK paper, we filtered out spots less than ten total read counts

and selected genes expressed at least 10% of the spots. For large data Stereo-seq,

genes were selected expressed at least 8% spots. Then permutation test was performed

on all genes after filtering criteria.

Comparison of methods

We compare our method SPRI with five existing algorithms for spatial expression

pattern recognition of genes, including SPARK [12], SpatialDE [9], Trendsceek [8],

MERINGUE [10] and Giotto [11]. SpatialDE, Trendsceek, MERINGUE and Giotto

are based on normalized data and SPARK is based on raw count data.

The first method we compared with is the SPARK (R package SPARK; v1.1.0),

we directly use SPARK's code on github for analysis [41]. Following the SPARK

paper, we performed the same data preprocessing. Specifically, genes that are

expressed in less than 10% of the spots were filtered out, and only spots containing at

least ten total read counts were retained. According to the SPARK paper, if the

adjusted P value (i.e., FDR) output by SPARK is below the threshold of 0.05, the

identified SE is significant.

The second method we compared with is the SpatialDE (python package

SpatialDE; v.1.1.3), we used SpatialDE's code downloaded from github for analysis

[42]. Following the SpatialDE paper, the SE gene is considered significant if the Q

value (i.e.,FDR) was below the threshold of 0.05.

The third method we compared with is the Trendsceek (R package trendsceek;

v.1.0.0). We also used the code provided from github for analysis [43]. Following the

Trendsceek paper, we performed the same data preprocessing. Specifically, genes that

express less than three spots were filtered out, and only spots containing at least five

read counts were retained. Then the raw count data were processed through log10

transformation. For the real data, the top 500 variable genes were taken for analysis.

The SE gene is considered significant if the p.bh value (i.e.,FDR) was below the

threshold of 0.05.

The fourth method we compared with is MERINGUE (R package MERINGUE;

v.1.0). The code provided on github was used for analysis [44]. Following

MERINGUE paper, poor spots (fewer than 100 read counts) and poor genes (fewer

than 100 read counts) were filtered out. Then Benjamini–Yekutieli correction was

performed to control FDR, the SE gene is regarded significant if the FDR was below

the threshold of 0.05.

The last method we compared with is the Giotto (R package Giotto; v.1.1.0). The

code provided from github was used in this study for analysis [45]. The same data

pre-processing in Visium data was adopted. Then Benjamini-Yekutieli correction was

performed to control FDR, the SE gene is regarded as significant if the FDR was

below the threshold of 0.05.

Declarations

Availability of data and materials

Code availability

SPRI is implemented in Python. All source code is available on Github at https://github.com/xiaoyeye/SPRI, and Zenodo [46].

Data availability

All public SRT datasets can be obtained as described in "**Data sets**" and "**Gene sets, cluster, visualization and functional enrichment analysis**" section in the "**Methods**" section. Specifically, the SRT data is available from the following websites: Human breast cancer data were downloaded from (https://www.spatialresearch.org/) [38]; DLPFC data [39] were downloaded from (http://research.libd.org/spatialLIBD/); Stereo-seq data [2] were downloaded from (https://github.com/JinmiaoChenLab/ SEDR_analyses); Adult Mouse Brain (FFPE) were downloaded from (https://www.10xgenomics.com/resources/datasets/) [40]. For enrichment analysis, a marker gene list in olfactory bulb was downloaded from [25], a marker gene list related to human breast cancer was downloaded from CancerMine database (http://bionlp.bcgsc.ca/cancermine/) [37], two brain gene lists were downloaded from Harmonizome database (https://amp.pharm.mssm.edu/Harmonizome/) [5].

Authors' contributions

Y.Y. and H.B. Shen designed the research, J.X. Hu and Y.Y. performed the research,

J.X. Hu and Y.Y. analyzed the data, and J.X. Hu, Z.R. Hu, Y.Y. and H.B. Shen wrote

the paper. The authors read and approved the final manuscript.

Funding

This work was supported by the National Natural Science Foundation of China (No.

61725302, 62073219, 62103262 to Y.Y.), Shanghai Pujiang Program (No.

21PJ1407700 to Y.Y.), and the Science and Technology Commission of Shanghai

Municipality (No. 22511104100).

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1.   Ji, A.L., et al., *Multimodal Analysis of Composition and Spatial Architecture in Human*

Squamous Cell Carcinoma. Cell, 2020.

2.   Chen, A., et al., *Large field of view-spatially resolved transcriptomics at nanoscale*

resolution. BioRxiv, 2021.

3.   Larsson, L., J. Frisén, and J. Lundeberg, *Spatially resolved transcriptomics adds a*

new dimension to genomics. Nature Methods, 2021. **18**(1): p. 15-18.

4.   Zhuang, X., *Spatially resolved single-cell genomics and transcriptomics by imaging.*

Nature Methods, 2021. **18**(1): p. 18-22.

5.   *Database - Harmonizome [cited 2022 May 5].* Available from:

https://amp.pharm.mssm.edu/Harmonizome/.

6.   Berglund, E., et al., *Spatial maps of prostate cancer transcriptomes reveal an*

unexplored landscape of heterogeneity. Nature communications, 2018. **9**(1): p. 1-13.

7.   Ji, A.L., et al., *Multimodal analysis of composition and spatial architecture in human*

squamous cell carcinoma. Cell, 2020. **182**(2): p. 497-514. e22.

8.   Edsgärd, D., P. Johnsson, and R. Sandberg, *Identification of spatial expression trends*

in single-cell gene expression data. Nature methods, 2018. **15**(5): p. 339-342.

9.   Svensson, V., S.A. Teichmann, and O. Stegle, *SpatialDE: identification of spatially*

variable genes. Nature methods, 2018. **15**(5): p. 343-346.

10.   Miller, B.F., et al., *Characterizing spatial gene expression heterogeneity in spatially*

resolved single-cell transcriptomic data with nonuniform cellular densities. Genome

research, 2021. **31**(10): p. 1843-1855.

11.   Dries, R., et al., *Giotto: a toolbox for integrative analysis and visualization of spatial*

630   *expression data.* Genome biology, 2021. **22**(1): p. 1-31.

631   12.   Sun, S., J. Zhu, and X. Zhou, *Statistical analysis of spatial expression patterns for*

632   *spatially resolved transcriptomic studies.* Nature methods, 2020. **17**(2): p. 193-200.

633   13.   Hu, J., et al., *SpaGCN: Integrating gene expression, spatial location and histology to*

634   *identify spatial domains and spatially variable genes by graph convolutional network.*

635   Nature methods, 2021: p. 1-10.

636   14.   Reshef, D.N., et al., *Detecting novel associations in large data sets.* science, 2011.

637   **334**(6062): p. 1518-1524.

638   15.   Reshef, Y.A., et al., *Measuring dependence powerfully and equitably.* The Journal of

639   Machine Learning Research, 2016. **17**(1): p. 7406-7468.

640   16.   Ståhl, P.L., et al., *Visualization and analysis of gene expression in tissue sections by*

641   *spatial transcriptomics.* Science, 2016. **353**(6294): p. 78-82.

642   17.   Lever, J., et al., *CancerMine: a literature-mined resource for drivers, oncogenes and*

643   *tumor suppressors in cancer.* Nature methods, 2019. **16**(6): p. 505-507.

644   18.   Zhang, X., et al., *Thymosin beta 10 is a key regulator of tumorigenesis and metastasis*

645   *and a novel serum marker in breast cancer.* Breast Cancer Research, 2017. **19**(1): p.

646   1-15.

647   19.   Dong, H., et al., *Long non-coding RNA SNHG14 induces trastuzumab resistance of*

648   *breast cancer via regulating PABPC1 expression through H3K27 acetylation.* Journal

649   of cellular and molecular medicine, 2018. **22**(10): p. 4935-4947.

650   20.   Luo, Y., et al., *Loss of ASAP3 destabilizes cytoskeletal protein ACTG1 to suppress*

651   *cancer cell migration.* Molecular medicine reports, 2014. **9**(2): p. 387-394.

21. Hong, S.W., et al., *RPS27a enhances EBV-encoded LMP1-mediated proliferation and*

invasion by stabilizing of LMP1. Biochemical and Biophysical Research

Communications, 2017. **491**(2).

22. Schoentgen, F. and S. Jonic, *PEBP1/RKIP behavior: A mirror of actin-membrane*

organization. Cellular and Molecular Life Sciences, 2020. **77**(5): p. 859-874.

23. Lin, Z.-H., et al., *PSAP intronic variants around saposin D domain and Parkinson's*

disease. Brain, 2021. **144**(1): p. e3-e3.

24. Yoo, S.-E., et al., *Gpx4 ablation in adult mice results in a lethal phenotype*

accompanied by neuronal loss in brain. Free Radical Biology and Medicine, 2012.

52(9): p. 1820-1827.

25. Tepe, B., et al., *Single-cell RNA-seq of mouse olfactory bulb reveals cellular*

heterogeneity and activity-dependent molecular census of adult-born neurons. Cell

reports, 2018. **25**(10): p. 2689-2703. e3.

26. Khan, A.T., et al., *Alzheimer's disease: are blood and brain markers related? A*

systematic review. Annals of clinical and translational neurology, 2016. **3**(6): p.

455-462.

27. Liu, N., et al., *CADM2 inhibits human glioma proliferation, migration and invasion.*

Oncology Reports, 2019. **41**(4): p. 2273-2280.

28. Gonzalez-Rodriguez, M., et al., *Neurodegeneration and Astrogliosis in the Human*

CA1 Hippocampal Subfield Are Related to hsp90ab1 and bag3 in Alzheimer's Disease.

International journal of molecular sciences, 2021. **23**(1): p. 165.

29. Wang, C., et al., *Novel role of COX6c in the regulation of oxidative phosphorylation*

and diseases. Cell death discovery, 2022. **8**(1): p. 1-8.

30.   Fujii, K., et al., *Controlling tissue patterning by translational regulation of signaling*

transcripts through the core translation factor eIF3c. Developmental Cell, 2021. **56**(21):

p. 2928-2937. e9.

31.   Sahu, S.K., et al., *FBXO32 promotes microenvironment underlying*

epithelial-mesenchymal transition via CtBP1 during tumour metastasis and brain

development. Nature communications, 2017. **8**(1): p. 1-18.

32.   Koch, J., et al., *Disturbed mitochondrial and peroxisomal dynamics due to loss of MFF*

causes Leigh-like encephalopathy, optic atrophy and peripheral neuropathy. Journal

of medical genetics, 2016. **53**(4): p. 270-278.

33.   Pan, Z., et al., *UBE2D3 Activates SHP-2 Ubiquitination to Promote Glycolysis and*

Proliferation of Glioma via Regulating STAT3 Signaling Pathway. Frontiers in

oncology, 2021. **11**: p. 2081.

34.   Marinov, G.K., et al., *From single-cell to cell-pool transcriptomes: stochasticity in gene*

expression and RNA splicing. Genome research, 2014. **24**(3): p. 496-510.

35.   Reshef, D.N., et al., *An empirical study of leading measures of dependence.* arXiv

preprint arXiv:1505.02214, 2015.

36.   Jin, S., et al., *Inference and analysis of cell-cell communication using CellChat.* Nature

communications, 2021. **12**(1): p. 1-20.

37.   *Datasets - CancerMine [cited 2022 May 5].* Available from:

http://bionlp.bcgsc.ca/cancermine/.

38.   *Datasets - Spatial Research [cited 2022 May 5].* Available from:

https://www.spatialresearch.org/resources-published-datasets/.

39.   Maynard, K.R., et al., *Transcriptome-scale spatial gene expression in the human*

dorsolateral prefrontal cortex. Nature neuroscience, 2021. **24**(3): p. 425-436.

40.   *Datasets - 10x Genomics [cited 2022 May 5].* Available from:

https://www.10xgenomics.com/cn/resources/datasets.

41.   un, S., J. Zhu, and X. Zhou. *SPARK, Github; [cited 2022 May 5].* Available from:

https://github.com/xzhoulab/SPARK-Analysis.

42.   Svensson, V., S.A. Teichmann, and O. Stegle. *SpatialDE, Github; [cited 2022 May 5].*

Available from: https://github.com/Teichlab/SpatialDE.

43.   Edsgärd, D., P. Johnsson, and R. Sandberg. *Trendsceek, Github; [cited 2022 May 5].*

Available from: https://github.com/edsgard/trendsceek.

44.   Miller, B.F., et al., *MERINGUE, Github; [cited 2022 May 5].* Available from:

https://github.com/JEFworks-Lab/MERINGUE.

45.   Dries, R., et al. *Giotto, Github; [cited 2022 May 5].* Available from:

https://rubd.github.io/Giotto_site/.

46.   Hu J.X., H.Z.R., Yuan Y, Shen H.B. . *SPRI: Spatial Pattern Recognition using*

Information based method for spatial gene expression data. Available from:

https://doi.org/10.5281/zendodo.7097246.

715