Very small spontaneous structural mutation rate in green algae

Matéo Léger-Pigout[1], Marc Krasovec[1*]

[1] CNRS, Biologie Intégrative des Organismes Marins (BIOM), Observatoire Océanologique, Banyuls sur mer, 66650, France

* marc.krasovec@obs-banyuls.fr

Running title

Structural mutation rate in green algae

Key words

mutation rate, structural mutation, mutation accumulation, phytoplankton

ABSTRACT

Structural mutations led to major innovations, but our knowledge of the spontaneous structural mutation rate is very limited. We cannot have a complete view of the adaptive potential of species and new variants in a population without addressing this question. Here, we used Illumina genomic data from mutation accumulation experiments of five green algae species and measured very low structural mutation rate, between 5×10^{-12} to 2×10^{-11} mutations per site, at least ten times lower than the nucleotide mutation rate in these species. Such low rates of structural mutations may be a consequence of selection. Assuming that structural mutations have higher fitness effect than point mutations because they impact a larger portion of the genome, selection toward a small structural mutation rate may be stronger than for point mutations. A second explanation could be the rate of true positive detection of de novo structural mutations, that is globally poor and variable between methods. This highlights the necessity of long read sequencing in future mutation accumulation studies.

SIGNIFICANCE STATEMENT

Mutations are the first source of all biodiversity. Biologists tried since a century to measure the spontaneous mutation rate, but our knowledge on the structural spontaneous mutation rate, μ_{st}, is still poor. Here we measured μ_{st} in five species of phytoplankton, and found a very small rate compared to nucleotide and short insertion-deletion mutations.

INTRODUCTION

Spontaneous mutations renew the standing genetic variation and permit adaptation through natural selection. The spontaneous mutation rate μ is so one of the most important parameters to understand evolution. Since decades, biologists tried to measure it using mutation accumulation (MA) experiment (Halligan & Keightley, 2009), first with a phenotypic approach by following fitness variation of MA lines (Mukai, 1964), and then directly by whole genome sequencing (Lynch *et al.*, 2016; Katju & Bergthorsson, 2019). Our knowledge made significant progresses the last fifteen years, particularly on the nucleotide (Lynch *et al.*, 2016; Katju & Bergthorsson, 2019) and short insertion-deletion (Sung *et al.*, 2016) mutation rates, noted μ_{bs} and μ_{id}, respectively. We observe 3-fold variation between species, from ~1×10^{-11} to ~1×10^{-8} mutations per site per haploid genome in ciliates and humans. This variation is in part explained by the drift barrier hypothesis (Sung *et al.*, 2012) where large effective population size (N_e) species reaches lower μ than small N_e species. As a consequence of the mutation load, selection pushes toward a small μ, that is more efficient in large N_e where drift is limited. This hypothesis is currently the most accepted, even if other factors play on the mutation rate but with smaller effect such as GC content (Krasovec *et al.*, 2017), genome size, generation time or metabolic rate (Martin & Palumbi, 1993; Mooers & Harvey, 1994; Thomas *et al.*, 2010; Weller & Wu, 2015). Both μ_{bs} and μ_{id} depends so mostly of N_e, but our knowledge about the spontaneous structural mutation rate μ_{st} is very poor (Press *et al.*, 2019; Ho & Schaack, 2021) and available in only few models (see Table 3 of Katju and Bergthorsson 2019 for copy number variant for example). Structural mutations, defined here as duplications, large insertions-deletions, inversions or chromosome rearrangements, may have stronger phenotypic effect than point mutations because they impact a larger portion of the genome. Thus, the pressure toward a small μ_{st} should be stronger than for μ_{bs} and μ_{id}. In addition, it is impossible to have a complete view of new variants and adaptive potential of a species with limited knowledge of μ_{st}. Structural mutations led to major innovations, for example the supergenes following inversions. In birds, this process saw the emergence of different morphotypes in a same species with different colors, behaviors and feather shapes (Küpper *et al.*, 2016; Tuttle *et al.*, 2016). Here, we measured μ_{st} for five species of green algae using mutation accumulation experiment Illumina reads from previous publications (Krasovec *et al.*, 2016, 2017, 2018).

MATERIALS AND METHODES

Illumina raw reads of 150 and 300 bp paired-end were mapped to the reference genomes with bwa mem (Li & Durbin, 2010), then bam files were treated with samtools (Li *et al.*, 2009) and structural variant called with delly (Rausch *et al.*, 2012), svaba (Wala *et al.*, 2018) and lumpy (Layer *et al.*, 2014) with standard parameters (excepted ploidy at 1). For svaba, we used -n option to input the T_0 genome bam file as control. VCF files were then sorted with several thresholds to increase the rate of true de novo mutations identification: the mutated site must be covered by 10 in the concerned MA line and the T0 genome; the mapping quality of the site must be higher than 20; the variant must be present in only one MA line, expected for few variants present in two or three MA lines because of non-independent lines (in that case common mutations and generations were count as one); the variant must totalize all the coverage of the site; last, some structural variants appeared only in one MA line, but were not count as mutation because other lines had same variants but at different positions within the confidence interval of start-end positions. De novo structural mutations candidates were then checked by coverage with samtools and by PCR and Sanger sequencing when possible.

RESULTS AND DISCUSSION

False - true positive rate of structural de novo mutation detection

Twenty-six de novo structural mutation candidates were found in the 5 species with 3 softwares (Table 1), but coverage and PCR checks reduced this number to 1, given a very low true positive rate of 4%. From Sanger sequencing, the 8 tested breaks from svaba and delly were false, so we considered all other non-tested breaks false. The other detected candidates were deletions and duplications, all larger than 300 pb apart one deletion of 215 pb. According to Wala et al (Wala *et al.*, 2018), the positive rates of deletions larger than 300 bp are 85% for svaba and delly, and 95% for lumpy; for insertion/duplication, it decreases to 37%, 24% and 29%. Here, the positive rate of deletion is 20% and the 3 duplication candidates were false. The positive rate of duplications was so 0%, but with only 3 candidates this rate should be taken with caution. Such low positive rates show the difficulties to identify structural de novo mutations with good accuracy, particularly with short reads. The de novo mutation candidates were different between the three softwares, and the single true positive was found with lumpy only, which implies that some true mutations have been possibly

105   missed. Further investigations of mutation accumulation experiments coupled with long
106   reads sequencing should be the next milestone to significantly increase the detection
107   accuracy.
108

μ_{st} is more than 10 time lower than μ_{bs} and μ_{id}
The single de novo mutation validated by PCR is a deletion of 1,382 bp identified in
Ostreoccocus mediterraneus on the outlier chromosome (Yau *et al.*, 2020), a particular
chromosome with low GC content. The mutated region is intergenic and the deleted
sequence appears elsewhere in the genome (Table S1). This makes this deleted
sequence a possible repeat region resulting from transposable elements, commonly
known to be implicated in this type of mutation. This single mutation gives a very low
μ_{st} for the 5 species, between 5×10^{-12} and 2×10^{-11} structural mutations per site per
haploid genome per generation (Figure 1), ten times lower than μ_{bs} and μ_{id} in these
species. We see three explanations for such low structural mutation rate. First is the
poor sensitivity of the method detection (see above), resulting in an artificially low
structural mutation rate. Second, low structural mutation rate may come from the
mutation mechanisms themself. Several mechanisms act directly on nucleotides such
as methylation and oxidation and are known to increase μ_{bs}, making it higher than μ_{st}
or μ_{id} just because single sites have more chance to mutate. Third is selection, for
which efficiency depends of N_e estimated around few millions to 10 million in
phytoplankton species (Blanc-Mathieu *et al.*, 2017; Krasovec *et al.*, 2019, 2020). With
such small μ_{st}, we may argue that selection pushes harder toward a low μ_{st} than for μ_{bs}
and μ_{id}. This is propable assuming that structural mutations have stronger and more
frequent deleterious effects than point mutations because they change a larger part of
the genome. The genomes of species studied here are small, very compacted with few
introns and small distance between genes. For example, *O. tauri* has a genome size
of 12.5 Mb, with 81.6% of coding sequences, 82% of genes without introns and an
average distance between genes of 254 bp. In that case, any structural mutation has
high chance to touch at least one gene or its regulatory region, making almost all
structural mutations deleterious. We can hypothesis that it is less the case in larger
genomes with a longer distance between genes and high proportion of repeats and
non-coding sequences.
137
138

Table 1. List of de novo structural mutation candidates. Only the deletion of Om21b is true.

MA line	Mutation type	Software	Start	End	PCR - Sanger	Coverage
Ot20	break	delly	NC_014444.2: 184889	NC_014431.2: 783883	false	na
Om14	break	delly	BCC102-800924: 233	BCC102-1186077: 1181287	false	na
Om17 - 18	break	delly	BCC102-594986: 60163	BCC102-768578: 224081	false	na
Om1	break	delly	BCC102_ML_SOC: 68272	BCC102-397956: 160264	false	na
Om2	break	delly	BCC102-466838: 2048	BCC102-867403: 392	false	false
Ot38	deletion	delly	NC_014427.2: 430027	NC_014427.2: 438281	no check	false
Bp8	deletion	delly	NC_023992.1: 297114	NC_023992.1: 301342	false	false
Bp20	duplication	delly	NC_023992.1: 296522	NC_023992.1: 297150	false	false
Ot38	break	svaba	NC_014427.2: 820183	NC_014428.2:994946	false	na
Ot38	break	svaba	NC_014433.2: 406614	NC_014437.2: 310849	false	na
Om13	break	svaba	BCC102-873724: 213431	BCC102_ML_SOC: 100063	false	na
Ot22	break	lumpy	NC_014426.2: 181294	NC_014428: 229429	no check	na
Ot23	break	lumpy	NC_014426.2: 929056	NC_014442.2: 399570	no check	na
Ot9 - 10	break	lumpy	NC_014435.2: 273589	NC_014439.2: 385664	no check	na
Ot20 - 31	break	lumpy	NC_014441.2: 17675	NC_014443.2: 58813	no check	na
Om38	break	lumpy	BCC102-993683: 535232	BCC102-768578: 224085	no check	na
Om12	break	lumpy	BCC102-637367: 375185	BCC102-637367: 319399	no check	na
Om3	break	lumpy	BCC102-873724: 329213	BCC102-768578: 224085	no check	na
Om13	break	lumpy	BCC102-873724: 173257	BCC102-546026: 350305	no check	na
Om28	break	lumpy	BCC102-867403: 378390	BCC102-496340: 275198	no check	na
Mp27 - 34 - 35	break	lumpy	chrom_01: 564769	chrom_01: 1360407	no check	na
Om9 - 29d	deletion	lumpy	BCC102-867403: 33955	BCC102-867403: 34170	false	false
Om31	deletion	lumpy	BCC102-466838: 274207	BCC102-466838: 276502	false	false
Om21b	deletion	lumpy	BCC102_ML_SOC: 571078	BCC102_ML_SOC: 572460	true	true
Om17	duplication	lumpy	BCC102-806365: 187888	BCC102-806365: 188243	no check	false
Om32	duplication	lumpy	BCC102_ML_SOC: 243207	BCC102_ML_SOC: 251990	no check	false

Figure 1. A. Spontaneous mutation rate per nucleotide per haploid genome per generation. Light blue: μ_{st}, spontaneous structural mutation rate. Blue: μ_{id}, spontaneous nucleotide mutation rate. Dark blue: μ_{bs}, spontaneous short insertion-deletion mutation rate. Horizontal bars are confidence interval following a Poisson distribution. μ_{id} and μ_{bs} come from previous studies (Krasovec *et al.*, 2017, 2018). μ_{st} for *O. tauri*, *M. pusilla*, *B. prasinos* et *P. costavermella* are calculated assuming one hypothetical mutation (non has been validated). **B.** PCR product migration of one large deletion (1,382 bp in intergenic region) identified in *O. mediterraneus*. The ladder is the 1 kb DNA ladder from Promega. No amplification is observed in the mutation accumulation line Om 21b because primers were designed insight the deleted sequence. OmML is the T0 genome of the mutation accumulation.

DATA AVAILABILITY STATEMENT

All genomic raw reads used in this study are available on the SRA database in the NCBI under the bioprojects PRJNA531882 (*Ostreoccocus tauri*, *O. mediterraneus*, *Micromonas pusilla*, *Bathycoccus prasinos*), PRJNA453760 and PRJNA389600 (*Picochlorum costavermella*).

ACKNOWLEDGEMENTS

We are grateful to all GENOPHY team of the Banyuls sur mer lab for discussions and advices.

AUTHOR CONTRIBUTIONS

MK designed and managed the study, MLP did bioinformatics analysis and wet lab works, MK wrote the draft of the manuscript, all authors contributed to editing the last manuscript version.

FUNDING

This work was funded by ANRJCJC-SVSE6-2013-0005.

COMPETING INTERESTS

The authors declare no conflicts of interest.

BIBLIOGRAPHY

Blanc-Mathieu, R., Krasovec, M., Hebrard, M., Yau, S., Desgranges, E., Martin, J., *et al.* 2017. Population genomics of picophytoplankton unveils novel chromosome hypervariability. *Sci. Adv.* **3**: e1700239.

Halligan, D.L. & Keightley, P.D. 2009. Spontaneous Mutation Accumulation Studies in Evolutionary Genetics. *Annu. Rev. Ecol. Evol. Syst.* **40**: 151–172.

Ho, E.K.H. & Schaack, S. 2021. Intraspecific variation in the rates of mutations causing structural variation in Daphnia magna. *Genome Biol. Evol.* evab241.

Katju, V. & Bergthorsson, U. 2019. Old Trade, New Tricks: Insights into the Spontaneous Mutation Process from the Partnering of Classical Mutation Accumulation Experiments with High-Throughput Genomic Approaches. *Genome Biol. Evol.* **11**: 136–165.

Krasovec, M., Eyre-Walker, A., Grimsley, N., Salmeron, C., Pecqueur, D., Piganeau, G., *et al.* 2016. Fitness Effects of Spontaneous Mutations in Picoeukaryotic Marine Green Algae. *G3 GenesGenomesGenetics* **6**: 2063–2071.

Krasovec, M., Eyre-Walker, A., Sanchez-Ferandin, S. & Piganeau, G. 2017. Spontaneous Mutation Rate in the Smallest Photosynthetic Eukaryotes. *Mol. Biol. Evol.* **34**: 1770–1779.

Krasovec, M., Rickaby, R.E.M. & Filatov, D.A. 2020. Evolution of Mutation Rate in

Astronomically Large Phytoplankton Populations. *Genome Biol. Evol.* **12**: 1051–1059.

Krasovec, M., Sanchez-Brosseau, S., Grimsley, N. & Piganeau, G. 2018. Spontaneous
mutation rate as a source of diversity for improving desirable traits in cultured microalgae.
Algal Res. **35**: 85–90.

Krasovec, M., Sanchez-Brosseau, S. & Piganeau, G. 2019. First Estimation of the
Spontaneous Mutation Rate in Diatoms. *Genome Biol. Evol.* **11**: 1829–1837.

Kucukyildirim, S., Behringer, M., Sung, W., Brock, D.A., Doak, T.G., Mergen, H., *et al.* 2020.
Low Base-Substitution Mutation Rate but High Rate of Slippage Mutations in the Sequence
Repeat-Rich Genome of Dictyostelium discoideum. *G3 Bethesda Md* **10**: 3445–3452.

Küpper, C., Stocks, M., Risse, J.E., dos Remedios, N., Farrell, L.L., McRae, S.B., *et al.* 2016.
A supergene determines highly divergent male reproductive morphs in the ruff. *Nat. Genet.*
48: 79–83. Nature Publishing Group.

Layer, R.M., Chiang, C., Quinlan, A.R. & Hall, I.M. 2014. LUMPY: a probabilistic framework
for structural variant discovery. *Genome Biol.* **15**: R84.

Li, H. & Durbin, R. 2010. Fast and accurate long-read alignment with Burrows–Wheeler
transform. *Bioinformatics* **26**: 589–595.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., *et al.* 2009. The
Sequence Alignment/Map format and SAMtools. *Bioinforma. Oxf. Engl.* **25**: 2078–2079.

Long, H., Doak, T.G. & Lynch, M. 2018. Limited Mutation-Rate Variation Within the
Paramecium aurelia Species Complex. *G3 GenesGenomesGenetics* **8**: 2523–2526.

Lynch, M., Ackerman, M.S., Gout, J.-F., Long, H., Sung, W., Thomas, W.K., *et al.* 2016.
Genetic drift, selection and the evolution of the mutation rate. *Nat. Rev. Genet.* **17**: 704–714.

Martin, A.P. & Palumbi, S.R. 1993. Body size, metabolic rate, generation time, and the
molecular clock. *Proc. Natl. Acad. Sci. U. S. A.* **90**: 4087–4091.

Mooers, A.O. & Harvey, P.H. 1994. Metabolic rate, generation time, and the rate of molecular
evolution in birds. *Mol. Phylogenet. Evol.* **3**: 344–350.

Mukai, T. 1964. The Genetic Structure of Natural Populations of Drosophila Melanogaster. I.
Spontaneous Mutation Rate of Polygenes Controlling Viability. *Genetics* **50**: 1–19.

Press, M.O., Hall, A.N., Morton, E.A. & Queitsch, C. 2019. Substitutions Are Boring: Some
Arguments about Parallel Mutations and High Mutation Rates. *Trends Genet. TIG* **35**: 253–
264.

Rausch, T., Zichner, T., Schlattl, A., Stütz, A.M., Benes, V. & Korbel, J.O. 2012. DELLY:
structural variant discovery by integrated paired-end and split-read analysis. *Bioinformatics*
28: i333–i339.

Sung, W., Ackerman, M.S., Dillon, M.M., Platt, T.G., Fuqua, C., Cooper, V.S., *et al.* 2016.
Evolution of the Insertion-Deletion Mutation Rate Across the Tree of Life. *G3 Bethesda Md*,
doi: 10.1534/g3.116.030890.

Sung, W., Ackerman, M.S., Miller, S.F., Doak, T.G. & Lynch, M. 2012. Drift-barrier hypothesis
and mutation-rate evolution. *Proc. Natl. Acad. Sci. U. S. A.* **109**: 18488–18492.

Thomas, J.A., Welch, J.J., Lanfear, R. & Bromham, L. 2010. A generation time effect on the
rate of molecular evolution in invertebrates. *Mol. Biol. Evol.* **27**: 1173–1180.

Tuttle, E.M., Bergland, A.O., Korody, M.L., Brewer, M.S., Newhouse, D.J., Minx, P., *et al.*
2016. Divergence and Functional Degradation of a Sex Chromosome-like Supergene. *Curr.*
Biol. CB **26**: 344–350.

Wala, J.A., Bandopadhayay, P., Greenwald, N.F., O'Rourke, R., Sharpe, T., Stewart, C., *et*
al. 2018. SvABA: genome-wide detection of structural variants and indels by local assembly.

Genome Res. **28**: 581–591.

Weller, C. & Wu, M. 2015. A generation-time effect on the rate of molecular evolution in
bacteria. *Evol. Int. J. Org. Evol.* **69**: 643–652.

Yau, S., Krasovec, M., Benites, L.F., Rombauts, S., Groussin, M., Vancaester, E., *et al.*
2020. Virus-host coexistence in phytoplankton through the genomic lens. *Sci. Adv.* **6**:
eaay2587. American Association for the Advancement of Science.

Zhu, Y.O., Siegal, M.L., Hall, D.W. & Petrov, D.A. 2014. Precise estimates of mutation rate
and spectrum in yeast. *Proc. Natl. Acad. Sci. U. S. A.* **111**: E2310-2318.

Table S1. Blastn results of the deleted sequence in *Ostrecoccus mediterraneus*. The blasted

sequence was the sequence amplified by PCR in the MA mother line (404 bp length).

Chromosome	Identity	Length	Start	End	Pvalue
BCC102-546026	98.916	369	358951	359319	0.0
BCC102-546026	93.590	156	356541	356696	3.17e-62
BCC102-546026	89.062	128	468750	468877	5.46e-40
BCC102-466838	91.617	167	435797	435963	9.76e-62
BCC102_ML_SOC	99.010	404	321019	321422	0.0
BCC102_ML_SOC	98.721	391	571052	571442	0.0
BCC102_ML_SOC	90.678	354	225814	226166	2.59e-133
BCC102_ML_SOC	90.571	350	257243	257591	4.34e-131
BCC102_ML_SOC	90.571	350	584735	585083	4.34e-131
BCC102_ML_SOC	90.299	134	208824	208955	2.28e-44