# Identification of Alkaloids and Related Intermediates of *Dendrobium officinale* by Solid-Phase Extraction Coupled with High-Performance Liquid Chromatography Tandem Mass Spectrometry

Cheng Song[1*], Yunpeng Zhang[2], Muhammad Aamir Manzoor[3], Guohui Li[1*]

[1] College of Biological and Pharmaceutical Engineering, West Anhui University, Lu'an 237012, China

[2] Shanghai Key Laboratory of regulatory biology, School of Life Science, East China Normal University, Shanghai 200241, China

[3] College of Life Science, Anhui Agricultural University, Hefei 230036, China

*Correspondence*:

Cheng Song (lanniao812329218@163.com)

Guohui Li (2352871552@qq.com)

**Abstract**：JA signaling plays a pivotal role in plant stress responses and secondary metabolism. Many studies have demonstrated that jasmonates effectively induce the expressions of alkaloid biosynthesis genes in various plants, which rendered to the accumulation of alkaloid to counteract stresses. Despite the multiple roles of jasmonate (JA) in the regulation of plant growth and different stresses, less studied involved in the regulatory role of JA in *D. officinale* alkaloids. A strategy for the rapid identification of alkaloid and the intermediates of *D. officinale* was established based on a solid-phase extraction coupled with high-performance liquid chromatography tandem mass spectrometry method. By using SPE-HPLC-LTQ-Orbitrap method, the potential compounds were tentatively identified by aligning the accurate molecular weight with the METLIN and Dictionary of Natural Products databases. The chemical structures and main characteristic fragments of the potential compounds were further confirmed by retrieving the multistage mass spectra from the MassBank and METLIN databases.The Mass Frontier software was used to speculate the fragmentation pathway of the identified compounds. Seven alkaloids were separated and identified

from *D. officinale*, which were mainly classified into five types (tropane alkaloids,

tetrahydroisoquinoline alkaloids, quinolizidine alkaloids, piperidine alkaloids and

spermidine alkaloids). Besides the alkaloids, forty-nine chemical substances,

including guanidines, nucleotides, dipeptides, sphingolipids and nitrogen-containing

glucosides, were concurrently identified. These findings gives the composition of

chemicals currently found in *D. officinale*, which could provide the scientific method

for the identification of alkaloids in other *Dendrobium* plants.

**Keywords**: Hormone elicitor, Liquid chromatography-mass spectrometry, Alkaloid,

*Dendrobium*

## Introduction

Traditional Chinese medicine (TCM) involves complex chemical constituents. An

effective determination of one group of metabolites from a certain plant entails a

comprehensive methodology for their purification and identification. Conventional

detection methods such as thin-layer chromatography (TLC) and high-performance

liquid chromatography (HPLC) generally provide such low accuracy and poor

sensitivity that leading to incapable of detecting some trace substances (Pan et al.,

2015). A separation and identification method with strong specificity, high efficiency,

and few interference is urgently needed. Solid-phase extraction (SPE) was usually

used for the complicated specimens by reducing matrix interference and raising

sensitivity (Bucar et al., 2013). Orbitrap mass spectrometry is one kind of high-

resolution mass analyzer containing high-resolution multistage mass fragment

monitoring, a selective fragmentation mode, and a personalized data acquisition

method (Qiu et al., 2015). Orbitrap has been widely applied in pharmacokinetic

analysis, food industry, environmental monitoring and proteomics (Quifer-Rada et al.,

2015; Li et al., 2013). Therefore, the SPE-HPLC-LTQ-Orbitrap technique is proved to

be a powerful means to elucidate the complicacy in TCM. *D. officinale*, as a type of

perennial herb, is attached to the Dendrobium genus of Orchidaceae. As a valuable

TCM, *D. officinale* contains many different types of active ingredients, including

polysaccharides, alkaloids, bibenzyls, phenanthrenes, sesquiterpenoids, fluorenones,

flavonoids, steroids, *etc.* (Xu et al., 2013). Some of these compounds had exhibited

versatile pharmacological and pharmacodynamic activity (Ng et al., 2012). The

earliest study on the alkaloids of the Dendrobium genus started from dendrobine of *D.*

*nobile*. For the decades, dozens of alkaloids have been subsequently identified from *D.*

*anosmum*, *D. chrysanthum*, *D. crepidatum*, *D. findlayanum*, *D. friedricksianum*, *D.*

*hilderbrandii*, *D. loddigessi*, *D. lohohense*, *D. moniliforme*, *D. parishii*, *D. pierardii*,

*D. primulimun*, *D. snowflake* and *D. wardianum* (Xu et al., 2013). At present, the

alkaloids identified from Dendrobiun plants had been divided into five categories

based on their chemical structures, including sesquiterpenoid alkaloids, indolizidine

alkaloids, pyrrolidine alkaloids, phthalide alkaloids, and imidazole alkaloids (Song et

al., 2022).

69           Thus far, the composition and biosynthetic pathway of the alkaloids in *D.*

*officinale* has not been entirely understood. The determination of the alkaloids

between *D. officinale* and *D. nobile* using an LC-MS method showed that 25 identical

compositions in the two Dendrobium plants were tentatively identified; however,

dendrobine was only detected in *D. nobile* (Song et al., 2020). Using a UPLC method,

a few coumarins, alkaloids and bibenzyl components from fifteen Dendrobium

species were quantitatively determined. Dendrobine was only detected in *D. nobile*

originated from the Guizhou and Yunnan Provinces (Xu et al., 2010). The genome and

transcriptome analyses of *D. officinale* implied that a biosynthetic pathway oriented to

the terpenoid indole alkaloids likely exists (Guo et al., 2013; Yan et al., 2015).

Recently, high-throughput sequencing revealed the identification of a total of 56 genes,

including 25 key enzymes potentially involved in the synthesis of terpenoid indole

alkaloids, tropine alkaloids and isoquinolines alkaloids in *D. officinale* (Niu et al.,

2021). Many studies have shown that *CrMYC2* and *ORCAs* transcription factors play

a positive role in elevating the TIAs content in *Catharanthus roseus* (Memelink et al.,

2007). With the help of transcriptome sequencing, 75 bHLH family genes and 66

AP2/ERF genes were identified in the DEGs of *D. officinale,* suggesting that those

transcription factors could be involved in the regulation of secondary metabolism,

especially in TIAs biosynthesis (Zhang et al., 2016; Song et al., 2022). Mevalonate

(MVA) and methylerythritol 4-phosphate (MEP) pathway are both core pathways

related to MeJA-induced accumulation of TIAs (Song et al., 2021). In *D. officinale*,

the key genes in MVA and MEP pathway and some post-modified enzymes were

significantly up-regulated by MeJA treatment (Chen et al., 2019). In the previous

study, high-yielding production of alkaloids from *D. officinale* were obtained with the

combination of tryptophan, secologanin and MeJA treatment (Jiao et al., 2018; Song

et al., 2022). Here, a technical scheme for the separation and determination of

alkaloids and precursors in *D. officinale* was established. The accurate mass weights

of the components separated by HPLC were determined using an Orbitrap mass

analyzer. The multistage fragmentation mass spectra of alkaloids were monitored to

speculate the potential substances based on a database-dependent retrieval method.

Alkaloid and their precursors related to alkaloid-synthetic pathway were associated to

construct the metabolic network between upstream and downstream. The results

would provide a reliable scientific reference and technical guidance for the

development and quality evaluation of *D. officinale*.

**Material and Methods**

Instruments and chemicals

The LTQ-Orbitrap XL mass spectrometer were manufactured by Thermo Fisher

Scientific (Bremen, Germany). The Agilent 1260 instrument used in the experiments

was manufactured by Agilent Technologies (Waldbronn, Germany). The Xcalibur (*v.*

2.1) and Mass Frontier (*v.* 6.0) were used for the analysis of mass spectrometry and

speculation of the fragmentation pathways . Acetonitrile and formic acid were of

HPLC grade. Other chemicals were domestically made analytical reagents. The

standards of reserpine and sarpagine was obtained from Aladdin (Shanghai, China)

and BioBioPha (Kunming, China), respectively.

Plant samples preparation

The *D. officinale* samples used in the experiment were obtained from the tissue

culture room of the Plant Cell Engineering Center in West Anhui University. The

tissue-cultural seedlings cultivated in MS medium were collected after 100 μM MeJA

treatment. The seedlings were firstly dried at 110 °C for 15 min and kept in a constant

weight at 60 °C. The dried materials were ground into powders by a universal grinder.

Using the ultrasonic extraction method, two gram of powder was placed in a

round-bottom flask with 100 mL of methanol and extracted three times at room

temperature. The bromocresol green colorimetric method was used for the

determination of total alkaloids. The extraction solution was centrifuged for 10 min at

6000 x $g$ to get the supernatant, then evaporated to dryness in a rotary evaporator. The

extraction was dissolved with 5 mL of dilute sulfuric acid, filtered through a 0.22 μm

membrane, and pretreated using MCX cartridges (60 mg, 3 mL) for purification. The

protocol was successively conducted by our previous method (Song et al., 2020).

LC-MS conditions

The Waters T3 column (150 mm × 4.6 mm, 3 μm) was used for the

chromatographic separation. The mobile phases acetonitrile (A) with 0.1% formic

acid and aqueous phase (B) with 0.1% formic acid were used for the experiment.

Gradient elution process was as follows: 0-2 min (0% A), 2-12 min (0 to 15% A),

12-22 min (15 to 35% A), 22-32 min (35 to 80% A), and 32-37 min (80 to 0% A). The

UV wavelength was set at 280 nm, and the column temperature was set at 25 °C. A

three-way splitter was connected with the mass analyzer at a 0.3 mL min$^{-1}$ flow rate.

The ion-source parameters were refereed to Kumar's method and slightly modified

(Kumar et al., 2016). The detailed parameters were as follows: sheath gas at 25 arb,

auxiliary gas at 3 arb, spray voltage at 4 kV, the capillary temperature at 320 °C, tube

lens at 120 V, and capillary voltage at 30 V. The MS data were collected at $100 \leq m/z$

$\leq 1000$ in positive-ion mode. Two data acquisition methods were used in the

experiments. In one method, a high-resolution scan was conducted using the Orbitrap

mass analyzer to acquire the MS data at a resolution at 30,000 FWHM and the MS$^2$

data at 15,000 FWHM. A data-dependent MS$^n$ scan was used for the analysis of the

MS$^2$ spectra generated from the most abundant ions of the MS spectra. In the other

method, a high-resolution scan was conducted first to acquire the MS data at a

resolution at 30,000 FWHM, then the LTQ dynode was used to scan the MS$^2$ and MS$^3$

spectra. The dynamic exclusion function was used to reduce repeat scans, and the

repeat count was 2. The exclusion duration was 20 s, and the exclusion mass width

was 3 $m/z$. The suitable collision-induced dissociation (CID) energy was set at 35%.

The minimum signal threshold was 500, and the isolation width of precursor ions was

2 $m/z$. Sarpagine and reserpine were respectively injected into the mass analyzer via a

syringe pump at a flow rate of 6 $\mu$L min$^{-1}$ to complete the direct infusion analysis. The

putative molecular formulae were calculated and obtained according to the accurate

molecular weight using Xcalibur (*v.* 2.1). Compounds were tentatively identified by

matching METLIN, Mass Bank, Dictionary of Natural Products and Chemspider

databases. For the compounds without multiple mass spectra, Mass Frontier (*v.* 6.0)

was helped to speculate and deduce the fragmentation pathway.

## Results and discussion

Biomass and alkaloid contents under the MeJA treatment

The effects of methyl jasmonate on the growth and alkaloid accumulation of *D.*

*officinale* was studied. The results showed that MeJA delayed the growth rate of *D.*

*officinale* and initiated the synthesis of a large number of alkaloids in logarithmic

growth phase (**Figure 1**). The highest content of alkaloids climaxed to 343 $\mu$g/g when

cultured on the 32 days. After that, the fresh weight tended to be stable, and the

alkaloid content began to fall gradually. The results suggested that jasmonates could

continuously promote the synthesis of alkaloids during the growth of *D. officinale*.

The growth trend and accumulating pattern were consistent with our previous study

(Jiao et al., 2018; Wang et al., 2021).

Mass spectra interpretation

Firstly, *D. officinale* samples were extracted by methanol and dissolved with

diluted sulfuric acid. The MCX cartridges were used for the purification of alkaloids

from the acidic aqueous extraction. To optimize the condition of mass spectrometry,

the suitable CID energy was achieved by the direct infusion experiment. Generally,

the strongest fragment ions and the smallest number of fragments were observed at

30-35% collision energy, whereas the abundances of the fragments decreased

substantially at 45% or greater collision energy. In this experiment, sarpagine and

reserpine achieved sufficient fragments both in quantity and intensity at 40% collision

energy. High-resolution mass analysis was conducted by a data-dependent $MS^n$ scan

from the workstation to produce abundant fragmentation spectra of the product ions.

Due to the high resolution, the scan rate of a single spectrum was reduced, and the

analytical analysis time of the entire sample was extended. An appropriate exclusion

duration time was required for the dynamic exclusion function to avoid repeat scans

of target ions and thereby shorten the analysis time of the samples. Another important

function of dynamic exclusion was to obtain the fragment information of the

co-eluting small peaks, which allowed compounds with low ionized potentials to be

detected (Pan et al., 2015). The total ion current chromatogram of the methanol

extract spiked with standards from *D. officinale* is presented, as obtained in the

positive-ion mode under the optimized mass parameters and analytical conditions

(**Figure S1a**). Generally, $[M+H]^+$ or $[M]^+$ was achieved through the soft ionization in

the positive mode. The molecular-ion peaks of sarpagine and reserpine were

separately extracted at *m/z* 311.1738 and 609.2772. The retention time of sarpagine

and reserpine were at 12.95 min and 25.41 min, respectively (**Figure S1b** and **S1c**).

The results indicated that the established method for the mass spectrometry was

suitable for the separation and characterization of alkaloids.

Using a data-dependent $MS^n$ scan method, the MS spectra of two standards were

obtained from the Orbitrap analyzer, and the $MS^n$ spectra were obtained from the LTQ.

It was indicated that the $\pi$ bonds on the indole groups of sarpagine and reserpine were

easily dissociated to produce $[M+H]^+$ peaks and produced massive hydrogen

rearrangement. The ionization and electron transfer were observed on the position of

the group cleavage because of the induced dissociation effect of the high-energy

collisions. The putative fragmentation modes of sarpagine and reserpine are presented

in Figure S2 and S3. The characteristic fragments with *m/z* 146 and 174 from the

indole group were produced by the cleavage of sarpagine and reserpine, respectively.

The characteristic ion *m/z* 195 cleaved on the ester bond was obtained to produce the

3,4,5-trimethoxybenzaldehyde group from the aglycone. Finally, forty-nine

nitrogen-containing compounds, including seven alkaloids, were separated and

tentatively identified from *D. officinale* (**Table S1**).

Identification of the alkaloids

The small molecular compounds previously identified from *D. officinale* mainly

include stilbenoids (bibenzyl and phenanthrene), phenols, lignans, nucleotides,

lactones, flavonoids, steroids, and fatty acids. Based on the high resolution and

sensitivity of LTQ-Orbitrap for the trace substances, coupled with the high selectivity

of MCX for the medium-strength alkaline compounds, seven alkaloids identified from

the methanol extract of *D. officinale* were divided into five classes based on the

chemical structure, including tropane alkaloids, tetrahydroisoquinoline alkaloids,

quinolizidine alkaloids, piperidine alkaloids, and spermidine alkaloids. The special

fragmentation modes of different alkaloids in the ESI source were discussed below.

The tropane alkaloids consist of pyridine conjugated with piperidine. The

experiment involved in DART-MS in-source CID at 90 V indicated that the

characteristic ions of tropine were *m/z* 158, 142, 124, 93, and 67 (Lesiak et al., 2015).

2,3-Dihydroxynortropane was a kind of nortropane-type alkaloid that lacks an

N-methyl group (Asano et al., 2001). Notably, the hydroxyl groups were present on its

C-2 and C-3 in both the α- and β-configurations, as confirmed by a benzoate chirality

method (Kusano et al., 2002). In this study, a strong $[M+H-\cdot OH]^{+}$ ion peak was

observed under the set CID conditions. In addition to the characteristic ions *m/z* 93

and 67,and the stronger fragments *m/z* 113, 98 and 82 were observed (**Figure 2a**). The

$MS^{2}$ and $MS^{3}$ spectra of 2,3-dihydroxynortropane were displayed, and $[M+H-\cdot OH]^{+}$

ion also appeared (**Figure 2b** and **2c**). However, the base peak of 2,3-dihydroxy-

nortropane was the 2-methylpyrrole ion, which was further fragmented to get *m/z* 56.

Finally, the compound was identified as 2,3-dihydroxynortropane or its isomers.Two

tetrahydroisoquinoline alkaloids were identified as alamaridine and

1,2-dihydro-*O*-methyltazettine. The characteristic ions of tetrahydroisoquinoline

group in alamaridine were *m/z* 177, 176, and 162 in the EI source, and the methyl

group on C-5 was easier to lost compared to that on C-10. The $MS^{2}$ and $MS^{3}$ spectra

of alamaridine are displayed respectively (**Figure 3a** and **3b**). These spectra indicated

that the methyl groups on C-5 and C-10 were lost to produce the base peaks

$[M+H-2CH_3-3H]^+$, which further lacked a hydroxyl group to generate $m/z$ 248 in the

$MS^3$ spectrum. The $C_2H_3$ and ·OH groups were successively cleaved to generate $m/z$

221 and 204 on ring B and $m/z$ 176 and 161 on ring C. 1,2-Dihydro-$O$-methyltazettine,

also known as ungvedine, is a tazettine-type alkaloid (Kadyrov et al., 1979). The main

fragments $m/z$ 207 and 142 were generated by 1,2-dihydro-$O$- methyltazettine with

equivalent intensity (**Figure 3c**). The $m/z$ 142 further lacked the methyl group and

simultaneously formed two double bonds to generate the base peak $m/z$ 124 (**Figure**

**3d**). Previous studies have shown that the fragmentation pattern of tazettine-type

alkaloid and its $C_3H_7N$ group was easily lost to produce the characteristic ion

$[M+H-57]^+$ (Zhang et al., 2009). In Fig. 3c, 1,2-dihydro-$O$- methyltazettine was

fragmented by RDA cleavage at both C-6, N-5 and C-4b, C-6a. 1,2-Dihydro-

$O$-methyltazettine was cleaved at C-4b, C-12a but not on C-3, C-4, and thus did not

produce $[M+H-C_3H_7N]^+$. There was a coordination bond between nitrogen and

oxygen presented in the quinolizidine alkaloid methyllagerine $N$-oxide. It was

indicated that this bond easily led to the generation of the fragment $[M-O]^+$ in the EI

source (Lee et al., 2011). The oxygen was easily lost as well to produce the fragment

$m/z$ 437 in the ESI source (**Figure 4a**). The characteristic ions $m/z$ 308 and 292 were

generated by the cleavage of C-11 and C-22. According to the different cleavage

positions on ring B, the characteristic ions $m/z$ 220 and 163 were also generated.

Meefarnine B was a spermidine alkaloid containing a polyamine heterocyclic ring.

These alkaloids are generally considered to be synthesized by putrescine, which is

produced by the degradation of L-arginine or L-ornithine (da Silva et al., 2015). The

$MS^2$ and $MS^3$ spectra of meefarnine B indicate that the $NH_3$ group on N-5 was

cleaved to produce the fragment $m/z$ 421 under the collision dissociation effect

(**Figure 4b** and **4c**). The fragment $m/z$ 292 was generated by the cleavage of the side

chain on N-9 and was further cleaved to produce $m/z$ 218. Besides, the fragment $m/z$

275 was generated by the cleavage of the $m/z$ 421 on N-1 and C-2. The fragment $m/z$

204 was generated by the cleavage of the $m/z$ 275 on N-9 and C-10, which further

produced the characteristic ion $m/z$ 147. In this study, a piperidine alkaloid anapheline

was tentatively identified from *D. officinale*. The retention time of anapheline was

27.88 min, and the molecular weight of the $[M+H]^+$ was 225.1945, which implies that

its molecular formula was $C_{13}H_{25}ON_2$ (**Figure S4**). Anapheline easily lost $H_2O$ to

produce the fragment $m/z$ 207 and the characteristic ion piperidine $m/z$ 100 during the

collision-induced dissociation. The $NH_3$ group was lost to produce the $m/z$ 83 in the

$MS^2$ spectrum. Based on the $MS^2$ and $MS^3$ spectra, a presumed fragmentation

pathway of anapheline was proposed (**Figure 5**).

Identification of the guanidines

Three guanidine compounds were separated and determined from the methanol

extract of *D. officinale* for the first time. The $MS^2$ and $MS^3$ spectra of

4-guanidinobutanoic acid were displayed respectively (**Figure 6a** and **6b**). The mass

spectra indicated that 4-guanidinobutanoic acid easily lost $H_2O$ to form the base peak

$m/z$ 128 and produced the characteristic ions $m/z$ 104, 87 and 60. The $m/z$ 128 was

further cleaved to produce the $m/z$ 111 and 86. These results were consistent with

those reported by the previous study (Nikolić et al., 2012). Cyclocimipronidine is a

cyclic guanidine compound separated from *Cimicifuga racemosa*. Some studies

showed that the characteristic ions of cyclocimipronidine, a guanidine alkaloid in

black cohosh, were the $m/z$ 112, 95, 94, 70 and 67 (Gödecke et al., 2009) . In addition

to the aforementioned characteristic ions, cyclocimipronidine easily lost $H_2O$ to

generate the fragments $m/z$ 136 and 84 (**Figure 6c** and **6d**). The $MS^2$ and $MS^3$ spectra

of *N-tert*-butyloxycarbonyl guanidine were displayed in Fig. 6e and 6f, respectively. It

was indicated that the characteristic ions were the $m/z$ 142, 125, 104 and 87.

Identification of the nucleotides and derivatives

Nucleotides were composed of purine or pyrimidine linked with ribose. adenosine,

uridine and guanosine had been separated from *D. officinale* in a previous study. In

this study, five nucleotide derivatives were tentatively identified. 2-Hydroxyadenine

was an adenine derivative that easily lost an $NH_3$ group to produce the characteristic

ions $m/z$ 135 and 110 (**Figure S5a**). The fragments $m/z$ 107, 92 and 77 were generated

in the MS$^3$ spectrum (**Figure S5b**). Besides 2-hydroxyadenine, another four

nucleotide derivatives were identified in the study. Their MS$^2$ and MS$^3$ spectra

indicated that the neutral loss of *m/z* 136 resulted from the loss of aglycones (**Figure**

**S5 c-j**). Comparison with the multistage mass spectra of the METLIN database

(https://metlin.scripps.edu/) resulted in the identification of adenosine, guanosine,

1-methyladenosine and cytidine.

Identification of the sphingolipids

Sphingolipid is a structurally complex lipid containing a long-chain sphingosine

skeleton, which forms the plasma and vacuole membranes of plant cells. Four

phytosphingosines were separated and identified from *D. officinale*. Because

sphingosine contained some hydroxyl groups, the characteristic ion $[M+H-H_2O]^+$ was

observed during the dissociation (**Figure S6a-h**). By aligning against the Dictionary

of Natural Products Database (http://dnp.chemnetbase.com), four sphingolipids,

including 2-amino-1,3-hexadecanediol, 2-amino-3-tetradecanol, 2-amino-1,3,4-

octadecanetriol and 2-amino-1,3,4-hexadecanetriol, were identified in the experiment.

Identification of the amino acids and their derivatives

Eight amino acids—L-lysine, L-histidine, L-arginine, L-valine, L-isoleucine,

L-tyrosine, L-phenylalanine and L-tryptophan—were identified from *D. officinale*

based on accurate molecular weights and the multistage mass spectra of standards

from the MassBank mass spectrum library (http://www.massbank.jp/QuickSearch.

html) (**Table S1**). 4-Hydroxy-L- threonine and urocanic acid were derivatives of

L-threonine and L-histidine that easily lost $H_2O$ to produce the base peaks *m/z* 118 and

121. The putative fragment ions generated during their cleavage were displayed in

**Figure S7a** and **S7b**, respectively. In addition to the derivatives of

L-threonine and L-histidine, four additional arginine derivatives were identified in this

study. $N^2$-Fructopyranosylarginine was an arginine monoglycoside and the byproduct

generated by a Maillard reaction between reducing sugars (fructose or glucose) and

arginine upon heating (Suzuki et al., 2004). Because fructose contained several

hydroxyl groups, $H_2O$ was easily lost to produce the characteristic ions *m/z* 319, 301

and 283. The fragments $m/z$ 257 and 239 were produced by the cleavage of the

guanidino group on the glycoside. $N^2$-Fructopyranosylarginine was also cleaved to

generate the characteristic ion $m/z$ 175 on the N-glucosidic band and further produced

the aglycone $m/z$ 158 via the neutral loss of fructose (**Figure S7c-d**). The $NH_3$ group

of $N$-methylarginine was lost to produce the base peak $m/z$ 172, as a result of the

dissociation. According to the different cleavages on the R group, the characteristic

ions $m/z$ 158, 133, 116 and 74 were generated (**Figure S7e**). The characteristic ions

$m/z$ 126, 115, 97 and 70 were also generated in the $MS^3$ spectrum by the cleavage of

the $m/z$ 172 fragment (**Figure S7f**). The ·OH group of $N^{\alpha}$-acetylarginine was lost to

produce the characteristic ion $m/z$ 200. The fragments $m/z$ 175 and 157 were

generated by the different cleavages on the R group. The characteristic ions $m/z$ 139,

115 and 70 were produced by the cleavage of amino and carboxyl groups linked with

the chiral carbons. The ·OH group of the $m/z$ 200 was lost to produce the $m/z$ 183 in

the $MS^3$ spectrum, and the other fragments were similar to those in the $MS^2$ spectrum

(**Figure S7g-h**). $N^2$-(3-Hydroxysuccinoyl)arginine was formed by the substitution of

the 3-hydroxysuccinoyl group on the N-terminal of arginine. The typical feature of its

cleavage was the loss of 3-hydroxysuccinoyl group to produce the characteristic ion

$m/z$ 175 (**Figure S7i**), which was further cleaved to generate the fragments $m/z$ 158,

139 and 60 (**Figure S7j**).

Identification of some isomers

The accurate molecular weights of the compounds were provided by

high-resolution mass spectrometry, which allowed us to deduce the speculative

molecular formulas. However, the identification of isomers was less effective. Two

pairs of isomers were identified in this study based on their different retention times

and distinct multistage fragments. The formula of both L-homophenylalanine and

β-phenyl-γ-aminobutyric acid was $C_{10}H_{13}O_2N$, but their retention times were at 6.79

and 12.47 min, respectively (**Table S1**). The $MS^2$ spectra of L-homophenylalanine and

β-phenyl-γ-aminobutyric acid are displayed in **Figure S8a** and **S8c**, respectively.

These mass spectra indicated that the $NH_3$ groups were easily lost to form the

characteristic ion $m/z$ 163. However, the fragment $m/z$ 120 produced in the spectrum

of β-phenyl-γ-aminobutyric acid was not observed in that of L-homophenylalanine.

The MS$^3$ spectra presented different fragments, which indicated two different

compounds (**Figure S8b** and **S8d**). The formula of both carbetamide and

$N^{\delta}$-benzoylornithine were $C_{12}H_{16}O_3N_2$, but their retention times were at 9.77 and

14.93 min, respectively (**Table S1**). The MS$^2$ and MS$^3$ spectra of carbetamide were

displayed in **Figure S8e** and **S8f**, respectively. The MS$^2$ and MS$^3$ spectra of

$N^{\delta}$-benzoylornithine were displayed in Fig. S8g and S8h, respectively. These two

isomers exhibited different fragmentation patterns. The characteristic ions of

carbetamide were the $m/z$ 120, 103 and 93 ions, whereas the characteristic ions of

$N^{\delta}$-benzoylornithine were the $m/z$ 219, 205, 175, 158 and 148 ions. The isomerism of

the dipeptides resulted from the chirality of the α-carbon and the different positions of

peptide bonds between two peptides. In the study, eight dipeptides were tentatively

identified. Most of these dipeptides had more than one retention time, suggesting that

the isomers existed.

Identification of the nitrogen-containing glycosides

By comparing our spectra with the Metlin and Dictionary of Natural Products

databases, we tentatively identified four nitrogen-containing glycosides from *D.*

*officinale*. The neutral loss of $H_2O$ in 5'-*O*-β-D-glucosylpyridoxine was easy to

produce the characteristic ion $m/z$ 314. The fragment $m/z$ 171 was generated by the

cleavage on the O-glucosidic bond and further produced the aglycone $m/z$ 152 by the

neutral loss of glucose (**Figure S9a**). The $H_2O$ on the aglycone of

4-*O*-β-D-glucopyranoside-2,4-benzoxazolediol was easily lost to produce the base

peak $m/z$ 296. The benzoxazole ion $m/z$ 136 was generated by the cleavage on the

O-glucosidic bond (**Figure S9b**). The $m/z$ 296 was cleaved to generate the

characteristic ions $m/z$ 119 and 94 (**Figure S9c**). The MS$^2$ and MS$^3$ spectra of

*O*-(tri-*O*-acetyl-α-L- rhamnopyranoside)- (4-hydroxybenzyl) methylcarbamic acid

were displayed in **Figure S9d** and **S9e**, respectively. These mass spectra indicated that

the characteristic ion $m/z$ 322 was generated by the cleavage of the ester bond and the

side chain. The *m/z* 322 was cleaved to produce the fragments *m/z* 304 and 160 in the MS$^3$ spectrum. Passicapsin was composed of one glucose and a nitrogen-containing aglycone that easily produced the aglycone *m/z* 238 via the neutral loss of glucose. The characteristic ions *m/z* 286 and 148 were cleaved at the O-glucosidic bond between the side chain and the dideoxy glucoside (**Figure S9f**). The fragments *m/z* 148 and 106 in the MS$^3$ spectrum were produced by the cleavage of the aglycone *m/z* 238 on the O-glucosidic bond. The H$_2$O of the aglycone *m/z* 238 was lost to produce the fragments *m/z* 220 and 212, and the dideoxy glucoside group was further cleaved to generate the *m/z* 190 and 174 (**Figure S9g**).

## Conclusions

The alkaloids and precursors of *D. officinale* with MeJA treatment were primarily studied by the SPE-HPLC–LTQ-Orbitrap technique. The extraction and purification of the target compounds was optimized, and the suitable conditions of ESI-CID-MS$^n$ for the mass spectrometric detection were established. Based on the accurate mass weight with MS$^n$ fragmentation information provided by the LTQ-Orbitrap mass spectrometer, forty-nine compounds, including amino acids derivatives, dipeptides, nucleotides, guanidines, alkaloids, sphingolipids and several nitrogen-containing glycosides, were tentatively identified from *D. officinale*. Seven alkaloids were identified from *D. officinale*. The combination of alkaloids and their precursors were associated to construct a metabolic network for the interpretation of biosynthetic pathways. These findings will serve as a technical guidance for the analyses and drug discoveries of *Dendrobium* alkaloids.

## Acknowledgement

We are grateful to Mrs. Airong Feng of University of Science and Technology of China for her support of the work on mass spectra interpretation.

## Funding

This work was supported by High-level Talents Research Initiation Fund of West Anhui University (WGKQ2022025, WGKQ2021079), and Quality Engineering Project of Anhui Provincial Department of Education (2020jyxm2137).

## Author contributions

CS and GHL discussed the writing plan. CS and YPZ drafted the manuscript. CS and MAM edited the manuscript. GHL and CS acquired the funding. All authors have read, reviewed, and approved the submitted version.

## Conflicts of interest

No interest to declare.

## References

Asano, N., Yokoyama, K., Sakurai, M., Ikeda, K., Kizu, H., Kato, A., et al. (2001). Dihydroxynortropane alkaloids from calystegine-producing plants. *Phytochemistry* 57, 721–726. doi:https://doi.org/10.1016/S0031-9422(01)00131-5.

Bhattacharjya, A., Mukhopadhyay, R., Pakrashi, S. C. (1986). Structure and synthesis of alamaridine, a novel benzopyridoquinolizine alkaloid from alangium lamarck II. *Tetrahedron Lett.* 27, 1215–1216. doi:10.1016/S0040-4039(00)84220-3.

Bucar, F., Wube, A., and Schmid, M. (2013). Natural product isolation--how to get from biological material to pure compounds. *Nat. Prod. Rep.* 30, 525–545. doi:10.1039/c3np20106f.

Chen, Y., Wang, Y., Lyu, P., Chen, L., Shen, C., and Sun, C. (2019). Comparative transcriptomic analysis reveal the regulation mechanism underlying MeJA-induced accumulation of alkaloids in Dendrobium officinale. *J. Plant Res.* 132, 419–429. doi:10.1007/s10265-019-01099-6.

da Silva, G., Martinho, A., Soengas, R. G., Duarte, A. P., Serrano, R., Gomes, E. T., et al. (2015). A new spermidine macrocyclic alkaloid isolated from Gymnosporia arenicola leaf. *Fitoterapia* 106, 7–11. doi:10.1016/j.fitote.2015.07.020.

Gödecke, T., Lankin, D. C., Nikolic, D., Chen, S.-N., van Breemen, R. B., Farnsworth, N. R., et al. (2009). Guanidine alkaloids and Pictet-Spengler adducts from black cohosh (Cimicifuga racemosa). *J. Nat. Prod.* 72, 433–437. doi:10.1021/np8006952.

Guo, X., Li, Y., Li, C., Luo, H., Wang, L., Qian, J., et al. (2013). Analysis of the

Dendrobium officinale transcriptome reveals putative alkaloid   biosynthetic

genes and genetic markers. *Gene* 527, 131–138. doi:10.1016/j.gene.2013.05.073.

Jiao, C., Song, C., Zheng, S., Zhu, Y., Jin, Q., Cai, Y., et al. (2018). Metabolic

profiling of Dendrobium officinale in response to precursors and methyl

jasmonate. *Int. J. Mol. Sci.* 19. doi:10.3390/ijms19030728.

Kalli, A., Smith, G. T., Sweredoski, M. J., and Hess, S. (2013). Evaluation and

optimization of mass spectrometric settings during data-dependent acquisition

mode: Focus on LTQ-orbitrap mass analyzers. *J. Proteome Res.* 12, 3071–3086.

doi:10.1021/pr3011588.

Kumar, S., Singh, A., Bajpai, V., and Kumar, B. (2016). Identification,

characterization and distribution of monoterpene indole alkaloids in Rauwolfia

species by Orbitrap Velos Pro mass spectrometer. *J. Pharm. Biomed. Anal.* 118,

183–194. doi:10.1016/j.jpba.2015.10.037.

Kusano, G., Orihara, S., Tsukamoto, D., Shibano, M., Coskun, M., Guvenc, A., et al.

(2002). Five new nortropane alkaloids and six new amino acids from the fruit of

Morus alba Linne growing in Turkey. *Chem. Pharm. Bull.* 50, 185–192.

doi:10.1248/cpb.50.185.

Lee, I., Youn, U., Kim, H., Min, B., Kim, J. S., and Bae, K. (2011). Biphenyl and

biphenyl ether quinolizidine N -oxide alkaloids from lagerstroemia indica L.

*Planta Med.* 77, 2037–2041. doi:10.1055/s-0031-1280064.

Lesiak, A. D., Cody, R. B., Dane, A. J., and Musah, R. A. (2015). Plant Seed Species

Identification from Chemical Fingerprints: A High-Throughput Application of

Direct Analysis in Real Time Mass Spectrometry. *Anal. Chem.* 87, 8748–8757.

doi:10.1021/acs.analchem.5b01611.

Li, X., Shen, B., Jiang, Z., Huang, Y., and Zhuo, X. (2013). Rapid screening of drugs

of abuse in human urine by high-performance liquid chromatography coupled

with high resolution and high mass accuracy hybrid linear ion trap-Orbitrap mass

spectrometry. *J. Chromatogr. A* 1302, 95–104.

doi:10.1016/j.chroma.2013.06.028.

Liu, W., Hua, Y., and Zhan, Z. (2007). Moniline, a New Alkaloid from Dendrobium

moniliforme. *ChemInform* 38, 317–318. doi:10.1002/chin.200748193.

Memelink, J., and Gantet, P. (2007). Transcription factors involved in terpenoid

indole alkaloid biosynthesis in Catharanthus roseus. *Phytochem. Rev.* 6, 353–362.

doi:10.1007/s11101-006-9051-z.

Morita, H., Fujiwara, M., Yoshida, N., and Kobayashi, J. (2000). New

Picrotoxinin-type and Dendrobine-type Sesquiterpenoids from Dendrobium

Snowflake 'Red Star.' *Tetrahedron* 56, 5801–5805.

doi:https://doi.org/10.1016/S0040-4020(00)00530-5.

Ng, T. B., Liu, J., Wong, J. H., Ye, X., Wing Sze, S. C., Tong, Y., et al. (2012).

Review of research on Dendrobium, a prized folk medicine. *Appl. Microbiol.*

*Biotechnol.* 93, 1795–1803. doi:10.1007/s00253-011-3829-7.

Nikolić, D., Gödecke, T., Chen, S. N., White, J., Lankin, D. C., Pauli, G. F., et al.

(2012). Mass spectrometric dereplication of nitrogen-containing constituents of

black cohosh (Cimicifuga racemosa L.). *Fitoterapia* 83, 441–460.

doi:10.1016/j.fitote.2011.12.006.

Niu, Z., Zhu, F., Fan, Y., Li, C., Zhang, B., Zhu, S., et al. (2021). The

chromosome-level reference genome assembly for Dendrobium officinale and its

utility of functional genomics research and molecular breeding study. *Acta*

*Pharm. Sin. B* 11, 2080–2092. doi:10.1016/j.apsb.2021.01.019.

Pan, H., Yang, W., Zhang, Y., Yang, M., Feng, R., Wu, W., et al. (2015). An

integrated strategy for the systematic characterization and discovery of

new  indole alkaloids from Uncaria rhynchophylla by

UHPLC/DAD/LTQ-Orbitrap-MS. *Anal. Bioanal. Chem.* 407, 6057–6070.

doi:10.1007/s00216-015-8777-0.

Qiu, S., Yang, W. Z., Shi, X. J., Yao, C. L., Yang, M., Liu, X., et al. (2015). A green

protocol for efficient discovery of novel natural compounds: Characterization of

new ginsenosides from the stems and leaves of Panax ginseng as a case study.

*Anal. Chim. Acta* 893, 65–76. doi:10.1016/j.aca.2015.08.048.

Quifer-Rada, P., Vallverdú-Queralt, A., Martínez-Huélamo, M., Chiva-Blanch, G.,

Jáuregui, O., Estruch, R., et al. (2015). A comprehensive characterisation of beer

polyphenols by high resolution mass spectrometry (LC-ESI-LTQ-Orbitrap-MS).

*Food Chem.* 169, 336–343. doi:10.1016/j.foodchem.2014.07.154.

Shi Qiu, Wen-zhi Yang, Xiao-jian Shi, Chang-liang Yao, Min Yang, Xuan Liu, Bao-

hong Jiang, Wan-ying Wu, D. G. (2015). A green protocol for efficient discovery

of novel natural compounds: characterization of new ginsenosides from the

stems and leaves of Panax ginseng as a case study. *Anal. Chim. Acta* 893, 65–76.

doi:10.1016/j.aca.2015.08.048.

Song, C., Cao, Y., Dai, J., Li, G., and Manzoor, M. A. (2022). The Multifaceted Roles

of MYC2 in Plants : Toward Transcriptional Reprogramming and Stress

Tolerance by Jasmonate Signaling. *Front. Plant Sci.* 13, 1–14.

doi:10.3389/fpls.2022.868874.

Song, C., Jiao, C., Jin, Q., Chen, C., Cai, Y., and Lin, Y. (2020). Metabolomics

analysis of nitrogen-containing metabolites between two Dendrobium plants.

*Physiol. Mol. Biol. Plants* 26, 1425–1435. doi:10.1007/s12298-020-00822-1.

Song, C., Li, G., Dai, J., and Deng, H. (2021). Genome-Wide Analysis of PEBP

Genes in Dendrobium huoshanense: Unveiling the Antagonistic Functions of

FT/TFL1 in Flowering Time. *Front. Genet.* 12, 1–11.

doi:10.3389/fgene.2021.687689.

Song, C., Ma, J., Li, G., Pan, H., Zhu, Y., Jin, Q., et al. (2022). Natural Composition

and Biosynthetic Pathways of Alkaloids in Medicinal Dendrobium Species.

*Front. Plant Sci.* 13, 1–15. doi:10.3389/fpls.2022.850949.

Song, C., Zhang, Y., Chen, R., Zhu, F., and Wei, P. (2022). Label-Free Quantitative

Proteomics Unravel the Impacts of Salt Stress on Dendrobium huoshanense.

*Front. Plant Sci.* 13, 1–12. doi:10.3389/fpls.2022.874579.

Wang, Y., Dai, J., Chen, R., Song, C., Wei, P., Cai, Y., et al. (2021). miRNA-Based

Drought Regulation in the Important Medicinal Plant Dendrobium huoshanense.

*J. Plant Growth Regul.* 3. doi:10.1007/s00344-021-10366-7.

Xu, J., Han, Q.-B., Li, S.-L., Chen, X.-J., Wang, X.-N., Zhao, Z.-Z., et al. (2013).

Chemistry, bioactivity and quality control of Dendrobium, a commonly used

tonic herb in traditional Chinese medicine. *Phytochem. Rev.* 12, 341–367.

doi:10.1007/s11101-013-9310-8.

Xu, J., Zhao, W., Qian, Z., Guan, J., and Li, S. (2010). Fast determination of five

components of coumarin, alkaloids and bibenzyls in   Dendrobium spp. using

pressurized liquid extraction and ultra-performance liquid chromatography. *J.*

*Sep. Sci.* 33, 1580–1586. doi:10.1002/jssc.201000034.

Yan, L., Wang, X., Liu, H., Tian, Y., Lian, J., Yang, R., et al. (2015). The Genome of

Dendrobium officinale Illuminates the Biology of the Important   Traditional

Chinese Orchid Herb. *Mol. Plant* 8, 922–934. doi:10.1016/j.molp.2014.12.011.

Zhang, J., He, C., Wu, K., Teixeira da Silva, J. A., Zeng, S., Zhang, X., et al. (2016).

Transcriptome analysis of dendrobium officinale and its application to the

identification of genes associated with polysaccharide synthesis. *Front. Plant Sci.*

7. doi:10.3389/fpls.2016.00005.

Zhang, X., Huang, H., Liang, X., Huang, H., Dai, W., Shen, Y., et al. (2009). Analysis

of Amaryllidaceae alkaloids from Crinum by high-performance liquid

chromatography coupled with electrospray ionization tandem mass spectrometry.

*Rapid Commun. Mass Spectrom.* 23, 2903–2916. doi:10.1002/rcm.4205.

**Figure captions**

**Figure. 1**. The fresh weight and alkaloid contents of *D. officinale* treated with 100 μM

MeJA.

**Figure. 2**. The mass spectra of tropine and 2,3-dihydroxynortropane. (a) The MS$^2$

spectrum of tropine, (b) the MS$^2$ spectrum of 2,3-dihydroxynortropane or an isomer,

and (c) the MS$^3$ spectrum of 2,3-dihydroxynortropane or an isomer.

**Figure. 3**. The mass spectra of alamaridine and 1,2-dihydro-*O*-methyltazettine. (a)

The MS$^2$ spectrum of alamaridine, (b) the MS$^3$ spectrum of alamaridine, (c) the MS$^2$

spectrum of 1,2-dihydro-*O*-methyltazettine, and (d) the MS$^3$ spectrum of

1,2-dihydro-*O*-methyltazettine.

**Figure. 4.** The mass spectra of methyllagerine *N*-oxide and meefarnine B. (a) The

$MS^2$ spectrum of methyllagerine *N*-oxide, (b) the $MS^2$ spectrum of meefarnine B, and

(c) the $MS^3$ spectrum of meefarnine B.

**Figure. 5**. The putative fragmentation pathway of anapheline.

**Figure. 6**. The mass spectra of the guanidines. (a) The $MS^2$ spectrum of

4-guanidinobutanoic acid, (b) the $MS^3$ spectrum of 4-guanidinobutanoic acid, (c) the

$MS^2$ spectrum of cyclocimipronidine, (d) the $MS^3$ spectrum of cyclocimipronidine, (e)

the $MS^2$ spectrum of *N-tert*-butyloxycarbonyl guanidine, and (f) the $MS^3$ spectrum of

*N-tert*-butyloxycarbonyl guanidine.

**Figure. S1**. The mass spectrum of *D. officinale* extract with the standards. (a) The

total ion current chromatogram of the methanol extract from *D. officinale* spiked with

standards, (b) the extracted ion current chromatogram of *m/z* 311.1738, and (c) the

extracted ion current chromatogram of *m/z* 609.2772.

**Figure. S2**. The putative fragmentation pathway of sarpagine. π denotes π-bond

dissociation; i denotes inductive cleavage; $rH_R$ denotes charge-remote rearrangement;

$rH_B$ denotes α,β-charge-site rearrangement; $rH_C$ denotes γ-charge-site rearrangement;

and $rH_{1,2}$ denotes radical-site rearrangement.

**Figure. S3**. The putative fragmentation pathway of reserpine. π denotes π-bond

dissociation; i denotes inductive cleavage; $rH_R$ denotes charge-remote rearrangement;

and $rH_C$ denotes γ-charge-site rearrangement.

**Figure. S4**. The chromatogram and mass spectra of anapheline. (a) The extracted ion

chromatogram of anapheline, (b) the MS spectrum of anapheline, (c) the $MS^2$

spectrum of anapheline, and (d) the $MS^3$ spectrum of anapheline.

**Figure. S5**. The mass spectra of nucleotides and derivatives. (a) The $MS^2$ spectrum of

2-hydroxyadenine, (b) the $MS^3$ spectrum of 2-hydroxyadenine, (c) the $MS^2$ spectrum

of adenosine, (d) the $MS^3$ spectrum of adenosine, (e) the $MS^2$ spectrum of guanosine,

(f) the $MS^3$ spectrum of guanosine, (g) the $MS^2$ spectrum of 1-methyladenosine, (h)

the $MS^3$ spectrum of 1-methyladenosine, (i) the $MS^2$ spectrum of cytidine, and (j) the

$MS^3$ spectrum of cytidine.

**Figure. S6**. The mass spectra of sphingolipids. (a) The $MS^2$ spectrum of 2-amino-1,3-

hexadecanediol, (b) the MS$^3$ spectrum of 2-amino-1,3- hexadecanediol, (c) the MS$^2$

spectrum of 2-amino-3-tetradecanol, (d) the MS$^3$ spectrum of 2-amino-3-tetradecanol,

(e) the MS$^2$ spectrum of 2-amino-1,3,4-octadecanetriol, (f) the MS$^3$ spectrum of

2-amino-1,3,4-octadecanetriol, (g) the MS$^2$ spectrum of 2-amino-1,3,4-

hexadecanetriol, and (h) the MS$^3$ spectrum of 2-amino-1,3,4-hexadecanetriol.

**Figure. S7**. The mass spectra of amino acids and their derivatives. (a) The MS$^2$

spectrum of 4-hydroxy-L-threonine, (b) the MS$^2$ spectrum of urocanic acid, (c) the

MS$^2$ spectrum of $N^2$-fructopyranosylarginine, (d) the MS$^3$ spectrum of

$N^2$-fructopyranosylarginine, (e) the MS$^2$ spectrum of $N$-methylarginine, (f) the MS$^3$

spectrum of $N$-methylarginine, (g) the MS$^2$ spectrum of $N^\alpha$-acetylarginine, (h) the

MS$^3$ spectrum of $N^\alpha$-acetylarginine, (i) the MS$^2$ spectrum of

$N^2$-(3-hydroxysuccinoyl)arginine, and (j) the MS$^3$ spectrum of

$N^2$-(3-hydroxysuccinoyl)arginine.

**Figure. S8**. The comparative mass spectrum of some isomers. (a) The MS$^2$ spectrum

of L-homophenylalanine, (b) the MS$^3$ spectrum of L-homophenylalanine, (c) the MS$^2$

spectrum of β-phenyl-γ-aminobutyric acid, (d) the MS$^3$ spectrum of

β-phenyl-γ-aminobutyric acid, (e) the MS$^2$ spectrum of carbetamide, (f) the MS$^3$

spectrum of carbetamide, (g) the MS$^2$ spectrum of $N^\delta$-benzoylornithine, and (h) the

MS$^3$ spectrum of $N^\delta$-benzoylornithine.

**Figure. S9**. The mass spectra of nitrogen-containing glycosides. (a) The MS$^2$

spectrum of 5'-$O$-$\beta$-D-glucosylpyridoxine, (b) the MS$^2$ spectrum of 4-$O$-$\beta$-

D-glucopyranoside-2,4-benzoxazolediol, (c) the MS$^3$ spectrum of

4-$O$-$\beta$-D-glucopyranoside-2,4-benzoxazolediol, (d) the MS$^2$ spectrum of

$O$-(tri-$O$-acetyl-α-L-rhamnopyranoside)-(4-hydroxy benzyl)methylcarbamic acid, (e)

the MS$^3$ spectrum of $O$-(tri-$O$-acetyl-$\alpha$-L-rhamnopyranoside)-(4-hydroxy

benzyl)methylcarbamic acid, (f) the MS$^2$ spectrum of passicapsin, and (g) the MS$^3$

spectrum of passicapsin.

**Table S1.** Identification of the main chemical compounds from *D.officinale*

**a**

**b**

**c**

**a**

**b**

**c**