Characterization of alkaloids and metabolic intermediates from *Dendrobium officinale* induced by methyl jasmonate

Cheng Song[1#], Yunpeng Zhang[2#], Muhammad Aamir Manzoor[3], Qing Jin[3*], Guohui Li[1*]

[1] *College of Biological and Pharmaceutical Engineering, West Anhui University, Luan, China*

[2] *Shanghai Key Laboratory of regulatory biology, School of Life Science, East China Normal University, Shanghai, China*

[3] *College of Life Science, Anhui Agricultural University, Hefei, China*

[#] *Contributed equally to this article*

*Correspondence:

Dr. Guohui Li

Affiliation: West Anhui University

Address: West Moon Island, Yunlu Bridge, Lu'an, Anhui, 237012, China

Email: 2352871552@qq.com

Prof. Qing Jin

Affiliation: Anhui Agricultural University

Address: No. 130 Changjiang West Road, Hefei, Anhui, 230036, China

Email: qingjin@ahau.edu.cn

Abstract

JA signaling plays a pivotal role in plant stress responses and secondary metabolism. Many studies have demonstrated that jasmonates effectively induce the expressions of alkaloid biosynthesis genes in various plants, which rendered to the accumulation of alkaloid to counteract stresses. Despite the multiple roles of JAs in the regulation of plant growth and different stresses, less studied involved in the regulatory role of JA in *Dendrobium officinale*. Here, a strategy for the rapid identification of alkaloid and precursors of *D. officinale* was established based on a solid-phase extraction-high performance liquid chromatography-linear ion trap- Orbitrap tandem mass spectrometry (SPE-HPLC-LTQ-Orbitrap) method. The potential compounds were tentatively identified by aligning the accurate molecular weight with the METLIN and Dictionary of Natural

Products databases. The chemical structures and main characteristic fragments of the potential

compounds were further confirmed by retrieving the multistage mass spectra from the MassBank

and METLIN databases.The Mass Frontier software was used to speculate the fragmentation

pathway of the identified compounds. As a result, seven alkaloids were separated and identified

from *D. officinale,* which were mainly classified into five types (tropane alkaloids,

tetrahydroisoquinoline alkaloids, quinolizidine alkaloids, piperidine alkaloids and spermidine

alkaloids). Besides the alkaloids, forty-nine chemical substances, including guanidines,

nucleotides, dipeptides, sphingolipids and nitrogen-containing glucosides, were concurrently

identified in the experiment. Taken together, this study revealed the composition of chemicals

currently found in *D. officinale*, which could provide the scientific method for the identification of

alkaloids in other Dendrobium plants.

Keywords: JA signaling, Liquid chromatography-mass spectrometry, Alkaloid, Fragmentation,

Dendrobium officinale

Introduction

Traditional Chinese medicine (TCM) involves complex chemical constituents. An effective

determination of one group of metabolites from a certain plant entails a comprehensive

methodology for their purification and identification. Conventional detection methods such as

thin-layer chromatography (TLC) and high-performance liquid chromatography (HPLC) generally

provide such low accuracy and poor sensitivity that leading to incapable of detecting some trace

substances (Pan et al., 2015). A separation and identification method with strong specificity, high

efficiency, and few interference is urgently needed. Solid-phase extraction (SPE) was usually used

for the complicated specimens by reducing matrix interference and raising sensitivity (Bucar et al.

2013). Orbitrap mass spectrometry is one kind of high- resolution mass analyzer containing

high-resolution multistage mass fragment monitoring, a selective fragmentation mode, and a

personalized data acquisition method (Qiu et al. 2015). Orbitrap has been widely applied in

pharmacokinetic analysis, food industry, environmental monitoring and proteomics (Anastasia et

al. 2013; Rada et al. 2015; Haddad and Kümmerer 2014; Li et al. 2013). Therefore, the

SPE-HPLC-LTQ-Orbitrap technique is proved to be a powerful means to elucidate the complicacy

in TCM. *Dendrobium officinale* Kimura et Migo, as a type of perennial herb, is attached to the

Dendrobium genus of Orchidaceae. As a valuable TCM, *D. officinale* contains many different

types of active ingredients, including polysaccharides, alkaloids, bibenzyls, phenanthrenes,

sesquiterpenoids, fluorenones, flavonoids, steroids, etc. (Xu et al. 2013). Some of these

compounds had exhibited versatile pharmacological and pharmacodynamic activity (Ng et al.

2012). The earliest study on the alkaloids of the Dendrobium genus started from dendrobine of *D.*

nobile. For the decades, dozens of alkaloids have been subsequently identified from *D. anosmum*

(Leander and Lüning 1968), *D. chrysanthum* (Ekevåg et al. 1973), *D. crepidatum* (Elander et al.

1973), *D. findlayanum* (Granelli et al. 1970), *D. friedricksianum* (Hedman et al. 1971), *D.*

hilderbrandii (Elander and Leander 1971), *D. loddigessi* (Li et al. 1991), *D. lohohense* (Inubushi

et al. 1964), *D. moniliforme* (Liu et al. 2007), *D. parishii* (Leander and Lüning 1968), *D. pierardii*

(Elander et al. 1971), *D. primulimun* (Blomqvist et al. 1972), *D. snowflake* (Morita et al. 2000)

and *D. wardianum* (Glomqvist et al. 1973). At present, the alkaloids identified from Dendrobiun

plants had been divided into five categories based on their chemical structures, including

sesquiterpenoid alkaloids, indolizidine alkaloids, pyrrolidine alkaloids, phthalide alkaloids, and

imidazole alkaloids (Xu et al. 2013).

Thus far, the composition and biosynthetic pathway of the alkaloids in *D. officinale* has not

been entirely understood. The determination of the alkaloids between *D. officinale* and *D. nobile*

using an LC-MS method showed that 25 identical compositions in the two Dendrobium plants

were tentatively identified; however, dendrobine was only detected in *D. nobile* (Chen et al. 2006).

Using a UPLC method, a few coumarins, alkaloids and bibenzyl components from fifteen

Dendrobium species were quantitatively determined. Dendrobine was only detected in *D. nobile*

originated from the Guizhou and Yunnan Provinces (Xu et al. 2010). The genome and

transcriptome analyses of *D. officinale* implied that a biosynthetic pathway oriented to the

terpenoid indole alkaloids likely exists (Guo et al. 2013; Yan et al. 2015). Recently,

high-throughput sequencing revealed the identification of a total of 56 genes, including 25 key

enzymes potentially involved in the synthesis of terpenoid indole alkaloids, tropine alkaloids and

isoquinolines alkaloids in *D. officinale* (Niu et al, 2021). Many studies have shown that *CrMYC2*

and *ORCAs* transcription factors play a positive role in elevating the TIAs content in *Catharanthus*

roseus (Memelink and Gantet 2007). With the help of transcriptome sequencing, 75 bHLH family

genes and 66 AP2/ERF genes were identified in the DEGs of *D. officinale,* suggesting that those

transcription factors could be involved in the regulation of secondary metabolism, especially in

TIAs biosynthesis (Zhang et al. 2016). Mevalonate (MVA) and methylerythritol 4-phosphate

(MEP) pathway are both core pathways related to MeJA-induced accumulation of TIAs. In *D.*

officinale, the key genes in MVA and MEP pathway and some post-modified enzymes were

significantly up-regulated by MeJA treatment (Chen et al. 2019). In the previous study,

high-yielding production of alkaloids from *D. officinale* were obtained with the combination of

tryptophan, secologanin and MeJA treatment (Jiao et al. 2018). Here, a technical scheme for the

separation and determination of alkaloids and precursors in *D. officinale* was established. The

accurate mass weights of the components separated by HPLC were determined using an Orbitrap

mass analyzer. The multistage fragmentation mass spectra of alkaloids were monitored to

speculate the potential substances based on a database-dependent retrieval method. Alkaloid and

their precursors related to alkaloid-synthetic pathway were associated to construct the metabolic

network between upstream and downstream. The results would provide a reliable scientific

reference and technical guidance for the development and quality evaluation of *D. officinale*.

Materials and Methods

Instruments and chemicals

The LTQ-Orbitrap XL mass spectrometer were manufactured by Thermo Fisher Scientific

(Bremen, Germany). The Agilent 1260 instrument used in the experiments was manufactured by

Agilent Technologies (Waldbronn, Germany). The Xcalibur (*v.* 2.1) and Mass Frontier (*v.* 6.0)

were used for the analysis of mass spectrometry and speculation of the fragmentation pathways .

Acetonitrile and formic acid were of HPLC grade. Other chemicals were domestically made

analytical reagents. The standards of reserpine and sarpagine was obtained from Aladdin

(Shanghai, China) and BioBioPha (Kunming, China), respectively.

Plant materials preparation

The *D. officinale* samples used in the experiment were obtained from the tissue culture room

of the Plant Cell Engineering Center in West Anhui University. The tissue-cultural seedlings

cultivated in MS medium were collected after 100 μM MeJA treatment. The seedlings were firstly

dried at 110 °C for 15 min and kept in a constant weight at 60 °C. The dried materials were ground

into powders by a universal grinder. Using the ultrasonic extraction method, two gram of powder

was placed in a round-bottom flask with 100 mL of methanol and extracted three times at room

temperature. The bromocresol green colorimetric method was used for the determination of total

alkaloids (Chen et al. 2019). The extraction solution was centrifuged for 10 min at 6000 x g to get

the supernatant, then evaporated to dryness in a rotary evaporator. The extraction was dissolved

with 5 mL of dilute sulfuric acid, filtered through a 0.22 μm membrane, and pretreated using

MCX cartridges (60 mg, 3 mL) for purification. The protocol was successively conducted by our

previous method (Song et al. 2020).

Experimental conditions

The Waters T3 column (150 mm × 4.6 mm, 3 μm) was used for the chromatographic

separation. The mobile phases acetonitrile (A) with 0.1% formic acid and aqueous phase (B) with

0.1% formic acid were used for the experiment. Gradient elution process was as follows: 0-2 min

(0% A), 2-12 min (0 to 15% A), 12-22 min (15 to 35% A), 22-32 min (35 to 80% A), and 32-37

min (80 to 0% A). The UV wavelength was set at 280 nm, and the column temperature was set at

25 °C. A three-way splitter was connected with the mass analyzer at a 0.3 mL min^{-1} flow rate. The

ion-source parameters were refereed to Kumar's method and slightly modified (Kumar et al.,

2015). The detailed parameters were as follows: sheath gas at 25 arb, auxiliary gas at 3 arb, spray

voltage at 4 kV, the capillary temperature at 320 °C, tube lens at 120 V, and capillary voltage at 30

V. The MS data were collected at $100 \leq m/z \leq 1000$ in positive-ion mode. Two data acquisition

methods were used in the experiments. In one method, a high-resolution scan was conducted using

the Orbitrap mass analyzer to acquire the MS data at a resolution at 30,000 FWHM and the MS^2

data at 15,000 FWHM. A data-dependent MS^n scan was used for the analysis of the MS^2 spectra

generated from the most abundant ions of the MS spectra. In the other method, a high-resolution

scan was conducted first to acquire the MS data at a resolution at 30,000 FWHM, then the LTQ

dynode was used to scan the MS^2 and MS^3 spectra. The dynamic exclusion function was used to

reduce repeat scans, and the repeat count was 2. The exclusion duration was 20 s, and the

exclusion mass width was 3 m/z. The suitable collision-induced dissociation (CID) energy was set

at 35%. The minimum signal threshold was 500, and the isolation width of precursor ions was 2

m/z. Sarpagine and reserpine were respectively injected into the mass analyzer via a syringe pump

at a flow rate of 6 μL min^{-1} to complete the direct infusion analysis. The putative molecular

formulae were calculated and obtained according to the accurate molecular weight using Xcalibur

(*v.* 2.1). Compounds were tentatively identified by matching METLIN, Mass Bank, Dictionary of

Natural Products and Chemspider databases. For the compounds without multiple mass spectra,

Mass Frontier (*v.* 6.0) was helped to speculate and deduce the fragmentation pathway.

Results and discussion

Biomass and alkaloid contents under the MeJA treatment

The effects of methyl jasmonate on the growth and alkaloid accumulation of *D. officinale*

was studied. The results showed that MeJA delayed the growth rate of *D. officinale* and initiated

the synthesis of a large number of alkaloids in logarithmic growth phase (**Fig.1**). The highest

content of alkaloids climaxed to 343 μg/g when cultured on the 32 days. After that, the fresh

weight tended to be stable, and the alkaloid content began to fall gradually. The results suggested

that jasmonates could continuously promote the synthesis of alkaloids during the growth of *D.*

officinale. The growth trend and accumulating pattern were consistent with our previous study

(Jiao et al. 2018).

Mass spectra interpretation

Firstly, *D. officinale* samples were extracted by methanol and dissolved with diluted sulfuric

acid. The MCX cartridges were used for the purification of alkaloids from the acidic aqueous

extraction. To optimize the condition of mass spectrometry, the suitable CID energy was achieved

by the direct infusion experiment. Generally, the strongest fragment ions and the smallest number

of fragments were observed at 30-35% collision energy, whereas the abundances of the fragments

decreased substantially at 45% or greater collision energy (Song et al. 2020). In this experiment,

sarpagine and reserpine achieved sufficient fragments both in quantity and intensity at 40%

collision energy. High-resolution mass analysis was conducted by a data-dependent MS^n scan

from the workstation to produce abundant fragmentation spectra of the product ions. Due to the

high resolution, the scan rate of a single spectrum was reduced, and the analytical analysis time of

the entire sample was extended. An appropriate exclusion duration time was required for the

dynamic exclusion function to avoid repeat scans of target ions and thereby shorten the analysis

time of the samples (Zhang et al. 2012). Another important function of dynamic exclusion was to

obtain the fragment information of the co-eluting small peaks, which allowed compounds with

low ionized potentials to be detected (Pan et al. 2015). The total ion current chromatogram of the

methanol extract spiked with standards from *D. officinale* is presented, as obtained in the

positive-ion mode under the optimized mass parameters and analytical conditions (**Fig.S1a**).

Generally, $[M+H]^+$ or $[M]^+$ was achieved through the soft ionization in the positive mode. The

molecular-ion peaks of sarpagine and reserpine were separately extracted at *m/z* 311.1738 and

609.2772. The retention time of sarpagine and reserpine were at 12.95 min and 25.41 min,

respectively (**Fig. S1b** and **S1c**). The results indicated that the established method for the mass

spectrometry was suitable for the separation and characterization of alkaloids.

Using a data-dependent MS^n scan method, the MS spectra of two standards were obtained

from the Orbitrap analyzer, and the MS^n spectra were obtained from the LTQ. It was indicated that

the π bonds on the indole groups of sarpagine and reserpine were easily dissociated to produce

$[M+H]^+$ peaks and produced massive hydrogen rearrangement. The ionization and electron

transfer were observed on the position of the group cleavage because of the induced dissociation

effect of the high-energy collisions. The putative fragmentation modes of sarpagine and reserpine

are presented in Fig. S2 and S3. The characteristic fragments with *m/z* 146 and 174 from the

indole group were produced by the cleavage of sarpagine and reserpine, respectively. The

characteristic ion *m/z* 195 cleaved on the ester bond was obtained to produce the

3,4,5-trimethoxybenzaldehyde group from the aglycone. Finally, forty-nine nitrogen-containing

compounds, including seven alkaloids, were separated and tentatively identified from *D. officinale*

(**Table S1**).

Identification of the alkaloids

The small molecular compounds previously identified from *D. officinale* mainly include

stilbenoids (bibenzyl and phenanthrene), phenols, lignans, nucleotides, lactones, flavonoids,

steroids, and fatty acids (Chen et al. 2013). Based on the high resolution and sensitivity of

LTQ-Orbitrap for the trace substances, coupled with the high selectivity of MCX for the

medium-strength alkaline compounds, seven alkaloids identified from the methanol extract of *D.*

officinale were divided into five classes based on the chemical structure, including tropane

alkaloids, tetrahydroisoquinoline alkaloids, quinolizidine alkaloids, piperidine alkaloids, and

spermidine alkaloids. The special fragmentation modes of different alkaloids in the ESI source

were discussed below.

The tropane alkaloids consist of pyridine conjugated with piperidine. The experiment

involved in DART-MS in-source CID at 90 V indicated that the characteristic ions of tropine were

m/z 158, 142, 124, 93, and 67 (Lesiak et al, 2015). 2,3-Dihydroxynortropane was a kind of

nortropane-type alkaloid that lacks an N-methyl group (Asano et al, 2001). Notably, the hydroxyl

groups were present on its C-2 and C-3 in both the α- and β-configurations, as confirmed by a

benzoate chirality method (Kusano et al. 2002). In this study, a strong [M+H-·OH]$^+$ ion peak was

observed under the set CID conditions. In addition to the characteristic ions m/z 93 and 67,and the

stronger fragments m/z 113, 98 and 82 were observed (**Fig. 2a**). The MS^2 and MS^3 spectra of

2,3-dihydroxynortropane were displayed, and [M+H-·OH]$^+$ ion also appeared (**Fig. 2b** and **2c**).

However, the base peak of 2,3-dihydroxy- nortropane was the 2-methylpyrrole ion, which was

further fragmented to get m/z 56. Finally, the compound was identified as

2,3-dihydroxynortropane or its isomers.Two tetrahydroisoquinoline alkaloids were identified as

alamaridine and 1,2-dihydro-O-methyltazettine. The characteristic ions of tetrahydroisoquinoline

group in alamaridine were m/z 177, 176, and 162 in the EI source, and the methyl group on C-5

was easier to lost compared to that on C-10 (Bhattacharjya et al. 1988). The MS^2 and MS^3 spectra

of alamaridine are displayed respectively (**Fig. 3a** and **3b**). These spectra indicated that the methyl

groups on C-5 and C-10 were lost to produce the base peaks [M+H-2CH$_3$-3H]$^+$, which further

lacked a hydroxyl group to generate m/z 248 in the MS^3 spectrum. The C$_2$H$_3$ and ·OH groups were

successively cleaved to generate m/z 221 and 204 on ring B and m/z 176 and 161 on ring C.

1,2-Dihydro-O-methyltazettine, also known as ungvedine, is a tazettine- type alkaloid (Kadyrov et

al. 1979). The main fragments m/z 207 and 142 were generated by 1,2-dihydro-O- methyltazettine

with equivalent intensity (**Fig. 3c**). The m/z 142 further lacked the methyl group and

simultaneously formed two double bonds to generate the base peak m/z 124 (**Fig. 3d**). Previous

studies have shown that the fragmentation pattern of tazettine-type alkaloid and its C$_3$H$_7$N group

was easily lost to produce the characteristic ion [M+H-57]$^+$ (Zhang et al. 2009). In Fig. 3c,

1,2-dihydro-O- methyltazettine was fragmented by RDA cleavage at both C-6, N-5 and C-4b,

C-6a. 1,2-Dihydro- O-methyltazettine was cleaved at C-4b, C-12a but not on C-3, C-4, and thus

did not produce [M+H-C$_3$H$_7$N]$^+$. There was a coordination bond between nitrogen and oxygen

presented in the quinolizidine alkaloid methyllagerine N-oxide. It was indicated that this bond

easily led to the generation of the fragment [M-O]$^+$ in the EI source (Lee et al. 2011). The oxygen

was easily lost as well to produce the fragment m/z 437 in the ESI source (**Fig. 4a**). The

characteristic ions m/z 308 and 292 were generated by the cleavage of C-11 and C-22. According

to the different cleavage positions on ring B, the characteristic ions m/z 220 and 163 were also

generated. Meefarnine B was a spermidine alkaloid containing a polyamine heterocyclic ring.

These alkaloids are generally considered to be synthesized by putrescine, which is produced by

the degradation of L-arginine or L-ornithine (Silva et al. 2015). The MS^2 and MS^3 spectra of

meefarnine B indicate that the NH$_3$ group on N-5 was cleaved to produce the fragment m/z 421

under the collision dissociation effect (**Fig. 4b** and **4c**). The fragment m/z 292 was generated by

the cleavage of the side chain on N-9 and was further cleaved to produce m/z 218. Besides, the

fragment m/z 275 was generated by the cleavage of the m/z 421 on N-1 and C-2. The fragment m/z

204 was generated by the cleavage of the m/z 275 on N-9 and C-10, which further produced the

characteristic ion m/z 147. In this study, a piperidine alkaloid anapheline was tentatively identified

from *D. officinale*. The retention time of anapheline was 27.88 min, and the molecular weight of

the [M+H]$^+$ was 225.1945, which implies that its molecular formula was $C_{13}H_{25}ON_2$ (**Fig. S4**).

Anapheline easily lost H_2O to produce the fragment m/z 207 and the characteristic ion piperidine

m/z 100 during the collision-induced dissociation. The NH$_3$ group was lost to produce the m/z 83

in the MS^2 spectrum. Based on the MS^2 and MS^3 spectra, a presumed fragmentation pathway of

anapheline was proposed (**Fig. 5**).

Identification of the guanidines

Three guanidine compounds were separated and determined from the methanol extract of *D.*

officinale for the first time. The MS^2 and MS^3 spectra of 4-guanidinobutanoic acid were displayed

respectively (**Fig. 6a** and **6b**). The mass spectra indicated that 4-guanidinobutanoic acid easily lost

H_2O to form the base peak m/z 128 and produced the characteristic ions m/z 104, 87 and 60. The

m/z 128 was further cleaved to produce the m/z 111 and 86. These results were consistent with

those reported by Nikolić et al. (Nikolić et al. 2012). Cyclocimipronidine is a cyclic guanidine

compound separated from *Cimicifuga racemosa*. Some studies showed that the characteristic ions

of cyclocimipronidine, a guanidine alkaloid in black cohosh, were the m/z 112, 95, 94, 70 and 67

(Gödecke et al. 2009) . In addition to the aforementioned characteristic ions, cyclocimipronidine

easily lost H_2O to generate the fragments m/z 136 and 84 (**Fig. 6c** and **6d**). The MS^2 and MS^3

spectra of *N-tert*-butyloxycarbonyl guanidine were displayed in Fig. 6e and 6f, respectively. It was

indicated that the characteristic ions were the m/z 142, 125, 104 and 87.

Identification of the nucleotides and derivatives

Nucleotides were composed of purine or pyrimidine linked with ribose. Adenosine, uridine

and guanosine had been separated from *D. officinale* in a previous study. In this study, five

nucleotide derivatives were tentatively identified. 2-Hydroxyadenine was an adenine derivative

that easily lost an NH_3 group to produce the characteristic ions m/z 135 and 110 (**Fig. S5a**). The

fragments m/z 107, 92 and 77 were generated in the MS^3 spectrum (**Fig. S5b**). Besides

2-hydroxyadenine, another four nucleotide derivatives were identified in the study. Their MS^2 and

MS^3 spectra indicated that the neutral loss of m/z 136 resulted from the loss of aglycones (**Fig. S5**

c-j). Comparison with the multistage mass spectra of the METLIN database

(https://metlin.scripps.edu/) resulted in the identification of adenosine, guanosine,

1-methyladenosine and cytidine.

Identification of the sphingolipids

Sphingolipid is a structurally complex lipid containing a long-chain sphingosine skeleton,

which forms the plasma and vacuole membranes of plant cells. Four phytosphingosines were

separated and identified from *D. officinale*. Because sphingosine contained some hydroxyl groups,

the characteristic ion $[M+H-H_2O]^+$ was observed during the dissociation (Fig. S6a-h). By aligning

against the Dictionary of Natural Products Database (http://dnp.chemnetbase.com), four

sphingolipids, including 2-amino-1,3-hexadecanediol, 2-amino-3-tetradecanol, 2-amino-1,3,4-

octadecanetriol and 2-amino-1,3,4-hexadecanetriol, were identified in the experiment.

Identification of the amino acids and their derivatives

Eight amino acids—L-lysine, L-histidine, L-arginine, L-valine, L-isoleucine, L-tyrosine,

L-phenylalanine and L-tryptophan—were identified from *D. officinale* based on accurate

molecular weights and the multistage mass spectra of standards from the MassBank

mass spectrum library (http://www.massbank.jp/QuickSearch. html) (**Table S1**). 4-Hydroxy-L-

threonine and urocanic acid were derivatives of L-threonine and L-histidine that easily lost H_2O to

produce the base peaks m/z 118 and 121. The putative fragment ions generated during their

cleavage were displayed in **Fig. S7a** and **S7b**, respectively. In addition to the derivatives of

L-threonine and L-histidine, four additional arginine derivatives were identified in this study.

N^2-Fructopyranosylarginine was an arginine monoglycoside and the byproduct generated by a

Maillard reaction between reducing sugars (fructose or glucose) and arginine upon heating

(Suzuki et al. 2004). Because fructose contained several hydroxyl groups, H_2O was easily lost to

produce the characteristic ions m/z 319, 301 and 283. The fragments m/z 257 and 239 were

produced by the cleavage of the guanidino group on the glycoside. N^2-Fructopyranosylarginine

was also cleaved to generate the characteristic ion m/z 175 on the N-glucosidic band and further

produced the aglycone m/z 158 via the neutral loss of fructose (**Fig. S7c-d**). The NH_3 group of

N-methylarginine was lost to produce the base peak m/z 172, as a result of the dissociation.

According to the different cleavages on the R group, the characteristic ions m/z 158, 133, 116 and

74 were generated (**Fig. S7e**). The characteristic ions m/z 126, 115, 97 and 70 were also generated

in the MS^3 spectrum by the cleavage of the m/z 172 fragment (**Fig. S7f**). The ·OH group of

N^α-acetylarginine was lost to produce the characteristic ion m/z 200. The fragments m/z 175 and

157 were generated by the different cleavages on the R group. The characteristic ions m/z 139, 115

and 70 were produced by the cleavage of amino and carboxyl groups linked with the chiral

carbons. The ·OH group of the m/z 200 was lost to produce the m/z 183 in the MS^3 spectrum, and

the other fragments were similar to those in the MS^2 spectrum (**Fig. S7g-h**).

N^2-(3-Hydroxysuccinoyl)arginine was formed by the substitution of the 3-hydroxysuccinoyl group

on the N-terminal of arginine. The typical feature of its cleavage was the loss of

3-hydroxysuccinoyl group to produce the characteristic ion m/z 175 (**Fig. S7i**), which was further

cleaved to generate the fragments m/z 158, 139 and 60 (**Fig. S7j**).

Identification of the isomers

The accurate molecular weights of the compounds were provided by high-resolution mass

spectrometry, which allowed us to deduce the speculative molecular formulas. However, the

identification of isomers was less effective. Two pairs of isomers were identified in this study

based on their different retention times and distinct multistage fragments. The formula of both

L-homophenylalanine and β-phenyl-γ-aminobutyric acid was $C_{10}H_{13}O_2N$, but their retention times

were at 6.79 and 12.47 min, respectively (**Table S1**). The MS^2 spectra of L-homophenylalanine

and β-phenyl-γ-aminobutyric acid are displayed in **Fig. S8a** and **S8c**, respectively. These mass

spectra indicated that the NH_3 groups were easily lost to form the characteristic ion *m/z* 163.

However, the fragment *m/z* 120 produced in the spectrum of β-phenyl-γ-aminobutyric acid was

not observed in that of L-homophenylalanine. The MS^3 spectra presented different fragments,

which indicated two different compounds (**Fig. S8b** and **S8d**). The formula of both carbetamide

and N^δ-benzoylornithine were $C_{12}H_{16}O_3N_2$, but their retention times were at 9.77 and 14.93 min,

respectively (**Table S1**). The MS^2 and MS^3 spectra of carbetamide were displayed in **Fig. S8e** and

S8f, respectively. The MS^2 and MS^3 spectra of N^δ-benzoylornithine were displayed in Fig. S8g and

S8h, respectively. These two isomers exhibited different fragmentation patterns. The characteristic

ions of carbetamide were the *m/z* 120, 103 and 93 ions, whereas the characteristic ions of

N^δ-benzoylornithine were the *m/z* 219, 205, 175, 158 and 148 ions. The isomerism of the

dipeptides resulted from the chirality of the α-carbon and the different positions of peptide bonds

between two peptides. In the study, eight dipeptides were tentatively identified. Most of these

dipeptides had more than one retention time, suggesting that the isomers existed.

Identification of the nitrogen-containing glycosides

By comparing our spectra with the Metlin and Dictionary of Natural Products databases, we

tentatively identified four nitrogen-containing glycosides from *D. officinale*. The neutral loss of

H_2O in 5'-*O*-β-D-glucosylpyridoxine was easy to produce the characteristic ion *m/z* 314. The

fragment *m/z* 171 was generated by the cleavage on the O-glucosidic bond and further produced

the aglycone *m/z* 152 by the neutral loss of glucose (**Fig. S9a**). The H_2O on the aglycone of

4-*O*-β-D-glucopyranoside-2,4-benzoxazolediol was easily lost to produce the base peak *m/z* 296.

The benzoxazole ion *m/z* 136 was generated by the cleavage on the O-glucosidic bond (**Fig. S9b**).

The *m/z* 296 was cleaved to generate the characteristic ions *m/z* 119 and 94 (**Fig. S9c**). The MS^2

and MS^3 spectra of *O*-(tri-*O*-acetyl-α-L- rhamnopyranoside)- (4-hydroxybenzyl) methylcarbamic

acid were displayed in **Fig. S9d** and **S9e**, respectively. These mass spectra indicated that the

characteristic ion *m/z* 322 was generated by the cleavage of the ester bond and the side chain. The

m/z 322 was cleaved to produce the fragments *m/z* 304 and 160 in the MS^3 spectrum. Passicapsin

was composed of one glucose and a nitrogen-containing aglycone that easily produced the

aglycone *m/z* 238 via the neutral loss of glucose. The characteristic ions *m/z* 286 and 148 were

cleaved at the O-glucosidic bond between the side chain and the dideoxy glucoside (**Fig. S9f**). The

fragments m/z 148 and 106 in the MS^3 spectrum were produced by the cleavage of the aglycone

m/z 238 on the O-glucosidic bond. The H$_2$O of the aglycone m/z 238 was lost to produce the

fragments m/z 220 and 212, and the dideoxy glucoside group was further cleaved to generate the

m/z 190 and 174 (**Fig. S9g**).

The metabolic network and putative route of *D. officinale* alkaloids

Generally, the biosynthetic pathways of alkaloids are derived from the catabolism of amino

acids. The biogenesis between the amino acids and the alkaloids has a relationship between

upstream and downstream upstream and downstream. In the experiment，many amino acids and

their derivatives were identified. The metabolic profile of the alkaloids in *D. officinale* was

constructed based on the KEGG pathway database. This profile indicated that the precursors of the

alkaloids were mainly L-phenylalanine, L-arginine, L-tyrosine and L-lysine (**Fig. 7**).

L-Phenylalanine was catalyzed by phenylalanine ammonialyase to form L-homophenylalanine.

3,4-Dihydroxybenzaldehyde was degraded by phenylalanine combined with tyramine to generate

norbelladine, which further formed 1,2-dihydro-*O*-methyltazettine via multistep biochemical

reactions. L-Arginine, as a major amino acid in *D. officinale*, was catalyzed by an

ethylene-forming enzyme to form guanidine, which was the precursor of three guanidine alkaloids.

L-arginine was also catalyzed by arginine amidinase to form L-ornithine, which was further

catalyzed by ornithine decarboxylase to generate putrescine. On the one hand, putrescine was the

precursor used for the synthesis of meefarnine B. On the other hand, putrescine was also catalyzed

by several enzymatic reactions to form tropine and 2,3-dihydroxynortropane. L-Lysine was

catalyzed by lysine decarboxylase to generate cadaverine, which further formed anapheline via a

four-step reaction.

Conclusion

The alkaloids and precursors of *D. officinale* with MeJA treatment were primarily studied by

the SPE-HPLC–LTQ-Orbitrap technique. The extraction and purification of the target compounds

was optimized, and the suitable conditions of ESI-CID-MS^n for the mass spectrometric detection

were established. Based on the accurate mass weight with MS^n fragmentation information

provided by the LTQ-Orbitrap mass spectrometer, forty-nine compounds, including amino acids
derivatives, dipeptides, nucleotides, guanidines, alkaloids, sphingolipids and several
nitrogen-containing glycosides, were tentatively identified from *D. officinale*. Seven alkaloids
were identified from *D. officinale*. The combination of alkaloids and their precursors were
associated to construct a metabolic network for the interpretation of biosynthetic pathways. These
findings will serve as a technical guidance for the analyses and drug discoveries of *Dendrobium*
alkaloids.

Author's contribution

CS and QJ discuss the writing plan. CS and YPZ draft the manuscript. CS, MAM and GHL edit
the manuscript. GHL, CS and QJ acquire the funding.

Funding

This work was supported by Anhui Natural Science Foundation (2108085MC80), High-level
Talents Research Initiation Fund of West Anhui University (WGKQ202001011, WGKQ2021079),
and Quality Engineering Project of Anhui Provincial Department of Education (2020jyxm2137).

Declarations

Conflicts of interest

No interest to declare.

Consent to participate

All of the authors in this paper are informed content to participate in this manuscript.

Availability of data and material

All of the data including raw data sheets can be found in supplementary materials.

References

Asano N, Yokoyama K, Sakurai M, et al (2001) Dihydroxynortropane alkaloids from
calystegine-producing plants. Phytochemistry 57:721–726.
https://doi.org/https://doi.org/10.1016/S0031-9422(01)00131-5

Bhattacharjya, A., Mukhopadhyay, R., Pakrashi SC (1986) Structure and synthesis of alamaridine,
a novel benzopyridoquinolizine alkaloid from alangium lamarck II. Tetrahedron Lett
27:1215–1216. https://doi.org/10.1016/S0040-4039(00)84220-3

Blomqvist L, Leander K, Lüning B, et al (1972) Studies on orchidaceae alkaloids. XXIX. The

absolute configuration of dendroprimine, an alkaloid from Dendrobium primulinum Lindl.

Acta chemica Scandinavica 26:3203-3206. doi:10.3891/acta.chem.scand.26-3203

Bucar F, Wube A, Schmid M (2013) Natural product isolation-how to get from biological material

to pure compounds. Nat Prod Rep 30:525–545. https://doi.org/10.1039/c3np20106f

Chen X, Xiao S, Guo S (2006) Comparison of chemical compositions between Dendrobium

candidum and Dendrobium nobile. Acta Academiae Medicinae Sinicae 28:524-529

Chen X, Wang C, Yang J, et al (2013) Research Progress on Chemical Composition and Chemical

Analysis of Dendrobium officinale. Journal of Chinese Pharmaceutical ences, 48:1634-1640

Chen Y, Wang Y, Lyu P, et al (2019) Comparative transcriptomic analysis reveal the regulation

mechanism underlying MeJA-induced accumulation of alkaloids in Dendrobium officinale. J

Plant Res 132:419–429. https://doi.org/10.1007/s10265-019-01099-6

Da Silva G, Martinho A, Soengas RG, et al (2015) A new spermidine macrocyclic alkaloid

isolated from Gymnosporia arenicola leaf. Fitoterapia 106:7–11.

https://doi.org/10.1016/j.fitote.2015.07.020

Ekevåg U, Elander M, Gawell L, et al (1973) Studies on orchidaceae alkaloids. Part 33. Two new

alkaloids, N-cis- and N-trans-cinnamoyl-norcuskhygrine from Dendrobium chrysanthum

Wall. Acta chemica Scandinavica 27:1982-1986. doi:10.3891/acta.chem.scand.27-1982

Elander M, Gawell L, Leander K (1971) Studies on orchidaceae alkaloids. XXII. Synthesis and

absolute configuration of pierardine. Lactone-betaine isomerization of shihunine. Acta

chemica Scandinavica 25:721-724. doi:10.3891/acta.chem.scand.25-0721

Elander M, Leander K (1971) Studies on orchidaceae alkaloids. XXI. 6-hydroxynobiline, a new

alkaloid from Dendrobium hildebrandii Rolfe. Acta chemica Scandinavica 25:717-720.

doi:10.3891/acta.chem.scand.25-0717

Elander M, Leander K, Rosenblom J, et al (1973) Studies on orchidaceae alkaloids. XXXII.

Crepidine, crepidamine and dendrocrepine, three alkaloids from Dendrobium crepidatum

Lindl. Acta chemica Scandinavica 27:1907-1913. doi:10.3891/acta.chem.scand.27-1907

Glomqvist L, Brandänge S, Gawell L, et al (1973) Studies on orchidaceae alkaloids. XXXVII.

Dendrowardine, a quaternary alkaloid from Dendrobium wardianum Wr. Acta chemica

Scandinavica 27:1439-1441. doi:10.3891/acta.chem.scand.27-1439

Granelli I, Leander K, Lüning B (1970) Studies on orchidaceae alkaloids. XVI. A new alkaloid,

2-hydroxydendrobine, from Dendrobium findlayanum par. et Rchb. f. Acta chemica

Scandinavica 24:1209-1212. doi:10.3891/acta.chem.scand.24-1209

Gödecke T, Lankin DC, Nikolic D, et al (2009) Guanidine alkaloids and Pictet-Spengler adducts

from black cohosh (Cimicifuga   racemosa). J Nat Prod 72:433–437.

https://doi.org/10.1021/np8006952

Guo X, Li Y, Li C, et al (2013) Analysis of the Dendrobium officinale transcriptome reveals

putative alkaloid   biosynthetic genes and genetic markers. Gene 527:131–138.

https://doi.org/10.1016/j.gene.2013.05.073

Hedman K, Leander K, Lüning B (1971) Studies on orchidaceae alkaloids. XXV. N-isopentenyl

derivatives of dendroxine and 6-hydroxydendroxine from Dendrobium friedricksianum

Lindl. and Dendrobium hildebrandii Rolfe. Acta chemica Scandinavica 25:1142-1144.

doi:10.3891/acta.chem.scand.25-1142

Inubushi Y, Tsuda Y, Konita T, et al (1964) Shihunine: a new phthalide-pyrrolidine alkaloid.

Chemical & pharmaceutical bulletin 12:749-750. doi:10.1248/cpb.12.749

Jiao C, Song C, Zheng S, et al (2018) Metabolic profiling of Dendrobium officinale in response to

precursors and methyl jasmonate. Int J Mol Sci 19:. https://doi.org/10.3390/ijms19030728

Kadyrov KA, Abdusamatov A, Yunusov SY (1979) Ungvedine — A new alkaloid fromUngernia

vvedenskyi. Chemistry of Natural Compounds 15:513-514. doi:10.1007/BF00565070

Kalli A, Smith GT, Sweredoski MJ, Hess S (2013) Evaluation and optimization of mass

spectrometric settings during data-dependent   acquisition mode: focus on LTQ-Orbitrap

mass analyzers. J Proteome Res 12:3071–3086. https://doi.org/10.1021/pr3011588

Kumar S, Singh A, Bajpai V, Kumar B (2016) Identification, characterization and distribution of

monoterpene indole alkaloids in Rauwolfia species by Orbitrap Velos Pro mass spectrometer.

J Pharm Biomed Anal 118:183–194. https://doi.org/10.1016/j.jpba.2015.10.037

Kusano G, Orihara S, Tsukamoto D, et al (2002) Five new nortropane alkaloids and six new

amino acids from the fruit of Morus alba Linne growing in Turkey. Chem Pharm Bull

50:185–192. https://doi.org/10.1248/cpb.50.185

Leander K, Lüning B (1968) Studies on Orchidaceae alkaloids. 8. An imidazolium salt from

Dendrobium anosmum Lindl. and Dendrobium parishii Rchb. f. Tetrahedron letters

8:905-908. doi:10.1016/s0040-4039(01)98823-9

Lee I, Youn U, Kim H, et al (2011) Biphenyl and biphenyl ether quinolizidine N -oxide alkaloids

from lagerstroemia indica L. Planta Med 77:2037–2041.

https://doi.org/10.1055/s-0031-1280064

Lesiak AD, Cody RB, Dane AJ, Musah RA (2015) Plant Seed Species Identification from

Chemical Fingerprints: A High-Throughput Application of Direct Analysis in Real Time

Mass Spectrometry. Anal Chem 87:8748–8757.

https://doi.org/10.1021/acs.analchem.5b01611

Li M, Hirata Y, Xu G, et al (1991) Studies on the chemical constituents of Dendrobium loddigesii

rolfe.   Acta pharmaceutica Sinica 26:307-310

Li X, Shen B, Jiang Z, et al (2013) Rapid screening of drugs of abuse in human urine by

high-performance liquid chromatography coupled with high resolution and high mass

accuracy hybrid linear ion trap-Orbitrap mass spectrometry. J Chromatogr A 1302:95–104.

https://doi.org/10.1016/j.chroma.2013.06.028

Liu W, Hua Y, Zhan Z (2007) Moniline, a New Alkaloid from Dendrobium moniliforme. J Chem

Res 6:317–318. https://doi.org/10.1002/chin.200748193

Memelink J, Gantet P (2007) Transcription factors involved in terpenoid indole alkaloid

biosynthesis in Catharanthus roseus. Phytochem Rev 6:353–362.

https://doi.org/10.1007/s11101-006-9051-z

Morita H, Fujiwara M, Yoshida N, Kobayashi J (2000) New Picrotoxinin-type and

Dendrobine-type Sesquiterpenoids from Dendrobium Snowflake 'Red Star.' Tetrahedron

56:5801–5805. https://doi.org/https://doi.org/10.1016/S0040-4020(00)00530-5

Ng TB, Liu J, Wong JH, et al (2012) Review of research on Dendrobium, a prized folk medicine.

Appl Microbiol Biotechnol 93:1795–1803. https://doi.org/10.1007/s00253-011-3829-7

Nikolić D, Gödecke T, Chen SN, et al (2012) Mass spectrometric dereplication of

nitrogen-containing constituents of black cohosh (Cimicifuga racemosa L.). Fitoterapia

83:441–460. https://doi.org/10.1016/j.fitote.2011.12.006

Niu Z, Zhu F, Fan Y, et al (2021) The chromosome-level reference genome assembly for

Dendrobium officinale and its utility of functional genomics research and molecular

breeding study. Acta Pharm Sin B. https://doi.org/10.1016/j.apsb.2021.01.019

Pan H, Yang W, Zhang Y, et al (2015) An integrated strategy for the systematic characterization

and discovery of new indole alkaloids from Uncaria rhynchophylla by

UHPLC/DAD/LTQ-Orbitrap-MS. Anal Bioanal Chem 407:6057–6070.

https://doi.org/10.1007/s00216-015-8777-0

Quifer-Rada P, Vallverdú-Queralt A, Martínez-Huélamo M, et al (2015) A comprehensive

characterization of beer polyphenols by high resolution mass spectrometry

(LC-ESI-LTQ-Orbitrap-MS). Food Chem 169:336–343.

https://doi.org/0.1016/j.foodchem.2014.07.154

Qiu S, Yang W, Shi X, et al (2015) A green protocol for efficient discovery of novel natural

compounds: characterization of new ginsenosides from the stems and leaves of Panax

ginseng as a case study. Anal Chim Acta 893:65–76.

https://doi.org/10.1016/j.aca.2015.08.048

Song C, Jiao C, Jin Q, et al (2020) Metabolomics analysis of nitrogen-containing metabolites

between two Dendrobium plants. Physiol Mol Biol Plants.

https://doi.org/10.1007/s12298-020-00822-1

Suzuki Y, Choi KJ, Uchida K, et al (2004) Arginyl-fructosyl-glucose and Arginyl-fructose,

Compounds Related to Browning Reaction in the Model System of Steaming and

Heat-drying Processes for the Preparation of Red Ginseng. J Ginseng Res 28:143-148.

doi:10.5142/JGR.2004.28.3.143

Xu J, Han Q, Li S, et al (2013) Chemistry, bioactivity and quality control of Dendrobium, a

commonly used tonic herb in traditional Chinese medicine. Phytochem Rev 12:341–367.

https://doi.org/10.1007/s11101-013-9310-8

Xu J, Zhao W, Qian Z, et al (2010) Fast determination of five components of coumarin, alkaloids

and bibenzyls in   Dendrobium spp. using pressurized liquid extraction and

ultra-performance liquid chromatography. J Sep Sci 33:1580–1586.

https://doi.org/10.1002/jssc.201000034

Yan L, Wang X, Liu H, et al (2015) The Genome of Dendrobium officinale Illuminates the

Biology of the Important Traditional Chinese Orchid Herb. Mol Plant 8:922–934.

https://doi.org/10.1016/j.molp.2014.12.011

Zhang J, He C, Wu K, et al (2016) Transcriptome analysis of dendrobium officinale and its

application to the identification of genes associated with polysaccharide synthesis. Front

Plant Sci 7:. https://doi.org/10.3389/fpls.2016.00005

Zhang X, Huang H, Liang X, et al (2009) Analysis of Amaryllidaceae alkaloids from Crinum by

high-performance liquid chromatography coupled with electrospray ionization tandem mass

spectrometry. Rapid Commun Mass Spectrom 23:2903–2916.

https://doi.org/10.1002/rcm.4205

Figure captions

Fig. 1. The fresh weight and alkaloid contents of *D. officinale* treated with 100 μM MeJA.

Fig. 2. The mass spectra of tropine and 2,3-dihydroxynortropane. (a) The MS^2 spectrum of tropine,

(b) the MS^2 spectrum of 2,3-dihydroxynortropane or an isomer, and (c) the MS^3 spectrum of

2,3-dihydroxynortropane or an isomer.

Fig. 3. The mass spectra of alamaridine and 1,2-dihydro-*O*-methyltazettine. (a) The MS^2 spectrum

of alamaridine, (b) the MS^3 spectrum of alamaridine, (c) the MS^2 spectrum of

1,2-dihydro-*O*-methyltazettine, and (d) the MS^3 spectrum of 1,2-dihydro-*O*-methyltazettine.

Fig. 4. The mass spectra of methyllagerine *N*-oxide and meefarnine B. (a) The MS^2 spectrum of

methyllagerine *N*-oxide, (b) the MS^2 spectrum of meefarnine B, and (c) the MS^3 spectrum of

meefarnine B.

Fig. 5. The putative fragmentation pathway of anapheline.

Fig. 6. The mass spectra of the guanidines. (a) The MS^2 spectrum of 4-guanidinobutanoic acid, (b)

the MS^3 spectrum of 4-guanidinobutanoic acid, (c) the MS^2 spectrum of cyclocimipronidine, (d)

the MS^3 spectrum of cyclocimipronidine, (e) the MS^2 spectrum of *N-tert*-butyloxycarbonyl

guanidine, and (f) the MS^3 spectrum of *N-tert*-butyloxycarbonyl guanidine.

Fig. 7. The putative metabolic network of alkaloids and precursors. The red words denote the

identified compounds. The blue words denote the metabolites. The purple words denote the

enzymes. The solid arrows denote one-step reactions, and the dashed arrows denote multiple

reactions.

F**ig. S1**. The mass spectrum of *D. officinale* extract with the standards. (a) The total ion current

chromatogram of the methanol extract from *D. officinale* spiked with standards, (b) the extracted

ion current chromatogram of *m/z* 311.1738, and (c) the extracted ion current chromatogram of *m/z*

609.2772.

Fig. S2. The putative fragmentation pathway of sarpagine. π denotes π-bond dissociation; i

denotes inductive cleavage; rH$_R$ denotes charge-remote rearrangement; rH$_B$ denotes

α,β-charge-site rearrangement; rH$_C$ denotes γ-charge-site rearrangement; and rH$_{1,2}$ denotes

radical-site rearrangement.

Fig. S3. The putative fragmentation pathway of reserpine. π denotes π-bond dissociation; i denotes

inductive cleavage; rH$_R$ denotes charge-remote rearrangement; and rH$_C$ denotes γ-charge-site

rearrangement.

Fig. S4. The chromatogram and mass spectra of anapheline. (a) The extracted ion chromatogram

of anapheline, (b) the MS spectrum of anapheline, (c) the MS^2 spectrum of anapheline, and (d) the

MS^3 spectrum of anapheline.

Fig. S5. The mass spectra of nucleotides and derivatives. (a) The MS^2 spectrum of

2-hydroxyadenine, (b) the MS^3 spectrum of 2-hydroxyadenine, (c) the MS^2 spectrum of adenosine,

(d) the MS^3 spectrum of adenosine, (e) the MS^2 spectrum of guanosine, (f) the MS^3 spectrum of

guanosine, (g) the MS^2 spectrum of 1-methyladenosine, (h) the MS^3 spectrum of

1-methyladenosine, (i) the MS^2 spectrum of cytidine, and (j) the MS^3 spectrum of cytidine.

Fig. S6. The mass spectra of sphingolipids. (a) The MS^2 spectrum of 2-amino-1,3- hexadecanediol,

(b) the MS^3 spectrum of 2-amino-1,3- hexadecanediol, (c) the MS^2 spectrum of

2-amino-3-tetradecanol, (d) the MS^3 spectrum of 2-amino-3-tetradecanol, (e) the MS^2 spectrum of

2-amino-1,3,4-octadecanetriol, (f) the MS^3 spectrum of 2-amino-1,3,4-octadecanetriol, (g) the MS^2

spectrum of 2-amino-1,3,4- hexadecanetriol, and (h) the MS^3 spectrum of

2-amino-1,3,4-hexadecanetriol.

Fig. S7. The mass spectra of amino acids and their derivatives. (a) The MS^2 spectrum of

4-hydroxy-L-threonine, (b) the MS^2 spectrum of urocanic acid, (c) the MS^2 spectrum of

N^2-fructopyranosylarginine, (d) the MS^3 spectrum of N^2-fructopyranosylarginine, (e) the MS^2

spectrum of N-methylarginine, (f) the MS^3 spectrum of N-methylarginine, (g) the MS^2 spectrum of

N^α-acetylarginine, (h) the MS^3 spectrum of N^α-acetylarginine, (i) the MS^2 spectrum of

N^2-(3-hydroxysuccinoyl)arginine, and (j) the MS^3 spectrum of N^2-(3-hydroxysuccinoyl)arginine.

Fig. S8. The comparative mass spectrum of some isomers. (a) The MS^2 spectrum of

L-homophenylalanine, (b) the MS^3 spectrum of L-homophenylalanine, (c) the MS^2 spectrum of

β-phenyl-γ-aminobutyric acid, (d) the MS^3 spectrum of β-phenyl-γ-aminobutyric acid, (e) the MS^2

spectrum of carbetamide, (f) the MS^3 spectrum of carbetamide, (g) the MS^2 spectrum of

N^δ-benzoylornithine, and (h) the MS^3 spectrum of N^δ-benzoylornithine.

Fig. S9. The mass spectra of nitrogen-containing glycosides. (a) The MS^2 spectrum of

5'-O-β-D-glucosylpyridoxine, (b) the MS^2 spectrum of 4-O-β-

D-glucopyranoside-2,4-benzoxazolediol, (c) the MS^3 spectrum of

4-O-β-D-glucopyranoside-2,4-benzoxazolediol, (d) the MS^2 spectrum of

O-(tri-O-acetyl-α-L-rhamnopyranoside)-(4-hydroxy benzyl)methylcarbamic acid, (e) the MS^3

spectrum of O-(tri-O-acetyl-α-L-rhamnopyranoside)-(4-hydroxy benzyl)methylcarbamic acid, (f)

the MS^2 spectrum of passicapsin, and (g) the MS^3 spectrum of passicapsin.

Table S1. Identification of the main chemical compounds from *D.officinale*

a

b

c