The interplay between developmental stage and environment determines the adaptive effect of a natural transposable element insertion

Miriam Merenciano[1] and Josefa González[1]

[1]Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra), Barcelona, Spain.

*Josefa González

Email: josefa.gonzalez@ibe.upf-csic.es

Author Contributions: MM and JG designed research; MM performed research; MM and JG analyzed data; and MM and JG wrote the paper

Competing Interest Statement: The authors declare no competing interest

Classification: Biological Sciences. Evolution.

Keywords: cold-stress, immune-stress, Drosophila

ABSTRACT

Establishing causal links between adaptive mutations and their ecologically relevant phenotypes is key to understanding adaptation, a central goal in evolutionary biology. Although progress has been made, the number of causal adaptive mutations identified so far is still limited as gene by gene, and gene by environment interactions, among others, complicates linking genetic variation with its fitness–related effects. Transposable elements, often ignored in the quest for the genetic basis of adaptive evolution, are known to be a genome-wide source of regulatory elements across organisms that at times can lead to adaptive phenotypes. In this work, we combine gene expression, *in vivo* reporter assays, CRISPR/Cas9 genome editing and survival experiments to characterize in detail the molecular and phenotypic consequences of a natural *Drosophila melanogaster* transposable element insertion: the *roo* solo-LTR *FBti0019985*. This transposable element provides an alternative promoter to the transcription factor *Lime,* involved in cold- and immune-stress responses, and acts as an enhancer in response to immune-stress. We found that the effect of *FBti0019985* on *Lime* expression depends on the interplay between the developmental stage and the environmental conditions. We further establish a causal link between the presence of *FBti0019985* and increased survival to cold- and immune-stress. Our results exemplify how several developmental stages and environmental conditions need to be considered to characterize the molecular and functional effects of a genetic variant, and add to the growing body of evidence that transposable elements can induce complex mutations with ecologically relevant effects.

SIGNIFICANT STATEMENT

Transposable elements (TEs) are repetitive DNA sequences with the ability to move along the genome. TEs are a genome-wide source of regulatory elements capable of regulating the expression of nearby genes. TEs can act as enhancers in specific tissues or developmental stages, and can also regulate the expression of nearby genes depending on the environmental context. In this work, we characterize a natural TE insertion whose effects on the expression of its nearby gene depends on the interplay between the developmental stage and the environmental conditions. A precise deletion of this TE insertion in natural populations using the CRISPR/Cas9 editing tool further proves that the insertion is the causal mutation leading to increased survival to cold- and immune-stress.

INTRODUCTION

Adaptive processes are pervasive in nature and its study is even more relevant in the current scenario of rapid environmental change (1). Establishing causal links between mutations and their relevant fitness-related phenotypes is crucial to understand how organisms adapt to the environment (1-3). Besides identifying the adaptive mutation at the DNA level, and the fitness-related trait in a relevant ecological context, pinpointing the molecular mechanism by which the adaptive mutation influences the phenotype is key. However, causal links are difficult to establish since the effect of a mutation can depend on the genetic background and on the environmental conditions (4-6). Moreover, a single mutation can affect different phenotypes, thus characterizing the effect of a mutation should imply exploring multiple phenotypes (1, 3, 7, 8).

Different types of mutations might differ in the complexity of their molecular and fitness-related effects, as has been suggested for transposable element (TE) induced mutations (9-11). TEs can affect gene structure and gene expression through many different molecular mechanisms (11, 12), and some of these changes have been associated with adaptive phenotypes (13-19). In Drosophila, a dual role as enhancer and promoter of an adaptive insertion conferring tolerance to bacterial infection has been recently reported (19). There are also several examples of TE-induced mutations affecting more than one phenotype, such as the disruption of the *Chkov1* gene leading to resistance to pesticides and to viral infection (20, 21). Indeed, *D. melanogaster* is an unrivaled model organism to investigate the complexity of genotype-phenotype links because it has one of the best functionally annotated genomes and powerful tools to genetically modify the organism *in vivo,* which allows not only to identify the molecular mechanism behind the adaptive effect of a particular TE insertion, but also to demonstrate its causality (22).

FBti0019985 is a *D. melanogaster roo* solo-LTR element inserted in the promoter region of *Lime*, which is located in the first intron of *cbx* gene (Fig. 1). *Lime* is a C2H2-type zinc finger transcription factor associated with chill-coma resistance and immune response (23, 24) and *cbx*, is a ubiquitin-conjugating enzyme that has also been associated with immune-stress (19, 25). There is evidence showing that *FBti0019985* has increased in frequency due to positive selection (26). Previous studies suggest that *FBti0019985* is associated with changes in expression of its two nearby genes: up-regulation of *Lime* in embryos under nonstress conditions, and

downregulation of *cbx* in adults in nonstress and immune-stress conditions (19, 26). However, these results were based on the analysis of a limited number of genetic backgrounds, and whether *FBti0019985* is indeed the causal mutation, and if that is the case, whether it also affects *Lime expression* under stress conditions are still open questions. The molecular mechanism underpinning *FBti0019985* effects on gene expression is also not clear yet. While there is evidence for *FBti0019985* providing an alternative transcription start site to *Lime* in nonstress conditions (26, 27), whether this new transcript is also present in stress conditions and if it contributes to the phenotypic effect of the mutation is unknown. Similarly, while *FBti0019985* has been reported to act as an enhancer in response to immune-stress in adults (28), whether the TE also acts as an enhancer in other environmental conditions is also unknown. Finally, the evidence for a role of *FBti0019985* in egg-to-adult viability is based on association analysis (26), thus whether *FBti0019985* is the causal mutation and whether it also affects immune-stress survival is still unknown.

In this work, we characterized in detail the molecular and phenotypic effects of *FBti0019985* under cold- and immune-stress conditions, both of them relevant for *D. melanogaster* in nature. We performed gene expression analysis, *in vivo* reporter assays, and CRISPR/Cas9 genome editing in natural strains to investigate the role of *FBti0019985* on the expression of its nearby genes and the plausible molecular mechanisms. Furthermore, to check whether *FBti0019985* plays a role in different stress responses, we also performed phenotypic experiments using laboratory outbred populations and CRISPR/Cas9-mutant strains which allowed us to establish a causal link between the presence of the insertion and increased survival in different environmental conditions.

Figure 1. **Schematic representation of the genomic region where *FBti0019985* is inserted**. *Lime* and *cbx* genes are represented in yellow and green, respectively. Light boxes represent UTR regions and darker boxes represent exons. *FBti0019985* insertion is depicted in red. Transcription start sites, including the one that *FBti0019985* adds, are represented as black arrows. Transgenic insertion sites are represented with blue triangles.

RESULTS

Lime expression changes affect immune- and cold-stress survival and *cbx* expression changes affect immune-stress survival

To further confirm the role of *Lime* in cold- and immune-stress and of *cbx* in immune-stress, we performed survival experiments in these two stress conditions. Previous evidence for a role of *Lime* in immune-stress was obtained in larvae after wasp infection (24). To test whether *Lime* down-regulation is also associated with increased sensitivity to bacterial infection in adult flies, we exposed *Lime* mutant flies to *P. entomophila*. We first confirmed that *Lime*[MI02952] mutant down-regulates *Lime* expression (Fig. 2A, Table S1A and S1B), and we also discarded that *Lime*[MI02952] affected the expression of *cbx*, as mutations in this genomic region have previously been suggested to affect expression of both genes (Fig. 2A, Table S1B) (19, 26). As expected, we found that *Lime* mutant flies, both male and females, were more sensitive to *P. entomophila* infection compared with wild-type flies (Fig. 2B, Table S1C).

Lime is up-regulated in strains selected for cold-tolerance, and up-regulation of *Lime* in embryos is associated with increased egg-to-adult viability in nonstress and cold-stress conditions (23, 26). Consistent with these results, we found that *Lime*[MI02952] flies, which down-regulate *Lime,* showed reduced egg-to-adult viability in nonstress and in cold-stress conditions (Fig. 2C, Table S1D).

Evidence for a role of *cbx* in immune response was based on the analysis on two mutant stocks: *cbx*[c00428] and *cbx*[0741-G4] (Fig 1). However, although *cbx*[c00428] mutant flies were associated with increased sensitivity to the gram positive bacteria *S. aureus* (25), this mutant does not affect *cbx* expression (Table S1B) (19). On the other hand, we found that *cbx*[0741-G4], a null mutant associated with increased tolerance to *P. entomophila* infection (19), also down-regulates *Lime* expression (Fig. 2D, Table S1B). These results suggest that *cbx* indeed plays a role in immune

response as *Lime* down-regulation alone is associated with increased sensitivity to *P. entomophila* infection (Fig. 2B, Table S1C), while *Lime* downregulation and knockout of *cbx* expression in the same genetic background was associated with increased tolerance to infection (Fig. 2E, Table S1C).

Overall, our results add to previous evidence suggesting that changes in *Lime* expression affect cold- and immune-stress responses, and changes in *cbx* expression affects immune-stress response (19, 23-25).

Figure 2. *Lime* mutants are more sensitive to *P. entomphila* infection and have reduced egg-to-adult viability under cold-stress. A) Normalized expression of *Lime* and *cbx* with *Act5C* in the *Lime^{MI02952}* mutant strain compared with the WT strain. **B)** Survival curves of *Lime^{MI02952}* mutant males and females compared with the WT strain. Survival curves in non-infected conditions are depicted as dotted lines while survival curves after *P. entomophila* infection are depicted as continuous lines. Error bars represent SEM. **C)** Egg-to-adult viability in nonstress and in cold-stress conditions of *Lime^{MI02952}* mutants compared with the WT strain. **D)** Normalized expression of *Lime* with *Act5C* in the *cbx$^{0741-G4}$* mutant strain compared with the WT strain. **E)** Survival curves of *cbx$^{0741-G4}$* mutant males and females compared with the WT strain. Survival curves in non-infected conditions are depicted as dotted lines while survival curves after *P. entomophila* infection are depicted as continuous lines. Error bars represent SEM.

FBti0019985 is the causal mutation inducing _Lime_ up-regulation in adult flies under immune-stress

Previous evidence for an association between _FBti0019985_ and changes in expression of _Lime_, and _cbx_ were based on the analysis of inbred and isofemale strains (19, 26). To further test the association between the presence of _FBti0019985_ and changes in the level of expression these two genes, we constructed two outbred populations, one with the _FBti0019985_ insertion and one without it, and re-assessed the level of expression of _cbx_ and _Lime_.

We found that _cbx_ expression was not significantly affected by the insertion genotype (presence/absence of _FBti0019985_), the experimental condition (nonstress _vs_. immune-stress) or the interaction between the genotype and the experimental condition (ANOVA p-values > 0.05;

Fig. 3A and Table S2), suggesting that the TE does not affect *cbx* expression. We thus focused on the effect of *FBti0019985* on *Lime* for the rest of this work.

While no differences in *Lime* expression levels between outbred flies with and without the insertion were found in nonstress conditions, the outbred population with *FBti0019985* had increased *Lime* expression under immune-stress conditions, both in male and female guts (Fig. 3B, and Table S2). To confirm that the *FBti0019985* is the mutation causing increased *Lime* expression, we measured *Lime* levels in a natural strain with a precise deletion of *FBti0019985* that was generated using the CRISPR/Cas9 technique (see Material and Methods). While no differences in expression levels of *Lime* were found in nonstress conditions, under immune-stress, the CRISPR/Cas9 mutant showed a *Lime* down-regulation compared with the outbred strain with *FBti0019985*, as expected if this insertion is the causal mutation (Fig. 3B and Table S2).

Overall, we confirmed that *FBti0019985* was associated with *Lime* up-regulation under immune-stress conditions, and we established using CRISPR/Cas9 to precisely delete the insertion, that *FBti0019985* was the causal mutation (Fig. 3B).

Figure 3. *FBti0019985* is associated with *Lime* up-regulation in guts under immune-stress conditions and with increased tolerance to *P. entomophila* infection. **A)** Normalized expression of *cbx* with *Act5C* in female guts in nonstress and after *P. entomophila* infection. Two-way ANOVA: Genotype, G (presence/absence of *FBti0019985*) p=0.687; Experimental condition, EC (nonstress *vs*. immune stress) p=0.138; Interaction, GxEC p=0.219. **B)** Normalized expression of *Lime* with *Act5C* in nonstress and immune-stress conditions of female and male guts from outbred populations with and without *FBti0019985,* and for CRISPR/Cas9-mutant flies. Two-way ANOVA outbred females: G p<0.001, EC p<0.001, GxEC p<0.001. Two-way ANOVA CRISPR/Cas9 mutant females: G p<0.001, EC p<0.001, GxEC p<0.001. Two-way ANOVA outbred males: G p=0.002, EC p<0.001, GxEC p=0.001. Two-way ANOVA CRISPR/Cas9 mutant males: G p=0.001, EC p<0.001, GxEC p=0.001. **C)** Normalized expression of the reporter gene *lacZ* with *Act5C* in embryos under nonstress and immune-stress conditions from transgenic flies. Two-way ANOVA *FBti0019985*: G p=0.009, EC p<0.065, GxEC p=0.026. Two-way ANOVA *FBti0019985* without TFBSs: G p=0.036, EC p<0.024, GxEC p=0.084. **D)** Normalized expression of the reporter gene *lacZ* with *Act5C* in transgenic

adult female flies under nonstress conditions and after an ethanol exposure. Two-way ANOVA *FBti0019985*: G p=0.272, EC p<0.669, GxEC p=0.494. **E)** Survival curves of outbred flies with and without *FBti0019985* and **F)** CRISPR/Cas9-mutant flies after *P. entomophila* infection. Survival curves in non-infected conditions are depicted as dotted lines while survival curves after *P. entomophila* infection are depicted as continuous lines. Error bars represent SEM. Log-rank p-values < 0.05.

***FBti0019985* harbors functional transcription factor binding sites related with immune response**

While *FBti0019985* was previously described to act as an enhancer in guts after an infection with *P. entomophila* (28), the molecular mechanism is still unknown. *FBti0019985* sequence has three predicted binding sites for *DEAF-1*, *tin*, and *Dorsal* transcription factors, which are related with immune response (26, 28). We thus tested whether these immune-related transcription factor binding sites (TFBSs) could be responsible for the previously described enhancer activity, by using directed mutagenesis to delete them from the *FBti0019985* sequence (see Material and Methods). As expected, the complete sequence of *FBti0019985* drives the expression of the reporter gene only in infected conditions (Fig. 3C, Table S3). When we deleted the three immune-related TFBSs, the expression of the reporter gene was reduced, suggesting that the mutated binding sites were responsible for the enhancer activity of *FBti0019985* in infected conditions (Fig. 4C, Table S3).

To further test that the enhancer effect is stress-specific, we measured the expression of the reporter gene in flies exposed to ethanol-stress. We found no differences in expression under nonstress and ethanol-stress conditions (Fig. 3D, Table S3). These results confirmed that *FBti0019985* is not acting as an enhancer in nonstress conditions and suggested that the enhancer effect of *FBti0019985* is stress-specific.

FBti0019985 has also been previously reported to provide an alternative transcription start site to *Lime* in embryos in nonstress conditions (26, 27). This alternative transcript includes 50 bp of the *FBti0019985* sequence and it is 75 bp longer compared with the transcript that does not start in the TE. We found that the alternative transcript was also present in the gut of adult flies under both nonstress and under immune-stress conditions (Table S2). A transcript-specific qRT-PCR showed that the expression level of the transcript that starts in the TE does not increase significantly under infected conditions (t-test p-value > 0.352). Moreover, if we compared the expression level of the transcript that starts in the TE with the total *Lime* expression, we found that its contribution, both in nonstress and immune-stress conditions, is very low: <1% and 4%, respectively (Table S2).

Overall, we demonstrate that *FBti0019985* harbors functional TFBSs related to the immune response that are responsible for its enhancer activity in infected conditions, and that this enhancer activity is stress-specific. Although *FBti0019985* is adding an alternative transcription start site, the transcript that starts in the TE does not significantly contribute to the increased expression of *Lime* in infected conditions.

Figure 4. *FBti0019985* is associated with *Lime* up-regulation in embryos and with increased viability in both nonstress and cold-stress. A) Normalized expression of *Lime* with *Act5C* in nonstress and cold-stress conditions in embryos from outbred populations*,* and for the CRISPR/Cas9-mutant strain. Two-way ANOVA outbred: G p=0.018, EC p=0.142, GxEC p=0.007. Two-way ANOVA CRISPR/Cas9 mutant: G p=0.003, EC p=0.056, GxEC p=0.029. **B)** Normalized expression of the reporter gene *lacZ* with *Act5C* in embryos under nonstress and cold-stress conditions from transgenic flies. Two-way ANOVA: G p=0.002, EC p=0.001, GxEC p=0.012. **C)** Normalized expression of the reporter gene *lacZ* with *Act5C* in embryos under cold-stress conditions from transgenic flies. T-test p-value=0.035. **D)** Egg-to-adult viability in nonstress and in cold-stress conditions of outbred populations and CRISPR/Cas9 mutant. Two-way ANOVA outbred: G p<0.001, EC p<0.001, GxEC p<0.001. Two-way ANOVA CRISPR/Cas9 mutant: G p<0.001, EC p<0.001, GxEC p<0.001.

FBti0019985 increases tolerance to _P. entomophila_ infection

To test whether the up-regulation of _Lime_ in outbred flies with _FBti0019985_ under immune-stress conditions affects fly survival, we performed infection tolerance assays with _P. entomophila_. In outbred populations, we found that both female and males flies with _FBti0019985_ were more tolerant to infection than flies without the insertion (Fig. 3E, Table S4). We repeated the infection tolerance assay including this time the CRISPR/Cas9-mutant strain which has a precise deletion of the _FBti0019985_ insertion. As before, we observed that outbred flies with _FBti0019985_ were more tolerant to infection compared with outbred flies without the insertion in both females and males (Fig. 3F, Table S4). And, as expected if _FBti0019985_ was the causal mutation, the CRISPR/Cas9-mutant strain was more sensitive to infection compared to flies with _FBti0019985_ both in females and males (Fig. 3F, Table S4).

FBti0019985 drives _Lime_ up-regulation in embryos under nonstress conditions

As mentioned above, _FBti0019985_ is associated with up-regulation of _Lime_ in nonstress conditions in embryos from different inbred and isofemale strains (26). We confirmed that embryos from the outbred population with _FBti0019985_ showed _Lime_ up-regulation in nonstress conditions compared to embryos from the outbred population without the insertion (Fig. 4A, Table S2). However, we found no differences in _Lime_ expression in cold-stress conditions associated with the presence of the TE (Fig. 4A, Table S2), further suggesting that the effect of the TE is stress-specific. As expected if _FBti0019985_ is the causal mutation, CRISPR/Cas9-mutant embryos showed reduced expression levels of _Lime_ in nonstress conditions (Fig. 4A, Table S2).

Lime up-regulation in embryos is likely not due to functional binding sites in _FBti0019985_ sequence

Consistent with the effect of _FBti0019985_ in outbred populations, we found that transgenic embryos containing the _FBti0019985_ sequence showed significantly increased reporter gene expression only in nonstress conditions (Fig. 4B, Table S3). In-silico predictions found TFBSs related with developmental processes in the _FBti0019985_ sequence that could be responsible for this enhancer activity of the TE in embryos (28). We thus performed an _in vivo_ reporter assay

deleting the seven development-related predicted binding sites from the TE sequence (*Dorsal*, *Nub*, *ara/mirr* (x2), *Bap*, *Vnd*, and *Btd*) (see Material and Methods). Contrary to our expectations, we did not find reduced expression of the reporter gene when the binding sites were removed from the *FBti0019985* sequence suggesting that, although some of these binding sites might be functional, they were repressing the expression of the reporter gene (Fig. 4C, Table S3).

Finally, the level of expression of the transcript that starts in the *FBti0019985* insertion was not significantly different in nonstress and cold-stress conditions (t-test p-value= 0.174.). In embryos, the contribution to the total *Lime* expression of the transcript that starts in the TE was higher than the one observed in adults (13.6% and 10.2% in nonstress and stress, respectively *vs.* <1% and 4%; Table S2), as expected since *roo* elements are known to provide embryonic promoters (27).

FBti0019985 increases egg-to-adult viability under nonstress and cold-stress conditions

FBti0019985 was previously associated with increased egg-to-adult viability in nonstress and in cold-stress conditions in three different genetic backgrounds (26). We thus subjected the outbred strains with and without the element to the same cold-stress conditions and measured egg-to-adult viability. We found that outbred flies with *FBti0019985* had increased egg-to-adult viability under cold-stress (Fig. 4D, Table S5). As expected if *FBti0019985* is the causal mutation, CRISPR/Cas9-mutants showed reduced viability compared with the outbred strain with the element in nonstress and cold-stress conditions (Fig. 4D, Table S5).

DISCUSSION

In this work, we characterized a naturally occurring TE-induced mutation and found that the effect of the mutation depends on the interplay between the environmental conditions and the developmental stage. While in nonstress conditions, the *FBti0019985* insertion only affects *Lime* expression in embryos, in stress conditions *FBti0019985* affects *Lime* expression in adults, and specifically in response to immune-stress (Fig. 3 and 4). CRISPR/Cas9-mediated deletion of the insertion in a natural population confirmed that the TE is the causal mutation leading to gene up-regulation in nonstress in embryos and in immune-stress in adults. Furthermore, we showed that

FBti0019985 is the causal mutation leading to increased bacterial infection tolerance and increased viability in nonstress and cold-stress conditions (Fig. 3 and 4). Previous evidence for the complex effects of TE-induced mutations were mostly associated with coding mutations (29, 30). For instance, *Syncytin-1*, a human-endogenous retrovirus (HERV) gene with a clear role in human placental formation (31, 32), has also been associated with the development of neuropsychological disorders and multiple sclerosis under certain environmental stress conditions such as infection, or drug application (29, 33). Similarly, *Supressyn*, another gene from a retroviral origin also involved in placental development, has been recently linked with resistance to viral infection, demonstrating that its effect is context-dependent (30). Our results thus showed that besides TE-induced coding mutations, the effect of TE-induced regulatory mutations is also complex, and further suggest that TEs could be more likely to have complex effects compared to SNPs (10, 34).

Besides adding an alternative transcription start site to *Lime, FBti0019985* also acts as an enhancer (26, 27). There is increasing evidence across organisms that regulatory regions can have a dual function as both promoters and enhancers (35, 36). Indeed, in Drosophila, 4.5% of enhancer sequences overlap with transcription start sites (37, 38). TE insertions have been documented to contribute to gene regulation by functioning as enhancers (39, 40) and promoters (27, 41), and recently a dual role as both enhancer and promoter in immune-stress conditions has been described for *FBti0019386*, a natural TE insertion in *D. melanogaster* (19). However, unlike *FBti0019386,* the alternative transcript starting in *FBti0019985* does not seem to have a significant contribution to the changes in gene expression in the stages and conditions analyzed in this work. Thus, characterization of other TEs with the potential to increase transcript diversity and to act as enhancers is needed to evaluate the contribution of this type of mutations to complex regulatory regions.

Our results also suggest that the precise molecular mechanism by which a TE increases expression of its nearby gene could be different in nonstress *vs.* stress conditions While we found that the insertion harbors functional binding sites for immune-related transcription factors (Fig. 3C), a mechanism other than adding binding sites appear to be responsible for the enhancer role in embryonic stages (Fig. 4C). Although we cannot discard that binding sites not identified in this work are responsible for the enhancer activity of *FBti0019985* in embryos, other

mechanisms such as adding histone marks could also be responsible for the observed changes in expression (19, 42, 43).

Finally, we speculate that the effect of *Lime* on sugar metabolism might underlay the increased immune- and cold-stress survival observed (Fig 3. and Fig. 4) (24). *Lime* affects the levels of glucose and trehalose, which has been shown to affect immune cell proliferation and activation (24). Moreover, changes in glucose and trehalose have been reported in cold-shock and rapid cold hardening treatments in *D. melanogaster* (44, 45). Thus, increased basal levels of *Lime* expression in embryos might be beneficial when we expose them to cold-stress. Indeed, distinct basal transcriptional states have been associated with different phenotypic responses to enteric infection and also to cold-stress in *D. melanogaster* (23, 46). However, further experiments are needed to link the effect of *FBti0019985* on *Lime* expression with changes in sugar metabolism and its subsequent effects on cold and immune tolerance.

Overall, this work provides evidence for *FBti0019985* being the causal mutation responsible for *Lime* up-regulation, whose effect is determined by an interplay between the developmental stage and the environment. We further demonstrate that *FBti0019985* increases viability in cold-stress and immune-stress. These results show the complexity of the genotype-phenotype mapping and the importance of considering the effect of a candidate mutation under different contexts to better characterize its adaptive effects.

MATERIAL AND METHODS

Fly stocks

Fly stocks were reared on standard fly food medium in a 12:12 hour light/dark cycle at 25ºC.

Laboratory mutant and RNAi knock-down strains

We used three laboratory mutant strains and one RNAi knock-down strain that were likely to affect *Lime* and/ or *cbx* genes. We used *Lime^{MI02952}* mutants (Bloomington Drosophila Stock Center (BDSC) stock number #36170) that contains a MiMIC insertion in the first exon of *Lime* gene and a RNAi knock-down strain for the same gene (stock number #33735) (Table S1A). We

also used *cbx^0741-G4* mutants (stock number #63767) and *cbx^c00428* mutants (stock number #10067) with a PiggyBac insertion in the first and third intron of *cbx*, respectively. *Lime* and *cbx* expression levels were measured in all the strains mentioned above and only *Lime^MI02952* mutants that affected *Lime* expression were used in the phenotypic assays.

Outbred strains

We used the outbred populations with and without *FBti0019985* generated in Merenciano et al. (2019) (47). Briefly, they were generated by a round-robin cross-design of inbred lines from the Drosophila Genetic Reference Panel (DGRP) (48) and isofemale lines from different European populations (47). Outbred populations were maintained by random mating with a large population size for over five generations before starting the first experiments.

CRISPR/Cas9 mutant strains

To generate a CRISPR/Cas9 strain with a precise deletion of *FBti0019985* in the natural outbred population we followed a two-step approach. First, we substituted the insertion by the DsRed fluorescence marker and then, in a second step, we removed the visual marker to avoid any possible effect of the introduced DsRed sequence in the results (Figure S1). For the substitution of *FBti0019985* by the DsRed fluorescence marker, guide RNAs (gRNA1 and gRNA2) were designed in the *FBti0019985* flanking region using the flyCRISPR target finder (http://targetfinder.flycrispr.neuro.brown.edu/) and cloned into pCFD5 plasmid following the pCFD5 cloning protocol (www.crisprflydesign.com) (49) using the primers 5'-gcggcccgggttcgattcccggccgatgcaagctagacttatttgagatagtttaga gctagaaatagcaag-3' and 5'-attttaacttgctatttctagctctaaaaccaga gaaacgtcgagctgcgtgcaccagccgggaatcgaaccc-3'. gRNA1 is 5'-tatctcaaataagtctagct-3', while gRNA2 is 5'-cagagaaacgtcgagctgcg-3'. A donor DNA containing two homology arms flanking the DsRed sequence for homology repair were cloned into the pHD-ScarlessDsRed plasmid (https://flycrispr.org/scarless-gene-editing/) using the Q5® High-Fidelity DNA Polymerase kit (New England Biolabs). Left homology arm contained the sequence in 2R:9870299-9871095 (FlyBase Release 6) (50) while the right homology arm contained the sequence in 2R:9871529-9872365 (FlyBase Release 6) (50) from the outbred population with *FBti0019985*. To avoid cleavage of the donor construct and mutagenesis after integration by CRISPR/Cas9, two single-nucleotide synonymous substitutions (C > G for

sgRNA1; G > T for sgRNA2) were introduced into the two sgRNA target site PAM sequences, respectively. The pCFD5 plasmid containing the gRNAs, the donor pHD-ScarlessDsRed plasmid containing the homology arms, and a plasmid containing Cas9 endonuclease were co-injected as a unique mix into approximately 550 embryos from the outbred population with *FBti0019985*. All the injections were performed using the following mix concentrations: pCFD5 plasmid at 100ng/ul, donor plasmid at 500ng/ul, and Cas9 plasmid at 250ng/ul. Offspring was screened for eye fluorescence. Flies with the desired mutation were backcrossed with the parental line for a minimum of five generations to remove potential off-targets generated during the process (51, 52). Then, an homozygous strain containing the deletion of *FBti0019985* was established. The substitution of *FBti0019985* by DsRed was checked by PCR with two primer pairs: 5'-aacaatgcaagtccgtgctc-3', 5'-gtggttcctccacccttgtg-3' and 5'-ggccgcgactctagatcataatc-3' and 5'-gtggttcctccacccttgtg-3'. PCR bands were confirmed by Sanger sequencing. Then, to remove the DsRed sequence, we applied again the CRISPR/Cas9 technique designing two gRNAs (gRNA3 and gRNA4) in the flanking regions of the DsRed sequence as mentioned before and cloning them into the pCFD5 plasmid using the primers 5'-gcggcccgggttcgattcccggccgatgcttgaacactaatgacaatttgttttagagctagaaatagcaag-3' and 5'-attttaacttgctatttctagctctaaaacagctcacaactgcgcagctctgcaccagccgggaatcgaaccc-3'. gRNA3 is 5'-ttgaacactaatgacaattt-3', while gRNA4 is 5'-gagctgcgcagttgtgagct-3'. A donor DNA containing only two homology arms for homology-directed repair were cloned into the pHD-ScarlessDsRed plasmid. Left homology arm contained the sequence in 2R:9870299-9871095 (FlyBase Release 6) (50) while the right homology arm contained the sequence in 2R:9871529-9872365 (FlyBase Release 6) (50) from the outbred population with *FBti0019985* adding two single-nucleotide synonymous substitutions (G > A for sgRNA3 site; C > T for sgRNA4 site). The pCFD5 plasmid containing the gRNAs 3 and 4, the donor pHD-ScarlessDsRed plasmid containing the homology arms, and a plasmid containing Cas9 endonuclease were co-injected as explained before into embryos from the CRISPR/Cas9 mutant strain containing the DsRed fluorescent marker. This time, the offspring was screened for the absence of eye fluorescence. An homozygous strain containing the DsRed deletion was established as mentioned above. The DsRed deletion was checked by PCR using the primer pair 5'-aacaatgcaagtccgtgctc-3' and 5'-cgtaggatcagtgggtgaaaatg-3', and finally PCR bands were confirmed by Sanger sequencing. No

polymorphism were found in the sequenced region except for the two single-nucleotide synonymous substitutions introduced into the two sgRNA target site PAM sequences.

Transgenic strains

For the *in vivo* reporter assays, we used the transgenic flies with the *FBti0019985* sequence cloned in front of the *lacZ* reporter gene generated in Ullastres et al. (2021) (19), and we compared them with transgenic strains with the placZ.attB empty vector to control for possible *lacZ* expression driven by the vector sequence itself. Transgenic flies with deleted TFBSs were generated using the placZ.attB vector containing the *FBti0019985* sequence (19), and performing directed mutagenesis for each TFBS with the Q5 Site-Directed Mutagenesis Kit (New England Biolabs). Primers used for deleting the TFBSs can be found in Table S6. The placZ.attB vectors containing the *FBti0019985* and the desired deletions in TFBSs were microinjected at 350-500 ng/ul into a *D. melanogaster* strain with a stable docking site (Bloomington Stock number: #24749). Offspring was screened for red eyes and the insertion of the construct was verified by PCR and Sanger sequencing with the primer pair 5'-ggtgggcataatagtgttgtttat-3' and 5'-cgacgtgttcactttgcttgt-'3. Three independent homozygous stocks were generated and used as biological replicates for the qRT-PCR experiments.

Expression analysis

Sample collection

***P. entomophila* infection**: 5-7 day-old flies from each strain tested were separated by sex and placed in vials with fresh food in groups of 25-35. We allowed flies to recover from CO_2 anesthesia for 24 hours at 25ºC. To expose the flies to the gram-negative bacteria *P. entomophila* infection, we followed the protocol described in Neyen et al. (2014) (53). Briefly, after two hours of starvation, we transfer 75-105 flies in groups of 25-35 into three vials with fresh food and a filter paper soaked with 120 ul of a solution containing 1.25% sucrose and bacterial preparation adjusted to a final OD600 = 50 for females and OD600 = 150 for males. Flies were kept at 29ºC, the optimal temperature condition for *P. entomophila* infection. Simultaneously, a total of 75-105 flies were also transferred in groups of 25-35 into three vials with fresh food and a filter paper soaked with 120ul of a solution containing sterile LB with 1.25% sucrose and kept at 29ºC

as a control. Guts from males and from females were dissected after 12 hours. Samples were then flash-frozen in liquid nitrogen and stored at -80ºC until sample processing.

Cold-stress treatment: 5-7 day-old flies from each strain tested were allowed to lay eggs at 25ºC in a fly cage with egg-laying medium (2% agar with apple juice and a piece of fresh yeast) for four hours. After these four hours, adults were removed and the plate containing embryos was kept at 1ºC for four additional hours. Simultaneously, another plate with embryos was kept at 25ºC for four additional hours as a control. 4-8 hour-old embryos were then collected from both plates using the method described in Schou et al. (2013) (54) adding an additional step of dechorionation during 10 minutes with 50% bleach. Finally, samples were flash-frozen in liquid nitrogen and stored at -80ºC until sample processing.

Ethanol exposure: 5-7 days-old flies from each strain tested were separated by sex and placed in vials with fresh food in groups of 25. We allowed flies to recover from CO_2 anesthesia for 24 hours at 25ºC. To expose the flies to ethanol we followed the method described in Maples et al. (2011) (55). Briefly, flies were transferred to an empty vial and a coated cotton ball with 0.5 ml ethanol was inserted into the exposure vial for 9 minutes ensuring that the alcohol was facing into the vial and not towards the wall. Simultaneously, coated cotton balls with 0.5 ml H_2O were inserted into the control vials. After that, flies were flash-frozen in liquid nitrogen and stored at -80ºC until sample processing.

RNA extraction and cDNA synthesis

RNA was extracted using the GeneEluteTM Mammalian Total RNA Miniprep Kit following manufacturer's instructions (Sigma). RNA was then treated with DNase I (Thermo). cDNA was synthesized from a total of 250-1,000 ng of RNA using the NZY Fisrt-Strand cDNA synthesis kit (NZYTech).

qRT-PCR analysis

Lime expression was measured using the forward primer 5' -gagcagttggaatcgggtttttac -3' and the reverse primer 5'-gtatgaatcgcagtccagccata-3' spanning 99 bp cDNA in the exon 1/exon 2 junction. *cbx* expression was measured with the forward primer 5'-gggaaaacgatctgggagca -3' and the reverse primer 5'- gtcggagaagttgagtggga -3' spanning 233 bp cDNA in the exon 2/exon 3 junction. *lacZ* reporter gene expression was measured using the forward primer 5'-cctgctgatgaagcagaacaact-3' and the reverse primer 5'-gctacggcctgtatgtggtg-3'. Gene expression

was normalized with *Act5C* (5' -gcgcccttactctttcacca-3' and 5' -atgtcacggacgatttcacg-3' primers).
We performed the qRT-PCR analysis with SYBR Green (BioRad) or with the qPCRBIO
SyGreen Mix Lo-Rox (PCRBiosystems) on iQ5 and CFX384 Thermal cyclers, respectively.
Results were analyzed using the dCT method (56).

Transcript start site detection

We performed RT-PCRs to detect whether *FBti0019985* is adding an alternative TSS to the *Lime*
gene in outbred flies carrying *FBti0019985* after different stress conditions. We used the forward
primer 5'- aaaactcaacgagtaaagtcttc -3' and the reverse primer 5'- tataaagttccaacgcccagc -3' to
detect the *Lime* transcript starting in the TE. The forward primer 5'- cgcagagaaacgtcgagctg -3'
and the reverse primer 5'-cacgttaaattcactagggtggc -3' were used to detect *Lime* total transcript.
Outbred population without *FBti0019985* was used as a control sample.

Phenotypic assays

P. entomophila **infection**

100 5-7 day-old male flies and 100 5-7 day-old females from each one of the strains tested were
infected with the gram-negative bacteria *P. entomophila* as described before. Simultaneously, a
total of 30 male and female flies were tested as controls. We counted the number of dead flies in
every vial at different time points for at least 6 days. Log-rank tests were performed to analyze
survival curves with SPSS v21 software.

Egg-to-adult viability under cold-stress

5-7 day-old flies from each strain tested were allowed to lay eggs for 4 h at 25ºC in a fly cage
with egg-laying medium (2% agar with apple juice and a piece of fresh yeast) for four hours.
Then, adults were removed from the cage and plates were kept for four additional hours at 25ºC.
After that, 4-8 hour-old embryos were collected using the method described in Schou et al.
(2013) (54) and placed in vials with fresh food in groups of 30. In total, 240-300 embryos were
tested for each strain. Cold-stressed vials were kept at 1ºC for 15 h and then maintained at 25ºC
until adult emergence. Simultaneously, control vials were kept at 25ºC and never exposed to
cold. Percentage egg-to-adult viability was calculated based on the number of emerged flies to
the total number of embryos placed in each vial. Statistical significance was calculated
performing ANOVA using SPSS v.21 combining all the data into a full model: experimental

condition (stress and nonstress), insertion genotype (presence/absence of the insertion) and interaction between these two factors.

ACKNOWLEDGEMENTS

We thank Fillip Port from the Division of Signaling and Functional Genomic led by Prof. Dr. Michael Boutros for his advice in the generation of the CRISPR/Cas9 mutants. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (H2020-ERC-2014-CoG-647900).

REFERENCES

1.   Nelson TC, Jones MR, Velotta JP, Dhawanjewar AS, Schweizer RM. UNVEILing connections between genotype, phenotype, and fitness in natural populations. Mol Ecol. 2019;28(8):1866-76.
2.   Otwinowski J, Nemenman I. Genotype to phenotype mapping and the fitness landscape of the E. coli lac promoter. PLoS One. 2013;8(5):e61570.
3.   Mackay TFC, Huang W. Charting the genotype-phenotype map: lessons from the Drosophila melanogaster Genetic Reference Panel. Wiley Interdiscip Rev Dev Biol. 2018;7(1).
4.   Kammenga JE. The background puzzle: how identical mutations in the same gene lead to different disease symptoms. FEBS J. 2017;284(20):3362-73.
5.   Young AI, Benonisdottir S, Przeworski M, Kong A. Deconstructing the sources of genotype-phenotype associations in humans. Science. 2019;365(6460):1396-400.
6.   Reddy KC, Andersen EC, Kruglyak L, Kim DH. A polymorphism in npr-1 is a behavioral determinant of pathogen susceptibility in C. elegans. Science. 2009;323(5912):382-4.
7.   Barrett RD, Hoekstra HE. Molecular spandrels: tests of adaptation at the genetic level. Nat Rev Genet. 2011;12(11):767-80.
8.   Orgogozo V, Morizot B, Martin A. The differential view of genotype-phenotype relationships. Front Genet. 2015;6:179.
9.   Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, et al. Ten things you should know about transposable elements. Genome Biol. 2018;19(1):199.
10.   Schrader L, Schmitz J. The impact of transposable elements in adaptive evolution. Mol Ecol. 2019;28(6):1537-49.
11.   Casacuberta E, González J. The impact of transposable elements in environmental adaptation. Mol Ecol. 2013;22(6):1503-17.
12.   Chuong EB, Elde NC, Feschotte C. Regulatory activities of transposable elements: from conflicts to benefits. Nat Rev Genet. 2017;18(2):71-86.
13.   Chuong EB, Elde NC, Feschotte C. Regulatory evolution of innate immunity through co-option of endogenous retroviruses. Science. 2016;351(6277):1083-7.

14.   Ding Y, Berrocal A, Morita T, Longden KD, Stern DL. Natural courtship song variation caused by an intronic retroelement in an ion channel gene. Nature. 2016;536(7616):329-32.

15.   Esnault C, Lee M, Ham C, Levin HL. Transposable element insertions in fission yeast drive adaptation to environmental stress. Genome Res. 2019;29(1):85-95.

16.   Huang C, Sun H, Xu D, Chen Q, Liang Y, Wang X, et al. ZmCCT9 enhances maize adaptation to higher latitudes. Proc Natl Acad Sci U S A. 2018;115(2):E334-E41.

17.   Li ZW, Hou XH, Chen JF, Xu YC, Wu Q, González J, et al. Transposable Elements Contribute to the Adaptation of Arabidopsis thaliana. Genome Biol Evol. 2018;10(8):2140-50.

18.   Van't Hof AE, Campagne P, Rigden DJ, Yung CJ, Lingley J, Quail MA, et al. The industrial melanism mutation in British peppered moths is a transposable element. Nature. 2016;534(7605):102-5.

19.   Ullastres A, Merenciano M, González J. Regulatory regions in natural transposable element insertions drive interindividual differences in response to immune challenges in Drosophila. Genome Biol. 2021;22(1):265.

20.   Aminetzach YT, Macpherson JM, Petrov DA. Pesticide resistance via transposition-mediated adaptive gene truncation in Drosophila. Science. 2005;309(5735):764-7.

21.   Magwire MM, Bayer F, Webster CL, Cao C, Jiggins FM. Successive increases in the resistance of Drosophila to viral infection through a transposon insertion followed by a Duplication. PLoS Genet. 2011;7(10):e1002337.

22.   Anholt RRH, Mackay TFC. The road less traveled: from genotype to phenotype in flies and humans. Mamm Genome. 2018;29(1-2):5-23.

23.   Telonis-Scott M, Hallas R, McKechnie SW, Wee CW, Hoffmann AA. Selection for cold resistance alters gene transcript levels in Drosophila melanogaster. J Insect Physiol. 2009;55(6):549-55.

24.   Mihajlovic Z, Tanasic D, Bajgar A, Perez-Gomez R, Steffal P, Krejci A. Lime is a new protein linking immunity and metabolism in Drosophila. Dev Biol. 2019;452(2):83-94.

25.   Ayres JS, Freitag N, Schneider DS. Identification of Drosophila mutants altering defense of and endurance to Listeria monocytogenes infection. Genetics. 2008;178(3):1807-15.

26.   Merenciano M, Ullastres A, de Cara MA, Barrón MG, González J. Multiple Independent Retroelement Insertions in the Promoter of a Stress Response Gene Have Variable Molecular and Functional Effects in Drosophila. PLoS Genet. 2016;12(8):e1006249.

27.   Batut P, Dobin A, Plessy C, Carninci P, Gingeras TR. High-fidelity promoter profiling reveals widespread alternative promoter usage and transposon-driven developmental gene expression. Genome Res. 2013;23(1):169-80.

28.   Villanueva-Cañas JL, Horvath V, Aguilera L, González J. Diverse families of transposable elements affect the transcriptional regulation of stress-response genes in Drosophila melanogaster. Nucleic Acids Res. 2019;47(13):6842-57.

29.   Wang X, Huang J, Zhu F. Human Endogenous Retroviral Envelope Protein Syncytin-1 and Inflammatory Abnormalities in Neuropsychological Diseases. Front Psychiatry. 2018;9:422.

30.   Frank JA, Singh M, Cullen HB, Kirou RA, Coyne CB, Feschotte C. Antiviral activity of a human placental protein of retroviral origin. bioRxiv. 2020:2020.08.23.263665.

31.   Senft AD, Macfarlan TS. Transposable elements shape the evolution of mammalian development. Nat Rev Genet. 2021;22(11):691-711.

32.   Mi S, Lee X, Li X, Veldman GM, Finnerty H, Racie L, et al. Syncytin is a captive retroviral envelope protein involved in human placental morphogenesis. Nature. 2000;403(6771):785-9.

33. Gröger V, Cynis H. Human Endogenous Retroviruses and Their Putative Role in the Development of Autoimmune Disorders Such as Multiple Sclerosis. Front Microbiol. 2018;9:265.

34. Baduel P, Quadrana L. Jumpstarting evolution: How transposition can facilitate adaptation to rapid environmental changes. Curr Opin Plant Biol. 2021;61:102043.

35. Andersson R, Sandelin A. Determinants of enhancer and promoter activities of regulatory elements. Nat Rev Genet. 2020;21(2):71-87.

36. Dao LTM, Spicuglia S. Transcriptional regulation by promoters with enhancer function. Transcription. 2018;9(5):307-14.

37. Arnold CD, Gerlach D, Spies D, Matts JA, Sytnikova YA, Pagani M, et al. Quantitative genome-wide enhancer activity maps for five Drosophila species show functional enhancer conservation and turnover during cis-regulatory evolution. Nat Genet. 2014;46(7):685-92.

38. Zabidi MA, Arnold CD, Schernhuber K, Pagani M, Rath M, Frank O, et al. Enhancer-core-promoter specificity separates developmental and housekeeping gene regulation. Nature. 2015;518(7540):556-9.

39. Barco B, Kim Y, Clay NK. Expansion of a core regulon by transposable elements promotes Arabidopsis chemical diversity and pathogen defense. Nat Commun. 2019;10(1):3444.

40. Sundaram V, Wysocka J. Transposable elements as a potent source of diverse. Philos Trans R Soc Lond B Biol Sci. 2020;375(1795):20190347.

41. Faulkner GJ, Carninci P. Altruistic functions for selfish DNA. Cell Cycle. 2009;8(18):2895-900.

42. Guio L, Vieira C, González J. Stress affects the epigenetic marks added by natural transposable element insertions in Drosophila melanogaster. Sci Rep. 2018;8(1):12197.

43. Rey O, Danchin E, Mirouze M, Loot C, Blanchet S. Adaptation to Global Change: A Transposable Element-Epigenetics Perspective. Trends Ecol Evol. 2016;31(7):514-26.

44. Koštál V, Korbelová J, Rozsypal J, Zahradníčková H, Cimlová J, Tomčala A, et al. Long-term cold acclimation extends survival time at 0°C and modifies the metabolomic profiles of the larvae of the fruit fly Drosophila melanogaster. PLoS One. 2011;6(9):e25025.

45. Overgaard J, Malmendal A, Sørensen JG, Bundy JG, Loeschcke V, Nielsen NC, et al. Metabolomic profiling of rapid cold hardening and cold shock in Drosophila melanogaster. J Insect Physiol. 2007;53(12):1218-32.

46. Bou Sleiman MS, Osman D, Massouras A, Hoffmann AA, Lemaitre B, Deplancke B. Genetic, molecular and physiological basis of variation in Drosophila gut immunocompetence. Nat Commun. 2015;6:7829.

47. Merenciano M, Iacometti C, González J. A unique cluster of *roo* insertions in the promoter region of a stress response gene in *Drosophila Melanogaster*. Mob DNA. 2019;10:10.

48. Mackay TF, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, et al. The Drosophila melanogaster Genetic Reference Panel. Nature. 2012;482(7384):173-8.

49. Port F, Bullock SL. Augmenting CRISPR applications in Drosophila with tRNA-flanked sgRNAs. Nat Methods. 2016;13(10):852-4.

50. Larkin A, Marygold SJ, Antonazzo G, Attrill H, Dos Santos G, Garapati PV, et al. FlyBase: updates to the Drosophila melanogaster knowledge base. Nucleic Acids Res. 2021;49(D1):D899-D907.

51. Bassett A, Liu JL. CRISPR/Cas9 mediated genome engineering in Drosophila. Methods. 2014;69(2):128-36.

52. Port F, Strein C, Stricker M, Rauscher B, Heigwer F, Zhou J, et al. A large-scale resource for tissue-specific CRISPR mutagenesis in. Elife. 2020;9.

53. Neyen C, Bretscher AJ, Binggeli O, Lemaitre B. Methods to study Drosophila immunity. Methods. 2014;68(1):116-28.

54. Schou MF. Fast egg collection method greatly improves randomness of egg sampling in Drosophila melanogaster. Fly (Austin). 2013;7(1):44-6.

55. Maples T, Rothenfluh A. A simple way to measure ethanol sensitivity in flies. J Vis Exp. 2011(48).

56. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res. 2001;29(9):e45.

Figure S1. Schematic representation of the CRISPR/Cas9 deletion of *FBti0019985*. A) Schematic representation of the substitution of *FBti0019985* by the *DsRed* visual marker. B) Schematic representation of the deletion of the *DsRed* visual marker.