Assessing Quantitative MRI Techniques using Multimodal Comparisons

Francis Carter[1,2], Alfred Anwander[3], Thomás Goucha[3], Helyne Adamson[3], Angela D. Friederici[3], Antoine Lutti[4], Claudine J Gauthier[5,6,7], Nikolaus Weiskopf[8,9], Pierre-Louis Bazin[10,11], Christopher J Steele[1,6,11]

[1]Department of Psychology, Concordia University, Montreal, QC, Canada
[2]Montreal Institute for Learning Algorithms, Université de Montréal, Montreal, QC, Canada
[3]Department of Neuropsychology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany
[4]Department of Clinical Neurosciences, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland
[5]Department of Physics, Concordia University, Montreal, QC, Canada
[6]Perform Center, Concordia University, Montreal, QC, Canada
[7]Montreal Heart Institute, Montreal, QC, Canada.
[8]Department of Neurophysics, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany
[9]Felix Bloch Institute for Solid State Physics, Faculty of Physics and Earth Sciences, Leipzig University, Leipzig, Germany
[10]Faculty of Social and Behavioral Sciences, University of Amsterdam, Amsterdam, Netherlands
[11]Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany
[1]Department of Psychology, Concordia University, Montreal, QC, Canada
[6]Perform Center, Concordia University, Montreal, QC, Canada
[11]Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

Abstract

 The study of brain change in neuroscience studies is commonly conducted using macroscopic morphological measures of the brain such as regional volume or cortical thickness, providing little insight into the microscopic mechanisms underlying brain disease. In contrast, quantitative MRI allows the monitoring of microscopic brain change non-invasively in-vivo, and provides normative values for comparisons between tissues, regions, and individuals. To support the development and common use of qMRI for cognitive neuroscience, we analysed a set of qMRI metrics (R1, R2*, Magnetization Transfer saturation, Proton Density saturation, Fractional Anisotropy, Mean Diffusivity) in 101 healthy young adults. Here we provide a comprehensive descriptive analysis of these metrics and their linear relationships to each other in grey and white matter to develop a more complete understanding of the relationship to tissue microstructure. Furthermore, we provide evidence that combinations of metrics may uncover informative microstructural cortical gradients across the brain by showing that dimensionality reduction through PCA provides cortical gradients that were hidden in individual metrics. We discuss these results and their links to microstructural and physiological neuroscience research.

Introduction

Magnetic Resonance Imaging (MRI) gives rise to a variety of image contrasts primarily driven by differences in longitudinal and transverse relaxation rates of the MRI signal between voxels. The estimates of these relaxation rates, provided by quantitative MRI (qMRI) (Guilfoyle et al., 2003; Weiskopf et al., 2021), have been shown to correlate with microstructural properties of brain tissues such as iron or myelin concentration (Fukunaga et al., 2010; Sereno et al., 2013) and show great potential for the monitoring of microscopic brain change in-vivo (Tardif et al., 2016; Weiskopf et al., 2015). However, the specificity of these biomarkers is limited by the combined contributions of multiple histological properties to qMRI estimates (Draganski et al., 2011; Tardif et al., 2017, 2016; Weiskopf et al., 2021, 2015). Improving the specificity of these biomarkers requires a detailed understanding of the impact of brain microscopic properties on qMRI estimates. Thus, the goal of this paper is to further our knowledge of the information contained in a set of quantitative MRI metrics to assist in the development of "in-vivo histology", the mapping of MRI signals to the microstructural properties of tissues (Weiskopf et al., 2021, 2015).

The assessment of microscopic brain tissue properties from in-vivo qMRI data requires a detailed understanding of how these properties impact the MRI signal (Weiskopf et al., 2021, 2015). Recent research includes evidence that the MR relaxation rates R1 and R2* are driven primarily by myelin and iron concentrations, respectively (Kirilina et al., 2020; Möller et al., 2019; Shams et al., 2019; Stüber et al., 2014; Trampel et al., 2019; Weiskopf et al., 2015) and that magnetization transfer (MT) measurements scale with myelin's contribution to the macromolecular content (Mangeat et al., 2015; Manning et al., 2017). Myelin concentration has been shown to be the dominant contributor to multiple qMRI measures (e.g. Callaghan et al., MRM 2015; Mancini et al., 2020), highlighting the need for complementary qMRI measures in order to achieve a complete description of brain microstructure.

One major issue is that a single voxel's value is potentially the result of many different combinations of molecules and their concentrations (i.e., there is a one-to-many mapping of specific magnetic properties to many different molecular arrangements) (Draganski et al., 2011; Tardif et al., 2017, 2016). Multimodal MRI data (i.e., including multiple MRI contrasts) may help address this issue through the extraction of latent variables from a combination of qMRI measures. If such latent variables can be shown to be informative, this suggests that they may be related to microstructural features and that "ground truth" molecular concentrations from histology may correspond to specific latent variables within the multimodal qMRI data. Extraction of such latent spaces could be a path forward for the development of in-vivo histology (Borsboom et al., 2003; Geeraert et al., 2019). Estimation of the latent variables needs to be conducted over structures that can be considered as homogeneous - which requires accurate segmentation and the reduction of tissue partial voluming - to simplify the task of identifying relevant latent variables. This segmentation step has a large but often overlooked effect on the interpretation of results (Lüsebrink et al., 2013), and is often difficult to properly implement, particularly for cortical grey matter tissues where partial voluming with CSF and WM is common at standard resolutions (Dahnke et al., 2013; Han et al., 2004; Roche and Forbes, 2014).qMRI data offers higher reproducibility and comparability between acquisitions than structural MRI data commonly used in neuroscience for the study of macroscopic morphological change (Caan et al., 2019; Leutritz et al., 2020; Weiskopf et al., 2015). The advent of more advanced qMRI sequences that simultaneously capture multiple quantitative metrics has led to shorter overall acquisition times (European Society of Radiology (ESR), 2015; Weiskopf et al., 2013) and removes the need for within-subject registration between metrics. Notably, this work has led to the development of Multi-Parametric Mapping, which simultaneously captures quantitative R1,

R2*, MT, and PD images (MPM; Weiskopf et al., 2013) and multi-echo MP2RAGE (T1, T2*, QSM; Caan et al., 2019; Metere et al., 2017).

While useful for acquiring multiple co-registered modalities, multiparametric methods likely contain redundancies in the information provided by each metric (Callaghan et al., 2015b; Weiskopf et al., 2021, 2015). Redundancies have the potential to be exploited to determine which combination of metrics accounts for the most variance in the tissue of interest (and could therefore form a basis for a minimum useful multiparametric acquisition profile) and, importantly, can then be used to map putative microstructural similarities across the brain (Weiskopf et al., 2015). Similar to the way that microstructure exhibits differences across different tissues and regions, the relationships between metrics are also expected to vary. Previous work has predominantly explored metric-metric covariance relationships with ROI-based or segmented network approaches (Seidlitz et al., 2018; Uddin et al., 2019), which rely on *a priori* regional delineations and averaging that may obscure subtle variability within regions. A voxel-wise approach could provide a more nuanced and comprehensive description of these relationships and help to determine if and/or how multiparametric combinations can provide additional information not found in individual metrics (Paquola et al., 2019).

The present study provides a statistical description of a set of six metrics, including four qMRI metrics from the MPM (Weiskopf et al., 2013) (R1, R2*, Magnetization Transfer saturation-MT, and Proton Density saturation - PD) and two diffusion MRI metrics (dMRI: Fractional Anisotropy - FA, and Mean Diffusivity - MD). We set out to identify potential redundancies between metrics, and additionally identify the set of normative relationships between metrics across different tissue types (sub/cortical grey matter, white matter) that can be used to develop and test hypotheses in the development of in-vivo histology. We also provide evidence that

multimodal imaging holds promise in describing the microstructural drivers of qMRI contrast by applying PCA to extract and map latent variables within the data (Geeraert et al., 2019).

Methods

2.1. Participants

Our sample consisted of 101 young healthy individuals (mean age 25.7 ± 4, range 18-34 years, 25 females) who were recruited in the area of Leipzig, Germany. All participants were right-handed, had a high-school-level education, no history of neurological or psychiatric disorders, and did not use any centrally effective medication. The study was approved by the ethics committee at the medical faculty of the University of Leipzig. All participants gave written informed consent and were paid for participation in the study.

2.2. MRI data acquisition and quantitative metric estimation

Quantitative multi-parametric maps (MPM, 1mm iso) and high spatial and angular resolution diffusion-weighted MRI (dMRI, 1.3mm iso) were acquired for all participants on a 3-Tesla PRISMAfit MRI system (Siemens Healthineers, Erlangen, Germany) using a standard 32-channel receive head coil. The MPM protocol consisted of three multi-echo 3D FLASH (fast low angle shot) acquisitions with predominant T1-, PD-, and MT-weighting by adapting the repetition time (TR) and the flip angle (T1w: 18.7 ms/20°, PDw and MTw: 23.7 ms/6°) following the setting previously published (Weiskopf et al., 2013). The MTw acquisition included an additional initial 4 ms long off-resonance MT saturation pulse (flip angle 220°, 2 kHz offset) (Helms et al., 2008). We acquired six gradient echoes (equidistant echo times, TE, from 2.2 to 14.7 ms) with alternating polarity for the T1w and the MTw volumes and eight datasets for the PDw volumes (TE from 2.2 to 19.7ms). The remaining imaging parameters were: field-of-view

256 × 240 mm, 176 slices, acceleration in phase-direction using GRAPPA 2 and in partition direction using 6/8 partial Fourier, non-selective RF excitation, high readout bandwidth = 425 Hz/pixel, RF spoiling phase increment = 50°, acquisition time ~21 min. The acquisition was preceded by additional calibration data to correct for spatial variations in the radio-frequency transmit field (B1+) as described by Lutti and colleagues (Lutti et al., 2012). The diffusion MRI protocol used the multi-band sequence developed at CMRR (https://www.cmrr.umn.edu/multiband) with the following parameters: 1.3 mm isotropic resolution, b-value = 1000 s/mm², 60 directions and seven images without diffusion weighting (b = 0), three repetitions to improve the SNR, TE = 75 ms, TR = 6 s, GRAPPA 2, CMRR-SMS 2, two b = 0 acquisitions with opposite phase encoding directions. The acquisition time for the dMRI protocol was 23 min.

The MPMs acquisitions were processed with the freely available hMRI toolbox (http://hmri.info, Tabelow et al., 2019) to estimate parameter maps of the longitudinal relaxation rate (R1), the effective transverse relaxation rate (R2*), the magnetization transfer (MT), the proton density (PD) for each participant. The dMRI images were corrected for motion, eddy currents, and susceptibility distortions (Andersson and Sotiropoulos, 2016), and the fractional anisotropy (FA), and mean diffusivity (MD) were computed from the diffusion tensor images using the FSL tools (Jenkinson et al., 2012). Co-registration between native diffusion and MPM space was accomplished by rigidly co-registering the b0 and R1 images within each individual, and applying the transform to FA and MD maps to bring them into R1 space (trilinear interpolation).

2.3. Preprocessing

After data collection and quantitative metric reconstruction, the following steps were performed to segment the brains into different tissue types (cerebral cortical GM, subcortical GM, WM),

and to create surface representations of the cortex. We obtained high-quality segmentations and cortical representations by employing previously established methods (Bazin et al., 2014; Huntenburg et al., 2017; Tardif et al., 2015). Other than the SPM skull stripping and segmentations described below, all other preprocessing was performed with the open-source neuroimaging toolbox Nighres v1.2 (Huntenburg et al., 2018).

2.3.1 Segmentation

Cortical, subcortical, and WM segmentations for each participant were performed with standard GM segmentation in SPM12 on the R1 image (Ashburner and Friston, 2005) and the Nighres implementation of Multiple object Geometric Deformable Model (MGDM) segmentation using FA, R1, MT, and PD (Bazin et al., 2014; Bogovic et al., 2013). MGDM allowed for the segmentation of the cerebral cortical and subcortical GM while preserving topology, and SPM segmentations were then combined as follows to ensure final segmentations that reduced tissue partial voluming. WM masks were defined as the overlap between SPM WM probability maps thresholded at 0.5 and the MGDM cerebral WM. GM masks were defined as the overlap between the SPM GM (probability maps thresholded at 0.95) and the MGDM labels for cerebral GM, including cortical and subcortical labels. In addition, the subcortical mask generated by MGDM was used to extract subcortical GM. The MGDM subcortical labels that corresponded with the SPM GM thresholded at 0.95 defined the subcortical mask. The resulting masks retained artifactual R2* signal (mainly in the medial temporal lobe) which we removed by excluding voxels containing R2* values above 25 in GM. We also used an MD threshold of 0.001 to reduce any partial voluming with CSF in the GM masks. Subsequent analyses were performed on voxel-wise values extracted from the cortical sheet, WM, and subcortical GM in

both hemispheres. A group surface co-registration step for both hemispheres was additionally used to project cortical results onto a common group surface for visualization as detailed below.

2.3.2 Cortex extraction, mesh generation, and surface projection

A 3D mesh representation of the cortex was used to map and visualize our results on the cortical surface. To mitigate against the effects of partial voluming on the pial and WM surfaces, we extracted a thin sheath of the middle layer of the cortex with Nighres as follows. The cortex was first extracted with CRUISE Cortex Extraction (Han et al., 2004), generating GM-WM and GM-CSF boundary levelset images (where a levelset is the distance of each voxel from each boundary; thus representing the inner and outer cortical surfaces in voxel space). These two boundaries were then combined to extract a levelset representation of the midline of the cortical sheet. The midline levelset, used later for group co-registration, was then thresholded at a distance of 0.5 mm to generate a binary mask, and converted to a 3D mesh for display. Details on segmentation and the levelset approach for the surface generation with Nighres can be found in our previous technical publications (Huntenburg et al., 2018), and all specific commands and parameters are included as code in our github repository (https://github.com/neuralabc/paper_QuantitativeMetricComparisons).

2.4 Group co-registration

Cortical midline levelsets were used to perform surface-based registration in voxel space with a method similar to that developed by Tardif and colleagues (Tardif et al., 2015). The ANTs (Avants et al., 2011; Tustison et al., 2014) multivariate template construction script was used to perform nonlinear registrations and generate a group template for both hemispheres. The levelsets representing each participant's middle cortical ribbon were thresholded at 10 mm from

the cortex and used as inputs to ANTs Multivariate Template Construction (v2, 3 iterative steps, with default linear and nonlinear registration parameters, Demons similarity metric). The final template levelset was then converted to a mesh representation for display (Huntenburg et al., 2018). To map the results of the analysis onto a mesh representation of the group template surface, participants' data from each metric was: 1) multiplied by a binarized midline ribbon, 2) nonlinearly transformed into the study-specific group space with the computed deformation (ANTs SyN, nearest-neighbor interpolation), and 3) a 2 mm intensity propagation step was performed to ensure that all participant's data overlapped with the final coregistered surface. In all cases, the vertex-wise median values across participants were presented for display.

2.5. Statistical analysis

2.5.1 Metric-metric density plots

Bivariate logarithmic density plots were created to describe the relationship between metrics in GM, WM, cortex, and subcortical GM. Linear regressions were used to calculate the coefficients of determination (R-squared) for each bivariate relationship. Univariate density plots (histograms) were also generated to further describe the data.

2.5.2 Dimensionality reduction

We performed model-free dimensionality reduction with Principal Component Analysis (PCA) in cortical tissue to visualize latent variables within the data. All six metrics in each of these tissues from all participants were z-scored and used as input to a PCA in scikit learn (Pedregosa et al., 2011) in Python (3.7), and all six PCs were assessed. Each participant's data was then transformed into the PCs' subspace for visualization, and transformed into the common group

space. This allowed us to compute the median across participants and visualize our results on the cortical surface template.

3. Results

3.1 Cortical segmentation

Our segmentations combined MGDM and SPM segmentations to reduce biases caused by partial voluming of GM with WM or CSF. Visual inspection showed that our results had limited partial voluming of the cortex with CSF and WM, with little to no gyral or sulcal bias (Dahnke et al., 2013; Figure 1). As an additional confirmatory step before using this segmentation in our analyses, we also checked to ensure that our cortical maps showed high correspondence with those presented in previous literature. As expected, R1 cortical maps exhibited a gradient with peaks in primary sensory and motor regions (Figure 1), a pattern that is in excellent correspondence with what has been shown in previous literature (Marques et al., 2017; Shams et al., 2019).

(a)

Figure 1: Overview of segmentation results. (a) The six metrics used were captured in 101 participants. (b) A sample brain with a red mask of the middle cortical ribbon as a semi-transparent overlay on R1, showing little to no partial voluming. (c) Using data from the cortical ribbon, we plotted the median R1 values across all participants to the group template. Other metric plots can be found in the supplemental data.

3.2 Descriptive statistics: GM and WM

All pairs of metrics, both in GM (combined cortical and subcortical) and WM, were found to be significantly correlated (Figure 2, all p<0.001). However, there were large fluctuations in these pairwise correlations, with R^2 values ranging from as low as 0.0001 in the MD-R2* GM correlation, to as high as 0.482 in the R1-MT GM correlation. These differences were not equally dispersed across metrics; R1 had the highest correlations with other metrics across all structures (R^2 average = 0.258), while MD had the weakest correlations (R^2 average = 0.051). Furthermore, the strengths of correlations varied across tissues, with some R^2 values differing by over 10% of explained variance. Notably, FA-MD, MD-MT, and R1-MD all had R^2 values that were more than 10% higher in GM than in WM, and almost all pairwise correlations were greater in GM than WM. However, FA-PD and FA-R2* had R^2 values slightly higher in WM than in GM. These relationships were reliable and consistent across participants, as shown by the low standard deviation of R^2 across participants in Supplementary Figure 3.

Figure 2: Pairwise correlations in GM (above the diagonal) and WM (below the diagonal). Data along the diagonal shows the histograms of particular metrics in GM (red) and WM (blue). GM includes values extracted from the midline cortical ribbon and subcortical nuclei.

3.3 Descriptive statistics: Cortical and subcortical GM

All pairs of metrics were found to be significantly correlated in cortical and subcortical GM (Figure 3, all p<0.001). However, there were large fluctuations in these pairwise correlations, with R^2 values ranging from as low as <0.005 in the PD-R2* cortical correlation, to as high as 0.447 in the R1-PD subcortical correlation. As with the GM and WM comparisons (Figure 2), differences were not equally dispersed across metrics; R1 had the highest correlations in all GM structures (R^2 average = 0.230), while MD had the weakest correlations (R^2 average = 0.071).

Furthermore, the strengths of correlations varied across tissues, with some R^2 values differing by over 10% of explained variance. Notably, FA-MD and FA-R1 correlations were 23.0 and 9.1% stronger in cortical GM than in subcortical GM, respectively. Conversely, the R1-PD and R1-R2* relationships were 11.2 and 9.1 % stronger in subcortical GM than cortical GM. These relationships were reliable and consistent across participants, as shown by the low standard deviation of R^2 across participants in Supplementary Figure 4.

Figure 3: Pairwise correlations in cortical GM (above the diagonal) and subcortical GM (below the diagonal). Data along the diagonal shows the histograms of particular metrics in cortical (red) and subcortical GM (blue).

3.4 Multimodal data visualization through dimensionality reduction

The results of the PCA revealed various patterns of cortical gradients. All components, when mapped back onto the cortex, provided informative gradients, often denoting primary sensory and motor areas (Figure 4). Interestingly, the components that explained the most variance in the metrics seemed to diverge the least from the gradients of the metrics themselves. For example, PC1, which explains the most variance, had a high weighting of R1 as well as metrics that had high correlations with R1. PC1's cortical gradient showed high correspondence with primary sensory and motor regions, similar to the R1 cortical map (Figure 1). Similarly, PC2 resembles an inverse mapping of R2* and FA gradients, while PC3 resembles PD gradients (see supplementary Figure 2). The most novel gradient was in the component that explained the least amount of variance in contrasts (Figure 4). This gradient least resembled individual quantitative metrics and showed a structural pattern having a degree of resemblance with the Default Mode Network, with high values in the posterior cingulate cortex, medial PFC, and posterior parietal cortex.

Figure 4: Overview of main results of the PCA. (a) radar plots showing the weighting of each component (ranging from -1 to 1) on each PC. In brackets is the percent variance accounted for

by individual PCs. (b) A plot of variance accounted for in each PC. (c) PC1 mapped back onto the brain, showing primary sensory and motor regions. (d) PC6 mapped back onto the brain. Although this only accounts for 3.1% of the variance in the data, this component shows an interesting cortical gradient.

Discussion

The goal of this study was to assess a set of quantitative metrics to provide a statistical description of their relationships. We used a set of previously established methods for extracting and co-registering the cortical ribbon (Bazin et al., 2014; Tardif et al., 2015) to identify core GM voxels that were less likely to suffer from partial volume effects. Bivariate correlations between metrics were performed for each tissue type, and our results showed that quantitative metrics have wide variations in their degree of correlation with each other, but are consistent across participants. Furthermore, we found that applying PCA resulted in the extraction of latent space components not obvious in individual unimodal metrics.

Firstly, we observed the strongest overall pairwise correlation between R1 and MT, and the strength of this correlation was highly stable across tissue types (R^2 = ~36-48%). This may be due to the strong link between R1 and myelin (Möller et al., 2019; Trampel et al., 2019; Weiskopf et al., 2015) and MT as a measure of macromolecular tissue content (Filo et al., 2019; Henkelman et al., 2001; Yarnykh et al., 2018); both of which are expected to be highly related throughout the brain (Callaghan et al., 2015a; Khodanovich et al., 2017). The normative R1-MT relationships that we have identified here could potentially be affected by pathological conditions that affect macromolecular myelin proportions, such as Alzheimer's tau pathologies and white matter hyperintensities, and baseline relationships would then be useful for research in these domains (Hentschel et al., 2004; van Es et al., 2006; Wardlaw et al., 2016).

Conversely, R2* had, on average, low correlations with other metrics (R^2 average ~= 7%), indicating little redundancy. Its strongest correlation was with R1 (GM R^2 ~= 21%, WM R^2 ~= 14%) which is consistent with previous research indicating a link between R2* and myeloarchitecture (Mangeat et al., 2015; Stüber et al., 2014), and an indirect link to myeloarchitecture through its relation with tissue iron concentration (Kirilina et al., 2020; Möller et al., 2019; Trampel et al., 2019; Weiskopf et al., 2015). Interestingly, the R1-R2* correlation was weaker in WM where we might expect it to be highest based on their shared link with myeloarchitecture. One potential reason for this is that R2* is known to be affected by white matter fiber orientation relative to the main magnetic field (B0) (de Pasquale et al., 2013; Kor et al., 2019). This orientation-dependent effect could lead to variation in R2* that does not arise from myeloarchitecture or iron and therefore decreases the strength of the correlation within the white matter. This effect is not expected to be significant within grey matter since there is no predominant net orientation of fibers at voxel resolutions. Overall, the differences in R1 and R2* correlations provide further support for the hypothesis that R1 is more strongly related to other aspects of microstructure, such as the cyto- and myeloarchitecture (Eickhoff et al., 2005) than R2*. Interestingly, the greater R1-R2* correlation in GM was predominantly driven by subcortical GM. Subcortical nuclei are relatively rich in iron (Drayer et al., 1986) and we collapsed across all identified subcortical GM structures (caudate, putamen, thalamus, and globus pallidus), which may lead to the greater variability seen in the subcortex than cortex for both metrics (as indicated by the spread in histograms along the diagonal of Figure 3). This greater variability may in turn drive the greater R1-R2* correlation by making it easier to identify a statistical correlation. This leads us to expect that changes in iron distribution throughout the brain and particularly in the basal ganglia, as is often seen in neurodegeneration and related pathologies (Carver, 2019; Daugherty and Raz, 2013; Khattar et al., 2021), may exhibit different R1-R2*

correlations than what we have observed. Future work could seek to link the relationship

between R1-R2* with other biomarkers of myelination and iron in patients diagnosed with

pathologies related to iron accumulation for use as a non-invasive marker.

dMRI metrics (FA, MD) showed higher correlations with other metrics in GM than WM,

as well as higher correlations in cortical GM than subcortical GM. This suggests that dMRI

metrics may be dependent on tissue parameters that vary more independently in WM such as

fiber orientation and directional uniformity, as would be expected from the diffusion tensor

(Pierpaoli and Basser, 1996). This then suggests that dMRI metrics in GM are more strongly

driven by non-axonal microstructural differences than in white matter. For example, it is possible

that differences in axonal radius and volume, as well as cell types and shapes could be a source

of this change in dMRI correlations between the cortical and subcortical GM (Szafer et al., 1995,

1995). While FA differences/changes have previously been inferred to be due to differences in

myelination, fiber orientation, and/or density (Beaulieu, 2014; Concha et al., 2010; Zatorre et al.,

2012), we found that there were low correlations between FA and R1 (GM R^2 ~ 13%; WM R^2 ~

8%; cortical R^2 ~ 12%; subcortical R^2 ~ 3%), which is arguably a more direct indicator of myelin

(Trampel et al., 2019). Past research has found that R1 and diffusion have differential variation

throughout the lifespan, further supporting a difference in the information captured by these

metrics, and suggesting that varying R1-dMRI correlations may be informative about health and

ageing related issues (Yeatman et al., 2014). The relatively low correlation in WM is also likely

to be partially driven by the mismatch between the tensor-based simplification of WM

architecture as FA, which best represents a single fibre population, and the reality that most

regions include multiple fibre populations. It may be necessary to combine more fibre-specific

parameters from multishell DWI (Raffelt et al., 2017) with fibre-specific T1/R1 estimates from

inversion-recovery DWI (De Santis et al., 2016; Leppert et al., 2021) to more accurately specify

this relationship. Our findings are in line with recent work characterizing the relationships between MR metrics in WM and support the idea that differences/changes in FA may not generally be attributable to myelin (Geeraert et al., 2019; Jones et al., 2013).

The correlation between R1 and PD had a stronger correlation in GM than WM, and this effect was most prominent in subcortical GM, similarly to the R1-R2* relationship. We do not have a clear interpretation for this R1-PD relationship. However, previous literature suggests that PD and T1/R1 are related through their dependence on cell densities - which affects free water content (Tardif et al., 2016). Furthermore, the echo averaging prior to the map calculation introduces an R2* bias on the PD estimates (Weiskopf et al., 2013), which may be a partial driver of the stronger correlation of R1 and PD in subcortical GM.

We were also interested in exploring how dimensionality reduction could be used to determine whether combining metrics enhances details that are difficult or impossible to detect with single metrics (Geeraert et al., 2019). Consistent with the pattern that we identified in the bivariate statistics, R1's high covariance with multiple metrics appeared in the PCA as a cluster of metrics defining PC1 (including high loadings on all metrics except R2* and accounting for 44% of the variance). This component could be a potential candidate for a more specific index of cortical myelination (Lutti et al., 2014; Sereno et al., 2013). PC1, along with most other PCs, resembled gradients found in individual metrics when mapped onto the cortex, such as that of primary sensory and motor cortices seen in R1. However, the last PC provided a gradient containing some regions similar to those in the default-mode network (DMN) commonly identified in functional resting-state MRI (Raichle, 2015; Yeo et al., 2011). Previous research has shown links between the DMN and structural connectivity exhibited by diffusion tensor imaging and tractography (Horn et al., 2014), and between functional connectivity and cortical qT1 as a proxy for intracortical myelin (Huntenburg et al., 2017). Following from this work, it would be

interesting to explore if the gradients exhibited by the PCs are more directly related to functional connectivity and the DMN than any single modality. Whether or not our last PC corresponds to the DMN, we believe that it is interesting because it was the most dissimilar PC to individual metric gradients, suggesting that such multimodal tools can uncover subtle gradients that are overshadowed by other sources of variation in the data (Geeraert et al., 2019).

Future directions

One major limitation of the present study is that it provides only a description of the relationship between qMRI metrics, but cannot comment on their fundamental relationship with physiological tissue parameters. Properly addressing this question will require more studies combining qMRI sequences with ground truth histological and biomarker data (Weiskopf et al., 2021, 2015), ideally in the same brains. While our study used linear dimensionality reduction to allow for clear interpretation, it is likely that proper mapping from MRI to histological variables will require non-linear statistical methods as well. An approach similar to the recently proposed Morphometric Similarity Networks (Seidlitz et al., 2018) could be applied to quantitative multiparametric imaging data to better specify structural covariance (Mechelli, 2005; Zielinski et al., 2010)) and eventually link it to the underlying physiological parameters. In addition, future work with ultra-high resolution MRI could be used to investigate multiparametric differences in contrasts across cortical layers (Eickhoff et al., 2005; Marques et al., 2017; Tardif et al., 2015; Waehnert et al., 2016).

The current study describes the bivariate relationships between qMRI metrics and an initial exploration of how the data is grouped in multidimensional space. We found that there was a wide range of covariance between metrics that differed across tissue type, but that these relationships were stable across individuals. Our findings are a step towards using multimodal

metric combinations to better identify tissue characteristics and develop a more direct link

between non-invasive MRI and physiology.

Supplementary Figures

Supplementary Figure 1: Surface plots for the six metrics. Median values from the cortical ribbon across all participants are plotted on the cortical surface. The range of values and units are described as well, with pu standing for percentage units.

Supplementary Figure 2: Surface plots and metric loadings for all PC's. Median values across participants for each principal component was plotted on the cortical surface. The normalized loading of the metrics for each PC is also shown.

Supplementary Figure 3: Distribution of R^2 values across participants for all pairwise correlations (GM in blue, WM in red). This shows that the relationships between metrics were highly consistent across individuals.

Supplementary Figure 4: Distribution of R^2 values across participants for all pairwise correlations (cortical GM in green, subcortical GM in indigo). This shows that the relationships between metrics were highly consistent across individuals.

References

Andersson, J.L.R., Sotiropoulos, S.N., 2016. An integrated approach to correction for off-resonance effects and subject movement in diffusion MR imaging. NeuroImage 125, 1063–1078. https://doi.org/10.1016/j.neuroimage.2015.10.019

Ashburner, J., Friston, K.J., 2005. Unified segmentation. NeuroImage 26, 839–851. https://doi.org/10.1016/j.neuroimage.2005.02.018

Avants, B.B., Tustison, N.J., Song, G., Cook, P.A., Klein, A., Gee, J.C., 2011. A reproducible evaluation of ANTs similarity metric performance in brain image registration. NeuroImage 54, 2033–2044. https://doi.org/10.1016/j.neuroimage.2010.09.025

Bazin, P.-L., Weiss, M., Dinse, J., Schäfer, A., Trampel, R., Turner, R., 2014. A computational framework for ultra-high resolution cortical segmentation at 7Tesla. NeuroImage 93, 201–209. https://doi.org/10.1016/j.neuroimage.2013.03.077

Beaulieu, C., 2014. The Biological Basis of Diffusion Anisotropy, in: Diffusion MRI. Elsevier, pp. 155–183. https://doi.org/10.1016/B978-0-12-396460-1.00008-1

Bogovic, J.A., Prince, J.L., Bazin, P.-L., 2013. A multiple object geometric deformable model for image segmentation. Comput. Vis. Image Underst. 117, 145–157. https://doi.org/10.1016/j.cviu.2012.10.006

Borsboom, D., Mellenbergh, G.J., van Heerden, J., 2003. The theoretical status of latent variables. Psychol. Rev. 110, 203–219. https://doi.org/10.1037/0033-295X.110.2.203

Caan, M.W.A., Bazin, P., Marques, J.P., Hollander, G., Dumoulin, S.O., Zwaag, W., 2019. MP2RAGEME: T_1, T_2^*, and QSM mapping in one sequence at 7 tesla. Hum. Brain Mapp. 40, 1786–1798. https://doi.org/10.1002/hbm.24490

Callaghan, M.F., Helms, G., Lutti, A., Mohammadi, S., Weiskopf, N., 2015a. A general linear relaxometry model of R_1 using imaging data: General Linear Relaxometry Model of R1. Magn. Reson. Med. 73, 1309–1314. https://doi.org/10.1002/mrm.25210

Callaghan, M.F., Josephs, O., Herbst, M., Zaitsev, M., Todd, N., Weiskopf, N., 2015b. An evaluation of prospective motion correction (PMC) for high resolution quantitative MRI. Front. Neurosci. 9. https://doi.org/10.3389/fnins.2015.00097

Carver, P.L. (Ed.), 2019. 4. IRONING OUT THE BRAIN, in: Essential Metals in Medicine: Therapeutic Use and Toxicity of Metal Ions in the Clinic. De Gruyter, pp. 87–122. https://doi.org/10.1515/9783110527872-004

Concha, L., Livy, D.J., Beaulieu, C., Wheatley, B.M., Gross, D.W., 2010. In Vivo Diffusion Tensor Imaging and Histopathology of the Fimbria-Fornix in Temporal Lobe Epilepsy. J. Neurosci. 30, 996–1002. https://doi.org/10.1523/JNEUROSCI.1619-09.2010

Dahnke, R., Yotter, R.A., Gaser, C., 2013. Cortical thickness and central surface estimation. NeuroImage 65, 336–348. https://doi.org/10.1016/j.neuroimage.2012.09.050

Daugherty, A., Raz, N., 2013. Age-related differences in iron content of subcortical nuclei observed in vivo: A meta-analysis. NeuroImage 70, 113–121. https://doi.org/10.1016/j.neuroimage.2012.12.040

de Pasquale, F., Cherubini, A., Péran, P., Caltagirone, C., Sabatini, U., 2013. Influence of white matter fiber orientation on R2* revealed by MRI segmentation. J. Magn. Reson. Imaging 37, 85–91. https://doi.org/10.1002/jmri.23801

De Santis, S., Barazany, D., Jones, D.K., Assaf, Y., 2016. Resolving relaxometry and diffusion properties within the same voxel in the presence of crossing fibres by combining inversion recovery and diffusion-weighted acquisitions: Resolving Myelin and Axonal Properties with IR-DTI. Magn. Reson. Med. 75, 372–380.

https://doi.org/10.1002/mrm.25644

Draganski, B., Ashburner, J., Hutton, C., Kherif, F., Frackowiak, R.S.J., Helms, G., Weiskopf, N., 2011. Regional specificity of MRI contrast parameter changes in normal ageing revealed by voxel-based quantification (VBQ). NeuroImage 55, 1423–1434. https://doi.org/10.1016/j.neuroimage.2011.01.052

Drayer, B., Burger, P., Darwin, R., Riederer, S., Herfkens, R., Johnson, G.A., 1986. MRI of brain iron. Am. J. Roentgenol. 147, 103–110. https://doi.org/10.2214/ajr.147.1.103.

Eickhoff, S., Walters, N.B., Schleicher, A., Kril, J., Egan, G.F., Zilles, K., Watson, J.D.G., Amunts, K., 2005. High-resolution MRI reflects myeloarchitecture and cytoarchitecture of human cerebral cortex. Hum. Brain Mapp. 24, 206–215. https://doi.org/10.1002/hbm.20082

European Society of Radiology (ESR), 2015. Magnetic Resonance Fingerprinting - a promising new approach to obtain standardized imaging biomarkers from MRI. Insights Imaging 6, 163–165. https://doi.org/10.1007/s13244-015-0403-3

Filo, S., Shtangel, O., Salamon, N., Kol, A., Weisinger, B., Shifman, S., Mezer, A.A., 2019. Disentangling molecular alterations from water-content changes in the aging human brain using quantitative MRI. Nat. Commun. 10, 3403. https://doi.org/10.1038/s41467-019-11319-1

Fukunaga, M., Li, T.-Q., van Gelderen, P., de Zwart, J.A., Shmueli, K., Yao, B., Lee, J., Maric, D., Aronova, M.A., Zhang, G., Leapman, R.D., Schenck, J.F., Merkle, H., Duyn, J.H., 2010. Layer-specific variation of iron content in cerebral cortex as a source of MRI contrast. Proc. Natl. Acad. Sci. 107, 3834–3839. https://doi.org/10.1073/pnas.0911177107

Geeraert, B.L., Lebel, R.M., Lebel, C., 2019. A multiparametric analysis of white matter maturation during late childhood and adolescence. Hum. Brain Mapp. 40, 4345–4356. https://doi.org/10.1002/hbm.24706

Guilfoyle, D.N., Dyakin, V.V., O'Shea, J., Pell, G.S., Helpern, J.A., 2003. Quantitative measurements of proton spin-lattice (T1) and spin-spin (T2) relaxation times in the mouse brain at 7.0 T. Magn. Reson. Med. 49, 576–580. https://doi.org/10.1002/mrm.10371

Han, X., Pham, D.L., Tosun, D., Rettmann, M.E., Xu, C., Prince, J.L., 2004. CRUISE: Cortical reconstruction using implicit surface evolution. NeuroImage 23, 997–1012. https://doi.org/10.1016/j.neuroimage.2004.06.043

Helms, G., Dathe, H., Kallenberg, K., Dechent, P., 2008. High-resolution maps of magnetization transfer with inherent correction for RF inhomogeneity and T_1 relaxation obtained from 3D FLASH MRI: Saturation and Relaxation in MT FLASH. Magn. Reson. Med. 60, 1396–1407. https://doi.org/10.1002/mrm.21732

Henkelman, R.M., Stanisz, G.J., Graham, S.J., 2001. Magnetization transfer in MRI: a review. NMR Biomed. 14, 57–64. https://doi.org/10.1002/nbm.683

Hentschel, F., Kreis, M., Damian, M., Krumm, B., 2004. Leistet die Magnetization-Transfer-Ratio (MTR) einen Beitrag zur Diagnostik und Differenzialdiagnostik der Demenzen? RöFo - Fortschritte Auf Dem Geb. Röntgenstrahlen Bildgeb. Verfahr. 176, 1743–1749. https://doi.org/10.1055/s-2004-813650

Horn, A., Ostwald, D., Reisert, M., Blankenburg, F., 2014. The structural–functional connectome and the default mode network of the human brain. NeuroImage 102, 142–151. https://doi.org/10.1016/j.neuroimage.2013.09.069

Huntenburg, J.M., Bazin, P.-L., Goulas, A., Tardif, C.L., Villringer, A., Margulies, D.S., 2017. A Systematic Relationship Between Functional Connectivity and Intracortical Myelin in the

Human Cerebral Cortex. Cereb. Cortex 27, 981–997.
https://doi.org/10.1093/cercor/bhx030

Huntenburg, J.M., Steele, C.J., Bazin, P.-L., 2018. Nighres: processing tools for high-resolution neuroimaging. GigaScience 7. https://doi.org/10.1093/gigascience/giy082

Jenkinson, M., Beckmann, C.F., Behrens, T.E.J., Woolrich, M.W., Smith, S.M., 2012. FSL. NeuroImage 62, 782–790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Jones, D.K., Knösche, T.R., Turner, R., 2013. White matter integrity, fiber count, and other fallacies: The do's and don'ts of diffusion MRI. NeuroImage 73, 239–254. https://doi.org/10.1016/j.neuroimage.2012.06.081

Khattar, N., Triebswetter, C., Kiely, M., Ferrucci, L., Resnick, S.M., Spencer, R.G., Bouhrara, M., 2021. Investigation of the association between cerebral iron content and myelin content in normative aging using quantitative magnetic resonance neuroimaging. NeuroImage 239, 118267. https://doi.org/10.1016/j.neuroimage.2021.118267

Khodanovich, M.Y., Sorokina, I.V., Glazacheva, V.Y., Akulov, A.E., Nemirovich-Danchenko, N.M., Romashchenko, A.V., Tolstikova, T.G., Mustafina, L.R., Yarnykh, V.L., 2017. Histological validation of fast macromolecular proton fraction mapping as a quantitative myelin imaging method in the cuprizone demyelination model. Sci. Rep. 7, 46686. https://doi.org/10.1038/srep46686

Kirilina, E., Helbling, S., Morawski, M., Pine, K., Reimann, K., Jankuhn, S., Dinse, J., Deistung, A., Reichenbach, J.R., Trampel, R., Geyer, S., Müller, L., Jakubowski, N., Arendt, T., Bazin, P.-L., Weiskopf, N., 2020. Superficial white matter imaging: Contrast mechanisms and whole-brain in vivo mapping. Sci. Adv. 6, eaaz9281. https://doi.org/10.1126/sciadv.aaz9281

Kor, D., Birkl, C., Ropele, S., Doucette, J., Xu, T., Wiggermann, V., Hernández-Torres, E., Hametner, S., Rauscher, A., 2019. The role of iron and myelin in orientation dependent R_2^* of white matter. NMR Biomed. e4092. https://doi.org/10.1002/nbm.4092

Leppert, I.R., Andrews, D.A., Campbell, J.S.W., Park, D.J., Pike, G.B., Polimeni, J.R., Tardif, C.L., 2021. Efficient whole-brain tract-specific T_1 mapping at 3T with slice-shuffled inversion-recovery diffusion-weighted imaging. Magn. Reson. Med. 86, 738–753. https://doi.org/10.1002/mrm.28734

Leutritz, T., Seif, M., Helms, G., Samson, R.S., Curt, A., Freund, P., Weiskopf, N., 2020. Multiparameter mapping of relaxation (R_1, R_2*), proton density and magnetization transfer saturation at $3T$: A multicenter dual-vendor reproducibility and repeatability study. Hum. Brain Mapp. 41, 4232–4247. https://doi.org/10.1002/hbm.25122

Lüsebrink, F., Wollrab, A., Speck, O., 2013. Cortical thickness determination of the human brain using high resolution 3T and 7T MRI data. NeuroImage 70, 122–131. https://doi.org/10.1016/j.neuroimage.2012.12.016

Lutti, A., Dick, F., Sereno, M.I., Weiskopf, N., 2014. Using high-resolution quantitative mapping of R1 as an index of cortical myelination. NeuroImage 93, 176–188. https://doi.org/10.1016/j.neuroimage.2013.06.005

Lutti, A., Stadler, J., Josephs, O., Windischberger, C., Speck, O., Bernarding, J., Hutton, C., Weiskopf, N., 2012. Robust and Fast Whole Brain Mapping of the RF Transmit Field B1 at 7T. PLoS ONE 7, e32379. https://doi.org/10.1371/journal.pone.0032379

Mancini, M., Karakuzu, A., Cohen-Adad, J., Cercignani, M., Nichols, T.E., Stikov, N., 2020. An interactive meta-analysis of MRI biomarkers of myelin. eLife 9, e61523. https://doi.org/10.7554/eLife.61523

Mangeat, G., Govindarajan, S.T., Mainero, C., Cohen-Adad, J., 2015. Multivariate combination of magnetization transfer, T_2* and B0 orientation to study the myelo-architecture of the

in vivo human cortex. NeuroImage 119, 89–102.
https://doi.org/10.1016/j.neuroimage.2015.06.033

Manning, A.P., Chang, K.L., MacKay, A.L., Michal, C.A., 2017. The physical mechanism of "inhomogeneous" magnetization transfer MRI. J. Magn. Reson. 274, 125–136.
https://doi.org/10.1016/j.jmr.2016.11.013

Margulies, D.S., Ghosh, S.S., Goulas, A., Falkiewicz, M., Huntenburg, J.M., Langs, G., Bezgin, G., Eickhoff, S.B., Castellanos, F.X., Petrides, M., Jefferies, E., Smallwood, J., 2016. Situating the default-mode network along a principal gradient of macroscale cortical organization. Proc. Natl. Acad. Sci. 113, 12574–12579.
https://doi.org/10.1073/pnas.1608282113

Marques, J.P., Khabipova, D., Gruetter, R., 2017. Studying cyto and myeloarchitecture of the human cortex at ultra-high field with quantitative imaging: R1, R2* and magnetic susceptibility. NeuroImage 147, 152–163.
https://doi.org/10.1016/j.neuroimage.2016.12.009

Mechelli, A., 2005. Structural Covariance in the Human Cortex. J. Neurosci. 25, 8303–8310.
https://doi.org/10.1523/JNEUROSCI.0357-05.2005

Metere, R., Kober, T., Möller, H.E., Schäfer, A., 2017. Simultaneous Quantitative MRI Mapping of T1, T2* and Magnetic Susceptibility with Multi-Echo MP2RAGE. PLOS ONE 12, e0169265. https://doi.org/10.1371/journal.pone.0169265

Möller, H.E., Bossoni, L., Connor, J.R., Crichton, R.R., Does, M.D., Ward, R.J., Zecca, L., Zucca, F.A., Ronen, I., 2019. Iron, Myelin, and the Brain: Neuroimaging Meets Neurobiology. Trends Neurosci. 42, 384–401. https://doi.org/10.1016/j.tins.2019.03.009

Paquola, C., Vos De Wael, R., Wagstyl, K., Bethlehem, R.A.I., Hong, S.-J., Seidlitz, J., Bullmore, E.T., Evans, A.C., Misic, B., Margulies, D.S., Smallwood, J., Bernhardt, B.C., 2019. Microstructural and functional gradients are increasingly dissociated in transmodal cortices. PLOS Biol. 17, e3000284. https://doi.org/10.1371/journal.pbio.3000284

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., n.d. Scikit-learn: Machine Learning in Python. Mach. Learn. PYTHON 6.

Pierpaoli, C., Basser, P.J., 1996. Toward a quantitative assessment of diffusion anisotropy. Magn. Reson. Med. 36, 893–906. https://doi.org/10.1002/mrm.1910360612

Raffelt, D.A., Tournier, J.-D., Smith, R.E., Vaughan, D.N., Jackson, G., Ridgway, G.R., Connelly, A., 2017. Investigating white matter fibre density and morphology using fixel-based analysis. NeuroImage 144, 58–73. https://doi.org/10.1016/j.neuroimage.2016.09.029

Raichle, M.E., 2015. The Brain's Default Mode Network. Annu. Rev. Neurosci. 38, 433–447.
https://doi.org/10.1146/annurev-neuro-071013-014030

Roche, A., Forbes, F., 2014. Partial Volume Estimation in Brain MRI Revisited 8.

Seidlitz, J., Váša, F., Shinn, M., Romero-Garcia, R., Whitaker, K.J., Vértes, P.E., Wagstyl, K., Kirkpatrick Reardon, P., Clasen, L., Liu, S., Messinger, A., Leopold, D.A., Fonagy, P., Dolan, R.J., Jones, P.B., Goodyer, I.M., Raznahan, A., Bullmore, E.T., 2018. Morphometric Similarity Networks Detect Microscale Cortical Organization and Predict Inter-Individual Cognitive Variation. Neuron 97, 231-247.e7.
https://doi.org/10.1016/j.neuron.2017.11.039

Sereno, M.I., Lutti, A., Weiskopf, N., Dick, F., 2013. Mapping the Human Cortical Surface by Combining Quantitative T1 with Retinotopy†. Cereb. Cortex 23, 2261–2268.
https://doi.org/10.1093/cercor/bhs213

Shams, Z., Norris, D.G., Marques, J.P., 2019. A comparison of in vivo MRI based cortical myelin mapping using T1w/T2w and R1 mapping at 3T. PLOS ONE 14, e0218089.

https://doi.org/10.1371/journal.pone.0218089

Stüber, C., Morawski, M., Schäfer, A., Labadie, C., Wähnert, M., Leuze, C., Streicher, M., Barapatre, N., Reimann, K., Geyer, S., Spemann, D., Turner, R., 2014. Myelin and iron concentration in the human brain: A quantitative study of MRI contrast. NeuroImage 93, 95–106. https://doi.org/10.1016/j.neuroimage.2014.02.026

Szafer, A., Zhong, J., Anderson, A.W., Gore, J.C., 1995. Diffusion-weighted imaging in tissues: Theoretical models. NMR Biomed. 8, 289–296. https://doi.org/10.1002/nbm.1940080704

Tabelow, K., Balteau, E., Ashburner, J., Callaghan, M.F., Draganski, B., Helms, G., Kherif, F., Leutritz, T., Lutti, A., Phillips, C., Reimer, E., Ruthotto, L., Seif, M., Weiskopf, N., Ziegler, G., Mohammadi, S., 2019. hMRI – A toolbox for quantitative MRI in neuroscience and clinical research. NeuroImage 194, 191–210. https://doi.org/10.1016/j.neuroimage.2019.01.029

Tardif, C.L., Gauthier, C.J., Steele, C.J., Bazin, P.-L., Schäfer, A., Schaefer, A., Turner, R., Villringer, A., 2016. Advanced MRI techniques to improve our understanding of experience-induced neuroplasticity. NeuroImage 131, 55–72. https://doi.org/10.1016/j.neuroimage.2015.08.047

Tardif, C.L., Schäfer, A., Waehnert, M., Dinse, J., Turner, R., Bazin, P.-L., 2015. Multi-contrast multi-scale surface registration for improved alignment of cortical areas. NeuroImage 111, 107–122. https://doi.org/10.1016/j.neuroimage.2015.02.005

Tardif, C.L., Steele, C.J., Lampe, L., Bazin, P.-L., Ragert, P., Villringer, A., Gauthier, C.J., 2017. Investigation of the confounding effects of vasculature and metabolism on computational anatomy studies. NeuroImage 149, 233–243. https://doi.org/10.1016/j.neuroimage.2017.01.025

Trampel, R., Bazin, P.-L., Pine, K., Weiskopf, N., 2019. In-vivo magnetic resonance imaging (MRI) of laminae in the human cortex. NeuroImage 197, 707–715. https://doi.org/10.1016/j.neuroimage.2017.09.037

Tustison, N.J., Cook, P.A., Klein, A., Song, G., Das, S.R., Duda, J.T., Kandel, B.M., van Strien, N., Stone, J.R., Gee, J.C., Avants, B.B., 2014. Large-scale evaluation of ANTs and FreeSurfer cortical thickness measurements. NeuroImage 99, 166–179. https://doi.org/10.1016/j.neuroimage.2014.05.044

Uddin, Md.N., Figley, T.D., Solar, K.G., Shatil, A.S., Figley, C.R., 2019. Comparisons between multi-component myelin water fraction, T1w/T2w ratio, and diffusion tensor imaging measures in healthy human brain structures. Sci. Rep. 9, 2500. https://doi.org/10.1038/s41598-019-39199-x

van Es, A.C.G.M., van der Flier, W.M., Admiraal-Behloul, F., Olofsen, H., Bollen, E.L.E.M., Middelkoop, H.A.M., Weverling-Rijnsburger, A.W.E., Westendorp, R.G.J., van Buchem, M.A., 2006. Magnetization transfer imaging of gray and white matter in mild cognitive impairment and Alzheimer's disease. Neurobiol. Aging 27, 1757–1762. https://doi.org/10.1016/j.neurobiolaging.2005.09.042

Waehnert, M.D., Dinse, J., Schäfer, A., Geyer, S., Bazin, P.-L., Turner, R., Tardif, C.L., 2016. A subject-specific framework for in vivo myeloarchitectonic analysis using high resolution quantitative MRI. NeuroImage 125, 94–107. https://doi.org/10.1016/j.neuroimage.2015.10.001

Wardlaw, J.M., Hernandez, M.C.V., Munoz-Maniega, S., 2016. What are White Matter Hyperintensities Made of?: Relevance to Vascular Cognitive Impairment. J. Am. Heart Assoc. 5. https://doi.org/10.1161/JAHA.115.002006

Weiskopf, N., Edwards, L.J., Helms, G., Mohammadi, S., Kirilina, E., 2021. Quantitative magnetic resonance imaging of brain anatomy and in vivo histology. Nat. Rev. Phys. 3,

570–588. https://doi.org/10.1038/s42254-021-00326-1

Weiskopf, N., Mohammadi, S., Lutti, A., Callaghan, M.F., 2015. Advances in MRI-based computational neuroanatomy: from morphometry to in-vivo histology. Curr. Opin. Neurol. 28, 313–322. https://doi.org/10.1097/WCO.0000000000000222

Weiskopf, N., Suckling, J., Williams, G., Correia, M.M., Inkster, B., Tait, R., Ooi, C., Bullmore, E.T., Lutti, A., 2013. Quantitative multi-parameter mapping of R1, PD*, MT, and R2* at 3T: a multi-center validation. Front. Neurosci. 7. https://doi.org/10.3389/fnins.2013.00095

Yarnykh, V.L., Prihod'ko, I.Y., Savelov, A.A., Korostyshevskaya, A.M., 2018. Quantitative Assessment of Normal Fetal Brain Myelination Using Fast Macromolecular Proton Fraction Mapping. Am. J. Neuroradiol. 39, 1341–1348. https://doi.org/10.3174/ajnr.A5668

Yeatman, J.D., Wandell, B.A., Mezer, A.A., 2014. Lifespan maturation and degeneration of human brain white matter. Nat. Commun. 5, 4932. https://doi.org/10.1038/ncomms5932

Yeo, B.T., Krienen, F.M., Sepulcre, J., Sabuncu, M.R., Lashkari, D., Hollinshead, M., Roffman, J.L., Smoller, J.W., Zöllei, L., Polimeni, J.R., Fischl, B., Liu, H., Buckner, R.L., 2011. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. J. Neurophysiol. 106, 1125–1165. https://doi.org/10.1152/jn.00338.2011

Zatorre, R.J., Fields, R.D., Johansen-Berg, H., 2012. Plasticity in gray and white: neuroimaging changes in brain structure during learning. Nat. Neurosci. 15, 528–536. https://doi.org/10.1038/nn.3045

Zielinski, B.A., Gennatas, E.D., Zhou, J., Seeley, W.W., 2010. Network-level structural covariance in the developing brain. Proc. Natl. Acad. Sci. 107, 18191–18196. https://doi.org/10.1073/pnas.1003109107