Conformational change in the monomeric alpha-synuclein imparts fibril polymorphs

Short title: The monomeric αSyn imparts fibril polymorphs

Cesar Aguirre[1,†], Kensuke Ikenaka[1*,†], Masatomo So[2], Takahiro Maruno[3], Keiichi Yamaguchi[4], Kichitaro Nakajima[4], Chi-Jing Choong[1], Kei Nabekura[1], Goichi Beck[1], Kentaro Tomii[5], Yu Yamamori[5], Junko Doi[1], Tomoyasu Matsubara[6], Maho Morishima[6], Keita Kakuda[1], Makoto Hideshima[1], Yasuyoshi Kimura[1], Seiichi Nagano[1], Kousuke Baba[1], Shigeo Murayama[6,7], Hirotsugu Ogi[8], Yoshitaka Nagai[9], Yasushi Kawata[10], Susumu Uchiyama[3], Yohei Miyanoiri[11], Yuji Goto[4], and Hideki Mochizuki[1*]

[1] Department of Neurology, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan.

[2] Institute for Protein Research, Osaka University, Japan, 3-2 Yamadaoka, Suita, Osaka 565-0871, Japan.

[3] Department of Biotechnology, Graduate School of Engineering, Osaka University, Japan, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan.

[4] Global Center for Medical Engineering and Informatics, Osaka University, Japan, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan.

[5] Artificial Intelligence Research Center (AIRC), National Institute of Advanced Industrial Science and Technology (AIST), 2-4-7 Aomi, Koto-ku, Tokyo, 135-0064, Japan.

[6] Department of Neuropathology (the Brain Bank for Aging Research), Tokyo Metropolitan Geriatric Hospital and Institute of Gerontology, 35-2 Sakae-cho, Itabashi-ku, Tokyo, 173-0015, Japan.

[7] Brain Bank for Neurodevelopmental, Neurological and Psychiatric Disorders, United Graduate School of Child Development, Osaka University, 2-2, Yamadaoka, Suita- shi, Osaka, 565-0871, Japan.

[8] Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan.

[9] Department of Neurology, Faculty of Medicine, Kindai University, Osaka-Sayama, Osaka, Japan.

[10] Department of Chemistry and Biotechnology, Graduate School of Engineering, Tottori University, Tottori 680-8552, Japan.

[11] Laboratory for Ultra-High magnetic field NMR spectroscopy, Research Center for Next-Generation Protein Sciences, Institute for Protein Research, Osaka University, Japan, 3-2 Yamadaoka, Suita, Osaka 565-0871, Japan.

*Correspondence to Dr. Kensuke Ikenaka, M.D, Ph.D., and Prof. Hideki Mochizuki, M.D., Ph.D.

Department of Neurology, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Osaka 565-0871, Japan. Phone: 81-6-6879-3571.

E-mail: ikenaka@neurol.med.osaka-u.ac.jp, hmochizuki@neurol.med.osaka-u.ac.jp

† These authors contributed equally to this work.

Abstract

α-Synuclein inclusion bodies are a pathological hallmark of several neurodegenerative diseases. Although it has been hypothesized a relationship between fibril polymorphism and different pathologies, the molecular origins of polymorphism are not understood. Here, employing biophysical approaches, we demonstrate that the conformational state of the monomeric αSyn is responsible for fibril polymorphism: αSyn can exist as a compact monomer that produces rod fibrils, and as extended monomers that generate twisted fibrils. Using NMR, we found that the compaction relies on a polar interaction between the initial part of the NAC region and a wide section of the C-terminus domain. The compaction can be commonly affected by changes in the chemical environment, like NaCl, the presence of Ca^{2+} or cellular components, like endotoxins, that alter the interaction NAC/C-terminus domain. Our compaction model also provides mechanistic insights that explain how the behavior of the C-terminus domain imparts the polymorphism during the fibril formation.

Introduction

The accumulation of alpha-synuclein (αSyn) is a pathological hallmark of Parkinson's disease (PD) and multiple system atrophy (MSA)[1–3]; however, although these synucleinopathies are produced by the same protein, they exhibit clinically distinctive features. Recently, we have developed experimental strategies to study directly the structural characteristics of inclusions in patient's brain employing Fourier-Transform Infrared microscopy (FTIR) and Small-Angle X-Ray Scattering (SAXS), proving that Lewy Bodies (LBs) contain cross β-sheet αSyn fibrils, while the amount of β-sheet structure in LBs and glial cell inclusions in MSA patients is different[4–6]. These studies indicate that αSyn aggregates associated with PD and MSA correspond to different structural strains. Supporting our data, evidence corroborating distinctive strains in PD and MSA was provided analyzing amplified fibrils from the patients' brain using the protein misfolded cyclic amplification (PMCA) method[7,8], and through the direct observation of accumulated fibrils in patient's brain by cryo-electron microscopy (Cryo-EM)[9]. Moreover, brain homogenates extracted from PD and MSA patients, as well as structurally different preformed fibrils, induced different cellular transmissibility, toxicity, membrane disruption, etc.[7,10–16], leading to the hypothesis that pathological differences among synucleinopathies can be explained in terms of morphological variations in the accumulated αSyn fibrils in the brain[17,18]. *In vitro*, different strains of αSyn fibrils can be produced when the fibrils are generated under conditions of low or high ionic strength, whose characteristics resemble the morphological properties of the fibrils observed in PD and MSA patients[16,19,20].

Despite the substantial progress in the analysis of strains amplified from patients' tissues and created *in vitro,* the molecular mechanisms that explain the origins of different strains from the same wild type αSyn remain poorly understood. To address this question, we focused on the conformational change of the monomeric αSyn. Several studies have reached a consensus about the disordered nature of αSyn[21–24]; however, αSyn can also adopt some partially folded structures under acidic pH and high temperature[25,26]. Furthermore, some groups have proposed that these conformations represent intermediate species during the formation of fibrils[27–29]. Nevertheless, it remains unclear whether these conformational variations exist in the physiological environment and what their relationship is to polymorphisms. In this study, we propose that the molecular mechanisms responsible for polymorphisms are linked to the monomeric conformational state, which could be altered in the cell by the environment, eventually leading to the development of specific synucleinopathies.

Results

αSyn distinct fibril strains related to synucleinopathies can be modulated by the conformational state of the monomer

To understand the morphological and biochemical characteristics of PD- and MSA-related fibril polymorphisms, we amplified amyloid fibrils from PD and MSA patients' brain by seeding, using amygdalae homogenates. The presence of αSyn aggregates in the amygdalae for all the cases was confirmed by immunohistochemical analysis (**Fig. S1**). First, we monitored the kinetics of fibril amplification by our HANABI assay[30,31] (**Figure 1.A**), an instrument that consists of an optimized sonoreactor equipment developed by our group, that applies ultrasonic irradiation pulses individually to each one of the wells of a 96-wells microplate, and records the ThT fluorescence in function of time, named HANABI-2000[31]. This instrument accelerates substantially the rate of fibril formation reaction compared with the traditional RT-QuIC methods[32,33], from several days to a few hours. Then, we analyzed the morphological properties by transmission electron microscopy (TEM, **Figure 1.B**). As previously reported[8], we observed that fibrils amplified from PD patients' brain exhibit a higher ThT intensity than those obtained from MSA. The microscopic visualization showed that PD-amplified fibrils are characterized by rod-like fibrils, with two parallelly aligned protofilaments; while MSA-derived fibrils depicted a twisted morphology, in which two protofilaments entangle over each other. These results highlight the importance of a deeper investigation to reveal the molecular mechanisms responsible for polymorphism.

Previous reports have shown that different fibril morphologies, each one with distinctive biophysical and biochemical properties, can be generated *in vitro* employing low or high ionic strength conditions[16,19,20]. A deeper knowledge of the fibril morphology phase diagram in function of the concentration of NaCl would enable us to understand the upstream molecular events related to the polymorphism of fibrils. Therefore, we examined the αSyn fibril formation by ThT assay in the presence of 50 mM TRIS pH 7.4 supplemented with 0 to 500 mM NaCl (**Figure 1.C**). Interestingly, below 300 mM NaCl, we found typical sigmoidal kinetic curves with relatively low maximal ThT intensity values (< 0.2 A. U.). At 300 mM NaCl, a wider distribution of curves with both low and high maximal ThT was detected, while above 400 mM NaCl, just fibrils with high values of ThT intensity were observed. To determine the phase diagram of ThT intensity of fibrils, we plotted the maximal values of ThT as a function of the concentration of NaCl, finding a region dominated by low maximal ThT values below 300 mM NaCl, and a region with high maximal ThT values above 300 mM. We defined the inflection point of the phase diagram as the concentration of NaCl, 300 mM, in which fibrils with both low and high ThT can be produced by chance. Moreover, we evaluated by TEM the morphological properties of the fibrils created at different concentrations of NaCl (**Figure 1.D, upper and lower-right**). First, it was found that the low-ThT fibrils obtained below 300 mM NaCl exhibit a

twisted morphology, with similar characteristics than those observed in the MSA-amplified fibrils. The fibrils produced above 300 mM NaCl with high-ThT exhibit a rod morphology, resembling the properties of the PD-derived fibrils.

In addition, we also confirmed by SDS-PAGE densitometry that the yield of both the low and high ThT fibrils created at low and high salt, respectively, is higher than 95 % (**Fig. S2.A**), discarding the possibility of incomplete reactions or the lack of fibrillation products when low ThT is observed. Finally, we recorded the FTIR spectra of both low-salt and high-salt fibrils, finding in both cases typical bands of β-sheet structure around 1630 cm^{-1}, demonstrating the presence of amyloid fibrils in both NaCl conditions, and excluding the formation of amorphous aggregates (**Fig. S2.B**). The deconvolution of the FTIR spectra showed that the contents of β-sheet structure is higher for the fibrils created at 50 mM NaCl, indicating structural differences between the fibrils (**Fig. S2.C**).

For the biochemical characterization of fibrils, we developed a proteinase K resistance assay (PKR) to evaluate and compare the digestion patterns of fibrils with different morphologies. Considering the first five bands with the highest molecular in the proteolysis patterns after 60 minutes of reaction, rod-like fibrils showed a digestion profile with stronger band intensities for the second (B2) and fourth (B4) bands, while twisted fibrils presented a weaker B2 band (**Fig. S2.D-E**). Employing this PKR assay, we evaluated the digestion patterns of fibrils obtained at 50 mM and 500 mM NaCl concentrations, where the differential digestion profiles for twisted and rod fibrils, respectively, were confirmed (**Figure 1.E**, left). Furthermore, we tested the PKR assay of selected fibrils created at 300 mM NaCl with low and high maximal ThT fibrils, whose digestion patterns are in good agreement with the morphology visualized by TEM (**Figure 1.E**, center). Finally, we also evaluated the digestion patterns of fibrils amplified from PD and MSA brain samples, in which degradation profiles of rod-like and twisted morphologies, respectively, were detected (**Figure 1.E**, right), corroborating the biochemical similarities between *in vitro* created fibrils and fibril derived from patients' brain. Taken together, these results corroborated that the transition of maximal ThT values in function of the concentration of NaCl reflects the morphological phase diagram of the fibrils, which might explain the origins of the fibril diversities in the patients of PD and MSA.

We then explored the upstream molecular events that influence the phase diagram of the fibrils. It has been reported previously that αSyn can adopt different conformational states in solution[21,23,26]. In order to investigate whether these conformations are related to the molecular events in the early monomeric state that lead to the formation of specific fibrils, we started solving a solution of αSyn by analytical size exclusion chromatography (SEC) at 300 mM NaCl (**Figure 1.F,** upper). At this ionic strength, αSyn can produce both rod and twisted morphologies. Interestingly, the full-length αSyn monomer, whose molecular weight is 14.6 kDa, was observed as a single asymmetric peak, widely

distributed in retention volume values ranging from molecular weights of 15 kDa to 60 kDa for hypothetical spherical-shaped globular proteins. The peak center of αSyn was observed around the retention volume that corresponds to 50 kDa globular proteins. This result suggests that αSyn monomers can adopt conformations from compact to more extended states in which the apparent hydrodynamic size increases. We confirmed by Western blotting that all the collected αSyn fractions separated by SEC consist of intact full-length monomers that have access to conformations with different hydrodynamic radii (**Figure 1.F**, lower). In addition, we also evaluated the retention volume of the monomeric αSyn between 50 and 500 mM NaCl, finding a constant increase in the retention volume between 50 mM and 300 mM NaCl, but no significant change above 300 mM NaCl (**Figure 1.G** and **Fig. S3.A**), indicating that the monomers have reached a plateau conformational phase around 300 mM NaCl, in which the protein is found in a more compact conformation than in lower concentrations of NaCl. Moreover, we evaluated the retention volume of solutions of bovine serum albumin (BSA), in the presence of NaCl at 150 mM and 500 mM (**Fig. S3.B, C**). No variation in the retention volume was detected, corroborating that the changes in the retention volume observed in αSyn are not an artifact of the differences in the ionic strength.

A more exhaustive evaluation of the hydrodynamic radius of the monomeric αSyn was performed using Analytical Ultracentrifugation (AUC) at low (150 mM) and high (500 mM) concentrations of NaCl. By AUC Sedimentation Velocity, we observed a significant decrease in the values of the sedimentation coefficient, from 1.136 ± 0.012 S at 150 mM NaCl to 1.108 ± 0.011 S at 500 mM NaCl (**Figure 1.H**), indicating a compaction from 150 mM to 500 mM NaCl. These measurements also confirmed that αSyn is found mainly as a monomer (> 98 %). Taken together, our results indicate that the size distribution of the monomeric αSyn changes from an extended conformation at low ionic strength to a more compact form above 500 mM NaCl.

The behavior of the C-terminus determines the compaction of αSyn

To determine the molecular events responsible for the compaction of αSyn monomers, we employed different NaCl concentrations (0-500 mM) to record ^{1}H-^{15}N HSQC spectra (**Figure 2.A** and **Fig S4.A**). The analysis of the peak intensities as a function of NaCl concentration, using the spectrum obtained at 0 mM NaCl as reference, allowed us to detect a constant decrease in the peak intensity at the initial part of the NAC region and in the majority of residues in the C-terminal domain residues (**Figure 2.B** and **Fig. S4.B**, left), detected by both difference ($I_x - I_{0mM}$) and ratio (I_x / I_{0mM}) of peak intensities. Additionally, some peaks disappeared upon the increase in NaCl concentration, located mostly in the NAC region (**Fig. S4.B**, marked with *). Remarkably, the peaks that experienced the most significant changes in the NAC domain were attributed to aliphatic and non-charged polar residues. Among the peaks showing changes in the C-terminal domain, half were attributed to

aliphatic, aromatic, or non-charged polar residues, and the others to negatively charged residues (**Figure 2.C**). The analysis of chemical shift perturbation (**Fig. S2.B**, right) did not show significant variations.

To determine the relative flexibility differences between the monomers at distinct ionic strength, we performed heteronuclear Overhauser effect (hetNOE) experiments with αSyn incubated in 50 mM TRIS pH 7.4 supplemented with 0 mM and 500 mM NaCl (**Fig. S4.C**). We found that most of the changes, but not the only ones, were seen in the NAC region (highlighted with a gray square), where several residues show more negative hetNOE values at 0 mM NaCl, while at 500 mM NaCl some of these residues exhibit a higher hetNOE values or they have even disappeared in ^{1}H-^{15}N HSQC spectrum with ^{1}H saturation. The C-terminal domain shows just few changes between the residues 120 and 130, where hetNOE is higher. These results suggest that although the monomers retain their intrinsically disordered nature (**Fig. S4.D**), at higher ionic strength, both the NAC region and the C-terminal domain suffer a decrease in the flexibility, probably as consequence of the compaction of the monomer.

Additionally, we explored 576 structures deposited in the Protein Ensemble Data Bank, PED00024[34] to corroborate the experimental HSQC spectra. In all the salt concentrations, the closest structure to our data, evaluated by the lowest root-mean-square (RMS), was an extended conformation (**Fig. S4.E**, left). Interestingly, the second lowest RMS value at 500 mM NaCl had a more compact conformation with the C-terminal domain interacts with the central region of the protein (**Fig. S4.E**, right), and the RMS value of this structure decreased as the salt concentration increased. These results are consistent with our experimental data that the formation of compact monomer increased when NaCl was added.

Calcium ions modulate the monomeric compaction and shift the inflection point of fibril morphologies to the rod fibril.

It has been reported that the cation Ca^{2+} binds to the negatively charged residues of the C-terminal domain, and alters the conformation of αSyn[35]. With this context, we assessed Ca^{2+} ions as modulators of the conformational state of αSyn. First, we examined the fibril formation in presence of 200 Eq. Ca^{2+} at different ionic strength, ranging from 21 to 500 mM (**Figure 3.A** and **Fig S5.A**). In the presence of Ca^{2+}, below 200 mM ionic strength (IS) a variety of ThT values was seen, while above 200 mM IS just high ThT fibrils were observed. These results indicate that calcium promotes the formation of fibrils with high ThT at lower IS, and even prevents the formation of exclusively low ThT fibrils at low IS. The morphology of the fibrils formed at an IS of 200 mM in the presence of Ca^{2+} was tested by TEM (**Figure 3.B**) and PKR (**Figure 3.C**), that confirmed the morphological properties of the rod fibrils. In order to verify the relationship between fibril morphology and monomeric

conformation, we solved solutions of αSyn in the presence and absence of 200 Eq. Ca^{2+} at an IS of 200 mM by SEC (**Figure 3.D**), finding a significant higher retention volume when calcium is present in the solution and suggesting a connection between the rod morphology and a compact monomeric state.

To determine the residues affected in the compact monomer induced by calcium, we recorded ^{1}H-^{15}N HSQC spectra in the presence and absence of 200 Eq. Ca^{2+} at an ionic strength equivalent to 200 mM NaCl (**Figure 3.E**). The analysis of the peak intensities by both difference ($I_{200Ca} - I_{0Ca}$) and ratio (I_{200Ca} / I_{0Ca}), using the spectrum recorded in the absence of Ca^{2+} at 200 mM NaCl as reference, showed again a decrease in the peak intensity at the initial part of the NAC region and in residues in the C-terminal domain residues (**Figure 3.F,** upper and middle). The peaks showing changes share similarities with those detected by NaCl dependence in both the NAC region and in the C-terminus domain. The analysis of chemical shift perturbation (**Figure 3.F**, lower) did not show significant variations. These results indicate that the compaction achieved by high ionic strength are also involved when the compaction is enhanced by calcium. A similar analysis employing the spectrum recorded at 0 mM IS as reference indicate more intense variations in the peak intensity values (**Fig. S5.B**, **C**), however this comparison cannot distinguish the contribution attributed to the ionic strength from the effect of calcium.

To determine the relative flexibility in the presence and absence of Ca^{2+} at 200 mM IS, we performed hetNOE (**Fig. S5.D**). The detected changes were observed at the initial part of the NAC region, where in the presence of Ca^{2+} several residues show less negative hetNOE values or zero. The C-terminal domain shows slight changes around the residue 110, and more clearly around the residue 120, where hetNOE value is higher. These results, in a similar way to the situation at higher ionic strength, confirm that both the NAC region and the C-terminal domain suffer a decrease in the flexibility, as consequence of the compaction of the monomer.

The cellular environment alters the conformational state of αSyn

It has been reported that different cellular environments and their components can promote the formation of specific polymorphs[12]. One interesting case is observed during the bacterial dysregulation of the microbiome in the gut, phenomenon that has been related to the development of PD. The release of microbial membrane metabolites, like Lipopolysaccharides (LPS), could represent a crucial mediator of αSyn amyloidogenesis via the gut-brain axis[36]. To investigate the effect of the bacterial cellular components in the conformational state of the monomeric αSyn, and fibril morphology, we designed an alternative purification protocol in which some bacterial elements remained in the protein preparation, namely IEX-αSyn. Our standard purification strategy of αSyn (see Materials and Methods section) includes SEC and reverse phase chromatography after ion exchange chromatography

(IEX) as final steps to completely remove molecules with similar charge than αSyn. From this point, we will call this protein HPLC-αSyn. However, IEX-αSyn preparation lacks the final steps of purification in order to keep in the protein mixture some components of the bacterial environment. Then, we analyzed and compared the chromatograms of both standard αSyn (HPLC-αSyn) and IEX-αSyn by analytical SEC (**Figure 4.A**) and analytical reverse phase chromatography (**Figure 4.B**), showing the presence of a component with small size (dashed circle) but constituted by elements with different hydrophobicity (black arrows). The peaks detected by reverse phase chromatography were grouped by retention time, those with a shorter and those with a longer retention time than the peak of αSyn, analyzed by electrospray ionization mass spectrometry (ESI-MS) and deconvoluted to estimate the mass of the molecular ion (**Fig. S6.A**). We found, for both groups, that the molecular weight of the bacterial components was ~12-14 kDa. Furthermore, employing the low molecular weight fraction marked in **Figure 4.A**, we recorded its FTIR spectrum (**Fig. S6.B**), finding two bands between 1700 and 1500 cm^{-1}. The molecular weight determined by ESI-MS is in good agreement with the value expected for bacterial endotoxins, that range between 10 and 20 kDa[37]. The bands detected by FTIR could correspond to the Amide I and Amide II of LPS[38], while the heterogeneity in the number of carbohydrate moieties in LPS could explain the different hydrophobicity peaks observed by analytical reverse phase chromatography. Finally, to corroborate the presence of LPS in IEX-αSyn, we employed a Toxin Sensor Kit (GenScript), finding an LPS-reactive result (**Fig. S6.C**). Taken together, these results clearly show that most of the bacterial elements in IEX-αSyn corresponds to LPS.

In order to understand the effect of the bacterial elements in the fibril morphology, we monitored the formation of IEX-αSyn fibrils by ThT assay in 50 mM TRIS pH 7.4 supplemented with NaCl ranging from 0 to 500 mM and compared with the morphology profile obtained in the **Figure 1.D** for the HPLC-αSyn (**Figure 4.C**). The results indicate that the bacterial components induce a ThT inflection point at a lower ionic strength (~150 mM) than the observed for HPLC-αSyn (~300 mM). The morphological consequences become evident at 200 mM NaCl, conditions in which IEX-αSyn forms a fibril with higher ThT that exhibits a typical rod morphology (**Figure 4.C**, red square), while pure αSyn produces a fibril with lower ThT and twisted polymorph (**Figure 4.C**, gray square). Since the LPS molecule was identified as the main bacterial component in IEX-αSyn, we prepared an artificial IEX-αSyn by adding LPS to HPLC-αSyn and tested the fibril morphology phase diagram by ThT assay (**Figure 4.D**), finding a ThT inflection point of ~150 mM, in good agreement with the value determined for IEX-αSyn. The morphological differences produced by the change in the inflection point of morphologies were also corroborated by TEM at 300 mM NaCl, ionic strength in which LPS added αSyn shows a rod morphology and the HPLC-αSyn a twisted fibril (**Figure 4.D**, purple and black squares, respectively).

Next, we performed ^{1}H-^{15}N HSQC experiments to determine the residues affected by the presence of bacterial endotoxins in IEX-αSyn and compared with the spectrum of HPLC-αSyn, at a concentration of NaCl of 200 mM, in which the differences between IEX-αSyn and HPLC-αSyn are clearer (**Figure 4.E**). The peak intensities difference ($I_{IEX} - I_{HPLC}$) shows a slight variation in all the three domains of αSyn, while the ratio (I_{IEX} / I_{HPLC}) indicates that the most intense changes are located in the N-terminus domain and the NAC region (**Figure 4.F**), while the C-terminus domain exhibits decreases in the values of ratio at the initial part of the C-terminus region and around the residue 130. The chemical shift perturbation shows no significant differences. Interestingly, using the artificial IEX-αSyn (HPLC-αSyn + LPS), its ^{1}H-^{15}N HSQC spectrum is very similar to that obtained for IEX-αSyn (**Figure 4.G**). The similarities are clearer if we consider the peak intensities difference ($I_{HPLC+LPS} - I_{HPLC}$) and ratio ($I_{HPLC+LPS} / I_{HPLC}$), where the regions mostly affected by the presence of LPS are almost identical to those observed in IEX-αSyn (**Figure 4.H**). No significant differences were seen by chemical shift perturbation. These results suggest that the N-terminus domain and some residues in the NAC region are responsible for the interaction with the bacterial endotoxins. However, in order to determine if there is induced by bacterial endotoxins, we established an algorithm of peak intensities based on the compact state observed at 300 mM NaCl (**Fig. S7.A**). First, we defined the total variation in peak intensity (v_{total}) of a residue (r) by subtracting from the highest peak intensity, found at 0 mM NaCl (I^{0mM}), the lowest peak intensity, at 500 mM NaCl (I^{500mM}):

$$v_{total}(r) = I^{0mM}(r) - I^{500mM}(r)$$

We also defined the variation of a peak intensity, at a certain ionic strength ([I.S.]), affected by the X agent, like Ca^{2+} or LPS, as:

$$v_X^{[I.S.]}(r) = I^{0mM}(r) - I_X^{[I.S.]}(r)$$

Then, we normalized the variation of a peak intensity using the total variation of peak intensity:

$$c_X^{[I.S.]}(r) = \frac{v_X^{[I.S.]}(r)}{v_{total}(r)}$$

Since above 300 mM NaCl we detected a plateau phase of compaction by SEC, as shown in **Figure 1.G**, we defined the threshold of compaction for the intensity of a peak as its $c(r)$ value at 300 mM NaCl:

$$c^{300mM}(r) = \frac{v^{300mM}(r)}{v_{total}(r)}$$

Thus, those residues in the condition X that can induce a normalized variation in their peak intensities higher than their $c^{300mM}(r)$, have been considered as relevant for compaction (marked as the red area in **Fig. S7.A**):

$$\frac{c_X^{[I.S.]}(r)}{c^{300mM}(r)} \geq 1.0$$

Additional constraints were introduced. For instance, those residues whose v_{total} is lower than 0.1 and

their normalized ratio of intensities ($I^{[I.S.]}_X / I^{300mM}$) is higher than 0.75 have been omitted to avoid

false positives.

Using the algorithm described above, we performed a global analysis of the peak intensities at

an ionic strength equivalent to 200 mM NaCl in the presence of bacterial endotoxins (IEX-αSyn), LPS

(HPLC-αSyn + LPS), and Ca^{2+} (**Fig. S7.C**). This analysis allowed us to recognize common residues

affected under the tested conditions. For instance, while the N-terminus domain showed as common

disturbed residue just the A11, the NAC region exhibits clearer variations at its initial part, specifically

the residues Q62 and N65. The most affected region for all the conditions was the C-terminus domain,

in which several residues between G106 and E139 surpassed the threshold established by $c^{300mM}(r)$

(**Fig. S7.B**), suggesting that the bacterial endotoxins induce a similar conformational compaction when

interacting with αSyn than the observed at high ionic strength or in the presence of calcium. An

alignment comparing the most relevant changes detected by the defined algorithm applied to all the

conditions of high NaCl, Ca^{2+} at 200 mM I.S., bacterial endotoxins in IEX-αSyn at 200 mM I.S., and

LPS-added to HPLC-αSyn at 200 mM I.S., is provided in **Figure 4.I.** The global analysis shows two

main affected regions for all the cases: the initial part of the NAC region and a wide area of the C-

terminal domain. Considering these results, it is clear that all the tested conditions share a common

molecular mechanism, in which there is a compaction enhanced by the interaction between the initial

part of the NAC region and the C-terminus domain that precedes the formation of rod fibrils.

The C-terminus domain is essential for the monomeric compaction and fibril polymorphism.

We have identified the C-terminus domain as a key factor in the compaction of the monomeric

αSyn. If we examine the sequence of the C-terminus domain, we will find four aromatic residues: one

phenylalanine and three tyrosine residues. Among the tyrosine residues, two of them were directly

identified by the global algorithm employed to analyze the NMR data (Y133 and Y136), suggesting

that the chemical environment of the tyrosine residues could change upon the compaction of αSyn

(**Fig. S8.A**). Since the aromatic residues can be studied relatively easily with spectroscopic tools, we

employed near circular dichroism (CD) spectroscopy to evaluate the alterations in the tyrosine

environment in the presence of components that induce the compaction of monomers, like high

concentration of NaCl and calcium (**Fig. S8.B**). The spectra showed some differences in the region of

the tyrosine residues (265-285 nm), whose ellipticity values become less negative in conditions that

promote the compaction, while the phenylalanine residues region (255-265 nm) remained unaltered.

These results indicate that spectroscopic tools that target the aromatic residues could be implemented as first approaches to detect the compact conformation in αSyn.

Finally, since a wide section of the C-terminus domain has been recognized as crucial for the compaction of the monomeric αSyn, we examined a truncated form of αSyn that lacks this region, from E105 to A140, namely N103t-αSyn, (**Figure 5.A**). First, we monitored the fibril formation using N103t-αSyn at different concentrations of NaCl, ranging from 50 mM to 300 mM, finding that, independently from the ionic strength, the ThT values for all the fibrils were low (**Figure 5.B**). A TEM visualization of the fibrils showed that just twisted fibrils were detected, even at high concentrations of salt (**Figure 5.C**). Additionally, we evaluated if there were changes in retention volume by analytical SEC, however, the retention volume did not show any significant variation in the interval from 50 mM to 500 mM NaCl. These results indicate that the lack of the key region of the C-terminal domain is enough to prevent not just the formation of rod fibrils but also any variation in the conformational state of the monomeric N103t-αSyn.

Discussion

The direct visualization of fibril polymorphs extracted from the brain of DLB and MSA patients represent an unprecedent evidence for the hypothesis that different polymorphs are related to the development of different diseases[9]. *In vitro*, the methods to create a variety of αSyn fibril morphologies, altering conditions like the ionic strength, are well known[7,19,20]; however, the molecular mechanisms that generate those diverse polymorphs from the same intrinsically disordered protein remain poorly understood. In this study, employing physicochemical approaches, we have found that, at neutral pH, the local environment of αSyn can induce a conformational compaction of the monomeric protein, which is related to the specific formation of polymorphs. Our results of SEC and AUC indicate that if the ionic strength increases, the protein suffers a compaction that reaches a stable level above 300 mM NaCl, that corresponds with the region in the morphology phase diagram in which the formation of rod fibrils begins. The structural elements of the monomeric compaction, induced by NaCl, were revealed by NMR, in which variations in the initial part of the NAC region, specifically the residues Q62 and N65, and a wide area of the C-terminus domain suggests the existence of a polar interaction between these two regions. The identification of the C-terminus domain as key factor for the compaction is particularly remarkable since this region is the target of several modifications, like phosphorylation, cross-linking, ubiquitination, truncation, interaction with metals, etc.[39], events that could potentially affect the conformational state of αSyn and the consequent fibril polymorph.

Recently, it was reported, at low ionic strength, the interaction between αSyn and Ca^{2+}, a biologically relevant cation that also accelerates the speed of fibril formation[35]. The authors found that

the interaction between the negative residues of the C-terminal domain and Ca^{2+} promotes a disruption in the electrostatic contacts with the positively charged N-terminus domain, leading to a higher exposure of the N-terminus domain and the initial part of the NAC region. Our results confirm the interaction between the C-terminus domain and Ca^{2+} (**Figure 3**), and also provide a more general mechanism based on the monomeric compaction applicable not only to the reported low ionic strength[35], but also for mild (physiological) and high ionic strength. First, in the presence of 200 Eq. Ca^{2+}, we found that αSyn can form fibrils with both low and high ThT values when the ionic strength is low. However, above an ionic strength of 200 mM, just fibrils with rod morphology and high ThT were observed. Furthermore, at this ionic strength, we detected a compaction in the monomer (Fig 3.D) accompanied by structural modifications, detected by NMR, in the initial part of the NAC region and in a wide area of the C-terminus domain, similarly to those residues observed at high concentration of NaCl. Taken together, our results suggest that, independently of the ionic strength, Ca^{2+} establishes electrostatic interactions with the residues of the C-terminus domain, titrating the negative charges of residues like Glu and Asp. However, the Ca^{2+}-bound-C-terminus domain behaves differently when the ionic strength changes. At low ionic strength, the titrated C-terminus domain will get released from the transient electrostatic contacts with the N-terminus region, however, the interaction with the initial part of the NAC region still might occur sporadically; in consequence, the fibrils can acquire both the rod and the twisted morphology. When the ionic strength reaches 200 mM, the concentration of ions in solution induces a tendency in αSyn to protect its hydrophobic NAC region from the solvent. Since Ca^{2+} has partially neutralized some negative charges of the C-terminus domain, the interaction with the NAC region gets facilitated with the concomitant compaction of the monomer. In the absence of calcium, the negative charges of the C-terminus domain need a higher ionic strength, like 300 mM, to get neutralized and allow the interaction with the initial part of the NAC region. Thus, calcium enhances the compaction of the monomer and promotes the formation of rod fibrils at lower ionic strength values than NaCl, in the range of the physiological ionic strength. The compaction of monomers induced by calcium could trigger the fibril accumulation of rod fibrils during the dysregulation of intracellular calcium homeostasis, event that has been found to play an important role in the pathogenesis of PD[40].

The effect of some components present in the cellular environment on the compaction of monomers was evaluated employing bacterial endotoxins (LPS), that are molecules associated to the αSyn amyloidogenesis through the gut-brain axis[36]. The ThT assay evaluation of fibril morphologies indicated the formation of rod fibrils at lower ionic strength than salt. For example, at an ionic strength of 200 mM, pure αSyn shows twisted fibrils while IEX-αSyn just shows rod fibrils. This decrease in the ThT inflection point of fibril morphologies suggests that an enhanced compaction of monomers at lower ionic strength could take place in the presence of bacterial endotoxins. The NMR measurements

indicate that the binding between LPS and αSyn occurs involving the N-terminus domain and some hydrophobic residues of the NAC region. Interestingly,The driving force of the interaction could be a series of hydrogen bonds established between the positively charged N-terminus domain and the hydroxyl groups in the carbohydrate moieties of LPS, while hydrophobic contacts might take place between the lipidic aliphatic chains of LPS and the hydrophobic NAC region. However, our algorithm could recognize affected regions in the initial part of the NAC region and a wide area of the C-terminus domain, that are in good agreement with the residues identified at high ionic strength and in the presence of Ca^{2+}, and implying that the phenomenon of compaction also happens in the presence of endotoxins. These results suggest a mechanism of compaction: when LPS binds the N-terminus domain and some residues of the NAC region, the C-terminus domain gets released from the electrostatic interaction with the N-terminus domain and, since the hydrophobic residues of the NAC region are compromised with the aliphatic region of LPS, the polar interaction between the residues Q62 and N65 with the C-terminus domain gets facilitated, leading to the formation of a compact monomer bound to LPS, which is also able to produce rod fibrils at lower ionic strength. These results have a critical importance since they provide a new mechanism of fibril formation when the microbiota in the gut is dysregulated, and the bacterial endotoxins concentration increases. In addition, our findings also suggest that the binding of αSyn to other cellular components through the N-terminus domain, like the cellular membranes, could induce not only the acceleration of fibril aggregation[41], but also might expose the NAC region to the interaction with the C-terminus domain and lead to the monomeric compaction and the subsequent formation of specific polymorphs.

The conditions examined here (differences in ionic strength, the presence of calcium, the effect of bacterial endotoxins) modified the phase diagram of fibrils by affecting the monomeric conformation of αSyn and the interaction between the C-terminus domain with the initial part of the NAC region. In physiological conditions of pH and ionic strength, our morphological phase diagram indicates that the preferred fibril would be the twisted polymorph; however, the dysregulation of metabolites like LPS and calcium, or, hypothetically, lipids like glucosylceramide[42], might alter this phase diagram and enable the aggregation of rod fibrils, like those observed in PD. The model of the monomeric compaction and the phase diagram of fibrils could be expanded to the complexity of the cellular environments. For instance, Peng et al[12] reported that the αSyn aggregates in the neuronal cells can produce PD-like aggregates, whereas the αSyn aggregated in the oligodendrocyte resembles the MSA-like ones. Further investigation is required to determine the fibril phase diagrams in the neuronal cells and in the oligodendrocytic cells and the factors that trigger the aggregation in these environments. This knowledge would provide insights to better understand the pathogenesis of each alpha synucleinopathies.

Finally, although our model of compaction allows the prediction of fibril morphologies, the
direct link explaining how the monomeric compaction and fibril morphology are related remains
unsolved. Considering our results and the cryo-EM polymorphs structures already reported[43,44], we
propose a mechanism for the formation of both twisted and rod fibrils from the scenario of the
monomeric compaction: 1) (**Figure 6**, left) at low ionic strength, the monomer adopts an ensemble of
extended conformations with structural fluctuations. During the fibril formation, the structural
monomeric plasticity allows the protein to acquire the most stable morphology: the twisted fibril. The
driving force of the twisted fibril would be the protection from the solvent of the hydrophobic residues
of the NAC region at the interface of the constituting protofilaments. 2) (**Figure 6**, right) Under
conditions that promote the compaction of monomers, like high ionic strength, calcium or LPS, the C-
terminus domain that interacts with the initial part of the NAC region would protect the hydrophobic
residues of this domain, preventing the hydrophobic contact between the protofilament interfaces
required for the twisted fibril. Instead, an interface characterized by the electrostatic interaction
between H50 and E57 and a hydrophobic zipper from G51 to A56 would be stabilized, leading to the
formation of the rod fibril.

493
494   **Materials and Methods**
495
All chemicals and reactants were obtained from Sigma-Aldrich or Nacalai Tesque (Kyoto, Japan). All
reagents were used without further purification. Water was purified to a resistivity of 18 MΩ cm^{-1}
using a Millipore Gradient deionizing system.

<u>Protein purification</u>

Human WT αSyn was purified from *Escherichia coli* as described previously (*34*). Briefly, a plasmid
containing WT human αSyn was expressed in *E. coli* BL21 (DE3) cells (Novagen, Merck, San Diego,
CA, USA). The cells were suspended in purification buffer, disrupted by sonication, and centrifuged.
Streptomycin sulfate (final 2.5% w/w) was added to the supernatant, and centrifugation was repeated.
The supernatant was heated to 85 °C in a water bath and centrifuged. The supernatant was precipitated
by the addition of solid ammonium sulfate to 70% saturation, centrifuged, dialyzed overnight, applied
to a Resource-Q column (GE Healthcare, Little Chalfont, UK) with 50 mM TRIS-HCl pH7.4 buffer
containing 0.1 mM dithiothreitol and 0.1 mM phenylmethylsulfonyl fluoride as a running buffer, and
eluted with a linear gradient of 0–1 M NaCl. αSyn-enriched fractions (as determined by sodium
dodecyl sulfate-polyacrylamide gel electrophoresis SDS-PAGE/Coomassie blue staining) were pooled
and further purified by size exclusion chromatography (SEC) using a Superdex 200 26/600 PG column
(GE Healthcare, Little Chalfont, UK) equilibrated with 50 mM TRIS-HCl pH 7.4 supplemented with
150 mM NaCl. The fractions containing αSyn (as determined by SDS-PAGE/Coomasie blue staining)
were joint, dialyzed versus deionized water, acidified with 5 mM HCl and loaded onto a Reverse Phase

Cosmosil Protein R x 250 mm Preparative Column (Nacalai-Tesque, Kyoto, Japan) and eluted with a linear gradient of 30 % - 90 % acetonitrile. The pure fractions were combined and flash-frozen in liquid nitrogen, lyophilized and stored at -80 ℃ until use. The protein purity was confirmed to be greater than 95% by SDS-PAGE and matrix-assisted laser desorption/ionization mass spectrometry. The protein concentration was determined by UV absorption at 280 nm using an extinction coefficient $\varepsilon_{1\%}$ of 3.54 [45].

C-terminal truncated N103t αSyn: The C-terminal truncated αSyn construct was prepared using a QuikChange site-directed mutagenesis kit (Stratagene, Santa Clara, California), employing the WT-αSyn plasmid of the previous section as a template. Protein expression, cell lysis and αSyn precipitation were performed as described for the WT full-length αSyn. The protein solution was solved using a cationic Resource-S column (Amersham Biosciences) and eluted with a linear gradient of 1 M NaCl. Size exclusion chromatography and reverse phase chromatography purification steps were performed as described above. The fractions containing the isolated protein were lyophilized and stored as indicated for full-length αSyn. The protein concentration was determined by UV absorption at 280 nm using an extinction coefficient $\varepsilon_{1\%}$ of 1.45, calculated using the ExPASY tool [46].

<u>Kinetics of fibril formation followed by thioflavin T (ThT)</u>

The formation of αSyn amyloid fibrils was studied by preparing solutions of α Syn (0.5 mg mL^{-1}) in fibrillation buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl) containing 10 μM ThT in a total volume of 200 μL, which were placed in each well of a 96-well microplate, at 37 °C. The ThT fluorescence intensity was monitored using a HANdai Amyloid Burst Inducer (HANABI-2000) equipment, developed by our group in collaboration with CORONA ELECTRIC, Ibaraki, Japan, consisting of a transducer driving system, an acoustic-intensity measurement system, and a fluorescence measure system, in which a microplate reader was combined with a multichannel ultrasonication system that applies ultrasonic irradiation independently to each individual well of the microplate via a miniaturized ultrasonic resonator attached to each well. Ultrasonication was applied to accelerate amyloid formation at an optimized frequency of 30 kHz[31,47] in cycles of 300 ms of irradiation and 500 ms of quiescence. The kinetic parameters of fibril formation was determined by fitting the experimental data to empirical equations[48].

<u>Patient information and tissue preparation.</u>

Brain tissues from three MSA patients (67.7 ± 8.5 years old, M: F = 3: 0) and three PD patients (79.3 ± 2.8 years old, M: F = 2: 1) were obtained from the autopsy series of the Brain Bank for Aging Research (BBAR) in the Tokyo Metropolitan Geriatric Hospital and Institute of Gerontology. The brains were processed according to the BBAR protocol. Briefly, half of the brain was sliced and frozen,

and the other half was fixed in 20% buffered formalin for 7-13 days, after which the representative anatomical areas were embedded in paraffin.

Brain lysate preparation

For this study, amygdala sections from the frozen side and amygdala sections from the formalin-fixed side were prepared. Amygdala slices of 100 mg from the brains of PD and MSA patients were placed into Precellys Lysing tubes (M&S Instruments, Osaka, Japan), resuspended in 1 mL of fibrillation buffer, and subjected to two cycles of high-speed shaking for 20 s in a Lysis and Homogenization System (Bertin Instruments, France). Then, the homogenates were transferred into Eppendorf tubes and centrifuged at 2000 x g for 2 min at room temperature. The concentration of total protein in the supernatant fractions was quantified using a MicroBCA Protein Assay Reagent Kit (Pierce), and the homogenates were aliquoted and stored at -80 °C until use.

Immunohistochemistry of brain samples.

Immunohistochemistry was performed using a mouse monoclonal antibody against phosphorylated α-synuclein (pSyn#64; a gift from T. Iwatsubo, Japan), a Ventana BenchMark GX autostainer (Ventana Medical Systems, Tucson, AZ, USA), and an I-View Universal DAB Detection Kit (Roche, Basel, Switzerland). All slides were counterstained with haematoxylin to visualize neurons and other cells.

Amplification of αSyn aggregates from amygdala brain homogenates

A volume of amygdala brain homogenate was added to solutions of αSyn to reach a final concentration of 20 µg mL^{-1} total protein. Two hundred microliters of 0.5 mg mL^{-1} monomeric αSyn in the fibrillation buffer containing the amygdala lysate was placed into a 96-multiplate and subjected to ultrasonic irradiation to accelerate the formation of amyloid fibrils, as described above. The ThT fluorescence intensity was recorded as a function of time.

Transmission electron microscopy (TEM)

TEM imaging was achieved on a Hitachi H-7650 transmission electron microscope (Hitachi, Tokyo) operated at 80 kV. Samples were diluted 1:10 with deionized water, and 10 µL of this solution was placed onto copper grids (400-mesh) covered with carbon-coated collodion film (Nisshin EM, Tokyo) and incubated for 1 min at room temperature. The samples were negatively stained with 10 µL of a 1% (w/v) solution of phosphotungstic acid (PTA), incubated for 1 min, and finally washed again with 10 µL of deionized water. The magnification working interval ranged from 5 000× to 30 000×.

Proteinase K resistance (PKR) assay

αSyn fibrils (0.5 mg mL^{-1}) in the fibrillation buffer were digested using proteinase K (1 µg mL^{-1}, Roche) at 37 °C and 400 rpm for different time intervals. To stop the reaction, the samples were incubated for 5 min at 95 °C, mixed with loading buffer (50 mM TRIS-HCl, pH 6.8, 4% SDS, 2% β-

mercaptoethanol, 12% glycerol and 0.01% bromophenol blue) and incubated at 95 °C for additional 10 min. The digestion patterns were analyzed by SDS-PAGE followed by Coomassie Brilliant Blue staining. The first five products of digestion, namely B1 to B5, were employed for the analysis. The PKR Score has been established as the band intensity ratio between the bands B2 and B1 (B2/B1).

Analytical size-exclusion chromatography (SEC)

αSyn solutions were filtered through a 0.22 µm size-pore membrane and loaded onto a Superdex 200 10/300 GL SEC column (GE Healthcare) coupled to an AKTA Explorer HPLC Instrument (Amersham Biosciences). The elution buffer was 50 mM TRIS-HCl (pH 7.4) supplemented with 150 mM NaCl. The flow rate was set to 0.5 mL min^{-1}. The protein absorption at 220, 250 and 280 nm was monitored.

Analytical reverse phase chromatography

αSyn solutions in 5 mM HCl were filtered through a 0.22 µm size-pore membrane and loaded onto a Cosmosil 5C$_4$-AR-300 4.6ID x 150 mm analytical column (Nacalai-Tesque, Kyoto, Japan) coupled to a Gilson HPLC Unipoint System Instrument (Gilson, USA) with a 30 % v/v acetonitrile in water buffer supplemented with 0.05 % v/v trifluoroacetic acid. The sample was eluted using a 30-90 % v/v gradient of acetonitrile. The flow rate was set to 0.5 mL min^{-1}. The protein absorption at 220 and 280 nm was monitored.

Analytical ultracentrifugation (AUC)

Sedimentation velocity (SV) experiments were performed at 20 °C using an Optima AUC (Beckman Coulter) analytical ultracentrifuge (Fullerton, CA, USA) equipped with an An-50Ti rotor using 12-mm double-sector charcoal-filled epon centerpiece with sapphire windows. The protein concentrations were adjusted to absorbance values of 0.8 in the working buffer at 1-cm path length. Sedimentation data were collected every 2 min at 40,000 rpm with a radial increment of 0.001 cm using absorbance optics. The detection wavelength was set at 230 nm.

The distribution of the sedimentation coefficient was determined using the continuous c(s) distribution model in the program SEDFIT[49]. The range of the sedimentation coefficients for fitting was 0–15 S, with a resolution of 300. The partial specific volume of the sample, the buffer density, and the buffer viscosity were calculated by the program SEDNTERP[50], respectively.

Multidimensional NMR

^{1}H-^{15}N heterogenous single-quantum coherence (HSQC) NMR measurements were performed by using 100 µM ^{15}N-labelled αSyn dissolved in fibrillation buffer prepared in H$_2$O/D$_2$O (9:1, v/v). The ^{15}N-labelled αSyn was expressed in M9 minimal media containing ^{15}NH$_4$Cl and purified as described for the unlabeled protein. NMR spectra were acquired at 310 K on a Bruker Advance III 600 MHz NMR spectrometer equipped with a QCI-P cryogenic probe (Bruker Biospin). The peak assignment at

pH 7.4 was achieved employing the assignment data reported by El Turk *et al*[51]. Chemical shift perturbation (CSP) was calculated as follows:

$$\Delta\delta = \sqrt{\Delta\delta_H^2 + \left(\frac{1}{8}\Delta\delta_N\right)^2}$$

where $\Delta\delta_H$ and $\Delta\delta_N$ are the chemical shift changes (in ppm) with respect to the H and N axes, respectively.

For backbone dynamics study, ^{1}H-^{15}N hetero NOE of ^{15}N αSyn were obtained by measured ^{1}H-^{15}N HSQC spectra with and without ^{1}H saturation for 3 seconds, and then calculating the ratios of the intensities of the ^{1}H-^{15}N peaks. The errors in hetero NOE values were estimated using root-mean-square value of the spectrum background noise.

All NMR spectra were processed with Topspin (Bruker Biospin), NMRPipe[52] and NMRFAM-sparky[53].

Structural Model Selection

576 disordered conformations of αSyn from the Protein Ensemble Database (https://proteinensemble.org/) (PED ID: PED00024 [34]) were used to measure deviations between the chemical shifts predicted based on conformations and the shifts observed in this study. Predicted shifts and root-mean-square (RMS) deviations between the predicted and experimental shifts were calculated using SPARTA+[54]. The ^{1}H- and ^{15}N- chemical shifts of αSyn in 0, 150 and 500 mM NaCl were used for the calculations.

Circular dichroism (CD) spectroscopy

CD measurements were performed using a Jasco J820 spectrophotometer at 37 °C. Quartz cuvettes of 1 cm were utilized. The employed protein concentration was 5.0 mg mL^{-1} in fibrillation buffer. Spectra were recorded in 350-250 nm (near-UV CD) ranges and reported as the mean residue ellipticity ($[\theta]_{MRW}$, deg cm^2 dmol^{-1}) after subtracting the baseline.

Statistical analysis

We used Statistical Package for the Social Sciences 23.0J software (IBM Japan, Tokyo, Japan) to perform statistical analysis. For the statistical analysis of 2 groups, the paired or unpaired t-test was used as described in the figure legends. For more than 3 groups, we used one-way analysis of variance (ANOVA) with the Bonferroni/Dunn post hoc test. In all experiments, data were expressed as the mean ± SD, and a *p* value of < 0.05 was considered to indicate a statistically significant difference between 2 groups (* *p* < 0.05. ** *p* < 0.001, *** *p* < 0.0001).

Ethical Statement

The collection of autopsied human tissues and their use for this study were approved by the Ethics Committee of Osaka University Graduate School of Medicine, and written informed consent was obtained from the patients' next of kin. Experimental procedures involving human subjects were conducted in conformance with the principles expressed in the Declaration of Helsinki.

References

1. Spillantini, M. G. *et al.* Alpha-synuclein in Lewy bodies. *Nature* **388**, 839–840 (1997).

2. Spillantini, M. G., Crowther, R. A., Jakes, R., Hasegawa, M. & Goedert, M. α-Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and dementia with Lewy bodies. *Proc. Natl. Acad. Sci. U. S. A.* (1998). doi:10.1073/pnas.95.11.6469

3. Gai, W. P., Power, J. H. T., Blumbergs, P. C. & Blessing, W. W. Multiple-system atrophy: A new α-synuclein disease? *Lancet* (1998). doi:10.1016/S0140-6736(05)79256-4

4. Araki, K. *et al.* Synchrotron FTIR micro-spectroscopy for structural analysis of Lewy bodies in the brain of Parkinson's disease patients. *Sci. Rep.* **5**, 1–8 (2015).

5. Araki, K. *et al.* Parkinson's disease is a type of amyloidosis featuring accumulation of amyloid fibrils of α-synuclein. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 17963–17969 (2019).

6. Araki, K. *et al.* The secondary structural difference between Lewy body and glial cytoplasmic inclusion in autopsy brain with synchrotron FTIR micro-spectroscopy. *Sci. Rep.* **10**, 1–6 (2020).

7. Van der Perren, A. *et al.* The structural differences between patient-derived α-synuclein strains dictate characteristics of Parkinson's disease, multiple system atrophy and dementia with Lewy bodies. *Acta Neuropathol.* (2020). doi:10.1007/s00401-020-02157-3

8. Shahnawaz, M. *et al.* Discriminating α-synuclein strains in Parkinson's disease and multiple system atrophy. *Nature* **578**, 273–277 (2020).

9. Schweighauser, M. *et al.* Structures of α-synuclein filaments from multiple system atrophy. *Nature* **585**, 464–469 (2020).

10. Prusiner, S. B. *et al.* Evidence for α-synuclein prions causing multiple system atrophy in humans with parkinsonism. *Proc. Natl. Acad. Sci. U. S. A.* (2015). doi:10.1073/pnas.1514475112

11. Yamasaki, T. R. *et al.* Parkinson's disease and multiple system atrophy have distinct -synuclein seed characteristics. *J. Biol. Chem.* (2019). doi:10.1074/jbc.RA118.004471

12. Peng, C. *et al.* Cellular milieu imparts distinct pathological α-synuclein strains in α-synucleinopathies. *Nature* 1 (2018). doi:10.1038/s41586-018-0104-4

13. Fusco, G. *et al.* Structural basis of membrane disruption and cellular toxicity by α-synuclein oligomers. *Science (80-.).* **358**, 1440–1443 (2017).

14. Suzuki, G. *et al.* α-synuclein strains that cause distinct pathologies differentially inhibit proteasome. *Elife* (2020). doi:10.7554/eLife.56825

15. Hayakawa, H. *et al.* Structurally distinct α-synuclein fibrils induce robust parkinsonian pathology. *Mov. Disord.* 1–12 (2019). doi:10.1002/mds.27887

16. Peelaerts, W., Bousset, L., Baekelandt, V. & Melki, R. α-Synuclein strains and seeding in Parkinson's disease, incidental Lewy body disease, dementia with Lewy bodies and multiple system atrophy: similarities and differences. *Cell Tissue Res.* **373**, 195–212 (2018).

17. Ikenaka, K., Suzuki, M., Mochizuki, H. & Nagai, Y. Lipids as Trans-Acting Effectors for α-

Synuclein in the Pathogenesis of Parkinson's Disease. *Front. Neurosci.* **13**, 1–9 (2019).

18. Brundin, P. & Melki, R. Prying into the Prion Hypothesis for Parkinson's Disease. *J.*
Neurosci. **37**, 9808–9818 (2017).

19. Bousset, L. *et al.* Structural and functional characterization of two alpha-synuclein strains.
Nat. Commun. **4**, 2575 (2013).

20. Peelaerts, W. *et al.* α-Synuclein strains cause distinct synucleinopathies after local and
systemic administration. *Nature* **522**, 340–344 (2015).

21. Bertoncini, C. W. *et al.* Release of long-range tertiary interactions potentiates aggregation of
natively unstructured α-synuclein. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 1430–1435 (2005).

22. Kessler, J. C., Rochet, J. C. & Lansbury, P. T. The N-terminal repeat domain of α-synuclein
inhibits β-sheet and amyloid fibril formation. *Biochemistry* **42**, 672–678 (2003).

23. Zhou, W. *et al.* Methionine oxidation stabilizes non-toxic oligomers of α-synuclein through
strengthening the auto-inhibitory intra-molecular long-range interactions. *Biochim. Biophys.*
Acta - Mol. Basis Dis. **1802**, 322–330 (2010).

24. Izawa, Y. *et al.* Role of C-terminal negative charges and tyrosine residues in fibril formation
of ??-synuclein. *Brain Behav.* **2**, 595–605 (2012).

25. Ranjan, P. & Kumar, A. Perturbation in Long-Range Contacts Modulates the Kinetics of
Amyloid Formation in α-Synuclein Familial Mutants. *ACS Chem. Neurosci.* **8**, 2235–2246
(2017).

26. Furukawa, K. *et al.* Isoelectric point-amyloid formation of α-synuclein extends the generality
of the solubility and supersaturation-limited mechanism. *Curr. Res. Struct. Biol.* **2**, 35–44
(2020).

27. Morar, A. S., Olteanu, A., Young, G. B. & Pielak, G. J. Solvent-induced collapse of α-
synuclein and acid-denatured cytochrome c. *Protein Sci.* **10**, 2195–2199 (2008).

28. Edwards, A. J. & Reid, D. Introduction to NMR of Proteins. *Curr. Protoc. Protein Sci.* **19**,
Unit 17.5 (2000).

29. Ulrih, N. P., Barry, C. H. & Fink, A. L. Impact of Tyr to Ala mutations on α-synuclein
fibrillation and structural properties. *Biochim. Biophys. Acta - Mol. Basis Dis.* **1782**, 581–585
(2008).

30. Kakuda, K. *et al.* Ultrasonication-based rapid amplification of α-synuclein aggregates in
cerebrospinal fluid. *Sci. Rep.* **9**, 6001 (2019).

31. Nakajima, K. *et al.* Optimized sonoreactor for accelerative amyloid-fibril assays through
enhancement of primary nucleation and fragmentation. *Ultrason. Sonochem.* **73**, 105508
(2021).

32. Atarashi, R. *et al.* Ultrasensitive human prion detection in cerebrospinal fluid by real-time
quaking-induced conversion. *Nat. Med.* **17**, 175–178 (2011).

33. Nakagaki, T., Nishida, N. & Satoh, K. Development of α-Synuclein Real-Time Quaking-
Induced Conversion as a Diagnostic Method for α-Synucleinopathies. *Front. Aging Neurosci.*
0, 637 (2021).

34. Allison, J. R., Varnai, P., Dobson, C. M. & Vendruscolo, M. Determination of the free energy
landscape of α-synuclein using spin label nuclear magnetic resonance measurements. *J. Am.*
Chem. Soc. (2009). doi:10.1021/ja904716h

35. Stephens, A. D. *et al.* Extent of N-terminus exposure of monomeric alpha-synuclein
determines its aggregation propensity. *Nat. Commun.* **11**, (2020).

36. Bhattacharyya, D. & Bhunia, A. Gut-Brain axis in Parkinson's disease etiology: The role of

lipopolysaccharide. *Chem. Phys. Lipids* **235**, 105029 (2021).

37.   Raetz, C. R. H. BIOCHEMISTRY OF ENDOTOXINS. *Annu. Rev. Biochem.* **59**, 129–170
(1990).

38.   Barkleit, A. *et al.* Coordination of uranium(vi) with functional groups of bacterial
lipopolysaccharide studied by EXAFS and FT-IR spectroscopy. *Dalt. Trans.* **40**, 9868–9876
(2011).

39.   Oueslati, A. Implication of Alpha-Synuclein Phosphorylation at S129 in Synucleinopathies:
What Have We Learned in the Last Decade? *J. Parkinsons. Dis.* **6**, 39–51 (2016).

40.   Schapira, A. H. V. Calcium dysregulation in Parkinson's disease. *Brain* **136**, 2015–2016
(2013).

41.   Doherty, C. P. A. *et al.* A short motif in the N-terminal region of α-synuclein is critical for
both aggregation and function. *Nat. Struct. Mol. Biol.* **27**, 249–259 (2020).

42.   Mazzulli, J. R. *et al.* Gaucher disease glucocerebrosidase and α-synuclein form a bidirectional
pathogenic loop in synucleinopathies. *Cell* (2011). doi:10.1016/j.cell.2011.06.001

43.   Li, B. *et al.* Cryo-EM of full-length α-synuclein reveals fibril polymorphs with a common
structural kernel. *Nat. Commun.* **9**, 3609 (2018).

44.   Guerrero-Ferreira, R. *et al.* Cryo-EM structure of alpha-synuclein fibrils. *Elife* **7**, 1–18 (2018).

45.   Narhi, L. *et al.* Both familial Parkinson's disease mutations accelerate alpha-synuclein
aggregation. *J. Biol. Chem.* **274**, 9843–9846 (1999).

46.   Gasteiger, E. *et al.* ExPASy: The proteomics server for in-depth protein knowledge and
analysis. *Nucleic Acids Res.* **31**, 3784–3788 (2003).

47.   Nakajima, K. *et al.* Drastic acceleration of fibrillation of insulin by transient cavitation bubble.
Ultrason. Sonochem. **36**, 206–211 (2017).

48.   Arosio, P., Knowles, T. P. J. & Linse, S. On the lag phase in amyloid fibril formation. *Phys.*
Chem. Chem. Phys. **17**, 7606–7618 (2015).

49.   Schuck, P. Size-distribution analysis of macromolecules by sedimentation velocity
ultracentrifugation and Lamm equation modeling. *Biophys. J.* **78**, 1606–1619 (2000).

50.   Laue & M., T. Computer-aided interpretation of analytical sedimentation data for proteins. in
Analytical ultracentrifugation in biochemistry and polymer science. 90–125 (Royal Society of
Chemistry, 1992).

51.   El Turk, F. *et al.* Exploring the role of post-translational modifications in regulating α-
synuclein interactions by studying the effects of phosphorylation on nanobody binding.
Protein Sci. **27**, 1262–1274 (2018).

52.   Delaglio, F. *et al.* NMRPipe: A multidimensional spectral processing system based on UNIX
pipes. *J. Biomol. NMR* **6**, 277–293 (1995).

53.   Lee, W., Tonelli, M. & Markley, J. L. NMRFAM-SPARKY: Enhanced software for
biomolecular NMR spectroscopy. *Bioinformatics* **31**, 1325–1327 (2015).

54.   Shen, Y. & Bax, A. SPARTA+: A modest improvement in empirical NMR chemical shift
prediction by means of an artificial neural network. *J. Biomol. NMR* (2010).
doi:10.1007/s10858-010-9433-9

Acknowledgments

We would like to thank Shizuka Sonoda and Miki Yoshida for the protein expression and purification and the excellent technical support received. CA thanks JSPS (postdoctoral fellowship P16388) for the financial support.

Funding: This work was supported by JSPS KAKENHI Grant JP16H06277, JP18H02741; JST CREST Grant Numbers JPMJCR18H4 and JPMJCR17H6; JSPS Core-to-Core Program A Advance Research Networks; AMED Grant Numbers JP18dm0107103, JP19dm0207001, JP20dm0107106, JP20dm0107158, JP20km0405206, JP20am0101110 and SENTAN16809202; and the Ministry of Education, Culture, Sports, Science and Technology Grant Number 17H06352. The Brain Mapping by Integrated Neuroethologies for Disease Studies (Brain/MINDS) from Japan Agency for Medical Research and development, AMED JP18dm0207020. CA was funded by JSPS KAKENHI Grant 16F16388.

Author contributions: CA, KI, YG, and HM designed the experiments. CA, KI, MS, TM, KY, KN, CJC, KN, GB, KT, YY, JD, TM, MM, KK, MH, YK, SM and YM performed the experiments. CA and KI analyzed the data. CA and KI wrote the original draft. MS, YK, SN, KB, HO, YN, YK, SU, YM, YG and HM reviewed the manuscript.

Competing interests: Authors declare no conflicts of interest.

Data and materials availability: All data is available in the main text or the supplementary materials.

Dedications: We would like to express our deep gratitude to Professor Kazuhiro Ikenaka for his support, precious advices, and discussions, especially for the glycolipids section.

Figures and Tables

Figure 1. αSyn fibril morphology is modulated by the conformational state of the monomeric αSyn. (A) PD-(red) and MSA-(blue)-amygdala-seeded fibril amplification monitored by ThT intensity (mean ± s.e.m. of three different patients). **(B)** TEM visualization of fibrils amplified from the brain of patients. **(C)** ThT-monitored αSyn fibril formation at different concentrations of NaCl. **(D)** Maximal ThT intensity values of fibrils created under the indicated concentration of NaCl. Representative amyloid fibrils obtained at 50 mM (low ThT, blue square), 300 mM (low ThT, green square; high ThT, orange square) and 500 mM (red square) were visualized by TEM. (Bar scale: 200 nm). **(E)** From left to right, proteinase-K resistance assay of five replicates (numbers 1-5) of fibrils produced at 50 mM, 500 mM, created at 300 mM NaCl with maximal low- or high-ThT values, and amplified from PD and MSA brain of patients. Arrows indicate five selected bands subjected to intensity analysis. Bands marked with (*) denote the most relevant differences. Molecular weight markers (kDa) are shown on the left of each panel. **(F)** A solution of αSyn in 50 mM TRIS pH 7.4 supplemented with 300 mM NaCl was solved by SEC (upper) and the fractions obtained were analyzed by Western Blotting (Anti-Syn211, lower). **(G)** Representative chromatograms of Solutions of monomeric αSyn at different concentrations of NaCl solved by Size Exclusion Chromatography. **(H)** Analytical Ultracentrifugation Sedimentation Velocity measurements of solutions of monomeric αSyn in the presence of 150 mM (upper-left) and 500 mM (upper-right) NaCl. The sedimentation coefficient values ($s_{20,w}$) were determined to be 1.136 ± 0.012 S and 1.092 ± 0.011 S for the monomers at 150 mM and 500 mM NaCl, respectively (lower). The SV measurements were performed by triplicate.

Figure 2. Compact αSyn monomers establish interactions between the C-terminus
domain and NAC region. (A) ^{1}H-^{15}N HSQC spectra of solutions of αSyn supplemented with 63 mM
(blue) and 500 mM (red) NaCl. **(B)** Peak intensity difference analysis of the spectra of αSyn of the
amide N-H nuclei recorded at 63 mM, 300 mM, and 500 mM NaCl, employing the spectrum obtained
at 0 mM NaCl as a reference. Gray bars represent the residues without signal assignments. **(C)**
Residues that exhibit the most relevant changes in peak intensity, marked as red lines. The blue lines
represent residues with slight increases in peak intensity.

Figure 3. Ca²⁺ enhances the formation of αSyn rod-like fibrils through the compaction of
monomers. (A) Maximal ThT intensity values of fibrils created under the indicated ionic strength in
the presence (green) or absence (gray) of 200 Eq. Ca²⁺. (The values of maximal ThT in the absence of
Ca²⁺ are the same as those shown in **Figure 1.D**. They were inserted here for comparative purposes).
(B) TEM visualization of αSyn fibrils created at an ionic strength of 200 mM in the presence of 200
Eq. Ca²⁺ (Bar scale: 200 nm). **(C)** PKR assay of representative fibrils obtained in the presence of 200
Eq. Ca²⁺ at an ionic strength of 200 mM. Arrows indicate five selected bands subjected to intensity
analysis. Bands marked with (*) denote the most relevant differences. Molecular weight markers (kDa)
are shown on the left of each panel. **(D)** Representative chromatograms of solutions of monomeric
αSyn in the presence or absence of 200 Eq. Ca²⁺, at an ionic strength of 200 mM, solved by Size
Exclusion Chromatography (top). Retention volume values of αSyn monomers analyzed by SEC in
the presence or absence of 200 Eq. Ca²⁺ at an ionic strength of 200 mM (bottom). **(E)** ¹H-¹⁵N HSQC
spectra of solutions of αSyn supplemented with or without 200 Eq Ca²⁺ at an ionic strength of 200
mM. **(F)** Analysis of intensity difference (top), ratio (middle) and Chemical Shift Perturbation
(bottom) of the N-H nuclei employing the spectrum obtained at in the absence of Ca²⁺, at 200 mM of
ionic strength, as reference. Gray bars represent the residues without signal assignments. (* Represents
signals that disappeared in the condition tested but that are present in the reference spectrum)

Figure 4. Bacterial components present in the purification of recombinant proteins affect the conformational state of the monomeric αSyn. IEX-αSyn and HPLC-αSyn were solved by analytical size exclusion chromatography **(A)** and analytical reverse phase chromatography **(B)**. **(C)** Left: Maximal ThT intensity values of fibrils created using IEX-αSyn (red) and HPLC-αSyn (gray, taken from **Fig. 1.D** for comparison). Right: TEM visualization of representative fibrils obtained at 200 mM NaCl using IEX-αSyn (upper) and HPLC-αSyn (lower). **(D)** Left: Maximal ThT intensity values of fibrils created using HPLC-αSyn in the presence (purple) or absence (black) of LPS. Right: TEM visualization of representative fibrils obtained at 300 mM NaCl using HPLC-αSyn in the presence (upper) or absence (lower) of LPS. **(E)** ^{1}H-^{15}N HSQC spectra of solutions of IEX-αSyn (red) and HPLC-αSyn (black) at an ionic strength of 200 mM. **(F)** Analysis of intensity difference (top), ratio (middle) and chemical shift perturbation (bottom) of the N-H nuclei of IEX-αSyn employing the spectrum of HPLC-αSyn as reference. Gray bars represent the residues without signal assignments. (* Represents signals that disappeared in the condition tested but that are present in the reference spectrum) **(G)** ^{1}H-^{15}N HSQC spectra of solutions of HPLC-αSyn in the presence (purple) or absence (black) of LPS at an ionic strength of 200 mM. **(H)** Analysis of intensity difference (top), ratio (middle) and chemical shift perturbation (bottom) of the N-H nuclei of HPLC-αSyn treated with LPS employing the spectrum of HPLC-αSyn without LPS as reference. Gray bars represent the residues without signal assignments. (* Represents signals that disappeared in the condition tested but that are present in the reference spectrum) **(I)** Alignment of the residues affected by the compaction of αSyn under the conditions indicated in the schematic diagram. The method to determine the affected residues is shown in the **Fig. S6.**

Figure 5. N103t-C terminus truncated αSyn does not adopt the compact conformation.
(A) Schematic representation of the truncated form N103t, in which the region detected to be relevant
for the compaction of the monomers has been removed. (B) ThT-monitored fibril formation of C-
terminus truncated N103t αSyn at 50, 150 and 300 mM NaCl. (C) TEM visualization of N103t αSyn
fibrils generated at 300 mM NaCl. (D) Representative chromatograms of solutions of N103t αSyn at
different concentrations of NaCl solved by Size Exclusion Chromatography.

894
895
896 **Figure 6. The conformational state of the monomeric αSyn provides differences between**
897 **fibril morphologies.** The initial state of monomeric αSyn and the factors that induce affect the C-
898 terminus domain determine the final morphology of the fibrils. In the brain, this phenomenon could
899 trigger the development of PD or MSA.

900
901