The midpoint of cortical thinning between late childhood and early adulthood differs between individuals and brain regions: Evidence from longitudinal modelling in a 12-wave neuroimaging sample

Abstract

Charting human brain maturation between childhood and adulthood is a fundamental prerequisite for understanding the rapid biological and psychological changes during human development. Two barriers have precluded the quantification of maturational trajectories: demands on data and demands on estimation. Using high-temporal resolution neuroimaging data of up to 12-waves in the HUBU cohort (N = 90, aged 7-21 years) we investigate changes in apparent cortical thickness across childhood and adolescence. Fitting a four-parameter logistic nonlinear random effects mixed model, we quantified the characteristic, s-shaped, trajectory of cortical thinning in adolescence. This approach yields biologically meaningful parameters, including the midpoint of cortical thinning (MCT), which corresponds to the age at which the cortex shows most rapid thinning - in our sample occurring, on average, at 14 years of age. These results show that, given suitable data and models, cortical maturation can be quantified with precision for each individual and brain region.

Keywords

brain maturation; adolescence; cortical thickness; nonlinear mixed models, sex differences, rostral middle frontal gyrus, rostral anterior cingulate

Introduction

The human cortex undergoes protracted microscopic and macroscopic structural changes between childhood and adulthood[1]. Individual differences in cortical structure have been associated with a range of phenotypic differences, including physical and mental health[2–5], neurodevelopmental disorders as well as cognitive performance in childhood and adolescence[2,6]. Although other measures of brain structural development such as brain volume or white matter connectivity provide complementary insights into brain maturation, cortical thinning is one of the most widely used proxies of brain maturation[7–9]. We here use the term *apparent* cortical thickness throughout to highlight that Magnetic Resonance Imaging (MRI) studies measure a proxy of cortical thickness, that follows a similar spatial patten to that observed in histological studies[10], but has absolute values that may be influenced by signal intensities and contrasts[11].

Previous work has generally observed developmental decreases in cortical thickness from childhood (after 2-3 years of age) to early adulthood[8], with longitudinal studies showing that the rate of cortical thinning increases in adolescence[9,12]. There is an emerging consensus on a characteristic s-shaped, non-linear trajectory at the population level (Figure 1)[9,12]. This process of cortical thinning is thought to reflect a range of underlying biological processes[13], including increasing myelination of the deeper cortical layers[14] and decreasing synaptic density[15].

Figure 1. Schematic illustration of cortical thinning during adolescence, showing the Midpoint of Cortical Thinning (MCT), i.e., the age of most rapid cortical thinning.

Hypotheses concerning individual differences in the process of cortical thinning and
their link to psychological development feature prominently in neurodevelopmental
theories. For instance, it has been hypothesized that a life history of adversity may lead to
accelerated[16,17] or delayed[18,19] cortical maturation. The developmental mismatch theory
suggests that a mismatch in maturity between subcortical and cortical brain regions (with
frontal regions commonly thought to thin last) may help explain the prevalence of risk-taking
behaviour during adolescence when this maturation disparity is thought to be maximal[20,21].
At the group comparison level, hypotheses posit that girls demonstrate earlier cortical
maturation than boys[22–24], with these differences hypothesized to underlie developmental
differences in behavioural and psychopathological phenotypes. Similarly, Nunes et al. (2020)
hypothesized that children with autism spectrum disorder are characterized by accelerated
brain maturation[25]. Overall, hypothesized differences in cortical maturation are central to
some of the most influential neurodevelopmental theories. However, the data and analytic
methods currently used to capture maturation are no match for ambitions in understanding
and applying the construct of cortical maturation.

Longitudinal data is costly and time-consuming to procure, therefore, most
neuroimaging studies to-date rely on cross-sectional data. Cross-sectional data precludes
the investigation of developmental changes and cross-sectional measures necessarily
conflates distinct sources of cortical thickness differences (baseline thickness, the onset of
maturation, as well as speed and total amount of thinning). Only under extremely restrictive
assumptions (e.g., identical brain thickness in early childhood and late adolescence, identical
rates of thinning) can a cross-sectional measure be used as a proxy for development. Given
that these assumptions are known to be empirically untrue[1], our empirical knowledge of
cortical maturation is likely to be extremely limited.

Where longitudinal data does exist, the average number of waves per subject is
generally below three, and the time between scans is 2.5 years on average[26]. This limits our
ability to observe more subtle brain changes in time periods with ongoing maturation such
as adolescence, as well as our ability to capture the trajectory of cortical thinning, which is
known to be non-linear[27]. This is particularly true at the individual level. That said, large
cohorts with multiple time points, like ABCD[28], are currently emerging.

Most longitudinal studies investigating individual differences to date have used linear
modelling approaches, such as linear mixed effects models, using the percentage of, or
absolute change, in cortical thickness between two ages as an indicator of maturation of a
given brain region, with larger changes commonly equated to more protracted
maturation[29,30]. However, these estimates are typically confounded by the initial thickness
of a region - thicker regions can thin more than thinner regions. Even when such confounds
are controlled for, the absolute change in thickness remains dependent on the precise age
range studied. Alternative, nonlinear approaches, such as Generalized Additive Mixed
Models (GAMMs)[31], can capture complex nonlinear relationships and are excellent tools for
predictive purposes. Nonlinear mixed models offer a similarly flexible approach for capturing
complex nonlinear relationships to GAMMs. A particular advantage of nonlinear mixed
models is that they yield readily interpretable parameters, informative, of, e.g., ages of rapid
development (Figure 1), making them an attractive, but currently underused, tool for
developmental neuroscientists.

To address the challenge of quantifying cortical maturation at the individual level, we
here leverage a unique dataset (with up to 12 longitudinal measurements between late
childhood and early adulthood) and a quantitative framework currently underutilized in
cognitive neuroscience (non-linear random effects modelling) to demonstrate that cortical
maturity can be defined, and estimated, at the individual level, offering a new window of
insight into cortical development across adolescence. Nonlinear mixed models are a
powerful tool that can capture developmental processes at the individual level[31] and yield
readily interpretable parameters. One of these parameters allows us to provide a novel
quantitative definition of cortical maturation during adolescence at the individual level: The
midpoint of cortical thinning (MCT, see Figure 1). The MCT is the point in adolescent
development where the rate of cortical thinning is at its peak for an individual. We show that
the MCT can be used to study the extent to which cortical maturation differs between
individuals, sexes, and brain regions.

Materials and Methods

Cohort

The present longitudinal study included data from 90 typically developing children and adolescents (53 females, 37 males), who were enrolled in the longitudinal HUBU ("*Hjernens Udvikling hos Børn og Unge*", Brain Maturation in Children and Adolescents) study. The HUBU study was initiated in 2007, where 95 participants (55 females, 40 males) aged 7 - 13 years and their families were recruited from three elementary schools in the Copenhagen, DK, suburban area[32]. All children whose families volunteered were included, except for children with a known history of neurological or psychiatric disorders or significant brain injury. Prior to participation, all children assented to the study procedures and informed written consent was obtained from parents. Informed written consent was also obtained from the participants themselves when they turned 18 years of age. The study was approved by the Ethical Committees of the Capital Region of Denmark (H-KF-01-131/03 and H-3-2013-037) and performed in accordance with the Declaration of Helsinki. Participants were scanned up to 12 times with scanning intervals of 6 months for the first 10 visits, one year between visits 10 and 11, and three years between visits 11 and 12 (Figure 2).

Here, we included data from the first 12 assessments of the HUBU study. All baseline MRI scans were evaluated by an experienced neuroradiologist, and all raw images were visually inspected to ensure sufficient quality. Five participants were excluded from the present study, due to receiving a psychiatric diagnosis after study initiation ($N = 2$), incidental clinical finding on the MRI scan ($N = 1$), or no MRI scans or FreeSurfer outcomes of sufficient quality ($N = 2$). Our final sample analysed here consisted of 90 participants (53 females, 37 males) aged 7.6 - 21.6 years. For this sample, we excluded 73 MRI sessions if one of the following criteria was met: participants did not finish the MRI session (2 participants, 2 scans), participants were not scanned due to metallic dental braces (14 participants, 31 scans), had poor MR-image quality (22 participants, 31 scans), or had acquired a brain injury after baseline (1 participant, 9 scans). A total of 745 valid MRI scans (scans per participant: range = 1 - 12, mean = 8.3, median = 10, interquartile range (*IQR*) = 8 – 12) were included in the statistical analyses. Data from the HUBU cohort has previously been used in cross-sectional[33–37] and longitudinal[32,38] studies examining brain-behavioural relationships.

133 **Figure 2**: Spacing and timing of scans for each participant (Panel A) and a histogram of the

134 number of scans included in the analysis (Panel B).

MRI protocol

Participants underwent structural MRI on a 3T Siemens Magnetom Trio MR scanner (Siemens, Erlangen, Germany) using an eight-channel head coil (Invivo, FL, USA). Two T1-weighted images were acquired using a 3D MPRAGE sequence (TR = 1550 ms, TE = 3.04 ms, matrix = 256 × 256, 192 sagittal slices, $1 \times 1 \times 1$ mm^3 voxels, acquisition time = 6:38). A T2-weighted image was acquired using a 3D turbo spin echo sequence (TR = 3000 ms, TE = 354 ms, FOV = 282 × 216, matrix = 256 × 196, 192 sagittal slices, $1 \times 1 \times 1$ mm^3 voxels, acquisition time = 8:29).

FreeSurfer pre-processing and extraction of cortical thickness

All T1-weighted and T2-weighted images were processed using tools available in the FreeSurfer (version 6.0) software suite[39–41]. Cortical surface reconstruction was implemented using the following procedures: skull stripping, non-uniformity correction, white matter segmentation, creation of initial mesh, correction of topological defects, and creation of optimal white and pial surfaces[39–41]. Images were then processed with the longitudinal stream[42] in FreeSurfer, to estimate changes in cortical thickness across time. Apparent cortical thickness was calculated as a measure of the shortest distance between the white and pial surfaces. Cortical grey matter parcellations were based on surface-based nonlinear registration to the Desikan-Killiany atlas based on gyral and sulcal patterns and Bayesian classification rules[41], yielding estimates for 34 ROIs in each hemisphere. Cortical thickness estimates were averaged across the hemispheres.

To check the quality of the FreeSurfer outputs, we followed the Enigma protocol (http://enigma.ini.usc.edu/protocols/imaging-protocols/). We used statistical outlier detection for average thickness and surface area of each cortical parcel. Statistical detection of both within- and between-subject outliers was performed based on the age, sex, and age-by-sex adjusted residuals derived using GAMMs, to account for age and sex differences in the brain measures. Scans of concern were then visually inspected. In line with the Enigma protocol, we did not perform manual editing on the FreeSurfer outputs and instead discarded data of questionable quality. Based on these quality checks, we excluded one participant from the entire study. Furthermore, we excluded the following cortical parcels

from statistical analysis for *N* participants: right temporal pole (*N* = 1), left frontal pole (*N* =

1), right paracentral (*N* = 1), right cuneus (*N* = 1), right lateral occipital (*N* = 1), and left fusiform

gyrus (*N* = 2).

Finally, the Euler number, which indicates the topological complexity of the

reconstructed cortical surface, was extracted from FreeSurfer for each scan and used as a

proxy for in-scanner motion and a quantitative measure of image data quality in our

statistical analyses, to account for potential systematical bias in data quality across age. The

Euler number has been correlated with visual image quality inspection scores as well as with

cortical thickness in a regionally heterogeneous pattern across datasets[43]. The Euler number

was extracted for each hemisphere and summed to produce one value per scan.

Statistical analyses

We here modelled cortical thinning between childhood and adulthood using nonlinear

mixed models implemented via the saemix[44] package (version 2.4) in R (version 4.1.0) and

RStudio (version 1.4.1717). To capture the characteristic s-shape of cortical thinning, we fit

the four-parameter logistic function[45], as defined in the Results section. See [BLINDED] for

our analysis code. Cortical thickness was modelled as the dependent variable and age as the

independent variable. We fit a model to mean cortical thickness, as well as for one for each

of the 34 ROIs of the Desikan-Killiany atlas[41]**.** We included sex at birth and in-scanner motion

as covariates in all models, to account for potential differences thereof in developmental

trajectories[46,47]. In-scanner motion was operationalized by the Euler number (*M* = -137.07, *SE*

= 5.33, range = -298.2 - -69.08). Females were coded as the reference group. While sex was

a covariate of interest, and motion a covariate of no interest, both are modelled the same

way in the NLMM framework. Covariate parameters can, in principle, be included to control

for the effect of motion and sex on any of the main fixed effect parameters of the model (in

our case, the two asymptotes, the MCT, and the hill). This yields a maximum of eight

covariates. To avoid potential overfitting with a highly parametrized model[48], we

determined the optimal covariate model via forward stepwise model selection using mean

thickness as the outcome variable. To the best of our knowledge, no consensus methodology

for model selection with covariates exists for nonlinear mixed models. We use forward

selection as a pragmatic and tractable solution in the context of demanding model
estimation. Although some have advocated for more complex model selection algorithms,
such as best subset selection[49], recent work suggests that forward selection can perform
similarly to these alternatives[50]. The process starts with no covariate parameters in the
model. It then iteratively adds the variable that best improves the fit. For all covariate
parameters not in the model (e.g., eight at the first iteration), we check their *p*-value when
added to the model one at a time. We chose the covariate parameter with the lowest *p*-value
less than 0.05. This process was continued until no new predictors could be added. Based on
this model selection process, we allowed the upper and lower asymptote, and MCTs to differ
between sexes and included the motion parameter as a covariate for the lower asymptote
(see Supplementary Table 1 for the full results). The remaining parameters were held
constant for sex and motion. Estimates were obtained for the four parameters of the logistic
function, as well as the two covariates. Precision was assessed by inspecting the coefficient
of variation (CV) for each parameter, as provided by saemix. CVs are standardized measures
of dispersion and are calculated as the ratio of the standard deviation to the mean. CVs <
20% are generally considered acceptable[51].

In a second step, we assessed differences in MCT estimates across different brain
regions. We assessed whether MCTs across brain regions could be constrained to equality
using Confirmatory Factor Analysis (CFA), as implemented in lavaan[52] for R (version 0.6-9).
We used full information maximum likelihood with robust standard errors to account for
missingness and nonnormality. We estimated a one-factor CFA model in which factor
loadings were freely estimated for each brain region. We then compared this model to one
where the factor loadings were still freely estimated, but the intercepts were constrained to
equality using the likelihood ratio test. A significant likelihood ratio test here indicates a loss
in model fit for a constrained model, providing evidence that brain regions differ in their
MCTs.

Third, we used Exploratory Factor Analysis to assess the dimensionality of MCTs across
regions and to identify maturational factors capturing developmental trends across regions.
This was implemented through parallel analysis via the psych[53] package for R (version 2.1.6),
using an oblique oblimin transformation.

Finally, we tested the popular hypothesis of a posterior-anterior gradient of

development[54] by correlating MCTs for each region with their y-coordinates in MNI space.

Coordinates were obtained from the brainGraph[55] package in R (version 3.0.0). In an

additional exploratory analysis we also correlated MCTs for each region with their z-

coordinates in MNI space to test for a potential dorsal-ventral gradient.

Results

233   Cortical thickness showed nonlinear changes between childhood and early adulthood across

234   cortical regions (Figure 3). The characteristic sigmoid, or s-shaped, curve, of cortical thinning,

235   was apparent across most brain regions. We captured this shape using the four-parameter

236   logistic function:

237
$$y = A_{Lower} + \frac{A_{Upper} - A_{Lower}}{1 + \left(\frac{x}{Inflection}\right)^{-Hill}}$$

The function yields four biologically meaningful parameters: The upper asymptote (A_{Upper},

maximal apparent thickness in mm), lower asymptote (A_{Lower}, minimal apparent thickness in

mm), and *Hill*, the slope of change, and the *Inflection*. We were particularly interested in the

latter parameter, which corresponds to the MCT, and was used here as an index of cortical

maturation (see Figure 1). The MCT can be compared across individuals, sexes, and brain

regions. To this end, we first modelled cortical thickness averaged across the cortex,

identified the average pattern of maturation, and assessed sex differences. Next, we

modelled cortical thickness for different brain regions to investigate patterns of maturation

across the cortex. Sex and in-scanner motion, as a proxy for image quality, was controlled for

in all analyses.

Because the sigmoid is asymptotic, there is no age at which the brain is mature.

Instead, the brain develops throughout the age range investigated here (7 – 21 years).

Parameters indicated that mean thickness showed high levels in late childhood, with an

upper asymptote of 2.95 mm (*SE* = 0.01 mm) and decreasing thereafter to a lower asymptote

of 2.62 mm (*SE* = 0.03 mm). Our central parameter of interest, the MCT, was estimated to be

14.36 years (*SE* = 0.28 years). See Table 1. We observed a substantial range of MCTs across

individuals, with a minimum and maximum of 12.25 and 19.54 years (Figure 4) and a variance

of 1.79 years (*SE* = 0.46 years). Together, our analysis demonstrates that we can estimate a novel, quantitative definition of cortical maturity which is independent of overall thickness and shows substantial differences between people.

Figure 3. Apparent cortical thickness across brain regions (defined by the Desikan-Killiany atlas) as a function of age. Average and individual trajectories for each participant are shown.

261 **Table 1**: NLMM Parameter estimates for mean cortical thickness

Parameter	Estimate	SE	CV (%)	p
A_{Lower}	2.62	0.03	1.3	-
$b_{Sex(ALower)}$	-0.07	0.04	55.1	0.035
$b_{Motion(ALower)}$	0.00	<0.01	52.0	0.027
A_{Upper}	2.95	0.01	0.4	-
$b_{Sex(AUpper)}$	-0.06	0.02	30.5	0.001
MCT	14.36	0.28	2.0	-
$b_{Sex(MCT)}$	1.92	0.51	26.7	<0.001
Hill	-6.34	0.42	6.6	-

262 *Note. CV* = Coefficient of variation – the ratio of the standard deviation to the mean. Used to
263 assess precision. *CV* values ≤ 20% are deemed acceptable. *SE* = standard error, NLMM =
264 Nonlinear Mixed Models. Females were coded as the reference group.

There are substantial individual differences in the MCT

We found that the nonlinear model fit the data well (Supplementary Figure 1). To test
whether our model fit better than a simpler alternative, we compared the model fit of the
four-parameter logistic nonlinear model to a simple linear model. We found that the four-
parameter logistic model fits better than the simple linear model (ΔAIC = -137.22),
suggesting that our more complex model is plausible across most regions. Mean cortical
thickness was estimated with very good precision for the asymptotes, MCT, and the hill, as
indicated by low coefficients of variation (*CV*; Table 1). The MCT was uncorrelated with the
upper asymptote, showing that it is independent of the initial thickness of a region (r = 0.01,
$t(88)$ = 0.09, p = .925). It was also independent of the mean cortical thickness in early
adulthood (A_{Lower}, r = -0.13, $t(88)$ = -1.22, p = .227). This independence means that cross-
sectional measurements of cortical thickness will likely not function well as an approximation
of cortical maturity. The rate of change (*Hill*) was also not associated with the MCT (r = -0.16,
$t(88)$ = -1.48, p = .141). A sensitivity analysis shows that results were very similar when only
participants with at least three scans were included (N = 84, Supplementary Table 2).

Significant, but noisy, sex differences in trajectories

There were significant sex differences: Females started with thicker cortices than males (b =
-0.06, p = .001), had thicker corteces in adulthood (b = -0.07, p = .035) and showed earlier MCT

(*b* = 1.92, *p* < .001, Figure 4). However, the *CVs* for these estimates were all well above 20%

(Table 1), indicating that these parameters were estimated with low precision. The low

precision was likely due to the small number of males in the sample (*N_Males* = 37).

Figure 4. *Panel A*: Predicted apparent mean cortical thickness as a function of age. Estimated

individual trajectories and the average trajectory is shown, as well as the average MCT. *Panel*

B: Density plot of MCTs showing individual differences in the sample. *Panel C:* Sex

differences in the sample. MCT = Midpoint of Cortical Thinning.

MCTs differ across the cortex

To examine the specificity in maturational timing across regions, we estimated the MCT for

each of the 34 cortical regions of the Desikan-Killiany atlas independently (Figure 5), again

using non-linear mixed models and the four-parameter logistic function. The four

parameters were estimated with good precision (*CV* < 20%) for all regions[51], except for the

entorhinal, lingual, pericalcarine, temporal pole, and transverse temporal regions, which

were all excluded from subsequent analyses (see Supplementary Tables 3 and 4 for all

estimates). For these regions, the logistic may not be a good approximation of the functional

relationship between age and cortical thickness. For the pericalcarine, for instance, we find

that a linear model fits marginally better than the four-parameter logistic (ΔAIC = 0.50),

potentially reflecting the early maturation for this region. The enthorinal is known to be more

likely to be affected by eye-movement artifacts and may therefore show a poor model fit.

Mean MCTs ranged from 13.51 years for the cuneus to 17.75 years for the rostral

anterior cingulate cortex (Figure 5). The brain regions that reached the MCT first were the

occipital (cuneus and lateral occipital) and parietal (precuneus, superior and inferior parietal)

cortices, the rostral middle frontal cortex, and the cortex of the banks of the superior

temporal sulcus (bankssts). The brain regions that reached the MCT last were the caudal and

rostral anterior cingulate and parahippocampal cortices, followed by the sensorimotor pre-

and postcentral. Moreover, the frontal lobe regions (precentral, caudal middle frontal, rostral

middle frontal, superior frontal, and frontal pole) showed the fastest rate of change (i.e.,

steepest hill), while the slowest rate of change (i.e., flattest hill) was observed in the

parahippocampal, cuneus, (caudal and rostral) anterior cingulate, and superior temporal

cortices. Most regions showed a relative independence of the MCT, upper and lower

asymptote and hill (Figure 6) and low parameter correlations (Supplementary Table 5).

Figure 5. Individual and regional differences in thinning shown in raincloud plots. The estimated kernel density is shown to visualize the distribution of values. The median is shown as the black vertical line within a bar, which itself shows the interquartile range. Black horizontal lines show the 1.5 interquartile range. Values beyond these lines can be considered outliers.

Figure 6. Parameter estimates for the MCT (Panel A), hill (Panel B), upper asymptote (Panel C) and lower asymptote (Panel D) plotted across the cortex. Darker shades reflect higher parameter estimates. Excluded regions are shown in grey.

Figure 7. Factor loading for Factor 1 (Panel A) and Factor 2 (Panel B) across the cortex. Only loadings > 0.6 are shown to facilitate interpretation. Darker colours represent higher loadings.

Next, we implemented a formal quantitative test to examine whether regions differed in their MCTs. To do so, we compared a confirmatory factor model with intercepts for MCTs across regions constrained to equality (reflecting the hypothesis that regions mature at the same approximate age) to a model where they are estimated freely, reflecting the hypothesis that regions mature at distinct ages). Despite the considerable added complexity, we found that allowing MCTs to differ between regions, substantially improved model fit ($\Delta X^2(29) = 930.69$, $p < .001$) indicating pronounced differences in maturational timing between regions independent of their overall thickness.

In addition to MCT-differences across regions globally, we also examined whether MCTs are linked between some or all regions: In other words, if a person is mature in one region, are they then also more mature in all other regions, or are there clusters of brain regions that covary in their relative maturity? To examine this question, we first examined the fit of a one-factor confirmatory model, testing the hypothesis that a single factor could capture MCTs across the brain. This model fit poorly ($X^2(377) = 1236.68$, $p < .001$, CFI = 0.787, SRMR = 0.059, RMSEA = 0.159 [0.150, 0.169]). To determine whether a more complex model might fit the data, we used Exploratory Factor Analysis. Eigenvalues of a parallel analysis suggested that a two-factor model yields the best, albeit imperfect, solution ($X^2(349)$ = 633.85, $p < .001$). The two-factor model explained 79.3% of the variance in MCTs. Several central , as well as cingulate, regions loaded strongly onto Factor 1. Parietal regions regions and the rostral middle frontal loaded strongly onto Factor 2 (Figure 7, Supplementary Table 6).

Finally, we explored whether there is a regional ordering in the timing of maturation in line with hypotheses of a posterior-to-anterior gradient across the cortex[7,56]. We correlated inflection ages of each region with the region's average y-coordinate in MNI space, as contained in mni.y in brainGraph[55]. We found no evidence for a significant correlation between the spatial location and the MCT ($t(27) = 0.12$, $p = .903$, $r = 0.02$, Bayes Factor = 0.35). In an additional exploratory analysis, we also analyzed whether there is a regional ordering in the timing of maturation in line with hypotheses of a dorsal-ventral gradient across the cortex[57]. We correlated inflection ages of each region with the region's average z-coordinate in MNI space, as contained in mni.z in brainGraph[55]. We found no evidence for a significant correlation between the spatial location and the inflection ages ($t(19) = -0.65$, $p = .522$, $r = -0.15$, Bayes Factor = 0.45).

Together, these analyses offer new insight into cortical maturation. We demonstrate, for the first time, that it is possible to estimate non-linear maturation independent of overall cortical thickness. Maturational trajectories differed between individuals and cortical regions. The ability to estimate these differences offers a new window into elucidating long-standing debates concerning the speed of maturation, its association with early adversity, and the implications for cognitive and mental health development.

Discussion

Using longitudinal data of up to 12-waves imaged between late childhood and early

adulthood and flexible nonlinear mixed models, we here show that cortical maturity as

indexed by the MCT can be separated from other cortical thickness parameters (i.e.,

asymptotes and slope – hill) and estimated precisely and reliably for each individual and most

brain regions. We identified a characteristic, s-shaped, trajectory of cortical thinning:

Cortical thickness was high in childhood, followed by decreases in early adolescence,

culminating around the age of 14 years, the average MCT. The reduction in cortical thickness

then decelerates to level off in late adolescence. This finding is in line with previous studies

showing s-shaped reductions in cortical thickness over adolescence[9,12] and extend the same

by providing estimates of upper limits in apparent cortical thickness (2.95 mm), lower limits

of apparent cortical thickness (2.62 mm), and, most importantly, an index of maturation: the

MCT (average of 14.36 years). This highlights that cortical thinning is protracted and shows

rapid changes in adolescence. This period of rapid brain development raises questions about

possible sensitive periods in adolescence[58,59]. Future research will be able to show whether

periods of structural change confer heightened plasticity in adolescence.

Developmental patterns differed between cortical regions, with the superior parietal

and precuneus showing some of the earliest MCTs, around 14 years, and cingulate regions

showing some of the latest MCTs, around 17 years. Early maturing regions were found in

lateral frontal, and parietal areas, while late-maturing regions were found in temporal and

dorsal central areas.

Our finding of an early MCT in several frontal areas is surprising, given previous, mostly

cross-sectional studies. There are several possible explanations for this result, which will

need to be tested in future research: First, while most frontal regions are usually imaged well,

the frontal pole is known to be difficult to image. We therefore advise to interpret the finding

for this region with caution. Second, we cannot rule out that data quality issues or model

misfit may have affected findings, although quality control procedures, statistical control for

in-scanner motion, diagnostic plots and precision estimates do not suggest that this was

likely to be the sole explanation. Third, while the scans were evenly distributed across most

ages, the interval between waves 11 and 12 was longer, and data was sparser at older ages.

This may cause the model fit to be poorer at older ages, although this would be more likely

to affect estimation of late maturing, rather than early maturing regions. Fourth, the late

maturation suggested by the, mostly, cross-sectional analyses to date, do not replicate in

appropriate longitudinal analyses like ours, because of the inherent limitation of cross-

sectional data for the examination of developmental patterns. Finally, it may be that cross-

sectional analyses have captured aspects of maturation that are independent of the MCT,

the time point of fastest thinning. It may be, for instance, that frontal regions asymptote

later than other region, even though the peak of thinning happens quite early. Altogether

this finding, along with recent work identifying structural brain networks and hubs[60–63],

supports a complex systems account, in which the brain matures in a distributed pattern.

Future studies could use complex systems approaches like network models to identify how

maturation across regions produces changes in cognition and mental states across

development.

We found evidence for pronounced individual differences, with MCTs differing by

several years between individuals. This supports the notion that brain maturation is highly

variable and invites questions of potential predictors and outcomes. We here investigated

sex as a potential predictor of individual differences. There was some evidence for sex

differences, with females showing earlier maturation than males, by about 2 years. This

supports initial evidence for earlier maturation in girls from older longitudinal studies[46,47] and

may be linked to earlier puberty and different socialization in girls[64,65]. Our estimates of sex

differences were relatively noisy, however. Future studies could use nonlinear mixed models

in larger cohorts, such as ABCD[28], to investigate the robustness of sex differences in

maturation. Future studies could also investigate other candidate predictors such as

environmental influences (e.g., adversity and education) to identify whether these

accelerate or decelerate maturation – a yet unsolved conundrum in developmental

science[18,19,66]. Investigations of potential outcomes of maturational differences, e.g.,

cognitive performance or psychiatric diagnosis, would be similarly fascinating, and could

eventually include distal effects, such as cognitive and brain changes during ageing[1,67].

It is worth noting that our analytical approach, using the four-parameter logistic

function, depends on the nature of cortical thinning: Other measures of brain maturation

(e.g., brain volume) likely show different developmental shapes and different maturational

timelines, not all of which will be amenable to estimating the MCT. The simple linear

decreases, reported for volume changes in some cortical regions, would not allow for a
quantitative 'midpoint.' However, nonlinear mixed models are extremely versatile and can
be fit, in principle, to any functional relationship. This also includes more complex
relationships than that captured by the four-parameter logistic fit here, allowing, e.g., for
asymmetries in development before and after adolescence. In the future, these nonlinear
mixed models can be used to model white matter trajectories and other morphometric
measures of cortical and subcortical development to better understand similarities and
differences across brain tissues. This will yield a more precise understanding of how changes
in grey and white matter work in concert to produce functional changes in the brain and
behaviour.

Readers may be interested in understanding the power prerequisites for using
nonlinear mixed models. Power in nonlinear mixed models depends on the interaction
between several factors, including the number of participants, number of timepoints,
spacing between time points, missingness and model complexity[68,69]. While past studies of
optimal design in pharmacology indicate that at least three time points may be sufficient to
estimate simple nonlinear mixed models[70], future studies will need to evaluate power for
plausible developmental functions and typical designs of neuroimaging studies in detail.

In conclusion, this study shows that apparent cortical thinning in adolescence is s-
shaped, with the most rapid changes occurring in mid-adolescence, at around 14 years of
age, on average. Further, we show that individuals vary substantially, with up to several
years, in the age at which the cortex undergoes most rapid changes. On a practical level, this
work shows that high-resolution temporal data, combined with nonlinear modelling
approaches, can be used to quantify brain maturation with unprecedented precision. This
will allow the field to provide rigorous tests of prominent theoretical models of adolescent
development, such as the structural mismatch hypothesis[20,21] or accelerated maturation
hypotheses[17].

Conflict of Interest Statement

The authors report no conflict of interest.

References

1. Bethlehem, R. A. I. *et al.* Brain charts for the human lifespan. *bioRxiv* 2021.06.08.447489

(2021) doi:10.1101/2021.06.08.447489.

2. Fuhrmann, D., Simpson-Kent, I. L., Bathelt, J., The CALM Team & Kievit, R. A. A

hierarchical watershed model of fluid intelligence in childhood and adolescence.

Cerebral Cortex (2019) doi:10.1093/cercor/bhz091.

3. Humphreys, K. L. *et al.* Stressful Life Events, ADHD Symptoms, and Brain Structure in

Early Adolescence. *Journal of Abnormal Child Psychology* **47**, 421–432 (2019).

4. Mofrad, S. A., Lundervold, A. J., Vik, A. & Lundervold, A. S. Cognitive and MRI

trajectories for prediction of Alzheimer's disease. *Scientific Reports* **11**, 2122 (2021).

5. Bos, M. G. N., Peters, S., van de Kamp, F. C., Crone, E. A. & Tamnes, C. K. Emerging

depression in adolescence coincides with accelerated frontal cortical thinning. *Journal*

of Child Psychology and Psychiatry **59**, 994–1002 (2018).

6. Shaw, P. *et al.* Intellectual ability and cortical development in children and adolescents.

Nature **440**, 676–679 (2006).

7. Squeglia, L. M., Jacobus, J., Sorg, S. F., Jernigan, T. L. & Tapert, S. F. Early adolescent

cortical thinning is related to better neuropsychological performance. *Journal of the*

International Neuropsychological Society : JINS **19**, 962–70 (2013).

8. Walhovd, K. B., Fjell, A. M., Giedd, J., Dale, A. M. & Brown, T. T. Through Thick and

Thin: a Need to Reconcile Contradictory Results on Trajectories in Human Cortical

Development. *Cerebral Cortex* **27**, (2017).

9. Tamnes, C. K. *et al.* Development of the cerebral cortex across adolescence: A

multisample study of interrelated longitudinal changes in cortical volume, surface area

and thickness. *Journal of Neuroscience* (2017) doi:10.1523/JNEUROSCI.3302-16.2017.

10. Cardinale, F. *et al.* Validation of FreeSurfer-Estimated Brain Cortical Thickness:

Comparison with Histologic Measurements. *Neuroinformatics* **12**, 535–542 (2014).

11. Furlong, C. *et al.* Application of stereological methods to estimate post-mortem brain

surface area using 3 T MRI. *Magnetic resonance imaging* **31**, 456–465 (2013).

12. Ducharme, S. *et al.* Trajectories of cortical thickness maturation in normal brain

development--The importance of quality control procedures. *Neuroimage* **125**, 267–279

(2016).

13. Jernigan, T. L., Baare, W. F., Stiles, J. & Madsen, K. S. Postnatal brain development:

structural imaging of dynamic neurodevelopmental processes. *Progress in brain*

research **189**, 77–92 (2011).

14. Natu, V. S. *et al.* Apparent thinning of human visual cortex during childhood is

associated with myelination. *Proc Natl Acad Sci USA* **116**, 20750 (2019).

15. Huttenlocher, P. R. Synaptic density in human frontal cortex - Developmental changes

and effects of aging. *Brain research* **163**, 195–205 (1979).

16. Keding, T. J. *et al.* Differential Patterns of Delayed Emotion Circuit Maturation in

Abused Girls With and Without Internalizing Psychopathology. *AJP* **178**, 1026–1036

(2021).

17. Belsky, J. Early-Life Adversity Accelerates Child and Adolescent Development. *Curr Dir*

Psychol Sci **28**, 241–246 (2019).

18. Tooley, U. A., Bassett, D. S. & Mackey, A. P. Environmental influences on the pace of

brain development. *Nature Reviews Neuroscience* **22**, 372–384 (2021).

19. Colich, N. L., Rosen, M. L., Williams, E. S. & McLaughlin, K. A. Biological aging in

childhood and adolescence following experiences of threat and deprivation: A

systematic review and meta-analysis. *Psychol Bull* **146**, 721–764 (2020).

20. Steinberg, L. Risk taking in adolescence: what changes, and why? *Ann. N. Y. Acad. Sci.*

1021, 51–58 (2004).

21. Casey, B. J., Getz, S. & Galván, A. The adolescent brain. *Dev Rev* **28**, 62–77 (2008).

22. Giedd, J. N. *et al.* Brain development during childhood and adolescence: A longitudinal

MRI study. *Nature neuroscience* **2**, 861–863 (1999).

23. Vijayakumar, N. *et al.* Thinning of the lateral prefrontal cortex during adolescence

predicts emotion regulation in females. *Soc Cogn Affect Neurosci* **9**, 1845–1854 (2014).

24. Peper, J. S., Burke, S. M. & Wierenga, L. M. Chapter 3 - Sex differences and brain

development during puberty and adolescence. in *Handbook of Clinical Neurology* (eds.

Lanzenberger, R., Kranz, G. S. & Savic, I.) vol. 175 25–54 (Elsevier, 2020).

25. Nunes, A. S. *et al.* Atypical age-related changes in cortical thickness in autism spectrum

disorder. *Scientific Reports* **10**, 11067 (2020).

26. Vijayakumar, N., Mills, K. L., Alexander-Bloch, A., Tamnes, C. K. & Whittle, S. Structural

brain development: A review of methodological approaches and best practices.

Developmental Cognitive Neuroscience **33**, 129–148 (2018).

27. Mills, K. L. *et al.* Inter-individual variability in structural brain development from late

childhood to young adulthood. *Neuroimage* **242**, 118450–118450 (2021).

28. Casey, B. J. *et al.* The Adolescent Brain Cognitive Development (ABCD) study: Imaging

acquisition across 21 sites. *Developmental Cognitive Neuroscience* **32**, 43–54 (2018).

29. Storsve, A. B. *et al.* Differential longitudinal changes in cortical thickness, surface area

and volume across the adult life span: regions of accelerating and decelerating change.

J Neurosci **34**, 8488–8498 (2014).

30. Schnack, H. G. *et al.* Changes in Thickness and Surface Area of the Human Cortex and

Their Relationship with Intelligence. *Cerebral Cortex* **25**, 1608–1617 (2015).

31. Davidian, M. & Giltinan, D. M. Nonlinear models for repeated measurement data: An

overview and update. *Journal of Agricultural, Biological, and Environmental Statistics* **8**,

387 (2003).

32. Madsen, K. S. *et al.* Maturational trajectories of white matter microstructure underlying

the right presupplementary motor area reflect individual improvements in motor

response cancellation in children and adolescents. *NeuroImage* **220**, 117105 (2020).

33. Angstmann, S. *et al.* Microstructural asymmetry of the corticospinal tracts predicts

right–left differences in circle drawing skill in right-handed adolescents. *Brain Structure*

and Function **221**, 4475–4489 (2016).

34. Gonzalez, M. R. *et al.* Brain structure associations with phonemic and semantic fluency

in typically-developing children. *Dev Cogn Neurosci* **50**, 100982–100982 (2021).

35. Madsen, K. S., Jernigan, T. L., Vestergaard, M., Mortensen, E. L. & Baaré, W. F. C.

Neuroticism is linked to microstructural left-right asymmetry of fronto-limbic fibre

tracts in adolescents with opposite effects in boys and girls. *Neuropsychologia* **114**, 1–10

(2018).

36. Madsen, K. S. *et al.* Brain microstructural correlates of visuospatial choice reaction time

in children. *NeuroImage* **58**, 1090–1100 (2011).

37. Klarborg, B. *et al.* Sustained attention is associated with right superior longitudinal

fasciculus and superior parietal white matter microstructure in children. *Hum Brain*

Mapp **34**, 3216–3232 (2013).

38. Plachti, A. *et al.* Only females show a stable association between neuroticism and

microstructural asymmetry of the cingulum across childhood and adolescence: A

longitudinal DTI study. *bioRxiv* 2021.08.31.458188 (2021)

doi:10.1101/2021.08.31.458188.

39. Dale, A. M. & Sereno, M. I. Improved Localizadon of Cortical Activity by Combining EEG

and MEG with MRI Cortical Surface Reconstruction: A Linear Approach. *Journal of*

Cognitive Neuroscience **5**, 162–176 (1993).

40. Dale, A. M., Fischl, B. & Sereno, M. I. Cortical Surface-Based Analysis: I. Segmentation

and Surface Reconstruction. *NeuroImage* **9**, 179–194 (1999).

41. Desikan, R. S. *et al.* An automated labeling system for subdividing the human cerebral

cortex on MRI scans into gyral based regions of interest. *NeuroImage* **31**, 968–980

(2006).

42. Reuter, M., Schmansky, N. J., Rosas, H. D. & Fischl, B. Within-subject template

estimation for unbiased longitudinal image analysis. *Neuroimage* **61**, 1402–1418 (2012).

43. Rosen, A. F. G. *et al.* Quantitative assessment of structural image quality. *Neuroimage*

169, 407–418 (2018).

44. Comets, E., Lavenu, A. & Lavielle, M. Parameter Estimation in Nonlinear Mixed Effect

Models Using saemix, an R Implementation of the SAEM Algorithm. *Journal of*

Statistical Software; Vol 1, Issue 3 (2017) (2017) doi:10.18637/jss.v080.i03.

45. An, H., Landis, J. T., Bailey, A. G., Marron, J. S. & Dittmer, D. P. dr4pl: A Stable

Convergence Algorithm for the 4 Parameter Logistic Model. *The R Journal* **11**, 171–190

(2019).

46. Lenroot, R. K. *et al.* Sexual dimorphism of brain developmental trajectories during

childhood and adolescence. *NeuroImage* **36**, 1065–1073 (2007).

47. Wierenga, L. M. *et al.* A Key Characteristic of Sex Differences in the Developing Brain:

Greater Variability in Brain Structure of Boys than Girls. *Cerebral Cortex* **28**, 2741–2751

(2017).

48. Bilger, M. & Manning, W. G. Measuring overfitting and mispecification in nonlinear

models. *Health, Econometrics and Data Group Working Paper* **11**, 25 (2011).

49. Bertsimas, D., King, A. & Mazumder, R. BEST SUBSET SELECTION VIA A MODERN

OPTIMIZATION LENS. *The Annals of Statistics* **44**, 813–852 (2016).

50. Trevor Hastie, Robert Tibshirani, & Ryan Tibshirani. Best Subset, Forward Stepwise or

Lasso? Analysis and Recommendations Based on Extensive Comparisons. *Statistical*

Science **35**, 579–592 (2020).

51. Reed, G. F., Lynn, F. & Meade, B. D. Use of coefficient of variation in assessing

variability of quantitative assays. *Clin Diagn Lab Immunol* **9**, 1235–1239 (2002).

52. Rosseel, Y. lavaan: An R package for Structural Equation Modeling. *Journal of Statistical*

Software **48**, 1–36 (2012).

53. The Personality Project's Guide to R. http://personality-project.org/r/psych/.

54. Shaw, P. *et al.* Neurodevelopmental trajectories of the human cerebral cortex. *J.*

Neurosci. **28**, 3586–3594 (2008).

55. Watson, C. G. *brainGraph: Graph Theory Analysis of Brain MRI Data.* (2020).

56. MacPherson, S. E. *et al.* Processing speed and the relationship between Trail Making

Test-B performance, cortical thinning and white matter microstructure in older adults.

Cortex **95**, 92–103 (2017).

57. Chen Chi-Hua *et al.* Genetic topography of brain morphology. *Proceedings of the*

National Academy of Sciences **110**, 17089–17094 (2013).

58. Fuhrmann, D., Knoll, L. J. & Blakemore, S. J. Adolescence as a sensitive period of brain

development. *Trends in cognitive sciences* (2015) doi:10.1016/j.tics.2015.07.008.

59. Frankenhuis, W. E. & Walasek, N. E. Evolution of sensitive periods: Bridging theory and

data. *Developmental Cogntitive Neuroscience* (in press).

60. Baum, G. L. *et al.* Modular segregation of structural brain networks supports the

development of executive function in youth. *Current Biology* (2017)

doi:10.1016/j.cub.2017.04.051.

61. Baum, G. L. *et al.* Development of structure–function coupling in human brain networks

during youth. *Proc Natl Acad Sci USA* **117**, 771 (2020).

62. Power, J. D., Fair, D. A., Schlaggar, B. L. & Petersen, S. E. The development of human

functional brain networks. *Neuron* **67**, 735–748 (2010).

63. Oldham, S. & Fornito, A. The development of brain network hubs. *Developmental*

Cognitive Neuroscience **36**, 100607 (2019).

64. Goddings, A.-L., Beltz, A., Peper, J. S., Crone, E. A. & Braams, B. R. Understanding the

Role of Puberty in Structural and Functional Development of the Adolescent Brain.

Journal of Research on Adolescence **29**, 32–53 (2019).

65. Hyde, J. S. Gender Similarities and Differences. *Annu. Rev. Psychol.* **65**, 373–398 (2014).

66. Ferschmann, L., Bos, M., Herting, M., Mills, K. L. & Tamnes, C. K. Contextualizing

adolescent structural brain development. (2021) doi:10.31234/osf.io/w8jq6.

67. Tamnes, C. K. *et al.* Brain development and aging: Overlapping and unique patterns of

change. *NeuroImage* **68**, 63–74 (2013).

68. Retout, S., Comets, E., Samson, A. & Mentré, F. Design in nonlinear mixed effects

models: optimization using the Fedorov-Wynn algorithm and power of the Wald test

for binary covariates. *Stat Med* **26**, 5162–5179 (2007).

69. Roy, A. & Ette, E. I. A pragmatic approach to the design of population pharmacokinetic

studies. *AAPS J* **7**, E408–E420 (2005).

70. al-Banna, M. K., Kelman, A. W. & Whiting, B. Experimental design and efficient

parameter estimation in population pharmacokinetics. *J Pharmacokinet Biopharm* **18**,

347–360 (1990).