Feasibility of enhancing carbon sequestration and stock capacity in temperate and boreal European forests via changes to management regimes

Authors: Dalmonech D.[1,†], Marano G.[1,2,†,*], Amthor J.S.[3], Cescatti A.[4], Lindner M.[5], Trotta C.[6], Collalti A.[1,7,†]

1. Forest Modelling Lab., Institute for Agriculture and Forestry Systems in the Mediterranean, National Research Council of Italy (CNR-ISAFOM), Via Madonna Alta 128, 06128, Perugia, Italy

2. Forest Ecology, Institute of Terrestrial Ecosystems, Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland

3. Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, Arizona, 86001, USA

4. European Commission, Joint Research Centre (JRC), Ispra, Italy

5. European Forest Institute, Platz der Vereinten Nationen 7, 53113, Bonn, Germany

6. Department of Innovation in Biological, Agro-food and Forest Systems (DIBAF), University of Tuscia, 01100 Viterbo, Italy

7. Department of Agriculture and Forest Sciences, University of Tuscia, 01100 Viterbo, Italy

corresponding author

[†]*contributed equally*

Corresponding author: E-mail: gina.marano@usys.ethz.ch

ORCID:

Dalmonech D.: 0000-0002-1932-5011

Marano G.: 0000-0003-2600-984X

Amthor J.: 0000-0001-8601-403X

Cescatti A.: 0000-0002-2769-2591

Lindner M.: 0000-0002-0770-003X

Trotta C.: 0000-0001-6377-0262

Collalti A.: 0000-0002-4980-8487

Abstract

Forest management practices might act as nature-based methods to remove CO_2 from the atmosphere and slow anthropogenic climate change and thus support an EU forest-based climate change mitigation strategy. However, the extent to which diversified management actions could lead to quantitatively important changes in carbon sequestration and stocking capacity at the tree level remains to be thoroughly assessed. To that end, we used a state-of-the-science bio-geochemically based forest growth model to simulate effects of multiple forest management scenarios on net primary productivity (NPP) and potential carbon woody stocks (pCWS) under twenty scenarios of climate change in a suite of observed and virtual forest stands in temperate and boreal European forests. Previous modelling experiments indicated that the capacity of forests to assimilate and store atmospheric CO_2 in woody biomass is already being attained under business-as-usual forest management practices across a range of climate change scenarios. Nevertheless, we find that on the long-term, with increasing atmospheric CO_2 concentration and warming, managed forests show both higher productivity capacity and a larger potential pool size of stored carbon than unmanaged forests as long as thinning and tree harvesting are of moderate intensity.

Keywords: adaptive forest management, modeling, virtual forests, climate change, carbon sequestration

1 | Introduction

Forest ecosystems have the capacity to slow anthropogenic climate change by reducing the rate of atmospheric carbon dioxide (CO_2) increase through increased photosynthetic CO_2 assimilation, tree growth, and carbon storage in plants and soil, including growth of wood that is harvested and converted into products such as paper and timber (Pugh et al., 2019; Friedlingstein et al., 2020). Reducing net CO_2 emissions is the central priority for achieving climatic stability in Europe by the target date of 2050 (European Green Deal). A key research question is whether current or future forest management practices can provide a concrete, cost-effective toolset for enhanced carbon storage at the ecosystem and landscape scales (Kauppi et al., 2001; Pussinen et al., 2002; Nolè et al., 2015; Tong et al., 2020) as well as greater wood harvesting for material substitution purposes (Leskinen et al., 2018; Howard et al., 2021).

European (EU) forests have been shaped by centuries of human activities that affect their present potential for carbon storage (Nabuurs et al., 2008). In the EU today, *circa* 165 Mha of forests are managed in ways that drive a net uptake of about 286 Mt CO_2 year^{-1} (Grassi et al., 2017, 2021). Past management strategies were designed to maximize yield of wood products (Leslie, 1966) rather than total biomass production or carbon storage (Tahvonen, 2016). Recent EU-level policies are directed toward sustainable and climate-resilient forests (MCPFE 1990-2003; Forest Europe 2010; Forest Europe 2015; EU Forest Strategy 2015; State of Mediterranean Forest 2018, 2020; New EU-Forest Strategy; 2021; EU Adaptation Strategy 2021) to secure ecosystem services (including carbon sequestration for climate change mitigation) under changing climatic conditions (Churkina et al., 2020; Favero et al., 2020). However, substantial uncertainty remains about the effectiveness of current management practices to maintain the current carbon sink under ongoing climate changes,

and their ability to enhance carbon storage in the future, because classical silvicultural

schemes were developed in the context of past environmental conditions (Bellassen et al.,

2011). In addition, in recent decades European forests achieved increased productivity that

exceeded harvesting rates (Ciais et al., 2008; State of Europe's forests 2020). These

productivity increases resulted from the combinations of several factors: changes to forest age

distribution, climate change (e.g. lengthening of the growing season; Peano et al., 2019),

increased atmospheric CO_2 concentration (stimulating photosynthesis through 'CO$_2$-

fertilization'; Chen et al., 2022), increased nitrogen deposition (stimulating growth through

nitrogen fertilization; Etzold et al., 2020; Chen et al., 2022) and forest management

(Bellassen et al., 2011; Piao et al., 2020; Walker et al., 2021). But these positive trends may

not continue at rapid rates in the future and could be offset by increased disturbance

frequency and severity (Seidl et al., 2017; Senf and Seidl, 2021a), raising concerns that recent

harvesting rates may be approaching, or even exceeding, net tree growth rates in temperate

and boreal European forests (Nabuurs et al., 2013; Ceccherini et al., 2020; Schulze et al.,

2020; State of Europe's forests 2020). Moreover, recent positive trends in gross primary

productivity (GPP; photosynthetic assimilation of atmospheric CO_2) and plant growth in the

northern hemisphere might not be sustained in the future if the CO_2 fertilization effect is not

persistent (Körner, 2005; Walker et al., 2021; Wang et al., 2021) or if they are counteracted

on by increased heat stress and drought (Yuan et al., 2019; Grossiord et al., 2020; Wang et

al., 2020), disturbance-related tree mortality (McDowell et al., 2020; Senf and Seidl, 2021b;

Gampe et al., 2021; Hlasny et al., 2021), or by tree-age-related effects on net primary

productivity (NPP; the balance of photosynthetic CO_2 assimilation and plant respiratory CO_2

release) (Ryan et al., 2006; Zaehle et al., 2006; Luyssaert et al., 2007; Tang et al., 2014; Pugh

et al., 2019). Hence, a reduction in the forest net carbon sink capacity and its transition to a

CO_2 source could be approaching (Duffy et al., 2015; Peñuelas et al., 2017; Wang et al.,

2020). This might reduce both the potential for forests to sequester additional carbon and

sustainably produce wood products.

Forest carbon cycle and productivity models are used to investigate and project the effects of

climate change and management scenarios on forest productivity across a range of spatial and

temporal scales to support science-driven policymaking (Temperli et al., 2012; Vacchiano et

al., 2012; Reyer et al., 2017; Maréchaux et al., 2021). This includes design of alternative

management strategies to support climate change mitigation policies (Maréchaux et al., 2021;

De Marco et al., 2022).

This study aimed to inform the debate about the role of forest management practices in

sustaining high forest productivity under future climate conditions. To that end, a validated

process-based modeling approach was designed to quantify controls on CO_2 uptake and C

storage in a composite matrix of managed forests, taking into account how combinations of

climate change and management scenarios may affect those controls. Specifically, we

questioned whether, relative to business-as-usual (BAU) management scenarios, alternative

forest management practices could maximize NPP while at the same time maintaining and/or

increasing potential Carbon Woody Stocks (pCWS; i.e., when no decay of harvested wood

products occurs) in response to a range of climate change scenarios.

2. | Materials and Methods

2.1 | 3D-CMCC-FEM Model

2.1.2 | Model description

The 3D-CMCC-FEM v.5.5 (Three Dimensional - Coupled Model Carbon Cycle - Forest

Ecosystem Model; Collalti et al., 2014, 2016; Marconi et al., 2017; Engel et al., 2021;

Mahnken et al., 2022) simulates daily GPP through the Farquhar-von Caemmerer-Berry

photosynthesis model (Farquhar et al., 1980), modified for sun and shaded leaves (de Pury

and Farquhar, 1997) and acclimation to temperature history (Kattge and Knorr, 2007). Plant

respiration (R_a) is simulated explicitly and partitioned into growth (R_g) and maintenance (R_m)

fractions as in the growth-and-maintenance-respiration paradigm (Amthor, 2000; McCree,

1970; Thornley, 2000). R_m is computed, for each functional-structural tree C pool (i.e. live

wood, leaves and fine roots), using a temperature-acclimated Q_{10} relationship (for details on

thermal acclimation see Tjoelker et al., 2001; Atkin and Tjoelker, 2003; Smith and Dukes,

2012; Collalti et al., 2018) and a N-based maintenance respiration (m_R) rate for living tissues

of 0.218 g C g N^{-1} day^{-1} (Ryan et al., 1991; Amthor and Baldocchi, 2001; Oleson et al., 2013;

Drake et al., 2016Collalti et al., 2016, 2020a). R_g is considered a fixed fraction (i.e. 30%) of

the remaining C once tissue R_m is accounted for and removed from GPP. The sum of daily R_g

(if any) and R_m gives R_a. Daily NPP is then GPP less R_a. Allocation of NPP among tree C

pools is performed daily, with preference to a non-structural carbon pool (NSC, i.e. storage in

starch and sugars), which is used directly to fuel R_m, up to a minimum NSC threshold level.

The minimum NSC-threshold level is a fraction (a model parameter) of the live wood C-

content (Collalti et al., 2020a). Once (and if) the minimum NSC threshold is reached, C is

allocated preferentially for biomass growth for the different tree structural C-pools depending

on the phenological phase as formerly described in Collalti et al. (2016). The only

phenological phase during which NSC has no priority in allocation is during bud break

(D'Andrea et al., 2020, 2021), when recent GPP is completely allocated for growth of leaves

up to a maximum annual leaf area index (LAI, m^2 m^{-2}), which is computed at the beginning

of each year of simulation through the pipe-model (Shinozaki et al., 1964; Mäkelä, 1997),

and growth of fine roots. This NSC allocation scheme reflects a quasi-active role of NSC,

with NSC usually having priority over growth of new structural tissues, as described by Sala

(2011), Merganičová et al. (2019) and Collalti et al. (2020a). This implies that any

asynchrony between C-demand (i.e. respiration and growth) and C-supply (photosynthesis) is

buffered by changes to the NSC pool. When NSC approaches zero and cannot be refilled by

GPP, carbon starvation occurs, and tree death is simulated. This overall C-allocation scheme

in the 3D-CMCC-FEM model follows the functional balance theory of allocation, similarly to

other models (Merganičová et al., 2019). Age-related mortality, carbon starvation, and a

background mortality (i.e. the as-yet unexplained mortality), represent the different types of

mortality simulated by the model; the last one is turned off when forest management is

applied. An in-depth description of the model's underlying characteristics, responses to

climate change, and sensitivity to model parameter values, as well as model limitations, is in

Collalti et al. (2019, 2020a, and references therein).

2.2.2 | Forest management routine

Historically, the majority of European forests were actively managed and the share of

coniferous species, particularly Scots pine and Norway spruce, was favored due to their high

growth increment rates (Naudts et al., 2016). Management techniques (thinning and clear-

cutting) aimed at maximizing productivity and reducing losses often resulted in even-aged,

mono-specific forest stands (Campioli et al., 2015, and references therein; State of Europe's

forests 2020; Figure S1 in Supplementary Material). In such stands, carbon pools and fluxes

strongly depend on rotation lengths (tree age-class distribution), thinning interval, and

thinning intensity (Kaipainen et al., 2004; Nabuurs et al., 2008).

In this study we varied these three key management factors: thinning intensity, thinning

interval and rotation age (following Reyer et al., 2020). Thinning intensity is represented in

the model by the percentage of stand basal area to be removed based on the total stand basal

area. Thinning interval is the number of years between two consecutive operations. Rotation

age is the stand age at which a full harvest occurs, after which the stand is replanted with

saplings of the species adopted in the Inter-Sectoral Impact Model Intercomparison Project

(ISIMPI, https://www.isimip.org, Warszawski et al., 2014) protocol. The model benchmark

was the BAU forest management scheme for the most common European species as

described in Reyer et al. (2020) and Mahnken et al. (2022) and applied in three contrasting

forest stands as in Collalti et al. (2018).

2.2. | Sites, data and experimental design

The model was parameterized for, and simulated C fluxes and tree growth in three even-aged,

long-monitored, managed European forest sites which are part of the Fluxnet network

(Pastorello et al., 2020), the ISIMIP initiative and the PROFOUND database (Reyer et al.,

2020; Mahnken et al. 2022): (1) the temperate European beech (*Fagus sylvatica* L.) forest of

Sorø, Denmark; (2) the Norway spruce (*Picea abies* (L.) H. Karst) stand of Bílý Křiž in

Czech Republic, and (3) the boreal Scots pine (*Pinus sylvestris* L.) forest of Hyytiälä, Finland

(Table 1). These sites were selected because they represent the dominant forest types in

Europe covering the temperate and boreal climatic zones (Figure S1),which contain more

than 50% of European forest biomass (Avitabile et al., 2020), and their management best

corresponds to 'the intensive even-aged forestry' as defined by Duncker et al. (2012).

Daily weather values input to the model for each site were derived from the climate scenario

data from the ISIMIP Fast Track initiative based on the Climate Model Intercomparison

Project 5 (CMIP5) in which five Earth System Models (ESMs; i.e.: HadGEM2-ES, IPSL-

CM5A-LR, MIROC-ESM-CHEM, GFDL-ESM2M, and NorESM1-M) were driven by four

Representative Concentration Pathways (RCPs) of atmospheric greenhouse gas concentration

trajectories, namely RCP 2.6, RCP 4.5, RCP 6.0, and RCP 8.5 (Moss et al., 2010; van Vuuren

et al., 2011). Daily meteorological forcing-data for each site used by 3D-CMCC-FEM were

available as bias-corrected/downscaled variables (air temperature, precipitation, solar

irradiance) and as non-corrected variables (relative humidity) according to Hempel et al.

(2013). The RCP atmospheric CO_2 concentration values for the period 2016 to 2500, based

on Meinshausen et al. (2011) as described in Reyer et al. (2020), were used to drive the

biogeochemical photosynthesis model with values varying at the end of this century from

421.4 µmol mol^{-1} (RCP 2.6) to 926.6 µmol mol^{-1} (RCP8.5).

2.2.1. | Virtual stands

Given that the European forested area is composed of a mix of differing-aged stands, and

since forest C cycle processes may respond differently to climate factors at different ages

(e.g. Collalti and Prentice, 2019; Collalti et al., 2020b; Huber et al., 2018, 2020; Migliavacca

et al., 2021), we developed a Composite Forest Matrix (CFM) consisting of a mixture of

stands of different age, structure and associated biomass. Starting from the real stands, we

generated a prescribed number of virtual stands in order to obtain representative model

outputs of a larger set of different age-classes (with their associated forest attributes) to cover

an entire rotation period (~140 years, depending on species), similar to the approach in in

Bohn and Huth (2017). Because 3D-CMCC-FEM, in accordance with experimental evidence,

is sensitive to the stand structure (see Collalti et al., 2019, 2020a), this procedure allows a

robust assessment of effects of management practices across the full range of applicable stand

ages . A CFM framework was then created by running at each site the model from 1997 to

2199 (to cover the entire rotation length for each species) under a contemporary climate

scenario (no climate change), consisting of de-trended and repeated cycles of 1996-2006

historical weather, with fixed atmospheric CO_2 concentration of 368.8 µmol mol^{-1} and BAU

management practices. From each of these simulations data needed to reinitialize the model

at every *rotation length*/10 were extracted (Figure 1). Thus, in total, ten virtual stands

representing different age classes of the composite matrix were selected and included into the

CFM. The management scenario analysis was carried out on this set of forest virtual stands.

2.2.2. | Alternative management (AM) schemes

For each real and virtual stand at each of the three sites we applied 28 management scenarios:

the BAU and 'no management' (NO-MAN: stands allowed to grow without thinning or

harvesting) schemes plus 26 alternative management schemes. These alternative forest

management scenarios represent all the possible combinations of two thinning *intensities*, two

thinning *intervals*, and two *rotation* durations than differ from those in the BAU scenario.

The schemes were grouped (Tables 1 and Table S1) into combinations of: (1) '*more*

intensive' ('AM+'), where at least one out of the three management variables reflect an

intensified management case relative to BAU (e.g. higher thinning intensity and/or shortened

interval and/or shortened rotation length than BAU), and the other one or two (or no)

variables are kept as in BAU; (2) '*less intensive*' ('AM–'), where at least one variable reflects

lower thinning intensity and/or prolonged interval and/or prolonged rotation length,

compared to the BAU case; and (3) '*mixed schemes*' ('MIX'), where at least one

management variable was more intensive and at least one management variable was less

intensive than the BAU scheme.

2.2.3 | Model runs and evaluation

The starting year for all simulations was 1997, consistently with the availability of measured

stand carbon flux data used for the model initialization and evaluation. After creation of the

virtual stands, based on de-trended historical weather time series for period 1996-2006,

climate change simulations were conducted for the period 2006 to 2100. By considering all

combinations of management and climate change scenarios from the different ESMs climate

forcings, 6,160 simulations were performed at each site (i.e., 5 ESMs * 4 RCPs * 11 stands *

28 management schemes) for a grand total of 18,480 model runs.

To gauge model sensitivity to the study variables we organized the analysis according to a

factorial design (Mason et al., 2003; Collalti et al., 2018) across a matrix of Stand, ESM and

RCP, generating seven possible combinations from each factor. This was used to identify the

most influential factor the modelled carbon cycle variables GPP, R_a, NPP, carbon-use

efficiency (CUE = NPP/GPP), net woody productivity (NPP_{woody}), and potential Carbon

Woody Stocks. The model was evaluated using 1997-2005 annual GPP and NPP_{woody} data for

Sorø and Hyytiälä and 2000-2005 for Bílý Křiž by comparing simulated GPP against eddy

covariance estimates (Pastorello et al., 2020), and compared modeled wood growth against

measured NPP_{woody} (Principal Investigator's site for Hyytiälä and Bílý Křiž under *personal*

communication, and Wu et al., 2013 for the site of Sorø) and stem diameter at breast height

(DBH; Reyer et al., 2020; Mahnken et al., 2022). Subsequent years were excluded from the

model evaluation since the scenario period in the ESMs started in 2006, and hence was driven

by different atmospheric CO_2 concentration trajectories after 2006.

2.3 | Effects of climate change and management on carbon fluxes and biomass

As carbon fluxes do not scale linearly to stocks (Schulze et al., 2020) data analyses focused

separately on the variables NPP and pCWS, the sum of standing and previously harvested

woody stocks. Net primary productivity can be a good proxy for forest carbon sink processes

(Sha et al., 2022), and the net biomass input to forest ecosystems (Trotsiuk et al., 2020), with

decomposition (decay, or heterotrophic respiration) processes representing the active carbon

source process. Net primary production is a dynamic balance between photosynthesis and

plant respiration, which respond separately and/or in combination to a range of climatic

factors and, in managed forests, to management practices (Collalti et al., 2020a; 2020b).

These age-dependent responses are generally not prone to *in situ* quantification over long

periods, especially for climate change issues, hence raising the need for process-based

modeling. Harvested wood products are considered here without decay; hence we aim at

evaluating only the potential maximum attainable total woody standing stocks under a wide

spectrum of possible management schemes without any consideration of the turn-over of

harvested wood products. Data were averaged over the emission scenario simulation period

2006-2099 and aggregated over virtual and real stands and over ESMs but distinguished

between RCPs. In spite of site-specific differences in magnitude of response to the different

management schemes and climate (see Supporting Material) the main emerging pattern with

increasing intensity of management intervention was of similar magnitude. For this reason,

data were also aggregated over sites. Therefore, alternative management practice results are

presented as aggregated according to the groups AM+ and AM– to highlight patterns of

directional changes when moving from the current management schemes toward a more

intensive or a less intensive scenario.

3. | Results

3.1 | Model evaluation

The 3D-CMCC-FEM model was evaluated at the three sites separately and at different

temporal scales with robust data for both the carbon fluxes GPP and NPP_{woody}, and the

structural variable average stand DBH (Figures S2-S4 in Supplementary Material).

Simulations forced with both observed local daily weather and with an ensemble of outputs

from climate models for the contemporary period compare well with the eddy-covariance

based estimated daily GPP values at the sites of Sorø (Root Mean Square Error, RMSE =

2.15 g C m^{-2} day^{-1} with local climate, and RMSE = 2.98 g C m^{-2} day^{-1} with the ensemble

across ESMs forcing; r > 0.86), Hyytiälä (RMSE = 1.48 g C m^{-2} day^{-1} with local climate,

and RMSE = 1.91 g C m^{-2} day^{-1} with the ensemble across ESMs forcing; r > 0.78) and Bílý

Křìž (RMSE = 2.07 g C m^{-2} day^{-1} with local climate, and RMSE = 2.69 g C m^{-2} day^{-1} with

the ensemble across ESMs forcing; r > 0.67) (see Supplementary Material Table S2-S3).

Modelled annual GPP was also consistent with site measurements at Sorø (1665 ± 171 g C

m^{-2} year^{-1} and 1585 ± 190 g C m^{-2} year^{-1} modelled from observed and modelled climate vs.

1731 ± 184 g C m^{-2} year^{-1} measured; here and elsewhere, ± denotes one standard deviation)),

Hyytiälä (894 ± 57 g C m^{-2} year^{-1} and 871 ± 52.6 g C m^{-2} year^{-1} modelled from observed and

modelled climate vs. 1028 ± 50 g C m^{-2} year^{-1} measured), and Bílý Křìž (893 g C ± 252 g C

m^{-2} year^{-1} and 893 ± 222 g C m^{-2} year^{-1} modelled from observed and modelled climate vs.

1024 ± 354 g C m^{-2} year^{-1} measured). Similarly, modelled values were similar to measured

values of tree woody pools (i.e. NPP$_{woody}$) at Sorø (351 ± 61 g C m^{-2} year^{-1} and 275 ± 63 g C

m^{-2} year^{-1} modelled from observed and modelled climate vs. 346 ± 36 g C m^{-2} year^{-1}

measured) and Bílý Křìž (442 ± 79 g C m^{-2} year^{-1} and 405 ± 36 g C m^{-2} year^{-1} modelled

from observed and modelled climate vs. 380 ± 38 g C m^{-2} year^{-1} measured). The NPP$_{woody}$

simulated at Hyytiälä was not quite as close to available data (317 ± 21 g C m^{-2} year^{-1} and

290 ± 24 g C m^{-2} year^{-1} modelled from observed and modelled climate vs. 228 ± 23 g C m^{-2}

year^{-1} measured). Mean DBH increase, which was only qualitatively compared, is slightly

underestimated at the site of Bílý Křìž (Figure S3). Comparisons of literature data for NPP,

R$_a$ and CUE with modelled values are in Table S3. Notably, results generated by3D-CMCC-

FEM forced with the EMSs' climates are close to those generated when forced with observed

climate data in the evaluation period.

3.2 | Less intensive management vs. BAU

Simulated average NPP in the less intensive management scenario group (i.e. AM−) is close

to the reference BAU values, ranging from 495 g C m^{-2} year^{-1} (−1.2%; here and elsewhere,

percentages refer to changes compared to BAU) to 525 g C m^{-2} year^{-1} compared to the range

from 502 g C m^{-2} year^{-1} to 542 g C m^{-2} year^{-1} for RCP 2.6 and 8.5 under BAU, increasing

slightly with warming and increased atmospheric CO_2 concentration in the early years of the

study period. Changes were larger toward the end of the century and without significant

differences across RCPs (Figure 2 and Figure S4, Table 2). Simulated pCWS values increase

steadily along the simulation time for all the alternative management scenarios, with time-

averaged values in the range 180-199 t C ha^{-1} compared to 193-199 t C ha^{-1} for RCP 2.6 and

8.5 under BAU, respectively (Figure 3 and Figure S4, Table 2). Relative to BAU, the ratio

pCWS/NPP decreased with less intensive management and with no management (Table 4;

results for each of the RCPs scenarios and all the alternative management options combined

are reported separately across the sites in the Supplementary Material Table S4).

On average, pCWS increased with less intensive management only in the case of a prolonged

rotation period under RCP 8.5, while NPP declined (~6%) in response to an extended rotation

and reduced thinning intensity under RCP 8.5. Conversely, pCWS shows the greatest

reduction with −13.9% when both thinning intensity and regime only are set to simulate

decreased management intensity under RCP 8.5 (Table S5). In summary, AM− simulations

generally reduced pCWS with little effect on NPP. For NPP some intra-site differences exist,

with the representative forest of Hyytiälä showing higher NPP values when applying a less

intensive management than the reference BAU. Differences between the emissions scenarios

were small.

3.3 | More intensive management vs. BAU

Across all stands and ESMs the AM+ simulations reduced NPP about 30% relative to the BAU scenario (Figure 2 and Figure S4, Table 2). Conversely, AM+ reduced pCWS only 4-5% relative to BAU across the climate change scenarios (Figure 3 and Figure S4, Table 2). (Results for each of the RCP scenarios and under all the alternative management options combined are reported separately across the sites in the Supplementary Material (Table S4 and S5).) The decreases in NPP with AM+ were larger than the decreases with AM- across RCPs, with even a slight increase in NPP in two of the seven AM- scenarios under RCP 2.6 (Table S5). Conversely, pCWS shows a net gain with +4.1%, compared to the BAU, which applies under an increased thinning intensity under RCP 8.5, while the greatest reduction in pCWS with −13.9% is modelled when both thinning intensity and frequency are increased and a shortened rotation turn are simulated under RCP 8.5 (Table S5). Generally, both NPP and pCWS results from the AM+ schemes are, on average, close to or lower than those from the reference BAU and this holds across the three representative forests (Figures S10-12). When considered during the early portion of the study period, the differences in NPP and pCWS between AM+ management scenarios and BAU are narrower. In particular, an average gain of 5% in pCWS could be achieved over a wide range of AM+ schemes, but at the expenses of an average NPP change of −30% (Figure S6).

For both NPP and pCWS, the differences across RCP scenarios are much smaller than across the different management scenarios indicating that management may be more important than near-term climate change in controlling in the C cycle of European forests.

3.4 | No management vs. BAU

The NPP values in the NO-MAN scenario are, on average, lower than the reference BAU scenario varying from −14.5% (RCP 2.6) to −19.5% (RCP 8.5) (Figure 2, Table 2). However,

a site-specific variability in the NPP response to the management scenarios applied exists,

with differences between NO-MAN and BAU options ranging from 9.0% to –41.7%, for

Hyytiälä and Bílý Křìž respectively, both under the warmest emission scenario (Table S4).

Differences between NO-MAN and BAU become more evident during the simulation period

and across RCPs scenarios, with the mean NPP value stabilizing or slightly increasing under

the BAU (and AM–) option over time. Conversely, in the NO-MAN scenario the values

steadily decreased (Figure 2).

The simulated pCWS, which is represented only by standing biomass (i.e., no harvested

wood) in the NO-MAN scenario was, on average, lower than in the BAU scenario, with

differences of order –30.0% (from –21.6% to –40% across the different sites). During the

simulation period, pCWS values in the NO-MAN option first increase slightly at the

beginning of the simulation and then decrease significantly toward the end of the century

(Figure 3, Table 2). The NO-MAN case returns the lowest average amount of total woody

stocks under every emission scenario (Figure 4 and Table 2).

3.5 | Mixed management alternatives and the factorial analysis

A mixed combination of management schemes (namely 'MIX') was analyzed for all possible

combinations (Figure 5 and Supplementary material Table S1, S4 and S5). There were no

'MIX' options that simultaneously increased both in NPP and pCWS compared to the BAU

scenarios. Values for NPP range from –1.38%, with a prolonged thinning regime and

rotation, and an increase in thinning intensity; to –58.4% under a prolonged thinning intensity

and rotation period but with a reduced thinning regime, both under RCP 2.6 (Table S5).

Similarly, with the same management schemes, pCWS declined 16.1% under RCP 6.0, when

compared to BAU, but increased 5.7% under RCP 8.5 in response to increased rotation length

in combination with increased thinning intensity.

The factorial analysis performed over all the main carbon fluxes and stock variables produced

by the model and separated by site, indicates that a significant fraction of the total variability

of the key carbon flux variables was driven by the stand factor, i.e. the forest structure as

generated by different management schemes, affecting the distribution of age classes,

including different above- and below-ground biomass (Figure S6 and Table S6 in

Supplementary Material).

4. | Discussion

4.1 | Limited leeway to increase carbon uptake and woody stocks with alternative

management scenarios

The variables NPP and pCWS represent two sides of the same coin since the former (i.e.

NPP) represents the short- to medium-term active carbon sequestration capacity while the

latter is the sequestered (and maintained) carbon over the medium- to long-term. The present

simulations clearly indicate that even under future climate change scenarios the options are

limited for managing forests to help trees maintain their carbon sequestration potential and

enhance their ability to cope with increasingly stressful conditions, while in addition

increasing their productivity compared to the no-management option. Reducing tree density

allows individual trees to benefit from less competition for potentially limiting resources such

as light and soil moisture, driving their potential to increase photosynthetic activity and

growth rate (Zeide et al., 2001). In addition, stand rejuvenation through harvesting and the

replanting of saplings, with less 'respiring' (live) biomass per unit of photosynthetic leaf area

(because of a shift toward younger stands), and combined with the fertilization effects of

increased atmospheric CO_2 concentration can potentially drives to more productive and

efficient forests. This is mirrored in the observed capability of trees to partition more of the

photosynthetically assimilated carbon into new woody biomass rather than into nonstructural

carbon pools (Campioli et al., 2015; Pappas et al., 2020; Collalti et al., 2020a; Huang et al.,

2021). This aspect is reflected in the increasing Harvested Wood Products (HWP; Figure S7)

over time with more frequent thinning, reduced tree density, replacement and presence of

younger forest stands that potentially can remain in the system as standing biomass over the

long-term (Figure S8). Overall, the model projects on average the highest pCWS under the

BAU management scheme, even in the future.

The potential to extract more wood and more often, i.e. to shorten the harvest interval, and at

the same time maintain at least the current forest biomass, depends on NPP under the

different scenarios. We found, however, that BAU management remains the most favorable

scheme under future environmental conditions and might already be a close-to-optimum

management approach for different RCP scenarios (Figure 4) and across the individual sites

(Figures S9-S11). This is an endorsement of past research arriving at today's management

practices and a coincidence that today's most favorable scheme might also be most favorable

in a future altered climate. With more frequent harvesting and replanting and increasing

intensity of intervention compared to the benchmark BAU, the NPP is not shown to increase

any further under any RCP scenario, in spite of an average younger and, in theory, more

productive forest stand. The net growth rate does not compensate for the increased fellings

(tree harvesting), while in parallel there is a limited yield in terms of increased woody carbon

stocks, as reflected in a low standing biomass that is likely a sign of a critically low tree

density. Albeit the BAU reference benchmark is already an intensive management approach

with tree fellings as a percentage of net annual increment of 84%, 77% and 101% for Czech

Republic, Finland and Denmark, respectively, as reported for 2005 (State of Europe's forests

2020), the first year of our RCP-based climate change response simulations. Similarly,

Pussinen et al. (2009) showed that increasing the total harvested products led to a decrease in

both NPP and forest standing biomass in some European areas. The difficulties associated

with simultaneously increasing both forest standing biomass and harvested wood products

were shown in the seminal modeling study of Thornley and Cannell (2000).

An important factor contributing to the apparent lack of significant differences in forest

responses across RCPs scenarios, compared to the differences across different management

schemes, might come from the combination of counteracting key drivers of plant physiology

(e.g. lengthening of the growing season by warming and, in parallel, an increased

maintenance respiration rate from that same warming) which are considered in the model

despite temperature acclimation. Although experimental evidence for the CO_2 fertilization

effect on plant carbon accumulation is strong and is typically predicted by vegetation models

albeit with different degrees of certainty, the probability for its persistence into the longer-

term future is a hotly debated issue (Nabuurs et al., 2013; Habau et al., 2020; Wang et al.,

2020; Gatti et al., 2021; Walker et al., 2021). The biochemical model of photosynthesis used

here (Farquhar et al., 1980) includes a nonlinear saturating response to CO_2, yet, other

environmental drivers, such as temperature, vapor pressure deficit (which scales

exponentially with warming) and water availability were shown to interact to down-regulate

the positive CO_2 effect on GPP (Grossiord et al., 2020). Data at the biome scale (see

Luyssaert et al., 2007) indicated a potentially high sensitivity of plant respiration to warming

that may stabilize NPP over a temperature threshold with no further gains. Warming in low-

temperature-limited forest biomes would be expected instead to have a positive effect on

annual GPP and NPP (Henttonen et al., 2017; Sedmáková et al., 2019). However a warming-

induced increased respiration cost might curb these trends and even offset a positive GPP

and/or NPP response to increasing atmospheric CO_2 concentration, as indicated by some

other modeling and experimental studies (Way et al., 2008; Gustavson et al., 2017; Collalti et

al., 2018; but see Reich et al., 2016). For example, Mathias and Trugman (2021) showed a

potential future unsustainable growth for boreal and temperate broadleaved forests, with the

net overall effect of decreased NPP. Other studies already indicated that combined impacts of

warming and increasing atmospheric CO_2 concentration might cause forests to grow faster

and mature earlier but also to die younger (Kirschbaum, 2005; Collalti et al., 2018, 2019).

With the increasing standing biomass and accumulation of more respiring tissue in older

trees, plant respiration might increase more quickly than GPP, as the canopy closure would

be reached earlier, capping GPP when all available solar radiation is intercepted, but with

sustained respiratory needs. The use here of many virtual stands of different ages in our

simulations might have (realistically) compensated for local stand-age effects on biomass and

stand structural responses to climate change that would be counteracting across a landscape.

To the extent that this is true, the simulated patterns obtained would be related to effects of

climate, forest management, and their multiple combinations only.

Ultimately these simulations indicate that increasing the harvest rate and at the same time

keeping the fellings below the growth rate will be challenging. As such, the possibility of

simultaneously increasing both carbon sequestration rate and tree carbon (standing biomass)

storage capacity while managing forests in a sustainable way may be very limited. A steady

intensification or intervention frequency – alone or in combination – compared to the

business-as-usual scheme might come at the price of a substantial loss of primary

productivity. While the amount of potential harvested woody products still would be

significant, we would *de facto* end up reducing the active forest carbon sink and thus the

forest's potential to assimilate and sequester CO_2 from the atmosphere.

4.2 | Role of forest management in the context of climate change

In the context of climate uncertainty and because of policy intentions, management practices

may no longer prioritize productivity only – which traditionally includes rotation times being

adjusted to maximize value of timber – without preserving the forest carbon sink and

ensuring the long-term functioning of forests and the continued provision of their many

ecosystem services (Krofcheck et al., 2019). The selection of adaptive management practices

has been suggested as a mechanism to potentially enhance the climate change mitigation

potential of forest ecosystems (Tahvonen, 2016; Yousefpour et al., 2017). Our model results

highlight, for Central and Northern temperate and boreal European forests, the importance of

forest structure to sustain or even enhance productivity and carbon storage. This in turn

indicates that management practices may be quantitatively as important as future climate and

atmospheric CO_2 concentration in regulating the carbon sink strength of forests, which is in

line with some previous modeling studies (Garcia-Gonzalo et al., 2007; Akujärvi et al., 2019;

de Wergifosse et al., 2022). These simulations indicate that silvicultural practices included in

the model would persist as key factors in the regulation of carbon sequestration through the

end of this century under any of the CMIP5 RCP scenarios. In accordance with the modeling

study of Kindermann et al. (2013), our results indicate the need to sustain NPP rather than

maximizing forest carbon stocks. Our results also point out, however, a narrow operational

space surrounding the BAU scheme which can be designated as near-optimal over a wide and

diversified portfolio of alternative management schemes across the broad range of

RCP/ESM-based climate change scenarios. Conversely, other studies (Garcia-Gonzalo et al.,

2007; Luyssaert et al., 2018) showed that harvest intensity should be loosened in order to

maximize the carbon sink which would lead to reductions in the wood harvesting rates.

However, under adverse climate change effects, reductions in wood harvesting rates may

correspond with declining carbon sequestration.

Even considering changes in species composition by replanting better-suited species under a

climate change adaptive framework, Schelhaas et al. (2015) found a reduction of the net

increments, although these did not affect wood product amounts. On the other hand, Pussinen

et al. (2009), found that under an increased felling rates regime (i.e. high harvested products

amount) maintaining the current forest standing biomass was possible even under future

climate scenarios. However, in their study NPP was simply assumed to increase

proportionally to the temperature, driving increased growth rates at boreal and temperate

forest sites. The present simulation study used a more realistic process-based representation

of climate change impact on forest growth. This reveals instead a more modest, or even

unfavorable, effect of climate change despite a CO_2 fertilization effect on NPP. Others have

suggested that past and/or future climate change did, or could, negatively affect NPP (Reich

and Oleksyn, 2008; Bastos et al., 2020) in a range of forested and non-forested ecosystems

through increased frequency and/or magnitude of large-scale disturbances (e.g. heat waves,

windstorms, weather-based pest outbreaks), with significant variation in effects in different

ecosystems or forest types and locations (e.g. Thom et al., 2017; Nabuurs et al., 2019;

McDowell et al., 2020; Senf and Seidl, 2021b; Gampe et al., 2021). Should such increases in

climatic extremes (e.g. drought and heatwaves) and disturbances (pests outbreaks, wildfires,

storms) occur -- and negatively affect the carbon sink capacity of a significant fraction of

European forests -- the robustness of the BAU management scheme specified in our

simulations might be questioned with an adaptive set of management options designed, tested

and ultimately put into place (Yousefpour et al., 2017). However, under the unmanaged-

forest scenario, our simulations resulted in significant and steady decline in NPP through the

year 2099 when averaged over all the climate change scenarios considered.

4.3 | Further consideration and outlook

In this study we found that intensification of forest management practices might lead to

declines in both CO_2 assimilation and the total woody stocks.

The focus was on a subset of forest types that currently play a significant role in the European

carbon balance and wood product markets by representing more than 50% of the EU standing

biomass (Avitabile et al., 2020).

In the current modelling framework, our results refer to potential values of carbon woody

biomass, with no estimated decay for harvested wood products.

In this context, our aim was primarily to quantify the efficiency of forest trees to actively

sequester and store carbon from the atmosphere and, thus, defining their potential

contribution to mitigation policy requirements (see e.g. Valade et al., 2017).

Pure stands were selected for this study as they represent most of the managed stands in

Europe, and no species migration under climate change was included, even in the no-

management scenario. However, we highlight that the overall climate change mitigation of

the forestry sector is further influenced by carbon sequestration in harvested wood products

(Brunet-Navarro et al., 2016) as well as energy and material substitution effects of wood

products, which is depending on dynamic changes in energy portfolios and greenhouse gas

emissions in other sectors and thus associated with substantial and additional uncertainty (e.g.

Leskinen et al., 2018; Howard et al., 2021; Soimakallio et al., 2021).

Since the rates of potential species migration and/or replacement may be incompatible with

the expected rates of climate change, at least for the high RCP scenarios (Settele et al., 2014),

the effectiveness of tree species change as a mitigation factor in unmanaged forests could be

limited. Future research with validated dynamic vegetation models is paramount for it might

provide needed insights into European forest carbon sequestration potential across a broad

geographic extent through mechanisms of seed dispersal, species migration and up-scaling

techniques (Fritsch et al., 2020).

In addition, despite simulating a broad and diversified range of different silvicultural

interventions aimed at testing the possibility to increase NPP and carbon stocks, additional

studies will be crucial to address changes in species selection (i.e. genetic selection) and

assisted species migration in adaptive forest management.

Finally, the focus of the present work was to quantify the capacity of trees to sequester

carbon in woody biomass and in turn to provide wood products in the face of expected

climate changes. A life-cycle analysis of the harvested wood products was not included, and

in essence implicitly assumed to be unaffected by climate change.

5. | Conclusions

To our knowledge this is one of the first modeling studies that systematically analyses, over a

wide range of scenarios, the possibilities and limitations of altering forest management

practices to achieve the twofold objective of maximizing forest C-sequestration capacity

while concomitantly maintaining and/or increasing pCWS in the face of future climate

change.

Even though the analysis was confined to three sites, the representativeness of those sites to

other European forests and the general consistency of the results indicates a broader

significance and applicability. In particular, the differences between AM+ and AM−, each

relative to BAU, indicates a relatively consistent response across species within different

climate change scenarios and management options.

Our results indicate that the scope to meet the above twofold objective may be limited,

because business-as-usual management practices may already be nearly optimal in terms of

carbon uptake, sequestration and storage, as testament to positive outcomes of previous

silvicultural research. A general conclusion of the modelled results is that NPP and/or pCWS

are likely to decline, relative to BAU, with any significant change in forest management practices.

Beside the economic value of the extractable wood and the potential for energy and material substitution, it is today crucial for EU countries to preserve forest's functionalities under the pressure of the rapidly changing climate conditions, in order to maintain the climate mitigation potential and the supply of wood products and many ecological goods and services. Forest management based on scientific principles remains a valuable tool for local, regional and global strategies to optimize the forest carbon sinks and provide desired products under a varying climate. However, the extensive modeling in this study and related research does not support an optimistic view that changes to management practices in European forests can be relied on to significantly increase carbon uptake and storage while increasing wood production above current rates.

Acknowledgements

We are thankful to A. Ibrom, P. Kolari and J. Krejza for providing us data for Sorø, Hyytiälä and Bílý Kříž sites. We thank J. Holder, B. Waring, H. Bugmann, L. Perugini, S. Luyssaert, M. Michetti, and A. Mäkelä for comments on a previous version of the manuscript and/or fruitful discussions. We also thank the ISI-MIP project (https://www.isimip.org/) and the COST Action FP1304 PROFOUND (Towards Robust Projections of European Forests under Climate Change), supported by COST (European Cooperation in Science and Technology) for providing us the climate historical scenarios and site data used in this work. This work used eddy covariance data acquired and shared by the FLUXNET community, including these networks: AmeriFlux, AfriFlux, AsiaFlux, CarboAfrica, CarboEurope-IP, CarboItaly, CarboMont, ChinaFlux, Fluxnet-Canada, GreenGrass, ICOS, KoFlux, LBA, NECC, OzFlux-

TERN, TCOS-Siberia, and USCCC. The ERA-Interim reanalysis data are provided by

ECMWF and processed by LSCE. The FLUXNET eddy covariance data processing and

harmonization was carried out by the European Fluxes Database Cluster, AmeriFlux

Management Project, and Fluxdata Project of FLUXNET, with the support of CDIAC and

ICOS Ecosystem Thematic Center, and the OzFlux, ChinaFlux, and AsiaFlux offices. We

acknowledge the World Climate Research Programme's Working Group on Coupled

Modelling, which is responsible for CMIP, and we thank the respective climate modeling

groups for producing and making available their model output. The U.S. Department of

Energy's Program for Climate Model Diagnosis and Intercomparison at Lawrence Livermore

National Laboratory provides coordinating support for CMIP and led development of

software infrastructure in partnership with the Global Organization for Earth System Science

Portals. A.C. was supported by the EFI-Funded FORMASAM (FORest MAnagement

Scenarios for Adaptation and Mitigation) project; D.D. and A.C. were also supported by

resources available from the Italian Ministry of University and Research (FOE-2019), under

projects 'Climate Changes' (CNR DTA. AD003.474.029); A.C. and G.M. were supported by

LANDSUPPORT Horizon 2020 research and innovation programme under grant agreement

No. 774234. The 3D-CMCC-FEM model code is publicly available and can be found on the

GitHub platform at: https://github.com/Forest-Modelling-Lab/3D-CMCC-FEM). All data

supporting this study are publicly available at the Zenodo repository

(https://zenodo.org/record/6026117#.YgTfXnXMJhE). Correspondence and requests for

additional materials should be addressed to corresponding authors.

623 Credit authorship contribution statement

References

Akujärvi, A., Shvidenko, A., and Pietsch, S. A. (2019). Modelling the impacts of intensifying
forest management on carbon budget across a long latitudinal gradient in Europe.
Environmental Research Letters, *14*(3).

Amthor, J.S. 2000. The McCree–de Wit–Penning de Vries–Thornley respiration paradigms:
30 years later. Ann. Bot. 86, 1-20. https://doi.org/10.1006/anbo.2000.1175.

Amthor, J.S., Baldocchi, D.D. (2001). Terrestrial higher plant respiration and net primary
production. In: Roy, J., Saugier, B., Mooney, H.A. (Eds.), Terrestrial Global Productivity.
Academic Press, San Diego, pp. 33–59.

Atkin OK and Tjoelker M (2003). Thermal acclimation and the dynamic response of plant
respiration to temperature. *Trends in Plant Science*, 8: 343-351.

Avitabile V., Pilli R., Camia A., The biomass of European forests, EUR 30462 EN,
Publications Office of the European Union, Luxembourg, 2020, ISBN 978-92-76-26100-1,
doi:10.2760/758855, JRC122635.

Bastos, A., Fu, Z., Ciais, P., Friedlingstein, P., Sitch, S., Pongratz, J., Weber, U., Reichstein,
M., Anthoni, P., Arneth, A., Haverd, V., Jain, A., Joetzjer, E., Knauer, J., Lienert, S.,
Loughran, T., McGuire, P.C., Obermeier, W., Padrón, R.S., Shi, H., Tian, H., Viovy, N.,
Zaehle, S. (2020) Impacts of extreme summers on European ecosystems: a comparative
analysis of 2003, 2010 and 2018. *Philosophical Transactions of the Royal Society B*:
Biological Sciences 375, 20190507.

Bellassen, V., Viovy, N., Luyssaert, S., Le Maire, G., Schelhaas, M. J., and Ciais, P. (2011).
Reconstruction and attribution of the carbon sink of European forests between 1950 and
2000. *Global Change Biology*, *17*(11), 3274–3292. https://doi.org/10.1111/j.1365-
2486.2011.02476.x

Bohn, FJ,, Huth, A. 2017 The importance of forest structure to biodiversity–productivity
relationships. *R. Soc. open sci*. 4: 160521.

Brunet-Navarro, P., Jochheim, H., Muys, B. (2016) Modelling carbon stocks and fluxes in the
wood product sector: a comparative review. *Global Change Biology*, 22, 2555-2569.

Campioli, M., Vicca, S., Luyssaert, S., Bilcke, J., Ceschia, E., Iii, F. S. C., and Ciais, P.
(2015). *Biomass production efficiency controlled by management in temperate and boreal
ecosystems*. *Nature Geoscience*, https://doi.org/10.1038/NGEO2553

Ceccherini, G., Duveiller, G., Grassi, G., Lemoine, G., and Avitabile, V. (2020). Abrupt
Increase in Forest Harvested Area over Europe After 2015. *Nature*, *583*.
https://doi.org/10.1038/s41586-020-2438-y

Chen, C., Riley, W.J., Prentice, I.C., Keenan, T.F. (2022) CO_2 fertilization of
terrestrial photosynthesis inferred from site to global scales. *Proceedings of the National*
Academy of Sciences, 119, e2115627119.

Churkina, G., Organschi, A., Reyer, C. P. O., Ruff, A., Vinke, K., Liu, Z., Reck, B. K.,
Graedel, T. E., & Schellnhuber, H. J. (2020). Buildings as a global carbon sink. *Nature*
Sustainability. https://doi.org/10.1038/s41893-019-0462-4

Ciais, P., Schelhaas, M. J., Zaehle, S., Piao, S. L., Cescatti, A., Liski, J., Luyssaert, S., Le-
Maire, G., Schulze, E. D., Bouriaud, O., Freibauer, A., Valentini, R., & Nabuurs, G. J.
(2008). Carbon accumulation in European forests. *Nature Geoscience*, *1*(7), 425–429.
https://doi.org/10.1038/ngeo233

Collalti, A., Perugini, L., Santini, M., Chiti, T., Nolè, A., Matteucci, G., & Valentini, R.
(2014). A process-based model to simulate growth in forests with complex structure:
Evaluation and use of 3D-CMCC Forest Ecosystem Model in a deciduous forest in Central
Italy. *Ecological Modelling*, *272*, 362–378. https://doi.org/10.1016/j.ecolmodel.2013.09.016

Collalti A., Marconi S., Ibrom A., Trotta C., Anav A., D'Andrea E., Matteucci G.,
Montagnani L., Gielen B., Mammarella I., Grünwald T., Knohl A., Berninger F., ZhaoY.,
Valentini R., Santini M., (2016). "Validation of 3D-CMCC Forest Ecosystem Model (v.5.1)
against eddy covariance data for ten European forest sites", *Geoscientific Model*
Development, 9, 1-26, https://doi.org/10.5194/gmd-9-479-2016

Collalti, A., Trotta, C., Keenan, T. F., Ibrom, A., Bond-Lamberty, B., Grote, R., Vicca, S.,
Reyer, C. P. O., Migliavacca, M., Veroustraete, F., Anav, A., Campioli, M., Scoccimarro, E.,
Šigut, L., Grieco, E., Cescatti, A., & Matteucci, G. (2018). Thinning Can Reduce Losses in
Carbon Use Efficiency and Carbon Stocks in Managed Forests Under Warmer Climate.
Journal of Advances in Modeling Earth Systems, *10*(10), 2427–2452.
https://doi.org/10.1029/2018MS001275

Collalti, A., Thornton, P. E., Cescatti, A., Rita, A., Borghetti, M., Nolè, A., Trotta, C., Ciais,
P., & Matteucci, G. (2019). The sensitivity of the forest carbon budget shifts across processes
along with stand development and climate change. *Ecological Applications*, *29*(2), 1–18.
https://doi.org/10.1002/eap.1837

Collalti, A., & Prentice, I. C. (2019). Is NPP proportional to GPP? Waring's hypothesis 20
years on. *Tree Physiology*, *39*(8), 1473–1483. https://doi.org/10.1093/treephys/tpz034

Collalti, A., Tjoelker, M. G., Hoch, G., Mäkelä, A., Guidolotti, G., Heskel, M., Petit, G.,
Ryan, M. G., Battipaglia, G., Matteucci, G., & Prentice, I. C. (2020a). Plant respiration:
Controlled by photosynthesis or biomass? *Global Change Biology*.
https://doi.org/10.1111/gcb.14857

Collalti, A., Ibrom, A., Stockmarr, A., Cescatti, A., Alkama, R., Fernández-Martínez, M.,
Matteucci, G., Sitch, S., Friedlingstein, P., Ciais, P., Goll, D. S., Nabel, J. E. M. S., Pongratz,
J., Arneth, A., Haverd, V., & Prentice, I. C. (2020b). Forest production efficiency increases
with growth temperature. *Nature Communications*, *under review*.
https://doi.org/https://doi.org/10.1101/2020.04.15.042275

de Pury DGG & Farquhar GD (1997) Simple scaling of photosynthesis from leaves to
canopies without the errors of big-leaf models. *Plant Cell and Environment*, 20, 537–557.

de Wergifosse L. et al. (2022). Simulating tree growth response to climate change in
structurally-diverse oak and beech forests. *Science of the Total Environment*, 806(2), 150422,
https://doi.org/10.1016/j.scitotenv.2021.150422

De Marco, A., Sicard, P., Feng, Z., Agathokleous, E., Alonso, R., Araminiene, V.,
Augustatis, A., Badea, O., Beasley, J. C., Branquinho, C., Bruckman, V. J., Collalti, A.,
David-Schwartz, R., Domingos, M., Du, E., Garcia Gomez, H., Hashimoto, S., Hoshika, Y.,
Jakovljevic, T. … Paoletti, E. (2022). Strategic roadmap to assess forest vulnerability under
air pollution and climate change. Global Change Biology, 28, 5062– 5085.
https://doi.org/10.1111/gcb.16278

D'Andrea A., Rezaie N., Prislan P., Gričar J., Collalti A., Muhr J., Matteucci G., "Cold and
drought: effects of extreme weather events on Stem Carbon Dynamic in a Mediterranean
beech forest", *Plant Cell & Environment*, 43(10):2365–2379,
https://doi.org/10.1002/PCE.13858, 2020

D'Andrea E., Scartazza A., Battistelli A., Collalti A., Proietti S., Rezaie N., Matteucci G.,
Moscatello S. (2021). Unravelling resilience mechanisms in forests: role of non-structural
carbohydrates in responding to extreme weather events. *Tree Physiology*, 41, 10, 1808-1818,
https://doi.org/10.1093/treephys/tpab044

Drake, J. E., & Tjoelker, M. G. (2016). Does physiological acclimation to climate warming
stabilize the ratio of canopy respiration to photosynthesis? *New Phytologist*, 211(3), 850–863.
https://doi.org/10.1111/nph.13978

Duffy, P. B., Brando, P., Asner, G. P., & Field, C. B. (2015). Projections of future
meteorological drought and wet periods in the Amazon. *Proceedings of the National
Academy of Sciences of the United States of America*.
https://doi.org/10.1073/pnas.1421010112

Duncker, P. S., Barreiro, S. M., Hengeveld, G. M., Lind, T., Mason, W. L., Ambrozy, S., &
Spiecker, H. (2012). Classification of forest management approaches: A new conceptual
framework and its applicability to European forestry. *Ecology and Society*, *17*(4).
https://doi.org/10.5751/ES-05262-170451

Engel M., Vospernik S., Toigo M., Morin X., Tomao A., Trotta C., Steckel M., Barbati A.,
Nothdurft A. Pretzsch H., del Rio M., Skrzyszewski J., Ponette Q., Lof M., Jansons A.,
Brazaitis G. (2021). Simulating the effects of thinning and species mixing on stands of oak
(Quercus petrea (Matt.) Liebl. / Quercus robur L.) and pine (Pinus sylvestris L.) across
Europe. *Ecological Modelling*, 442 (109406),
https://doi.org/10.1016/j.ecolmodel.2020.109406

Etzold, S., Ferretti, M., Reinds, G.J., Solberg, S., Gessler, A., Waldner, P., Schaub, M.,
Simpson, D., Benham, S., Hansen, K., Ingerslev, M., Jonard, M., Karlsson, P.E., Lindroos,
A.-J., Marchetto, A., Manninger, M., Meesenburg, H., Merilä, P., Nöjd, P., Rautio, P.,

Sanders, T.G.M., Seidling, W., Skudnik, M., Thimonier, A., Verstraeten, A., Vesterdal, L., Vejpustkova, M., de Vries, W. (2020) Nitrogen deposition is the most important environmental driver of growth of pure, even-aged and managed European forests. *Forest Ecology and Management* 458, 117762. https://doi.org/10.1016/j.foreco.2019.117762

European Commission. "Multi-annual implementation plan of the new EU forest strategy." SWD (2015) 164 Final (2015).

Farquhar GD, et al. (1980) A biochemical model of photosynthetic CO2 assimilation in leaves of C3 species. *Planta*. 149: 78-90.

Favero, A., Daigneault, A., & Sohngen, B. (2020). Forests: Carbon sequestration, biomass energy, or both? *Science Advances*, *6*(13), 1–14. https://doi.org/10.1126/sciadv.aay6792

FAO and PlanBleu (Eds.). "State of Mediterranean Forests 2018". Food and Agriculture Organization of the United Nations, Rome and Plan Bleu, Marseille, ISBN 978-92-5-131047-2, http://www.fao.org/3/CA2081EN/ca2081en.PDF, 2018.

FOREST EUROPE, & FAO, U. and. (2020). *FOREST EUROPE, 2020: State of Europe's Forests 2020*.

Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C., Arneth, A., Arora, V., Bates, N. R., … Zaehle, S. (2020). Global Carbon Budget 2020. *Earth System Science Data*, *12*(4), 3269–3340. https://doi.org/10.5194/essd-12-3269-2020

Fritsch, M, Lischke, H, Meyer, KM. Scaling methods in ecological modelling. *Methods Ecol Evol*. 2020; 11: 1368– 1378. https://doi.org/10.1111/2041-210X.13466

Gampe, D., Zscheischler, J., Reichstein, M., O'Sullivan, M., Smith, W. K., Sitch, S., & Buermann, W. (2021). Increasing impact of warm droughts on northern ecosystem productivity over recent decades. *Nature Climate Change*, 11(9), 772-779.

Garcia-Gonzalo, J., Peltola, H., Briceño-Elizondo, E., & Kellomäki, S. (2007). Changed thinning regimes may increase carbon stock under climate change: A case study from a Finnish boreal forest. *Climatic Change*, *81*(3–4), 431–454. https://doi.org/10.1007/s10584-006-9149-8

Gatti, L.V., Basso, L.S., Miller, J.B. et al. Amazonia as a carbon source linked to deforestation and climate change. *Nature* 595, 388–393 (2021). https://doi.org/10.1038/s41586-021-03629-6

Grassi, G., House, J., Dentener, F., Federici, S., Den Elzen, M., & Penman, J. (2017). The key role of forests in meeting climate targets requires science for credible mitigation. *Nature Climate Change*, *7*(3), 220–226. https://doi.org/10.1038/nclimate3227

Grassi, G., Stehfest, E., Rogelj, J., van Vuuren, D., Cescatti, A., House, J., Nabuurs, G. J., Rossi, S., Alkama, R., Viñas, R. A., Calvin, K., Ceccherini, G., Federici, S., Fujimori, S., Gusti, M., Hasegawa, T., Havlik, P., Humpenöder, F., Korosuo, A., … Popp, A. (2021).

Critical adjustment of land mitigation pathways for assessing countries' climate progress. *Nature Climate Change*, *11*(5), 425–434. https://doi.org/10.1038/s41558-021-01033-6

Grossiord, C., Buckley, T. N., Cernusak, L. A., Novick, K. A., Poulter, B., Siegwolf, R. T. W., Sperry, J. S., & McDowell, N. G. (2020). Plant responses to rising vapor pressure deficit. *New Phytologist*, *226*(6), 1550–1566. https://doi.org/https://doi.org/10.1111/nph.16485

Hempel, S., Frieler, K., Warszawski, L., Schewe, J., and Piontek, F.: A trend-preserving bias correction – the ISI-MIP approach, *Earth Syst. Dynam.*, 4, 219–236, https://doi.org/10.5194/esd-4-219-2013, 2013.

Henttonen, H. M., Nöjd, P., & Mäkinen, H. (2017). Environment-induced growth changes in the Finnish forests during 1971–2010 – An analysis based on National Forest Inventory. *Forest Ecology and Management*, *386*, 22–36. https://doi.org/10.1016/j.foreco.2016.11.044

Howard, C., Dymond, C. C., Griess, V. C., Tolkien-Spurr, D., & van Kooten, G. C. (2021). Wood product carbon substitution benefits: a critical review of assumptions. *Carbon Balance and Management*, *16*(1), 1–11. https://doi.org/10.1186/s13021-021-00171-w

Huang J., Hammerbacher A., Gershenzon J., van Dam, N.M., Sala, A., McDowell N.G., et al. (2021). Storage of carbon reserves in spruce trees is prioritized over growth in the face of carbon limitation. *PNAS*, 118(33), e2023297118; DOI: 10.1073/pnas.2023297118

Huber, N., Bugmann, H., & Lafond, V. (2018). Global sensitivity analysis of a dynamic vegetation model: Model sensitivity depends on successional time, climate and competitive interactions. *Ecological Modelling*, *368*, 377–390. https://doi.org/10.1016/j.ecolmodel.2017.12.013

Huber, N., Bugmann, H., & Lafond, V. (2020). Capturing ecological processes in dynamic forest models: why there is no silver bullet to cope with complexity. *Ecosphere*, *11*(5). https://doi.org/10.1002/ecs2.3109

Kauppi, P. E., Sedjio, R., Apps, M., Cerri, C., Fujimori, T., Janzen, H., Krankina, O., Makundi, W., Marland, R., Masera, O., & Nabuurs. (2001). Technological and economic potential of options to enhance, maintain, and manage biological carbon reservoirs and Geo-engineering. In IPCC (Ed.), *Climate Change 2001: Mitigation* (pp. 302–343). Cambridge University Press.

Kattge, J. and Knorr, W. (2007), Temperature acclimation in a biochemical model of photosynthesis: a reanalysis of data from 36 species. *Plant, Cell & Environment*, 30: 1176-1190. https://doi.org/10.1111/j.1365-3040.2007.01690.x

Kaipainen, T., Liski, J., Pussinen, A., Karjalainen, T. (2004) Managing carbon sinks by changing rotation length in European forests. *Environmental Science & Policy* 7, 205-219.

Kindermann, G.E., Schörghuber, S., Linkosalo, T. et al. Potential stocks and increments of woody biomass in the European Union under different management and climate scenarios. *Carbon Balance Management* 8, 2 (2013). https://doi.org/10.1186/1750-0680-8-2

Kirschbaum, M. (2005). A model analysis of the interaction between forest age and forest
responsiveness to increasing CO_2 concentration. *Tree Physiology*, 25(7), 953–963.
https://doi.org/10.1093/treephys/25.7.953

Körner, C. (2005). Worldwide positions of alpine treelines and their causes. *The Impacts of*
Climate Variability on Forests, 221–229. https://doi.org/10.1007/bfb0009775

Krofcheck, D. J., Remy, C. C., Keyser, A. R., & Hurteau, M. D. (2019). Optimizing Forest
Management Stabilizes Carbon Under Projected Climate and Wildfires. *Journal of*
Geophysical Research: Biogeosciences, 3075–3087. https://doi.org/10.1029/2019JG005206

Leskinen, P., Cardellini, G., González García, S., Hurmekoski, E., Sathre, R., Seppälä, J.,
Smyth, C. E., Stern, T., & Verkerk, H. (2018). Substitution effects of wood-based products in
climate change mitigation. *From Science to Policy*, *7*(November), 28.
https://www.efi.int/publications-bank/substitution-effects-wood-based-products-climate-
change-mitigation

Leslie, A. J. (1966). A review of the concept of the normal forest. *Australian Forestry*, *30*(2),
139–147. https://doi.org/10.1080/00049158.1966.10675407

Luyssaert, S., Inglima, I., Jung, M., Richardson, A. D., Reichstein, M., Papale, D., Piao, S. L.,
Schulze, E. D., Wingate, L., Matteucci, G., Aragao, L., Aubinet, M., Beer, C., Bernhofer, C.,
Black, K. G., Bonal, D., Bonnefond, J. M., Chambers, J., Ciais, P., … Janssens, I. A. (2007).
CO2 balance of boreal, temperate, and tropical forests derived from a global database. *Global*
Change Biology, *13*(12), 2509–2537. https://doi.org/10.1111/j.1365-2486.2007.01439.x

Luyssaert, S., Marie, G., Valade, A., Chen, Y. Y., Njakou Djomo, S., Ryder, J., Otto, J.,
Naudts, K., Lansø, A. S., Ghattas, J., & McGrath, M. J. (2018). Trade-offs in using European
forests to meet climate objectives. *Nature*, *562*(7726), 259–262.
https://doi.org/10.1038/s41586-018-0577-1

Mahnken, M., Cailleret, M., Collalti, A., Trotta, C., Biondo, C., D'Andrea, E., Dalmonech,
D., Marano, G., Mäkelä, A., Minunno, F., Peltoniemi, M., Trotsiuk, V., Nadal-Sala, D.,
Sabaté, S., Vallet, P., Aussenac, R., Cameron, D. R., Bohn, F. J., Grote, R. … Reyer, C. P. O.
(2022). Accuracy, realism and general applicability of European forest models. Global
Change Biology, 00, 1– 23. https://doi.org/10.1111/gcb.16384

Marconi, S., Chiti, T., Nolè, A., Valentini, R., & Collalti, A. (2017). The role of respiration in
estimation of net carbon cycle: Coupling soil carbon dynamics and canopy turnover in a
novel version of 3D-CMCC forest ecosystem model. *Forests*.
https://doi.org/10.3390/f8060220

Maréchaux I., Langerwish F., Huth A., Bugmann H., Morin X., Reyer C.P.O., Seidl R.,
Collalti A., Dantas de Paula M., Fischer R., Gutsch M., Lexer M.J., Lischke H., Rammig A.,
Rödig E., Sakschewski B., Taubert F., Thonicke K., Vacchiano G., Bohn F.J. (2021),
"Modelling forests to address key ecological questions: lessons learned from different
modelling communities and possible future paths", Ecology and Evolution, 11(9):3746–3770,
https://doi.org/10.1002/ece3.7391

Mathias, J.M. & Trugman, A.T. (2021) Climate change impacts plant carbon balance, increasing mean future carbon use efficiency but decreasing total forest extent at dry range edges. *Ecology Letters*, 00, 1– 11. https://doi.org/10.1111/ele.13945

Mason, R. L., Gunst, R. F., & Hess, J. L. (2003). Factorial Experiments in Completely Randomized Designs. In *Statistical Design and Analysis of Experiments*. https://doi.org/10.1002/0471458503.ch5

McCree, K. (1970). An equation for the rate of respiration of white clover plants grown under controlled conditions. In I. Šetliik (Ed.), Prediction and measurement of photosynthetic productivity (pp. 221–229). Wageningen, The Netherlands: Centre for Agricultural Publishing and Documentation.

McDowell, N. G., Allen, C. D., Anderson-teixeira, K., Aukema, B. H., Bond-Lamberty, B., Chini, L., Clark, J. S., Dietze, M., Grossiord, C., Hanbury-brown, A., Hurtt, G. C., Jackson, R. B., Johnson, D. J., Kueppers, L., Lichstein, J. W., Ogle, K., Poulter, B., Pugh, T. A. M., Seidl, R., … Xu, C. (2020). Pervasive shifts in forest dynamics in a changing world. *Science*. https://doi.org/10.1126/science.aaz9463

Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J.-F., Matsumoto, K., Montzka, S. A., Raper, S. C. B., Riahi, K., Thomson, A., Velders, G. J. M., and van Vuuren, D. P. P. (2011). The RCP greenhouse gas concentrations and their extensions from 1765 to 2300, *Clim. Change*, 109, 213, https://doi.org/10.1007/s10584-011-0156-z.

Merganičová, K., Merganič, J., Lehtonen, A., Vacchiano, G., Sever, M. Z. O., Augustynczik, A. L. D., Grote, R., Kyselová, I., Mäkelä, A., Yousefpour, R., Krejza, J., Collalti, A., & Reyer, C. P. O. (2019). Forest carbon allocation modelling under climate change. *Tree Physiology*, *39*(12), 1937–1960. https://doi.org/10.1093/treephys/tpz105

Migliavacca, M., Musavi, T., Mahecha, M.D. et al. The three major axes of terrestrial ecosystem function. *Nature* 598, 468–472 (2021). https://doi.org/10.1038/s41586-021-03939-9

Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., Van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P., & Wilbanks, T. J. (2010). The next generation of scenarios for climate change research and assessment. *Nature*. https://doi.org/10.1038/nature08823

Nabuurs, G. J., Thürig, E., Heidema, N., Armolaitis, K., Biber, P., Cienciala, E., Kaufmann, E., Mäkipää, R., Nilsen, P., Petritsch, R., Pristova, T., Rock, J., Schelhaas, M. J., Sievanen, R., Somogyi, Z., & Vallet, P. (2008). Hotspots of the European forests carbon cycle. *Forest Ecology and Management*, *256*(3), 194–200. https://doi.org/10.1016/j.foreco.2008.04.009

Nabuurs, Gert Jan, Lindner, M., Verkerk, P. J., Gunia, K., Deda, P., Michalak, R., & Grassi, G. (2013). First signs of carbon sink saturation in European forest biomass. *Nature Climate Change*, *3*(9), 792–796. https://doi.org/10.1038/nclimate1853

Nabuurs, Gert Jan, Verweij, P., Van Eupen, M., Pérez-Soba, M., Pülzl, H., & Hendriks, K. (2019). Next-generation information to support a sustainable course for European forests. *Nature Sustainability*, *2*(9), 815–818. https://doi.org/10.1038/s41893-019-0374-3

Naudts, K., Chen, Y., McGrath, M. J., Ryder, J., Valade, A., Otto, J., & Luyssaert, S. (2016). Europe's forest management did not mitigate climate warming. *Science*, 351(6273), 597-600

Nolè, A., Collalti, A., Borghetti, M., Chiesi, M., Chirici, G., Magnani, F., et al. (2015). The role of managed forest ecosystems: A modelling based approach. In R. Valentini & F. Miglietta (Eds.), The greenhouse gas balance of Italy (pp. 71–85). Berlin-Heidelberg: Springer-Verlag.

Oleson, K. W., Lawrence, D. M., Bonan, G. B., Drewniak, B., Huang, M., Koven, C. D., ... Yang, Z.L. (2013). Technical description of version 4.5 of the community land model (CLM). Ncar Technical Note NCAR/TN- 503+STR. Boulder, CO: National Center for Atmospheric Research, 422 pp.

Pappas, C., Fatichi, S., Leuzinger, S., Wolf, A., & Burlando, P. (2013). Sensitivity analysis of a process-based ecosystem model: Pinpointing parameterization and structural issues. *Journal of Geophysical Research*, 118, 502–528. https://doi.org/10.1002/jgrg.20035

Pappas, C., Maillet, J., Rakowski, S., Baltzer, J. L., Barr, A. G., Black, T. A., Fatichi, S., Laroque, C. P., Matheny, A. M., Roy, A., Sonnentag, O., & Zha, T. (2020). Aboveground tree growth is a minor and decoupled fraction of boreal forest carbon input. *Agricultural and Forest Meteorology*, *290*, 108030. https://doi.org/10.1016/j.agrformet.2020.108030

Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y. W., Poindexter, C., Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Ribeca, A., van Ingen, C., Zhang, L., Amiro, B., Ammann, C., Arain, M. A., Ardö, J., … Papale, D. (2020). The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data. *Scientific Data*, *7*(1), 225. https://doi.org/10.1038/s41597-020-0534-3

Peano D., Materia S., Collalti A., Alessandri A., Anav A., Bombelli A., Gualdi S. (2019). Global variability of simulated and observed vegetation growing season. *Journal of Geophysical Research: Biogeosciences*, 124, 3569-3587, https://doi.org/10.1029/2018JG004881

Peñuelas, J., Ciais, P., Canadell, J. G., Janssens, I. A., Fernández-Martínez, M., Carnicer, J., Obersteiner, M., Piao, S., Vautard, R., & Sardans, J. (2017). Shifting from a fertilization-dominated to a warming-dominated period. *Nature Ecology and Evolution*, *1*(10), 1438–1445. https://doi.org/10.1038/s41559-017-0274-8

Piao, S., Wang, X., Park, T., Chen, C., Lian, X., He, Y., Bjerke, J. W., Chen, A., Ciais, P., Tømmervik, H., Nemani, R. R., & Myneni, R. B. (2020). Characteristics, drivers and feedbacks of global greening. *Nature Reviews Earth and Environment*, *1*(1), 14–27. https://doi.org/10.1038/s43017-019-0001-x

Pugh, T. A. M., Lindeskog, M., Smith, B., Poulter, B., Arneth, A., Haverd, V., & Calle, L. (2019). Role of forest regrowth in global carbon sink dynamics. *Proceedings of the National*

Academy of Sciences of the United States of America. https://doi.org/10.1073/pnas.1810512116

Pussinen, Ari, Karjalainen, T., Mäkipää, R., Valsta, L., & Kellomäki, S. (2002). Forest carbon sequestration and harvests in Scots pine stand under different climate and nitrogen deposition scenarios. *Forest Ecology and Management.* https://doi.org/10.1016/S0378-1127(00)00675-7

Pussinen, A., Nabuurs, G. J., Wieggers, H. J. J., Reinds, G. J., Wamelink, G. W. W., Kros, J., Mol-Dijkstra, J. P., & de Vries, W. (2009). Modelling long-term impacts of environmental change on mid- and high-latitude European forests and options for adaptive forest management. *Forest Ecology and Management*, *258*(8), 1806–1813. https://doi.org/10.1016/j.foreco.2009.04.007

Reich, P. B. et al. Scaling of respiration to nitrogen in leaves, stems and roots of higher land plants. *Ecol. Lett.* 11, 793–801 (2008).

Reich, P. B., & Oleksyn, J. (2008). Climate warming will reduce growth and survival of Scots pine except in the far north. *Ecology Letters*, **1**(6), 588–597.

Reich, P., Sendall, K., Stefanski, A., Wei, X., Rich, R. L., & Montgomery, R. A. (2016). Boreal and temperate trees show strong acclimation of respiration to warming. *Nature*, 531(7596), 633–636. https://doi.org/10.1038/nature17142

Reyer, C. P. O., Silveyra Gonzalez, R., Dolos, K., Hartig, F., Hauf, Y., Noack, M., Lasch-Born, P., Rötzer, T., Pretzsch, H., Meesenburg, H., Fleck, S., Wagner, M., Bolte, A., Sanders, T. G. M., Kolari, P., Mäkelä, A., Vesala, T., Mammarella, I., Pumpanen, J., … Frieler, K. (2020). The PROFOUND Database for evaluating vegetation models and simulating climate impacts on European forests. *Earth System Science Data*, *12*(2), 1295–1320. https://doi.org/10.5194/essd-12-1295-2020

Reyer, C.P.O., Bathgate, S., Blennow, K., Borges, J.G., Bugmann, H., Delzon, S., Faias, S.P., Garcia-Gonzalo, J., Gardiner, B., Gonzalez-Olabarria, J.R., Gracia, C., Hernández, J.G., Kellomäki, S., Kramer, K., Lexer, M.J., Lindner, M., van der Maaten, E., Maroschek, M., Muys, B., Nicoll, B., Palahi, M., Palma, J.H.N., Paulo, J.A., Peltola, H., Pukkala, T., Rammer, W., Ray, D., Sabaté, S., Schelhaas, M.-J., Seidl, R., Temperli, C., Tomé, M., Yousefpour, R., Zimmermann, N.E., Hanewinkel, M. (2017) Are forest disturbances amplifying or canceling out climate change-induced productivity changes in European forests? *Environmental Research Letters* 12, 034027.

Ryan, M. G. (1991). Effects of climate change on plant respiration. *Ecological Applications*, 1(2), 157–167.

Ryan, M.G., Phillips, N., & Bond, B. J. (2006). The hydraulic limitation hypothesis revisited. *Plant, Cell & Environment*, *29*(3), 367–381. https://doi.org/https://doi.org/10.1111/j.1365-3040.2005.01478.x

Sala, A., Fouts, W., & Hoch, G. (2011). Carbon storage in trees: Does relative carbon supply decrease with tree size? In F. C. Meinzer, B. Lachenbruch, & T. E. Dawson (Eds.), Size-

and age-related changes in tree structure and function (pp. 287–306). Dordrecht, the Netherlands: Springer.

Schelhaas, MJ., Nabuurs, GJ., Hengeveld, G. et al. Alternative forest management strategies to account for climate change-induced productivity and species suitability changes in Europe. *Reg. Environ. Change* 15, 1581–1594 (2015). https://doi.org/10.1007/s10113-015-0788-z

Schulze, E. D., Sierra, C. A., Egenolf, V., Woerdehoff, R., Irslinger, R., Baldamus, C., Stupak, I., & Spellmann, H. (2020). The climate change mitigation effect of bioenergy from sustainably managed forests in Central Europe. *GCB Bioenergy*, *12*(3), 186–197. https://doi.org/10.1111/gcbb.12672

Sedmáková, D., Sedmák, R., Bosela, M., Je, M., Bla, M., & Maru, R. (2019). *Dendrochronologia Growth-climate responses indicate shifts in the competitive ability of European beech and Norway spruce under recent climate warming in East- Central Europe*. *54*, 37–48. https://doi.org/10.1016/j.dendro.2019.02.001

Seidl, R., Thom, D., Kautz, M., Martin-Benito, D., Peltoniemi, M., Vacchiano, G., Wild, J., Ascoli, D., Petr, M., Honkaniemi, J., Lexer, M.J., Trotsiuk, V., Mairota, P., Svoboda, M., Fabrika, M., Nagel, T.A., Reyer, C.P.O. (2017) Forest disturbances under climate change. *Nature Climate Change* 7, 395-402.

Senf, C., Seidl, R. (2021a) Mapping the forest disturbance regimes of Europe. *Nature Sustainability* 4, 63–70.

Senf, C., & Seidl, R. (2021b). *Storm and fire disturbances in Europe : Distribution and trends*. 3605–3619. https://doi.org/10.1111/gcb.15679

Settele J, Scholes R, Betts R, Bunn S, Leadley P, Nepstad D, Overpeck JT, Taboada MA (2014) Terrestrial and inland water systems. In: Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, Chatterjee M, Ebi KL, Estrada YO, Genova RC, Girma B, Kissel ES, Levy AN, MacCracken S, Mastrandrea PR,White LL (eds) Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp 271–359

Sha, Z., Bai, Y., Li, R. et al. The global carbon sink potential of terrestrial vegetation can be increased substantially by optimal land management. Commun Earth Environ 3, 8 (2022). https://doi.org/10.1038/s43247-021-00333-1

Shinozaki K, et al. (1964) A quantitative analysis of plant form – the pipe model theory. I. Basic analyses. *Japanese Journal of Ecology* 14, 97–105

Smith N & Dukes J (2012). Plant respiration and photosynthesis in global-scale models: incorporating acclimation and CO_2. *Global Change Biology*, 19: 45-63.

Tahvonen, O. (2016). Economics of rotation and thinning revisited: The optimality of clearcuts versus continuous cover forestry. *Forest Policy and Economics*. https://doi.org/10.1016/j.forpol.2015.08.013

Tang, J., Luyssaert, S., Richardson, A. D., Kutsch, W., & Janssens, I. A. (2014). Steeper declines in forest photosynthesis than respiration explain age-driven decreases in forest growth. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(24), 8856–8860. https://doi.org/10.1073/pnas.1320761111

Temperli, C., Bugmann, H., & Elkin, C. (2012). Adaptive management for competing forest goods and services under climate change. *Ecological Applications*, *22*(8), 2065–2077. https://doi.org/10.1890/12-0210.1

Thornley, J. H. M., & Cannell, M. G. R. (2000). Modelling the components of plant respiration: Representation and realism. *Annals of Botany*, 85, 55–67. https ://doi.org/10.1006/anbo.1999.0997

Thom, D., Rammer, W., & Seidl, R. (2017). Disturbances catalyze the adaptation of forest ecosystems to changing climate conditions. *Global Change Biology*. https://doi.org/10.1111/gcb.13506

Tong, X., Brandt, M., Yue, Y., Ciais, P., Rudbeck Jepsen, M., Penuelas, J., Wigneron, J. P., Xiao, X., Song, X. P., Horion, S., Rasmussen, K., Saatchi, S., Fan, L., Wang, K., Zhang, B., Chen, Z., Wang, Y., Li, X., & Fensholt, R. (2020). Forest management in southern China generates short term extensive carbon sequestration. *Nature Communications*, *11*(1), 1–10. https://doi.org/10.1038/s41467-019-13798-8

Trotsiuk, V, Hartig, F, Cailleret, M, et al. (2020). Assessing the response of forest productivity to climate extremes in Switzerland using model–data fusion. *Global Change Biology*, 26: 2463– 2476. https://doi.org/10.1111/gcb.15011

Tjoelker M, et al. (2001). Modelling respiration of vegetation: evidence for a general temperature-dependent Q10. Global Change Biology, 7: 223-230.

Walker, A. P., De Kauwe, M. G., Bastos, A., Belmecheri, S., Georgiou, K., Keeling, R. F., McMahon, S. M., Medlyn, B. E., Moore, D. J. P., Norby, R. J., Zaehle, S., Anderson-Teixeira, K. J., Battipaglia, G., Brienen, R. J. W., Cabugao, K. G., Cailleret, M., Campbell, E., Canadell, J. G., Ciais, P., … Zuidema, P. A. (2021). Integrating the evidence for a terrestrial carbon sink caused by increasing atmospheric CO2. *New Phytologist*, *229*(5), 2413–2445. https://doi.org/https://doi.org/10.1111/nph.16866

Wang, Y., Zhang, C., Meng, F. R., Bourque, C. P. A., & Zhang, C. (2020). Evaluation of the suitability of six drought indices in naturally growing, transitional vegetation zones in Inner Mongolia (China). *PLoS ONE*, *15*(5), 1–15. https://doi.org/10.1371/journal.pone.0233525

Wang et al. (2021) Recent global decline of CO_2 fertilization effects on vegetation photosynthesis" Science, https://doi.org/10.1126/science.abb7772

Warszawski, L., Frieler, K., Huber, V., Piontek, F., Serdeczny, O., & Schewe, J. (2014). The inter-sectoral impact model intercomparison project (ISI-MIP): Project framework. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(9), 3228–3232. https://doi.org/10.1073/pnas.1312330110

Way, D. A. & Sage, R. F. Elevated growth temperatures reduce the carbon gain of black
spruce [Picea mariana (Mill.) B.S.P.] (2008). *Glob. Chang. Biol* 14, 624–636.

Wu, J., Larsen, K., van der Linden, L., Beier, C., Pilegaard, K., & Ibrom, A. (2013).
Synthesis on the carbon budget and cycling in a Danish, temperate deciduous forest.
Agricultural and Forest Meteorology, 181, 94–107.
https://doi.org/10.1016/j.agrformet.2013.07.012

Vacchiano G, Magnani F, Collalti A (2012) Modeling Italian forests: state of the art and
future challenges. *iForest*, e1–e8

Valade, A., Bellassen, V., Magand, C., & Luyssaert, S. (2017). Forest Ecology and
Management Sustaining the sequestration efficiency of the European forest sector. *Forest
Ecology and Management*, *405*, 44–55. https://doi.org/10.1016/j.foreco.2017.09.009

Vanninen P., Mäkelä A., Carbon budget for Scots pine trees: effects of size, competition and
site fertility on growth allocation and production (2005), *Tree Physiology*, Volume 25, 17–30,
https://doi.org/10.1093/treephys/25.1.17

van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt,
G. C., Kram, T., Krey, V., Lamarque, J. F., Masui, T., Meinshausen, M., Nakicenovic, N.,
Smith, S. J., & Rose, S. K. (2011). The representative concentration pathways: An overview.
Climatic Change. https://doi.org/10.1007/s10584-011-0148-z

Yousefpour, R., Temperli, C., Jacobsen, J. B., Thorsen, B. J., Meilby, H., Lexer, M. J.,
Lindner, M., Bugmann, H., Borges, J. G., Palma, J. H. N., Ray, D., Zimmermann, N. E.,
Delzon, S., Kremer, A., Kramer, K., Reyer, C. P. O., Lasch-Born, P., Garcia-Gonzalo, J., &
Hanewinkel, M. (2017). A framework for modeling adaptive forest management and decision
making under climate change. *Ecology and Society*, *22*(4). https://doi.org/10.5751/ES-09614-
220440

Yuan, Z., Ali, A., Jucker, T., Ruiz-Benito, P., Wang, S., Jiang, L., Wang, X., Lin, F., Ye, J.,
Hao, Z., & Loreau, M. (2019). Multiple abiotic and biotic pathways shape biomass
demographic processes in temperate forests. *Ecology*, *100*(5), e02650.
https://doi.org/https://doi.org/10.1002/ecy.2650

Zaehle, S., Sitch, S., Prentice, I. C., Liski, J., Cramer, W., Erhard, M., Hickler, T., & Smith,
B. (2006). The importance of age-related decline in forest NPP for modeling regional carbon
balances. *Ecological Applications*, *16*(4), 1555–1574. https://doi.org/10.1890/1051-
0761(2006)016[1555:TIOADI]2.0.CO;2

Zeide B. (2001). Thinning and growth: A full turnaround, *Journal of Forestry*, 99(1), 20-25.
https://doi.org/10.1093/jof/99.1.20

Figures

Figure 1 | Conceptual scheme of the virtual stands creation: in Phase 1 the model is initialized with data from the actual forest stands and then simulations are carried out for 202 years of contemporary (1996-2006) weather and atmospheric CO_2 concentration. In Phase 2, multiple stands are drawn from the simulations in Phase 1 and used to build the Composite Forest Matrix (CFM) composed of representative forest stands. The climate change (RCPs) and management scenarios (BAU, Alternative Managements, No-Management) simulations are then applied to the CFM.

1093

1094

Figure 2 | NPP (Net primary productivity, g C m^{-2} year^{-1}) simulations under different
management scenarios (AM+, BAU, AM–) and the NO-MAN scenario for each of the four
atmospheric CO_2 concentration pathways (RCPs). NPP, solid line, is averaged across the
representative forests, different ESMs and aggregated according to the management regime.
Shaded areas represent the maximum and minimum values (5[th] and 95[th] percentiles) across the
representative forests, different ESMs and aggregated according to the management regime.

1101

Figure 3 | pCWS (potential Carbon Woody Stock = standing and potential harvested woody
biomass; t C ha^{-1}) simulations under different management scenarios (AM+, BAU, AM–) and
the NO-MAN scenario divided by different emission scenario RCPs. pCWS, solid line, is
averaged across the representative forests, different ESMs and aggregated according to the
management regime. Shaded areas represent the maximum and minimum values (5[th] and 95[th]
percentiles) across the representative forests, different ESMs and aggregated according to the
management regime. Carbon sequestration rates (as annual increase of CWS, t C ha^{-1} year^{-1})
in the potential total woody stocks (mean and standard deviation) are reported in the bar
plots.

1111

1112

1113

1114

1115

1116

1117

Figure 4 | Average NPP (net primary productivity, g C m^{-2} year^{-1}) vs. pCWS (the sum of standing and potential harvested woody products; t C ha^{-1}) over the period 2006-2099, for the three management scenarios: AM+, AM−, BAU; and the NO-MAN for the 4 RCPs. Reported values refer to data averaged across real and virtual stands and across species. Data ellipses are also reported in shaded colors and refer to all data. NOTE: each single scenario according to Table S1 is reported here (16 in total excluding the mixed ones). In the subplot the differences are expressed as % and are reported along a parametric curve (third order polynomial) with the point (0, 0) representing the reference BAU. Arrows indicate the increasing intensity of management intervention. No significant differences across RCPs were detected.

1128

1129

1130

1131

1132

Figure 5 | Percentage of changes for Mean NPP (net primary production) and pCWS (potential Carbon Woody Stocks) over the period 2006-2099 between the managed scenarios AM+, AM–, MIX, NO-MAN (AM+, more intensive than BAU; AM–, less intensive than BAU (business as usual); MIX: mixes options according to Table S1, NO-MAN: no management option) and the BAU scenario reported for the four representative concentration pathways (RCPs). Values refer to data averaged across real and virtual stands and across species. Note: each single scenario according to Table S1 is here reported (28 in total). Parametric curve fitting (polynomial of order 3) is also reported for each RCP and it is computed according to the averaged data for each management intervention scenario and RCP combination.

1143

Tables

Table 1 | Site description for model initialization data (corresponding to the year 1997 for Sorø and Hyytiälä and 2000 for Bílý Kříž) to the real stands' characteristics, and management variables used in simulations (see also Collalti et al., 2018; Reyer et al., 2020). Values in brackets represent bounds of variability (the maximum and the minimum absolute values) adopted for alternative management simulations. Re-planting information for the sites in the simulation experiments, according to ISIMIP protocol as in Reyer et al. (2020). The real stands refer to the monitoring sites in Sorø (*F. sylvatica*, Denmark), Bílý Kříž (*P. abies*, Czech Republic) and Hyytiälä (*P. sylvestris*, Finland).

Species	DBH (cm)	Age (years)	Tree height (m)	Density (trees ha^{-1})	Thinning intensity (% basal area)	Thinning interval (years)	Rotation age (years)	Replanting Species	Density (trees ha^{-1})	Age (years)	Tree height (m)
Fagus sylvatica	25	80	25	400	30 (20-40)	15 (5-25)	140 (120-160)	*Fagus sylvatica*	6000	4	1.3
Pinus sylvestris	10.3	36	10	1800	20 (10-30)	15 (5-25)	140 (120-160)	*Pinus sylvestris*	2250	2	1.3
Picea abies	7.1	16	5.6	2408	30 (20-40)	15 (5-25)	120 (100-140)	*Picea abies*	4500	4	1.3

Table 2 | NPP and pCWS computed as average over the simulation period 2006-2099, across all stands and ESMs climate forcing but grouped across RCPs. Mean differences (in percentage) are reported in parenthesis for NPP and pCWS between the alternative management scenarios and the *Business-As-Usual* (BAU) practices used here as the benchmark scenario.

MANAGEMENT type	RCP	NPP gC m^{-2} y^{-1}	pCWS tC ha^{-1}
BAU	RCP 2.6	501.7	192.9
BAU	RCP 4.5	522.1	195.6
BAU	RCP 6.0	530.5	196.0
BAU	RCP 8.5	542.0	198.5
AM+	RCP 2.6	350.0 (−30.2%)	184.3 (−4.4%)
AM+	RCP 4.5	366.8 (−29.7%)	186.5 (−4.6%)
AM+	RCP 6.0	372.0 (−29.8%)	186.4 (−4.8%)
AM+	RCP 8.5	388.1 (−28.4%)	189.7 (−4.4%)
AM−	RCP 2.6	495.4 (−1.2%)	179.8 (−6.6%)
AM−	RCP 4.5	510.6 (−2.1%)	181.6 (−7.1%)
AM−	RCP 6.0	519.9 (−1.9%)	182.2 (−6.9%)
AM−	RCP 8.5	524.7 (−3.1%)	183.8 (−7.3%)
NO-MAN	RCP 2.6	429.1 (−14.4%)	136.2 (−29.3%)
NO-MAN	RCP 4.5	436.4 (−16.4%)	136.6 (−30.1%)
NO-MAN	RCP 6.0	444.8 (−16.1%)	137.1 (−30.0%)
NO-MAN	RCP 8.5	436.5 (−19.4%)	136.8 (−31.0%)

Supplementary Material

1179

"Feasibility of enhancing carbon sequestration and stock capacity in temperate and boreal European forests via changes to management regimes"

1180

1181

1182

Dalmonech et al.

1183

1184

1185

1186

1187

1188

Figure S1 | Evaluation of seasonal GPP (gross primary productivity) fluxes. Top row shows
the seasonal course of average daily GPP (g C m^{-2} day^{-1}) over the years (DoY = Day of the
Year). Blue shaded area represents the maximum and minimum bounds of GPP values among
the five ESMs used to force the model (GPP ESM clim.), red line and the red shaded area
represent the average and the maximum and minimum GPP values when the model is forced
by observed climate (GPP Obs. clim.), and black dots represent the quality-checked and -
filtered GPP values evaluated at the sites by the eddy covariance technique (GPP Meas.) and
the shaded area of the inter-annual variability. ESMs = model forced with Earth System
Models climate. Performance statistics are reported in Table S2.

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

Figure S2 | Comparison of the predicted annual GPP (gross primary productivity, gC m^{-2}
day^{-1}, above panel) and NPP$_{woody}$ (the annual NPP allocated to woody pools, gC m^{-2} day^{-}1,
below panel) with site observations at the three sites modelling real stands. The blue shaded
area represents the maximum and minimum bounds of GPP and NPP$_{woody}$ values among the
five ESMs used to force the model; the red line represents the average NPP$_{woody}$ values when
the model is forced by observed climate. Black dots represent the measured GPP and
NPP$_{woody}$ values (i.e. GPP Meas. and NPP$_{woody}$ Meas.), and the gray shaded area represents
the relative uncertainty bounds. At the Hyytiälä site NPP$_{woody}$ observed data for the year 2002
was missing. OT = observed thinning at the site; ESM = model forced with Earth System
Models climate.

1218

1219

1220

1221

1222

1223

Figure S3 | Comparison of the predicted annual DBH (stem diameter at breast height in cm) with site observations at the three sites. The shaded area represents the maximum and minimum bounds of DBH values among the five ESMs used to force the model; the red line represents the average DBH values when the model is forced by observed climate (DBH Obs. clim.). Black dots represent the measured DBH values. OT = observed thinning at the sites; ESMs = model forced with Earth System Models climate.

1231

1232

1233

Figure S4 | Box plots of the differences for pCWS (potential C Woody Stocks), (a) computed
with respect to the unmanaged scenario NO-MAN (dashed line) under different RCPs and
NPP (net primary production) (b). Management scenarios are reported according to changes
in rotation turn (*r*), thinning frequency (*th*) and thinning intensity (*thi*) and according to the
AM+ scenarios (more intensive than BAU) and AM− (less intensive than BAU) reported in
Table S1. The management key variables are changed with respect to the BAU scenario
(r+20 th+10 thi −10, r+00 th+10 thi 00 , r+00 th+00 thi −10, r+20 th+00 thi+00 belongs to the
less intensive scenario AM−; r–20 th+00 thi+00, r+00 th+00 thi+10, r+00 th–10 thi+00, r–20
th–10 thi+10 belongs to the more intensive scenario AM+). Scenarios from the BAU are
ordered following the most important management variable driving the fluxes and stocks

variability, i.e. thinning frequency and rotation (data not shown). For the sake of clarity we
reported 8 out of 14 scenarios in the figure. However, the main pattern across intensities does
not change.

1247

Figure S5 | Percentage of changes for Mean NPP (net primary productivity) and pCWS
(potential Carbon Woody Stocks) over the period 2006-2099 between the managed scenarios
AM+, AM–, MIX, NO-MAN (AM+ , more intensive than BAU; AM–, less intensive than
BAU; MIX: mixes options according to Table S1, NO-MAN: no management option) and the
BAU (business–as–usual) scenario reported for the 4 RCPs. Values refer to data averaged
across real and virtual stands and across species. Note: each single scenario according to
Table S1 is here reported (28 in total). Parametric curve fitting (Polynomial of order 3) is also
reported for each RCP and it is computed according to the averaged data for each
management intervention scenario and RCP.

1257

1258

1259

1260

Figure S6 | Factorial analysis of the factors driving the variability of GPP, NPP, R_a, CUE, NPP_{woody} and pCWS, for the three study sites Bílý Křìž (square), Hyytiälä (triangle), Sorø (circle) and an average among sites (rhombus); ESMs = model forced with Earth System Models climate, RCP = emission scenarios, 'Stand' = virtual stands of the Composite Forest Matrix, i.e. simulations from each individual virtual stand, characterized by a specific age class and structure, are considered.

1267

1268

Figure S7 | Potential harvested woody Biomass (pHPW) simulations under the management scenarios (AM+, BAU, AM–) and the NO-MAN scenario divided by different emission scenario RCPs. pHWP, solid line, is averaged across the representative forests, different ESMs and aggregated according to the management regime. Yearly mean values and 5[th] and 95[th] percentiles are reported.

1274

1275

1276

1277
1278 **Figure S8** | Standing woody biomass simulations under the management scenarios (AM+,
1279 BAU, AM–) and the NO-MAN scenario divided by different emission scenario RCPs.
1280 Standing woody biomass, solid line, is averaged across the representative forests, different
1281 ESMs and aggregated according to the management regime. Yearly mean values and 5[th] and
1282 95[th] percentiles are reported.

1283

1284

1285

1286

Figure S9 | Average NPP (net primary productivity, g C m^{-2} year^{-1}) and pCWS (the sum of
standing and potential harvested woody products; t C ha^{-1}) over the period 2006-2099, for the
three groups of management scenarios: AM+, AM−, BAU; and the NO-MAN reported for the
4 RCPs at the site of Sorø. Reported values refer to data averaged across real and virtual
stands. Data ellipses are also reported in shaded colors and refer to all data. NOTE: each
single scenario according to Table S1 is reported here (16 in total). In the subplot the
differences are expressed as % and are reported along a parametric curve (third order
polynomial) with the point (0, 0) representing the reference BAU. Arrows indicate the
increasing intensity of management intervention.

1296

1297

Figure S10 Average NPP (net primary productivity, g C m^{-2} year^{-1}) and pCWS (the sum of standing and potential harvested woody products; t C ha^{-1}) over the period 2006-2099, for the three groups of management scenarios: AM+, AM−, BAU; and the NO-MAN reported for the 4 RCPs at the site of Hyytiälä. Reported values refer to data averaged across real and virtual stands. Data ellipses are also reported in shaded colors and refer to all data. NOTE: each single scenario according to Table S1 is reported here (16 in total). In the subplot the differences are expressed as % and are reported along a parametric curve (third order polynomial) with the point (0, 0) representing the reference BAU. Arrows indicate the increasing intensity of management intervention.

1307

1308

Figure S11 Average NPP (net primary productivity, g C m^{-2} year^{-1}) and pCWS (the sum of standing and potential harvested woody products; tC ha^{-1}) over the period 2006-2099, for the three groups of management scenarios: AM+, AM−, BAU; and the NO-MAN reported for the 4 RCPs at the site of Bílý Kříž. Reported values refer to data averaged across real and virtual stands. Data ellipses are also reported in shaded colors and refer to all data. NOTE: each single scenario according to Table S1 is reported here (16 in total). In the subplot the differences are expressed as % and are reported along a parametric curve (third order polynomial) with the point (0, 0) representing the reference BAU. Arrows indicate the increasing intensity of management intervention.

1318

1319

1320

1321

1322

Supp. Tables

Table S1 | Full experimental design of management scenarios: 26 alternative scenarios, plus an unmanaged scheme (NO-MAN) where trees are removed only through mortality, each relative to the business-as-usual scheme. Scenarios ranged from more limited interventions, grouped under the label AM–, to more intensive silvicultural practice, grouped under the label AM+. The cutting scenarios are a combination of three different thinning intensities at three different thinning intervals and three different rotation ages (e.g. rotation = +20, thinning intensity = 10, thinning regime = –10 indicate increasing the rotation period by 20 years with respect to the BAU setting, increasing the thinning interval by 10 years relative to the BAU setting, and decreasing the intensity of removal by 10% relative to the BAU setting (Table 1). A mixed combination of scenarios, grouped under the label MIX, was also simulated and indicated where the parameter changes follow opposite directions (e.g.

1335 increase thinning intensity in combination with prolonged rotation period).

Management type	rotation years	thinning regime years	thinning intensity %
AM−	20	10	−10
AM−	0	10	−10
AM−	0	10	0
AM−	20	10	0
AM−	20	0	−10
AM−	0	0	−10
AM−	20	0	0
BAU	0	0	0
AM+	−20	0	0
AM+	0	0	10
AM+	−20	0	10
AM+	−20	−10	0
AM+	0	−10	0
AM+	0	−10	10
AM+	−20	−10	10
NO−MAN	/	/	/
MIX	−20	−10	−10
MIX	−20	0	−10
MIX	−20	10	−10
MIX	−20	10	0
MIX	−20	10	10
MIX	0	−10	−10
MIX	0	10	10
MIX	20	−10	−10
MIX	20	−10	0
MIX	20	−10	10
MIX	20	0	10
MIX	20	10	10

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348 **Table S2 |** Performance statistics (Pearson correlation coefficient r, absolute and relative root
1349 mean square error RMSEabs (g C m^{-2} day^{-1}) and RMSErel (%); mean absolute error, MAE
1350 (g C m^{-2} day^{-1}) computed from daily series of model gross primary productivity, GPP against
1351 eddy covariance data when model is forced with measured (LOCAL) and modelled climate
1352 data (including the ensemble mean across ESMs climate data).

Site	Period	Forcing	RMSEabs gC m^{-2} day^{-1}	RMSErel %	MAE gC m^{-2} day^{-1}	n	r
Hyytiälä	1997-2005	ESM1	1.98	0.44	1.34	3084	0.82 ***
Hyytiälä	1997-2005	ESM2	1.79	0.4	1.21	3084	0.86 ***
Hyytiälä	1997-2006	ESM3	1.75	0.39	1.2	3084	0.85 ***
Hyytiälä	1997-2007	ESM4	2.12	0.48	1.38	3084	0.78 ***
Hyytiälä	1997-2008	ESM5	1.93	0.43	1.3	3084	0.84 ***
Hyytiälä	1997-2009	ENSEMBLE	1.91	0.43	1.29	3084	
Hyytiälä	1997-2005	LOCAL	1.48	0.33	1.05	3084	0.92 ***
Bílý Křiž	2000-2005	ESM1	2.65	0.46	1.98	1390	0.74 ***
Bílý Křiž	2000-2005	ESM2	2.6	0.45	1.93	1390	0.75 ***
Bílý K?iž	2000-2006	ESM3	2.63	0.45	1.97	1390	0.72 ***
Bílý K?iž	2000-2007	ESM4	2.89	0.5	2.1	1390	0.67 ***
Bílý K?iž	2000-2008	ESM5	2.68	0.46	2	1390	0.72 ***
Bílý K?iž	2000-2009	ENSEMBLE	2.69	0.46	1.99	1390	
Bílý Křiž	2000-2005	LOCAL	2.07	0.36	1.52	1390	0.87 ***
Sorø	1997-2005	ESM1	2.78	0.37	1.78	3092	0.89 ***
Sorø	1997-2005	ESM2	2.81	0.37	1.77	3092	0.89 ***
Sorø	1997-2006	ESM3	3.07	0.41	1.98	3092	0.88 ***
Sorø	1997-2007	ESM4	3.25	0.43	2.08	3092	0.86 ***
Sorø	1997-2008	ESM5	3.01	0.4	1.91	3092	0.88 ***
Sorø	1997-2009	ENSEMBLE	2.98	0.4	1.91	3092	
Sorø	1997-2005	LOCAL	2.15	0.29	1.43	3092	0.94 ***

1353

1354

1355

1356

1357

1358

1359

1360

1361

1362

Table S3 | Modeled and Estimated (bold) and modeled values of net primary productivity
(NPP), plant autotrophic respiration (R_a), and carbon use efficiency (CUE, i.e. NPP/GPP).
Model output reports the annual average of the variables over the observation period.
'LOCAL-Climate' and 'ESM-Climate' refer to model output when driven by locally
observed or climate-modeled data. (a = Wu et al., 2013; Knohl et al., 2008; Vanninen and
Mäkelä 2005; Tang et al., 2014; b = Granier et al., 2008; Luyssaert et al., 2007; c = Xiao et
al., 2003; Vanninen and Mäkelä 2005; Luyssaert et al., 2007; Tang et al., 2014)

	Sorø	Bílý Křiž	Hyytiälä
NPP [a] gC m^{-2} year^{-1}	708±65	611±45	434±55
LOCAL	860.35	547.31	454
ESMs' ENSEMBLE	734±32	517±28	416±26
R_a [b] gC m^{-2} year^{-1}	751±52	726±110	558±24.5
LOCAL	945.11	662.3	464.8
ESMs' ENSEMBLE	973±16	697±30	471±14
CUE [c] adim.	0.45-0.50	0.45-0.5	0.32/0.45-0.54
LOCAL	0.47	0.45	0.49
ESMs' ENSEMBLE	0.42±0.01	0.42±0.01	0.47±0.01

1381

1382

1383

Table S4 | NPP and pCWS at the sites of Sorø, Bílý Křiž and Hyytiälä computed as
percentage differences between the alternative management scenarios and the *Business-As-*
Usual one are reported in parenthesis. Values refer to data averaged over the simulation
period 2006-2099, across real and virtual standard climate for each RCPs. AM+ = more
intensive than the *businnes-as-usual*; AM– = less intensive than the *businnes-as-usual*; NO-
MAN = no management; RCPs = Representative Concentration Pathways. In bold the
maximum and the minimum values across the RCPs and management type scenarios divided
per site.

1392

Site	Management type	RCP	NPP %	pCWS %
Sorø	AM+	2.6	-30.9	**-6.7**
	AM+	4.5	-30.2	-7.2
	AM+	6.0	-30.3	-7.2
	AM+	8.5	-28.8	-7.4
	AM-	2.6	**-2.2**	-7.2
	AM-	4.5	-2.7	-7.4
	AM-	6.0	-2.6	-7.3
	AM-	8.5	-3.4	-7.5
	NO-MAN	2.6	-18.8	-30.0
	NO-MAN	4.5	-19.7	-29.9
	NO-MAN	6.0	-19.3	-29.9
	NO-MAN	8.5	-21.7	-30.2
Bílý Křiž	AM+	2.6	-22.6	1.5
	AM+	4.5	-21.6	1.9
	AM+	6.0	-21.7	1.4
	AM+	8.5	-19.1	**3.0**
	AM-	2.6	**-8.2**	-8.9
	AM-	4.5	-10.3	-10.0
	AM-	6.0	-9.5	-9.6
	AM-	8.5	-11.2	-10.4
	NO-MAN	2.6	-36.6	-35.8
	NO-MAN	4.5	-39.6	-38.1
	NO-MAN	6.0	-39.0	-38.0
	NO-MAN	8.5	**-41.7**	**-40.0**
Hyytiälä	AM+	2.6	-37.7	-7.0
	AM+	4.5	-37.6	-7.4
	AM+	6.0	**-37.9**	-7.7
	AM+	8.5	-37.1	-7.1
	AM-	2.6	9.2	-3.8
	AM-	4.5	7.7	-4.1
	AM-	6.0	7.6	**-3.9**
	AM-	8.5	5.6	-4.2
	NO-MAN	2.6	**21.6**	-21.6
	NO-MAN	4.5	16.4	-22.5
	NO-MAN	6.0	15.9	-22.1
	NO-MAN	8.5	9.0	**-23.7**

1393

 Table S5 | Percentage of changes for Mean NPP (net primary productivity) and pCWS
 (potential C woody stocks) over the period 2006-2099, between each alternative management
 scenario, the NO-MAN (no-management) scenario and the *business-as-usual scenarios*
 (BAU). Values refer to data averaged across real and virtual stands, species and climate for
 each RCPs. Positive values indicate a positive effect of one management scheme compared to
 BAU. ESMs = Earth System Models; RCPs = Representative Concentration Pathways, r =
 rotation turn, *th* = thinning frequency, *thi*= thinning intensity expressed as year or %
 differences compared to the BAU scheme, MIX = mixed scenario. In bold the maximum and
 the minimum values across the RCPs and management type scenarios.

Management type	Rotation	Thinning regime	Thinning intensity	NPP%				pCWS%			
				RCP 2.6	RCP 4.5	RCP 6.0	RCP 8.5	RCP 2.6	RCP 4.5	RCP 6.0	RCP 8.5
	years	years	%								
NO-MAN	/	/	/	**-14.47**	-16.42	-16.16	**-19.46**	**-29.40**	-30.16	-30.04	**-31.08**
AM−	20	10	−10	-3.33	-4.81	-4.61	-6.17	-11.49	-12.19	-12.10	-12.38
AM−	0	10	−10	-3.52	-5.17	-4.75	**-6.65**	-12.86	-13.60	-13.27	**-13.92**
AM−	0	10	0	-1.04	-2.08	-1.94	-3.23	-7.66	-8.11	-8.03	-8.48
AM−	20	10	0	-1.10	-2.03	-1.81	-2.88	-5.81	-6.27	-6.12	-6.42
AM−	20	0	−10	0.50	-0.24	-0.05	-1.28	-4.93	-5.20	-5.03	-5.49
AM−	0	0	−10	**0.66**	-0.16	-0.04	-1.39	-7.23	-7.46	-7.33	-7.82
AM−	20	0	0	-0.98	-0.87	-0.80	-0.70	2.46	2.41	2.46	**2.58**
AM+	−20	0	0	-4.60	-4.41	-4.41	**-4.26**	0.59	0.61	0.63	0.79
AM+	0	0	10	-6.36	-5.69	-5.75	-4.51	3.57	3.66	3.56	**4.14**
AM+	−20	0	10	-10.97	-10.37	-10.70	-9.34	3.23	3.34	2.88	3.72
AM+	−20	−10	0	-40.21	-39.58	-39.75	-37.63	-4.75	-5.14	-5.36	-4.83
AM+	0	−10	0	-38.90	-38.20	-38.27	-36.16	-5.51	-5.87	-6.05	-5.51
AM+	0	−10	10	**-55.46**	-55.09	-55.25	-53.56	-14.91	-15.43	**-15.65**	-15.50
AM+	−20	−10	10	-55.20	-54.82	-54.99	-53.36	-13.46	-13.97	-14.22	-14.02
BAU	0	0	0	/	/	/	/	/	/	/	/
MIX	−20	−10	−10	-18.73	-17.83	-18.10	-16.05	3.54	3.49	3.01	4.03
MIX	−20	0	−10	-3.05	-3.83	-3.45	-4.65	-5.20	-5.50	-5.21	-5.75
MIX	−20	10	−10	-6.51	-7.98	-7.25	-9.46	-10.15	-10.85	-10.23	-11.53
MIX	−20	10	0	-4.58	-5.46	-5.16	-6.49	-5.51	-5.96	-5.79	-6.53
MIX	−20	10	10	-5.07	-5.40	-5.03	-5.87	-1.71	-1.95	-1.67	-2.50
MIX	0	−10	−10	-16.11	-15.19	-15.73	-13.29	3.25	3.25	2.36	3.87
MIX	0	10	10	**-0.86**	-1.36	-1.09	-1.95	-3.00	-3.27	-3.05	-3.62
MIX	20	−10	−10	-19.57	-18.88	-18.83	-17.19	3.72	3.62	3.56	4.12
MIX	20	−10	0	-42.47	-41.97	-42.11	-40.15	-5.67	-6.12	-6.31	-5.95
MIX	20	−10	10	**-58.40**	-58.12	-58.25	-56.72	-15.32	-15.93	**-16.13**	-16.08
MIX	20	0	10	-8.69	-7.92	-8.15	-6.52	5.08	5.13	4.72	**5.67**
MIX	20	10	10	-1.38	-1.56	-1.46	-1.99	-0.80	-0.93	-0.82	-1.18

1412

1413

1414

1415

1416

1417

1418

1419

1420 **Table S6 |** Factorial analysis (values range from 0 to 100) for the described variables (GPP,
1421 R_a, NPP, CUE, NPP_{woody} and potential Carbon woody stock) across sites and the mean
1422 between sites.

Factor	Bílý Kříž - GPP	Bílý Kříž - NPP	Bílý Kříž - R_a	Bílý Kříž - CUE	Bílý Kříž - NPP_{woody}	Bílý Kříž - pCWS
ESM	11.79	19.33	7.75	11.93	13.63	11.90
ESM-RCP	3.78	12.47	3.76	11.57	8.39	1.16
ESM–RCP–Stand	2.94	2.37	2.51	2.28	2.13	1.75
ESM–Stand	2.00	2.00	1.84	1.40	1.98	1.11
RCP	1.12	3.40	1.34	4.57	2.19	0.25
RCP–Stand	0.80	0.76	0.71	0.60	0.67	0.48
Stand	77.56	59.67	82.10	67.64	71.01	83.35
Error	0.00	0.00	0.00	0.00	0.00	0.00
	Hyytiälä - GPP	Hyytiälä - NPP	Hyytiälä - R_a	Hyytiälä - CUE	Hyytiälä - NPP_{woody}	Hyytiälä - pCWS
ESM	3.21	5.15	4.31	9.55	7.04	2.67
ESM-RCP	0.90	4.45	1.16	7.08	6.46	0.23
ESM–RCP–Stand	0.15	0.53	0.18	0.38	0.91	0.05
ESM–Stand	0.39	0.92	0.51	0.57	1.37	0.26
RCP	3.11	2.41	3.65	4.11	3.17	0.36
RCP–Stand	0.33	0.51	0.44	0.19	0.72	0.05
Stand	91.90	86.04	89.76	78.12	80.33	96.38
Error	0.00	0.00	0.00	0.00	0.00	0.00
	Sorø - GPP	Sorø - NPP	Sorø - R_a	Sorø - CUE	Sorø - NPP_{woody}	Sorø - pCWS
ESM	21.07	21.54	15.11	8.47	14.13	15.03
ESM-RCP	10.07	17.88	5.22	9.42	11.84	0.42
ESM–RCP–Stand	0.31	0.68	0.22	0.49	0.60	0.03
ESM–Stand	0.83	1.33	0.94	0.85	1.45	0.27
RCP	8.69	8.90	6.99	3.25	6.45	1.54
RCP–Stand	0.41	0.64	0.51	0.37	0.76	0.07
Stand	58.64	49.03	71.01	77.15	64.77	82.65
Error	0.00	0.00	0.00	0.00	0.00	0.00
	ALL - GPP	ALL - NPP	ALL - R_a	ALL - CUE	ALL - NPP_{woody}	ALL - pCWS
ESM	12.02	15.34	9.06	9.98	11.60	9.87
ESM-RCP	4.92	11.60	3.38	9.36	8.90	0.60
ESM–RCP–Stand	1.13	1.19	0.97	1.05	1.22	0.61
ESM–Stand	1.07	1.42	1.09	0.94	1.60	0.54
RCP	4.31	4.90	4.00	3.98	3.94	0.72
RCP–Stand	0.51	0.63	0.55	0.39	0.72	0.20
Stand	76.03	64.91	80.96	74.31	72.04	87.46
Error	0.00	0.00	0.00	0.00	0.00	0.00

1423