TITLE

A multispecies framework for modeling adaptive immunity and immunotherapy in cancer

AUTHORS

Timothy Qi[1], Benjamin Vincent[2,3], and Yanguang Cao[1,2]*

AFFILIATIONS

[1]Division of Pharmacotherapy and Experimental Therapeutics, Eshelman School of Pharmacy,

The University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA

[2]Lineberger Comprehensive Cancer Center, The University of North Carolina at Chapel Hill,

Chapel Hill, NC 27599, USA

[3]Division of Hematology/Oncology, School of Medicine, The University of North Carolina at

Chapel Hill, Chapel Hill, NC 27599, USA

***Lead Contact**. Email: yanguang@unc.edu. Address: 301 Pharmacy Ln, Chapel Hill, NC 27599

WORD COUNT: 4,986

ABSTRACT

Predator-prey theory is commonly used to approximately describe tumor growth in the presence of selective pressure from the adaptive immune system. These interactions are mediated by the tumor immunopeptidome (what the tumor "shows" the body) and the T-cell receptor (TCR) repertoire (how well the body "sees" cancer cells). Importantly, both tumor and T cell populations are dynamic and in competition with each other and their fraternal lineages. In particular, the immunopeptidome comprises neoantigens which can be gained and lost throughout tumorigenesis and treatment. Heterogeneity in the immunopeptidome is predictive of poorer survival and response to T-cell dependent immunotherapy in some tumor types, suggesting the TCR repertoire is unable to support a fully polyclonal response against every neoantigen. Whether between-lineage competition among T cells plays a role, and in what contexts, is unknown; moreover, longitudinal TCR profiling studies are expensive and logistically complex to conduct. In silico models may offer an inexpensive way to interrogate these phenomena ex ante and deepen our understanding of the tumor-immune axis. Here, we construct and calibrate a predator-prey-like model to preclinical and clinical data to describe tumor growth and immunopeptidome diversification. Simultaneously, we model the expansion of neoantigen-specific T cell lineages and their consumption of both lineage-specific and shared, lineage-agnostic resources. This predator-prey-like framework accurately described clinically observed immunopeptidomes as well as correlates and consequences of response to immunotherapy, including immunoediting. The model was also suitable for exploring treatment of tumors with varying growth and mutation rates.

INTRODUCTION

Cancer is a disease with parallels to predator-prey theory [1]. Control of prey (cancer

cells) by predators (cytotoxic T cells, CTLs) depends upon the latter recognizing and killing the

former. This process is mediated by the interaction between neoantigens on the surface of

cancer cells and T cell receptors (TCRs) on the surface of CTLs. Neoantigens and TCRs that

bind each other are called cognate; cognate interactions trigger CTLs to clonally expand,

producing daughter cells that express the same TCR. At the molecular level, neoantigens arise

from tumor-specific alterations to DNA or RNA and are bound by major histocompatibility

complex (MHC) molecules for extracellular presentation to TCRs (Figure 1A) [2], [3]. The myriad

neoantigens presented by a population of cancer cells, exclusive of self antigens, is referred to

here as the immunopeptidome, while the universe of TCRs expressed by all CTLs in the body is

referred to here as the TCR repertoire. Importantly, while one cancer cell can present a

theoretically unlimited number of neoantigens, individual T cells express TCRs of only a single

specificity.

CTLs kill cancer cells that present their cognate neoantigen. The propensity of a

neoantigen to cause this is referred to as its immunogenicity. Numerous factors influence

immunogenicity, including several key parameters reported by the global community-based

Tumor Neoantigen Selection Alliance [4]. These parameters comprise (1) MHC binding affinity,

(2) MHC binding stability, or duration of peptide-MHC interaction, (3) agretopicity, or the ratio of

mutant to wild-type binding affinity, (4) RNA transcript abundance within the tumor, and (5)

foreignness, or homology to epitopes of known pathogens. These parameters and their

associated cutoffs associated with immunogenicity are referred to here as the TESLA criteria.

The heritability of both neoantigens and their cognate TCRs over cellular generations is

important to consider when evaluating anti-neoantigen immunity. For one, clonal expansion of

distinct cancer and T cell lineages is common. Tumors enriched in clonally expressed

neoantigens are more highly infiltrated by CTLs, particularly those with cognate TCR

clonotypes. Highly clonal neoantigen burden correlates with longer survival in patients treated

with T cell-dependent cancer immunotherapies, such as immune checkpoint blockade (ICB) [5],

[6]. Furthermore, a recent pan-cancer analysis found clonal tumor mutation burden (TMB), a

measure closely related to clonal neoantigen burden, to be among the strongest predictors of

response to ICB [7].

Neoantigen clonality plays a particularly important role in differentiating between intrinsic

and apparent immunogenicity. When an antigen is subclonally expressed below a certain

fraction, lineages presenting these antigens may not be efficiently eliminated by the immune

system even if the antigen has high intrinsic immunogenicity (e.g., chicken ovalbumin antigen, a

highly foreign model antigen commonly used in murine studies) (Figure 1B) [8], [9]. Mechanisms

for subclonal neoantigen include not only de novo generation, but also through (epi)genomic

silencing of extant neoantigens and/or their associated MHC alleles [10], [11]. In humans,

studies across multiple cancer types have found that among hundreds of putative neoantigens

identified by bioinformatics pipelines, very few (generally < 4) are bona fide immunogenic when

tested in ex vivo assays; moreover, those that are immunogenic tend also to be clonal [4], [5],

[12]–[15].

It is unclear why the TCR repertoire does not support a fully polyclonal response against

every neoantigen. We speculate this limitation could be due to the spatiotemporally restricted

availability of resources and nutrients critical to CTL survival and proliferation [16]. For example,

cytokines, co-stimulatory signals from antigen-presenting cells, and other factors beyond direct

antigen-mediated signaling through TCRs are a shared resource among CTL lineages (Figure

1C) [16], [17]. The asymmetric and possibly contemporaneous presentation of multiple

neoantigens might exhaust these nonspecific stimuli, leading to an inadequate immune

response to subclonal neoantigens. In humans, the clonal expansion of antigen-reactive CTL

lineages can indeed suppress the expansion of other CTL lineages, a phenomenon called

immunodominance that is well characterized within virology [18]–[20]. In cancer, ICB has been

widely reported to affect the diversity of the CTL TCR repertoire, although whether it biases CTL

expansion toward polyclonal or oligoclonal states appears to vary by ICB target and tumor

histology [21]. However, given the shared nature of nonspecific stimuli, highly clonal tumors

might be more susceptible to CTL-based immunotherapies due to reduced interclonal

competition for nonspecific stimuli as compared to highly subclonal tumors rife with lower-

quality, "distracting" neoantigens [16].

Neoantigen quantity is therefore critical to consider alongside immunogenicity for

immune recognition and ICB treatment efficacy [15], [22]. However, the quantitative relationship

between immunopeptidome diversity, CTL diversity, and response to ICB is open to further

characterization. In this work, we construct a mathematical model of nonlinear predator-prey-like

dynamics to describe the relationship between immunopeptidome diversity and CTL responses

while accounting for interclonal CTL competition for nonspecific stimuli. Tumor growth and

immunopeptidomics are modeled and calibrated with experimental data from two different

species and three types of cancer, including an orthotopic murine model of bladder cancer, a

murine model of colorectal cancer, a 100-patient cohort of NSCLC patients who underwent

extensive immunopeptidome profiling [10], [11], [23]–[25]. Simulated NSCLC-like

immunopeptidomes and TCR repertoires are cross-validated on an independent cohort [26] and

subject to a local sensitivity analysis. Finally, tumor response dynamics in the NSCLC-like

cohort, as well as in two additional cohorts resembling other tumor types, are simulated under

ICB treatment to evaluate pre-treatment biomarkers of response within the immunopeptidome

and the TCR repertoire.

1 **METHODS**

2 **Tumor growth dynamics**

3   Tumors grew according to a modified logistic growth model, where N is the number of

4 cells in the tumor, k_g is the crude birth rate, K is the population carrying capacity, $k_{d,i}$ is the

5 lineage-specific death rate (see *Lineage-specific killing*), and j is number of unique cancer cell

6 lineages in the tumor:

8     **Equation 1**: Modified logistic growth model describing tumor growth

9
$$\frac{dN}{dt} = \sum_{i=1}^{j} k_g \left(1 - \frac{N_i}{K}\right)\left(1 - k_{d,i}\right)$$

10
$$N_{t=0} = \begin{cases} 4.77 \ x \ 10^7, if \ mouse \\ 1, if \ human \end{cases}$$

12   All cancer cell lineages were assigned the same birth rate and shared population-level

13 carrying capacity. Death rates varied by lineage and depended on the number of CTLs cognate

14 to the lineage's neoantigens (see respective sections on *CTL clonal dynamics* and *Neoantigen*

15 *features*). Cancer cell numbers were converted into tumor diameters assuming spherical tumors

16 and an approximate cell volume of 4 pL [27].

17   During mouse simulations, tumors were initiated from a single lineage ~$4.77 \ x \ 10^7$ cells,

18 corresponding to an approximate volume of 200 mm^3 at which tumor measurements began in

19 [23], [28]. During human simulations, tumors were initiated from a single founder cell with a

20 gamma-distributed number of neoantigens [10]. Furthermore, the founder cell was assigned a

21 14% chance of having an MHC loss event [11] in one of its three MHC alleles. Further details on

22 neoantigen immunogenicity, expression, and MHC presentation are provided in the *Neoantigen*

23 *features* section. Parameter values, definitions, and their sources are provided in Table S1.

CTL clonal dynamics

At baseline, we assumed a tumor-ignorant and nonspecific CTL population equal to the steady-state population of T cells CTL_{ss} in either mice or humans (Table S1). With the number of unique neoantigens k known from either a founder lineage (mice) or cell (humans), we then created k tumor-reactive CTL lineages (Figure 2). For simplicity, we assume TCRs are specific to a single neoantigen, which aligns with the high specificity of TCRs despite their considerable cross-reactive potential [29]. For each neoantigen among k neoantigens, a CTL lineage of 2^n cells was created, where n is a Poisson-distributed number $n \sim P(\mu)$. This represents the random number of divisions the founder CTL cell underwent prior to thymic efflux [30]. The remaining nonspecific CTLs were lumped into a tumor-ignorant, nonspecific population with a starting cell number equal to CTL_{ss} minus the total number of neoantigen-reactive CTLs.

Tumor-specific CTL lineages grew or shrank according to the quantity of their cognate antigenic stimulus and the availability of shared nonspecific stimuli, where CTL_k is the kth CTL lineage reactive to the kth neoantigen, T_d is its baseline turnover rate, C is the abundance of nonspecific stimulus, S_k is the immunogenicity-weighted number of cancer cells expressing the kth neoantigen (see *Neoantigen features*), and T_a is a scaling factor relating the rate of CTL_k expansion per cancer cell expressing the kth neoantigen:

Equation 2: Clonal dynamics of neoantigen-reactive CTL lineages

$$\frac{dCTL_k}{dt} = CTL_k \left(T_d(C-1) + \frac{S_k T_a C}{CTL_k} \right)$$

$$CTL_{k,t=0} = 2^{P(\mu)}$$

Nonspecific CTLs presented with no antigen stimulus grew or shrank according only to the availability of shared nonspecific stimuli, where $NSCTL$ is the number of non-specific CTLs:

Equation 3: Growth dynamics of nonspecific CTLs

$$\frac{dNSCTL}{dt} = NSCTL\big(T_d(C-1)\big)$$

$$NSCTL_{t=0} = \begin{cases} 10^7, if\ mouse \\ 10^{11}, if\ human \end{cases}$$

Nonspecific stimulus C was consumed by T cells and non-T cells and replenished at rate k_{in}. The fraction of C consumed by non-T cells was f_{out}. C therefore decreased when the total number of nonspecific and tumor-reactive CTLs was greater than CTL_{ss} and increased when it was less than CTL_{ss}:

Equation 4: Availability of nonspecific stimulus

$$\frac{dC}{dt} = k_{in} - k_{in}f_{out} - k_{in}(1 - f_{out})\left(\frac{NSCTL + \sum_{i=1}^{k} CTL_k}{CTL_{ss}}\right);$$

$$C_{t=0} = 1;\ C = \{C \in \mathbb{R} : C \geq 0\}$$

Shannon entropy among the tumor-reactive CTL lineages was calculated using MATLAB's *wentropy* function.

Neoantigen features

Each neoantigen was assigned a non-negative real number R between 0 and 1 to describe its immunogenicity. In the mouse simulations, R for each of the 34 neoantigens identified in BBN963 cells in [23] was assigned by scaling the baseline-corrected ELISpot scores to the range [0, 1].

In the human simulations, R was assigned for each neoantigen in the founder cell by first sampling values for each of the TESLA criteria. Neoantigens were first assigned random values for MHC binding affinity, MHC binding stability, agretopicity, transcript abundance, and

foreignness drawn from distributions defined by 146 NSCLC-derived peptides (Figure S1) [4].

To achieve this, we fit an exponential distribution to transcript abundance and a zero- and one-

inflated binomial distribution to foreignness. To account for correlation among MHC binding

affinity, MHC binding stability, and agretopicity values, we fit a Gaussian copula to log-

transformed values using MATLAB's *copulafit* function. Parameters associated with these fits

are provided in Table S1. Transcript abundance was discarded due to conflation with clonality

and the remaining four parameters were assessed against the following TESLA criteria: MHC

binding affinity < 34 nM, MHC binding stability > 1.4 hours, agretopicity > 0.1, and foreignness <

10^{-16}. Neoantigens that failed to satisfy these cutoffs were assigned $R = 0$, or no

immunogenicity. Neoantigens that passed these cutoffs were assigned a value for R from an

exponential distribution:

13   **Equation 5:** Empirically determined neoantigen immunogenicity distribution

14                                   $$R \sim Exp(-6x)$$

15                                   $$1 \geq x \geq 0$$

17   In addition to intrinsic immunogenicity, each neoantigen was also assigned values to

indicate its expression status and MHC binding partner. In all cases, neoantigens were

assumed to be expressed at baseline. Murine simulations assumed each neoantigen could bind

to one of two MHC alleles (H2-Db and H2-Kb), both of which were intact and expressed at

baseline. Human simulations assumed neoantigen binding to one of three MHC alleles (HLA-A,

HLA-B, and HLA-C), all of which were expressed at baseline, except in ~14% of tumors where

loss of one of three alleles at random conferred clonal MHC loss [11]. Thus, inclusion of a

neoantigen in its parent cell's immunopeptidome depended upon its own expression status as

well as the intactness of its binding MHC allele (Figure S1).

S_k, the immunogenicity-weighted number of cancer cells with neoantigen k in their

immunopeptidome, was used to stimulate the proliferation of CTL_k per *CTL clonal dynamics*. It

was calculated by multiplying the intrinsic immunogenicity of a neoantigen k by the number of

cells whose immunopeptidomes include it, where E_k is the lineage's expression status of

neoantigen k, MHC_k is the lineage's expression status of the MHC binding partner of neoantigen

k, and R_k is the intrinsic immunogenicity of neoantigen k:

8   **Equation 6**: Neoantigen-driven CTL expansion

$$S_k = E_k MHC_k R_k$$

In mouse simulations, cancer cell lineage's immunopeptidomes were held constant

within the short simulated timeframe due to the lack of data to inform event rates. In human

simulations, stochastic modifications to a cancer cell lineage's immunopeptidome were allowed

to occur. These events were one of three types: (1) neoantigen gain, (2) neoantigen loss, and

(3) MHC allele loss. Neoantigen gain events, representing the generation of new and heritable

neoantigens within the DNA or RNA, involved drawing TESLA parameters and assigning R

immunogenicity values for each new neoantigen as described above. Neoantigen loss events,

representing the genetic, epigenetic, transcriptomic, or metabolic suppression of a neoantigen,

were represented as an irreversible change in the neoantigen's expression from 1 to 0;

functionally, this removed the neoantigen from the lineage's immunopeptidome and reduced its

S_k to 0. MHC loss events, analogous to neoantigen loss events for MHC alleles, were

represented as an irreversible change in one of the three MHC alleles' expression from 1 to 0.

Functionally, this removed all neoantigens presented by that MHC allele from the lineage's

immunopeptidome and reduced their S_k to 0. Lineages that experienced one of these stochastic

events were deducted 1 cell from their population size. An identical lineage save the stochastic

event was then created at N=1.

Lineage-specific killing

Each unique clone in the tumor was killed at an independent and saturable rate

calculated from the contents of its immunopeptidome and the abundance of cognate, reactive

CTLs. Here, $k_{d,i}$ is a dimensionless value used to modify the logistic growth described in

Equation 1; f_{die} is the fraction of cell death not attributed to CTL killing, k_{max} and k_{kill} are shape

parameters describing the saturable killing function, and j is the number of unique neoantigens

harbored by the lineage:

Equation 8: Empirically derived function for cancer cell lineage-specific death rates

11
$$k_{d,i} = f_{die} + (k_{max} + (1 - f_{die})) \frac{\emptyset * k_{kill}}{\sqrt{1 + (\emptyset * k_{kill})^2}}$$

12
$$\emptyset = \frac{\sum_{k=1}^{j} E_k MHC_k CTL_k}{NSCTL + \sum_{k=1}^{j} E_k MHC_k CTL_k}$$

14   Thus, the total CTLs reactive against j neoantigens is summed to calculate antigen-

15   specific killing. CTL_k is only counted if the kth neoantigen and its MHC are expressed (E_k and

16    $MHC_k = 1$).

Simulated tumor growth in mice

BBN963 orthotopic bladder cancer cells were previously established by chronic

exposure of C57BL/6 mice to 0.05% N-Butyl-N-(4-hydroxybutyl) nitrosamine [23]. Neoantigens

were predicted computationally and screened for bona fide immunogenicity in an ex vivo

ELISpot assay (Figure S2A). Additional details can be found in [20]. BBN963 cells were

implanted in immunocompetent C57BL/6 or immunodeficient NSG mice [26]. Tumor volumes

were measured weekly for five weeks once the tumors reached 200 mm^3. Tumor growth

volumes were digitized using WebPlotDigitizer [31] and used to calibrate mouse-specific model parameters in Table S1.

An independent dataset describing the growth dynamics of MC38 murine colon cancer cells in immunodeficient and immunocompetent mice were used to validate the mouse model. MC38 tumors were subcutaneously established in immunocompetent C57BL/6 or immunodeficient RAG1 KO mice [25]. Whole exome and RNA sequencing were used to identify 489 expressed neoantigens. In the absence of reported MHC allele specificity and ex vivo-validated immunogenicity, each neoantigen was assigned a random for MHC allele specificity and immunogenicity from the distribution reported for BBN963 cells [23]. Tumor growth in both immunodeficient and immunocompetent mice was successfully recapitulated by using the same parameters as BBN963 cells, except growth rate, which was increased 50% (Figure S2B-D).

Simulated tumor growth in humans

The TRACERx 100 cohort [24] contained 100 patients with varying stages of NSCLC: 26 stage I, 36 stage IB, 13 stage IIA, and 25 stage IIB/IIIA/IIIB. These patients underwent multiregion biopsy for extensive immunopeptidome profiling [10], [11]. Tumor diameters were estimated for each stage using conventional NSCLC staging criteria [32]. Clonal and subclonal neoantigen burden digitized from [10] with WebPlotDigitizer, as well as rates of neoantigen expression loss and MHC clonal/subclonal loss of heterozygosity reported in these studies, were used to calibrate simulated immunopeptidomes. In total, 100 tumors were simulated to target diameters mirroring the TRACERx 100 cohort. Clonal and subclonal neoantigen burden for each tumor was then calculated by determining the number of neoantigens present in 100% or < 100% of lineage-specific immunopeptidomes.

Human immunopeptidome cross-validation and local sensitivity analysis

Simulated immunopeptidomes were evaluated against an independent cohort of 12

patients [26] (Figure S3). Putative neoantigens from single-region biopsies collected from 12

patients prior to ICB treatment were assessed. Total neoantigens were calculated by filtering for

positive netCTL classification. To apply the TESLA criteria, we calculated agretopicity by taking

the ratio of somatic and wild-type MHC binding affinity. For predictions containing somatic

peptides of identical sequences but different predicted MHC binding alleles, the more strongly

binding prediction was used. Binding stabilities were calculated with NetMHCStabPan using

default parameters [33]. The number of neoantigens that passed three of five the TESLA criteria

(MHC binding affinity < 34 nM, MHC binding stability > 1.4 hours, and agretopicity > 0.1) was

multiplied, in the absence of reported foreignness, by a correction factor equal to the ratio of

neoantigens in the TRACERx 100 [10] cohort passing the MHC binding affinity, stability, and

agretopicity criteria that also did or did not pass the foreignness cutoff of $< 10^{-16}$. This reduced

the number of predicted neoantigens in the immunopeptidome by 80.6%. Neoantigen clonality

was not assessed due to potential sampling biases intrinsic to single-region biopsies. For each

of the seven discrete tumor stages, tumor diameters were estimated for each stage using

conventional NSCLC staging criteria [32]. A cohort of 40 tumors was simulated for each of the

seven diameters. The simulated and experimentally reported immunopeptidomes are shown in

Figure S3.

Local sensitivity analysis on the simulated TRACERx 100 cohort was performed by re-

simulating each tumor after varying one estimated parameter up or down 20%. This was

repeated for each estimated parameter in the model; the results are shown in Figure S4.

Exploratory simulations of ICB treatment and other tumor histologies

A cohort of 100 tumors was simulated until a target population roughly equivalent to a ~5

cm diameter tumor. ICB treatment of varying strength was implemented to increase the

sensitivity of CTLs to antigenic stimuli as well as the population-level carrying capacity, as below

(where *ICB* is a positive real number representing simulated treatment strength):

Equation 9: Implementation of ICB within the model

$$\frac{dCTL_k}{dt} = CTL_k \left(T_d(C-1) + \boldsymbol{ICB} \left(\frac{S_k T_a C}{CTL_k} \right) \right)$$

$$\frac{dC}{dt} = k_{in} - k_{in} f_{out} - k_{in}(1 - f_{out}) \left(\frac{NSCTL + \sum_{i=1}^{k} CTL_k}{\boldsymbol{ICB} * CTL_{ss}} \right)$$

To emulate renal cell carcinoma (RCC)-like tumors, a cohort of 100 tumors with 50% lower growth rate and 90% lower neoantigen gain, neoantigen loss, and MHC loss rates was simulated. To emulate microsatellite instability-high colorectal cancer (MSI)-like tumors, a cohort of 100 tumors with 100% higher growth rate and 200% higher clonal neoantigen burden was simulated. ICB treatment was simulated as above.

Statistical analyses

All simulations and statistical analyses were performed in MATLAB R2020b. Pearson's and Spearman's ρ were used as appropriate to assess linear correlation among continuous or ranked variables. Continuous variables in the local sensitivity were assessed for statistical significance at α = 0.05 using one-way ANOVA and a post-hoc pairwise Student's t-test, with Bonferroni correction for multiplicity. In cases of post-hoc multiple hypothesis testing, a Bonferroni correction was applied. Details of specific statistical analyses are provided in their corresponding figure captions.

RESULTS

Model recapitulates tumor growth in immunocompetent and immunodeficient mice

BBN963 orthotopic bladder cancer cells generated in [23] were implanted in

immunocompetent C57BL/6 and immunodeficient NSG mice [28] (Figure 3A). Thirty-four

neoantigens putatively expressed by the cells were identified and tested for immunogenicity ex

vivo [28] (see Methods). We used the results of these immunogenicity assays to scale in silico

immunogenicity and assign one of two murine MHC alleles for presentation (Figure 3B). Five-

week simulations of tumor growth in the absence of CTL expansion and CTL-mediated killing

accurately reproduced BBN963 growth dynamics in NSG mice (Figure 3C). Similarly, five-week

simulations with CTL expansion and cytotoxicity restored accurately reproduced BBN963

growth dynamics in C57BL/6 mice (Figure 3D).

To test model fidelity against an independent dataset, we used the growth dynamics of

MC38 murine colon cancer cells in immunodeficient and immunocompetent mice reported by

Efremova et al. [25]. Assuming the same distribution of neoantigen immunogenicity as BBN963

cells and changing no parameters except growth rate, the model accurately captured MC38

tumor growth in both immunocompetent and immunodeficient mice (Figure S2B-D). These

preliminary results suggested the generalizability of our predator-prey-like framework and

motivated further simulation of human tumor-immune dynamics.

Model recapitulates the tumor immunopeptidome of NSCLC patients

We substituted human parameter values for steady-state T cell abundance and lifespan

from the mouse model, while also re-estimating tumor carrying capacity and T cell sensitivity to

antigenic stimulus. No other parameters related to tumor or T cell dynamics were changed

(Table S1). This enabled successful simulation of the growth and immunopeptidome of 100

human NSCLC tumors (Figure 4A) [24]. Tumor growth was simulated until target population

sizes approximating the tumor diameters of TRACERx 100 NSCLC patients were reached. In

contrast to the murine simulations, cancer cells in the human simulations were allowed to

stochastically gain new neoantigens, lose expression of existing neoantigens, and lose MHC

expression (see Methods, Figure S2A) [10], [11]. Simulations recapitulated the distribution of

clonal, subclonal, and total neoantigen burden within this cohort (Figure 4B).

Interestingly, we observed substantial tumor and CTL diversification over time, as

depicted in Figures 4C-D. The number of unique tumor lineages was consistently higher than

the number of newly tumor-reactive CTL lineages due to lineage branching events caused by

neoantigen and MHC loss. In these cases, a new cancer lineage was formed without

introducing a new neoantigen and its cognate predator CTL. There was also a mild positive

correlation between the number of unique neoantigens and CTL diversity (Figure 4E).

Simulated ICB highlights pretreatment clonal neoantigen burden as a biomarkers of

response

Prolonged antigen exposure results in CTL exhaustion, diminishing their proliferative and

cytolytic capabilities. This process is accelerated by checkpoint molecules, many of which are

overexpressed in the tumor microenvironment. ICB blocks these checkpoint molecules,

reinvigorating the immune response and promoting the expansion of a select subset of CTL

lineages (Figure 5A) [34]. To simulate ICB of late-stage NSCLC, we first constructed a 100-

tumor virtual cohort comprising tumors of at least ~5 cm in diameter. This roughly corresponds

to the advanced stage of NSCLC for which ICB is indicated. Similar to the TRACERx 100 virtual

cohort, we observed a mild positive correlation between neoantigen burden and CTL diversity

prior to treatment (Figure 5B).

Various strengths of ICB treatment were explored within the model by simulating tumor

growth for one year in the presence of 2-, 6-, and 10-fold increases in CTL sensitivity to antigen

stimulus with concurrent 2-, 6-, and 10-fold increases in steady-state CTL carrying capacity. As

expected, the proportion of patients with maximal tumor shrinkage of at least 30% –

"responders" – increased with treatment strength [35]. Notably, response rates in the 10-fold

treatment simulation approached levels observed for ICB in Phase 3 clinical trials (Figure 5C)

[36].

Pretreatment neoantigen burden correlated positively with deeper responses, in line with

previous reports. (Figure 5D) [5]. Interestingly, this was driven primarily by clonal neoantigen

burden. No significant relationship was found between response depth and the number of

unique subclonal neoantigens. We also examined whether the immunogenicity or clonality of

the most immunogenic neoantigen (i.e., rank 1) could predict response. In this case, clonality

outperformed immunogenicity in predictive sensitivity and specificity; however, both were

outperformed by the product of the two, termed clonality-weighted immunogenicity (Figure 5E).

Responding tumors also had consistently lower Shannon entropy in their immunopeptidomes,

corresponding to more homogeneous sets of neoantigens (Figure 5F). These data support the

importance of neoantigen clonality and highlight intratumoral heterogeneity as a key detractor

from ICB response [5], [37].

CTL dynamics correlate with ICB response and immunoediting

While responders tended to have greater pretreatment CTL diversity, further CTL

diversification induced by ICB treatment strongly correlated with shallower responses (Figure

6A-B). Instead, expansion and maintenance of CTL lineages that were highly abundant before

ICB initiation tended to elicit more profound responses (Figure 6C) [21], [38], [39]. Responders

also had more stable pools of nonspecific stimulus during treatment (Figure 6C). These findings

support the notion that prognoses are most favorable when ICB induces focused, oligoclonal

responses against highly clonal neoantigens, at least for certain tumor types [21], [40].

Finally, we evaluated the potential for immunoediting against the rank 1 neoantigen

during treatment. Immunoediting is the phenomenon by which cell lineages bearing certain

neoantigens are selected against and killed by CTLs, resulting in diminishment or even

eradication of certain neoantigens from the immunopeptidome [6]. Across the simulated tumors,

we observed no correlation between the immunogenicity nor clonality-weighted immunogenicity

of the rank 1 neoantigen and on-treatment immunoediting. There was, however, a very modest

negative correlation between the immunoediting and the clonality of the rank 1 neoantigen alone

(Figure 6D-F). This suggests that clonality may supersede immunogenicity in provoking lineage-

specific selection from CTLs; however, given the very modest effect and borderline p-value,

further work in larger patient populations – simulated or otherwise – is needed.

Model enables exploration of diverse tumor immunopeptidomes

To evaluate model generalizability, RCC-like and MSI-like immunopeptidomes with

adjusted tumor growth and mutation rates were simulated and subjected to ICB (Figure 7A-B)

[37], [41]. As expected, lower tumor growth and mutation rates led to lower neoantigen burden

in RCC-like tumors (Figure 7B). This resulted in poorer predicted responses in these tumors

when compared to NSCLC-like and MSI-like tumors, in alignment with clinical experience

(Figure 7C) [36], [42], [43]. Even in these neoantigen-poor tumors, however, clonal neoantigen

burden correlated with response depth (Figure 7D).

This was also the case for neoantigen-rich MSI-like tumors. However, despite their

higher neoantigen burden, the most heterogeneous MSI-like tumors were less likely to respond

than most homogenous tumors (Figure 7E). Furthermore, the CTL lineages that were most

abundant prior to treatment in MSI-like tumors expanded less profoundly upon ICB treatment

(Figure 7F). This is in sharp contrast to NSCLC-like tumors, in which oligoclonal CTL expansion

was still observed even in nonresponders. The nonlinear relationship between CTL expansion

and neoantigen burden warrants further investigation.

DISCUSSION

Characterizations of the quantitative relationship between the immunopeptidome and CTL clonal dynamics remain lacking. This work provides a mathematical, predator-prey-like framework capable of modeling the co-evolution of the tumor immunopeptidome and TCR repertoire. Built initially upon mouse data, this framework was straightforward to adapt for human tumor simulations. This allowed us to interrogate several clinically observed phenomena, including both pre- and on-treatment changes to the immunopeptidome and TCR repertoire. While several noteworthy computational models of neoantigen evolution have been published these models consider either a fixed cancer cell population size or calculate death rates as a function of the cumulative immunogenicity of neoantigens expressed by a lineage or clone [2], [15], [44], [45]. No work, to our knowledge, has modeled a growing tumor with dynamic immunopeptidomes and CTL populations. Our results support the directive that neoantigen discovery methods should focus not only on immunogenicity, but also on clonality [46].

One key insight from this work is the shift away from neoantigen "dilution" as the predominant sequelae of intratumoral heterogeneity and toward CTL "distraction". Several studies report the ability of subclonally expressed neoantigens to dilute an otherwise strongly immunogenic neoantigen; that is, to shield it from immunoediting and dampen anti-tumor killing [8], [14]. Our model supports the importance of neoantigen dilution, showing that neoantigen clonality may be more important than immunogenicity in predicting immunoediting and conceiving biomarkers for responders to ICB. Considering clonality and immunogenicity in tandem is especially informative to ICB response (Figure 6E). However, while these studies and mechanisms are seminally informative to this work, they largely ignore the CTL compartment. Holistically characterizing the anti-immunopeptidome response requires concomitantly assessing how the TCR repertoire evolves across space, time, and tumor types.

Here, we propose the competition for nonspecific stimulus by different CTL lineages as a mechanism for ICB resistance and provide a mathematical modeling framework for its investigation. Our simulations support interclonal CTL competition as a potential reason why

highly heterogeneous, subclonal tumors are associated with poor outcomes. We show that CTL

diversification during treatment correlates with shallower responses, and that responders who

do not diversify have more replete pools of nonspecific stimulus. In addition, many responding

MSI-like tumors that responded to ICB have noticeably lower tumor-specific CTL expansion

despite greater clonal neoantigen burden. Given these findings, "neoantigen dilution" may be a

slight mischaracterization that diverts attention from the fact that CTLs responding to

neoantigens are the primary mediators of tumor debulking.

Limitations of this work include its spatially ignorant modeling of metastatic disease as

one lumped tumor. Neoantigens can differ between patients and among metastases, potentially

leading to increased systemic heterogeneity but decreased local heterogeneity [47]. Moreover,

whether the immunopeptidome of individual metastases influences the already spatially biased

trafficking of CTLs remains an open question. Are neoantigen-reactive CTLs more likely to

linger in tumors where their target neoantigen is present? What if the neoantigen is highly

subclonal – what is the probability they will physically encounter it? Is there a critical clonality

threshold? These are critical questions that the model is not currently equipped to address.

Our work is primarily informed by data from two mouse models and two NSCLC studies.

While the role of neoantigen clonality in NSCLC is generally consistent with its role in melanoma

and colorectal cancer, it could differ in other tumor types such as low-grade glioblastoma and

prostate cancer [37]. Moreover, the simulated response rate in MSI-like tumors was above that

observed in the clinic, suggesting variability in treatment or CTL potency across tumor types.

Ongoing clinical studies profiling the effect of ICB on TCR repertoire diversity in CRC may shed

additional light on the veracity of our observations [48]. More generally, studies considering the

coevolution of the cancer immunopeptidome and TCR repertoire diversity (or lack thereof) will

be critical to achieving long-term disease control and cure.

In summary, this predator-prey-like framework accurately described clinically observed

immunopeptidome diversity and recapitulated clinical evidence supporting the correlation

between clonal neoantigen burden and ICB response. By including a dynamic CTL

compartment, the model could also describe the effects of TCR diversity on predicted

responses. Analysis of the tumor-immune axis also analyses also suggested that neoantigen

immunoediting may correlate more strongly with pretreatment clonality than immunogenicity.

This model can be readily adapted to future data published in other tumor types through simple,

biologically motivated adjustments to parameter values. Further refinement could facilitate low-

cost theoretical explorations of histology-specific therapeutic strategies in advance of

developing and deploying them in the clinic. This work supports the further development of

predator-prey-like models to elucidate nonlinear tumor-immune dynamics for clinical translation

and utility.

DECLARATIONS

Patient consent for publication

Not applicable.

Ethics approval

Not applicable.

Availability of data and material

All original code has been deposited at Zenodo (10.5281/zenodo.6596442) and is

publicly available. All data collected for simulation benchmarking have been made available in

the Supplement.

Competing interests

T.Q. is a contractor for Hatteras Venture Partners. B.V. is a shareholder of and consultant for GeneCentric. Y.C. is a consultant for Janssen Research & Development.

Funding

This work was supported by the National Institute of Health (R35 GM119661).

Authors' contributions

Conceptualization, T.Q. and Y.C.; Methodology, T.Q.; Software, T.Q.; Validation, T.Q.; Formal Analysis, T.Q.; Investigation, T.Q.; Resources, Y.C.; Data Curation, T.Q.; Writing – Original Draft, T.Q.; Writing – Review & Editing – T.Q., Y.C., and B.V.; Visualization – T.Q.; Supervision – Y.C.; Project Administration – T.Q. and Y.C.; Funding Acquisition – Y.C.

Acknowledgements

The authors thank Amber Cipriani and J. Justin Milner for their helpful input during study design and manuscript preparation.

SUPPLEMENTAL MATERIAL

Table S1: Model parameters and sources

Figure S1: Modeled neoantigen parameters and stochastic events

Figure S2: Overview of BBN963 cell line characterization and MC38 growth simulations

Figure S3: Validation of simulated neoantigen burden in an independent cohort

Figure S4: Local sensitivity analysis of model parameters

REFERENCES

[1] P. T. Hamilton, B. R. Anholt, and B. H. Nelson, "Tumour immunotherapy: lessons from predator–prey theory," *Nat. Rev. Immunol.*, May 2022, doi: 10.1038/s41577-022-00719-y.

[2] E. Lakatos *et al.*, "Evolutionary dynamics of neoantigens in growing tumors," *Nat. Genet.*, vol. 52, no. 10, pp. 1057–1066, Oct. 2020, doi: 10.1038/s41588-020-0687-1.

[3] C. C. Smith, S. R. Selitsky, S. Chai, P. M. Armistead, B. G. Vincent, and J. S. Serody, "Alternative tumour-specific antigens," *Nat. Rev. Cancer*, vol. 19, no. 8, pp. 465–478, Aug. 2019, doi: 10.1038/s41568-019-0162-4.

[4] D. K. Wells *et al.*, "Key Parameters of Tumor Epitope Immunogenicity Revealed Through a Consortium Approach Improve Neoantigen Prediction," *Cell*, vol. 183, no. 3, pp. 818-834.e13, Oct. 2020, doi: 10.1016/j.cell.2020.09.015.

[5] N. McGranahan *et al.*, "Clonal neoantigens elicit T cell immunoreactivity and sensitivity to immune checkpoint blockade," *Science*, vol. 351, no. 6280, pp. 1463–1469, 2016.

[6] V. Anagnostou *et al.*, "Evolution of Neoantigen Landscape during Immune Checkpoint Blockade in Non–Small Cell Lung Cancer," *Cancer Discov.*, vol. 7, no. 3, pp. 264–276, Mar. 2017, doi: 10.1158/2159-8290.CD-16-0828.

[7] K. Litchfield *et al.*, "Meta-analysis of tumor- and T cell-intrinsic mechanisms of sensitization to checkpoint inhibition," *Cell*, vol. 184, no. 3, pp. 596-614.e14, Feb. 2021, doi: 10.1016/j.cell.2021.01.002.

[8] R. S. Gejman *et al.*, "Rejection of immunogenic tumor clones is limited by clonal fraction," *eLife*, vol. 7, p. e41090, Nov. 2018, doi: 10.7554/eLife.41090.

[9] M. L. Burger *et al.*, "Antigen dominance hierarchies shape TCF1+ progenitor CD8 T cell phenotypes in tumors," *Cell*, vol. 184, no. 19, pp. 4996-5014.e26, Sep. 2021, doi: 10.1016/j.cell.2021.08.020.

[10] R. Rosenthal *et al.*, "Neoantigen-directed immune escape in lung cancer evolution," *Nature*, vol. 567, no. 7749, pp. 479–485, Mar. 2019, doi: 10.1038/s41586-019-1032-7.

[11] N. McGranahan *et al.*, "Allele-Specific HLA Loss and Immune Escape in Lung Cancer Evolution," *Cell*, vol. 171, no. 6, pp. 1259-1271.e11, Nov. 2017, doi: 10.1016/j.cell.2017.10.001.

[12] C. C. Smith *et al.*, "Machine-Learning Prediction of Tumor Antigen Immunogenicity in the Selection of Therapeutic Epitopes," *Cancer Immunol. Res.*, vol. 7, no. 10, pp. 1591–1604, Oct. 2019, doi: 10.1158/2326-6066.CIR-19-0155.

[13] D. Miao *et al.*, "Genomic correlates of response to immune checkpoint blockade in microsatellite-stable solid tumors," *Nat. Genet.*, vol. 50, no. 9, pp. 1271–1281, Sep. 2018, doi: 10.1038/s41588-018-0200-2.

[14] Y. Wolf *et al.*, "UVB-Induced Tumor Heterogeneity Diminishes Immune Response in Melanoma," *Cell*, vol. 179, no. 1, pp. 219-235.e21, Sep. 2019, doi: 10.1016/j.cell.2019.08.032.

[15] M. Łuksza *et al.*, "Neoantigen quality predicts immunoediting in survivors of pancreatic cancer," *Nature*, May 2022, doi: 10.1038/s41586-022-04735-9.

[16] C. Leitão, A. A. Freitas, and S. Garcia, "The role of TCR specificity and clonal competition during reconstruction of the peripheral T cell pool," *J. Immunol. Baltim. Md 1950*, vol. 182, no. 9, pp. 5232–5239, May 2009, doi: 10.4049/jimmunol.0804071.

[17] D. Nandi *et al.*, "T cell costimulation, checkpoint inhibitors and anti-tumor therapy," *J. Biosci.*, vol. 45, p. 50, 2020.

[18] H. Schreiber, T. H. Wu, J. Nachman, and W. M. Kast, "Immunodominance and tumor escape," *Semin. Cancer Biol.*, vol. 12, no. 1, pp. 25–31, Feb. 2002, doi: 10.1006/scbi.2001.0401.

[19] J. L. Urban, C. van Waes, and H. Schreiber, "Pecking order among tumor-specific antigens," *Eur. J. Immunol.*, vol. 14, no. 2, pp. 181–187, 1984, doi: 10.1002/eji.1830140214.

[20] J. W. Yewdell, "Confronting complexity: real-world immunodominance in antiviral CD8+ T cell responses," *Immunity*, vol. 25, no. 4, pp. 533–543, Oct. 2006, doi: 10.1016/j.immuni.2006.09.005.

[21] J. Kidman *et al.*, "Characteristics of TCR Repertoire Associated With Successful Immune Checkpoint Therapy Responses," *Front. Immunol.*, vol. 11, p. 587014, 2020, doi: 10.3389/fimmu.2020.587014.

[22] C. Bailey *et al.*, "Tracking Cancer Evolution through the Disease Course," *Cancer Discov.*, vol. 11, no. 4, pp. 916–932, Apr. 2021, doi: 10.1158/2159-8290.CD-20-1559.

[23] R. Saito *et al.*, "Molecular Subtype-Specific Immunocompetent Models of High-Grade Urothelial Carcinoma Reveal Differential Neoantigen Expression and Response to Immunotherapy," *Cancer Res.*, vol. 78, no. 14, pp. 3954–3968, Jul. 2018, doi: 10.1158/0008-5472.CAN-18-0173.

[24] M. Jamal-Hanjani *et al.*, "Tracking the Evolution of Non-Small-Cell Lung Cancer," *N. Engl. J. Med.*, vol. 376, no. 22, pp. 2109–2121, Jun. 2017, doi: 10.1056/NEJMoa1616288.

[25] M. Efremova *et al.*, "Targeting immune checkpoints potentiates immunoediting and changes the dynamics of tumor evolution," *Nat. Commun.*, vol. 9, no. 1, p. 32, Jan. 2018, doi: 10.1038/s41467-017-02424-0.

[26] P. M. Forde *et al.*, "Neoadjuvant PD-1 Blockade in Resectable Lung Cancer," *N. Engl. J. Med.*, vol. 378, no. 21, pp. 1976–1986, May 2018, doi: 10.1056/NEJMoa1716078.

[27] H. Lodish *et al.*, *Molecular cell biology*, Ninth edition. New York: Macmillan Learning, 2021.

[28] A. S. Truong *et al.*, "Entinostat induces antitumor immune responses through immune editing of tumor neoantigens," *J. Clin. Invest.*, vol. 131, no. 16, p. 138560, Aug. 2021, doi: 10.1172/JCI138560.

[29] C. H. Lee, M. Salio, G. Napolitani, G. Ogg, A. Simmons, and H. Koohy, "Predicting Cross-Reactivity and Antigen Specificity of T Cell Receptors," *Front. Immunol.*, vol. 11, p. 565096, 2020, doi: 10.3389/fimmu.2020.565096.

[30] G. Lythe, R. E. Callard, R. L. Hoare, and C. Molina-París, "How many TCR clonotypes does a body maintain?," *J. Theor. Biol.*, vol. 389, pp. 214–224, Jan. 2016, doi: 10.1016/j.jtbi.2015.10.016.

[31] A. Rohatgi, *WebPlotDigitizer*. 2022. [Online]. Available: https://automeris.io/WebPlotDigitizer

[32] M. B. Amin, American Joint Committee on Cancer, and American Cancer Society, Eds., *AJCC cancer staging manual*, Eight edition / editor-In-Chief, Mahul B. Amin, MD, FCAP ; Editors, Stephen B. Edge, MD, FACS [and 16 others] ; Donna M. Gress, RHIT, CTR-Technical editor ; Laura R. Meyer, CAPM-Managing editor. Chicago IL: American Joint Committee on Cancer, Springer, 2017.

[33] M. Rasmussen *et al.*, "Pan-Specific Prediction of Peptide–MHC Class I Complex Stability, a Correlate of T Cell Immunogenicity," *J. Immunol.*, vol. 197, no. 4, pp. 1517–1524, Aug. 2016, doi: 10.4049/jimmunol.1600582.

[34] J. Zhang *et al.*, "Compartmental Analysis of T-cell Clonal Dynamics as a Function of Pathologic Response to Neoadjuvant PD-1 Blockade in Resectable Non–Small Cell Lung Cancer," *Clin. Cancer Res.*, vol. 26, no. 6, pp. 1327–1337, Mar. 2020, doi: 10.1158/1078-0432.CCR-19-2931.

[35] E. A. Eisenhauer *et al.*, "New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1)," *Eur. J. Cancer Oxf. Engl. 1990*, vol. 45, no. 2, pp. 228–247, Jan. 2009, doi: 10.1016/j.ejca.2008.10.026.

[36] M. Reck *et al.*, "Pembrolizumab versus Chemotherapy for PD-L1–Positive Non–Small-Cell Lung Cancer," *N. Engl. J. Med.*, vol. 375, no. 19, pp. 1823–1833, Nov. 2016, doi: 10.1056/NEJMoa1606774.

[37] Y. Wolf and Y. Samuels, "Intratumor Heterogeneity and Antitumor Immunity Shape One Another Bidirectionally," *Clin. Cancer Res.*, pp. OF1–OF8, Apr. 2022, doi: 10.1158/1078-0432.CCR-21-1355.

[38] E. Cha *et al.*, "Improved survival with T cell clonotype stability after anti-CTLA-4 treatment in cancer patients," *Sci. Transl. Med.*, vol. 6, no. 238, p. 238ra70, May 2014, doi: 10.1126/scitranslmed.3008211.

[39] L. Au *et al.*, "Determinants of anti-PD-1 response and resistance in clear cell renal cell carcinoma," *Cancer Cell*, vol. 39, no. 11, pp. 1497-1518.e11, Nov. 2021, doi: 10.1016/j.ccell.2021.10.001.

[40] M. Casarrubios *et al.*, "Pretreatment Tissue TCR Repertoire Evenness Is Associated with Complete Pathologic Response in Patients with NSCLC Receiving Neoadjuvant Chemoimmunotherapy," *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.*, vol. 27, no. 21, pp. 5878–5890, Nov. 2021, doi: 10.1158/1078-0432.CCR-21-1200.

[41] F. Raynaud, M. Mina, D. Tavernari, and G. Ciriello, "Pan-cancer inference of intra-tumor heterogeneity reveals associations with different forms of genomic instability," *PLoS Genet.*, vol. 14, no. 9, p. e1007669, Sep. 2018, doi: 10.1371/journal.pgen.1007669.

[42] D. F. McDermott *et al.*, "Open-Label, Single-Arm, Phase II Study of Pembrolizumab Monotherapy as First-Line Therapy in Patients With Advanced Non-Clear Cell Renal Cell Carcinoma," *J. Clin. Oncol. Off. J. Am. Soc. Clin. Oncol.*, vol. 39, no. 9, pp. 1029–1039, Mar. 2021, doi: 10.1200/JCO.20.02365.

[43] T. André *et al.*, "Pembrolizumab in Microsatellite-Instability-High Advanced Colorectal Cancer," *N. Engl. J. Med.*, vol. 383, no. 23, pp. 2207–2218, Dec. 2020, doi: 10.1056/NEJMoa2017699.

[44] G. Aguadé-Gorgorió and R. Solé, "Tumour neoantigen heterogeneity thresholds provide a time window for combination immunotherapy," *J. R. Soc. Interface*, vol. 17, no. 171, p. 20200736, Oct. 2020, doi: 10.1098/rsif.2020.0736.

[45] M. Łuksza *et al.*, "A neoantigen fitness model predicts tumour response to checkpoint blockade immunotherapy," *Nature*, vol. 551, no. 7681, pp. 517–520, Nov. 2017, doi: 10.1038/nature24473.

[46] N. McGranahan and C. Swanton, "Neoantigen quality, not quantity," *Sci. Transl. Med.*, vol. 11, no. 506, p. eaax7918, Aug. 2019, doi: 10.1126/scitranslmed.aax7918.

[47] T. Jiang *et al.*, "Heterogeneity of neoantigen landscape between primary lesions and their matched metastases in lung cancer," *Transl. Lung Cancer Res.*, vol. 9, no. 2, pp. 246–256, Apr. 2020, doi: 10.21037/tlcr.2020.03.03.

[48] N. Huyghe, E. Benidovskaya, P. Stevens, and M. Van den Eynde, "Biomarkers of Response and Resistance to Immunotherapy in Microsatellite Stable Colorectal Cancer: Toward a New Personalized Medicine," *Cancers*, vol. 14, no. 9, p. 2241, Apr. 2022, doi: 10.3390/cancers14092241.

1   **MAIN FIGURES**

2   **Figure 1.** Intratumoral predator-prey dynamics depend on the immunopeptidome

3   A. Schematic of neoantigen presentation. Neoantigens arriving from tumor-specific

4   aberrations in DNA or RNA are presented on MHC molecules. CTLs with TCRs that

5   recognize the neoantigen subsequently proliferate and attack the neoantigen-presenting

6   cell.

7   B. Schematic of immunodominance. Highly immunogenic neoantigens that are also

8   subclonal can be shielded from CTLs due to the abundance of more clonal neoantigens.

9   C. Schematic of interclonal competition. CTLs reactive against different neoantigens

10   compete for the same pool of shared nonspecific stimulus, including cytokines and co-

11   stimulation from antigen-presenting cells.

A

- ○ Self antigen
- ● Neoantigen
- MHC (different alleles)
- Neoantigen-reactive TCR

Cancer cell

Mutation

T cell

B

Highly immunogenic

Weakly immunogenic

Self antigen

C

Nonspecific stimulus
(e.g., cytokines)

1   **Figure 2.** Distinct tumor-reactive CTL lineages were modeled to capture immunopeptidome

2   interactions

3   A. Schematic of the TCR-specific CTL lineage model. Cancer cells present neoantigens,

4   providing pro-proliferative stimulus to neoantigen-specific populations of CTLs. A fraction

5   of cancer cells dies to CTL-mediated killing, while the remaining fraction dies to

6   unrelated causes. In parallel, CTL expansion depends on sufficient nonspecific stimulus,

7   which is consumed by all tumor-reactive CTL lineages as well as a pooled lineage

8   representing all tumor-ignorant CTLs. A fraction of nonspecific stimulus turns over

9   irrespective of CTL consumption.

Figure 3. Murine tumor growth dynamics were recapitulated in divergent immune contexts

A. NSG and C57BL/6 mice were inoculated with subcutaneous BBN963 tumors. Tumor

volumes were measured over five weeks [28].

B. Saito et al. identified putative neoantigens expressed by BBN963 and the MHC alleles

predicted to present them (H2-Db, H2-Kb). Baseline-corrected ELISpot scores were

used to scale neoantigen immunogenicity to the range [0,1].

C. Simulated tumor volumes (left, black) normalized to baseline volume (dashed gray line)

compared against growth dynamics in NSG mice reported by Truong et al. (gray circles

and error bars). Nonspecific stimulus (center) and CTL populations (right) are shown.

Dashed lines, 1.

D. Simulated tumor volumes (left, black) normalized to baseline volume (dashed gray line)

compared against growth dynamics in C57BL/6 mice reported by Truong et al. (gray

circles and error bars). Nonspecific stimulus (center) and CTL populations (right) are

shown. Black dashed lines, 1; Red dashed lines, approximate total abundance of 3 Ag-

specific T cell lineages observed in mice [28]

Figure 4. Human NSCLC immunopeptidomes were recapitulated from clinical data

A. Schematic of multiregion biopsies collected in the TRACERx 100 cohort. 100 NSCLC tumors of various stages were sampled, corresponding to target diameters of 1.5 cm (stage IA), 3.5 cm (stage IB), 4.5 cm (stage IIA), and 5 cm (stage IIB/III).

B. Simulated neoantigen burden compared against clonal and subclonal neoantigen burden digitized from [10].

C. Emergence of unique tumor lineages over the course of simulated tumor growth.

D. Emergence of new tumor-reactive CTL lineages over the course of simulated tumor growth.

E. Pearson's ρ correlation between unique neoantigen number and tumor-reactive CTL Shannon entropy in the simulated TRACERx 100 cohort. Excludes one tumor with > 600 neoantigens for visualization purposes.

1   **Figure 5.** Pretreatment tumor-immune status correlated with response to immune checkpoint

2   blockade

3   A. Schematic of the proposed mechanism of action of PD-1 blockade, a common ICB.

4   B. Pearson's ρ correlation between unique neoantigen number and tumor-reactive CTL

5   Shannon entropy in the simulated 100-tumor cohort prior to treatment. Excludes one

6   tumor with > 600 neoantigens for visualization purposes.

7   C. Maximum tumor shrinkage under increasing levels of treatment strength: 2- (left), 6-

8   (center), and 10- (center) fold increases in sensitivity to antigen stimulus and systemic

9   carrying capacity were simulated. Each bar corresponds to a tumor. The dashed line at

10   corresponds to the 30% or greater tumor shrinkage required to classify as partial

11   response per RECIST 1.1 criteria [35].

12   D. Pearson's ρ correlation between response and pretreatment total, clonal, and subclonal

13   neoantigen burden. Excludes one tumor with > 600 neoantigens for visualization

14   purposes.

15   E. Response status (colored bars) relative to rank 1 neoantigen immunogenicity (top),

16   clonality (middle), and clonality-weighted immunogenicity (bottom). Sensitivity and

17   specificity were calculated from the correctly classified fraction of responders (N=36) and

18   non-responders (N=64), respectively.

19   F. Fraction of responses among the X% most and least entropic immunopeptidomes, X

20   being indicated by the X axis value. Entropy refers to Shannon entropy. Dashed lines,

21   means.

Figure 6. Immunopeptidome and CTL dynamics correlated with ICB response

A. Pearson's ρ correlation between response and pretreatment tumor-reactive CTL Shannon entropy.

B. Pearson's ρ correlation between response and new tumor-reactive CTL lineages that emerged upon simulated ICB.

C. Summed abundance of the top 5 most abundant TCR lineages prior to ICB (top) and concurrent pool of nonspecific stimulus (bottom). Each line represents a tumor. Abundances are normalized to the pretreatment baseline average of all 100 simulated tumors in the cohort.

D. Spearman's ρ rank correlation between pre-treatment immunogenicity of the rank 1 most immunogenic neoantigen prior to treatment and the extent of immunoediting after one year of ICB treatment. Immunoediting was determined by calculating the fold change of the number of cells expressing the rank 1 neoantigen between the first and last day of ICB.

E. Spearman's ρ rank correlation between pre-treatment clonality of the rank 1 most immunogenic neoantigen prior to treatment and the extent of immunoediting after one year of ICB treatment.

F. Spearman's ρ rank correlation between pre-treatment clonality-weighted immunogenicity of the rank 1 most immunogenic neoantigen prior to treatment and the extent of immunoediting after one year of ICB treatment.

Figure 7. Model describes diverse tumor immunopeptidomes and their response to treatment

A. Emergence of unique tumor lineages over the course of simulated tumor growth in RCC-

like (left), NSCLC-like (center), and MSI-like (right) tumors.

B. Simulated clonal and subclonal neoantigen burden of tumors simulated in (A). Each bar

corresponds to a tumor

C. Maximum shrinkage of tumors simulated in (A) under treatment constituting 10- fold

increases in sensitivity to antigen stimulus and systemic carrying capacity; each bar

corresponds to a tumor. The dashed line at corresponds to the 30% or greater tumor

shrinkage required to classify as partial response per RECIST 1.1 criteria [35].

D. Pearson's ρ correlation between response and pretreatment clonal neoantigen burden of

tumors simulated in (A).

E. Fraction of responses among the X% most and least entropic immunopeptidomes of

tumors simulated in (A), with X being indicated by the X axis value. Entropy refers to

Shannon entropy. Dashed lines, means.

F. Summed abundance of the top 5 most abundant TCR lineages prior to ICB (top row) and

concurrent pool of nonspecific stimulus (bottom row) of the tumors simulated in (A).

Each line represents a tumor. Abundances are normalized to the pretreatment baseline

average of all 100 simulated tumors in the cohort.

SUPPLEMENTAL TABLES AND FIGURES

Table S1. Model parameters

Figure S1. Model-simulated neoantigens account for stochastic events and TESLA criteria

A. Each neoantigen was assigned values for the TESLA parameters drawn from the

distributions for NSCLC tumors by Wells et al. in [4]. TPM, transcripts per million.

Distributions (top) and values (bottom) of sampled and experimental parameters from

146 NSCLC-derived peptides are shown.

B. Schematic of stochastic neoantigen events. A founder cell is established with a random

number of neoantigens and assigned a neoantigen gain rate from a lognormal

distribution. Each neoantigen was also assigned to be presented by one of three MHC

alleles. Neoantigen and MHC expression could also be lost by a cell during tumor

growth.

A

Figure S2. Characterization of the BBN963 immunopeptidome and validation in MC38 cells

A.   BBN963 cells were established by chronic exposure of C57BL/6 mice to 0.05% N-Butyl-

N-(4-hydroxybutyl) nitrosamine. Neoantigens were called and screened for bona fide

immunogenicity in an ex vivo ELISpot assay. Additional details can be found in [23].

B. Growth dynamics of MC38 were simulated using the same parameters as BBN963 cells,

except growth rate, which was increased 50% [25].

C. Simulated tumor volumes (left, black) normalized to baseline volume (dashed gray line)

compared against growth dynamics in RAG1 KO mice reported by Efremova et al. (gray

circles and error bars). Nonspecific stimulus (center) and CTL populations (right) are

shown. Dashed lines, 1.

D. Simulated tumor volumes (left, black) normalized to baseline volume (dashed gray line)

compared against growth dynamics in C57BL/6 mice reported by Efremova et al. (gray

circles and error bars). Nonspecific stimulus (center) and CTL populations (right) are

shown. Dashed lines, 1.

Figure S3. Simulated neoantigen burdens were within range of an independent cohort

A. Neoantigen predictions reported by Forde et al. 2018 in 12 patients with sufficient

pretreatment tissue to characterize candidate neoantigens were assessed. Total

neoantigens were calculated by filtering positive netCTL classification. To apply Wells'

criteria, we calculated agretopicity by taking the ratio of somatic and wild-type MHC

binding affinity. For predictions containing somatic peptides of identical sequences but

different HLA alleles, the stronger predicted binding was used. Binding stabilities were

calculated with NetMHCStabPan using default parameters. As biopsies were single-

region, neoantigen clonality was not assessed. Tumor diameters were estimated for

each tumor using conventional NSCLC staging criteria. For each of seven discrete

estimated tumor diameters, a cohort of 40 tumors was simulated.

Figure S4. Local sensitivity analysis of parameters indicates model robustness

A. Mean number of unique relative to baseline (N=100 simulations per condition). No significant fold differences in neoantigen number were found from pairwise Student's t-tests with neither nominal nor Bonferroni-corrected significance thresholds.

B. Mean TCR entropy relative to baseline (N=100 simulations per condition). No significant fold differences in Shannon entropy were found from pairwise Student's t-tests with neither nominal nor Bonferroni-corrected significance thresholds.

A

B

2