RESOURCE ARTICLE

Single cell analysis of lymphatic endothelial cell fate specification and differentiation during zebrafish development

Lin Grimm[1,2,3,*], Elizabeth Mason[1,2*,#], Oliver Yu[1,2], Stefanie Dudczig[1,2], Virginia Panara[4], Tyrone Chen[1,2], Neil I. Bower[3], Scott Paterson[1,2,3], Kazuhide Okuda[1,2], Maria Rondon Galeano[1,2], Sakurako Kobayashi[1,2], Anne Senabouth[3,5], Anne K. Lagendijk[3], Joseph Powell[3,5,6,7], Kelly A. Smith[8], Katarzyna Koltowska[4] and Benjamin M. Hogan[1,2,3,8 #].

[1] Organogenesis and Cancer Program, Peter MacCallum Cancer Centre, Melbourne, VIC 3000, Australia.
[2] Sir Peter MacCallum Department of Oncology, University of Melbourne, Melbourne, VIC 3000, Australia.
[3] Division of Genomics of Development and Disease, Institute for Molecular Bioscience, The University of Queensland, St Lucia, Brisbane, Queensland 4072, Australia.
[4] Department of Immunology, Genetics and Pathology, Uppsala University, Uppsala, Sweden.
[5] Garvan Institute of Medical Research, 384 Victoria Street, Darlinghurst, Sydney, 2010, Australia.
[6] School of Medical Sciences, 18 High Street, University of New South Wales, Kensington, Sydney, Australia.
[7] Garvan-Weizmann Centre for Cellular Genomics, Garvan Institute of Medical Research, 384 Victoria Street, Darlinghurst, Sydney, 2010, Australia.
[8] Department of Anatomy and Physiology, University of Melbourne, Melbourne, VIC 3000, Australia.

[#] *Authors for Correspondence:*

Professor Ben Hogan
Organogenesis and Cancer Program,
Peter MacCallum Cancer Centre,
Melbourne, VIC 3000, Australia
E-mail: ben.hogan@petermac.org

Dr Elizabeth Mason (computational biology lead)
Organogenesis and Cancer Program,
Peter MacCallum Cancer Centre,
Melbourne, VIC 3000, Australia
E-mail: elizabeth.mason@petermac.org

Keywords: Lymphatics, Lymphangiogenesis, Prox1, Single cell, RNA Seq, ATAC seq.
Summary: 181 words
Main Text: 4,700 words

During development, the lymphatic vasculature forms as a second, new vascular network
derived from blood vessels. The transdifferentiation of embryonic venous endothelial cells
(VECs) into lymphatic endothelial cells (LECs) is the first step in this process. Specification,
differentiation and maintenance of LEC fate are all driven by the transcription factor Prox1,
yet downstream mechanisms remain to be elucidated. We present a single cell
transcriptomic atlas of lymphangiogenesis in zebrafish revealing new markers and
hallmarks of LEC differentiation over four developmental stages. We further profile single
cell transcriptomic and chromatin accessibility changes in zygotic *prox1a* mutants that are
undergoing a VEC-LEC fate reversion during differentiation. Using maternal and zygotic
***prox1a/prox1b* mutants, we determine the earliest transcriptomic changes directed by**
Prox1 during LEC specification. This work altogether reveals new transcriptional targets and
regulatory regions of the genome downstream of Prox1 in LEC maintenance, as well as
showing that Prox1 specifies LEC fate primarily by limiting blood vascular and
hematopoietic fate. This extensive single cell resource provides new mechanistic insights
into the enigmatic role of Prox1 and the control of LEC differentiation in development.

Introduction

Lymphatic vasculature plays crucial physiological roles that include the drainage of interstitial

fluids, trafficking of immune cells and drainage of dietary lipids. The formation of new

lymphatic vessels from pre-existing vessels (lymphangiogenesis) occurs in both development

and disease. Signalling through Vegfr3 (Flt4) can be triggered by Vegfc or Vegfd and drives

lymphangiogenesis in settings as diverse as development, cancer metastasis, inflammation

and ocular disease [1]. In the embryo, lymphangiogenesis begins when the first LEC progenitors

depart the cardinal veins (CVs) from E9.5 in mice and 32 hours post fertilisation (hpf) in

zebrafish [2]. Vegfc-Flt4 signalling drives LEC progenitor sprouting but also up-regulates Prox1

expression in both zebrafish and mice [3-7]. The transcription factor (TF) Prox1 acts as the

master regulator of LEC fate [8] and is exclusively expressed in developing LECs in early

embryonic vasculature. Loss of Prox1 (Prox1a and Prox1b in zebrafish) leads to a loss of

developing lymphatic vessels [4,9]. Following departure from the CV, LEC progenitors go on to

colonise embryonic tissues and organs and remodel to form functional lymphatic vessels [10].

While at later stages in mammals there are contributions of LECs from non-venous origins,

early embryonic lymphanigogenesis occurs chiefly from the CVs [11-17]. In mice, both the earliest

stages of LEC progenitor sprouting from CVs and maintenance of LEC fate in stable lymphatics

are dependent on the function of Prox1 [9,18].

Despite over two decades of study of this enigmatic developmental process, the

transcriptomic changes that occur as embryonic venous endothelial cells (VECs)

transdifferentiate into LEC progenitors, and further differentiate into mature lymphatics,

have not been transcriptionally profiled *in vivo*. In the absence of Prox1 in conditional

knockout mice, LECs have been shown to lose the expression of some LEC markers and to gain

expression of some blood vascular endothelial cell (BEC) markers[18]. Prox1 is known to

autoregulate its own expression and to also regulate Flt4 expression in a positive regulatory

loop during early development[7]. Yet how Prox1 controls the transcriptome during LEC

specification, differentiation and maintenance has not been described in detail. This is in part

because of the technical challenge of accessing early mouse endothelial cells (ECs) in wildtype

and mutant embryos, a problem that is not limiting when using the zebrafish embryo.

As recent studies have demonstrated highly conserved expression and function of Prox1

homologues in zebrafish[4,6,19,20], we here took advantage of the accessibility of the zebrafish

embryo to examine developmental lymphangiogenesis using single cell transcriptomics. We

provide a resource of new markers of VEC-LEC transdifferentiation, differentiating and

mature cell types and validate several new markers with transgenic approaches. We analysed

zebrafish zygotic *prox1a* mutants with single cell RNA sequencing and single cell ATAC

sequencing. This identified a VEC-LEC fate reversion in the absence of zygotic Prox1a, defined

key Prox1-dependent genes in fate maintenance and discovered regulatory regions of

chromatin (enhancers) controlled by Prox1. Finally, profiling maternal-zygotic double

prox1a/prox1b (null) mutant vasculature with single cell RNA sequencing revealed that Prox1

reduces gene expression in early LEC progenitors, including for a core network of conserved

haematopoietic and blood vascular fate regulators. Overall, this single cell resource reveals

embryonic lymphangiogenesis with Prox1-dependent mechanisms central in specification,

differentiation and maintenance of LEC fate.

Results

A single cell RNA-seq atlas of embryonic lymphangiogenesis

Zebrafish secondary angiogenesis occurs when Prox1-positive LECs and VECs both sprout

from the cardinal vein (CV) in the trunk or the head in a progressive process between ~32 and

~48hpf. In the trunk, sprouting LECs migrate dorsally and invest the horizontal myoseptum,

where they form a transient pool of parachordal LECs (PLs) from approximately 48hpf[21,22]. In

craniofacial regions of the embryo, LECs sprout from the CVs at several locations[23,24]. After

this, LECs throughout the embryo proliferate and migrate extensively (between ~56hpf-

80hpf) to colonise new regions and tissues[24-27]. In the trunk, LECs anastomose to form the

first lymphatic vessels at around 4 days post fertilisation; forming the thoracic duct (TD),

dorsal longitudinal lymphatic vessels (DLLVs) and intersegmental lymphatic vessels (ISLVs). In

craniofacial regions, they assemble from disparate sources into lateral (LFL), medial (MFL),

otolithic lymphatic vessels (OLV) and lymphatic branchial arches (LBA), as well as forming a

lymphatic loop (LL) in the head that will later give rise to a unique mural LEC population in the

brain (muLECs, also known as FGPs or brain LECs)[28-30]. By 5dpf, major lymphatics in the

craniofacial and trunk regions of the embryo are functional and drain dyes and fluids

deposited in the peripheral tissues[22,31].

To profile stages of development spanning key steps in lymphatic differentiation, we selected:

40hpf, when specification and initial sprouting of LECs are occurring; 3dpf, when immature

LECs are migrating through the embryo; 4dpf when LECs are assembling into vessels; and 5dpf

when lymphatics are functional and maturing[32] (**Fig 1a**). We used transgenic zebrafish strains

that specifically labelled embryonic vasculature to allow for fluorescence activated cell sorting

(FACS, full details in methods). We dissociated whole transgenic embryos and FAC sorted ECs,

for profiling on the 10X Chromium scRNA-seq platform. We sequenced 35,634 cells across 3

runs, then merged and normalised the data[33,34], filtering low-quality libraries (**Extended data**

Fig 1a, full details in methods). To define the cellular identity of each cluster, we

systematically evaluated the expression of known markers summarised in **Extended data**

Table 1a. We identified 9,771 lymphatic and venous ECs that comprise our single cell atlas of

lymphangiogenesis, which is accessible in an interactive CellXGene explorer web app at the

link http://115.146.95.206:5006/ (**Fig 1b**, **Extended data Fig 1b-c**).

This dataset is displayed[35] as a UMAP in **Fig 1b** coloured according to developmental stage

and cell phenotype, with venous (VEC) to lymphatic (LEC) trajectory of differentiation

confirmed by RNA-velocity analysis[36]. Comparison of cells at 40hpf with later stages revealed

that the earlier populations are transcriptionally distinct from VECs and LECs at 3, 4 and 5dpf.

Notably, the 40hpf "VEC preLEC" cluster contained both *prox1a+* and *prox1a-* cells forming a

single population rather than discrete clusters. This indicates that LECs are not

transcriptionally distinct from VECs at 40hpf, suggesting that while early *prox1a+* cells are

specified they have not yet differentiated (**Fig 1c**)[4]. We identified 3 main classes of LECs at 3,

4 and 5dpf: muLECs (1,511 cells) marked by expression of *osr2*[28], canonical LECs marked by

expression of *cdh6* (2,669 cells), and a smaller sub-population of LECs expressing low levels

of *prox1a* (LEC_low_prox1a, 310 cells) (**Fig 1c**, **Extended data Fig 1d-e**). To define markers of

differentiating canonical LECs at each developmental stages, we applied differential

expression (DE) analysis (**Extended data Table 1b**,). This analysis not only captured known

LEC markers including *prox1a, angpt2a, tbx1* and *cldn11b*, but also uncovered new genes

commonly expressed in canonical LECs across all developmental stages including *grp156,*

hapln3, cdh6 and *tspan18a* (**Fig 1d, Extended data Fig 1f**). We extended this approach and

evaluated global differences between all canonical LECs and all VECs (n=1,240 genes,

Extended data Table 1c). GO analysis[37] confirmed the association of biological processes

known to be associated with lymphatics with genes upregulated in LEC (n=752 LEC gene set),

these terms included "lymphangiogenesis", "glycolytic process", "lymph vessel development"

and "ameboidal-type cell migration" (**Fig 1e**, **Extended data Table 1d**).

To provide further confidence in the specificity of the new LEC markers defined by this scRNA-

seq resource, we used DE analysis to define cluster specific gene expression. We noted that

this analysis suggested higher expression of the well-known marker *lyve1b* in muLECs than in

LECs, which we validated by examining expression levels of *lyve1b* using the established

transgenic line *Tg(lyve1b:DsRed2)* (heat map in **Fig 2a**). We further identified expression of

the known kidney epithelial solute transporter *slc7a7* as uniquely expressed in muLEC cells

and *fabp11a* as a canonical LEC and VEC marker excluded from muLECs. We generated both

new *slc7a7a*–Citrine and a *fabp11a-Citrine* BAC transgenic strains, which confirmed the

muLEC and vascular expression patterns and further validated our scSeq atlas (**Fig 2b-c**).

Overall, this atlas therefore identifies n=752 specific markers of lymphangiogenesis, the

majority of which are new markers (**Extended data Table 1c**), representing new candidate

regulators and spanning four developmental stages.

Prox1 maintains LEC identity by repressing blood vascular fate and promoting lymphatic

vascular fate at the level of the transcriptome

Prox1 is both necessary and sufficient to drive VEC to LEC trans-differentiation in mammals

and this is proposed to occur by Prox1 simultaneously initiating LEC fate and repressing the

VEC fate program[8,18]. Prox1 expression is also necessary for the maintenance of LEC identity

during development[18,38,39]. Despite the role for Prox1 being well-studied in mouse models,

the transcriptomic program controlled by Prox1 *in vivo* has never been profiled. Zebrafish

prox1a zygotic and maternal zygotic mutants have been previously described[4,20]. The zygotic

mutants retain maternal deposition of *prox1a* in the oocyte, sufficient to drive normal PL

formation, LEC migration to the horizontal myoseptum (HM) and initial assembly of

lymphatics by 4dpf[4,19], however mutants have a reduction in total LEC numbers throughout

the face and trunk[4]. When the maternal contribution of *prox1a* is removed, lymphatic

development is more severely impaired[4]. We hypothesised that the zygotic mutants which

form lymphatic vessels, likely have abnormal vessel identity in the absence of zygotic Prox1.

Thus, we applied scRNA-seq to ECs FAC sorted from *Zprox1a[-/-]* mutants[20] and WT sibling

zebrafish at 4dpf, during lymphatic vessel assembly (n=8,075 cells, **Extended data Fig 2a**,

Extended data Table 2a). Cluster analysis revealed 3 populations of LECs (n=2,068) and a

single population of VECs (n=1,051) comprising both mutant and wild type cells, and perhaps

most striking, a single population of mutant cells (n=484) transcriptionally similar to VECs,

marked by the expression of *aqp1a.1* (**Fig 3a-d**, **Extended data Figs 2b-e**). All three LEC

clusters (LEC, LEC_S1, LEC_S2) showed graded expression of lymphatic markers *prox1a*,

cldn11b, *cdh6* and *angpt2a*, such that "LEC" most closely resembles canonical LECs from our

WT atlas. LEC_S1 and S2 represent alternative LEC sub-types with different proliferative

potential based on S-phase occupancy and cell cycle marker *mki67* (**Fig 3b,e**, **Extended data**

Fig 2e). The expression of *prox1a* was lowest in the more proliferative LEC_S1 cells, and

highest in the less proliferative LEC_S2 (**Fig 3b,e**). RNA-velocity analysis suggested a

trajectory between the mutant cluster and the LEC cluster (**Fig 3c** lower). Taken together,

these findings suggest that the mutant specific cluster sits on a trajectory between LEC and

VEC fate, representing cells either failing to fully differentiate from VEC to LEC or LECs

undergoing dedifferentiation and fate reversion.

To survey how Prox1 maintains normal LEC differentiation, we performed DE analysis

comparing the mutant cluster with WT LEC (**Fig 3f**, **Extended data Table 2b**). Overall, a much

larger set of genes were downregulated than upregulated in the mutant cluster (n=1,034 vs

n=294) with almost half of the most downregulated genes (AvLogFC) highly enriched for LEC

markers (eg. *tbx1*, *cdh6*, *cldn11b*). Of the upregulated genes, almost 75% were VEC markers

(eg. *sox7*, *kdrl*, *cdh5*) and again this was enriched in the most highly upregulated genes (**Fig

3f, Extended data Fig 2f**). Comparing the change in gene expression between the wildtype

LEC cluster and VEC cluster, with the change between wildtype LEC and mutant cluster, we

found striking concordance suggesting the mutant cluster is shifted along a LEC fate to VEC

fate trajectory (**Fig 3g**). Overall, there is a simultaneous loss of lymphatic fate and re-

acquisition of blood vascular gene expression, consistent with work in mouse and with Prox1

function being highly conserved between vertebrates. To validate these observations, we

used confocal imaging of Z*prox1a*$^{-/-}$ mutants and observed upregulation of blood vascular

markers *cdh5* (**Fig 3h-i, Extended data Fig 3a,c**) and *kdrl* (**Fig 3j-k, Extended data Fig 3d-g**) in

lyve1b-positive lymphatics vessels. We saw a coincident reduction in the expression of *lyve1b*

in these vessels (**Extended data Fig3b).** Interestingly, the difference in *kdrl* levels (relative

intensity) in lymphatics between mutants and wildtype was more significant at 5dpf than

4dpf, suggesting that progressive dedifferentiation may be occurring in the mutant (**Fig 3k**).

This single cell profile thus demonstrates that Prox1 alone is sufficient to maintain LEC fate

and identifies the gene expression maintained by Prox1 during lymphatic differentiation.

Single cell ATAC sequencing reveals chromatin accessibility signatures in LECs and VECs,

identifying lymphatic specific enhancers and predicting key LEC TF families

At the level of gene expression, Prox1 function is essential for VEC to LEC differentiation,

however it is unknown if Prox1 controls chromatin accessibility during this process. Thus, we

next profiled Z*prox1a*$^{-/-}$ mutants at 4dpf during vessel assembly using single nuclei (sn) ATAC-

seq (n=3,731 nuclei, **Extended data Fig 4a, Extended data Table 3a**). Cluster analysis (**Fig 4a**)

and overall accessibility of key markers (**Fig 4b**) identified similar populations to the scRNA-

seq profiling: canonical LEC (LEC_01 n=114 nuclei, LEC_02 n=213 nuclei) and VEC (VEC_01

n=157 nuclei) clusters, and a small but discrete population comprised almost entirely of

mutant cells (Mutant Cluster n=47 nuclei). Notably, we found that there was almost no

contribution of mutant cells to the LEC clusters (**Fig 4a** lower bar plot), indicating a loss of fate

when cell identity is determined at the level of chromatin state **(Extended data Fig 4b)**.

Differential accessibility (DA) analysis revealed unique sets of peaks that mark individual VEC,

LEC and AEC (arterial EC) clusters (**Fig 4c**).

To identify phenotype specific chromatin accessibility, we performed DA analysis between

WT LEC and WT VEC. This revealed the more accessible regions in the LEC clusters were

associated with LEC genes identified in our atlas, and the less accessible regions with VEC

genes (**Fig 4d, Extended data table 3b**). DA identified n=1,561 LEC specific peaks and n=2,624

VEC specific peaks representing putative lineage specific enhancers or regulatory elements

(**Extended data table 3b**). To test if these regions identified enhancers, we used the zebrafish

enhancer detector plasmid system (ZED vector[40]) and tested peaks that were uniquely open

5' of the newly identified LEC marker gene *cdh6.* To exclude non-specific reporter expression,

we also generated F1 embryos using a ZED vector only control. This did display mosaic

neuronal and dorsal root ganglion expression **Extended data Fig 4h**, suggesting that the

expression from the *cdh6* enhancer in neural structures using this vector should be taken with

some caution. No vascular expression was detected. Transgenesis identified one new

functional and LEC specific enhancer upstream of *cdh6*, validating the use of this dataset for

enhancer discovery (**Fig. 4e-f**).

We next aimed to use this new set of putative enhancer regions in an unbiased manner to

identify key TF families likely to regulate LEC development. We performed motif enrichment

analysis using HOMER[41] for all LEC enriched DA peaks identifying n=64 TF family motifs

(**Extended data table 3c**). Notably, we found motifs for TCF, ETS, SCL (TAL1), NFAT, TBX, MAF,

SLUG and RBPJ family TFs to be enriched in peaks more accessible in LECs (**Fig. 4g**). Analysis

of the human homologues in ENCODE data[42] revealed these TFs regulate a highly connected

network of genes expressed in our LEC atlas (**Fig. 4h**). Importantly, a number of these TFs are

already known to play important roles in lymphatics (e.g. NFAT[43,44], MAFB[45,46], TBX1[47],

TCF[48,49]), supporting the prediction that members of these TF families will play important

function roles in LEC development.

Prox1 reduces the accessibility of chromatin peaks associated with blood vascular and

haematopoetic TF motifs

Given that cells in the mutant specific cluster analysed by scRNA sequencing demonstrate loss

of LEC gene expression and acquisition of VEC gene expression, we expected chromatin

accessibility to change in an equally coordinated manner. However, analysis of accessibility

at individual genes (gene score) for the Mutant Cluster revealed changes inconsistent with a

simple fate shift (**Extended data tables 4a-b**). Some LEC specific genes (eg. *prox1a*, *cdh6* and

tbx1) showed loss of transcription in the mutant cluster but increased chromatin accessibility,

and some specific VEC genes with increased transcription in the mutant cluster (eg. *cdh5, flt1,*

gata6) also showed discordant chromatin changes (**Fig 5a**-**b, Extended data Fig 4g**). A DAP

analysis revealed little concordance between regions with increased accessibility, and

expression in LEC or VEC in the scSeq atlas (**Fig 5c**). We identified a subset of genes with more

accessible chromatin overall in the Mutant Cluster than either WT LEC or VEC settings (**Fig**

5d). Notably, at the level of individual peaks we found that n=1,726 peaks displayed a striking

increase in accessibility in the mutant cluster compared with WT LEC and n=1,794 peaks an

increase compared with WT VEC (**Fig 5e**, **Extended data tables 4a-b**). Of these, 431 were

common peaks identifying more accessible chromatin regions in the mutant cluster than in

either WT LEC or WT VEC (**Fig 5e,** examples in **Fig. 5f**). This suggests that some regions of

chromatin open up more than usual during LEC or VEC differentiation in the absence of Prox1.

To investigate the nature of the peaks that were opening in the absence of Prox1, we used an

unbiased assessment of TF motifs within these regions.   The more open regions were highly

enriched for motifs of early acting TFs involved in embryonic vasculogenesis and

haematopoiesis, including Erg, Etv2, Etv4, Ets1, Fli1 and Spi1/Pu.1 (**Fig 5g**, **Extended data**

table 4c). We examined the human homologues of these TFs in ENCODE[42] data together with

homologues of genes expressed in our atlas and identified a highly connected putative gene

regulatory network (GRN) made up of early blood and blood vascular TFs (including ETV2,

TAL1, SOX7, SOX17 ERG and other key fate regulators) driving target genes including each

other, FOS, JUN, MYC and STAT3 (**Fig. 5h**). We take this to suggest that in the absence of

Prox1, TFs that are part of a GRN normally suppressed by Prox1 function are reactivated to

drive blood vascular and blood fates. This demonstrates that Prox1 coordinates the correct

accessibility of the chromatin and likely controls other TF functions while maintaining LEC fate

in ECs. Furthermore, we note that a large number of the genes with increased chromatin

accessibility are known regulators of lymphangiogenesis and so this increased accessibility

may be a sensitive way to identify key regulatory genes (**Extended data table 4c**).

Prox1 is required cell autonomously for the normal sprouting of lymphatic progenitors and

their contribution to the lymphatic lineage

While the above data identifies the role of Prox1 at stages when it is maintaining LEC identity

in assembling LECs, Prox1 is essential for the very earliest decision made when a VEC becomes

specified to develop into a LEC[9]. To ask how Prox1 controls the earliest stages of VEC to LEC

transdifferentiation, we first needed to generate a complete Prox1 loss of function model in

zebrafish. We generated *prox1a*$^{+/-}$, *prox1b*$^{+/-}$ double heterozygous animals (lacking both Prox1

homologues) and then used germline transplantation approaches to produce animals

carrying a double mutant germline (**Extended data Fig 5a**). Genetic crossing of these animals

generated embryos that were maternal zygotic (MZ) *MZprox1a*$^{-/-}$, *MZprox1b*$^{-/-}$ mutant

zebrafish completely lacking *prox1a/b* transcript expression or maternal deposition (**Fig 6a;**

Extended data Fig 5b-c). A quantitative phenotypic analysis revealed double MZ mutants

show a severe reduction of facial lymphatics and a near complete loss of lymphatic vessels in

the trunk by 4dpf (**Fig 6e**, **Extended data Fig 5d-g**) [4]. Furthermore, *MZprox1a*$^{-/-}$, *MZprox1b*$^{-/-}$

mutants initially show a delay in the formation of PLs as these cells emigrate the CV to invest

the HM, but despite this delay, these cells eventually seed the HM before failing to undertake

any further migration (**Fig 6d,f**). We saw evidence for genetic interaction between *prox1a* and

prox1b in PL formation and no change in the overall number of cells sprouting from the PCV

or number of cells contributing to the venous ISVs (blood vessels) suggesting both *prox1a* and

prox1b contribute specifically to early PL development (**Fig 6g-k**). To test whether LECs require

Prox1a in a cell autonomous manner, we performed embryonic transplantation to generate

chimeric embryos. Due to challenges generating and maintaining large numbers of double MZ

mutant embryos, we performed transplantion of *MZprox1a*$^{-/-}$ mutant cells into wildtype hosts

only and assessed the contributions of vascular grafts to arteries, veins and lymphatics

(**Extended data Fig 6a**). MZ*prox1a*$^{-/-}$ mutant cells efficiently contributed AECs, VECs but not

LECs to developing vessels in otherwise wildtype hosts at 5dpf (**Extended data Fig 6b-d**). Thus,

we confirmed that Prox1 is necessary cell autonomously for zebrafish lymphatic development

and that the *MZprox1a^-/-^, MZprox1b^-/-^* mutant phenotype is more severe than any previously

described zebrafish Prox1 mutants.

Prox1 functions to suppress blood and blood vascular fate during the early specification of

LEC fate in the embryo

To understand the very earliest role that Prox1 plays in LEC development, we profiled the

endothelium of *MZprox1a^-/-^, MZprox1b^-/-^* mutant zebrafish at 40hpf (when cells are both

being specified to LEC fate and also actively sprouting from various regions of the PCV[3,4,6])

using single cell RNA seq as described above (**Extended data table 5a, Extended data Fig 7a-**

c). Using key marker expression and DE analysis, we identified populations of cells that

included VECs, endocardium, mixed populations of VECs and Prox1+ LEC progenitors (named

LEC_VEC) and clusters with expression of some VEC and AEC markers that were likely still

differentiating (**Fig 7a-c, Extended data Fig 7d-h**). Based on expression of known markers and

genes associated with EC sprouting (*mki67, pcna*), we defined two populations of secondary

sprouts of venous origin (LEC VEC 01 n=713 cells, LEC VEC 02 n=677 cells) and a single

population of cells representing the cardinal vein (PCV n=812 cells; **Fig 7d-e**). Consistent with

our observations in **Fig 1,** cells expressing *prox1a* at 40hpf were not transcriptionally distinct

from sprouting VECs and failed to form a "lymphatic progenitor" cluster, suggesting they are

specified and express *prox1a* but not yet differentiated. RNA velocity analysis suggested that

these three clusters remained closely related, consistent with little differentiation between

these populations at this stage (**Fig 7a** right).

To identify the earliest transcriptional changes controlled by Prox1 in lymphatic development,

we used DE to evaluate the global differences between *MZprox1a^-/-^, MZprox1b^-/-^* and WT in

the LEC VEC 01, LEC VEC 02 and PCV clusters of sprouting cells (**Fig 7f, Extended data table**

5b). We found a significant upregulation of a large set of genes in the absence of Prox1 that

were identified as enriched for blood vascular and haematopoietic genes by GO-term and

marker analysis (**Fig 6g-h,** n=1,137 genes, **Extended data table 5c**). We also saw increased

expression of mitochondrial metabolism genes and of pre-mRNA splicing genes in the

absence of Prox1 (**Fig. 7f, Extended data table 5c**). At this stage of development, in contrast

to the 4 dpf stage, we saw little evidence of positively regulated genes downstream of Prox1

(just n=49 genes downregulated in *MZprox1a$^{-/-}$, MZprox1b$^{-/-}$*). We next examined the genes

up-regulated in Prox1 mutants that we identified as key early blood and vascular

developmental regulators. This includes *esm1, flt4, mef2c, hdac1, lmo2, lmo4, rasip1, egfl7,*

dusp5, clec14a, fli1b and others (**Fig. 7h-i**). STRING[50] analysis suggested the presence of inter-

related gene networks and so we further examined up-regulated TFs in this network by

leveraging human homologues in ENCODE data and identifying known target genes expressed

in our dataset. This identified a GRN containing SOX7, FLI1, LMO2, ETV2, TCF12, LMO4 and

MAFB as well as known targets of these TFs, that is repressed in response to Prox1-function

during LEC specification (**Fig. 7j**). This is highly concordant with observations at later stages of

up-regulated TF activity in 4-day old mutants (**Fig. 5**). Overall, we believe this suggests that in

the earliest stages of LEC specification Prox1 is not actively driving a fate program but rather

is acting primarily to down-regulate blood vascular and hematopoietic fate. It seems likely

that the negative regulation of core blood and blood vascular fate genes (those in **Fig. 7j**),

would be sufficient to drive the differentiation in these earliest specified LECs.

Discussion

In this study, we first present a single cell RNA seq analysis of four key stages of embryonic

lymphangiogenesis in a vertebrate embryo, revealing new markers and potential regulators

of LEC differentiation. We further profile the ECs of *Zprox1a* mutants, which form lymphatic

structures, but these "lymphatics" dedifferentiate or revert their fate to blood vascular in a

mutant specific, fate-shifted, single cell cluster. This analysis identified the transcriptional

code maintained by Prox1 in order to maintain LEC identity and also shows the highly

conserved nature of Prox1 function comparing zebrafish with mice. Overall, the striking fate

shift identified across the whole transcriptome confirms that the function of just one

transcription factor (Prox1) is sufficient to maintain cellular identity, validating its status as

the master regulator of LEC identity.

As well as single cell profiling of the developing LEC transcriptome, we also performed snATAC

seq on 4 dpf zygotic *prox1a* mutants and wildtype, again identifying a mutant specific, fate

shifted cluster in this analysis. Analysis of the wildtype LECs and VECs revealed strong

concordance between chromatin accessibility at LEC enhancers and the transcriptional profile

for LEC and VEC specific genes. Interestingly however, analysis of the mutant cluster

chromatin identified ectopically open regions and peaks with a distinct discordance between

chromatin accessibility at enhancers and transcriptional profile. In the mutant cluster at 4dpf,

a large number of opened chromatin regions were enriched for TF motifs for earlier acting

vasculogenic and haematopoietic TFs. This suggests that regulators of earlier vascular fate

become more active and increase chromatin accessibility at specific targets in the absence of

Prox1. Overall, this revealed that at the level of chromatin accessibility the fate-shifted ECs

display a more immature state, perhaps a consequence of regulatory "confusion" due to a

failed fate transition. It seems likely that Prox1 has a combinatorial function as part of a larger

GRN of developmental TFs, which has yet to be studied in detail.

Additional biological insights from this study come from our analysis of double maternal

zygotic *prox1a, prox1b* mutants. These mutants are presumed "null" mutants for Prox1

orthologues and they revealed (through scRNA seq) that during its earliest role in vascular

development, in LEC fate specification and VEC-LEC transdifferentiation, Prox1 functions

primarily to negatively regulate blood and blood vascular fate. While this may represent a

downstream program rather than direct repression of gene expression by Prox1 orthologues,

it is notable that Prox1 and *Drosophila* Prospero have been reported to be able to function as

repressors in a context dependent manner [51-55]. Interestingly, the analysis of upregulated

genes identified a set of TFs that are known early regulators of blood and blood vascular fates

during embryonic haematopoiesis and vasculogenesis. These included Sox7, Etv2, Lmo2 and

Lmo4 and we take this to suggest that Prox1 functions during VEC-LEC transdifferentiation at

least in part by blocking expression of early acting blood vascular fate driving TFs. This is in

line with maintaining a negative regulatory relationship with these and other fate driving TFs

at 4 dpf as described above. It will be interesting in the future to understand at a mechanistic

level if Prox1 is actively repressing gene expression at bound targets to block alternative cell

fates and if so, if the activity of Prox1 switches between repressor or activator at different

stages or target genes.

Altogether this study describes the process of developmental venous to lymphatic

transdifferentiation, early LEC differentiation and LEC maintenance in detail, *in vivo*. We

describe the role of Prox1 during this process, revealing a conserved and dynamic regulatory

process with unprecedented resolution. This resource will help to understand

lymphangiogenesis in contexts beyond the embryo, such as in pathological

lymphangiogenesis in metastasis, inflammation and tissue repair. Finally, we suggest that

combining zebrafish mutant models, with their inherent ease of cellular accessibility, and the

resolution afforded by single cell profiling, will be an approach that yields powerful new

insights in many areas of developmental biology in the future.

Acknowledgements

This project was supported in part by NHMRC Ideas grant 2004300 and ARC Discovery project

DP180102846. B.M.H. was supported by NHMRC Fellowship 1155221. We thank Angelika

Christ and the IMB Sequence Facility (University of Queensland), Tim Semple and Peter Mac

Genomics Facility, the Research Computing Facility at Peter Mac, FACS facilities at UQ and

Petermac and Isaac Virshup for assistance.

Author contributions

L.G. and E.M. performed, analysed experiments and co-wrote manuscript. B.M.H, K.K and

N.L.H conceptualised experiments, analysed data and co-wrote manuscript. S.D, T.C, O.Y,

N.I.B, S.P, K.O, A.S and A.L. performed and analysed experiments. J.P, and K.A.S, provided key

unpublished reagents and analysed data.

Conflicts of interest

The authors declare no conflict of interests.

Online Methods

Zebrafish husbandry

Zebrafish work was conducted in compliance with animal ethics committees at the Peter

MacCallum Cancer Centre, the University of Melbourne and the University of Queensland.

Published transgenic lines used were: *Tg(fli1a:nEGFP)^{y7}* [56]; *Tg(-5.2lyve1b:DsRed)^{nz101}* [24]; and

Tg(kdrl:Has.HRAS-mCherry)^{s916} [21]. Published mutant lines used were *prox1a^{i278}* [20], *prox1b* [4,19].

Tg(lyve1b:Venus)^{uq51bh} was generated as previously described[28] but here using an

independent genomic integration with the same construct.

Generation of maternal zygotic mutants

Germline replacement was performed using embryonic transplantation as previously described [4,57]. Maternal zygotic (MZ) *prox1a* mutant embryos were made by crossing germline replaced *prox1a^{i278−/−};prox1b^{sa0035+/−}* females with *prox1a^{i278+/−};prox1b^{sa0035+/−}* males [4,57]. Genotyping of individual embryos during transplantation and phenotypic analysis was performed as previously described [4].

Transgenesis, genome-editing and genotyping

All microinjections were performed as previously described [58]. *slc7a7a^{BAC}:slc7a7a-Citrine^{uom10}* and *fabp11a^{BAC}:fabp11a-Citrine^{uom10}* recombineering was performed as previously described [59].

Primers for BAC recombineering:

slc7a7a-BAC-Citrine-forward:5'-
AACTGCTTTAGACAGTGTTTTTTGGTACCATCCCATATATTTAAAAAACAGCCACCATGGTGA
GCAAGGGCGAGGAG-3'

slc7a7a-BAC-Citrine-reverse:5'-
TTCGACACCTCAGGGGATGCCTCTTCTGCAGGCGTAGGGCTGTAGGACGCTCAGAAGAACT
CGTCAAGAAGGCG-3'

fabp11a-BAC-Citrine-forward:5'-
TTACAGCTGTTGCGAGATTGAAAAGTAGAGGAGCATCATTATTCGGGAAAGCCACCATGGT
GAGCAAGGGCGAGGAG-3'

fabp11a-BAC-Citrine-reverse:5'-
TCAAAGTTGTCGCTGGTGGTCATTTTCCACGTTCCTACGAATTTGTCAACTCAGAAGAACTC
GTCAAGAAGGCG-3'

Enhancer detection transgenesis and analysis

A 501bp PCR fragment of *cdh6* enhancer (*chr2:28150709-28151209*) was cloned into the zebrafish enhancer detection (ZED) vector[40]. Empty ZED vector was injected as previously described[40]. Briefly, 1nL of construct at 40 ng/µl or 45ng/µL and tol2 transposase mRNA at 100 ng/µl or 55ng/µl was injected into the one-cell stage wild type zebrafish embryos. All F0 embryos were screened for skeletal muscle DsRed2 expression. Stable F1 embryos were imaged on a Zeiss LSM 780 or Leica TCS SP8 DLS microscope. Images were processed using

490   ImageJ 2.0.0. or 2.3.0.

Imaging and quantification

Imaging was conducted at the Centre for Advanced Histology and Microscopy (Peter MacCallum Cancer Centre). Imaging of live samples was performed using a Zeiss LSM 710 FCS confocal microscope, a Zeiss LSM 780 FCS confocal microscope, or an Olympus FV3000 confocal microscope. Mounting and imaging were performed as previously described [60]. In **Figs 6e-f,i-k**, and **Extended data Fig 5f-h, k-m** quantification of vascular phenotypes was performed as previously described [60,61]. In Figs 3h-k, *cdh5:Venus* or *kdrl:GFP* intensity in the TD was measured using Imaris software (Bitplane) and normalised to fluorescence intensity of the DA for *cdh5* and to the fluorescence intensity PCV for *kdrl* in the same embryos.

Fluorescence activated cell sorting

FACS was performed at the Peter MacCallum Cancer Centre and the University of Queensland. For **Fig 1**, we isolated cells using the following transgenic lines: 40hpf, *Tg(fli1a:nEGFP)^{y7}*; 3,4 and 5dpf, *Tg(-5.2lyve1b:Venus)^{uq47bh}*, *Tg(kdrl:Has.HRAS-mCherry)^{s916}* For **Fig 3 and Fig 4**, we isolated cells using the following transgenic lines: 4dpf, *Tg(fli1a:nEGFP)^{y7}*, *Tg(-5.2lyve1b:DsRed)^{nz101}*. For **Fig 6**, we isolated cells using the following transgenic lines: 40hpf, *Tg(fli1a:nEGFP)^{y7}* .To dissociate embryos and obtain single cell suspensions, we followed published protocols [62]. Briefly, at the desired developmental stage we deyolked embryos by pipetting up and down and rinsing in calcium free ringers solution, we centrifuged at 2000rpm for 5' at 4°C and remove supernatant and dissociated the cells by incubating in liberase [2.5mg/mL] (Cat #5401119001 Sigma-Aldrich) diluted at a 1:35 ratio in DPBS at 28.5°C for approximately 5', homogenizing the samples during and after the incubation. To stop the reaction, we added CaCl2 to a final concentration of 1-2mM and FBS to a final concentration of 5-10%. We centrifuged at 2000rpm for 5' at 4°C. discarded the supernatant, in order to be able to asses live vs. dead cells, we re-suspend the cells solution in Zombie Violet TM Viability Dye (Cat# 423113, BioLegend) and incubated for 20' at RT softly rocking, we rinsed the cells by centrifuging and resuspending in DPBS/EDTA, for ATAC-seq experiments samples were resuspended in 2%BSA/PBS. Suspension was filtered through a strainer and taken to the FAC sorting facility. In the Flow Cytometry facility, we used the BD FACS Aria Fusion sorter (BD Biosciences), we based the selection for the desired population

on FSC and SCC, alive cells were selected based on the Zombie Violet profile and double

positive cells for the desired transgenics were targeted according to the expression profiles

of single cells. Double positive cells were sorted in 300uL 100% FBS in a cold block and taken

immediately to the sequencing facility.

scRNA-seq library preparation

Library preparation and sequencing was performed at the Institute for Molecular Bioscience

Sequencing Facility (University of Queensland) or Peter Mac Genomics Facility. Single cell

suspensions were sorted by FACS, spun down to concentrate and a cell count was performed

to determine post-sort viability and cell concentration. Single cell suspension was partitioned

and barcoded using the 10X Genomics Chromium Controller (10X Genomics) and the Single

Cell 3' Library and Gel Bead Kit (V2 10X Genomics PN-120237; V3.1; 10X Genomics; PN-

1000123). The cells were loaded onto the Chromium Single Cell Chip A (10X Genomics; PN-

120236), B (10X Genomics; PN-1000073 or PN-1000074) or G (10X Genomics; PN-1000120)

to target 10,000 cells. GEM generation and barcoding, cDNA amplification, and library

construction was performed according to the 10X Genomics Chromium User Guide. The

resulting single cell transcriptome libraries contained unique sample indices for each sample.

The libraries were quantified on the Agilent BioAnalyzer 2100 using the High Sensitivity DNA

Kit (Agilent, 5067-4626). Libraries were pooled in equimolar ratios, and the pool was

quantified by qPCR using the KAPA Library Quantification Kit - Illumina/Universal (KAPA

Biosystems, KK4824) in combination with the Life Technologies Viia 7 real time PCR

instrument. After the initial sequencing run, libraries were re-pooled according to estimated

captured cells as determined using the Cell Ranger software (10X Genomics).

Sequencing of scRNA-seq libraries

At the IMB(UQ) genomics facility, denatured libraries were loaded onto an Illumina NextSeq-

500 and sequenced using a 150-cycle High-Output Kit as follows: 26bp (Read1), 8bp (i7 index),

98bp (Read2). Read1 supplies the cell barcode and UMI, i7 the sample index, and Read2 the

3' sequence of the transcript. At the Peter Mac Molecular Genomics facility, single cell

transcriptome libraries were sequenced on an Illumina NovaSeq 6000 using S4 300-cycle

chemistry. Read1 supplies the cell barcode and UMI, i7 the sample index, and Read2 the 3'

sequence of the transcript. Sequencing read lengths were trimmed to 28bp (Read1), 8bp (i7

index), 91bp (Read2), ensuring compatibility with the 10X Genomics analysis software, Cell

Ranger.

scRNA-seq data processing and analysis

All code and documentation associated with this analysis is publicly available under an open

source software license at:

https://atlassian.petermac.org.au/bitbucket/users/tyrone.chen/repos/hogan_lab/browse/2

022_Grimm_Mason_et_al_PROX1/

Relevant functions are in italics for reference. Where necessary fastq files were made using

Cell Ranger [63] (version 3.1.0 or 3.0.2) *mkfastq*. Sequencing QC was assessed using *FastQC*

0.11.6 and *MultiQC* viewer for aggregated reports. *Cell Ranger count* and *aggr* were used to

generate aggregated count files mapped to GRCz11 (Ensembl 101), without depth

normalisation. Doublets were identified from the filtered aggregated count files using

Scrublet [64] in *Python* version 3.6 and filtered from subsequent analyses. For the *MZprox1$^{-/-}$*

mutant and *Zprox1$^{-/-}$* mutant datasets filtered aggregated count files were processed, sc-

transform normalised, filtered and clustered (louvain) using *Seurat* version 2.0 [65] and 3.0 [34]

respectively for *R statistical software* version 4.0.2. QC was evaluated before and after

normalisation using plot functions in *Seurat* and *scater* 1.20.1 [66], and all thresholds and

settings are described in scripting. Cluster solutions were evaluated using *ClusTree* [67].

Datasets used in the atlas of lymphangiogenesis were processed, filtered, merged and log-

normalised using *Seurat* version 3.0 [34], with QC and settings as above. Merged data was

clustered and normalised using *CSS simspec* [33] and clustering and cluster evaluation

performed on this object only, as described above. For all scRNA-seq datasets cluster

phenotype was determined using the expression of key markers (Supplementary Table), with

the aid of *CellXGene* visualisation software [68]. All downstream analysis and plotting were

performed using *Seurat* version 3.0 [34] with default settings. All gene ontology analyses were

performed using Panther.db [37] (Biological Process Complete).

Preparation of single nuclei for snATAC-seq

Single cell suspensions were sorted by FACS and prepared for nuclei isolation as previously

described by 10x Genomics Demonstrated Protocol for Single Cell ATAC Sequencing

(CG000169 - Rev D). Cell suspensions were pelleted (300 x g for 5 minutes) and rinsed with

PBS + 0.04% BSA. Cells were resuspended in 95uL of freshly prepared lysis buffer (10mM Tris-

HCl pH 7.4, 10 mM NaCl, 3 mM MgCl2, 0.1% Tween-20, 0.1% NP40 Substitute, 0.01%

Digitonin, and 1% BSA) and incubated on ice for 1 minute. 100uL of chilled wash buffer

(10 mM Tris-HCl, pH 7.4, 10 mM NaCl, 3 mM MgCl2, 0.1% Tween-20, 1% BSA) was used to

neutralise the reaction, before the nuclei were pelleted (500 x g for 5 minutes) and

resuspended again in 7uL of 1x Nuclei Buffer (10X Genomics Cat# PN-2000153/2000207).

Presence of healthy and intact nuclei was assessed by visual inspection on a brightfield

microscope using Trypan Blue staining (Thermo Fisher Cat# T10282) and Countess Cell

Counting Chamber Slides (Thermo Fisher Cat# C10228).

snATAC-seq library preparation and sequencing

Single nuclei suspensions were resuspended at approximately 5000 nuclei per μL before

undergoing tagmentation for 60min at 37°C. After tagmentation nuclei were partitioned and

barcoded using the 10X Genomics Chromium Controller (10X Genomics) and the Single Cell

ATAC Reagent Kit (V1.1; 10X Genomics; PN-1000176). Tagmented nuclei were loaded onto

the Chromium Single Cell Chip H (10X Genomics; PN-1000162), GEM generation, barcoding

and library construction was performed according to the 10X Genomics Chromium User

Guide. The resulting single cell ATAC libraries contained unique sample indices for each

sample. The libraries were quantified on the Agilent BioAnalyzer 2100 using the High

Sensitivity DNA Kit (Agilent, 5067-4626) and pooled in equimolar ratios. Sequencing was

performed on an Illumina NextSeq 500 using a 150-cycle High-Output Kit as follows: 50bp

(Read1), 8bp (i7 index), 16bp (i5 index), 50bp (Read2) achieving a read depth of 25,000 read

pairs per nucleus.

snATAC-seq processing and analysis

FASTQ files generated from sequencing were used as inputs to 10X Genomics *Cell Ranger*

ATAC 2.0.0. *cellranger-atac count* was used to generate count files mapped to GRCz11

(ENSEMBL 101), without depth normalisation. Resulting fragment files were read into *ArchR*

1.0.1 for *R statistical software 4.0.5* as a tile matrix with 500-bp bins. All remaining steps in

the ATAC-Seq analysis were performed within *ArchR 1.0.1*. QC filtering was performed, and

only high-quality cells with a TSS enrichment score greater than 4 and greater than 1,000

unique nuclear fragments were retained. Doublets were predicted using *addDoubletScores*

and filtered using *filterDoublets.* Data normalization and dimensionality reduction were

performed using iterative Latent Semantic Indexing (LSI) and Uniform Manifold

Approximation and Projection (UMAP) embeddings were used for visualisation in reduced

dimension space. Separate from the *ArchR 1.0.1* package, cluster solutions were

independently evaluated using *clustree 0.4.3*. A Gene Score Matrix that stores predicted gene

expression was then generated based on the accessibility of regulatory elements in the

vicinity of the gene. We used gene scores of endothelial markers for cluster annotation and

subsetting. Differentially accessible genes were identified by differential testing using

getMarkerFeatures. Local chromatin accessibility of the marker genes was visualised using

the *plotBrowserTrack*.

Gene regulatory network analyses

For gene regulatory network construction we leveraged the ENCODE transcription factor

targets gene-attribute edge list from the Harmonize database

https://maayanlab.cloud/Harmonizome/dataset/ENCODE+Transcription+Factor+Targets

that includes information for n=181 transcription factors from ChIP-seq analyses[42]. Gene lists

generated from scRNA-seq or snATAC-seq analysis were mapped from Zebrafish genome

version GRCz11 to Human genome version GRCh38.p13 using ENSEMBL Biomart [69]. These

mapped gene sets were used to select relevant edges, that were visualised as a degree-sorted

circular network in Cytoscape[70].

639
640

FIGURES AND FIGURE LEGENDS

Figure 1: A single cell RNAseq developmental atlas of lymphangiogenesis in zebrafish

    a. Schematic representation of four key stages of zebrafish lymphatic development in head and trunk: 40hpf encompasses both specification (Prox1-induction) and sprouting; 3dpf migration of LECs; 4dpf assembly of lymphatic vessels; 5dpf maturation functional lymphatics.

b. UMAP visualization of n=9,771 cells filtered for VEC and LEC populations (n=6 samples;
see **Extended data table 1a** for cluster identification and **Extended data Fig 1** for
whole dataset) coloured according to developmental stage (top), predicted cell
phenotype (middle), and RNA-velocity (bottom).

c. UMAP visualization of key marker gene expression. Colour scale represents log-
normalised expression. LEC markers: *prox1a, cdh6*. BEC markers: *cdh5*. LEC and VEC
markers: *mrc1a, flt4*. AEC marker: *flt1*. muLEC markers: *osr2, hexa*.

d. Dot plot of top genes commonly marking LECs at 3, 4 and 5dpf, with expression
displayed in VECs (all stages), VEC_preLEC (40hpf) and LECs at 3, 4 and 5dpf (see also
Extended data table 1b, **Extended data Fig 1f**). Colour scale represents average log-
normalised expression and point size represents percentage of cells expressing gene.

e. Bar plot summarizing GO BP analyses of genes DE between 3, 4 and 5dpf LEC and VEC
populations (**Extended data table 1c-d**). Y-axis represents enriched BP term, x-axis
represents the -log10(*adjusted p value*), bars are coloured and ordered according to
fold enrichment of the GO term in LEC. GO terms enriched in LEC are coloured purple,
terms enriched in VEC are coloured red.

Hpf, Hours post-fertilisation.dpf, Days post fertilisation. BEC, Blood vascular endothelial cell.
VEC, Venous endothelial cell. LEC, Lymphatic endothelial cell. muLEC, mural Lymphatic
endothelial cell (a.k.a FGP or brain LEC).

Figure 2: Transgenic marker strains confirm cluster identity and identify vessel specific markers

a. UMAP of *lyve1b* expression predicts higher expression in muLECs than LEC populations (left), confirmed by a heat map of a *Tg(lyve1b:DsRed2)* zebrafish larvae at 6dpf showing high *lyve1b* expression in craniofacial (middle) and moderate expression in trunk lymphatic vessels (right).

b. UMAP of *slc7a7* expression predicts restricted expression in muLEC populations (left), confirmed in lateral confocal projections showing co-expression of *lyve1b* (magenta) and a new BAC transgenic strain for *slc7a7a* (green) in the zebrafish head at 5dpf (middle) which is not expressed in veins and lymphatic vessels of the trunk (right).

c. UMAP of *fabp11a* expression predicts expression in LEC and VEC but not muLEC populations (left), confirmed in lateral confocal projections showing co-expression of *lyve1b* and a new BAC transgenic strain for *fabp11a* marking venous and lymphatic vessels in the trunk (right) without showing expression in the vasculature of the head (middle) at 5dpf.

Lateral confocal images, anterior to the left. muLECs, mural lymphatic endothelial cells; facial LV, facial lymphatic vessels; vISV, venous intersegmental vessel; PCV, posterior cardinal vein; TD, thoracic duct; ISLV, intersomitic lymphatic vessel. Scale bars, 80 μm for head (middle) in (a) and 50 μm for trunk images (right).

Figure 3: Single cell RNA seq analysis reveals a fate shift from LEC to VEC in the absence of Prox1 in zygotic *prox1a* mutants

a. UMAP visualization of n=8,075 cells filtered for VEC and LEC subpopulations (n=2 samples; see **Extended data table 2a** for cluster annotation, **Extended data Fig3** for full dataset) coloured according to predicted cell phenotype.

b. UMAP visualization of marker gene expression. Colour scale represents SCT-normalised expression. LEC markers: *prox1a, cldn11b, cdh6, angpt2a*. LEC and VEC markers: *flt4, mrc1a, lyve1b*. BEC markers: *cdh5, kdrl*. Mutant cluster markers: *frem1b, aqp1a.1.*

c. UMAP visualization with cells coloured according to genotype (*Zprox1a*^{-/-} or WT) identifies a mutant specific cluster and RNA velocity analysis suggests a trajectory between the mutant cluster and the LEC cluster, likely indicative of a fate shift.

d. Stacked bar plot representing the genotype composition of cells in a given phenotypic cluster.

e. Dot plot of marker expression across defined clusters (indicated on Y-axis). Colour scale represents average SCT-normalised expression and point size represents percentage of cells expressing gene.

f. Bar plot of average log fold change in gene expression comparing WT LECs with the Mutant Cluster. Y-axis represents average log fold change, x-axis represents differentially expressed genes (Wilcoxin Rank Sum *adjusted p value* < 0.05) and bars are coloured according to status as a LEC (purple) or VEC (red) marker in **Fig. 1**. Pie charts (right) indicate the LEC and VEC marker composition of genes up-regulated (n=365) and down regulated (n=1,922) in the Mutant Cluster, demonstrating a fate shift.

g. Concordance in the fate shift between Mutant Cluster and WT LEC with the WT LEC and VEC trajectory. Each point represents significant DE genes (n=2,287) between Mutant Cluster and LEC WT, coloured according to LEC or VEC marker status. X-axis represents average log fold change relative to the Mutant Cluster, Y-axis represents average log fold change relative to LEC WT.

h. Lateral confocal images of *cdh5* (green) and *lyve1b* (magenta) expression in the developing trunk in WT and *Zprox1a* mutants at 4dpf.

i. Quantification of *cdh5* intensity in the thoracic duct in WT and mutants (relative to expression in the DA).

j. Lateral confocal images of *kdrl* (green) and *lyve1b* (magenta) expression at 4dpf (left) and 5dpf (right).

k. Quantification of *kdrl* intensity in the thoracic duct in WT and mutants (relative to expression in the PCV) at 4 and 5dpf.

WT, wildtype. Z, zygotic. TD, thoracic duct, DA, Dorsal Aorta. PCV posterior cardinal vein. ISLV, intersomitic lymphatic vessel. DLLV, Dorsal longitudinal lymphatic vessel. PL, parachordal LEC. Scale bars, 50 μm. *, The error bars represent mean ± s.e.m.; *p=0.01 and **p=0.0094 from an unpaired, two-sided t-test.

Figure 4: Single nuclei ATAC seq identifies lineage specific regulatory regions in VECs and

LECs

a. UMAP visualization of snATAC-seq from *Zprox1a*$^{-/-}$ mutant and WT endothelial cells at

4dpf (n=3,731 nuclei), coloured by cell phenotype (top) and genotype (middle) (see

Extended data Fig 4 for full dataset). Stacked bar plot (bottom) summarising the

composition of wildtype and mutants in each cluster. A mutant specific cluster is

identified, similar to **Fig 3**.

b. UMAP visualization indicating accessibility of key marker genes (gene accessibility

score with imputation) that confirm predicted cluster phenotypes. LEC markers:

prox1a, cdh6. VEC markers: *cdh5*, *flt1*, *kdrl.*

c. Heatmap of cluster specific accessible peaks defined using DAP analysis, for all

endothelial cells. Colour reflects a column-wise Z-score, rows represent clusters

defined in **Extended data Fig 4a** and columns are peaks.

d. Bar plot of log2 fold change for the differentially accessible peaks (n=4,185 peaks,

Wilcoxon Rank Sum, *FDR* < 0.05) between WT LECs and WT VECs. LEC/VEC markers

identified in scRNA-seq are coloured in red/purple respectively and demonstrate a

strong correlation between chromatin state and the VEC to LEC fate shift. 228/1,561

more accessible peaks in the LEC cluster were associated with LEC markers and

398/2,624 more accessible peaks in VEC cluster associated with VEC markers. Pie

charts (right) summarising the proportion of LEC/VEC markers in the differentially

accessible peaks (top) and the type of peak (bottom).

e. Genome accessibility track of LEC marker *cdh6*. Red bars represent peaks in the

reproducible peak set from snATAC-seq. The peak highlighted blue indicates a

potential enhancer of *cdh6* (*chr2:28150709-28151209*) with significantly more

accessible chromatin in WT LECs compared to WT VECs (Wilcoxon Rank Sum, *FDR* <

0.05).

f. Overall GFP expression of *cdh6* enhancer (*chr2:28150709-28151209*) reporter at 4dpf

(top). Lateral confocal image of *cdh6* enhancer reporter in the trunk at 4dpf (bottom

middle). Co-expression of *cdh6* enhancer reporter (green) and *lyve1b* (red) in the

facial lymphatics (bottom left) and trunk lymphatics (bottom right) at 5dpf.

g. Vascular TF motifs (n=10 from top n=50) enriched in peaks that are more permissive

in WT LEC compared to WT VEC. The depth of bar colour represents the -

log10(*RawPVal*), y-axis displays individual motifs and schematic (left) and x-axis

represents the percentage of target regions enriched for the motif in the n=1561 peak

set more open in WT LEC.

h. Degree-sorted gene regulatory network displaying known TF binding at genes with

more permissive chromatin in WT LECs compared to WT VECs. TFs are represented

by red circles (nodes), target genes by white circles (nodes), and known binding of TF

to target by a grey arrow (edges). Nodes with a larger number of edges are more

highly connected (bottom right), and nodes with fewer edges are less connected (top

left).

DAP, differentially accessible peaks. FDR, false discovery rate. snATAC-seq, single nuclei

ATAC-seq. TF, transcription factor.

Figure 5: Zygotic *prox1a* mutants display a unique chromatin accessibility state consistent with increased activity of early blood and blood vascular fate transcription factors

a. Schematic illustrates DAP analyses between Mutant Cluster vs. LEC WT and Mutant

Cluster vs. VEC WT.

b. Dot plots of scRNA-seq (top) and snATAC-seq accessibility (bottom) data summarising

the behaviour of key LEC (*prox1a*, *cdh6*, *tbx1, kdr*) and VEC (*cdh5*, *flt1*, *gata6*) markers

in WT LECs, WT VECs and the Mutant Cluster at 4dpf. LEC genes are upregulated

(scRNA-seq) and chromatin is more permissive (snATAC-seq) in WT LEC, and VEC genes

are upregulated and more permissive in WT VEC. This concordance between gene

expression and chromatin accessibility is lost in the Mutant Cluster. The size of the

dots represents the proportion of cells or nuclei, and colour represents either SCT-

normalised expression or gene score of accessibility.

c.   Bar plot of log2 fold change for the differentially accessible peaks (n=2,688 peaks,
794   Wilcoxon Rank Sum, *FDR* < 0.05) between Mutant Cluster and WT LECs. LEC/VEC
795   markers identified in scRNA-seq are coloured in purple/red respectively and
796   demonstrate regions more open in the Mutant are associated with more vascular than
797   lymphatic genes. Pie charts (bottom) summarising the proportion of LEC/VEC markers
798   in the differentially accessible peaks.

799   **d.**   Heatmaps of accessibility (gene accessibility score) for all genes (n=32,020) in WT LECs,
800   WT VECs and Mutant Cluster showing that mutant cluster cells display a unique
801   chromatin state at many genes. Red box indicates genes that are more permissive in
802   the Mutant Cluster than either WT LEC or WT VEC. Colour indicates level of
803   accessibility.

804   **e.**   Venn diagram (left) indicates all individual peaks with increased in accessibility in the
805   Mutant Cluster vs LEC or VEC, with n=431 DAP commonly more open than in both LEC
806   WT and VEC WT (*FDR* < 0.05, log2 fold change > 1.5). Pie chart (middle) indicates the
807   proportion of peak types for n=431 DAP classified as distal, intronic, promoter and
808   exonic respectively. Dot plot (right) summarizing the scRNA-seq expression level of
809   n=13 genes with DAP more open in the Mutant Cluster at 4dpf, demonstrating that
810   accessibility changes for these genes did not correlate with changes in transcription.
811   The size of the dot represents the proportion of cells that express the markers in the
812   cluster, and colour represents SCT-normalised expression.

813   **f.**   Genome accessibility tracks for key markers with DAP more permissive in the Mutant
814   Cluster: *prox1a, gata2a* and *ets1* (left to right). Red bars represent peaks in the
815   reproducible peak set from snATAC-seq. Blue bars highlight DAP (Wilcoxon Rank Sum,
816   *FDR* < 0.05).

817   **g.**   Top 10 enriched motifs (HOMER analysis, *adjusted p value* < 0.05) in the n=431 peaks
818   that are more open in the Mutant Cluster than WT LEC and VEC. The depth of colour
819   represents the -log10(*RawPVal*), y-axis displays individual motifs and schematics (left),
820   and x-axis represents the percentage of peaks enriched for the motif in the n=431
821   peak set.

822   **h.**   Degree-sorted gene regulatory network displaying known TF binding at genes with
823   more permissive chromatin in Mutant Cluster compared to WT LECs. TFs are
824   represented by red circles (nodes), target genes by white circles (nodes), and known

825   binding of TF to target by a grey arrow (edges). Nodes with a larger number of edges

826   are more highly connected (bottom right), and nodes with fewer edges are less

827   connected (top left).

829 DAP, differentially accessible peaks. FDR, false discovery rate. snATAC-seq, single nuclei

830 ATAC-seq. TF, transcription factor.

Figure 6: maternal zygotic *prox1a;prox1b* double mutants display a specific loss of lymphatic

vessels throughout the developing embryo

    a. Overall morphology of control and *MZprox1a⁻/⁻;MZprox1b⁻/⁻* mutants at 5dpf. Arrows

        indicate oedema around eyes, heart and intestine. Scale bars 500 μm.

    b. Lateral confocal images of zebrafish heads at 4dpf showing endothelial cell nuclei

        (green) and venous and lymphatic vessels (white) in control (upper) and *MZprox1a⁻/⁻*

        ;MZprox1b⁻/⁻ embryos (lower). Facial lymphatics are absent or shorter or missing

        (asterisk) in the mutants (asterisk). Scale bar, 80 μm.

c. Lateral confocal images of zebrafish trunks at 4dpf in control (upper) and *MZprox1a^-/-*
;MZprox1b^-/- mutants (lower), showing absent lymphatic vessels (asterisk) but
retained PLs (arrows in lower). Scale bars, 50 μm.

d. Lateral confocal images of PLs in the horizontal myoseptum at 2, 3 and 4dpf. PLs form
later in *MZprox1a^-/-;MZprox1b^-/-* mutants and accumulate in the horizontal
myoseptum while the PLs of control embryos emigrate the HM by 4dpf. Scale bars,
50 μm.

e. Quantification of lymphatic endothelial cell (LEC) number from 4dpf heads control:
n=19, *MZprox1a^-/-;MZprox1b^-/-* mutants: n=18) and trunks (control: n=29, *MZprox1a^-/-*
;MZprox1b^-/- mutants:n=25). Error bars represent mean ± s.e.m.; **** p < 0.0001,
from an unpaired, two-sided t-test.

f. Quantification of number of PLs per somite in trunks at 2dpf in controls (n=41) and
MZprox1a^-/-;MZprox1b^-/- mutants (n=38), at 3dpf in controls (n=11) and *MZprox1a^-/-*
;MZprox1b^-/- mutants (n=10) and at 4dpf in controls (n=29) and *MZprox1a^-/-*
;MZprox1b^-/- mutants (n=25). Error bars represent mean ± s.e.m.; **** p < 0.0001, *
p < 0.05, from an unpaired, two-sided t-test.

g. Lateral confocal images showing defects in the formation of PLs in mutants upon loss
of *prox1a* compared with controls in the trunk. Genotypes are indicated.

h. Colour coded list of analysed genotypes abbreviated in (**i-k**). Each embryo has a
defined genotype for *prox1b* which is represented in colour code as displayed in j.

i. The number of PLs formed (per somite) at 2dpf in genotypes indicated. Decreasing
gene dosage for *prox1a* and *prox1b* progressively reduced the initial seeding of the
HM by PLs.

j. The number of cells in sprouts departing the PCV is unchanged in mutants showing
that sprouting occurs normally.

k. The number of cells in vISVs is unchanged in mutants indicating no effect on the
venous endothelium.

LL, lymphatic loop. OLV, otholitic lymphatic vessel. LFL, lateral facial lymphatic vessel. MFL,
medial facial lymphatic vessel. LBA, lymphatic branchial arches. DLLV, dorsal longitudinal
lymphatic vessel. PL, parachordal LEC. ISLV, intersomitic lymphatic vessel. TD, thoracic duct.
DA, dorsal aorta. PCV, posterior cardinal vein.

Figure 7: Single cell profiling of maternal zygotic *prox1a;prox1b* double mutants reveals that

Prox1 homologues initially suppress gene expression to drive a fate transition

a. UMAP visualisation of endocardial, venous and arterial endothelial cells (Level 02
n=5,347) coloured according to predicted cell phenotype (left), genotype (middle) and
RNA-velocity (right).

b. Heatmap displaying expression of phenotype specific genes defined using differential
expression analysis. Columns are cells grouped by phenotype assignment, rows are
genes, and colour indicates the average level of expression.

c. Dot plot indicating expression of key markers used to define cell phenotype. Colour
scale represents average SCT-normalised expression and point size represents
percentage of cells expressing gene.

d. UMAP visualisation of venous endothelial cells (Level 03 n=2,747) coloured according
to predicted cell phenotype (top) and genotype (bottom).

e. UMAP visualisation of key marker gene expression. Colour scale represents SCT-
normalised expression.

f. Bar plot indicating average log fold change of n=1,186 significantly different genes
between all *MZprox1a/b^{-/-}* and WT venous endothelial cells (LEC VEC 01, LEC VEC 02
and PCV combined) at 40hpf (Wilcoxin Rank Sum *adjusted p value* < 0.05). Colour
indicates genes associated with one or more mitochondrial, mRNA processing and
blood and blood vascular GO terms.

g. Bar plot summarizing GO BP analysis of the n=1,137 genes upregulated in the
MZprox1a/b^{-/-} venous endothelial cells. Y-axis represents enriched GO BP term, x-axis
represents the -log10(*adjusted p value*) and bars are coloured according to fold
enrichment.

h. Dot plot of n=36 key blood vascular markers upregulated in the *MZprox1a/b^{-/-}* venous
endothelial cells, indicating genotype specific expression in LEC VEC and PCV cell
phenotypes. Colour scale represents average SCT-normalised expression and point
size represents percentage of cells expressing gene.

i. STRING analysis of n=36 key blood vascular markers upregulated in the *MZprox1a/b^{-/-}*
venous endothelial cells.

j. Degree-sorted gene regulatory network displaying known TF binding at n=1,137 genes
upregulated in the *MZprox1a/b^{-/-}* venous endothelial cells. TFs are represented by red
circles (nodes), target genes by white circles (nodes), and known binding of TF to

target by a grey arrow (edges). Nodes with a larger number of edges are more highly

connected (bottom right), and nodes with fewer edges are less connected (top left).

 EXTENDED DATA FIGURES AND FIGURE LEGENDS

Extended data Fig 1: Full dataset and analysis for data in Fig 1.

a. UMAP visualization of n=35634 endothelial cells sequenced for the single cell atlas of lymphangiogenesis (n=6 samples; see also **Extended data table 1a** for markers used to identify clusters) coloured according to developmental stage (top left), cell cycle phase (top right), predicted cell phenotype (bottom left), and cluster solution (RNA-snn-res.1).

b. Clustree analysis demonstrating the relationship between different cluster resolutions for the n=9771 venous and lymphatic endothelial cells in Figure 1a. Resolution 0.4 (boxed) was used for all downstream analyses.

c. UMAP visualization of cluster resolution 0.4 (as appears in **Fig 1**).

d. Stacked bar plot of resolution 0.4 cluster composition cell cycle phase.

e. Stacked bar plot of resolution 0.4 cluster composition developmental stage.

925   **f.** Venn diagram of significant genes commonly upregulated in LEC compared to VEC at

926   3, 4 and 5dpf.

Extended data Fig 2: Full dataset and analysis for data in Fig 3.

a. UMAP visualization of n=8075 endothelial cells sequenced comparing lymphangiogenesis in WT and *Zprox1a* mutants at 4dpf (n=2 samples; see also **Extended data table 2a** for markers used to identify clusters) coloured according to developmental stage (top left), cell cycle phase (top right), predicted cell phenotype (bottom left), and cluster solution (SCT-snn-res.8)

b. Clustree analysis demonstrating the relationship between different cluster resolutions

for the n=3063 venous and lymphatic endothelial cells in **Figure 3a**. Resolution 0.2

(boxed) was used for all downstream analyses.

c. UMAP visualization of cluster resolution 0.2

d. Stacked bar plot of resolution 0.2 cluster composition genotype

e. Stacked bar plot of resolution 0.2 cluster composition cell cycle phase

f. Venn diagram indicates the overlap between genes significantly upregulated in VEC

compared to LEC at each developmental stage of the single cell atlas, and those

upregulated in the *Zprox1a$^{-/-}$* Mutant Cluster at 4dpf compared to WT LECs. N=85

genes (coloured red) are uniquely associated with the Mutant Cluster and WT VEC at

4dpf, compared to N=4 genes in WT VEC at 3 and 5dpf.

Extended data Fig 3: Additional phenotypic analysis of blood vascular markers in _Zprox1a_−/−

mutants associated with Fig 3

a. Lateral confocal images of zebrafish trunks at 4dpf showing _cdh5_ expression (green)

and _lyve1b_ expression in lymphatic vessels (magenta) in control (upper) and _Zprox1a_

mutant embryos (lower). Transgenic lines are indicated. Scale bar, 50 μm.

b. Quantification of _lyve1b_ expression intensity in the TD relative to the PCV in sibling

and _Zprox1a_ mutant embryos at 4dpf.

c. Quantification of *cdh5* expression intensity in the TD relative to the DA in sibling and
Zprox1a mutant embryos at 4dpf.

d. Lateral confocal images of zebrafish trunks at 4dpf showing *kdrl* expression (green)
and *lyve1b* expression in lymphatic vessels (magenta) in control (upper) and *Zprox1a*
mutants (lower). Transgenic lines are indicated. Scale bars, 50 μm.

e. Quantification of *kdrl* expression intensity in the TD relative to the DA in sibling and
Zprox1a mutant embryos at 4dpf and 5dpf (upper).

f. Lateral confocal images of zebrafish trunks at 5dpf showing *kdrl* expression (green)
and *lyve1b* expression in lymphatic vessels (magenta) in control (upper) and *Zprox1a*
mutants (lower). Transgenic lines are indicated. Scale bars, 50 μm.

g. Quantification of the number of TD segments displaying *kdrl* expression in *Zprox1a*
mutant embryos at 4dpf and 5dpf (lower).

DLLV, dorsal longitudinal lymphatic vessel. ISLV, intersomitic lymphatic vessel. TD, thoracic
duct. DA, Dorsal aorta. PCV, posterior cardinal vein. vISV, venous intersegmental vessel. PL,
Parachoral lymphatic endothelial cells.

Extended data Fig 4: Additional data and analysis for data in Fig 4.

a. UMAP visualization of n=3,731 single nuclei ATAC-seq data comparing lymphangiogenesis in WT and *Zprox1a* mutants at 4dpf. Nuclei coloured according to

cluster (Res-0.8, left), predicted cell phenotype (middle) and genotype (right). Cell

phenotype was predicted based on accessibility changes at the genes defined in

Extended data table 3a.

b. Stacked bar plot indicating the genotype composition of nuclei captured per cell type.

c. Density plot indicating the majority of snATAC-seq peaks fall around the TSS. Y-axis

represents distance from TSS (bp) and y-axis represents normalised Tn5 insertion

profile.

d. Heatmap of cluster specific accessible peaks for all cell clusters. Colour reflects a

column-wise Z-score, each row represents a cluster defined in **Extended Data Fig. 4a**,

columns are peaks.

e. Genome accessibility track of key VEC markers *cdh5* (top) and *flt1* (bottom) indicating

these genes are more accessible in WT VEC than WT LEC cell phenotypes. Red bars

represent peaks in the reproducible peak set from snATAC-seq. DAP between WT VECs

and WT LECs (Wilcoxon Rank Sum, *FDR* < 0.05) are highlighted blue.

f. Genome accessibility track of key LEC markers *prox1a* (top) and *cdh6* (bottom)

indicating these genes are more accessible in WT LEC than WT VEC cell phenotypes.

Red bars represent peaks in the reproducible peak set from snATAC-seq. DAP between

WT LECs and WT VECs (Wilcoxon Rank Sum, *FDR* < 0.05) are highlighted blue.

g. Heatmaps of accessibility (gene score) for DEG (**Fig 3f**) upregulated (left, n=300

mapped genes) and downregulated (right, n=1,052 mapped genes) in the Mutant

Cluster compared to WT LECs. Colour indicates row-wise Z-score, each row represents

a differentially expressed gene in scRNA-seq data, columns are snATAC-seq clusters.

Boxed genes downregulated in Mutant display more permissive chromatin in Mutant

than WT LEC, showing that mutant cluster cells display a unique chromatin state at

many genes.

h. Confocal projections of zebrafish head (left) labelled with *Tg(empty:EGFP;*

XCA:DsRed2) at 5 dpf, showing GFP expression in neuronal tissues. Confocal

projections of zebrafish trunk (right) labelled with *Tg(empty:EGFP; XCA:DsRed2)* at 5

dpf, showing GFP expression in neuronal tissues. All scale bars = 100μm.

1004

1005

Extended data Fig 5: **Additional phenotypic analysis of *MZprox1a⁻/⁻;MZprox1b⁻/⁻* mutants**

associated with Fig 6

a. Schematic explanation of the germline replacement transplantation method used to

generate *MZprox1a⁻/⁻;MZprox1b⁻/⁻* mutants.

b-c. Lateral confocal images showing vascular defects in control and mutant in the

craniofacial (**b**) and trunk (**c**) regions of the larvae at 4dpf. Genotypes and transgenic labels

are indicated. *MZprox1a-/-* and *MZprox1a+/-, MZprox1b-/-* animals both showed a strong

loss of lymphatic vessels in craniofacial and trunk regions. The *MZprox1a-/-, MZprox1b-/-*

mutants showed a robust loss of craniofacial vessels equivalent to the other mutant

phenotypes but a more severe loss of trunk lymphatics and an accumulation of PLs in the HM.

1016

    d. Genotypes that are shown in the quantification. Colour codes are used to identify individual embryonic genotypes in the dot plots showing quantification in **e-g**.

    e. Quantification of the number of craniofacial (head) LECs within facial lymphatic vessels showing evidence for contributions from *prox1a* and *prox1b*.

    f. Quantification of the number of mural LECs (muLECs) showing a major role for *prox1a* and a role for *prox1b* in muLEC development.

    g. Quantification of the number of trunk LECs within vessels showing a dominant role for *prox1a* in trunk lymphangiogenesis.

Z, zygotic; MZ, maternal and zygotic; HM, horizontal myoseptum; LL, lymphatic loop; OLV, otholitic lymphatic vessel; LFL, lateral facial lymphatic vessel; MFL, medial facial lymphatic vessel; LBA, lymphatic branchial arches; DLLV, dorsal longitudinal lymphatic vessel; PL, parachordal LEC; ISLV, intersomitic lymphatic vessel; TD, thoracic duct; DA, dorsal aorta; PCV, posterior cardinal vein.

1030

1031

Extended data Fig 6: **Embryonic transplantation data demonstrating endothelial cell**

autonomy of *prox1a* maternal zygotic mutant phenotype associated with Fig 6.

a. Schematic showing the endothelial cell transplantation approach used to test cell

autonomy for the vascular phenotypes seen in *MZprox1a* mutants. Cell identities were

traced following transplantation in unlabelled host embryos using expression of the

pan -endothelial marker *fli1a* and venous/lymphatic marker *lyve1b* (indicated).

b. Examples of 5dpf wildtype hosts engrafted with wildtype donor cells (upper) that

contribute to arteries (aISVs, DA), veins (PCV, vISVs) and lymphatic vessels (TD, ISLVs)

and examples of wildtype embryos engrafted with *MZprox1a* mutant cells

contributing arteries (aISVs, DA) and veins (vISV, PCV) but not lymphatic vessels. Note

the PL accumulation in the horizontal myoseptum..

c. The number of embryos imaged and graft sizes scored post-tranplantation. As the

contribution to the different EC types is affected by the graft size, we analysed grafts

of comparable sizes and position from wildtype and mutant embryos. Grafts were

categorised as large, medium or small. Small (spanning 1-2 somites), medium

(spanning 2-3 somites) and large grafts (spanning 4-5 somites). The number of ECs

contributing to each EC type was scored at single cell resolution and the percentage

of grafts with a contribution to each lineage calculated accordingly. The numbers

within the different graft groups indicate the number of analysed embryos within that

group.

d. The percentage of embryos with vascular grafts that display contributions to arteries,

veins and lymphatics. *MZprox1a$^{-/-}$* mutant cells contribute to arteries and veins but

showed a near complete loss of contribution to lymphatics. The size of the grafts did

not influence the outcome. *Z1b** indicates a mixed prox1b background. p < 0.0001

from an unpaired, two-sided t-test.

DA, dorsal aorta. PCV, posterior cardinal vein. PL, parachordal LEC. TD, thoracic duct. aISV,

arterial intersegmental vessel. vISV, venous intersegmental vessel. ISLV, intersegmental

lymphatic vessel.

1060

1061
Extended data Fig 7: Additional data and analysis for data in Fig 7.

a. UMAP visualization of n=26,289 endothelial cells sequenced comparing
lymphangiogenesis in WT and *MZprox1a/b* mutants at 40hpf (n=4 samples; see also
Extended data table 5a for markers used to identify clusters) coloured according to
cluster solution (SCT-snn-res.8, left), predicted phenotype per cluster (middle), and
predicted cell phenotype (right). Endothelial cells selected for further analysis are
indicated by a red circle.

b. Dot plot summarizing the scRNA-seq expression level of n=20 key genes used to
predict cell phenotypes of each cluster in the complete dataset (Level 01), for each
cluster defined in **Extended Data Fig. 7a** (SCT-snn-res.8). The size of the dot
represents the proportion of cells that express the markers in the cluster, and colour
represents SCT-normalised expression.

c. Dot plot summarizing the scRNA-seq expression level of n=20 key genes used to
predict cell phenotype in the complete dataset. The size of the dot represents the
proportion of cells that express the markers in the cluster, and colour represents SCT-
normalised expression.

d. Clustree analysis demonstrating the relationship between different cluster resolutions
for the n=5,347 endocardial, venous and arterial endothelial cells in **Fig. 7a** and
Extended Data Fig. 7a (Level 02). Cluster resolution SCT_snn_0.7 (boxed) was selected
for all downstream analyses.

e. UMAP visualization of cluster resolution SCT_snn_0.7 (left), predicted phenotype of
each cluster (middle) and phenotypic group (right).

f. Stacked bar plot of cluster resolution SCT_snn_0.7 cluster composition genotype (top)
and cell cycle phase (bottom).

g. Dot plot summarizing the scRNA-seq expression level of n=14 key genes used to
predict cell phenotype in the endothelial cell populations (Level 02, n=5,347 cells).
The size of the dot represents the proportion of cells that express the markers in the
cluster, and colour represents SCT-normalised expression.

References

1   Oliver, G., Kipnis, J., Randolph, G. J. & Harvey, N. L. The Lymphatic Vasculature in
the 21(st) Century: Novel Functional Roles in Homeostasis and Disease. *Cell* **182**,
270-296, doi:10.1016/j.cell.2020.06.039 (2020).
2   Koltowska, K., Betterman, K. L., Harvey, N. L. & Hogan, B. M. Getting out and
about: the emergence and morphogenesis of the vertebrate lymphatic vasculature.
Development **140**, 1857-1870, doi:10.1242/dev.089565 (2013).
3   Baek, S. *et al.* The Alternative Splicing Regulator Nova2 Constrains Vascular Erk
Signaling to Limit Specification of the Lymphatic Lineage. *Dev Cell* **49**, 279-292
e275, doi:10.1016/j.devcel.2019.03.017 (2019).

4 Koltowska, K. *et al.* Vegfc Regulates Bipotential Precursor Division and Prox1
Expression to Promote Lymphatic Identity in Zebrafish. *Cell Rep* **13**, 1828-1841,
doi:10.1016/j.celrep.2015.10.055 (2015).
5 Deng, Y., Atri, D., Eichmann, A. & Simons, M. Endothelial ERK signaling controls
lymphatic fate specification. *The Journal of clinical investigation* **123**, 1202-1215,
doi:10.1172/JCI63034 (2013).
6 Shin, M. *et al.* Vegfc acts through ERK to induce sprouting and differentiation of
trunk lymphatic progenitors. *Development* **143**, 3785-3795, doi:10.1242/dev.137901
(2016).
7 Srinivasan, R. S. *et al.* The Prox1-Vegfr3 feedback loop maintains the identity and the
number of lymphatic endothelial cell progenitors. *Genes Dev* **28**, 2175-2187,
doi:10.1101/gad.216226.113 (2014).
8 Oliver, G. & Srinivasan, R. S. Endothelial cell plasticity: how to become and remain a
lymphatic endothelial cell. *Development* **137**, 363-372, doi:10.1242/dev.035360
(2010).
9 Wigle, J. T. & Oliver, G. Prox1 function is required for the development of the
murine lymphatic system. *Cell* **98**, 769-778, doi:10.1016/s0092-8674(00)81511-1
(1999).
10 Hagerling, R. *et al.* A novel multistep mechanism for initial lymphangiogenesis in
mouse embryos based on ultramicroscopy. *EMBO J* **32**, 629-644,
doi:10.1038/emboj.2012.340 (2013).
11 Klotz, L. *et al.* Cardiac lymphatics are heterogeneous in origin and respond to injury.
Nature **522**, 62-67, doi:10.1038/nature14483 (2015).
12 Martinez-Corral, I. *et al.* Nonvenous origin of dermal lymphatic vasculature. *Circ Res*
116, 1649-1654, doi:10.1161/CIRCRESAHA.116.306170 (2015).
13 Stanczuk, L. *et al.* cKit Lineage Hemogenic Endothelium-Derived Cells Contribute to
Mesenteric Lymphatic Vessels. *Cell Rep* **10**, 1708-1721,
doi:10.1016/j.celrep.2015.02.026 (2015).
14 Srinivasan, R. S. *et al.* Lineage tracing demonstrates the venous origin of the
mammalian lymphatic vasculature. *Genes Dev* **21**, 2422-2432,
doi:10.1101/gad.1588407 (2007).
15 Srinivasan, R. S. *et al.* The nuclear hormone receptor Coup-TFII is required for the
initiation and early maintenance of Prox1 expression in lymphatic endothelial cells.
Genes Dev **24**, 696-707, doi:10.1101/gad.1859310 (2010).
16 Yang, Y. *et al.* Lymphatic endothelial progenitors bud from the cardinal vein and
intersomitic vessels in mammalian embryos. *Blood* **120**, 2340-2348,
doi:10.1182/blood-2012-05-428607 (2012).
17 Stone, O. A. & Stainier, D. Y. R. Paraxial Mesoderm Is the Major Source of
Lymphatic Endothelium. *Dev Cell* **50**, 247-255 e243,
doi:10.1016/j.devcel.2019.04.034 (2019).
18 Johnson, N. C. *et al.* Lymphatic endothelial cell identity is reversible and its
maintenance requires Prox1 activity. *Genes Dev* **22**, 3282-3291,
doi:10.1101/gad.1727208 (2008).
19 Tao, S. *et al.* Zebrafish prox1b mutants develop a lymphatic vasculature, and prox1b
does not specifically mark lymphatic endothelial cells. *PLoS One* **6**, e28934,
doi:10.1371/journal.pone.0028934 (2011).
20 van Impel, A. *et al.* Divergence of zebrafish and mouse lymphatic cell fate
specification pathways. *Development* **141**, 1228-1238, doi:10.1242/dev.105031
(2014).

21   Hogan, B. M. *et al.* Ccbe1 is required for embryonic lymphangiogenesis and venous
1155   sprouting. *Nat Genet* **41**, 396-398, doi:10.1038/ng.321 (2009).
1156   22   Yaniv, K. *et al.* Live imaging of lymphatic development in the zebrafish. *Nat Med* **12**,
1157   711-716, doi:10.1038/nm1427 (2006).
1158   23   Eng, T. C. *et al.* Zebrafish facial lymphatics develop through sequential addition of
1159   venous and non-venous progenitors. *EMBO Rep* **20**, doi:10.15252/embr.201847079
1160   (2019).
1161   24   Okuda, K. S. *et al.* lyve1 expression reveals novel lymphatic vessels and new
1162   mechanisms for lymphatic vessel development in zebrafish. *Development* **139**, 2381-
1163   2391, doi:10.1242/dev.077701 (2012).
1164   25   Bussmann, J. *et al.* Arteries provide essential guidance cues for lymphatic endothelial
1165   cells in the zebrafish trunk. *Development* **137**, 2653-2657, doi:10.1242/dev.048207
1166   (2010).
1167   26   Cha, Y. R. *et al.* Chemokine signaling directs trunk lymphatic network formation
1168   along the preexisting blood vasculature. *Dev Cell* **22**, 824-836,
1169   doi:10.1016/j.devcel.2012.01.011 (2012).
1170   27   Jung, H. M. *et al.* Development of the larval lymphatic system in zebrafish.
1171   *Development* **144**, 2070-2081, doi:10.1242/dev.145755 (2017).
1172   28   Bower, N. I. *et al.* Mural lymphatic endothelial cells regulate meningeal angiogenesis
1173   in the zebrafish. *Nat Neurosci* **20**, 774-783, doi:10.1038/nn.4558 (2017).
1174   29   van Lessen, M. *et al.* Intracellular uptake of macromolecules by brain lymphatic
1175   endothelial cells during zebrafish embryonic development. *Elife* **6**,
1176   doi:10.7554/eLife.25932 (2017).
1177   30   Venero Galanternik, M. *et al.* A novel perivascular cell population in the zebrafish
1178   brain. *Elife* **6**, doi:10.7554/eLife.24369 (2017).
1179   31   Kuchler, A. M. *et al.* Development of the zebrafish lymphatic system requires
1180   VEGFC signaling. *Curr Biol* **16**, 1244-1248, doi:10.1016/j.cub.2006.05.026 (2006).
1181   32   Hogan, B. M. & Schulte-Merker, S. How to Plumb a Pisces: Understanding Vascular
1182   Development and Disease Using Zebrafish Embryos. *Dev Cell* **42**, 567-583,
1183   doi:10.1016/j.devcel.2017.08.015 (2017).
1184   33   He, Z., Brazovskaja, A., Ebert, S., Camp, J. G. & Treutlein, B. CSS: cluster similarity
1185   spectrum integration of single-cell genomics data. *Genome Biol* **21**, 224,
1186   doi:10.1186/s13059-020-02147-4 (2020).
1187   34   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902
1188   e1821, doi:10.1016/j.cell.2019.05.031 (2019).
1189   35   Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP.
1190   *Nat Biotechnol*, doi:10.1038/nbt.4314 (2018).
1191   36   La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498,
1192   doi:10.1038/s41586-018-0414-6 (2018).
1193   37   Thomas, P. D. *et al.* Applications for protein sequence-function evolution data:
1194   mRNA/protein expression analysis and coding SNP scoring tools. *Nucleic Acids Res*
1195   **34**, W645-650, doi:10.1093/nar/gkl229 (2006).
1196   38   Wigle, J. T. *et al.* An essential role for Prox1 in the induction of the lymphatic
1197   endothelial cell phenotype. *EMBO J* **21**, 1505-1513, doi:10.1093/emboj/21.7.1505
1198   (2002).
1199   39   Kang, J. *et al.* An exquisite cross-control mechanism among endothelial cell fate
1200   regulators directs the plasticity and heterogeneity of lymphatic endothelial cells.
1201   *Blood* **116**, 140-150, doi:10.1182/blood-2009-11-252270 (2010).

40   Bessa, J. *et al.* Zebrafish enhancer detection (ZED) vector: a new tool to facilitate
transgenesis and the functional analysis of cis-regulatory regions in zebrafish. *Dev*
Dyn **238**, 2409-2417, doi:10.1002/dvdy.22051 (2009).

41   Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors
prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell*
38, 576-589, doi:10.1016/j.molcel.2010.05.004 (2010).

42   Rouillard, A. D. *et al.* The harmonizome: a collection of processed datasets gathered
to serve and mine knowledge about genes and proteins. *Database (Oxford)* **2016**,
doi:10.1093/database/baw100 (2016).

43   Kulkarni, R. M., Greenberg, J. M. & Akeson, A. L. NFATc1 regulates lymphatic
endothelial development. *Mech Dev* **126**, 350-365, doi:10.1016/j.mod.2009.02.003
(2009).

44   Norrmen, C. *et al.* FOXC2 controls formation and maturation of lymphatic collecting
vessels through cooperation with NFATc1. *J Cell Biol* **185**, 439-457,
doi:10.1083/jcb.200901104 (2009).

45   Dieterich, L. C. *et al.* DeepCAGE Transcriptomics Reveal an Important Role of the
Transcription Factor MAFB in the Lymphatic Endothelium. *Cell Rep* **13**, 1493-1504,
doi:10.1016/j.celrep.2015.10.002 (2015).

46   Koltowska, K. *et al.* mafba is a downstream transcriptional effector of Vegfc
signaling essential for embryonic lymphangiogenesis in zebrafish. *Genes Dev* **29**,
1618-1630, doi:10.1101/gad.263210.115 (2015).

47   Chen, L. *et al.* Tbx1 regulates Vegfr3 and is required for lymphatic vessel
development. *J Cell Biol* **189**, 417-424, doi:10.1083/jcb.200912037 (2010).

48   Nicenboim, J. *et al.* Lymphatic vessels arise from specialized angioblasts within a
venous niche. *Nature* **522**, 56-61, doi:10.1038/nature14425 (2015).

49   Cha, B. *et al.* Complementary Wnt Sources Regulate Lymphatic Vascular
Development via PROX1-Dependent Wnt/beta-Catenin Signaling. *Cell Rep* **25**, 571-
584 e575, doi:10.1016/j.celrep.2018.09.049 (2018).

50   Szklarczyk, D. *et al.* The STRING database in 2017: quality-controlled protein-
protein association networks, made broadly accessible. *Nucleic Acids Res* **45**, D362-
D368, doi:10.1093/nar/gkw937 (2017).

51   Armour, S. M. *et al.* An HDAC3-PROX1 corepressor module acts on HNF4alpha to
control hepatic triglycerides. *Nat Commun* **8**, 549, doi:10.1038/s41467-017-00772-5
(2017).

52   Liu, X. *et al.* Mitotic Implantation of the Transcription Factor Prospero via Phase
Separation Drives Terminal Neuronal Differentiation. *Dev Cell* **52**, 277-293 e278,
doi:10.1016/j.devcel.2019.11.019 (2020).

53   Qin, J. *et al.* Prospero-related homeobox (Prox1) is a corepressor of human liver
receptor homolog-1 and suppresses the transcription of the cholesterol 7-alpha-
hydroxylase gene. *Mol Endocrinol* **18**, 2424-2439, doi:10.1210/me.2004-0009 (2004).

54   Song, K. H., Li, T. & Chiang, J. Y. A Prospero-related homeodomain protein is a
novel co-regulator of hepatocyte nuclear factor 4alpha that regulates the cholesterol
7alpha-hydroxylase gene. *J Biol Chem* **281**, 10081-10088,
doi:10.1074/jbc.M513420200 (2006).

55   Takeda, Y. & Jetten, A. M. Prospero-related homeobox 1 (Prox1) functions as a novel
modulator of retinoic acid-related orphan receptors alpha- and gamma-mediated
transactivation. *Nucleic Acids Res* **41**, 6992-7008, doi:10.1093/nar/gkt447 (2013).

56   Lawson, N. D. & Weinstein, B. M. In vivo imaging of embryonic vascular
development using transgenic zebrafish. *Dev Biol* **248**, 307-318,
doi:10.1006/dbio.2002.0711 (2002).

57   Ciruna, B. *et al.* Production of maternal-zygotic mutant zebrafish by germ-line
replacement. *Proc Natl Acad Sci U S A* **99**, 14919-14924,
doi:10.1073/pnas.222459999 (2002).

58   Westerfield, M. *The Zebrafish Book. A Guide for the Laboratory use of Zebrafish
(Danio rerio)*. 4 edn, (University of Oregorn Press, 2000).

59   Bussmann, J. & Schulte-Merker, S. Rapid BAC selection for tol2-mediated
transgenesis in zebrafish. *Development* **138**, 4327-4332, doi:10.1242/dev.068080
(2011).

60   Okuda, K. S., Baek, S. & Hogan, B. M. Visualization and Tools for Analysis of
Zebrafish Lymphatic Development. *Methods Mol Biol* **1846**, 55-70, doi:10.1007/978-
1-4939-8712-2_4 (2018).

61   Bower, N. I. *et al.* Vegfd modulates both angiogenesis and lymphangiogenesis during
zebrafish embryonic development. *Development* **144**, 507-518,
doi:10.1242/dev.146969 (2017).

62   Kartopawiro, J. *et al.* Arap3 is dysregulated in a mouse model of hypotrichosis-
lymphedema-telangiectasia and regulates lymphatic vascular development. *Hum Mol
Genet* **23**, 1286-1297, doi:10.1093/hmg/ddt518 (2014).

63   Zheng, G. X. *et al.* Massively parallel digital transcriptional profiling of single cells.
Nat Commun **8**, 14049, doi:10.1038/ncomms14049 (2017).

64   Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of
Cell Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291 e289,
doi:10.1016/j.cels.2018.11.005 (2019).

65   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nat
Biotechnol* **36**, 411-420, doi:10.1038/nbt.4096 (2018).

66   McCarthy, D. J., Campbell, K. R., Lun, A. T. & Wills, Q. F. Scater: pre-processing,
quality control, normalization and visualization of single-cell RNA-seq data in R.
Bioinformatics **33**, 1179-1186, doi:10.1093/bioinformatics/btw777 (2017).

67   Zappia, L. & Oshlack, A. Clustering trees: a visualization for evaluating clusterings at
multiple resolutions. *Gigascience* **7**, doi:10.1093/gigascience/giy083 (2018).

68   Kejie Li, Z. O., Dongdong Lin, Michael Mingueneau, Will Chen, David Sexton,
Baohong Zhang. cellxgene VIP unleashes full power of interactive visualization,
plotting and analysis of scRNA-seq data in the scale of millions of cells. *bioRxiv*
2020.08.28.270652, doi:https://doi.org/10.1101/2020.08.28.270652 (2020).

69   Smedley, D. *et al.* The BioMart community portal: an innovative alternative to large,
centralized data repositories. *Nucleic Acids Res* **43**, W589-598,
doi:10.1093/nar/gkv350 (2015).

70   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of
biomolecular interaction networks. *Genome Res* **13**, 2498-2504,
doi:10.1101/gr.1239303 (2003).

1292

1293

1294